1

Title: Better null models for assessing predictive accuracy of disease models
Running Head: Null Models for Disease Modeling

Article Type: Letter

Authors: Alexander C. Keyel[1,2] and A. Marm Kilpatrick[3]

[1] Division of Infectious Diseases, Wadsworth Center, New York State Department of Health, Albany, NY,

United States of America

[2] Department of Atmospheric and Environmental Sciences, University at Albany, SUNY, Albany, NY,

United States of America

Contact information: ACK: alexander.keyel@health.ny.gov; 120 New Scotland Road, Albany, NY 12208.

+1 518-485-6466, fax: 518-473-1326, AMK: akilpatr@ucsc.edu

[3] Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA

Statement of Authorship: ACK and AMK designed the study, drafted and revised the manuscript, and

performed the analyses.

Data Accessibility Statement: No new data were used in this study. Data used in this manuscript are

available from the Centers for Disease Control.

Abstract Word Count: 144

Manuscript Word Count: 1808

Text Box Word Count: 333 + 293 = 626

Number of References: 27

Number of Figures: 2

Number of Tables: 1

Number of Text Boxes: 2

Number of Supplemental Tables: 3

Abstract

Null models provide a critical baseline for the evaluation of predictive disease models. Many studies consider only the grand mean null model (i.e. R^2) when evaluating the predictive ability of a model, which is often misleading in conveying the predictive power of a model. We evaluated ten null models for human cases of West Nile virus (WNV), a zoonotic mosquito-borne disease introduced to the United States in 1999. The Negative Binomial, Historical (i.e. using previous cases to predict future cases) and Always Absent null models were the strongest overall, and the majority of null models significantly outperformed the grand mean. Somewhat surprisingly, the length of the training timeseries did not strongly affect the performance of most null models. We argue that a combination of null models is needed to evaluate the forecasting performance of predictive models and the grand mean is the lowest bar.

Keywords: arbovirus, continuous ranked probability score, ecological forecasting, forecast, mosquito, null model, probabilistic, statistics, vector-borne disease, West Nile virus

Introduction

Forecasting infectious disease dynamics is a key challenge for the 21[st] century (Johansson *et al.* 2019). Climate and land use change, combined with the introduction of pathogens to new regions, has created an urgent need for predicting future disease threats (Kilpatrick & Randolph 2012). Large data sets and new modeling and statistical techniques have opened up possibilities for ecological forecasting (Dietze 2017). A key step in the evaluation of predictive models is assessing their improvement over null models. The use of null models to provide a baseline in the absence of specific mechanisms has a long history in ecology (Gotelli & Graves 1996). Such baselines are important, as in some cases, predictive models may appear to be informative, but may be no better than a simple and uninformative null model

(Olden *et al.* 2002; Beale *et al.* 2008). For example, when dealing with rare events, if a predictive model

is outperformed by a null model that predicts the event to never occur, it is not providing much useful

information about the process being studied (Olden *et al.* 2002).

West Nile Virus (WNV) is an excellent system in which to examine null models in a probabilistic context.

WNV is a flavivirus that cycles between mosquito and avian populations (Work *et al.* 1955; Komar *et al.*

2003; Kilpatrick 2011). WNV was introduced to the United States (US) in 1999 (Lanciotti *et al.* 1999) and

rapidly spread to the conterminous US and throughout the Americas (Kramer *et al.* 2019). As a

nationally notifiable disease in the US, long-term data sets (>20 years) exist on human cases (CDC

2019b). Many models have been built for predicting WNV risk (Barker 2019). Most studies of WNV, and

many other pathogens, have included only a very simplistic null model (e.g. R^2, which uses the grand

mean of the training data) for assessment of model accuracy.

Our aim was to examine a range of null models (Box 1) to provide guidance on null model selection and

performance in disease forecasting for locations with frequent (≥50% of years with disease) and

infrequent cases (disease present, but <50% of years). We tested 10 null models using the number of

WNV cases in each county in the US in each year in a probabilistic framework. Where cases were

frequent and timeseries were long, we hypothesized the Negative Binomial model would perform the

best due to its ability to model count distributions with a variable rate parameter. Where cases were

infrequent and time series were short, we predicted that no models would significantly outperform the

Always Absent mode.

Material and methods

Data set

We compared the accuracy of 10 null models using the CDC Neuroinvasive WNV Case records (Source:

ArboNET, Arboviral Diseases Branch, Centers for Disease Control and Prevention, Fort Collins, Colorado;

contact the CDC for data access). This is a national data set of the number of WNV neuroinvasive disease

cases in each county in each year from 3108 counties in the conterminous United States (US) from 2000

– 2021. We used WNV neuroinvasive cases, because there is less variability across different states in

detection of these cases compared to WNV fever cases.

We divided the data set into two groups: 159 counties that have had 11 or more years with WNV

(frequent WNV set), and 1880 counties that had 1 – 10 years with WNV (infrequent WNV set). The 1069

counties that never had WNV cases were excluded to avoid zero-inflation. The first year a state reported

a case of WNV (per (Kilpatrick 2011)) was used as the first year of training data for all counties within

that state. As a result, the number of counties included in the analysis increased over time (S1 Table).

Model predictions were made using at least 4 years of training data. We used the Continuous Ranked

Probability Score, a probabilistic scoring approach that can evaluate a distribution of predicted

outcomes (Box 2). Population data for each year for the incidence-based null models came from the

United States Census Bureau (US Census Bureau 2017, 2019). Population data from 2019 were used for

2020 and 2021 as well, due to missing data for these years.

We also tested whether our model results were sensitive to the length of time series for selected

models. Model years were selected at random (without replacement) to use as training data to predict a

randomly selected focal year. This allowed us to disentangle length of time series from the specific order

of observation of results. However, models that required a temporal structure were excluded from this

analysis (i.e. Prior Year and Autoregressive). The Incidence and Pooled Incidence models were also

excluded from this analysis as they used the prior-year's population for converting incidence to case

counts. Only data from 2005 and later were used in this analysis to ensure that WNV had already been

established in all counties.

Null Models:

The ten null models are described in Box 1. Note that using case counts versus incidence does not make

a large difference to the outcome when stratifying by county, because population is relatively consistent

from one year to another. However, the choice of incidence or case counts as the model basis does lead

to different outcomes when pooling across counties with different population sizes.

Scoring Method

We used the Continuous Ranked Probability Score (CRPS), which is a proper scoring method (Jordan *et*

al. 2019; Bracher *et al.* 2021) We chose the CRPS rather than the Logarithmic Score because the former

scores forecasts based on the distance from each predicted probability to the observation, whereas the

latter only scores whether an observation is within a bin or outside of a bin, with no consideration of

how far outside the bin the prediction was (Matheson & Winkler 1976; Hersbach 2000; Wilks 2011). As

the CRPS scores are based on distance from the observed value, the models above were allowed to

predict fractional cases of WNV (e.g. if the mean number of cases was 2.5 cases, that would be used as

the prediction rather than rounding up or down to the nearest whole number). For null models that

required sampling from a probabilistic distribution, we used 100 random draws. Data analyses were

performed in R (R Core Team 2017). Code for running the null models is available via the probnulls

package on GitHub (www.github.com/akeyel/probnulls/R/NullModels.R).

Results

The Mean Value null model (R^2) that is frequently used as the baseline for prediction accuracy was

among the weakest of the null models (Fig. 1). It performed worse than 5 null models (significantly

worse than 4; Fig. 1) for frequent WNV counties and worse than 6 null models (significantly worse than

5) for infrequent WNV counties (Fig. 1). In contrast, the Negative Binomial null model was significantly

better than other null models for predicting neuroinvasive cases of WNV in the frequent WNV analysis

(Fig. 1a, paired t-tests using a Holm correction for multiple comparisons). The Negative Binomial was

also significantly better than eight other null models (all except the Always Absent model which was

equally accurate) in the infrequent WNV analysis (Fig. 1b). The Negative Binomial was the top model in 8

individual years (out of 18) for the frequent WNV analysis, and in 9 of 18 years for the infrequent WNV

analysis (Table 1). The Historical Null model also performed very well in both frequent and infrequent

WNV analyses across all years (Fig 1), and outperformed all other models in 5 individual years for

frequent WNV counties (Table 1). Finally, in the infrequent WNV analysis, the Always Absent model was

tied for the best model for all years combined, and outperformed seven other models (Fig 1b) and was

the best model for 8 individual years (Table 1).

The length of the training time series had only weak effects on null model performance (Fig. 2). For

infrequent WNV counties there was no support for an improvement in model score with the length of

the training time series (Table S2). For frequent WNV counties, model score improved with the length of

the training time series for four of the six models examined, but the effect was small relative to the

difference between most models (Table S3). For the two remaining models, increasing the length of the

training time series had no effect on model score or actually made the score worse (Table S3).

Discussion

At least five null models significantly outperformed a county-based grand mean and many did far better

(Figs. 1, 2). A grand mean calculated for the entire US (Pooled Mean model) performed even worse.

Thus, when evaluating the performance of new statistical or mechanistic models of disease incidence,

there are far better null models than the grand mean (i.e. R^2). These null models can be easily calculated

for time-series data (e.g., using the probnulls package from GitHub in R), and our results suggest that the

length of time series was not critical for developing a robust null model across a range of 4-16 years. The

Negative Binomial and Historical nulls were the strongest null models overall, with the Always Absent

null performing well where disease cases were infrequent. The strong performance of the Always

Absent null in regions where WNV was infrequent (statistically tied with Negative Binomial, Fig. 1; top

model in 8 of 18 years, Table 1) is a reminder that basic accuracy statistics for rare events can appear

high.

The structure and scale of the underlying data may affect the performance of the different null models.

The WNV data set here does not have a clear temporal trend. A strong temporal trend would likely have

changed which model performed the best. Specifically, null models that use the recent past to predict

future cases (e.g. autoregressive models) would perform much better. Seasonal patterns, as examined in

recent dengue forecasts (Johansson *et al.* 2019), could also affect which null model performs best.

Future work could explore the performance of different models under different magnitudes of temporal

trend and stochastic variation. Additionally, county-annual scales may be more relevant to academic

study than to vector control and public health responses (Keyel *et al.* 2021). Research on null model

performance is needed at finer spatial and temporal scales.

Conclusion

We strongly recommend the inclusion of multiple null models when testing predictive models of vector-

borne diseases. A grand mean calculated from the training data set is an inadequate null model given

the suite of probabilistic alternatives available. The Negative Binomial and Historical nulls performed

especially well for WNV and simple autoregressive models performed moderately well and would likely

perform even better for data with temporal trends. Negative Binomial and Historical null models

performed well both when WNV cases were frequent and when they were infrequent, and their

performance did strongly not depend on the length of the training time series. Researchers proposing

mechanistic models should determine if their models are an improvement over a simple statistical

description of historical patterns.

Acknowledgements

We thank L. F. Chaves for constructive discussion. This publication was supported by cooperative

agreement 1U01CK000509-01, funded by the Centers for Disease Control and Prevention and by the

National Institutes of Health grant R01AI168097 and National Science Foundation grants DEB 1911853,

DEB-1717498 and CNH-1115069. Its contents are solely the responsibility of the authors and do not

necessarily represent the official views of the Centers for Disease Control and Prevention or the

Department of Health and Human Services.

References

Barker, C.M. (2019). Models and Surveillance Systems to Detect and Predict West Nile Virus Outbreaks. *J
Med Entomol*, 56, 1508–1515.
Beale, C.M., Lennon, J.J. & Gimona, A. (2008). Opening the climate envelope reveals no macroscale
associations with climate in European birds. *Proceedings of the National Academy of Sciences*,
105, 14908–14912.
Bracher, J., Ray, E.L., Gneiting, T. & Reich, N.G. (2021). Evaluating epidemic forecasts in an interval
format. *PLOS Computational Biology*, 17, e1008618.
CDC. (2019a). Final Cumulative Maps & Data for 1999-2018.
CDC. (2019b). Nationally notifiable arboviral diseases reported to ArboNET: Data release guidelines.
Dietze, M. (2017). *Ecological Forecasting*. Princeton University Press.

Gotelli, N.J. & Graves, G.R. (1996). *Null models in ecology*.
Hersbach, H. (2000). Decomposition of the continuous ranked probability score for ensemble prediction
systems. *Weather and Forecasting*, 15, 559–570.
Johansson, M.A., Apfeldorf, K.M., Dobson, S., Devita, J., Buczak, A.L., Baugher, B., *et al.* (2019). An open
challenge to advance probabilistic forecasting for dengue epidemics. *Proceedings of the*
National Academy of Sciences, 116, 24268–24274.
Jordan, A., Krüger, F. & Lerch, S. (2019). Evaluating Probabilistic Forecasts with scoringRules. *Journal of*
Statistical Software, 90, 1–37.
Keyel, A.C., Gorris, M.E., Rochlin, I., Uelmen, J.A., Chaves, L.F., Hamer, G.L., *et al.* (2021). A proposed
framework for the development and qualitative evaluation of West Nile virus models and their
application to local public health decision-making. *PLOS Neglected Tropical Diseases*, 15,
e0009653.
Kilpatrick, A.M. (2011). Globalization, land use, and the invasion of West Nile virus. *Science*, 334, 323–
327.
Kilpatrick, A.M. & Randolph, S.E. (2012). Drivers, dynamics, and control of emerging vector-borne
zoonotic diseases. *LANCET*, 380, 1946–1955.
Komar, N., Langevin, S., Hinten, S., Nemeth, N., Edwards, E., Hettler, D., *et al.* (2003). Experimental
infection of North American birds with the New York 1999 strain of West Nile virus. *Emerging*
infectious diseases, 9, 311.
Kramer, L.D., Ciota, A.T. & Kilpatrick, A.M. (2019). Introduction, Spread, and Establishment of West Nile
Virus in the Americas. *Journal of Medical Entomology*, 56, 1448–1455.
Kwan, J.L., Kluh, S. & Reisen, W.K. (2012). Antecedent Avian Immunity Limits Tangential Transmission of
West Nile Virus to Humans. *PLOS ONE*, 7, e34127.
Lanciotti, R.S., Roehrig, J.T., Deubel, V., Smith, J., Parker, M., Steele, K., *et al.* (1999). Origin of the West
Nile virus responsible for an outbreak of encephalitis in the northeastern United States. *Science*,
286, 2333–2337.
Matheson, J.E. & Winkler, R.L. (1976). Scoring rules for continuous probability distributions.
Management Science, 22, 1087–1096.
Olden, J.D., Jackson, D.A. & Peres-Neto, P.R. (2002). Predictive Models of Fish Species Distributions: A
Note on Proper Validation and Chance Predictions. *null*, 131, 329–336.
R Core Team. (2017). *R: A Language and Environment for Statistical Computing*. R Foundation for
Statistical Computing, Vienna, Austria.
Ripley, B.D. (2002). Time series in R 1.5.0. *R News*, 2, 2–7.
Smith, K.H., Tyre, A.J., Hamik, J., Hayes, M.J., Zhou, Y. & Dai, L. (2020). Using Climate to Explain and
Predict West Nile Virus Risk in Nebraska. *GeoHealth*, 4, e2020GH000244.
US Census Bureau. (2017). Intercensal estimates of the resident population for counties and states: April
1, 2000 to July 1, 2010. Suitland, MD: US Census Bureau. Retrieved from:
https://www.census.gov/data/datasets/time-series/demo/popest/intercensal-2000-2010-
counties.html.
US Census Bureau. (2019). Population, Population Change, and Estimated Components of Population
Change: April 1, 2010 to July 1, 2019 (CO-EST2019-alldata). Suitland, MD: US Census Bureau.
Retreived from: https://www.census.gov/data/tables/time-series/demo/popest/2010s-
counties-total.html.
Venables, W.N. & Ripley, B.D. (2002). *Modern Applied Statistics with S*. Fourth. Springer, New York.
Wilks, D.S. (2011). *Statistical Methods in the Atmospheric Sciences*. Academic Press.
Work, T.H., Hurlbut, H.S. & Taylor, R. (1955). Indigenous Wild Birds of the Nile Delta as Potential West
Nile Virus Circulating Reservoirs. *The American Journal of Tropical Medicine and Hygiene*, 4,
872–888.

Box 1. Meet the 10 Null Models

Always Absent Null (non-probabilistic): WNV is always predicted to be absent (i.e., the model always predicts 0 cases or incidence).

Pooled Mean Value Null[1] (non-probabilistic): WNV is predicted to have its mean value of cases.

Mean Value Null (non-probabilistic): As in the Pooled Mean Value Null, but stratified by county.

Prior Year Null (non-probabilistic): The results from the prior year are used to predict the current year (Smith *et al.* 2020).

Historical Null: Prior observations for the location are sampled with replacement, creating a probabilistic distribution of future cases.

Pooled Incidence Null: The number of cases were predicted by sampling a binomial distribution, using incidence as the probability, and the county's population as the size parameter. Incidence was calculated for the entire region of interest (i.e. the United States).

Incidence Null: As in the Pooled Incidence Null, but incidence is calculated stratified by county, allowing the model to capture local hotspots of disease.

Negative Binomial Null[2]: We estimated the mean and dispersion parameters of a negative binomial distribution using the human case counts in each year from each county Random draws for each county were then drawn using the estimated mean and dispersion parameter.

Autoregressive AR1 Null[3]: An autoregressive model was fit. The model produced a mean estimate for the next time step and a normally-distributed error around that mean. We randomly sampled the normal distribution around the mean. All values less than zero were assigned to 0, as negative human cases are not possible.

Uniform Null: Random draws were taken from a uniform distribution of cases bounded by 0 to the maximum number of cases observed in a county.

[1] Model predictions would differ if a pooled mean incidence null were used. However, the Pooled Incidence model represents a more sophisticated version of that null model, hence we did not evaluate a non-probabilistic pooled mean incidence model.
[2] Using the MASS package in R (Venables & Ripley 2002).
[3] Using the arima function in R (Ripley 2002).

Box 2. A quick introduction to
probabilistic models

A probabilistic forecast can convey
substantially more information than a
point forecast. Here, (a) a dashed line
indicates the point estimate, while the
histogram of 1000 samples from a
negative binomial distribution with μ = 5
(mean) and size = 100 (dispersion
parameter) show the expected range of
possible outcomes. (b) Note that
because the CRPS score is based on
distance between the observed value
and the entire distribution of the
prediction, it can give non-intuitive
results where a distribution that includes
the observed value (Dist. 2) can score
worse than a distribution that assigns
the observation a zero or near-zero
probability (Dist. 1). (c) Negative
binomial fits for a county with a high
number of annual neuroinvasive West Nile virus (WNV) cases and (d) a county with very few WNV cases.
Shading shows data used to train the models (2002 - 2012) and unshaded shows testing data used to
evaluate the models (2013 - 2019). The contribution of each year to the overall CRPS score is given in
the table below for the two locations for three null models: Always Absent, Historical, and Negative
Binomial. The CRPS value for the Always Absent Null is also the number of WNV cases for that year
because the CRPS becomes the mean absolute error when applied to a point estimate and the Always
Absent Null always predicts 0 cases. Lower CRPS scores indicate better model performance, with 0 being
a perfect prediction. Here, Historical performed the best on average for Cook, IL, while Always Absent
performed the best on average for Barbour, AL. Visually, the Negative Binomial fit for Cook County
appears poor, due to over-estimation in the lowest case bin, and under-estimation of most of the
remaining bins.

Year	Cook, IL			Barbour, AL		
	Always Absent	Historical	Negative Binomial	Always Absent	Historical[1]	Negative Binomial[1]
2013	46	15.9	16.1	0	0.008	0.008
2014	23	8.3	14.6	0	0.008	0.008
2015	21	8.2	14.8	0	0.008	0.008
2016	65	24.9	21.9	0	0.008	0.008
2017	41	14.1	15.3	0	0.008	0.008
2018	76	31.5	26.7	0	0.008	0.008
2019	9	11.3	19.4	0	0.008	0.008
Mean	40.1	16.3	18.4	0	0.008	0.008

286 [1] By chance, these two models predicted the same distribution of zero and one case occurrences.
287 Different random draws would cause these scores to vary slightly.

289 **Table 1.** Frequency of each model having the lowest CRPS score for a year in counties where WNV is
290 frequent (>50% of time series) or infrequent (present < 50% of the time series).

Model	Frequent WNV	Infrequent WNV
Negative Binomial	8	9
Historical Null	5	0
Always Absent	2	8
Incidence	2	1
Autoregressive	1	0
Mean Value + 4 others[1]	0	0

291 [1] Mean Value, Pooled Mean Value, Pooled Incidence, Uniform, and Prior Year null models were never
292 the top model in either analysis

Fig. 1. Continuous Ranked Probability Scores (CRPS) for 2004 – 2021 for 10 null models for a) counties

with at least 50% of the time series with WNV ("Frequent") and b) for counties with at least one case,

but cases < 50% of the time series ("Infrequent"). The mean CRPS scores was calculated across all

counties for each model and year, and the plot shows the median value of these mean annual CRPS

scores by model, with the box showing the 25% and 75% quartiles, whiskers corresponding to +/- 1.5

times the Interquartile range, and circles corresponding to values outliers outside this range.

Fig. 2. The Negative Binomial and Historical were generally the top two models, independent of length of time series used to train the models for both a) the counties with frequent WNV and c) the counties with infrequent WNV cases. Individual training replicates showed a range of variation in CRPS score (b, d) depending on which replicates were selected (only Negative Binomial and Historical null models shown). Training years were randomly selected from the entire time series, and a random focal year was selected for evaluation. Only a subset of null models was evaluated over time.

310 **S1 Table.** Number of counties used for the null model analysis by prediction year for 2004 – 2021. After

311 2008, the sample size remained constant for all following years as all states had the minimum 4 years of

312 training data with WNV by that point.

Year	N (frequent WNV)	N (infrequent WNV)
2004	25	213
2005	79	1042
2006	154	1831
2007	159	1865
2008 – 2021	159	1880

313
314

315 **Table S2. Analysis of the length of the training time series on the CRPS score for six null models for**
316 **infrequent WNV counties. An additive model of null model and time series length fit better than an**
317 **interactive model (ΔAIC = 50).**

Predictor	Estimate	SE	t value	P-value
Intercept (Neg Binomial)	0.18	0.015	12.18	<2e-16
Length of time series	0.00072	0.0011	0.68	0.50
Always absent	0.014	0.014	1.01	0.31
Historical	0.0047	0.014	0.34	0.73
Mean	0.13	0.014	9.82	<2e-16
Pooled mean	0.17	0.014	12.09	<2e-16
Uniform	0.26	0.014	18.94	<2e-16

319 **Table S3. Analysis of the length of the training time series on the CRPS score for six null models for**
320 **frequent WNV counties. A model with an interaction between model and time series length fit better**
321 **than an additive model (ΔAIC = 21).**

Predictor	Estimate	SE	t value	P-value
Intercept (Neg Binomial)	4.61	0.51	9.09	< 2e-16
Length of time series	-0.13	0.044	-2.90	0.0038
Always absent	1.68	0.72	2.35	0.019
Historical	0.29	0.72	0.41	0.68
Mean	1.39	0.72	1.94	0.053
Pooled mean	1.61	0.72	2.25	0.025
Uniform	-0.52	0.72	-0.73	0.47
Length*Always absent	0.0070	0.062	0.11	0.91
Length*Historical	-0.021	0.062	-0.35	0.73
Length*Mean	0.0076	0.062	0.12	0.90
Length*Pooled mean[1]	0.090	0.062	1.46	0.15
Length*Uniform	0.27	0.062	4.31	1.86E-05

322 [1] If the analysis is releveled to the Pooled Mean model, the length of time series variable is not
323 significant for this model (p = 0.40).