Interactive effects of herbivory and the level and fluctuations of nutrient

availability on dominance of alien plants in synthetic native communities

Yanjun Li[1,2], Yingzhi Gao[2], Mark van Kleunen[3,4], Yanjie Liu[1,3*]

[1]Key Laboratory of Wetland Ecology and Environment, Northeast Institute of

Geography and Agroecology, Chinese Academy of Sciences, Changchun, 130102,

China

[2]Key Laboratory of Vegetation Ecology, Northeast Normal University, Changchun,

130024, China

[3]Ecology, Department of Biology, University of Konstanz, 78464 Konstanz, Germany

[4]Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and

Conservation, Taizhou University, Taizhou 318000, China

*Email of corresponding author: liuyanjie@iga.ac.cn +86 431 82536096

Running title: Herbivory mediates nutrient-change effects on plant invasion

Abstract

Numerous studies have highlighted the role of nutrient availability and fluctuations therein for invasion success of alien plants. Others also highlighted the role of herbivores in invasion success. However, how herbivory and the level and fluctuations in nutrient availability interact in driving alien plant invasion into resident communities remains largely unexplored.

We grew eight invasive alien species as target species in pot☐mesocosms with five different synthetic native communities in a three-factorial design with two levels of nutrient availability (low *vs* high), two levels of nutrient fluctuations (constant *vs* pulsed) and two levels of herbivory (with *vs* without).

The relative biomass production of the alien target plants decreased in response to an increase in nutrient availability, and increased in response to the presence of herbivores. Furthermore, herbivory could interact with changes in nutrient availability and nutrient fluctuations to affect the dominance of the alien target species (a marginally significant interaction; 90% CIs: [0.125, 2.712]).

Our multispecies experiment indicates that herbivory could mediate the interactive effect of nutrient enrichment and variability in nutrient supply on invasion of alien plants into native communities. Therefore, we recommend that studies testing the fluctuating resources hypothesis should also consider interactive effect of other trophic levels.

Keywords: biological invasions, exotic plants, global change, plant-herbivore interactions, resource variability, trophic level

Introduction

Invasion by alien plants could reduce native biodiversity, influence ecosystem functions, and degrade ecosystem services [1]. Due to the rapid globalization, the increase in the number of naturalized alien plant species does not show any sign of saturation, and it was recently predicted that their numbers may increase on average by 18% from 2005 to 2050 [2-4]. Therefore, a major research objective in the field of ecology is to identify the mechanisms that underlie alien plant invasion [5-9].

It has frequently been suggested that an increase in soil-nutrient availability is one of the most important drivers of alien plant invasion [8, 10, 11]. As successful alien plant species are often introduced from anthropogenic and more nutrient-rich environments, they might be more likely to be adapted to high-nutrient environments [12, 13]. Indeed, both a recent meta-analysis [14] and a globally replicated study of 64 grasslands [15] showed that successful alien plants respond more strongly to nitrogen enrichment than native plants do. So far, empirical studies that tested how soil-nutrient changes affect alien plant invasion mainly focused on changes in the mean nutrient level [16-22]. However, due to increasing occurrences of extreme events (i.e., droughts, floodings, heat waves, fires), soil-nutrient changes also entail changes in their variability, and this may affect plant invasions [20, 23]. Therefore, it is important to test how changes in soil-nutrient levels, as well as fluctuations in nutrient availability over time, drive alien plant invasion in resident communities.

The fluctuating resource hypothesis proposes that temporal fluctuations in nutrient supply could promote alien plant invasion in resident communities [10]. However, empirical studies testing the hypothesis found mixed results. For example, Parepa et al. found that a pulsed nutrient supply, compared to a constant nutrient supply, increased the dominance of the invasive plants *Fallopia japonica* and *F.* ×

bohemica in experimental plant communities [23]. In contrast, Liu et al. showed that a

pulsed nutrient supply decreased the dominance of invasive alien plants [24]. Thus,

more studies are needed to test the hypothesis, and why the results might vary. The

studies that tested the effect of temporal changes in nutrient availability did so under

overall nutrient-rich conditions. Under more nutrient-limiting conditions, however,

the effect of temporal fluctuations may be even stronger. In other words, mean

nutrient availability may interact with temporal fluctuations in nutrient availability to

affect alien plant invasion into resident communities. However, very few studies have

tested whether this expectation holds [25].

Although the fluctuating resource hypothesis has become a key theory in

invasion ecology, previous tests only used study systems consisting of a single trophic

level (i.e., only considered plant-plant interactions). Plant growth, however, can be

strongly regulated by other trophic levels, such as herbivores. This might be relevant

for the fluctuating resource hypothesis as alien species are likely to be released from

most of their native enemies, and thus should suffer less herbivory than native species

in their introduced regions [26-28]. Following this logic, the presence of herbivores,

just like increases in resource availability and fluctuations therein [20, 23, 29, 30],

could promote alien plant invasion in resident communities. Moreover, as herbivore

effects on plants are often regulated by soil-nutrient availability, because plants

growing in relatively high-nutrient conditions might be better able to compensate or

tolerate herbivory [31-34], the effects of herbivores might interact with the effects of

nutrients. Therefore, it is reasonable to assume that the presence of herbivores might

amplify the positive effect of increases in resource availability and fluctuations on

alien plant invasion.

To test the individual effects of nutrient availability, nutrient fluctuations,

herbivory, and their interactions on alien plant invasion into resident communities, we

grew eight invasive alien species as target species in pot□mesocosms with five

different synthetic native communities, each consisting of three grassland species.

Then, we exposed the plants to eight combinations of two nutrient availability (low *vs*

high), two nutrient-fluctuation (constant *vs* pulsed) and two herbivory (with *vs*

without) treatments. By comparing the absolute aboveground biomass production of

the alien target species as well as their biomass production relative to the biomass

production of the native competitors, we addressed the following specific questions:

1) Do nutrient availability, nutrient fluctuations and the presence of herbivores

promote the absolute and relative biomass of alien plants? 2) Does the effect of

nutrient fluctuations on absolute and relative biomass of alien plants depend on the

overall nutrient availability level? 3) Does the presence of herbivores interact with

increases in nutrient availability and fluctuations therein to affect the absolute and

relative biomass of alien plants?

Material and methods

Study species

To investigate the individual and interactive effects of nutrient availability, nutrient fluctuations and the presence of herbivores on alien plant invasion into resident communities, we chose eight invasive alien species as targets, and 15 native species as native community members from the herbaceous flora of China (Table S1). We classified these species as invasive alien or native to China based on information in the book "The Checklist of the Alien Invasive Plants in China" [35] and the Flora of China database (www.efloras.org). To cover a wide taxonomic breadth, we selected the eight alien target species from seven genera of three families. As plants with different life histories (i.e., annuals or perennials) may respond differently to nutrient availability and fluctuations [20, 36], we assured that both the alien targets and native community members included annuals and perennials (Table S1). Seeds of these species were collected in natural populations in grasslands in China or ordered from a commercial seed company (see Table S1).

To impose a herbivory treatment, we selected two aboveground insect herbivores. As natural systems usually include both generalist and specialist insect herbivores, we chose the generalist grasshopper *Stenocatantops splendens* and the specialist grasshopper *Locusta migratoria* (i.e., grass-feeder) as the shoot herbivores. The grasshoppers were acquired from a commercial insect company (Cangzhou Grasshoppers Breeding Center, China). As both grasshoppers and all plant species occur mainly in grasslands, and according to the GBIF database (www.gbif.org), overlap in their distributions, they are very likely to co-occur in nature.

Pre-cultivation and experimental setup

We conducted the experiment at the Northeast Institute of Geography and

Agroecology, Chinese Academy of Sciences (43°5'49"N, 125°24'40"E). From 7 May

to 26 June 2020, we sowed each of the invasive alien and native species separately

into plastic circular trays (diameter = 25.5 cm, height = 4 cm) filled with potting soil

(Pindstrup Plus, Pindstrup Mosebrug A/S, 103 Denmark; pH 6; 120 mg/L N; 12 mg/L

P; 400 mg/L K; 28 mg/L Mg; 0.4 mg/L B; 2 mg/L Mo; 1042 mg/L Cu; 3 mg/L Mn;

0.9 mg/L Zn; 8 mg/L Fe). Because the time required for germination varies among the

species, we sowed them at different times (Table S1) to ensure that at transplanting

the seedlings were in a similar developmental stage. All trays with seeds were kept in

a greenhouse (temperature: 18-27^{o}C; natural lighting with an intensity of ~75% of the

light outdoors; and ~68% relative humidity).

On 10 July 2020, we selected similar-sized seedlings from each of the eight

invasive alien and 15 native species, and transplanted them into 2.5 L circular plastic

pots (top diameter: 18.5 cm, bottom diameter: 12.5 cm, height: 15 cm) filled with a

1:1 mixture of sand and fine vermiculite. We transplanted one seedling of an alien

target species in the center of each pot. For each of the eight alien target species, we

transplanted a total of 40 seedlings into 40 pots (i.e., one individual per pot), resulting

in total of 320 pots. Immediately after transplanting the alien target species, we

equally distributed the 40 pots of each alien target over five different native

communities (i.e., eight pots per native community). To create the five different

native grassland communities, we randomly assigned the 15 native species into five

groups of three species (Table S1). We planted two seedlings of each native

community member, so that each pot included six individuals of native species

equally spaced in a circle (diameter = 10 cm) around the alien target seedling. The

two individuals of the same species were planted at opposite positions of the circle

(Fig. 1a). After transplanting, we randomly assigned all pots to two cages (3.5 × 4.5 ×

2.5 m) located outside of the pre-cultivation greenhouse. The cages were covered with

transparent plastic roofs and white nylon net (mesh size: 0.25 × 0.25 mm) all around.

Half of the pots of each combination of an alien target species and a specific native

community were assigned to one cage, and the remaining ones to the other cage (Fig.

1a). In other words, each cage included 160 pots in total (8 alien target species × 5

native communities × 4 nutrient-supply treatments [2 nutrient availability × 2 nutrient

fluctuation treatments]). To avoid the loss of water and nutrient solution, we put a

plastic tray under each pot. We re-randomized the positions of the pots in each cage

after five weeks.

On 14 July 2020 (i.e., one week after transplanting), we started to apply the

nutrient treatments at weekly intervals for a total of 10 weeks (Fig. 1b). We applied

two nutrient availability treatments (low *vs* high) crossed with two nutrient-fluctuation

treatments (constant *vs* pulsed), using a 400%￼strength Hoagland solution (Methods

S1). During the experiment, we added a total of 50 and 400 ml of the Hoagland

solution to the low and high-nutrient availability pots, respectively. Although it is

difficult to compare the absolute nutrient levels in a pot experiment to those found in

a natural system, Liu et al. previously showed that both the low- and high-nutrient

availabilities are limiting plant growth, as is usually the case in nature [24]. Within

each nutrient-availability treatment, we created two different nutrient-supply patterns,

a constant nutrient supply and a pulsed nutrient supply (Fig. 1b). We supplied 5 ml of

the nutrient solution for the constant low-nutrient supply each week, and 40 ml of the

same nutrient solution for the constant high-nutrient supply. The pulsed treatment at

low-nutrient availability consisted of three weeks of 2 ml per week, followed by four

weeks of 9.5 ml per week, and again three weeks of 2 ml per week (Fig. 1b). The

pulsed treatment at high-nutrient availability consisted of three weeks of 10 ml a week,

followed by four weeks of 85 ml, and again three weeks of 10 ml (Fig. 1b). To avoid

differences in water supply among the four treatments, we added extra water to the

amount of nutrient solution in each treatment to ensure that each pot received a total

of 85 ml of water per nutrient application. In each cage, there were five replicates per

alien target species for each of the four nutrient-supply treatments (i.e., one replicate

for each of the five native communities). We watered all plants regularly by filling the

dish under each pot to avoid water limitation.

The herbivory treatment started on 14 August 2020, and ended six weeks later on

20 September 2020. We added the grasshoppers in one of the two cages, and treated

the other cage as control (Fig. 1a). As the commercial company hatched these two

species of grasshoppers at different times, we added *Locusta migratoria* from 14

August to 6 September 2020, and *Stenocatantops splendens* from 31 August to 6

September 2020. We checked the herbivory pressure every day to determine whether

we would add more grasshoppers. During the experiment, we added *Locusta*

migratoria seven times (three times for 3–4th instars and four times for adults), and

Stenocatantops splendens two times (one time for 3–4th instars and the other time for

adults). In total, we added 354 individuals of *Locusta migratoria*, and 450 individuals

of *Stenocatantops splendens* in the herbivory-treatment cage.

Measurements

On 25 September 2020 (i.e., eleven weeks after transplanting), we harvested the

aboveground biomass of all pots. For each pot, we first harvested the alien target

species and then the three native competitor species. As some alien target and native

plants died and three pots had accidentally been treated with the wrong nutrient

solution, we only harvested 274 instead of 320 pots (see the raw data on

https://doi.org/10.5061/dryad.fj6q573vn) [37]. We did not harvest the roots of the

alien and native species, because the roots of the species were intertwined, and it was

impossible to separate them. All aboveground biomass samples were dried at 65°C for

72 hours and then weighed. We calculated total aboveground biomass per pot by

summing the biomass of the alien target species and the three native competitors. We

also calculated the biomass proportion of the alien target species in each pot as the

ratio between the biomass of the alien target species and the total aboveground

biomass per pot.

Statistical analysis

To test the effects of nutrient availability (low *vs* high), nutrient fluctuations

(constant *vs* pulsed), herbivory treatments (with *vs* without) and their interactions on

aboveground biomass production of the alien target species, biomass production of

the native communities and biomass proportion of the alien target species in each pot,

we fitted Bayesian multilevel models using the function brm of the R package brms

[38] in R 4.0.2 [39]. In all models, we included the following explanatory variables

nutrient availability (low *vs* high), nutrient fluctuations (constant *vs* pulsed) and

herbivory treatment (with *vs* without) as fixed factors. To account for phylogenetic

non-independence of species belonging to the same family and for non-independence

of replicates of the same species, we included identity of the alien target species

nested in their family as random factors in all models. To account for variation among

the five different native communities, we also included identity of the native

community as random factor in all models. To relax the homogeneity of variance

assumption in all models, we allowed the residual standard deviation sigma to vary by

the identity of alien target species [40].

For all models, we used the default priors set by the brms package, and ran four

independent chains. The number of total iterations per chain was 5000, and the

number of warm-up samples was 2500. To directly test hypotheses about the main

effects and interactive effects based on each coefficient's posterior distribution, we

used the sum coding, which effectively 'centers' the effects to the grand mean (i.e.,

the mean value of all data observations) [41]. To implement this in brms, we used the

functions contrasts and contr.sum of the stats package in R. We considered the fixed

effects nutrient availability, nutrient fluctuation and herbivory treatments, and their

interactions as significant when their 95% credible interval of the posterior

distribution did not overlap zero, and as marginally significant when the 90% credible

interval did not overlap zero. As we had only two cages available, one with herbivores

and one without herbivores, our herbivory treatment is obviously pseudoreplicated

[42]. This means that the main effect of herbivory should be interpreted with care

[43]. However, it still allows us to test whether the effects of the nutrient availability

and nutrient fluctuation treatments differ between the cage with and without

herbivores.

Results
Averaged across the nutrient fluctuation and herbivory treatments, an increase in

nutrient availability significantly increased the biomass production of alien target

species (+600.7%; Table 1; Fig. 2 and Fig. S1) and the biomass production of native

communities (+601.0%; Table 1; Fig. 2 and Fig. S1). The presence of herbivores,

however, significantly decreased the biomass production of native communities

(-39.4%; Table 1; Fig. 2 and Fig. S2). The similar pattern was found for the biomass

production of alien target species (-44.3%; Table 1; Fig. 2 and Fig. S2), although it

was only marginally significant (90% CIs: [-0.170, -0.010]). The negative effect of

herbivory on the biomass production of both alien target species and native

communities was stronger under high-nutrient availability (alien: -46.4%; native:

-39.7%) than under low-nutrient availability (alien: -28.0%; native: -36.6%;

significant NA × H interactions in Table 1; Fig. 2 and Fig. S3).

We found that an increase in nutrient availability significantly decreased (-6.5%;

Table 1; Fig. 2 and Fig. S1), whereas the presence of herbivores significantly

increased the relative biomass production of the alien target species in the native

communities (+3.6%; Table 1; Fig. 2 and Fig. S2). Additionally, under low-nutrient

availability, the pulsed nutrient supply tended to increase the relative biomass

production of the alien target plants in the absence of herbivores (+7.9%), whereas the

reverse was true for plants in the presence of herbivores (-18.8%; Fig. 2). Under

low-nutrient availability, however, pulsed nutrient supply and the presence of

herbivores did not affect the relative biomass production of the alien target species

(Fig. 2). The dependency of the effect of the nutrient fluctuation treatment on the

levels of the other treatments was indicated by a marginally significant interaction

between herbivory, nutrient availability and fluctuation therein (90% CIs: [0.125,

2.712]) in Table 1.

Discussion

Our multispecies experiment showed that an increase of nutrient availability promoted the biomass production for both the alien target species and native communities. However, herbivory suppressed the biomass production of plants, in particular under high-nutrient availability. We found that increases in nutrient availability suppressed, whereas herbivory promoted the dominance of the alien target species in native resident communities. Interestingly, we also found tentative evidence that herbivory could interact with changes in nutrient availability and nutrient fluctuations to affect the dominance of the alien target species. In other words, herbivory could mediate the interactive effect of nutrient enrichment and fluctuations in nutrient supply on alien plant invasion into native communities.

Although each pot had one invasive alien and six native plants, the invasive plant accounted for about one quarter of the aboveground biomass in each pot (Fig. 2c). This suggests that the invasive alien species were more dominant than the native species. We also found that this dominance decreased with an increase of the average nutrient availability. This finding does not support the idea that increased nutrient availability could promote alien plant invasion in resident communities, although many theoretical [10, 44] and empirical studies [14, 15, 20, 23, 25] found evidence for this. A potenial reason could be that in our study the native species themselves are also quite common, and it are usually the rare species that take less advantage of increased nutrient availability [45].

Previous empirical studies testing the nutrient-fluctuation effect on alien plant invasion found mixed results [20, 23-25, 46, 47]. We hypothesized that this might be because the effect of temporal fluctuations may be even stronger under more nutrient-limiting conditions than under less nutrient-limiting conditions. However, we

found no significant interactive effect of nutrient availability and fluctuations therein

on the aboveground biomass and dominance of alien target plants in native

communities. This is in line with the results of a recent case study by Gao et al. [25].

It is worth noting though that in the absence of herbivory, the absolute and relative

aboveground biomass of the invasive alien plants tended to be higher in the low,

pulsed nutrient treatment than in the low, constant treatment, while the reverse was

true in the presence of herbivory.

Not surprisingly, herbivory decreased the biomass production of plants (also see

the total biomass production per pot in Fig. S4a and Fig. S4b). In particular, the

biomass suppression of herbivory was stronger under high-nutrient availability than

under low-nutrient availability (Fig. S4c). This seems inconsistent with previous

findings that plants compensate or tolerate herbivory more when growing in high

nutrient conditions [31, 32, 48-51]. However, it should be noted that in our

experiment, the herbivores could choose between the plants. As the plants grown at

high-nutrient availabilities might be more nutritious [52-55], and have decreased plant

secondary metabolite concentrations such as tannins [56, 57], the herbivores might

have preferably fed on the plants with high nutrient availabilities [53, 58-61]. On the

other hand, we found that herbivory increased the dominance of the alien target

species in the native resident communities. This is also not surprising, because

according to the enemy release hypothesis, alien plants often have escaped from many

of their herbivorous enemies, and therefore could in their introduced range

outcompete resident plants [26].

When growing under low-nutrient availability and also in the absence of

herbivores, the nutrient pulse promoted the dominance of alien plants, which supports

the fluctuating resource hypothesis [10]. However, when growing under low-nutrient

availability but in the presence of herbivores, the nutrient pulse suppressed the

dominance of alien plants in pots. This may because herbivores often reduce the

abundance (biomass, cover) of dominant species [62, 63]. On the other hand, under

high-nutrient availablity, the nutrient pulse and the herbivory treatment hardly

affected the dominance of alien plants. One plausibility explanation for this finding is

that the overall high-nutrient availability reduces, or even cancels, the

nutrient-limitation shifts over time caused by nutrient fluctuations [20, 24, 64],

resulting in very weak effects of the nutrient pulse on dominance changes in plant

communities [65]. In addition to the weak evidence that herbivory mediated the

effects of changes in nutrient availability and fluctuations on alien plant invasion, a

recent case study also found that the parasitic plant *Cuscuta australis* could also

regulate the effects of nutrient availablity and fluctuations on the invasion success of

the alien plant *Bidens pilosa* [25](Gao et al. 2021). Therefore, if organisms from other

trophic levels frequently mediate the effects of nutrient fluctuations on alien plant

invasion, we recommend that studies testing the fluctuating resources hypothesis

should more frequently consider the interactive effect of other trophic levels.

Conclusions

The fluctuating resource hypothesis has become a key theory for explaining

invasion success of alien plants. However, our study is, to the best of our knowledge,

the first multi-species experiment that tested how another trophic level influences the

effects of nutrient fluctuations on alien plant invasion. Partly in line with the

fluctuating resource hypothesis, we found tentative evidence that nutrient variability

promotes alien plant invasion only under overall low-nutrient conditions, and only in

the absence of herbivores. Therefore, other trophic levels, such as herbivores in our

study, might mediate the interactive effect of nutrient enrichment on alien plant

invasion into resident communities.

Acknowledgements

We thank Xue Zhang, Mingxin Pan, Chunling Chang, Huifei Jin, Lichao Wang,

Lingxi Wang, Liping Shan, Han Yu for set-up of the experiment and plant harvest,

and we thank Mingxin Pan for weighing the biomass. This work was supported by

funding from the Chinese Academy of Sciences (Y9B7041001).

Author contributions

Y Liu conceived the idea and designed the experiment. Y Li and Y Gao

performed the experiment. Y Li and Y Liu analyzed the data. Y Li and Y Liu wrote

the first draft of the manuscript, with further inputs from Y Gao and M van Kleunen.

Data availability

All data and code available from the Dryad Digital Repository

https://doi.org/10.5061/dryad.fj6q573vn (Li et al., 2021).

References

1. Vilà M, Espinar JL, Hejda M, Hulme PE, Jarošík V, Maron JL, Jan P, Schaffner U,

Sun Y, Pyšek P. 2011 Ecological impacts of invasive alien plants: a meta-analysis of

their effects on species, communities and ecosystems. *Ecol. Lett.* **14**, 702-708. (doi:

10.1111/j.1461-0248.2011.01628.x)

2. Seebens H *et al.* 2017 No saturation in the accumulation of alien species

worldwide. *Nat. Commun.* **8**, 14435. (doi: 10.1038/ncomms14435)

3. Seebens H *et al.* 2020 Projecting the continental accumulation of alien species

through to 2050. *Global Change Biol.* (doi: 10.1111/gcb .15333)

4. Gippet JMW, Bertelsmeier C. 2021 Invasiveness is linked to greater commercial

success in the global pet trade. *P. Natl. Acad. Sci. USA* **118**, e2016337118. (doi:

10.1073/pnas.2016337118)

5. Catford JA, Jansson R, Nilsson C. 2009 Reducing redundancy in invasion ecology

by integrating hypotheses into a single theoretical framework. *Divers Distrib* **15**,

22-40. (doi: 10.1111/j.1472-4642.2008.00521.x)

6. Schultheis EH, Berardi AE, Lau JA. 2015 No release for the wicked: enemy release

is dynamic and not associated with invasiveness. *Ecology* **96**, 2446-2457. (doi:

10.1890/14-2158.1)

7. Liu YJ, van Kleunen M. 2019 Nitrogen acquisition of Central European herbaceous

plants that differ in their global naturalization success. *Funct. Ecol.* **33**, 566-575. (doi:

10.1111/1365-2435.13288)

8. Enders M *et al.* 2020 A conceptual map of invasion biology: Integrating hypotheses

into a consensus network. *Global Ecol. Biogeogr.* **29**, 978-991. (doi:

10.1111/geb.13082)

9. Speißer B, Liu YJ, van Kleunen M. 2021 Biomass responses of widely and less-

widely naturalized alien plants to artificial light at night. *J. Ecol.* **109**, 1819-1827.

(doi: 10.1111/1365-2745.13607)

10. Davis MA, Grime JP, Thompson K. 2000 Fluctuating resources in plant

communities: a general theory of invasibility. *J. Ecol.* **88**, 528-534. (doi:

10.1046/j.1365-2745.2000.00473.x)

11. Bradley BA, Blumenthal DM, Wilcove DS, Ziska LH. 2010 Predicting plant

invasions in an era of global change. *Trends Ecol. Evol.* **25**, 310-318. (doi:

10.1016/j.tree.2009.12.003)

12. Dostál P, Dawson W, van Kleunen M, Keser LH, Fischer M. 2013 Central

European plant species from more productive habitats are more invasive at a global

scale. *Global Ecol. Biogeogr.* **22**, 64-72. (doi: 10.1111/j.1466-8238.2011.00754.x)

13. Kalusová V *et al.* 2017 Naturalization of European plants on other continents: The

role of donor habitats. *P. Natl. Acad. Sci. USA,* 13756. (doi:

10.1073/pnas.1705487114)

14. Liu YJ, Oduor AMO, Zhang Z, Manea A, Tooth IM, Leishman MR, Xu XL, van

Kleunen M. 2017 Do invasive alien plants benefit more from global environmental

change than native plants? *Global Change Biol.* **23**, 3363-3370. (doi:

10.1111/gcb.13579)

15. Seabloom EW *et al.* 2015 Plant species' origin predicts dominance and response

to nutrient enrichment and herbivores in global grasslands. *Nat. Commun.* **6**, 1-8. (doi:

10.1038/ncomms8710)

16. Pyšek P, Jarošík V, Hulme PE, Pergl J, Hejda M, Schaffner U, Vilà M. 2012 A

global assessment of invasive plant impacts on resident species, communities and

ecosystems: the interaction of impact measures, invading species' traits and

environment. *Global Change Biol.* **18**, 1725-1737. (doi:

10.1111/j.1365-2486.2011.02636.x)

17. Manea A, Leishman MR. 2015 Competitive interactions between established

grasses and woody plant seedlings under elevated CO_2 levels are mediated by soil

water availability. *Oecologia* **177**, 499-506. (doi: 10.1007/s00442-014-3143-z)

18. Jia JJ, Dai ZC, Li F, Liu YJ. 2016 How will global environmental changes affect

the growth of alien plants? *Front. plant sci.* **7**, 1623. (doi: 10.3389/fpls.2016.01623)

19. Heckman RW, Halliday FW, Wilfahrt PA, Mitchell CE. 2017 Effects of native

diversity, soil nutrients, and natural enemies on exotic invasion in experimental plant

communities. *Ecology* **98**, 1409-1418. (doi: 10.1002/ecy.1796)

20. Liu YJ, van Kleunen M. 2017 Responses of common and rare aliens and natives

to nutrient availability and fluctuations. *J. Ecol.* **105**, 1111-1122. (doi:

10.1111/1365-2745.12733)

21. Sardans J, Bartrons M, Margalef O, Gargallo-Garriga A, Janssens IA, Ciais P,

Obersteiner M, Sigurdsson BD, Chen HYH, Peñuelas J. 2017 Plant invasion is

associated with higher plant-soil nutrient concentrations in nutrien-poor

environments. *Global Change Biol.* **23**, 1282-1291. (doi: 10.1111/gcb.13384)

22. Kelso MA, Wigginton RD, Grosholz ED. 2020 Nutrients mitigate the impacts of

extreme drought on plant invasions. *Ecology* **101**, e02980. (doi: 10.1002/ecy.2980)

23. Parepa M, Fischer M, Bossdorf O. 2013 Environmental variability promotes plant

invasion. *Nat. Commun.* **4**, 1-4. (doi: 10.1038/ncomms2632)

24. Liu YJ, Zhang XQ, van Kleunen M. 2018 Increases and fluctuations in nutrient

availability do not promote dominance of alien plants in synthetic communities of

common natives. *Funct. Ecol.* **32**, 2594-2604. (doi: 10.1111/1365-2435.13199)

25. Gao FL, He QS, Xie RQ, Hou JH, Shi CL, Li JM, Yu FH. 2021 Interactive effects

of nutrient availability, fluctuating supply, and plant parasitism on the post-invasion

success of *Bidens pilosa*. *Biol. Invasions*, 1-12. (doi: 10.1007/s10530-021-02555-y)

26. Keane RM, Crawley MJ. 2002 Exotic plant invasions and the enemy release

hypothesis. *Trends Ecol. Evol.* **17**, 164-170. (doi: 10.1016/S0169-5347(02)02499-0)

27. Blumenthal DM. 2006 Interactions between resource availability and enemy

release in plant invasion. *Ecol. Lett.* **9**, 887-895. (doi:

10.1111/j.1461-0248.2006.00934.x)

28. Zhang ZJ, Pan XY, Blumenthal D, van Kleunen M, Liu M, Li B. 2018 Contrasting

effects of specialist and generalist herbivores on resistance evolution in invasive

plants. *Ecology* **99**, 866-875. (doi: 10.1002/ecy.2155)

29. Pearson DE, Ortega YK, Villarreal D, Lekberg Y, Cock MC, Eren Ö, Hierro JL.

2018 The fluctuating resource hypothesis explains invasibility, but not exotic

advantage following disturbance. *Ecology* **99**, 1296-1305. (doi: 10.1002/ecy.2235)

30. Dawson W, Rohr RP, van Kleunen M, Fischer M. 2012 Alien plant species with a

wider global distribution are better able to capitalize on increased resource

availability. *New Phytol.* **194**, 859-867. (doi: 10.1111/j.1469-8137.2012.04104.x)

31. Meyer GA. 2000 Interactive effects of soil fertility and herbivory on *Brassica*

nigra. *Oikos* **88**, 433-441. (doi: 10.1034/j.1600-0706.2000.880221.x)

32. Hawkes CV, Sullivan JJ. 2001 The impact of herbivory on plants in different

resource conditions: a meta-analysis. *Ecology* **82**, 2045-2058. (doi: 10.2307/2680068)

33. Gianoli E, Salgado-Luarte C. 2017 Tolerance to herbivory and the resource

availability hypothesis. *Biol. Lett.* **13**, 20170120. (doi: 10.1098/rsbl.2017.0120)

34. Hu XT, Dong BC. 2019 Herbivory and nitrogen availability affect performance of

an invader *Alternanthera philoxeroides* and its native congener *A. sessilis*. *Flora* **257**,

151412. (doi: 10.1016/j.flora.2019.05.011)

35. Ma JS, Li HR. 2018 The checklist of alien invasive plants in china. Beijing,

China: Higher education press.

36. Roumet C, Urcelay C, Díaz S. 2006 Suites of root traits differ between annual and

perennial species growing in the field. *New Phytol.* **170**, 357-368. (doi:

10.1111/j.1469-8137.2006.01667.x)

37. Li YJ, Gao YZ, van Kleunen M, Liu YJ. 2021 Data from: Interactive effects of

herbivory and the level and fluctuations of nutrient availability on dominance of alien

plants in synthetic native communities, Dryad, Dataset. (doi:

10.5061/dryad.fj6q573vn)

38. Bürkner PC. 2017 An R package for bayesian multilevel models using Stan. *J.*

Statist. Software **80**. (doi: 10.18637/jss.v080.i01)

39. R Core Team. 2020. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

40. Zuur A, Ieno EN, Walker N, Saveliev AA, Smith GM. 2009 Mixed effects models

and extensions in ecology with R. New York: USA Springer.

41. Schad DJ, Hohenstein S, Vasishth S, Kliegl R. 2020 How to capitalize on a priori

contrasts in linear (mixed) models: A tutorial. *J. Mem. Lang.* **110**, 104038 (doi:

10.1016/j.jml.2019.104038)

42. Hurlbert SH. 1984 Pseudoreplication and the Design of Ecological Field

Experiments. *Ecol. Monogr.* **54**, 187-211. (doi: 10.2307/1942661)

43. Colegrave N, Ruxton GD. 2018 Using Biological Insight and Pragmatism When

Thinking about Pseudoreplication. *Trends Ecol. Evol.* **33**, 28-35. (doi:

10.1016/j.tree.2017.10.007)

44. Sher AA, Hyatt LA. 1999 The disturbed resource-flux invasion matrix: a new

framework for patterns of plant invasion. *Biol. Invasions* **1**, 107-114. (doi:

10.1023/A:1010050420466)

45. Dawson W, Fischer M, van Kleunen M. 2012 Common and rare plant species

respond differently to fertilisation and competition, whether they are alien or native.

Ecol. Lett. **15**. (doi: 10.1111/j.1461-0248.2012.01811.x)

46. Dener E, Kacelnik A, Shemesh H. 2016 Pea plants show risk sensitivity. *Curr.*

Biol. **26**, 1763-1767. (doi: 10.1016/j.cub.2016.05.008)

47. Zhao CY, Liu YY, Shi XP, Wang YJ. 2020 Effects of soil nutrient variability and

competitor identify on growth and co-existence among invasive alien and native

clonal plants. *Environ. Pollut.* **261**, 113894. (doi: 10.1016/j.envpol.2019.113894)

48. Alvarenga R, Auad AM, Moraes JC, da Silva SEB, Rodrigues BS. 2019 Tolerance

to nymphs and adults of *Mahanarva spectabilis* (Hemiptera: Cercopidae) by forage

plants in fertilized soils. *Pest Manage. Sci.* **75**, 2242-2250. (doi: 10.1002/ps.5361)

49. Garcia LC, Eubanks MD. 2019 Overcompensation for insect herbivory: a review

and meta-analysis of the evidence. *Ecology* **100**, e02585. (doi: 10.1002/ecy.2585)

50. Getman-Pickering ZL, Stack GM, Thaler JS. 2021 Fertilizer quantity and type

alter mycorrhizae-conferred growth and resistance to herbivores. *J. Appl. Ecol.* **58**,

931-940. (doi: 10.1111/1365-2664.13833)

51. Zhong ZW, Li XF, Sanders D, Liu YM, Wang L, Ortega YK, Pearson DE, Wang

DL. 2021 Soil engineering by ants facilitates plant compensation for large herbivore

removal of aboveground biomass. *Ecology*, e03312. (doi: 10.1002/ecy.3312)

52. Deng Q, Hui DF, Dennis S, Reddy KC. 2017 Responses of terrestrial ecosystem

phosphorus cycling to nitrogen addition: A meta-analysis. *Global Ecol. Biogeogr.* **26**,

713-728. (doi: 10.1111/geb.12576)

53. Sardans J, Grau O, Chen HYH, Janssens IA, Ciais P, Piao S, Peñuelas J. 2017

Changes in nutrient concentrations of leaves and roots in response to global change

factors. *Global Change Biol.* **23**, 3849-3856. (doi: 10.1111/gcb.13721)

54. Scalon MC, Haridasan M, Franco AC. 2017 Influence of long-term nutrient

manipulation on specific leaf area and leaf nutrient concentrations in savanna woody

species of contrasting leaf phenologies. *Plant Soil* **421**, 233-244. (doi:

10.1007/s11104-017-3437-0)

55. Oldroyd GED, Leyser O. 2020 A plant's diet, surviving in a variable nutrient

environment. *Science* **368**, eaba0196. (doi: 10.1126/science.aba0196)

56. Li FR, Dudley T, Chen BM, Chang XY, Liang LY, Peng SL. 2016 Responses of

tree and insect herbivores to elevated nitrogen inputs: A meta-analysis. *Acta Oecol.*

77, 160-167. (doi: 0.1111/geb.12576)

57. Jamieson MA, Burkle LA, Manson J, Runyon JB, Trowbridge AM, Zientek J.

2017 Global change effects on plant-insect interactions: The role of phytochemistry.

Curr. Opin. Insect Sci. **70**. (doi: 10.1016/j.cois.2017.07.009)

58. Kudo G. 2003 Variations in leaf traits and susceptibility to insect herbivory within

a *Salix miyabeana* population under field conditions. *Plant Ecol.* **169**, 61-69. (doi:

10.1023/A:1026209017627)

59. Borgström P, Strengbom J, Marini L, Viketoft M, Bommarco R. 2017 Above-and

belowground insect herbivory modifies the response of a grassland plant community

to nitrogen eutrophication. *Ecology* **98**, 545-554. (doi: 10.1002/ecy.1667)

60. Anderson TM *et al.* 2018 Herbivory and eutrophication mediate grassland plant

nutrient responses across a global climatic gradient. *Ecology* **99**, 821-833. (doi:

10.1002/ecy.2175)

61. Hernán G, Castejón I, Terrados J, Tomas F. 2019 Herbivory and resource

availability shift plant defense and herbivore feeding choice in a seagrass system.

Oecologia **189**, 719-732. (doi: 10.1007/s00442-019-04364-6)

62. Olff H, Ritchie ME. 1998 Effects of herbivores on grassland plant diversity.

Trends Ecol. Evol. **13**, 261-265. (doi: 10.1016/S0169-5347(98)01364-0)

63. Koerner SE, Smith MD, Burkepile DE, Hanan NP, Avolio ML, Collins SL,

Knapp AK, Lemoine NP, Forrestel EJ, Eby S. 2018 Change in dominance determines

herbivore effects on plant biodiversity. *Nature Ecol. Evol.* **2**, 1925-1932. (doi:

10.1038/s41559-018-0696-y)

64. Allington GRH, Koons DN, Morgan Ernest SK, Schutzenhofer MR, Valone TJ.

2013 Niche opportunities and invasion dynamics in a desert annual community. *Ecol.*

Lett. **16**, 158-166. (doi: 10.1111/ele.12023)

65. Sakavara A, Tsirtsis G, Roelke DL, Mancy R, Spatharis S. 2018 Lumpy species

coexistence arises robustly in fluctuating resource environments. *P. Natl. Acad. Sci.*

USA **115**, 738-743. (doi: 10.1073/pnas.1705944115)

Table 1 Output of the models testing effects of nutrient availability (low *vs* high), nutrient fluctuations (constant *vs* pulsed), herbivory

treatments (with *vs* without), and their interactions on aboveground biomass production of the alien target species, biomass production

of the native communities and biomass proportion of the alien target species in each pot. Shown are the model estimates and standard

errors as well as the lower (L) and upper (U) values of the 95% and 90% credible intervals (CI).

		Estimate	SE	L95% CI	U95% CI	L90% CI	U90% CI
Biomass production of alien target plants	Intercept	1.105	0.786	-0.506	2.634	-0.105	2.279
	Nutrient availability (NA)	0.416*	0.058	0.303	0.532	0.322	0.515
	Nutrient fluctuation (NF)	0.046	0.047	-0.047	0.139	-0.032	0.124
	Herbivory treatment (H)	-0.090†	0.049	-0.185	0.005	-0.170	-0.010
	NA×H	-0.113*	0.048	-0.208	-0.018	-0.190	-0.035
	NF×H	-0.009	0.048	-0.104	0.086	-0.087	0.071
	NA×NF	0.031	0.049	-0.066	0.127	-0.050	0.111
	NA×NF×H	0.020	0.049	-0.076	0.116	-0.060	0.100
Biomass production of native communities	Intercept	3.529*	1.154	1.114	5.759	1.609	5.321
	Nutrient availability (NA)	3.252*	0.110	3.033	3.467	3.069	3.434
	Nutrient fluctuation (NF)	0.023	0.111	-0.191	0.238	-0.158	0.204
	Herbivory treatment (H)	-1.150*	0.109	-1.361	-0.939	-1.328	-0.973
	NA×H	-0.735*	0.110	-0.951	-0.520	-0.913	-0.554
	NF×H	-0.144	0.109	-0.359	0.071	-0.324	0.035
	NA×NF	-0.044	0.109	-0.254	0.167	-0.224	0.133
	NA×NF×H	-0.086	0.109	-0.302	0.129	-0.267	0.095
Biomass proportion of alien target plants	Intercept	23.129*	11.495	0.112	46.144	4.487	41.768
	Nutrient availability (NA)	-1.890*	0.766	-3.387	-0.380	-3.127	-0.623
	Nutrient fluctuation (NF)	0.649	0.753	-0.820	2.125	-0.592	1.897
	Herbivory treatment (H)	4.229*	0.882	2.470	5.937	2.754	5.647
	NA×H	-1.064	0.766	-2.554	0.435	-2.321	0.182
	NF×H	-0.57	0.765	-2.079	0.921	-1.835	0.665
	NA×NF	0.918	0.770	-0.636	2.419	-0.343	2.188
	NA×NF×H	1.432†	0.786	-0.123	2.959	0.125	2.712

Parameters whose 95% credible intervals do not overlap with zero are indicted with asterisks (*), and whose 90% credible intervals do

not overlap with zero are indicated with daggers (†). Residual standard deviations *sigma* for individual alien species are found in

Table S2.

Figure legends

Figure 1 Graphical illustration of the experimental design. Overview of the herbivory treatment,

the herbivory-treatment cage during the experiment, and the pot design (a); the amount of

nutrient solution supplied each week during the ten weeks of the experiment (b). The constant

and pulsed nutrient supply under low- or high-nutrient availability received the same total

amount of nutrients during the ten weeks.

Figure 2 Mean values (±SE) of biomass production of alien target species (a) and native

communities (b) and biomass proportion of the alien target species (c) under each combination of

two nutrient availability (low *vs* high), two nutrient-fluctuation (constant *vs* pulsed) and two

herbivory (with *vs* without) treatments. Parameters whose 95% credible intervals do not overlap

with zero are indicted with asterisks (*), whose 90% credible intervals do not overlap with zero

are indicated with daggers (†), and whose 90% credible intervals do not overlap with zero are

indicated with "ns".

Figure 1 Graphical illustration of the experimental design. Overview of the herbivory treatment,

the herbivory-treatment cage during the experiment, and the pot design (a); the amount of

nutrient solution supplied each week during the ten weeks of the experiment (b). The constant

and pulsed nutrient supply under low- or high-nutrient availability received the same total

amount of nutrients during the ten weeks.

Figure 2 Mean values (±SE) of biomass production of alien target species (a) and native

communities (b) and biomass proportion of the alien target species (c) under each combination of

two nutrient availability (low *vs* high), two nutrient-fluctuation (constant *vs* pulsed) and two

herbivory (with *vs* without) treatments. Parameters whose 95% credible intervals do not overlap

with zero are indicted with asterisks (*), whose 90% credible intervals do not overlap with zero

are indicated with daggers (†), and whose 90% credible intervals do not overlap with zero are

indicated with "ns".

Supporting information

Methods S1

Recipe for 400% strength Hoagland's Complete Nutrient Solution

The preparation of Hoagland nutrient solution essentially followed recommendations by

Hoagland and Arnon (1950) and Liu et. al (2017), with the exception of the form in which iron

was added (see below). We prepared the stock solutions 1-6 and micronutrient-stock solution

given below, and used the amounts indicated to prepare 1 liter of nutrient solution:

(1) 4 mL of 1.00 M/L KH_2PO_4

(2) 20 mL of 1.00 M/L KNO_3

(3) 20 mL of 1.00 M/L $Ca(NO_3)_2$

(4) 8 mL of 1.00 M/L $MgSO_4$

(5) 4 ml of micronutrient stock solution (see recipe below)

(6) 10 ml of 1000 mg/liter iron from iron chelate (Fe-EDTA)

* Micronutrient-stock solution per liter:

2.86 g H_3BO_3

1.81 g $MnCl_2·4H_2O$

0.22 g $ZnSO_4·7H_2O$

0.08 g $CuSO_4·5H_2O$

0.018 g H_2MoO_4 (Assaying 94% MoO_3)

* Hoagland's recipe called for 1 ml of 0.5% iron tartrate stock per liter of nutrient solution but we

used iron chelate instead.

Reference: Hoagland, D.R. & Arnon, D.I. 1950. The water-culture method of growing plants without soil. California

Agricultural Experiment Station, Circular 347.

Table S1 Detailed information on the eight alien target species and 15 native competitor species used in this study

Species	Family	Status	Life history	Seed sources	Sowing date
Bidens frondosa L.	Compositae	Alien	Annual	Zhejiang	26.06.2020
Bidens pilosa L.	Compositae	Alien	Annual	Zhejiang	26.06.2020
Solidago canadensis L.	Compositae	Alien	Perennial	Zhejiang	26.06.2020
Xanthium strumarium L.	Compositae	Alien	Annual	Nei Mongol	26.06.2020
Abutilon theophrasti Medik.	Malvaceae	Alien	Annual	Jilin	26.06.2020
Hibiscus trionum L.	Malvaceae	Alien	Annual	Jilin	05.07.2020
Lolium perenne L.	Poaceae	Alien	Perennial	Greenwood flower seed company	26.06.2020
Paspalum notatum Flüggé	Poaceae	Alien	Perennial	Zhejiang	15.05.2020
Catolobus pendulus (L.) Al-Shehbaz[5]	Brassicaceae	Native	Biennial	Nei Mongol	26.06.2020
Cynanchum chinense R. Br.[2]	Apocynaceae	Native	Annual	Nei Mongol	26.06.2020
Lappula myosotis V. Wolf[1]	Boraginaceae	Native	Annual or Biennial	Nei Mongol	26.06.2020
Artemisia rubripes Nakai[4]	Compositae	Native	Perennial	Nei Mongol	26.06.2020
Achillea asiatica Serg.[2]	Compositae	Native	Perennial	Nei Mongol	26.06.2020
Convolvulus arvensis L.[3]	Convolvulaceae	Native	Perennial	Jilin	26.06.2020
Gypsophila licentiana Hand.-Mazz.[4]	Caryophyllaceae	Native	Perennial	Nei Mongol	26.06.2020
Euphorbia humifusa Willd.[5]	Euphorbiaceae	Native	Annual	Nei Mongol	26.06.2020
Chloris virgata Sw.[2]	Poaceae	Native	Annual	Nei Mongol	26.06.2020
Cleistogenes squarrosa (Trin.) Keng[5]	Poaceae	Native	Perennial	Jilin	26.06.2020
Digitaria sanguinalis (L.) Scop.[1]	Poaceae	Native	Annual	Jilin	26.06.2020
Echinochloa crus-galli (L.) P.Beauv.[4]	Poaceae	Native	Annual	Jilin	26.06.2020
Setaria pumila (Poir.) Roem. & Schult.[3]	Poaceae	Native	Annual	Jilin	27.06.2020
Portulaca oleracea L.[3]	Portulacaceae	Native	Annual	Nei Mongol	26.06.2020
Rumex crispus L.[1]	Polygonacea	Native	Perennial	Nei Mongol	26.06.2020

614 [1,2,3,4,5]Allocation of native species to the five native communities; each group included three different native species

615 **Table S2** Output of the models residual standard deviations sigma for individual alien species. Shown are the model estimates and

616 standard errors as well as the lower (L) and upper (U) values of the 95% and 90% credible intervals (CI).

		Estimate	SE	L95% CI	U95% CI	L90% CI	U90% CI
Biomass production of alien target plants	*Abutilon theophrasti* Medik.	-0.785*	0.138	-1.033	-0.499	-1.001	-0.547
	Bidens frondose L.	1.074*	0.127	0.839	1.341	0.874	1.290
	Bidens pilosa L.	0.327*	0.126	0.092	0.590	0.128	0.544
	Hibiscus trionum L.	-0.730*	0.175	-1.072	-0.388	-1.018	-0.443
	Lolium perenne L.	0.667*	0.138	0.413	0.953	0.452	0.907
	Paspalum notatum Flüggé	0.444*	0.123	0.217	0.700	0.249	0.659
	Solidago canadensis L.	-0.488*	0.125	-0.718	-0.232	-0.688	-0.277
	Xanthium strumarium L.	0.681*	0.127	0.445	0.944	0.480	0.900
Biomass production of native communities	*Abutilon theophrasti* Medik.	0.373*	0.133	0.135	0.652	0.169	0.602
	Bidens frondose L.	0.591*	0.129	0.354	0.862	0.388	0.810
	Bidens pilosa L.	0.511*	0.125	0.275	0.766	0.310	0.725
	Hibiscus trionum L.	0.603*	0.127	0.370	0.869	0.403	0.822
	Lolium perenne L.	0.484*	0.146	0.216	0.788	0.253	0.737
	Paspalum notatum Flüggé	0.960*	0.122	0.731	1.209	0.768	1.168
	Solidago canadensis L.	0.721*	0.113	0.510	0.949	0.539	0.911
	Xanthium strumarium L.	0.563*	0.123	0.335	0.816	0.369	0.773
Biomass proportion of alien target plants	*Abutilon theophrasti* Medik.	2.471*	0.134	2.219	2.749	2.258	2.700
	Bidens frondose L.	2.559*	0.133	2.314	2.833	2.353	2.783
	Bidens pilosa L.	2.206*	0.128	1.971	2.470	2.004	2.426
	Hibiscus trionum L.	2.624*	0.141	2.355	2.912	2.398	2.860
	Lolium perenne L.	2.962*	0.144	2.698	3.262	2.737	3.211
	Paspalum notatum Flüggé	2.908*	0.123	2.679	3.162	2.712	3.116
	Solidago canadensis L.	2.416*	0.125	2.176	2.669	2.218	2.626
	Xanthium strumarium L.	2.596*	0.127	2.355	2.858	2.394	2.811

Figure S1 Mean values (±SE) of biomass production of the alien target species (a) and the native

communities (b) and biomass proportion of the alien target species (c) under different nutrient

availability conditions.

Figure S2 Mean values (±SE) of biomass production of the alien target species (a) and the native

communities (b) and biomass proportion of the alien target species (c) growing with and without

herbivory.

Figure S3 Mean values (±SE) of biomass production of native communities (a) and alien target

species (b) under each combination of two nutrient availability (low *vs* high) and two herbivory

(with *vs* without) treatments.

Figure S4 Mean values (±SE) of total biomass per pot under each combination of two nutrient

availability (low *vs* high), two nutrient fluctuation (constant *vs* pulsed) and two herbivory (with

vs without) treatments (a), with and without herbivory (b), and under each combination of two

nutrient availability (low *vs* high) and two herbivory (with *vs* without) treatments (c). Parameters

whose 95% credible intervals do not overlap with zero are indicted with asterisks (*).