Modelling and assessing additional transmission routes for porcine reproductive and respiratory

syndrome virus: vehicle movements and feed ingredients

Jason A. Galvis [1], Cesar A. Corzo[2], Gustavo Machado[1]*

[1]Department of Population Health and Pathobiology, College of Veterinary Medicine, Raleigh, North

Carolina, USA.

[2]Veterinary Population Medicine Department, College of Veterinary Medicine, University of Minnesota,

St Paul, MN, USA.

*Corresponding author: gmachad@ncsu.edu

Summary

Accounting for multiple modes of livestock disease dissemination in epidemiological models remains a

challenge. We developed and calibrated a mathematical model for transmission of porcine reproductive and

respiratory syndrome virus (PRRSV), tailored to fit nine modes of between-farm transmission pathways

including: farm-to-farm proximity (local transmission), contact network of batches of pigs transferred

between farms (pig movements), re-break probabilities for farms with previous PRRSV outbreaks, with the

addition of four different contact networks of transportation vehicles (vehicles to transport pigs to farms,

pigs to markets, feed and crew) and the amount of animal by-products within feed ingredients (e.g. animal

fat or meat and bone meal). The model was calibrated on weekly PRRSV outbreaks data. We assessed the

role of each transmission pathway considering the dynamics of specific types of production (i.e., sow,

nursery). Although our results estimated that the networks formed by transportation vehicles were more

densely connected than the network of pigs transported between-farms, pig movements and farm proximity

were the main PRRSV transmission routes regardless of farm types. Among the four vehicle networks,

vehicles transporting pigs to farms explained a large proportion of infections, sow = 20.9%; nursery = 15%;

and finisher = 20.6%. The animal by-products showed a limited association with PRRSV outbreaks through

descriptive analysis, and our model results showed that the contribution of animal fat contributed only 2.5%

and meat and bone meal only 0.03% of the infected sow farms. Our work demonstrated the contribution of

multiple routes of PRRSV dissemination, which has not been deeply explored before. It also provides strong

evidence to support the need for cautious, measured PRRSV control strategies for transportation vehicles

and further research for feed by-products modeling. Finally, this study provides valuable information and

opportunities for the swine industry to focus effort on the most relevant modes of PRRSV between-farm

transmission.

Keywords: PRRSV transmission, swine disease dynamics, truck, animal by-product, swine disease

transmission, contact networks.

1. Introduction

Porcine reproductive and respiratory syndrome virus (PRRSV) remains a major economic burden in North America (Holtkamp et al., 2013) as it continues to spread across multiple pig-producing companies (Sanhueza et al., 2019; Jara et al., 2020; Galvis et al., 2021). A recent study developed a mathematical model to reconstruct the between-farm PRRSV dynamics to reveal the role of between-farm pig movements, farm-to-farm proximity, and the continued circulation of PRRSV within infected sites (named as re-break) on PRRSV transmission (Galvis et al., 2021). Despite the promising results, the study did not fully consider indirect contacts through between-farm transportation vehicles (e.g. vehicles transporting pigs, feed or farm personnel) contact networks, which has been previously described as one of the major modes of between-farm transmission of diseases in swine (Büttner and Krieter, 2020; Porphyre et al., 2020; Niederwerder, 2021), such as PRRS (Dee et al., 2002, 2004; Pitkin et al., 2009; Thakur, Sanchez et al., 2015) and African swine fever (ASF) (Gao et al., 2021; Gebhardt et al., 2021).

Detailed data regarding transportation vehicle movement and routes, coming in and out pig premises, can be difficult to obtain, which indeed could help explaining the lack of models considering this transmission pathway (Bernini et al., 2019; EFSA Panel on Animal Health and Welfare (AHAW) et al., 2021). Previous studies which have approached indirect transmission by transportation vehicles, have either used simulated probabilities to define indirect contact between farms such as the studies by Thakur et al., 2015 and Wiltshire, 2018, or observed truck movements such as the study by Buttner, 2020 using between farms movements in Germany. The potential of viral stability on vehicle surfaces has been demonstrated to be dependent on environmental conditions such as temperature, pH, moisture, and vehicle disinfection procedures (Dee et al., 2002, 2003; Jacobs et al., 2010). For example, Dee et al., isolated PRRSV under field conditions from surfaces such as concrete, floor mats and fomites between 2h and 4h after surface contamination (Dee et al., 2002). PRRSV has been shown to be more stable at low temperatures (-20° C to -70° C), surviving for long periods of time (>4 months) (Benfield et al., 1992), and becoming unstable as the temperature increases (Jacobs et al., 2010). In addition, dry conditions, low pH ranging between 5 and 7 (Benfield et al., 1992), iodine, quaternary ammonium or chlorine compounds used in vehicle disinfection

were successful in inactivating PRRSV (Shirai et al., 2000). Thus, if environmental conditions favor

PRRSV survivability and the vehicles are poorly cleaned and disinfected, the potential for pathogens to

disseminate within highly connected networks could indeed play a major role in disease spread (Büttner

and Krieter, 2020; Gebhardt et al., 2021).

In addition to transportation movements, contaminated feed could represent a possible route for

pathogen transmission (Gebhardt et al., 2021; Niederwerder, 2021), but the probability of PRRSV

transmission through feed has been described as relatively low (Cochrane et al., 2017; Ochoa et al., 2018;

Blázquez et al., 2020). However, a recent report by Dee et al. 2020, demonstrated under experimental

conditions that pigs which consumed pellet feed contaminated with 1×10^5 TCID50/ml of PRRSV became

infected. As such, cross-contamination of pellet feed may occur when coming into direct contact with

contaminated fomites or feed mill workers after the pelleting process (Niederwerder, 2021). In addition, it

is important to acknowledge that inadequate temperature applied during the pelleting process could reduce

the probability to inactivate PRRSV from contaminated feed ingredients, as observed in an experimental

study with porcine epidemic diarrhea virus (Cochrane et al., 2017). In North America, most feed

formulations include some animal by-products, such as animal fat, dried plasma, or bone meal in order to

increase growth performance (Lewis and Southern, 2001). Without adequate processes to inactivate

PRRSV, these ingredients could potentially be a source of contamination and later infection. Magar and

Larochellle in 2004 surveyed two Canadian slaughterhouses and found 4.3% of animal serum samples and

1.2% of the meat samples were positive for PRRSV by polymerase chain reaction. The same study also

demonstrated that the consumption of contaminated animal by-products caused infection in pigs (Magar

and Larochelle, 2004).

In this study, we built a novel mathematical model of PRRSV transmission tailored to nine modes

of between-farm propagation: local transmission by the farm-to-farm proximity, between-farm animal and

re-break for farms with previous PRRSV outbreaks (Galvis et al., 2021), with the addition of between-farm

vehicle movements (feed, shipment of live pigs between farms and to slaughterhouses, and farm personnel

[crews]), and the quantity of animal by-products which was restricted to animal fat, meat and bone meal in

pig feed ingredients. The model was used to estimate the weekly number of new PRRSV outbreaks and

their spatial distribution, which were compared to available data, and to quantify the contribution of each

transmission route.

Material and Methods

Databases

In this study, we used weekly PRRSV records captured by the Morrison Swine Health Monitoring Program

(MSHMP) (MSHMP, 2020). Data included outbreaks between January 22, 2009 and December 31, 2020,

from 2,294 farms from three non-commercially related pig production companies (coded as A, B, and C)

in a U.S. region (not disclosed due to confidentiality). Additional details about the study population and

PRRSV classification in the U.S. are available in (Galvis et al., 2021, Holtkamp et al., 2011). Individually,

each PRRSV record was classified as a new or recurrent outbreak according to the time between consecutive

outbreaks per farm (Galvis et al., 2021). A list of pig farms was available from the MSHMP database

(MSHMP, 2020), which included individual national premises identification number, farm type (sow

[which included farrow, farrow-to-wean, and farrow-to-feeder farms], nursery, finisher [which included

wean-to-feeder, wean-to-finish, feeder-to-finish], gilt development unit [which could be either part in

finisher or sow farms depending upon farm type used by pig production company], isolation and boar stud),

pig spaces per farm, and geographic coordinates. Between-farm pig movement data from January 01, 2020,

to December 31, 2020, were used to reconstruct directed weekly contact networks. Each movement batch

included movement date, farm of origin and destination, number of pigs transported, and purpose of

movement (e.g., weaning). Movement data missing either the number of animals transported, farm type,

farm of origin, or destination were excluded prior to analysis (731 [1.2%] unique movements were

excluded). In addition, four networks formed by transportation vehicles were recorded from the global

positioning system (GPS) vehicle tracker for which included tracking of all farms from company A (76%

of all farms within the study region for 2020) from January 01, 2020, to December 31, 2020. These

movements comprise near-real time GPS records of each transporting vehicle, which include geographic

coordinates for every 5 seconds, of any vehicle. Overall, 398 vehicles were monitored which included: (i)

159 trucks used to deliver feed to farms, (ii) 118 trucks utilized in the transportation of live pigs between

farms, (iii) 89 truck used in the transportation of pigs to markets (slaughterhouse), and (iv) 32 vehicles used

in the transportation of crew members, which by the information we collected correspond to the movement

of additional personnel needed for vaccination, pig loading and unloading among other activities which

included power washing (Figure 1). Each movement batch included a unique identification number, speed,

date and time along with coordinates of each vehicle location recorded every 5 seconds. A vehicle visit was

defined as a vehicle coordinate (latitude and longitude) and speed of zero km/h for at least 5 minutes within

a radius of 1.5 km of any farm or cleaning station (time and distance radius selected after discussion with

personnel in charge of vehicle logistics and data observation). In case more than one farm was located

within a 1.5 km radius, we assumed all these farms were at-risk of transmission, and therefore the vehicle

contacted all of them. It is worth noting that in some cases where farms are located at distances <1.5 km

from other farms it is because they belong to the same pig production (personal communication). We

calculated the time in minutes the vehicle remained within each farm's perimeter and the vehicle contact

networks between the farms was built considering the elapsed time a vehicle visited two or more different

farms (Figure 1). To accommodate PRRSV survivability in the environment, we considered two seasons

(cold and warm weather) based on previous literature (Dee et al., 2002; Jacobs et al., 2010). Under

laboratory conditions, it was reported that PRRSV preserved stability for more than 72h when temperatures

oscillated between 4° C and 10° C (cold temperatures) and less than 24h when temperatures were equal or

higher than 20° C (warm temperatures) (Jacobs et al., 2010). Thus, an edge among two different farms was

recorded if the elapsed time the vehicle visited both farms was less than 72 hours or 24 hours, for the cold

and warm seasons, respectively. However, we did not consider the formation of edges between consecutive

farms after vehicles were observed via GPS driving through clean stations (Figure 1). The edges for all four

vehicle networks were weighted by the elapsed time each vehicle visited two different farms, which was

later transformed to a probability assuming a decreasing linear relationship of PRRSV stability in the

environment (Figure 1 and Supplementary Material Figure S1). Additionally, we collected feed load out

records from all (three) feed mills of company A for 2020, with each feed record including feed mill

identification with individualized feed formulation (ingredients), amount of feed delivered, destination farm

identification, and destination farm delivery data. From the feed records, we collected the amount in pounds

(lb) of animal by-products (parts of a slaughtered animal that included animal fat, pig plasma and meat and

bone meal) of each feed formulation received by the farms for each week of 2020 (Supplementary Material

Figure S2 and S3). Although company B and C data about vehicle movements and feed delivery was not

available, we kept the farms from both companies in the transmission model to complement the PRRSV

dissemination by the local transmission (Jara et al., 2020).

Descriptive analysis

Between farm animal and transportation vehicles movement

The networks formed by movement of live pigs transported between farms and four types of transportation

vehicles visiting farms were reconstructed and analyzed. A set of network metrics including: size, general

properties, and heterogeneity at node-level were evaluated for each directed static and temporal network

(Supplementary Material Table S1 for terminology and network metric description). To determine if the

static networks of pig and vehicle movements could represent the temporal variation of the between farm

contacts over a year, we calculated the causal fidelity (Lentz et al., 2016). Briefly, causal fidelity quantifies

the error of the static representation of temporal networks through a ratio among the number of paths

between both representations. Thus, a casual fidelity of 100% would mean that a temporal network is well

represented by its static counterpart, and conversely, a value close to 0% means the network should not be

considered as a static system (Lentz et al., 2016). We then estimated whether farms with PRRSV outbreaks

records were more frequently connected with other infected farms through the ingoing and outgoing contact

chain, compared with farms without PRRSV records. Finally, we estimated if the time transportation

vehicles remained within farms premises was higher in the farms with PRRSV outbreaks compared with

farms without PRRSV breaks. The association for the contact chain and the time the vehicles stayed in the

farms with PRRSV outbreaks were evaluated through a Mann-Whitney test.

Animal by-products in feed ingredients

We calculated the total amounts of animal fat, pig plasma, animal protein blend 58%, protein blend (animal

protein blend and protein blend was made of a combination of ingredients such as meat meal, corn germ

meal, hominy and dried distillers grains with solubles), and meat and bone meal present in each of the 23

feed formulations delivered to farms with and without PRRSV outbreaks in 2020 (Supplementary Material

Figure S2 and S3). In order to further evaluate the association between PRRSV outbreaks and the delivery

of feed with animal by-products, we performed a logistic regression analysis for each farm type and

ingredient in which the response variable was positive or negative for PRRSV from January 1^{st} , 2020 to

December 31^{st} , 2020, and the predictor was the amount of animal by-product divided by the farm's pig

capacity to avoid confusion by the farm size.

Transmission model

The analysis of spatiotemporal distribution of farm-level PRRSV outbreaks was based on our previously

developed stochastic model (Galvis et al., 2021), which here was extended to include vehicle transportation

networks and the delivery of animal by-products. The model was calibrated on the weekly PRRSV

outbreaks and considering nine transmission modes including: (1) contact network of discrete pig

movements; (2) the local transmission events between neighboring farms driven by distances among farms;

(3) re-break by a previous exposure to PRRSV; indirect contact by vehicles coming into farms, including

for (4) feed, animal delivery to (5) farms and (6) market, and (7) vehicles used by personnel (crew) involved

in the loading and unloading of pigs; amount of (8) animal fat and (9) meat and bone meal in feed

formulation delivered to farms (Figure 2). The model simulates between farm transmission among three

farm-level infectious states, Susceptible-Infected-Outbreak (SIO), and we defined Susceptible status as

farms free of PRRSV, Infected status as farms with PRRSV but not yet detected, and Outbreaks status as

infected farm that detected PRRSV. Thus, farms in a susceptible state (i) receive the force of infection of

infected and outbreak farms (j) in each time step t and become infected at probability Y_{it} (Figure 2). It is

worth noting that the latent period of PRRSV is not explicitly modeled, as it is typically a few days after

infection, and often viral shedding starts within 7 days post-infection (Pileri and Mateu, 2016; Chase-

Topping et al., 2020), thus it is embedded in the weekly timestep. Local transmission was modelled through

a gravity model where the probability of infection is proportional to the animal capacity of the farms and

inversely related to the distance between the two farms (i.e. lower transmission at longer distances), with

the maximum distance set at 35 km, similar to our previous study to facilitate the comparison of results

(Galvis et al., 2021). Local transmission is also dependent on the enhanced vegetation index (EVI) around

the farm i (Jara et al., 2020; Galvis et al., 2021), such that the probability of transmission decreases with

high EVI values (Supplementary Material Figure S4). The transmission associated with between farm pig

movements is modeled by the number of all infected and outbreak farms sending pigs to susceptible farms.

The dissemination via transportation vehicle networks (e.g. vehicles transporting pigs to farms) is modeled

by the edge weight (E) and the time the vehicles remained on the susceptible farm premise (Z_{it}) (Figure 1

and Supplementary Material Figure S1 and S5). The transmission via animal fat and meat and bone meal

was only considered to sow farms and modulated simply by the amounts delivered to susceptible farms

(A_{it}). Pig plasma, animal protein blend meal 58% and protein blend were only delivered to nursery and

finisher farms, thus were not considered. For the re-break rate which is only considered for sow farms, we

assumed that subsequent new infections at an individual farm, within a time period of two years, were

associated with the same strain as the previous outbreak, and the probability was based on a survival

analysis evaluating the time farms re-break after recovering (W_{it}) (Holtkamp et al., 2010) (Supplementary

Material Figure S6). Then, for each transmission route, the force of infection (λ) of infected and outbreak

farms varies with a seasonality derived from analysis of the PRRSV records from 2015 to 2019

(Supplementary Material Figure S7). In addition, sow farms without a record of PRRSV outbreaks since

2009 were assumed to have high biosecurity levels (H) that reduce the force of infection received by

infected and outbreak farms, being H higher than zero and calibrated according to the observed outbreaks

(Supplementary Material Table S2). Otherwise, farms with outbreaks records were assumed to have low

biosecurity levels and H was defined as zero.

The transition from infected to an outbreak farm is estimated through a detection rate $f(x)$ (Figure

2). Thus, the probability that farms transit to outbreak state is assumed to be dependent on the maximum

detection probability (L), considered equal to cases reported to MSHMP (MSHMP, 2020), and the average

time it takes a farm to detect the disease (*x0*), assumed to be 4 weeks (estimated from information provided

by local swine veterinarians and previous literature) (Neira et al., 2017). The proportion of Infected and

Outbreak sow farms that return to a susceptible state is drawn from a Poisson distribution with mean of 41

weeks, which is the average time to stability described elsewhere (Sanhueza et al., 2019). Nursery and

finisher farms' transition to susceptible status are driven by pig production movement scheduling of the all-

in all-out management schemes of closeouts or by incoming or outgoing movements, whichever came first

(Galvis et al., 2021). Briefly, nurseries and finisher farms become susceptible within 7 and 25 weeks of pig

placement, respectively, or when at least one new pig movement is recorded before the farm reaches the

scheduled production phase timeline described earlier. A detailed description of the model can be found in

Figure 2, and previous work describes in greater detail other model parameters (Galvis et al., 2021). Finally,

we used an Approximate Bayesian Computation (ABC) rejection algorithm (Minter and Retkute, 2019), to

estimate the posterior distribution of unknown model parameters (list of model parameters available in

Supplementary Material Table S3) by selecting the 100 particles (number of particles accepted defined

according to our computational resources) that best fitted the temporal and spatial distribution of observed

PRRSV outbreaks (Supplementary Material Figures S8-S10).

Model outputs

The model outputs included (a) the force of infection for each farm type and transmission route, (b) the

weekly number of infected undetected and detected farms (outbreaks) and (c) the sensitivity performance

to detect PRRSV outbreak locations (Supplementary Material Section 2). We carried out 1,000 simulations,

each simulation choosing at random one of the 100 accepted particles, to estimate the relative contribution

of each transmission route and the weekly number of cases. For the sensitivity performance, 100 simulations

were used for each particle to estimate the risk of infection, and only 100 particles were used due to

computational resources (additional information see Supplementary Information Figure S10). For the

contribution of the routes, we evaluated the number of infected farms resulting from each transmission

route individually, which were then divided by the number of simulated infected farms from all the

combined routes and commercial companies. In addition, the average contribution for each route was

estimated by summing the weekly contributions divided by the number of simulated weeks, and credible

intervals (CI) using an equal-tailed interval method were estimated from the weekly contribution

distribution. The model was developed in the R (3.6.0 R Core Team, Vienna, Austria) environment, and all

simulations were run in RStudio Pro (1.2.5033, RStudio Team, Boston, MA) and transmission model

framework is available at https://github.com/machado-lab/pigspread.

Results

The comparison among the five networks showed that vehicles transporting feed, pigs to farms, pigs to

markets and movement of crews were significantly more connected than the pig movement network (Table

1). The network density formed by vehicles transporting feed exhibited the highest density (edge density =

0.2). Comparing the paths between the static and temporal networks of pig movement and transportation

vehicles movements, we found that causal fidelity was above 32% for all networks (Table 1), which means

that a significant number of the causal paths described by the static networks can be found in the temporal

networks. The networks of vehicles transporting feed and pigs to farms exhibited the highest causal fidelity

values (causal fidelity >90%), which means these networks are the closest representation of causal paths

between the static and temporal network. Analyzing the network components of each vehicle network for

company A, we found that the Largest Strongly Connected Component (LSCC) had between 976 and 1,591

farms, thus these vehicle movement networks connected between 55% and 90% of company A farms (Table

1). For the pig movement network, company B showed the highest number of farms in the LSCC with 61

farms, which represented 27% of the farms from that company, while company A and C showed a low level

of connectivity with the LSCC representing less than 1% of the farms. Comparing pig movement networks

also showed that company B had the highest edge density of 0.013, followed by company C (edge density

of = 0.005) and company A the lowest (edge density of = 0.002).

The vehicles transporting feed static network had a median in-degree and out-degree of 319 and

304, respectively, which was the highest in comparison with vehicles transporting pigs to farms, pigs to

markets and vehicles used in the transportation of crew (Table 1, Supplementary Material Figure S11 and

S12). The networks of vehicles transporting pigs to farms and crew had a median in-degree and out-degree

ranging between 14 and 21, while vehicles transporting pigs to markets and pig movements for all three

companies showed median in-degree and out-degree less than 7 (Table 1). The network of vehicles

transporting pigs to farms showed the highest median betweenness centrality, followed by vehicles

transporting feed, vehicles transporting pigs to markets and then crew networks, whilst pig movements had

the lowest betweenness (Supplementary Material Figure S13). Interestingly, despite the network of vehicles

transporting feed being more densely connected, the network of vehicles transporting pigs to farms showed

the highest median betweenness centrality values, indicating that this network has the highest number of

shortest paths by farm to connect other farms (Table 1 and Supplementary Material Figure S13).

Considering the result from the temporal networks, vehicles transporting feed showed the highest median

ingoing (ICC) and outgoing contact chain (OCC) (Supplementary Material Table S1), which indicates that

feed routes create the largest sequential paths over time that allow for more connections between farms than

any transportation or pig movements network (Table 1 and Supplementary Material Figure S14 and S15).

The ICC and OCC from vehicles transporting crew showed the second highest values, followed by vehicles

transporting pigs to farms and then vehicles transporting pigs to markets. When considering pig movement,

company A had the highest median ICC among the different companies (ICC = 34), and a median OCC of

zero. Companies B and C showed lower ICC values, but a higher median OCC of 15 and 4, respectively

(Table 1 and Supplementary Material Figure S14 and S15).

Furthermore, we evaluated the association of PRRSV outbreaks frequency within both ICC and

OCC from infected and non-infected farms. The results showed that PRRSV infected farms were more

frequently found within the ICC and OCC of infected farms for the networks of vehicles transporting feed,

pigs to farms and pigs to markets networks ($p < 0.05$), while such association was not observed for vehicles

transporting crew, and for pig movement networks it was only significant for OCC (Supplementary Material

Figures S16 and S17). We also evaluated the association between the time transportation vehicles remained

within infected and non-infected farms. The vehicles transporting feed and crew members spent more time

within infected nursery farms when compared with the non-infected farms ($p < 0.05$), while no association

was found with PRRSV at sow and finisher farms for these vehicles ($p > 0.05$) (Supplementary Material

Figures S18). In addition, no differences were found among the time spent on infected farms compared

with the non-infected farms for the vehicles transporting pigs to farms and pigs to markets for any farm

type ($p > 0.05$). Finally, the amount of animal by-products in feed (animal fat, pig plasma, protein blend

and meat and bone meal) were not significantly associated with PRRSV outbreaks ($p > 0.05$)

(Supplementary Material Figures S19-S21).

From the weekly model simulations from January 1th until December 31th, 2020, we estimated the

total number of infected farms at 1,790 (95% CI: 1,776–1,804), 113 (95% CI: 112–113) of which

corresponded to infected sow farms, 715 (95% CI: 704–726) to nursery farms and 960 (95% CI: 952–967)

to finisher farms (isolation and board stud farm were excluded from the results because no outbreaks were

reported in the studied period). It is worth noticing that, just as in the data, in our simulations the same farm

could have been at infected state more than once over the simulated year. Overall, results showed a good

agreement between the weekly observed number of PRRSV outbreaks and simulated outbreaks

(Supplementary Material Figure S9). The model inferred that, at the end of the 52 weeks, on average 158

(8.8%) of all PRRSV infected farms would be detected, in which 90% of the infected sow farms were

detected, while a much lower proportion of detection was estimated for nurseries (4.8%) and finishers farms

(2%). The model's predictive performance to correctly identify the weekly spatial distribution of known

PRRSV outbreaks showed an area under the ROC curve of 0.7 (More details are available in Supplementary

Material Section 2).

Evaluating the contribution of nine transmission routes over the simulated PRRSV spread for

company A's farms demonstrated that for sow farms the most important route was the local transmission

contributing to an average of 32.4% (95% CI 15%-67%) of the farm infections, followed by pig movements

with 28.3% (95% CI 1.9%-68%), vehicles transporting pig to farms with 20.9% (95% CI 5%-45%), vehicles

transporting feed 12% (95% CI 0,5%-32%), re-break 3.2% (95% CI 0%-6%), amount of animal fat within

feed formulation 2.5% (95% CI 0,7%-6%), vehicles transporting pigs to markets 0.4% (95% CI 0%-2.5%),

vehicles transporting crew 0.2% (95% CI 0%-1.5%) and amount of meat and bone meal within feed

formulation 0.03% (95% CI 0%-0,5%) (Figure 3). For nursery farms, pig movements were the most

important route contributing to 76.4% (95% CI 57%-89%) of the farm infections, followed by vehicles

transporting pigs to farms 15% (95% CI 5%-26%), local transmission with 5.8% (95% CI 3%-16%),

vehicles transporting feed 2.3% (95% CI 0.3%-6%), vehicles transporting pigs to markets 0.44% (95% CI

0%-1.4%) and vehicles transporting crew 0.1% (95% CI 0%-0,5%). For finisher farms, local transmission

was also the most important route contributing to 35.5% (95% CI 17%-58%) of the farm infections,

followed by pig movements with 30.1% (95% CI 12%-52%), vehicles transporting pigs to farms 20.6%

(95% CI 7%-38%), vehicles transporting feed 9.2% (95% CI 3%-19%), vehicles transporting pigs to

markets 3.8% (95% CI 0.13%-11%) and vehicles transporting crew 0.61% (95% CI 0.01%-1.8%). As

transportation vehicle data were not available for companies B and C, the results were restricted to three

transmission pathways (pig movements, local transmission and re-break) and are available in

Supplementary Material section 4, Figure S22.

Discussion

In this study, we demonstrated the contribution of nine pathways in PRRSV dissemination dynamics which

included pig movements network, farm-to-farm proximity, different types of transportation vehicle

networks (vehicles transporting feed, pigs to farms, pigs to markets, and crew), the delivery of animal by-

products, in particular animal fat and meat and bone meal in the feed meal and re-break. We demonstrated

that the transportation vehicle networks connected more farms than the pig movement networks, therefore

between farm contacts by transportation vehicles have the potential to propagate PRRSV to more sites and

more quickly than moving live pigs between-farms. Thus, we remark that the greater potential for PRRSV

transmission via transportation vehicle networks pose a great challenge to surveillance and effective control

of endemic disease in North America and future eradication of possible introduction of foreign animal

diseases such as ASF (Brown et al., 2020; Gao et al., 2021). Our regression results also suggest that the

overall volume of animal by-product delivery to the farms was not associated with the PRRSV outbreaks,

and our mathematical model results indicated that animal fat and meat and bone meal delivered to sow

farms combined contributed to <2,6% farm infection in the simulations.

Our study addresses major gaps in the understanding of how PRRSV propagates between-farms by

modelling multiple modes of transmission, which expand the understanding of PRRSV propagation, thus

far restricted to studies that have only considered the dissemination of PRRSV throughout animal

movement networks (Thakur, Revie et al., 2015; Lee et al., 2017; Makau et al., 2021). Our previous study

(Galvis et al., 2021) and the information about the transmission dynamics of PRRSV (Dee et al., 2002;

Perez et al., 2015; Thakur, Revie et al., 2015; Pileri and Mateu, 2016; Silva et al., 2019; Jara et al., 2020)

highlighted limitations of model accuracy and paucity evaluation of PRRSV dissemination by not

considering well known modes of transmission, such as indirect contacts formed by transportation vehicles.

In this study, we have further extended the previous transmission model (Galvis et al., 2021), by including

six additional transmission routes, including the contact networks formed by transportation vehicles and

animal by-products delivered to sow farms via feed ingredients. Our results demonstrate that pig

movements and local transmission were the main transmission routes, regardless of farm types (sow,

nursery, and finisher) (Figure 3). However, the contribution of transportation vehicles used to transfer pigs

to farms explained a significant number of infected farms as follows, 20.9% of sow farms, 15% of nurseries

and 20.6% of finisher farms. Our examination of feed delivery to farms, more specifically the volume of

animal by-products animal fat and meat and bone meal, indicated that this mode did not contribute

significantly to PRRSV transmission, contributing to only 2.5% and 0.03% of the infection in sow farms,

respectively. Although our results tend to prove a small role of feed, the approach used in this study is

limited by several factors that affect the contribution of feed in the PRRSV dissemination, therefore these

results must be taken with caution (more details in a section below). Finally, for companies B and C, the

dominant routes of transmission were similar to the results for company A, with pig movements having the

greatest contribution to infection at nursery farms, while local transmission mainly affected finishers

(Figure 3). However, re-break contributed to a high number of infections that was not observed in our

previous study (Galvis et al., 2021). Overall, our results have also reinforced the findings from previous

studies which highlight the role of vehicles in the transmission of infectious diseases that pose a significant

threat to or have a particular impact on the swine industry (Dee et al., 2003, 2004, 2007; Melmer et al.,

2020) and the potential risk of contaminated animal by-products in the feed meals to contribute to between-

farm transmission of PRRSV (Dee et al., 2020; Niederwerder, 2021).

From individual vehicle GPS movement data, we reconstructed networks while considering the

elapsed time between farm visits, and the time vehicles spent within each farm to define effective contacts

among them. In addition, we considered contact between farms to end once a truck drove through a cleaning

station (Figure 1), as previous studies suggest that vehicle disinfection can reduce the probability of

pathogens' introduction into pig farms, e.g., PRRSV (Thakur et al., 2017) and ASFV (Yoo et al., 2021). In

the current work, the inclusion of cleaning stations reduced around half the edges of the four types of

transportation vehicle networks, which represent valuable information for the analysis of disease

transmission given that including or not cleaning stations provide different results of the pathogen

dissemination among the farms through vehicles. It is worth to notice that we assumed that the cleaning

process was always effective to inactivate PRRSV (Shirai et al., 2000), even though the literature about

probability of PRRSV survival on vehicle surfaces after cleaning and disinfection remains limited (Dee et

al., 2004). Thus, further studies evaluating the presence and infectivity of PRRSV after vehicle cleaning

and disinfection are necessary.

Previous studies attempting to model transportation vehicle networks were limited to either static

or simulated between-farm contact networks, thus limiting our ability to further compare our networks

results (Thakur, Sanchez et al., 2015; Wiltshire, 2018; Sterchi et al., 2019; Porphyre et al., 2020; Yang et

al., 2020). However, our result demonstrated that for the four vehicle networks described in this study, the

static networks provided a close representation of the actual temporal networks, in which the ratio of the

number of path between the static and temporal network varied from 58% for vehicles transporting pigs to

markets to 98.7% for vehicles delivering feed (Table 1). Because of the high fidelity of the vehicle networks,

our results demonstrate that the static networks may be sufficient to explain causal paths among farms

formed by vehicle movements (Lentz et al., 2016). Although these results were only evaluated by

comparing the static and temporal network from a single commercial company and with one year of GPS

data, our results strongly suggest that static views of transportation vehicles networks can be used either to

describe such networks or to be used in disease transmission models when researchers only have access to

a static view of a network.

The transportation vehicle networks were more densely connected than the networks of between-

farm pig movements, ranging from 3 to 100 times more connected considering the network of vehicles

transporting pigs to markets and feed, respectively. Other studies have also found that vehicle movements

between-farms used for animal hauling increased the indirect contacts among farms by more than 50%

(Porphyre et al., 2020), consequently the network weak connected component became 50% larger (Sterchi

et al., 2019). Among the transportation vehicle networks analyzed here, the network of feed deliveries was

the most connected; for instance more than 85% of all farms of company A were connected through

sequential paths in the temporal network (ICC and OCC Table 1). On the contrary, the network of trucks

transporting live pigs between-farms was less connected, but in our transmission model it represented

between 15% and 20% of the total transmissions. In turn, the probability of transportation vehicles being

contaminated with PRRSV is more likely in vehicles transporting animals than feed, mainly because of the

direct contact with infected animals. Vehicles transporting pigs to markets could in the same way play a

similar risk of contamination as vehicles transporting pigs to farms, however the former vehicles tend to

visit mostly finisher farms with pigs ready for slaughter, thus less likely to re-introduce PRRSV back into

the transmission chain by visiting breeding farms (Passafaro et al., 2020). It is worth noting that this is not

the case for farms with complete production cycles, such as farrow to finisher farms, in which movements

returning from slaughterhouses are likely to pose a great risk (Henry et al., 2018). Our study was also the

first to collect and consider in a transmission model the contact networks of vehicles transporting crew

between-farms. It is known that the number of farm visitors increase the probability of introduction of

infectious pathogens (e.g. PRRSV), thus the movement of additional personnel often involved in loading

and unloading pigs have been previously associated with PRRSV dissemination and outbreaks (Dee et al.,

2002; Pitkin et al., 2009; Rossi et al., 2017). It is worth noting that in our study we modeled the consecutive

contacts between farms formed by the vehicles used by crew as a route of PRRSV dissemination, by

evaluating individual vehicles connecting farms that were visited consecutively rather than the group of

crew members, which is known to vary between each farm visit. Therefore, the risk of infection by vehicles

transporting crew could be more related to the vehicle itself as fomite, which in turn is likely to represent a

lower risk of infection compared to the risk of infection related to the personnel. In general, all four

transporting vehicle networks evaluated in this study contributed to PRRSV transmission, similarly to

previous studies (Thakur, Sanchez et al., 2015; Rossi et al., 2017; Porphyre et al., 2020; Yang et al., 2020).

It is important to remark that the contribution of the networks described above may have been

underestimated, mainly because we have not fully considered the effectiveness of cleaning and disinfection

to reduce PRRSV contamination in vehicles driving through cleaning stations. The lack of studies

measuring the effectiveness of cleaning and disinfections stations in reducing PRRSV and the contribution

of additional on-farm biosecurity cleaning and disinfection such as the presence of automatic cleaning

stations at the entrance of farms (Dee et al., 2004, 2007; Silva et al., 2019), is clearly needed in order to

better specify the reduction of such procedures in the force of PRRSV transmission.

The potential propagation of infectious diseases within feed ingredients has been of concern not

only to the swine industry but across other livestock systems; more recently studies have attempted to relate

different feed categories: blood products from livestock animals (animal by-products), cereal grains (i.e.

soybeans, corn, wheat), oil (canola, corn, soybean), forage, pellets (complete compound feed) and straw

(bedding material) with the propagation of ASFV (Gordon et al., 2019; EFSA Panel on Animal Health and

Welfare (AHAW) et al., 2021; Niederwerder, 2021). Despite the concerns about feed as a route of PRRSV

transmission, many uncertainties remain, including the minimal infection dosage required to cause disease,

the effectiveness of feed processing such as pelleting, extruding, and roasting and the use of feed additives

(Dee et al., 2020; Niederwerder, 2021). In addition, it is important to mention that feed contamination may

also occur within the feed mill facility either by contaminated environments, personnel, equipment, birds,

or rodents, or even contaminated trailers coming in and out feed mill facilities (Dee et al., 2020; Gebhardt

et al., 2021; Niederwerder, 2021), our model does not explicitly consider such uncertainties nor attempt to

account for such complexity. In this study we assumed that all feed meals with any amount of animal by-

products were still able to cause infection once delivered. In addition, we also assumed that the pelleting

process did not eliminate PRRSV contamination, and feed was delivered with enough viral load to cause

infection. Even though our results showed a relatively small role of animal by-products (less than 2.6%),

there are limitations to our approach and also limited information about the risk and pathways of collateral

contamination during or after feed manufacturing. For example, remain unknown the effect of high

temperatures and pressure used during pelleting in inactivating PRRSV (Benfield et al., 1992; Van Alstine

et al., 1993; Bloemraad et al., 1994; Cochrane et al., 2017), once such data becomes available it will be

possible to better assess the role of animal by-products in PRRSV transmission.

Limitations and final remarks

We identify a number of limitations related to the simulated scenarios and data availability. First, our

calculation approaches for the contribution of each transmission route was based on simulations that best

calibrated to the observed PRRSV cases in space and time. However, while the sensitivity of the final

simulations to detect PRRSV outbreaks was good, results may change once more data about other routes

of transmission (e.g. rendering networks) or other relevant disease control interventions are considered such

as vaccination programs or on-farm biosecurity. Additionally, the lack of data on transportation vehicles

movements and feed delivery of companies B and C limited our ability to examine possible differences and

similarities about the contribution of those routes among companies. Even with such limitations, it is

important to mention that companies B and C were included in the model because of the relevance of the

local transmission in the company-to-company spread of PRRSV, described in details elsewhere (Jara et

al., 2020). The 1.5 km radius used to define when a vehicle visited a farm or cleaning station was a limitation

for reconstructing the vehicle networks for sites that were 1.5 km from each other, thus in some cases, the

contacts were counted towards two or more instead of a single farm (median number of neighbors into 1.5

km = 1). A future alternative to reduce such events would require geographic information of each farms'

feed bins, for example. Additionally, the on-farm model parameters were oversimplified, since we have

based those estimations through historical records of PRRSV outbreaks, in which farms with fewer

infections were considered to have better biosecurity levels. Although there are several ways that the current

version of PigSpread model can be expanded, the inclusions of specific on-farm biosecurity practices and

infrastructure (e.g. present of cleaning and disinfection stations) could not only improve model calibration,

but to analyze the role of individualized and combined biosecurity on PRRSV dissemination (Sykes et al.,

2021). Another important limitation of our modelling work was the lack of information about the

vaccination strategies used by each farm, which could have contributed to the probability of new PRRSV

outbreaks (Galvis et al., 2021). Finally, we highlight that additional studies are necessary to understand the

effect of EVI and seasons on the local transmission, and to evaluate the performance of the gravity model

against other transmission methods, such as density kernels, to fit PRRSV outbreak locations. Despite the

limitations, this is the first study modelling simultaneously nine routes involved in PRRSV dissemination

dynamics over an entire year of outbreak data. This study is unique because it provides the swine industry

and regulatory agencies with robust and essential results about the dynamics of between-farms swine

disease transmission and the most relevant routes of transmission, offering an unique opportunity to

enhance the control of endemic disease and also prepare for future threats (Herrera-Ibatá et al., 2018; Jurado

et al., 2019; Brown et al., 2020).

Conclusion

We expanded a previously developed stochastic PRRSV simulation model (Galvis et al., 2021) to account

and quantify the contribution of nine different routes of between-farm transmission, including for the first

time the role of animal by-products delivered via feed meals and multiple transportation vehicle networks.

Our results demonstrate that transportation vehicle networks have a greater potential to spread PRRSV

when compared with the movement of pigs between-farms. In addition, vehicles transporting feed

represented the highest risk for PRRSV propagation in comparison with other vehicle networks, connecting

around 85% of farms. The temporal network was well represented by its static view for the networks of

vehicles transporting feed, and pigs to farms, with causal fidelity values >89%, thus we infer that studies

using a static view for vehicle networks are well supported when temporal data is not accessible. Our model

demonstrated that pig movements and local transmission remained the main routes of PRRSV transmission

regardless of farm types, but vehicles transporting pigs to farms also explained a significant proportion of

the farm infections: sow = 20.9%; nursery = 15%; and finisher = 20.6%. As expected, vehicles transporting

pigs to markets were more important for PRRSV introduction into finisher farms (3.8%), while vehicles

transporting feed showed the highest transmission contribution to sow farms (12%), while the vehicles

transporting crew had limited contribution in the propagation of PRRSV regardless of farm types. Finally,

animal fat and meat and bone meal delivered via feed contributed to 2.5% and 0.03% of sow farm infections,

respectively. Even though we were able to uncover the contribution of by-products and networks of several

transportation vehicles in the dissemination of PRRSV, we highlight the need for experimental or

observational studies able to measure the viability of PRRSV within feed formulation and the exterior or

transportation vehicles. Ultimately, this study provides a better understanding of the role of several

transmission routes for PRRSV dissemination, and can provide bases to the swine industry to evaluate and

strengthen the surveillance of transportation vehicles and feed delivery to better contain the propagation of

PRRSV.

Acknowledgements

The primary funding support of this project is from the Fat and Proteins Research Foundation. The Morrison

Swine Health Monitoring Project is a Swine Health Information Center funded project. Authors would like

to acknowledge participating companies, veterinarians and Scott Dee for his useful discussion about the

role of feed in the transmission of swine diseases.

Authors' contributions

JAG and GM conceived the study. JAG and GM participated in the design of the study. CC coordinated the

disease data collection by the Morrison Swine Health Monitoring Program (MSHMP). JAG and GM

conducted data processing, cleaning, designed the model, and simulated scenarios. JAG and GM designed

the computational analysis. JAG and GM wrote and edited the manuscript. All authors discussed the results

and critically reviewed the manuscript. GM secured the funding.

Conflict of interest

All authors confirm that the funding agency or other third parties had no role in the study design,

interpretation of results, writing manuscript and publication process.

Ethical statement

The authors confirm the ethical policies of the journal, as noted on the journal's author guidelines page.

Since this work did not involve animal sampling nor questionnaire data collection by the researchers, there

was no need for ethics permits.

Data Availability Statement

The data that support the findings of this study are not publicly available and are protected by confidential

agreements, therefore, are not available.

Funding

This project was funded by the Fats and Proteins Research Foundation.

References

Benfield, D.A., E. Nelson, J.E. Collins, L. Harris, S.M. Goyal, D. Robison, W.T. Christianson, R.B.

Morrison, D. Gorcyca, and D. Chladek, 1992: Characterization of Swine Infertility and

Respiratory Syndrome (SIRS) Virus (Isolate ATCC VR-2332). *J. Vet. Diagn. Invest.* **4**, 127–133,

DOI: 10.1177/104063879200400202.

Bernini, A., L. Bolzoni, and R. Casagrandi, 2019: When resolution does matter: Modelling indirect

contacts in dairy farms at different levels of detail. (Javier Sanchez, Ed.)*PLOS ONE* **14**,

e0223652, DOI: 10.1371/journal.pone.0223652.

Blázquez, E., C. Rodríguez, J. Ródenas, J. Segalés, J. Pujols, and J. Polo, 2020: Biosafety steps in the

manufacturing process of spray-dried plasma: a review with emphasis on the use of ultraviolet

irradiation as a redundant biosafety procedure. *Porc. Health Manag.* **6**, 16, DOI: 10.1186/s40813-

020-00155-1.

Bloemraad, M., E.P. de Kluijver, A. Petersen, G.E. Burkhardt, and G. Wensvoort, 1994: Porcine

reproductive and respiratory syndrome: temperature and pH stability of Lelystad virus and its

survival in tissue specimens from viraemic pigs. *Vet. Microbiol.* **42**, 361–371, DOI:

10.1016/0378-1135(94)90067-1.

Brown, V.R., R.S. Miller, S.C. McKee, K.H. Ernst, N.M. Didero, R.M. Maison, M.J. Grady, and S.A.

Shwiff, 2020: Risks of introduction and economic consequences associated with African swine

fever, classical swine fever and foot-and-mouth disease: A review of the literature. *Transbound.*

*Emerg. Dis.*tbed.13919, DOI: 10.1111/tbed.13919.

Büttner, K., and J. Krieter, 2020: Illustration of Different Disease Transmission Routes in a Pig Trade

Network by Monopartite and Bipartite Representation. *Animals* **10**, 1071, DOI:

10.3390/ani10061071.

Chase-Topping, M., J. Xie, C. Pooley, I. Trus, C. Bonckaert, K. Rediger, R.I. Bailey, H. Brown, V.

Bitsouni, M.B. Barrio, S. Gueguen, H. Nauwynck, and A. Doeschl-Wilson, 2020: New insights

about vaccine effectiveness: Impact of attenuated PRRS-strain vaccination on heterologous strain

transmission. *Vaccine* **38**, 3050–3061, DOI: 10.1016/j.vaccine.2020.02.015.

Cochrane, R.A., L.L. Schumacher, S.S. Dritz, J.C. Woodworth, A.R. Huss, C.R. Stark, J.M. DeRouchey,

M.D. Tokach, R.D. Goodband, J. Bia, Q. Chen, J. Zhang, P.C. Gauger, R.J. Derscheid, D.R.

Magstadt, R.G. Main, and C.K. Jones, 2017: Effect of pelleting on survival of porcine epidemic

diarrhea virus–contaminated feed. *J. Anim. Sci.* **95**, 1170–1178, DOI: 10.2527/jas.2016.0961.

Dee, S. A., M. Torremorell, R. Thompson, P. Cano, J. Deen, and C. Pijoan, 2007: Evaluation of the

thermo-assisted drying and decontamination system for sanitation of a full-size transport vehicle

contaminated with porcine reproductive and respiratory syndrome virus. *J Swine Health Prod* **15**,

12–18.

Dee, S., J. Deen, D. Burns, G. Douthit, and C. Pijoan, 2004: An assessment of sanitation protocols for

commercial transport vehicles contaminated with porcine reproductive and respiratory syndrome

virus. *Can. J. Vet. Res. Rev. Can. Rech. Veterinaire* **68**, 208–214.

Dee, S., J. Deen, K. Rossow, C. Weise, R. Eliason, S. Otake, H.S. Joo, and C. Pijoan, 2003: Mechanical

transmission of porcine reproductive and respiratory syndrome virus throughout a coordinated

sequence of events during warm weather. *Can. J. Vet. Res. Rev. Can. Rech. Veterinaire* **67**, 12–

19.

Dee, S., J. Deen, K. Rossow, C. Wiese, S. Otake, H.S. Joo, and C. Pijoan, 2002: Mechanical transmission

of porcine reproductive and respiratory syndrome virus throughout a coordinated sequence of

events during cold weather. *Can. J. Vet. Res. Rev. Can. Rech. Veterinaire* **66**, 232–239.

Dee, Scott A., J. Deen, S. Otake, and C. Pijoan, 2004: An experimental model to evaluate the role of

transport vehicles as a source of transmission of porcine reproductive and respiratory syndrome

virus to susceptible pigs. *Can. J. Vet. Res. Rev. Can. Rech. Veterinaire* **68**, 128–133.

Dee, Scott A., M.C. Niederwerder, R. Edler, D. Hanson, A. Singrey, R. Cochrane, G. Spronk, and E.

Nelson, 2020: An evaluation of additives for mitigating the risk of virus-contaminated feed using

an ice-block challenge model. *Transbound. Emerg. Dis.* **n/a**, DOI:

https://doi.org/10.1111/tbed.13749.

Duinhof, T.F., G. van Schaik, E.J.B. van Esch, and G.J. Wellenberg, 2011: Detection of PRRSV

circulation in herds without clinical signs of PRRS: Comparison of five age groups to assess the

preferred age group and sample size. *Vet. Microbiol.* **150**, 180–184, DOI:

10.1016/j.vetmic.2011.01.001.

EFSA Panel on Animal Health and Welfare (AHAW), S.S. Nielsen, J. Alvarez, D.J. Bicout, P. Calistri, E.

Canali, J.A. Drewe, B. Garin-Bastuji, J.L. Gonzales Rojas, C. Gortázar Schmidt, M. Herskin,

M.Á. Miranda Chueca, V. Michel, B. Padalino, P. Pasquali, L.H. Sihvonen, H. Spoolder, K.

Stahl, A. Velarde, A. Viltrop, C. Winckler, A. Boklund, A. Botner, A. Gervelmeyer, O. Mosbach-

Schulz, and H.C. Roberts, 2021: Ability of different matrices to transmit African swine fever

virus. *EFSA J.* **19**, DOI: 10.2903/j.efsa.2021.6558.

Galvis, J.A., J.M. Prada, C.A. Corzo, and G. Machado, 2021: Modeling the transmission and vaccination

strategy for porcine reproductive and respiratory syndrome virus. *Transbound. Emerg.*

*Dis.*tbed.14007, DOI: 10.1111/tbed.14007.

Gao, L., X. Sun, H. Yang, Q. Xu, J. Li, J. Kang, P. Liu, Y. Zhang, Y. Wang, and B. Huang, 2021:

Epidemic situation and control measures of African Swine Fever Outbreaks in China 2018–2020.

Transbound. Emerg. Dis. **68**, 2676–2686, DOI: 10.1111/tbed.13968.

Gebhardt, J.T., S.S. Dritz, C.G. Elijah, C.K. Jones, C.B. Paulk, and J.C. Woodworth, 2021: Sampling and

detection of African swine fever virus within a feed manufacturing and swine production system.

*Transbound. Emerg. Dis.*tbed.14335, DOI: 10.1111/tbed.14335.

Gordon, R.K., I.K. Kotowski, K.F. Coulson, D. Link, A. MacKenzie, and J. Bowling-Heyward, 2019:

The Role of Non-animal Origin Feed Ingredients in Transmission of Viral Pathogens of Swine: A

Review of Scientific Literature. *Front. Vet. Sci.* **6**, 273, DOI: 10.3389/fvets.2019.00273.

Henry, A.E., A. Letellier, J.-C. Côté, G. Desmarais, V. Lachapelle, N. Bergeron, S.L. Lewandowsky, and

P. Fravalo, 2018: Overlooked sources of Salmonella contamination in the pig production

network: Slaughterhouse yard pathways and mudguards and carpets from transport trucks. *Can.*

Vet. J. Rev. Veterinaire Can. **59**, 1105–1108.

Herrera-Ibatá, D.M., B. Martínez-López, D. Quijada, K. Burton, and L. Mur, 2018: Correction:

Quantitative approach for the risk assessment of African swine fever and Classical swine fever

introduction into the United States through legal imports of pigs and swine products. *PLOS ONE*

13, e0208065, DOI: 10.1371/journal.pone.0208065.

Holtkamp, D., J. Kliebenstein, and E. Neumann, 2013: Assessment of the economic impact of porcine

reproductive and respiratory syndrome virus on United States pork producers. *J. Swine Health*

Prod. **21**, 13.

Holtkamp, D. J., Morrison, B., Classen, D. M., Becton, L., Henry, S., Rodibaugh, M. T., Rowland, R. R.,

Snelson, H., Straw, B., Yeske, P., & Zimmerman, J, 2011: Terminology for classifying swine

herds by PRRS status. *J. Swine Health Prod.* **19**, 44– 56.

Holtkamp, D.J., P.E. Yeske, D.D. Polson, J.L. Melody, and R.C. Philips, 2010: A prospective study

evaluating duration of swine breeding herd PRRS virus-free status and its relationship with

measured risk. *Prev. Vet. Med.* **96**, 186–193, DOI: 10.1016/j.prevetmed.2010.06.016.

Jacobs, A.C., J.R. Hermann, C. Muñoz-Zanzi, J.R. Prickett, M.B. Roof, K.-J. Yoon, and J.J. Zimmerman,

2010: Stability of Porcine Reproductive and Respiratory Syndrome virus at Ambient

Temperatures. *J. Vet. Diagn. Invest.* **22**, 257–260, DOI: 10.1177/104063871002200216.

Jara, M., D.A. Rasmussen, C.A. Corzo, and G. Machado, 2020: Porcine reproductive and respiratory

syndrome virus dissemination across pig production systems in the United States. *Transbound.*

*Emerg. Dis.*tbed.13728, DOI: 10.1111/tbed.13728.

Jurado, C., L. Mur, M.S. Pérez Aguirreburualde, E. Cadenas-Fernández, B. Martínez-López, J.M.

Sánchez-Vizcaíno, and A. Perez, 2019: Risk of African swine fever virus introduction into the

United States through smuggling of pork in air passenger luggage. *Sci. Rep.* **9**, 14423, DOI:

10.1038/s41598-019-50403-w.

Lee, K., D. Polson, E. Lowe, R. Main, D. Holtkamp, and B. Martínez-López, 2017: Unraveling the

contact patterns and network structure of pig shipments in the United States and its association

with porcine reproductive and respiratory syndrome virus (PRRSV) outbreaks. *Prev. Vet. Med.*

138, 113–123, DOI: 10.1016/j.prevetmed.2017.02.001.

Lentz, H.H.K., A. Koher, P. Hövel, J. Gethmann, C. Sauter-Louis, T. Selhorst, and F.J. Conraths, 2016:

Disease Spread through Animal Movements: A Static and Temporal Network Analysis of Pig

Trade in Germany. (Thierry Boulinier, Ed.)*PLOS ONE* **11**, e0155196, DOI:

10.1371/journal.pone.0155196.

Lewis, A.J., and L.L. Southern (Eds.), 2001: Swine Nutrition2nd ed. Boca Raton: CRC Press.

Magar, R., and R. Larochelle, 2004: Evaluation of the presence of porcine reproductive and respiratory

syndrome virus in pig meat and experimental transmission following oral exposure. *Can. J. Vet.*

Res. **68**, 259–266.

Makau, D.N., M.A. Alkhamis, I.A.D. Paploski, C.A. Corzo, S. Lycett, and K. VanderWaal, 2021:

Integrating Animal Movements With Phylogeography to Model the Spread of PRRS Virus in the

U.S. *Virus Evol.*veab060, DOI: 10.1093/ve/veab060.

Melmer, D.J., T.L. O'Sullivan, A.L. Greer, and Z. Poljak, 2020: An investigation of transportation

practices in an Ontario swine system using descriptive network analysis. (Quan Yuan, Ed.)*PLOS*

ONE **15**, e0226813, DOI: 10.1371/journal.pone.0226813.

Minter, A., and R. Retkute, 2019: Approximate Bayesian Computation for infectious disease modelling.

Epidemics **29**, 100368, DOI: 10.1016/j.epidem.2019.100368.

MSHMP, 2020: PRRS Charts [Online].

Neira, V., B. Brito, J. Mena, M. Culhane, M.I. Apel, V. Max, P. Perez, V. Moreno, C. Mathieu, M.

Johow, C. Badia, M. Torremorell, R. Medina, and R. Ortega, 2017: Epidemiological

investigations of the introduction of porcine reproductive and respiratory syndrome virus in

Chile, 2013-2015. (Sanjay B. Jadhao, Ed.)*PLOS ONE* **12**, e0181569, DOI:

10.1371/journal.pone.0181569.

Niederwerder, M., 2021: Risk and Mitigation of African Swine Fever Virus in Feed. *Animals* **11**, 792,

DOI: 10.3390/ani11030792.

Ochoa, L., L. Greiner, and T. Pacion, 2018: 119 Feed As a Vehicle for Prrs Virus Transmission and the

Effects of Formaldehyde on Porcine Reproductive and Respiratory Virus in Feed: Proof of

Concept. *J. Anim. Sci.* **96**, 63–63, DOI: 10.1093/jas/sky073.117.

Passafaro, T.L., A.F.A. Fernandes, B.D. Valente, N.H. Williams, and G.J.M. Rosa, 2020: Network

analysis of swine movements in a multi-site pig production system in Iowa, USA. *Prev. Vet. Med.*

174, 104856, DOI: 10.1016/j.prevetmed.2019.104856.

Perez, A.M., P.R. Davies, C.K. Goodell, D.J. Holtkamp, E. Mondaca-Fernández, Z. Poljak, S.J.

Tousignant, P. Valdes-Donoso, J.J. Zimmerman, and R.B. Morrison, 2015: Lessons learned and

knowledge gaps about the epidemiology and control of porcine reproductive and respiratory

syndrome virus in North America. *J. Am. Vet. Med. Assoc.* **246**, 1304–1317, DOI:

10.2460/javma.246.12.1304.

Pileri, E., and E. Mateu, 2016: Review on the transmission porcine reproductive and respiratory syndrome

virus between pigs and farms and impact on vaccination. *Vet. Res.* **47**, 108, DOI:

10.1186/s13567-016-0391-4.

Pitkin, A., J. Deen, and S. Dee, 2009: Further assessment of fomites and personnel as vehicles for the

mechanical transport and transmission of porcine reproductive and respiratory syndrome virus.

Can. J. Vet. Res. Rev. Can. Rech. Veterinaire **73**, 298–302.

Porphyre, T., B.M. de C. Bronsvoort, G.J. Gunn, and C. Correia-Gomes, 2020: Multilayer network

analysis unravels haulage vehicles as a hidden threat to the British swine industry. *Transbound.*

Emerg. Dis. **67**, 1231–1246, DOI: https://doi.org/10.1111/tbed.13459.

Rossi, G., R.L. Smith, S. Pongolini, and L. Bolzoni, 2017: Modelling farm-to-farm disease transmission

through personnel movements: from visits to contacts, and back. *Sci. Rep.* **7**, 2375, DOI:

10.1038/s41598-017-02567-6.

Sanhueza, J.M., C. Vilalta, C. Corzo, and A.G. Arruda, 2019: Factors affecting Porcine Reproductive and

Respiratory Syndrome virus time-to-stability in breeding herds in the Midwestern United States.

Transbound. Emerg. Dis. **66**, 823–830, DOI: 10.1111/tbed.13091.

Shirai, J., T. Kanno, Y. Tsuchiya, S. Mitsubayashi, and R. Seki, 2000: Effects of Chlorine, Iodine, and

Quaternary Ammonium Compound Disinfectants on Several Exotic Disease Viruses. *J. Vet. Med.*

Sci. **62**, 85–92, DOI: 10.1292/jvms.62.85.

Silva, G.S., G. Machado, K.L. Baker, D.J. Holtkamp, and D.C.L. Linhares, 2019: Machine-learning

algorithms to identify key biosecurity practices and factors associated with breeding herds

reporting PRRS outbreak. *Prev. Vet. Med.* **171**, 104749, DOI: 10.1016/j.prevetmed.2019.104749.

Sterchi, M., C. Faverjon, C. Sarasua, M.E. Vargas, J. Berezowski, A. Bernstein, R. Grütter, and H.

Nathues, 2019: The pig transport network in Switzerland: Structure, patterns, and implications for

the transmission of infectious diseases between animal holdings. (Esteban Hernandez-Vargas,

Ed.)*PLOS ONE* **14**, e0217974, DOI: 10.1371/journal.pone.0217974.

Sykes, A.L., G.S. Silva, D.J. Holtkamp, B.W. Mauch, O. Osemeke, D.C.L. Linhares, and G. Machado,

2021: Interpretable machine learning applied to on-farm biosecurity and porcine reproductive and

respiratory syndrome virus. *ArXiv210606506 Q-Bio*.

Thakur, K., C. Revie, D. Hurnik, and J. Sánchez, 2017: Modelling contamination of trucks used in the

shipment of pigs infected with porcine reproductive and respiratory syndrome virus. *J. Swine*

Health Prod. **25**, 183–193.

Thakur, K.K., C.W. Revie, D. Hurnik, Z. Poljak, and J. Sanchez, 2015: Simulation of between-farm

transmission of porcine reproductive and respiratory syndrome virus in Ontario, Canada using the

North American Animal Disease Spread Model. *Prev. Vet. Med.* **118**, 413–426, DOI:

710 10.1016/j.prevetmed.2015.01.006.

711 Thakur, K.K., J. Sanchez, D. Hurnik, Z. Poljak, S. Opps, and C.W. Revie, 2015: Development of a

712 network based model to simulate the between-farm transmission of the porcine reproductive and

713 respiratory syndrome virus. *Vet. Microbiol.* **180**, 212–222, DOI: 10.1016/j.vetmic.2015.09.010.

714 Van Alstine, W.G., C.L. Kanitz, and G.W. Stevenson, 1993: Time and Temperature Survivability of

715 PRRS Virus in Serum and Tissues. *J. Vet. Diagn. Invest.* **5**, 621–622, DOI:

716 10.1177/104063879300500421.

717 Wiltshire, S.W., 2018: Using an agent-based model to evaluate the effect of producer specialization on

718 the epidemiological resilience of livestock production networks. *PLoS ONE* **13**, DOI:

719 10.1371/journal.pone.0194013.

720 Yang, Q., D.M. Gruenbacher, J.L.H. Stamm, D.E. Amrine, G.L. Brase, S.A. DeLoach, and C.M. Scoglio,

721 2020: Impact of truck contamination and information sharing on foot-and-mouth disease

722 spreading in beef cattle production systems. *PLOS ONE* **15**, e0240819, DOI:

723 10.1371/journal.pone.0240819.

724 Yoo, D.S., Y. Kim, E.S. Lee, J.S. Lim, S.K. Hong, I.S. Lee, C.S. Jung, H.C. Yoon, S.H. Wee, D.U.

725 Pfeiffer, and G. Fournié, 2021: Transmission Dynamics of African Swine Fever Virus, South

726 Korea, 2019. *Emerg. Infect. Dis.* **27**, 1909–1918, DOI: 10.3201/eid2707.204230.

728 **Table list**

729 **Table 1**. Summary of the network metric of pig movements and vehicle movements of the three different

730 pig producing companies, with data from January 2020 until December 2020.

Company	A		B	C
		Vehicles transporting:		

Network metric/type	Pig movement*	Feed to farms	Pigs to farms	Pigs to markets	Crew	Pig movement*	Pig movement*
Number of nodes (farms)	1,745	1,745	1,745	1,745	1,745	228	321
Number of edges, weekly temporal network	34,833	3,182,144	386,730	35,854	109,675	4,344	3,882
Number of edges, static network	5,664	603,977	43,968	18,836	53,880	692	553
Network density	0.002	0.2	0.014	0.006	0.018	0.013	0.005
Causal fidelity	43.2%	98.7%	89.6%	58%	76.9%	32.9%	61.9%
Number of strong connected components (SCC) groups[1]	9	2	6	7	3	1	0
Number of farms in the Largest SCC	11	1,591	1,479	976	1,058	61	1
In-degree[2]	3	319	19	5	14	2	1
Out-degree[2]	1	304	21	6	16	1	0
Betweenness centrality[2]	0	552.3	920.66	111.25	51.32	0.53	0
Ingoing contact chain[2]	34	1,515	27	12	131	15	4
Outgoing contact chain[2]	0	1,502	32	14	119	15	4

731   [1]Number of SCC groups with more than two farms.

732   [2]Median values

733   *Pig movement stands for the movement of live pigs between farms.

Figure legends

Figure 1. Framework of the indirect farm contacts formed by transportation vehicle movements.
The transportation vehicle networks were reconstructed based on consecutive farm visits of each vehicle.
Because the stability of PRRSV in the environment is directly impacted by the environmental conditions,
a contact network was reconstructed considering all edges between farm visits that happened within 72
hours for cold months (from October until March) and within 24 hours for warm months (from April until
September).

Figure 2. Transmission model framework. (a) Model flowchart of the farm's infectious status and routes
of PRSV transmission, and (b) description of the model transmission parameters. In the example, for the
local infection force, j takes the values from 1 and 2 because these are the only two farms within the
maximum radius distance.

Figure 3. Farm infection contribution for each transmission route of each farm type (rows). The *y*-
axis represents the proportion of transmission by each transmission route, while the x-axis shows each week
in the simulation. Weekly proportions of transmission were calculated by dividing the number of simulated
infected farms for the total number of routes combined.

Figure 1.

(b) Model parameters

β_n = Transmission rate of between farm pig movements

β_l = Local transmission rate

βr = Re-break rate

β_f = Transmission rate of between farm movements of vehicles transporting feed

β_p = Transmission rate of between farm movements of vehicles transporting pig to farm

β_m = Transmission rate of between farm movements of vehicles transporting pig to market

β_c = transmission rate of between farm movements of vehicles transporting crew to farms

β_a = Fat in the delivered feed rate

β_b = Meat and bone in the delivered feed rate

N = Number of asymptomatic and infected farms that sent pigs to "i"

g = Gravity model with barrier effect

E = Edge weight

Z = Time vehicle stay on the farm

A = Amount of fat in the meal

B = Amount of meat and bone in the meal

F = Pig population in the farm

W = Re-break probability based on the time after last outbreak

T = Monthly seasonality index

H = Biosecurity index

L = Detection probability

x0 = Average time detection

x = Time post virus introduction

k = Logistic growth rate

Figure 2.

Figure 3.

Supplementary Material

Modelling and assessing additional transmission routes for porcine reproductive and respiratory

syndrome virus: vehicle movements and feed ingredients

Jason A. Galvis [1], Cesar A. Corzo[2], Gustavo Machado[1]*

[1]Department of Population Health and Pathobiology, College of Veterinary Medicine, Raleigh, North

Carolina, USA.

[2]Veterinary Population Medicine Department, College of Veterinary Medicine, University of Minnesota,

St Paul, MN, USA.

*Corresponding author: gmachad@ncsu.edu

Section 1. A descriptive analysis of model parameters used in the PRRSV simulated transmission.

Figure S1. The distribution of PRRSV stability probability on vehicle surface over time. In A), we show

the cold season decay which included the months between October to March, and B) warm season decay

in PRRSV suitability which included the months between April to September. In summary, here we

assumed that PRRSV suitability decreased linearly over time after a vehicle visited a farm, down to zero

after 72 hours for cold months and 24 hours for warm months, when PRRSV was no longer viable. This

approach was used for all vehicle related contact networks including vehicles transporting feed, pigs-to-

farm, pigs-to-market and crew.

Figure S2. The overall proportion of animal by-products delivered to farms from January 2020 until

December 2020. The x-axis shows the number of unique feed formulations and the y-axis each animal by-

product utilized by company A.

Figure S3. The weight (lb) distribution of animal by-products in shipments of feed formulation delivered

to the farms of company A in the study area over from January 2020 until December 2020.

Table S1. Terminology and definition from the social network analysis.

Network terminology	Definition	Reference
Node	Element of the network representing the farms.	-
Edge	Link among two nodes.	-
Static network	Once an edge exists between two nodes, it is present for the whole time period.	(Kao et al., 2007)
Temporal network	The edges between two nodes only exist at different time steps.	(Lentz et al., 2016)
Density	Represent the proportion of edges among nodes in the network that are actually present.	(Wasserman and Faust, 1994)
Causal fidelity	Quantify the error of the static representation of a temporal network by comparing the number of paths in the static and temporal networks.	(Lentz et al., 2016)
Strongly	It is a subset of nodes for which a *directed* path exists	(Lentz et al., 2016)

Connected Component (SCC)	between all pairs of them allowing them to be mutually accessible by following the direction of the links in the network.	
Larger Strongly Connected Component (LSCC)	It is the larger number of nodes in a strong connected component.	(Lentz et al., 2016)
In-degree	Number of nodes providing animals to a specific node.	(Wasserman and Faust, 1994)
Out-degree	Number of nodes obtaining animals from a specific node.	(Wasserman and Faust, 1994)
Betweenness	The frequency at which a node is in the shortest path between pairs of other nodes in the network.	(Freeman, 1978)
Ingoing contact chain (ICC)	Subsets of nodes that can reach a specific node by direct contact or indirect contacts through a sequential order of edges through other nodes using the temporal network.	(Nöremark and Widgren, 2014)
Outgoing contact chain (OCC)	Subsets of nodes that can be reached by a specific node by direct contact or indirect contacts through a sequential order of edges through other nodes using the temporal network.	(Nöremark and Widgren, 2014)

To calculate the barrier index (vegetation level, utilized to modulate the probability of local transmission),

we used a linear regression, to express PRRSV infected farms from 2020 as a function of the Enhanced

Vegetation Index (EVI) and yearly seasonality (spring, summer, fall and winter). We found that PRRSV

frequency decreased as EVI increased, with a stronger association in winter and fall seasons (Figure S4).

Here we used the regression coefficients to predict weekly PRRSV incidence, which then were transformed

into parameter a, which was scaled into values between [0, 1], later utilized to modulate the local

transmission.

Figure S4. Linear regression of PRRSV infected farms. The y-axis is the number of PRRSV outbreaks

and in the x-axis EVI. Fall: coefficient -0.00029, standard error 0.00018, *p*-value 0.12, Spring: coefficient

-0.00001, standard error 0.00023, *p*-value 0.96, Summer: coefficient 0.00013, standard error 0.00018, *p*-

value 0.46. Winter: coefficient -0.00103, standard error 0.0004, *p*-value 0.02.

Figure S5. The time vehicles spent on each farm visit. The boxplot shows the distribution in minutes that

each vehicle remained within farms premises in A) and at cleaning stations in B).

Figure S6. A survival analysis of infected and recovered farms from 2009 to 2019. In this example we

show the distribution used for each farm to calibrate the re-break probability

Figure S7. Monthly seasonality index calculated from the frequency of PRRSV, calculated by an additive

moving average decomposition derived from analysis of the PRRSV records from 2015 to 2019.

Section 2: Model calibration and main model outputs

The model is calibrated in two steps, first fitting the frequency of observed outbreaks on time, and second

fitting the location of the observed outbreaks. In Table S2 we show the summary statistics used in step 1 of

the Approximate Bayesian Computation (ABC) rejection algorithm, where the tolerance interval represents

the square error allowed from the simulation to the observed values. In the model calibration, a particle is

defined as a set of transmission parameters (Table S3). For the model calibration in step 1, we defined as

an accepted particle the set of transmission parameters, whose simulation results are lower than the

tolerance interval for all the variables when compared with the observed values. Thus, the range of error

accepted (tolerance interval) defines the number of particles accepted in the step 1 of the model calibration.

In addition, low tolerance interval values produce particles with similar results to the observed values, but

at the same time, the number of particles accepted decreases. Thus, in this study the tolerance interval values

were chosen through initial exploratory analysis, selecting the minimum tolerance interval values to accept

an adequate number of particles to be analyzed in the step 2 of the model calibration, in this case for each

100,000 particles tested around 20 were accepted in the step 1.

Table S2. Summary statistics used by the ABC Sequential Monte Carlo rejection algorithm for the model.

Summary statistics	Observed values	Tolerance interval (ϵ)
Total number of sow farms with detected cases	96	20
The weekly average number of sow farms with detected cases	1.8	0.5
The weekly maximum number of sow farms with detected cases	9	5
Total number of nursery farms with detected cases	37	20

The weekly average number of nursery farms with detected cases	0.7	0.5
The weekly maximum number of nursery farms with detected cases	6	5
Total number of finisher farms with detected cases	17	20
The weekly average number of finisher farms with detected cases	0.3	0.5
The weekly maximum number of finisher farms with detected cases	2	5
Expected prevalence in finisher and nursery farms	30% (expert opinion)	100

To assess the model performance, we evaluated the probability to predict cells (10 x 10 km squares) with

true infected cells (cells where at least one sow farm outbreak was recorded) at time t. Each sow farm was

allocated to a cell in the spatial grid (total of 154 cells in the study area). For each particle accepted in the

step 1 model fitting, the risk of each cell was calculated by the sum of times at least one farm within a cell

was identified with infected status after 100 simulations; we utilized a percentiles thresholds (r) approach

to determine cells at high and low risk, where high risk cells were compared with the true infected cells at

time t. Subsequently we estimate the model sensitivity and specificity, for all thresholds, as follows:

$$S_r = TP_r/(TP_r + FN_r)$$

$$E_r = TN_r/(TN_r + FP_r)$$

where true positives (TP) was the subset of cells with observed outbreaks and the estimated risk was above

the r threshold; false negatives (FN) was the subset of cells with observed outbreaks and the estimated risk

was below the r threshold; true negative (TN) was the subset of cells without observed outbreaks and the

estimated risk was below the r threshold; and false positives (FP) was the subset of cells without observed

outbreaks and the estimated risk above the r threshold.

In step 2 of the ABC rejection algorithm, the sensitivity and specificity were calculated for each

particle accepted in the step 1 of model fitting. The particles accepted were those with sensitivity values

$\geq$30% with r = 85th and $\geq$50% with r = 70^{th} (percentile and sensitivity thresholds values were chosen

arbitrarily after authors discussed the minimum performance of the model). The priors for each parameter

were drawn from a uniform distribution that ranged between 0 and 1.5 for pig movements transmission

rate, 0 and 0.001 for local transmission, the four transporting vehicles and amount of animal fat and meat

and bone meal in the feed meals transmission rates, and finally between 0 and 0.01 for re-break transmission

rate. These range values were chosen according to model performance to fit the temporal and spatial

distribution of PRRSV cases through some test simulations, thus reducing the number of simulations and

processing time in the model calibration. It is worth noting that farm' biosecurity was only included for

sow farms, thus there are no parameters representing nursery, finisher or other farms. Finally, here we were

not able to describe the number simulation necessary to accept particles, similar how it was described in

step 1, because we did not track the number of simulation used in step 2.

Table S3. List of transmission parameters used in the simulations and values estimated from the posterior

distribution of the 100 particles accepted in the model calibration.

Model parameter	Symbol	Average values	95% credible interval*	Details & references
Transmission rate of pig movements	β_n	0.428	0.11-1.29	ABC fitting
Local transmission rate	β_l	0.00055	0.0001-0.0009	ABC fitting
Transmission rate of	β_f	0.000014	0.0000004-0.00004	ABC fitting

vehicles transporting feed				
Transmission rate of vehicles transporting pigs to farms	β_p	0.00026	0.000006-0.0009	ABC fitting
Transmission rate of vehicles transporting pigs to market	β_m	0.00049	0.00003-0.0009	ABC fitting
Transmission rate of vehicles transporting crew	β_c	0.00027	0.00003-0.0008	ABC fitting
Transmission rate of animal fat in the feed meal	β_a	0.00042	0.000002-0.0009	ABC fitting
Transmission rate of meat and bone meal in the feed meal	β_b	0.00042	0.000005-0.0009	ABC fitting
Transmission rate of re-break	β_r	0.0045	0.0001-0.009	ABC fitting
Farm' biosecurity	H(sow)	0.57	0.02-0.99	ABC fitting
Maximum effective surveillance	L(sow)	0.95	-	Expert opinion
	L(nurseries)	0.048	0.016-0.085	ABC fitting
	L(finisher)	0.0015	0.0001-0.003	ABC fitting
	L(others)	0.045	0.002-0.086	ABC fitting
PRRSV seasonality	T	Weekly values calculated	-	Figure S7

Average time for PRRSV detection	X0	4 weeks	-	Expert opinion
Average infectious time sow farms	-	41 weeks	-	(Sanhueza et al., 2019)

*Credible intervals calculated with method equal-tailed interval.

Figure S8. Posterior distribution of the calibrated transmission parameters derived from 100 accepted

particles, red lines represent 95% credible intervals calculated through equal-tailed interval method.

Figure S9. The simulated weekly number of infected farms in A) and infected detected farms (PRRSV

outbreaks) in B). The black line represents the median, the dark shade areas represent a 75% credible

interval and the light shade areas maximum and minimum generated by the model, and the red dots the

frequency of true outbreaks reported in our data. Uncertainty in the estimated model parameters is reflected

by 1,000 repeated simulations.

Figure S10. The average sensitivity and specificity for the weekly forecasts in A) and the average of all

weeks in B). These values were calculated from 100 model calculations with each model calculation having

100 individual model iterations per predicted week to estimate the spatial location of observed outbreaks

Section 3: Descriptive analysis of the between-farm pig movements and transportation vehicle movement

networks, and the quantity of animal by-product in feed ingredients.

Figure S11. Boxplot with the distribution of in-degree for between-farm pig movements of each

transportation vehicle movement networks.

Figure S12. Boxplot with the distribution of out-degree for between-farm pig movements of each

transportation vehicle movement networks.

Figure S13. Boxplot with the distribution of betweenness for between-farm pig movements of each

transportation vehicle movement networks.

Figure S14. Boxplot with the distribution of ingoing contact chains for between-farm pig movements of

each transportation vehicle movement networks.

Figure S15. Boxplot with the distribution of outgoing contact chains for between-farm pig movements of

each transportation vehicle movement networks.

Figure S16. Boxplot comparing the frequency of infected farms in the ingoing contact chain of infected

and non-infected farms of each transportation vehicle and pig movement networks. Infected farms are more

frequent in the ingoing contact chain of other infected farms for the vehicles transporting feed, pigs to farms

and pigs to market (Mann Whitney test $p < 0.05$). It is worth noticing that the statistical differences found

among the groups can be influenced by the presence of outliers in the distribution, as well as the large

number of farms analyzed (n = 2,294) that increase the test power and make it more likely to reject the null

hypothesis.

Figure S17. Boxplot comparing the frequency of infected farms in the outgoing contact chain of infected and non-infected farms of each transportation vehicle and pig movement networks. Infected farms are more frequent in the outgoing contact chain of other infected farms for pig movements and the vehicles transporting feed, pigs to farms and pigs to market (Mann Whitney test $p < 0.05$). It is worth noticing that the statistical differences found among the groups can be influenced by the presence of outliers in the

distribution, as well as the large number of farms analyzed (n = 2,294) that increase the test power and

make it more likely to reject the null hypothesis.

Figure S18. Boxplots compare the time vehicles remain on the farms of infected and non-infected farms

for the different transportation vehicles (rows) and production types (columns). Vehicles transporting feed

and crew to nursery farms were the only vehicles that showed a higher average of time on infected farms

(Mann Whitney test $p < 0.05$).

Figure S19. Boxplot comparing the distribution of **A**) animal fat and **B**) meat and bone meal in the feed

meal received by the sow farms with and without PRRSV records in 2020 (in red the result from the logistic

regression).

Figure S20. Boxplot comparing the distribution of **A**) animal fat and **B**) meat and bone meal, **C**) Protein

blend meal 58%, **D**) Protein blend and **E**) plasma in the feed meal received by the nursery farms with and

without PRRSV records in 2020 (in red the result from the logistic regression).

Figure S21. Boxplot comparing the distribution of **A**) animal fat and **B**) meat and bone meal, **C**) Protein

blend meal 58% and **D**) protein blend in the feed meal received by the finisher farms with and without

PRRSV records in 2020 (in red the result from the logistic regression).

 Section 4:contribution of the transmission routes.

For company B, re-break was the main source of farm infections in sow farms with a contribution of 42.6% (95% CI 0%-93%) to PRRSV transmission, followed by local transmission with 28.1% (95% CI 0.8%-88%) and pig movements with 27.3% (95% CI 0%-84%), for nursery 81.6% (95% CI 31%-96%) of PRRSV transmission was related to pig movements and 18.3% (95% CI 4%-68%) to local transmission, while in finishers 51.6% (95% CI 0%-83%) was related to pig movements and 48.4% (95% CI 16%-100%) to local transmission (Figure S22). Finally, for farms of company C, local transmission was the most important transmission route for sow farms contributing with 68.9% (95% CI 8%-97%) of the farm infections, followed by re-break with 26.1% (95% CI 0%-45%) and pig movements 1.2% (95% CI 0%-0%), in nursery 53.2% was related to local transmission (95% CI 8%-100%) and 46.8% (95% CI 0%-92%) to pig movements, and for finishers 78% was related to local transmission (95% CI 31%-100%) and 22% (95% CI 0%-69%) to pig movements (notice that the contribution of transmission routes from the same company and farm type may not reach 100% because for some weeks none of the transmission routes used in the simulations was able to infect farms) (Figure S22).

Figure S22. Farm infection contribution for each transmission route of each company (columns) and farm types (rows). The *y*-axis represents the proportion of transmission by each transmission route, while the x-axis shows each week in the simulation. Weekly proportions of transmission were calculated by dividing the number of simulated infected farms for each transmission route by the number of simulated

infected farms for the total number of routes combined. White areas represent weeks without farm
infections.

References

Freeman, L.C., 1978: Centrality in social networks conceptual clarification. *Soc. Netw.* **1**, 215–239, DOI:
10.1016/0378-8733(78)90021-7.

Kao, R.R., D.M. Green, J. Johnson, and I.Z. Kiss, 2007: Disease dynamics over very different time-
scales: foot-and-mouth disease and scrapie on the network of livestock movements in the UK. *J.*
R. Soc. Interface **4**, 907–916, DOI: 10.1098/rsif.2007.1129.

Lentz, H.H.K., A. Koher, P. Hövel, J. Gethmann, C. Sauter-Louis, T. Selhorst, and F.J. Conraths, 2016:
Disease Spread through Animal Movements: A Static and Temporal Network Analysis of Pig
Trade in Germany. (Thierry Boulinier, Ed.)*PLOS ONE* **11**, e0155196, DOI:
10.1371/journal.pone.0155196.

Nöremark, M., and S. Widgren, 2014: EpiContactTrace: an R-package for contact tracing during livestock
disease outbreaks and for risk-based surveillance. *BMC Vet. Res.* **10**, 71, DOI: 10.1186/1746-
6148-10-71.

Wasserman, S., and K. Faust, 1994: Social Network Analysis: Methods and Applications. Cambridge ;
New York: Cambridge University Press.