Directed yeast genome evolution by controlled introduction of trans-chromosomic structural variations

Bin Jia[1,2,*] , Jin Jin[1, 2,*] , Ming-Zhe Han[1, 2], Bing-Zhi Li[1, 2] & Ying-Jin Yuan[1,2,†]

[1] Frontier Science Center for Synthetic Biology and Key Laboratory of Systems Bioengineering (Ministry of Education), School of Chemical Engineering and Technology, Tianjin University, 300072 Tianjin, China.

[2] Collaborative Innovation Center of Chemical Science and Engineering (Tianjin), Tianjin University, 300072 Tianjin, China.

[*] Authors contributed equally to this work.

[†] Corresponding author (email: yjyuan@tju.edu.cn)

Abstract

Background: Naturally occurring structural variations (SVs) are a considerable source of genomic variation and can reshape chromosomes 3D architecture. The synthetic chromosome rearrangement and modification by loxP-mediated evolution (SCRaMbLE) system has been proved to generate random SVs to impact phenotypes and thus constitutes powerful drivers of directed genome evolution. However, controllable methods to introduce complex SVs and revealing the related molecular mechanism has so far remained challenging.

Results: We develop a SV-prone yeast strain by using SCRaMbLE with two synthetic chromosomes, synV and synX. An heterologous biosynthesis pathway allowing a high throughput screen for increased yield of astaxanthin is used as readout and a proof of

concept for the application of SV in industry. We report here that complex SVs, including a pericentric inversion and a trans-chromosomes translocation between synV and synX, result in two neochromosomes and a 2.7-fold yield of astaxanthin. We mapped genetic targets contributing to higher astaxanthin yield and demonstrated the SVs can led to large reorganization of the genetic information along the chromosomes. We also used the model learned from the aforementioned random screen and successfully harnessed the precise introduction of trans-chromosomes translocation and pericentric inversions by rational design.

Conclusions: Our work provides an effective tool to not only accelerate the directed genome evolution but also reveal mechanistic insight of complex SVs for altering phenotypes.

Key words: Directed genome evolution, Structural variations, SCRaMbLE, Synthetic genome, Synthetic biology

Introduction

Chromosomes are highly dynamic objects that often undergo large structural variations(SVs), which in turn can reshape chromosomes and impact phenotypes[1-11]. Naturally occurring SVs comprised both unbalanced SVs that are copy number variations including deletions and duplications, and balanced SVs that are copy number neutral including inversions and translocations[12]. Both the unbalanced SVs and balanced SVs can modify the 3D architecture of the chromosomes and potentially affect genetic functioning, which contributed to the genome evolution. For example,

Selmecki.et al showed tetraploids yeast undergo significantly faster adaptation in a poor

carbon source environment with larger deletion of chromosomes. [13]. Li.et al found

that deletion of left arm of synthetic chromosome X and duplication of chromosome

VIII (trisomy) lead to increased rapamycin resistance in yeast[14]. With the

development of genome medicine, it is indicated reciprocal translocation between

chromosomes have contributed to several human diseases, for instance, chronic

myeloid leukemia (chromosomes 9 with 22) and Mobius-like syndrome (chromosomes

1 with 2)[15] [16]. In addition, pericentric inversion of chromosome 9 is one of the

most common chromosomal abnormalities and can be associated with ambiguous

genitalia in children[17]. It is challenge to map the balanced SVs than unbalanced SVs

due to the copy number neutral. So far, the detailed molecular mechanism underlying

balanced SVs remained unexplored due to lack of cell models.

The baker yeast (Saccharomyces cere*visiae*) appreciated for the availability of its

powerful genetic tools, biosafety and fast growth rate has been used extensively in

synthetic biology for industrial purposes. High-throughput phenotypic screenings have

been employed to identify genome-wide deletions and duplications that increase the

yield of yeast synthetic biology products. The completion of the yeast gene-knockout

collection (YKOC) has enabled the analysis of synthetic genetic interactions[18]. The

construction of segmental duplications covering the whole *S. cerevisiae* genome has

been used to produce strains with enhanced tolerance to stresses[19]. Many rational

design strategies, including overexpression and downregulation, are time-consuming

because identifying gene targets remains an challenge due to the complexity of genetic

networks [20][Rapid prototyping of microbial cell factories via genome-scale engineering. Biotechnol Adv 2015, 33:1420-1432.]. Many successful examples demonstrate that the synthetic yeast chromosome could be reprogrammed for various purposes with greatly enhanced metabolic capacity[21-23]. The Synthetic Chromosome Rearrangement and Modification by LoxP-mediated Evolution (SCRaMbLE) technology have been developed to generate larger-scale SVs[14, 21-29]. SCRaMbLE is a powerful approach to study how chromosomal architecture impacts phenotypes and can be to mimic the Darwinian evolution process in the test tube for targeted phenotypes. Thus, SCRaMbLE can be viewed as natural molecular tools for directed genome evolution by iterative rounds of mutagenesis and screening or selection on the genome scale. So far, SCaMbLE has been used for directed genome evolution mainly with signal synthetic chromosome.

In this work we report that one by using SCaMbLE and two synthetic chromosomes, can for the first time creates chromosome inversion and translocation events on a larger scale and at a higher frequency. Introducing the same yeast cell with two synthetic chromosomes, each containing hundreds of non-directional loxPsym sites positioned downstream of non-essential genes and major landmarks provides a novel approach to genome reorganization especially SVs. We evaluate the effects of SVs on the enhanced production of an exogenous but commercially valuable terpenoid, astaxanthin. Two cycles of SCRaMbLE were used to iteratively generate complex SVs and improve the astaxanthin yield. We used the high-resolution chromosome conformational capture (Hi-C) sequencing approach to investigate the 3D conformations of chromosomes with

complex SVs[30], and observed an inversion spanning the centromere and a

translocation between synV and synX. We developed a method to precisely reconstitute

the translocations and inversions, and recaptured their combined effects on increasing

the yield of astaxanthin by 2.7-fold. Our work thus provides a new toolkit for directed

evolution of genomic assembly and reorganization.

Results

SCRaMbLE with two synthetic chromosomes for directed genome evolution

To investigate more complex SVs and biosynthesis pathways, we used the haploid

strain SYN510 containing both synV with 176 loxPsym sites[31] and synX with 245

loxPsym sites[32] for the biosynthesis of astaxanthin. Astaxanthin is the end metabolite

of the MVA pathway; it has tremendous antioxidant activity and is widely used in

nutraceuticals, cosmetics and aquaculture[33, 34]. The astaxanthin biosynthesis

pathway consists of 5 genes. *CrtE*, *CrtI* and *CrtYB* are involved in carotenoid

biosynthesis, after which a reticular metabolic pathway with β-carotene ketolase (CrtW)

and β-carotene hydroxylase (CrtZ) performs a two-step hydroxylation and ketolation

reaction(Supplementary information, Fig. 1a). Our previous study demonstrated that

the combination of Aa_CrtZ–BDC263_CrtW showed lower intermediate accumulation

and achieved better astaxanthin yield[35]. To increase the stability of the astaxanthin

biosynthesis pathway, all five genes were assembled together and integrated at the

YEL063C/*CAN1* locus of synV to generate the yeast strain YJJ001 (Fig. 1a). It has been

reported that the pCRE4 plasmid (pGal1-Cre-EBD-tCYC1) has tight control of Cre

recombinase and can be used for Multiplex SCRaMbLE Iterative Cycling (MuSIC).

Thus, the pCRE4 plasmid with a Ura3 marker was transformed into YJJ001 for the

control of SCRaMbLE.

In the first cycle of SCRaMbLE, strain YJJ168 was selected from the SCRaMbLEd

yeast pool due to its dark red pigmentation (Supplementary information, Fig. 1b). To

generate more complicated SVs and the desired phenotype, we used YJJ168 for the

second cycle of SCRaMbLE to yield another SCRaMbLEd strain, YJJ432. To identify

the chromosome SVs in haploids caused by SCRaMbLE, we deep sequenced YJJ168

and YJJ432. As shown in Fig. 1a and Supplementary information, Fig. 2, a 3 kb deletion

(YJL218W-YJL217W), a 1 kb inversion (YER044C) and a 210 kb pericentric inversion

(YJL052C-A-YJR071C) with 127 genes were observed in the SCRaMbLEd strain

YJJ168. It is interesting that the upstream and downstream of inversions of YJL052C-

A-YJR071C were on the left arm and right arm of synX, respectively. In other words,

this inversion spanned the centromere of synX. As an iterated SCRaMbLEd strain from

YJJ168, YJJ432 was observed to contain not only all the same SVs as YJJ168 but also

a inversion of YJL170C-YJL158C and a trans-chromosomictranslocation. It is noted

that the trans-chromosomictranslocation caused an exchange of 59 kb from the right

arm of synV and 74 kb from the right arm of synX that had not previously been reported.

From the sequencing data above, we find that the second-generation strains can inherit

genotypic SVs from first-cycle SCRaMbLEd strains. PCR verification of novel

junctions also confirmed that inversion and translocation events occurred in YJJ168 and

YJJ432 (Supplementary information, Fig. 3).

To investigate the structural organization of the chromosome SVs, Hi-C contact maps

(Fig. 1b) and 3D representations (Fig. 1c) of synV and synX were generated from

YJJ001, YJJ168 and YJJ432. The contact maps of synV and synX in YJJ168 and

YJJ432 showed several new interactions, indicating large-scale rearrangements in both

SCRaMbLEd strains. YJJ168 exhibited a new interaction indicating a large inversion

(YJL052C-A-YJR071C) event. The contact map of YJJ432 revealed three new

interactions caused by the large inversion (YJL052C-A-YJR071C), one inversion

(YJL170C-YJL158C) in synX and one translocation between the right arm of synV

(YJR130C) and the right arm of synX (YER164W). A three-dimensional model of the

16 yeast chromosomes and amplified chromosomes of synV and synX are provided in

Fig. 1c. The centromeres gather at the spindle pole, and all chromosomes are clustered

by the centromeres at one pole of the nucleus. The telomeres are clustered at the other

end. The 3D representations show that the average trajectories of chromosomes in

YJJ168 and YJJ432 did not appear to be substantively altered compared with those in

YJJ001, with the synthetic chromosomes neighboring the same chromosomes as the

native ones. New connections were generated in synX of strain YJJ168 and strain

YJJ432 due to the inversion spanning the centromere. In strain YJJ432, the right arms

of synV and synX generate two new connections because of the translocation between

synV and synX (the area within the dashed oval). It is indicated that the translocation

on the right arm of synV has a closer spatial distance with synX and consequently has

stronger interactions.

Meanwhile, the two SCRaMbLEd strains and YJJ001 were subjected to fermentation

experiments in YPD medium, and the astaxanthin yield was analyzed by high-

performance liquid chromatography (HPLC). As illustrated in Fig. 1d, the astaxanthin

yield in strain YJJ001 was 0.61 mg/g DCW, and the astaxanthin yield of the two

SCRaMbLEd strains (YJJ168 and YJJ432) was increased, to 0.90 and 2.7 mg/g DCW,

respectively. The astaxanthin yield was increased 1.48- to 4.43-fold in the SCRaMbLEd

strains compared with the nonSCRaMbLEd parent strain. These results revealed the

new interactions of synthetic chromosomes in SCRaMbLEd strains via Hi-C

technology. SCRaMbLE strains provide sufficiently detectable SVs, indicating that the

SCRaMbLE system provides a driving force to alter cell phenotypes through

rearranging gene order and number.

Reorganization of the genetic information on the regions of SVs

The pericentric inversion and trans-chromosomic translocation alter the spatial

positions of the breakpoint-flanking regions, which may which may led to large

reorganization of the genetic information along the chromosomes. To explore whether

the genetic information in the breakpoint-flanking regions were affected, we examined

transcription profile on the 20 kb regions upstream and downstream of the two loxPsym

breakpoints. As shown in Fig. 2a, the upstream and downstream regions of the

breakpoints of the pericentric inversion in YJJ168 were named A, B, C, and D,

respectively. We first observed changes in the transcription levels of YJL053W,

YJL052C-A and YJL052W near the left breakpoint and the transcription levels of

YJL067C, YJL070C and YJL071W near the right breakpoint. In the whole 80 kb region

consisting of A, B, C and D, 19 of 51 genes showed changes in expression

(Log2FoldChange > 1-fold) compared with the control strain YJJ001, with 1 and 18

gene downregulated and upregulated, respectively. Around the left breakpoint, the

changes in YJL052C-A and YJL052W were the most obvious, whereas around the right

arm recombination site, the change in YJR070C was the most obvious. Similarly, the

regions upstream and downstream of the breakpoint in the right arm of synV and the

breakpoint in the right arm of synX in YJJ432 were labeled E, F, G, and H, respectively

(Fig. 2b). Compared with the control strain, 15 of the 38 genes in these regions

exhibited changed expression (Log2FoldChange > 1-fold), with 2 and 13 genes

downregulated and upregulated, respectively. The results indicated that the expression

of genes in the breakpoint-flanking regions was indeed affected by different types of

SVs. To further quantify the interactions of the recombination sites, we counted the

interaction values of the four points on the interaction matrix of YJJ168 and YJJ432

and counted the differences between different interaction domains (areas marked by

boxes in Supplementary information, Fig. 4). As the results show, regions A and C and

regions B and D had significant interactions in inversion strains YJJ168 and YJJ432. In

translocation strain YJJ432, the interactions of regions E and H and F and G were

increased as a result of the greater proximity of the right arm of synV (YJR130C) to

synX and the right arm of synX (YER164W) to synV. These results further indicate that

two sites that are closer in space have greater interaction. It is spectacular that the

pericentric inversions and trans-chromosomes translocation changed larger number of

genes position and the transcription direction, which affected the transcription levels of

genes in the regions nearby the breakpoint of SVs[36].

Mapping of genetic targets contributing to higher astaxanthin yield

To identify the genetic targets with regard to the improvement of astaxanthin yield, the

above down-regulated genes and up-regulated genes in the regions nearby the

breakpoint of SVs were knocked out and overexpressed in the strain YJJ001 for

fermentation assay, respectively. As illustrated in Fig. 3a, compared to the control strain,

astaxanthin yields were significantly increased in strains with YJR086W, YJR123W

and YJR135C over-expression strain. The function of these three genes are briefly

described in Table 2. Although the exact reason for these genes to be beneficial to

astaxanthin yield remains to be elucidated, it is possible to speculate the mechanism

resulting in the favorable changes of this phenotype from functions previously

described in the literature. The result indicated upregulated these genes have

contributed to higher astaxanthin yield.

To further explore the global affection of chromosomal rearrangements on metabolic

pathways, enrichment analysis of transcriptomics was used to identify differentially

transcribed genes with known functions involved in KEGG pathways. Several genes

involved in terpenoid backbone biosynthesis, ergosterol biosynthesis and citrate cycle

were upregulated in the YJJ168 and YJJ432 (Fig. 3b). In SCRaMbLEd strain YJJ168,

the upregulated genes ERG24, ERG25, ERG26 and ERG27 were involved in ergosterol

biosynthesis, while ERG10, ERG20 and BTS1 were involved in terpenoid backbone

biosynthesis. In YJJ432, ERG24, ERG26 and ERG27, which are involved in the

biosynthesis of ergosterol, and ERG10, ERG12, ERG20 and BTS1, which are involved

in terpenoid backbone biosynthesis, were found to be upregulated. As a consequence,

these two pathways could increase the flux of the MVA pathway and cell storage

capacity of hydrophobic carotenoids[37, 38]. The improvement of citrate cycle could

provide sufficient reducing power and energy for the MVA pathway. These

transcriptome analyses indicated that the improved yield of astaxanthin was caused by

the upregulation of these genes, further clarifying the direct effect of the SVs on

astaxanthin accumulation. The results demonstrated the SVs can led to large

reorganization of the genetic information along the chromosomes and accelerate the

directed genome evolution with selection of targeted phenotypes.

Rational introduction of inversion and translocation in yeast

To verify the effect of the pericentric inversion and trans-chromosomic translocation,

we developed a method to rationally generate these inversions and translocations. First,

we constructed an astaxanthin-producing strain named YJJ468 from the wild-type

strain BY4741 by integrating the biosynthesis pathway of astaxanthin at the

YEL063C/*CAN1* locus of chromosome V. Then, we designed an on/off switch

consisting of two markers (URA3 and Hyg). As shown in Fig. 4a, the URA3 promoter

was inserted upstream of loxP site 1, and the open reading frame of URA3 was inserted

downstream of loxP site 1, allowing the expression of URA3. Hyg without a promoter

was positioned immediately downstream of loxP site 2. A complete HIS3 marker was

positioned upstream of loxP site 2 as a selection marker to help loxP-Hyg integrate into

the chromosome. The pGal1-Cre-EBD-tCYC1 plasmid with a G418 marker was

transformed into strain YJJ468 (Supplementary Fig. 7). The activation of Cre

recombinase will catalyze the recombination of the two loxP sites, leading to a change

in the phenotype of the yeast from Hyg- to Hyg+. Consequently, hygromycin media

could be used to select translocation and inversion strains. Using the above approach,

strain YJJ473 with pericentric inversion and strain YJJ474 with trans-chromosomic

translocation were generated and isolated by screening on SC-hygromycin agar plates.

Strains YJJ473 and YJJ474 were further analyzed using whole-genome sequencing to

verify the genomic variations produced by the recombination. As Fig. 4a shows, the

inversion and translocation region had breakpoints at the loxP sites in YJJ473 and

YJJ474, indicating that the desired inversion and translocation were artificially

recreated from the wild-type strain YJJ468.

Meanwhile, PCR was used to further verify the inversion and translocation. Before

recombination, sequences in YJJ468 could be amplified by primer pairs 1/2, 3/4, 5/6,

and 7/8. After recombination, sequences in YJJ473 could be amplified by primer pairs

1/3 and 2/4, while sequences in YJJ474 could be amplified by primer pairs 5/8 and 6/7

(Fig. 4b). Therefore, we can confirm that inversion and translocation events occurred

in YJJ473 and YJJ474. Compared to the initial strain yJJ468, YJJ473 and YJJ474 had

enhanced astaxanthin production phenotypes, yielding 1.11 mg/g DCW and 1.28 mg/g

DCW, respectively, which were similar to those of the SCRaMbLEd strains (Fig. 4c).

YJJ473 and YJJ474 increased the astaxanthin yield 1.8- and 2.1-fold relative to the parent strain YJJ468. These results indicated that the YJL052C-A-YJR071C inversion and YJR130C-YER164W translocation increased the production of astaxanthin and were responsible for the enhanced astaxanthin production phenotypes of SCRaMbLE strains yJJ168 and yJJ432. The results indicated that the YJL052C-A-YJR071C inversion and YJR130C-YER164W translocation contributed to the improvement of astaxanthin production. It is anticipated that we can used the model learned from the random screen of SCRaMbLE and successfully harnessed the precise introduction of pericentric inversions and trans-chromosomic translocation by rational design.

Discussion

Our study focused on investigating complex chromosome SVs by enabling haploid yeast containing multiple synthetic chromosomes to be SCRaMbLEd, which has significant advantages over pre-existing methods for laboratory evolution. First, traditional evolution approaches, such as physical mutagenesis or chemical mutagenesis, typically cause hundreds of single nucleotide polymorphisms (SNPs) or insertions and deletions (indels)[39], which make it difficult to confirm the targets responsible for observed changes in phenotypes. In comparison, SCRaMbLE has been demonstrated to rapidly generate larger scale SVs, providing clear targets to verify and understand more deeply. In this study, we have demonstrated that pathway-doubling is not the only way to increase yield [17], changes in map position and direction of endogenous genes can also affect yield. Second, the synthetic yeast project (Sc2.0) aims

to synthesize 16 yeast chromosomes, each of which contains hundreds of loxPsym sites,

downstream of every nonessential gene. Our research has shown that SCRaMbLE with

two synthetic chromosomes can generate translocations between two synthetic

chromosomes. Thus, it is anticipated that the integration of multiple synthetic

chromosomes will extend the diversity and quantity of SVs. Moreover, SCRaMbLE

with multiple synthetic chromosomes provided a powerful model in which to

investigate the trajectory of chromosome SVs during iterative evolution. In this study,

YJJ432 not only inherited the inversion spanning the centromere from YJJ168 but also

acquired new genetic changes, namely, a translocation between two synthetic

chromosomes.

The SCRaMbLE system provided sufficiently large structural variations to allow us to

analyze novel interactions and structures via Hi-C technology. The translocation strain

YJJ474 on the right arm of synV (YJR130C) has a closer spatial distance with synX

and consequently has stronger interactions. Chromosomal SVs cause changes in the 3D

architecture of the genome and potentially alter cellular functions. The development of

the 3C technique and its derivatives has provided a toolbox that enables the systematic

spatial interrogation of multiple loci or even the entire genome[40, 41]. Based on

genome-wide high-throughput chromosome conformation capture (Hi-C) technology,

a 2D heat map can be constructed to indicate the frequency of interaction between two

points in the genome. Depending on their positions, SVs often change the Hi-C profile

of a locus, leaving specific signatures that can be analyzed further[42, 43]. For example,

deletions can result in novel interactions between two regions that were previously separated, whereas inversions result in a characteristic 'bow tie' configuration when mapped onto a reference genome. Furthermore, owing to its proximity-ligation nature, the Hi-C is also suitable for the identification of SVs without a priori knowledge[43-46].

SCRaMbLEing two synthetic chromosomes generated random and complex SVs, which reshaped chromosomes 3D architecture. Duan.et al reported that the genome 3D structure affected genes expressions[47] It is known that telomeres and centromeres could inhibit the expression of genes nearby [48], while the autonomously replicating sequence (ARS) could increase the expression of genes nearby[49]. In the study, the pericentric inversions and trans-chromosomes translocation changed the map position of centromeres in chromosome X and caused the exchange of the right telomeres of chromosome V with chromosome X, respectively. Thus, larger number of genes position and the transcription direction, which affected the transcription levels of genes in the regions nearby the breakpoint of SVs. It is demonstrated that over-expression of YJR086W(STE18), YJR123W(RPS5) and YJR135C(MCM22) improved astaxanthin yield. Choudhury.et al show that phosphorylation of the yeast Gg subunit (Ste18) can serve as intrinsic regulators of G protein signaling and differentially activates mitogen-activated protein kinases (MAPKs) pathway [50], which potentially mediates the activation of some transcription factors and metabolic network related to the astaxanthin biosynthesis. Ghosh.et al reported Rps5 is an essential gene coding a

protein component of the small (40S) ribosomal subunit. Over-expression of Rps5

potentially impact the translation initiation of yeast S. cerevisiae [51], which may

increase the translation of some genes being beneficial to the astaxanthin biosynthesis.

Poddar.et al demonstrated MCM22 involved in chromosome segregation, mutations of

which caused a decrease in the stability of the minichromosome [52]. Over-expression

of MCM22 potentially maintained the stability of chromosome during segregation and

enhanced the cell viability during fermentation. The exact mechanism of over-

expression of STE18, RPS5 and MCM22 contributing to astaxanthin biosynthesis is

unknown, however, upregulation of these genes may be beneficial for tuning metabolic

network, the translation initiation process and stability of chromosome during

segregation, which may explain why this genes affects astaxanthin biosynthesis.

In this manuscript, we have developed both a random method and a rational method to

generate complex SVs. The random method uses SCRaMbLE in yeast with two

synthetic chromosomes to generate yeast libraries containing complex SVs for

screening desired phenotypes. The rational method enables a selection marker to switch

from transcriptionally "off" to transcriptionally "on". The rational method provides a

convenient way to determine the effects of translocation and inversion targets on strain

phenotypes. In addition, inversion and translocation strains generated from wild-type

strains can be used as materials to study open genetic questions, such as whether

inversion and translocation strains become reproductively isolated from control strains.

By this method, we validated a larger fragment translocation (right arm of synV

(YJR130C) and right arm of synX (YER164W)) and an inversion (YJL052C-A-

YJR071C) in wild-type strains and found that both the translocation and inversion

increased the yield of astaxanthin. This strategy is potentially a powerful tool for

inducing the recombination of any two positions on chromosomes. How nature evolves

with great biodiversity and nurtures organisms containing various numbers of

chromosomes is not known. Our work may spark interest and provide a new handle for

researchers to study this fundamental biological problem.

Methods

Strains and media.

All yeast strains used in this study are described in Table 1. The strain SYNVX carries a synV, and a synX is derived from BY4741 (*MATa leu2Δ0 met15Δ0 ura3Δ0 his3Δ1*). BY4741 was used as the initial strain to reconstruct the translocations and inversions caused by SCRaMbLE of the strains and verify targets of astaxanthin yield improvement.

Yeast strains were cultured in YPD medium (10 g l^{-1} yeast extract, 20 g l^{-1} peptone, and 20 g l^{-1} glucose). SGal-Ura (synthetic media lacking uracil with 20 g l^{-1} galactose) with 1 μM β-estradiol was used to induce SCRaMbLE. SC medium containing 1 g L−1 5-FOA was used to screen strains without the URA3 marker. All yeast solid media contained 20 g L−1 agar. β-Estradiol and 5-FOA were purchased from Sigma-Aldrich. *E. coli* DH5α purchased from BEIJING Biomed Co., Ltd. was used for plasmid construction and replication. *E. coli* were cultivated at 37 °C in Luria–Bertani (LB) complete medium. Kanamycin (50 μg/mL) or ampicillin (100 μg/mL) was added to the medium for selection.

Construction of plasmids and strains.

YJJ001 (astaxanthin-producing control strain) was constructed by homologous recombination in SYNVX, directed by 500-bp CAN1 and 500-bp delta site genomic sequences flanking crtE-crtI-crtYB-crtZ-crtW-LEU2. YJJ002 was constructed by transforming YJJ001 with pCRE4 (Supplementary Fig. 1, Supplementary Data 2),

followed by selection on SC-URA agar. Overexpression plasmids were constructed by

Gibson assemble. The gene knockout cassette (left homologous arm-URA3-right

homologous arm) was assembled by overlap extension PCR (OE-

PCR).Transformations were performed using the standard lithium acetate

procedure[53].

SCRaMbLE.

YJJ002 containing the inducible Cre plasmid pGal1-Cre-EBD-tCYC1 was grown

overnight in 5 mL SC-Ura media (30 °C, 250 r.p.m. shaking). Then, the cells were

harvested and washed three times with sterile water to wash out glucose, and the culture

was diluted to an OD600 of 0.6-1.0 in 3 mL SGal-Ura medium. Then, 1 μmol L^{-1} β-

estradiol was added to the cultures to induce SCRaMbLE for 6 h (30 °C, 250 r.p.m.

shaking). Cells were washed twice by centrifugation, resuspended in sterile water,

diluted 5,000-fold, and plated on SC-Ura. The plates were incubated at 30 °C for 60 h.

HPLC analysis of astaxanthin production.

SCRaMbLE strains with darker colors were selected for fermentation in shake flasks.

Three independent colonies of each strain were inoculated into 5 ml of YPD medium

and grown at 30 °C until the OD 600 ≈ 8.0 (approximately 24 h). Then, the seed culture

was transferred into 50 mL fresh YPD with 40 g l^{-1} glucose medium at an initial OD

600 of 0.1 and grown until ready to harvest.

Astaxanthin was extracted from HCl–heat treated cells with acetone according to Zhou

et al. [54] and Wang et al. [35] . Carotenoids were extracted as described below. Cells

from 2 mL culture were collected and washed with distilled water. Then, the cells were

resuspended in 1 mL of 3 M HCl, boiled for 5 min, and subsequently cooled in an ice

bath for 5 min. After that, the cell debris was washed twice with distilled water and

resuspended in 0.5 mL of acetone containing 1% (wv^{-1}) butylated hydroxytoluene.

Then, the mixture was vortexed until colorless (approximately 20 min) and incubated

at 30 °C for 10 min. This was followed by centrifugation at 12,000 rpm for 5 min. The

acetone phase containing the extracted astaxanthin was filtered through a 0.22-μm

membrane for HPLC analysis.

PCRTags analysis.

All the synV and synX PCRTags involved in this study are listed in Supplementary

Table 2. Amplification of PCRTags was performed using 7.5 μL 2 × Rapid Taq Master

Mix (Vazyme Biotech Co., Ltd), 0.4 μL each of forward and reverse primers, 1 μL

genomic DNA, and 5.7 μL ddH2O. The reaction procedure was as follows: 95 °C/1 min,

35 cycles of 95 °C/20 s, 53 °C/20 s, 72 °C/15 s and a final extension of 72 °C/5 min.

Detection of PCRTags was performed by gel electrophoresis.

Whole-genome sequencing (WGS) and analysis.

WGS was performed at BGI (Beijing Genomic Institute in Shenzhen, China), and cells

were harvested at exponential phase. Libraries were prepared and analyzed using an

Illumina HiSeq X-Ten system. The sequencing data were filtered with SOAPnuke

(v1.5.2)[55]. and clean reads were stored in FASTQ format for downstream analysis.

The read comparison was performed using BWA software with the reference sequence.

SVs including insertions, deletions, inversions, intrachromosomal translocations, and

interchromosomal translocations were detected using Break Dancer software.

Transcriptional analysis.

Yeast cells were harvested from YPD medium at 24 h (exponential phase). Total RNA

was extracted using the TRIzol® method following the NEB Next Ultra™ RNA

protocol. The concentration of the extracted RNA samples was determined using a

NanoDrop system (NanoDrop, Madison, USA), and the integrity of the RNA was

examined based on the RNA integrity number (RIN) determined using an Agilent 2100

Bioanalyzer (Agilent, Santa Clara, USA). RNA sequencing was carried out by the

BGISeq500 platform (BGI-Shenzhen, China). The sequencing data were filtered with

SOAPnuke (v1.5.2).[55]

Clean reads were obtained and stored in FASTQ format. The clean reads were mapped

to the reference genome using HISAT2 (v2.0.4)[56]. Bowtie2 (v2.2.5)[57] was applied

to align the clean reads to the reference coding gene set, and the expression level of

each gene was calculated by RSEM (v1.2.12)[58]. The heatmap of the gene expression

in different samples was drawn by GraphPad Prism (v8.0.1). Essentially, differential

expression analysis was performed using DESeq2 (v1.4.5)[59] with a Q value $\leqslant$ 0.05.

To gain insight into the change in phenotype, GO (http://www.geneontology.org/) and

KEGG (https://www.kegg.jp/) enrichment analyses of annotated differentially

expressed   genes   were   performed   by   Phyper

(https://en.wikipedia.org/wiki/Hypergeometric_distribution)   based   on   the

hypergeometric test. Significant terms and pathways were identified as those with

Bonferroni-corrected *P*-values below a rigorous threshold (P-value $\leqslant$ 0.05). Triplicate

samples were used for transcriptional analysis. The Saccharomyces Genome Database

(SGD)[60] was used to obtain gene information.

Hi-C library generation and sequencing.

Hi-C library generation and sequencing were carried out by Frasergen (Wuhan, China),

and cells were harvested at exponential phase. Cells were resuspended in 1×PBS to an

OD600 of 1.0. Cells were cross-linked in a 3% final concentration of fresh

formaldehyde and quenched with glycine (0.15 M final concentration) for 5 min. The

cells were resuspended in 1 mL of 1× NEBuffer 2.1 (NEB) and homogenized by

grinding to a fine powder in liquid nitrogen. Then, the homogenized yeast material was

washed with 25 mL of 1× NEBuffer 2.1 and suspended in 2.5 mL of 1× NEBuffer 2.1.

Cells were split into aliquots (V =456 µL) and solubilized in 0.1% SDS for 10 min at

65 °C. Cross-linked DNA was digested with 200 U MboI (NEB) per tube at 37 °C

overnight. Restriction fragment ends were labeled with biotinylated cytosine

nucleotides by biotin-14-dCTP (TriLINK). Blunt-end ligation was carried out at 16 °C

overnight in the presence of 100 Weiss units of T4 DNA ligase (Thermo, 10.0 mL final

volume per tube).

DNA purification was achieved through overnight incubation at 65 °C with 200 µg/mL

proteinase K (Thermo). Purified DNA was sheared to a length of ~400 bp. Point ligation

junctions were pulled down by Dynabeads® MyOne™ Streptavidin C1 (Thermo

Fisher). The Hi-C library for Illumina sequencing was prepared using the NEBNext®

Ultra™ II DNA library Prep Kit for Illumina (NEB) according to the manufacturer's

instructions. Fragments of between 400 and 600 bp were paired-end sequenced on an

Illumina HiSeq X10 platform (San Diego, CA, United States) in 150PEmode. Two

replicates were generated for each group of materials.

Construction of contact map.

The contact maps were generated using the ICE software package (version

1f8815d0cc9e)[61], and the Hi-C data of YJJ001, YJJ168 and YJJ432 cells were

iteratively mapped to their own genomes. Dangling ends and other unusable data were

filtered out, and the valid pairs were binned into 10 kb nonoverlapping genomic

intervals to generate contact maps. The contact maps were normalized using an iterative

normalization method to eliminate systematic biases.

Cre/loxP induced chromosome translocation and inversion.

Translocations and inversions were detected in YJJ168 and YJJ432 and verified in the

wild-type strain BY4741 by introducing two Cre/loxP sites to the corresponding

position. LoxP sites were integrated by homologous recombination. Homologous arms

upstream and downstream were amplified from the genome of BY4741. Two loxP sites

(loxP site 1 and loxP site 2) were integrated along with URA3+ and HIS3+/Hyg- as

selection markers. The URA3 promoter was inserted upstream of loxP site 1, and open reading frames of URA3 were inserted downstream of the loxP site, allowing the expression of URA3. Hyg without a promoter was positioned adjacent to downstream loxP site 2 and the HIS3 promoter, and open reading frames of HIS3 were positioned upstream of loxP site 2. Adding 1 μmol L−1 β-estradiol to the cultures to switch on Cre-mediated recombination between the two loxP sites results in translocation and inversion, which subsequently turn on the expression of the Hyg gene and simultaneously shut down URA3 expression. Translocation and inversion strains could be selected on SC-LEU-5-FOA plates and SC-LEU+Hyg.

Acknowledgements

This work was funded by Ministry of science and technology the National Key Research and Development Program of China (2021YFC2100800), and the National Natural Science Foundation of China (31800719, 31861143017 and 21621004).

Author contributions

B.J., and J.J. contributed equally to this work. B.J., J.J., and Y.J.Y. designed the experiments. B.J., J.J., and M.Z.H performed the experiments. B.J., J.J., and Y.J.Y. wrote the manuscript and all authors edited the manuscript. This project was supervised by Y.J.Y.

Conflict of Interest

The authors declare no competing financial interests.

Reference

517   1.   Dujon B, Sherman D, Fischer G, Durrens P, Casaregola S, Lafontaine I, De Montigny J,

518   Marck C, Neuvéglise C, Talla E, et al: Genome evolution in yeasts. *Nature* 2004, 430:35-

519   44.

520   2.   Redon R, Ishikawa S, Fitch KR, Feuk L, Perry GH, Andrews TD, Fiegler H, Shapero MH,

521   Carson AR, Chen WW, et al: Global variation in copy number in the human genome.

522   *Nature* 2006, 444:444-454.

523   3.   Kidd JM, Cooper GM, Donahue WF, Hayden HS, Sampas N, Graves T, Hansen N, Teague

524   B, Alkan C, Antonacci F, et al: Mapping and sequencing of structural variation from eight

525   human genomes. *Nature* 2008, 453:56-64.

526   4.   Dujon B: Yeast evolutionary genomics. *Nature Reviews Genetics* 2010, 11:512-524.

527   5.   Peter J, De Chiara M, Friedrich A, Yue JX, Pflieger D, Bergstrom A, Sigwalt A, Barre B,

528   Freel K, Llored A, et al: Genome evolution across 1,011 Saccharomyces cerevisiae

529   isolates. *Nature* 2018, 556:339-+.

530   6.   Conrad DF, Pinto D, Redon R, Feuk L, Gokcumen O, Zhang YJ, Aerts J, Andrews TD,

531   Barnes C, Campbell P, et al: Origins and functional impact of copy number variation in

532   the human genome. *Nature* 2010, 464:704-712.

533   7.   Pang AW, MacDonald JR, Pinto D, Wei J, Rafiq MA, Conrad DF, Park H, Hurles ME, Lee C,

534   Venter JC, et al: Towards a comprehensive structural variation map of an individual

535   human genome. *Genome Biology* 2010, 11.

536   8.   Darling AE, Miklos I, Ragan MA: Dynamics of Genome Rearrangement in Bacterial

537   Populations. *Plos Genetics* 2008, 4:16.

9.   Yue JX, Li J, Aigrain L, Hallin J, Persson K, Oliver K, Bergstrom A, Coupland P, Warringer J,

Lagomarsino MC, et al: Contrasting evolutionary genome dynamics between

domesticated and wild yeasts. *Nature Genetics* 2017, 49:913-+.

10.   Pevzner P, Tesler G: Genome Rearrangements in mammalian evolution: Lessons from

human and mouse genomes. *Genome Research* 2003, 13:37-45.

11.   Henssen AG, Koche R, Zhuang J, Jiang E, Reed C, Eisenberg A, Still E, MacArthur IC,

Rodriguez-Fos E, Gonzalez S, et al: PGBD5 promotes site-specific oncogenic mutations

in human tumors. *Nature Genetics* 2017, 49:1005-+.

12.   Fleiss A, O'Donnell S, Fournier T, Lu W, Agier N, Delmas S, Schacherer J, Fischer G:

Reshuffling yeast chromosomes with CRISPR/Cas9. *PLoS Genet* 2019, 15:e1008332.

13.   Selmecki AM, Maruvka YE, Richmond PA, Guillet M, Shoresh N, Sorenson AL, De S,

Kishony R, Michor F, Dowell R, Pellman D: Polyploidy can drive rapid adaptation in yeast.

Nature 2015, 519:349-352.

14.   Li Y, Wu Y, Ma L, Guo Z, Xiao W, Yuan Y: Loss of heterozygosity by SCRaMbLEing.

Science China Life Sciences 2019, 62:381-393.

15.   Eyupoglu D, Bozkurt S, Haznedaroglu I, Buyukasik Y, Guven D: The Impact of Variant

Philadelphia Chromosome Translocations on the Clinical Course of Chronic Myeloid

Leukemia. *Turkish Journal of Hematology* 2016, 33:60-65.

16.   Nishikawa M, Ichiyama T, Hayashi T, Furukawa S: Mobius-like syndrome associated with

a 1;2 chromosome translocation. *Clinical genetics* 1997, 51:122-123.

17.   Sotoudeh A, Rostami P, Nakhaeimoghadam M, Mohsenipour R, Rezaei N: Pericentric

Inversion of Chromosome 9 in an Infant With Ambiguous Genitalia. *Acta medica Iranica*

2017, 55:655-657.

18.   Puddu F, Herzog M, Selivanova A, Wang SY, Zhu J, Klein-Lavi S, Gordon M, Meirman R,

562   Millan-Zambrano G, Ayestaran I, et al: Genome architecture and stability in the

563   Saccharomyces cerevisiae knockout collection. *Nature* 2019, 573:416-+.

564   19.   Natesuntorn W, Iwami K, Matsubara Y, Sasano Y, Sugiyama M, Kaneko Y, Harashima S:

565   Genome-wide construction of a series of designed segmental aneuploids in

566   Saccharomyces cerevisiae. *Scientific Reports* 2015, 5.

567   20.   Si T, Xiao H, Zhao H: Rapid prototyping of microbial cell factories via genome-scale

568   engineering. *Biotechnol Adv* 2015, 33:1420-1432.

569   21.   Jia B, Wu Y, Li B-Z, Mitchell LA, Liu H, Pan S, Wang J, Zhang H-R, Jia N, Li B, et al: Precise

570   control of SCRaMbLE in synthetic haploid and diploid yeast. *Nature Communications*

571   2018, 9.

572   22.   Wu Y, Zhu R-Y, Mitchell LA, Ma L, Liu R, Zhao M, Jia B, Xu H, Li Y-X, Yang Z-M, et al: In

573   vitro DNA SCRaMbLE. *Nature Communications* 2018, 9.

574   23.   Shen MJ, Wu Y, Yang K, Li Y, Xu H, Zhang H, Li B-Z, Li X, Xiao W-H, Zhou X, et al:

575   Heterozygous diploid and interspecies SCRaMbLEing. *Nature Communications* 2018, 9.

576   24.   Luo Z, Wang L, Wang Y, Zhang W, Guo Y, Shen Y, Jiang L, Wu Q, Zhang C, Cai Y, Dai J:

577   Identifying and characterizing SCRaMbLEd synthetic yeast using ReSCuES. *Nature*

578   *Communications* 2018, 9.

579   25.   Blount BA, Gowers GOF, Ho JCH, Ledesma-Amaro R, Jovicevic D, McKiernan RM, Xie ZX,

580   Li BZ, Yuan YJ, Ellis T: Rapid host strain improvement by in vivo rearrangement of a

581   synthetic yeast chromosome. *Nature Communications* 2018, 9.

26.   Wang J, Jia B, Xie ZX, Li YX, Yuan YJ: Improving prodeoxyviolacein production via

multiplex SCRaMbLE iterative cycles. *Frontiers of Chemical Science and Engineering*

2018, 12:806-814.

27.   Ma L, Li Y, Chen X, Ding M, Wu Y, Yuan Y-J: SCRaMbLE generates evolved yeasts with

increased alkali tolerance. *Microbial Cell Factories* 2019, 18.

28.   Liu W, Luo Z, Wang Y, Pham NT, Tuck L, Pérez-Pi I, Liu L, Shen Y, French C, Auer M, et

al: Rapid pathway prototyping and engineering using in vitro and in vivo synthetic

genome SCRaMbLE-in methods. *Nature Communications* 2018, 9.

29.   Gowers GOF, Chee SM, Bell D, Suckling L, Kern M, Tew D, McClymont DW, Ellis T:

Improved betulinic acid biosynthesis using synthetic yeast chromosome recombination

and semi-automated rapid LC-MS screening. *Nature Communications* 2020, 11.

30.   Akdemir KC, Le VTT, Chandran S, Li YL, Verhaak RG, Beroukhim R, Campbell PJ, Chin L,

Dixon JR, Futreal PA, et al: Disruption of chromatin folding domains by somatic genomic

rearrangements in human cancer. *Nature Genetics* 2020, 52:294-+.

31.   Xie ZX, Li BZ, Mitchell LA, Wu Y, Qi X, Jin Z, Jia B, Wang X, Zeng BX, Liu HM, et al:

"Perfect" designer chromosome V and behavior of a ring derivative. *Science* 2017, 355:8.

32.   Wu Y, Li BZ, Zhao M, Mitchell LA, Xie ZX, Lin QH, Wang X, Xiao WH, Wang Y, Zhou X, et

al: Bug mapping and fitness testing of chemically synthesized chromosome X. *Science*

2017, 355:7.

33.   Ambati RR, Phang SM, Ravi S, Aswathanarayana RG: Astaxanthin: Sources, Extraction,

Stability, Biological Activities and Its Commercial Applications-A Review. *Marine Drugs*

2014, 12:128-152.

34.   Igielska-Kalwat J, Goscianska J, Nowak I: Carotenoids as natural antioxidants. *Postepy*

higieny i medycyny doswiadczalnej (Online) 2015, 69:418-428.

35.   Wang R, Gu X, Yao M, Pan C, Liu H, Xiao W, Wang Y, Yuan Y: Engineering of β-carotene

hydroxylase and ketolase for astaxanthin overproduction in Saccharomyces cerevisiae.

Frontiers of Chemical Science and Engineering 2017, 11:89-99.

36.   Wu X-L, Li B-Z, Zhang W-Z, Song K, Qi H, Dai J-b, Yuan Y-J: Genome-wide landscape

of position effects on heterogeneous gene expression in Saccharomyces cerevisiae.

Biotechnology for Biofuels 2017, 10.

37.   Wu T, Ye LJ, Zhao DD, Li SW, Li QY, Zhang BL, Bi CH, Zhang XL: Membrane engineering

- A novel strategy to enhance the production and accumulation of beta-carotene in

Escherichia coli. *Metabolic Engineering* 2017, 43:85-91.

38.   Gao SL, Tong YY, Zhu L, Ge M, Zhang YA, Chen DJ, Jiang Y, Yang S: Iterative integration

of multiple-copy pathway genes in Yarrowia lipolytica for heterologous beta-carotene

production. *Metabolic Engineering* 2017, 41:192-201.

39.   Jin J, Wang Y, Yao M, Gu X, Li B, Liu H, Ding M, Xiao W, Yuan Y: Astaxanthin

overproduction in yeast by strain engineering and new gene target uncovering.

Biotechnology for Biofuels 2018, 11.

40.   Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I,

Lajoie BR, Sabo PJ, Dorschner MO, et al: Comprehensive Mapping of Long-Range

Interactions Reveals Folding Principles of the Human Genome. *Science* 2009, 326:289-

293.

41.   Dekker J, Marti-Renom MA, Mirny LA: Exploring the three-dimensional organization of

genomes: interpreting chromatin interaction data. *Nature Reviews Genetics* 2013,

14:390-403.

42.   Bianco S, Lupianez DG, Chiariello AM, Annunziatella C, Kraft K, Schopflin R, Wittler L,

Andrey G, Vingron M, Pombo A, et al: Polymer physics predicts the effects of structural

variants on chromatin architecture. *Nature Genetics* 2018, 50:662-+.

43.   Burton JN, Adey A, Patwardhan RP, Qiu RL, Kitzman JO, Shendure J: Chromosome-scale

scaffolding of de novo genome assemblies based on chromatin interactions. *Nature*

Biotechnology 2013, 31:1119-+.

44.   Rickman DS, Soong TD, Moss B, Mosquera JM, Dlabal J, Terry S, MacDonald TY, Tripodi

J, Bunting K, Najfeld V, et al: Oncogene-mediated alterations in chromatin

conformation. *Proceedings of the National Academy of Sciences of the United States of*

America 2012, 109:9083-9088.

45.   Harewood L, Kishore K, Eldridge MD, Wingett S, Pearson D, Schoenfelder S, Collins VP,

Fraser P: Hi-C as a tool for precise detection and characterisation of chromosomal

rearrangements and copy number variation in human tumours. *Genome Biology* 2017,

18:11.

46.   Rao SSP, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn AL,

Machol I, Omer AD, Lander ES, Aiden EL: A 3D Map of the Human Genome at Kilobase

Resolution Reveals Principles of Chromatin Looping. *Cell* 2014, 159:1665-1680.

47.   Duan Z, Andronescu M, Schutz K, McIlwain S, Kim YJ, Lee C, Shendure J, Fields S, Blau

CA, Noble WS: A three-dimensional model of the yeast genome. *Nature* 2010, 465:363-

367.

48.   Chen X, Zhang J: The Genomic Landscape of Position Effects on Protein Expression Level

and Noise in Yeast. *Cell Syst* 2016, 2:347-354.

49.   Flagfeldt DB, Siewers V, Huang L, Nielsen J: Characterization of chromosomal integration

sites for heterologous gene expression in Saccharomyces cerevisiae. *Yeast* 2009, 26:545-

551.

50.   Choudhury S, Baradaran-Mashinchi P, Torres MP: Negative Feedback Phosphorylation

of Ggamma Subunit Ste18 and the Ste5 Scaffold Synergistically Regulates MAPK

Activation in Yeast. *Cell Rep* 2018, 23:1504-1515.

51.   Saliba AE, Westermann AJ, Gorski SA, Vogel J: Single-cell RNA-seq: advances and future

challenges. *Nucleic Acids Res* 2014, 42:8845-8860.

52.   MCM21and MCM22, two novel genes of the yeast Saccharomyces cerevisiae are

required for chromosome transmission.

53.   Gietz RD, Schiestl RH, Willems AR, Woods RA: Studies on the transformation of intact

yeast cells by the LiAc/SS-DNA/PEG procedure. *Yeast (Chichester, England)* 1995,

11:355-360.

54.   Zhou PP, Ye LD, Xie WP, Lv XM, Yu HW: Highly efficient biosynthesis of astaxanthin in

Saccharomyces cerevisiae by integration and tuning of algal crtZ and bkt. *Applied*

Microbiology and Biotechnology 2015, 99:8419-8428.

55.   Li RQ, Li YR, Kristiansen K, Wang J: SOAP: short oligonucleotide alignment program.

Bioinformatics 2008, 24:713-714.

56.   Kim D, Landmead B, Salzberg SL: HISAT: a fast spliced aligner with low memory

requirements. *Nature Methods* 2015, 12:357-U121.

57.   Langmead B, Salzberg SL: Fast gapped-read alignment with Bowtie 2. *Nature Methods*

671   2012, 9:357-U354.

672 58. Li B, Dewey CN: RSEM: accurate transcript quantification from RNA-Seq data with or

673   without a reference genome. *Bmc Bioinformatics* 2011, 12:16.

674 59. Love MI, Huber W, Anders S: Moderated estimation of fold change and dispersion for

675   RNA-seq data with DESeq2. *Genome Biology* 2014, 15:38.

676 60. Cherry JM, Hong EL, Amundsen C, Balakrishnan R, Binkley G, Chan ET, Christie KR,

677   Costanzo MC, Dwight SS, Engel SR, et al: Saccharomyces Genome Database: the

678   genomics resource of budding yeast. *Nucleic Acids Research* 2012, 40:D700-D705.

679 61. Imakaev M, Fudenberg G, McCord RP, Naumova N, Goloborodko A, Lajoie BR, Dekker J,

680   Mirny LA: Iterative correction of Hi-C data reveals hallmarks of chromosome

681   organization. *Nature Methods* 2012, 9:999-+.

682 62. Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P, Boeke JD: Designer deletion

683   strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and

684   plasmids for PCR-mediated gene disruption and other applications. *Yeast* 1998, 14:115-

685   132.

Table 1 *S. cerevisiae* **strains used in this study**

Strains	Description	Sources
BY4741	*MATa, HIS3Δ1, LEU2Δ0, MET15Δ0, URA3Δ0*	[62]
SYNVX	*MATa, HIS3Δ1, LEU2Δ0, MET15Δ0, URA3Δ0*	This study
YJJ001	yXZX573, CAN1:: astaxanthin pathway with Leu2 marker	This study
YJJ002	Introducing plasmid pGAL1-Cre-EBD-GFP-tCYC1 into strain YJJ001	This study
YJJ168	SCRaMbLEd strain from the YJJ002	This study
YJJ432	SCRaMbLEd strain from the YJJ168	This study
YJJ468	Astaxanthin producing strain, BY4741	This study
YJJ473	YJJ468 with inversion (YJL052C-A-YJR071C) strain	This study
YJJ474	YJJ468 with translocation (YJR130C and YER164W) strain	This study
YJJ522	YJJ001 with YJL053Woverexpression	This study
YJJ523	YJJ001 with YJL052C-A overexpression	This study
YJJ524	YJJ001 with YJL052W deletion	This study
YJJ525	YJJ001 with YJL049W overexpression	This study
YJJ526	YJJ001 with YJL046W overexpression	This study
YJJ527	YJJ001 with YJL045W overexpression	This study
YJJ528	YJJ001 with YJR067C overexpression	This study
YJJ529	YJJ001 with YJR068W overexpression	This study
YJJ530	YJJ001 with YJR070C overexpression	This study
YJJ531	YJJ001 with YJR071W overexpression	This study
YJJ532	YJJ001 with YJR072C overexpression	This study
YJJ533	YJJ001 with YJR076C overexpression	This study
YJJ534	YJJ001 with YJR079W overexpression	This study
YJJ535	YJJ001 with YJR080C overexpression	This study
YJJ536	YJJ001 with YJR082C overexpression	This study
YJJ537	YJJ001 with YJR085C overexpression	This study
YJJ538	YJJ001 with YJR086W overexpression	This study
YJJ539	YJJ001 with YJR087W overexpression	This study
YJJ540	YJJ001 with YJR088C overexpression	This study
YJJ541	YJJ001 with YER156C overexpression	This study
YJJ542	YJJ001 with YER158C deletion	This study
YJJ543	YJJ001 with YER163C overexpression	This study
YJJ544	YJJ001 with YER168C overexpression	This study
YJJ545	YJJ001 with YER170W overexpression	This study
YJJ546	YJJ001 with YJR122W overexpression	This study
YJJ547	YJJ001 with YJR123W overexpression	This study
YJJ548	YJJ001 with YJR127 deletion	This study
YJJ549	YJJ001 with YJR125C overexpression	This study
YJJ551	YJJ001 with YJR133W overexpression	This study
YJJ552	YJJ001 with YJR135C overexpression	This study
YJJ553	YJJ001 with YJR135W-A overexpression	This study
YJJ554	YJJ001 with YJR136C overexpression	This study

690   **Table 2** Known gene functions for beneficial over-expression

Gene	Description
YJR086W(STE18)	G protein gamma subunit; forms a dimer with Ste4p to activate the mating signaling pathway, forms a heterotrimer with Gpa1p and Ste4p to dampen signaling; C-terminus is palmitoylated and farnesylated, which are required for normal signaling
YJR123W(RPS5)	Protein component of the small (40S) ribosomal subunit; least basic of non-acidic ribosomal proteins; phosphorylated in vivo; essential for viability; homologous to mammalian ribosomal protein S5 and bacterial S7
YJR135C(MCM22)	Outer kinetochore protein and component of the Ctf3 subcomplex; binds to centromeric DNA in a Ctf19p-dependent manner; involved in chromosome segregation and minichromosome maintenance; orthologous to human centromere constitutive-associated network (CCAN) subunit CENP-K and fission yeast sim4

Fig. 1 SCRaMbLE with two synthetic chromosomes generating complex SVs. a,

Schematic representation of structural variations in YJJ001, YJJ168 and YJJ432.

Synthetic chromosome V and synthetic chromosome X are abbreviated as synV

and synX, respectively. b, Normalized contact maps (bin size, 2 kb) of synthetic

chromosomes in three different strains: YJJ117, YJJ168 and YJJ432. All Hi-C

reads are mapped against the reference genome of the parental strain YJJ001. c,

Three-dimensional model of the YJJ001, YJJ168 and YJJ432 genomes. Synthetic

chromosomes and wild chromosomes are indicated with graduated color and gray,

respectively. The SV area are indicated with the dashed oval. d, The astaxanthin

yield of control strain YJJ001 and SCRaMbLEd strains YJJ168 and YJJ432.

Fig. 2 Quantified interactions of loxP breakpoint-flanking regions. a, Triangle heat maps representing the contact frequencies among 20 bin regions (length 20 kb) upstream and downstream of two recombination sites (YJL052C-A and YJR071C) in YJJ168, labeled A, B, C, and D. Dots show fold changes in expression in the A, B, C, and D regions. b, Triangle heat maps representing the contact frequencies among 20 bin regions (length 20 kb) upstream and downstream of the two recombination sites (YER164W and YJR130C) in YJJ432, labeled E, F, G, and H. Dots show fold changes in expression in the E, F, G, H regions.

Fig. 3 a, The astaxanthin yield of control strain YJJ001 and genes overexpression

and deletion around pericentric inversion strain breakpoint-flanking regions and

genes overexpression and deletion around trans-chromosomic translocation strain

breakpoint-flanking regions. (Student's t-test; *P < 0.05). b, Enrichment analysis

of transcriptomics of YJJ168 and YJJ432 were used to identify differentially

transcribed genes with known functions involved in KEGG pathways.

Fig. 4 Rational translocation and inversion of chromosomes. a, The process of inversion and translocation was validated in wild-type strains. Wild-type chromosome V and wild-type chromosome X are abbreviated as wt V and wt X, respectively. Validation strains exhibited inversion and translocation events between two LoxP sites. These events were identified by aligning all raw reads. b, Verification of inversion and translocation by junction PCR. c, The astaxanthin yield of the wild-type control strain YJJ468, inversion strain YJJ473 and translocation strain YJJ474. Serial dilution assay of YJJ468, YJJ473, YJJ474.

a

synV
synX

SCRaMbLE   Deletion
LoxPsym   Inversion
Astaxanthin Translocation
pathway

YJJ001   YJJ168   YJJ432

b

YJJ001   YJJ168   YJJ432

200 kbp

c

synV
synX
Wild-type
chromosomes

Telomere

YJJ001   YJJ168   YJJ432

d

Astaxanthin yield (mg/g DCW)

YJJ001   YJJ168   YJJ432
Strains

a

Inversion

Translocation

Chromosome
337,886 570,806 wtX
509,801 wtV 665,190 wtX

Before
P1 *pTEF* *URA3* P2 P3 *LoxP* *pHIS* P4
LoxP *Hyg* *HIS3*

P5 *pTEF* *URA3* P6 P7 *pHIS* P8
LoxP *HIS3* *LoxP* *Hyg*

Recombination

After
P1 *pTEF* *Hyg* P3 P2 *LoxP* *pHIS* P4
LoxP *URA3* *HIS3*

P5 *pTEF* *Hyg* P8 P7 *pHIS* P6
LoxP *HIS3* *LoxP* *URA3*

Sequencing Coverage
LoxP *LoxP* *LoxP* *LoxP*

Reads alignment

b

YJJ468

YJJ473 YJJ474

YJJ473
P1 P3 P1 P2
P2 P4 P3 P4
8 kbp
2 kbp
750 bp
200 bp

YJJ474
P5 P7 P5 P7
P6 P8 P8 P6
8 kbp
2 kbp
750 bp
200 bp

c

Astaxanthin yield (mg/g DCW)

1.5

1.0

0.5

0.0

YJJ468 YJJ473 YJJ474

Strains