Optogenetic actuator/ERK biosensor circuits identify MAPK network nodes that shape ERK dynamics

Coralie Dessauges[1], Jan Mikelson[2], Maciej Dobrzyński[1], Marc-Antoine Jacques[1], Agne Frismantiene[1], Paolo Armando Gagliardi[1], Mustafa Khammash[2], Olivier Pertz[1]

[1]Institute of Cell Biology, University of Bern, Baltzerstrasse 4, 3012 Bern, Switzerland
[2]Department of Biosystems Science and Engineering, ETH Zurich, Mattenstrasse 26, 4058 Basel, Switzerland

Key words

ERK dynamics, MAPK network, signaling robustness, optogenetics, single cell biology

Abstract

Combining single-cell measurements of ERK activity dynamics with perturbations provides insights into the MAPK network topology. We built circuits consisting of an optogenetic actuator to activate MAPK signaling and an ERK biosensor to measure single-cell ERK dynamics. This allowed us to conduct RNAi screens to investigate the role of 50 MAPK proteins in ERK dynamics. We found that the MAPK network is robust against most node perturbations. We observed that the ERK-RAF and the ERK-RSK2-SOS negative feedbacks operate simultaneously to regulate ERK dynamics. Bypassing the RSK2-mediated feedback, either by direct optogenetic activation of RAS, or by RSK2 perturbation, sensitized ERK dynamics to further perturbations. Similarly, targeting this feedback in a human ErbB2-dependent oncogenic signaling model increased the efficiency of a MEK inhibitor. The RSK2-mediated feedback is thus important for the ability of the MAPK network to produce consistent ERK outputs and its perturbation can enhance the efficiency of MAPK inhibitors.

Introduction

The extracellular signal-regulated kinase (ERK) is part of the mitogen-activated protein kinase (MAPK) signaling network and regulates a large variety of fate decisions. While ERK can be activated by several extracellular inputs, ERK signaling has mostly been studied in the context of receptor tyrosine kinases (RTKs). Upon binding of their cognate growth factors (GFs), RTKs activate a complex signaling cascade with the following hierarchy: (1) recruitment of adaptor molecules such as GRB2 (Schlessinger 2000), (2) activation of RAS GTPases through Guanine nucleotide exchange factors (GEFs) and GTPase activating proteins (GAPs) (Cherfils and Zeghouf 2013), (3) triggering of a tripartite RAF, MEK, ERK kinase cascade that is further regulated by a variety of binding proteins (Lavoie et al. 2020), (4) ERK-mediated phosphorylation of a large number of substrates. Due to its central role in fate decisions, MAPK network dysregulation is causative for a large number of diseases including cancer (Rauen 2013; Samatar and Poulikakos 2014).

As for other signaling pathways (Purvis and Lahav 2013), temporal patterns of ERK activity, hereafter referred to as ERK dynamics, rather than steady states control fate decisions (Santos et al. 2007; Avraham and Yarden 2011; Ryu et al. 2015; Albeck et al. 2013). These specific ERK dynamics have been shown to arise from feedbacks in the MAPK network. For example, a negative feedback (NFB) from ERK to RAF can produce adaptive or oscillatory ERK dynamics (Santos et al. 2007; Kholodenko et al. 2010; Avraham and Yarden 2011). The ERK-RAF NFB was also shown to buffer against MAPK node perturbations (Sturm et al. 2010; Fritsche-Guenther et al. 2011). This property might allow cells to produce consistent ERK outputs despite heterogeneous node expression (Blüthgen and Legewie 2013). In this work, we specifically refer to the ability of the MAPK network to produce consistent ERK dynamics in presence of node perturbations as signaling robustness. While several NFBs have been mapped experimentally in the MAPK network (Lake et al. 2016), their contribution to this signaling robustness and shaping ERK dynamics remains largely unknown.

Single-cell biosensor imaging has provided new insights into MAPK signaling that were not accessible with biochemical, population-averaged measurements. This showed that the MAPK network can produce a wide variety of ERK dynamics such as transient (Ryu et al. 2015), pulsatile (Albeck et al. 2013), oscillatory (Shankaran et al. 2009) and sustained dynamics (Ryu et al. 2015; Blum et al. 2019). Mathematical modeling has provided insights into the network's structures that decode different signaling inputs into specific ERK dynamics (Santos et al. 2007; Shankaran et al. 2009; Nakakuki et al. 2010; Ryu et al. 2015). Combined modeling/experimental approaches helped to shed light on various subparts of the MAPK network, including the epidermal growth factor receptor (EGFR) module (Koseska and Bastiaens 2020), the RAS module (Schmick et al. 2015; Erickson et al. 2019), and the tripartite RAF/MEK/ERK cascade (Ferrell and Bhatt 1997; Kholodenko 2000; Orton et al. 2005; Santos et al. 2007; Ryu et al. 2015; Kochańczyk et al. 2017; Arkun and Yasemi 2018). However, the low experimental throughput to measure ERK dynamics, or other MAPK

network nodes, has precluded a global understanding of the specific functions of the
nodes present in the network.

Here, we built multiple genetic circuits consisting of optogenetic actuators together
with an ERK biosensor to simultaneously activate ERK from different nodes in the
MAPK network and report single-cell ERK dynamics. These circuits allowed us to
investigate the role of 50 MAPK signaling nodes in ERK dynamics regulations with
RNA interference (RNAi). We observed that most perturbations of individual nodes
resulted in mild ERK dynamics phenotypes despite targeting major MAPK signaling
nodes. Further, the ERK dynamics induced by various perturbations suggest that two
NFBs (ERK-RAF and ERK-RSK2-SOS) act simultaneously to regulate ERK dynamics.
Targeting the RSK2-mediated NFB increased the efficiency of additional MAPK
network perturbations both in our optogenetic systems and in an ErbB2-driven
oncogenic ERK signaling model. This suggests that the RSK2-mediated feedback
plays a role in MAPK signaling robustness and can be targeted for potent inhibition of
oncogenic ERK signaling.

Results

An optogenetic actuator-biosensor genetic circuit to study input-dependent ERK dynamics

In order to measure ERK dynamics in response to dynamic RTK input, we built a genetically-encoded circuit made of an optogenetic RTK actuator and an ERK biosensor (Figure 1A). We chose optoFGFR, which consists of a myristoylated intracellular domain of the fibroblast growth factor receptor 1 (FGFR1) fused to a CRY2 domain and tagged with mCitrine (Kim et al. 2014). Upon stimulation with blue light, optoFGFR dimerizes and trans-autophosphorylates, leading to the activation of the MAPK/ERK, phosphoinositide 3-kinase (PI3K)/AKT, and phospholipase C (PLC)/Ca^{2+} pathways. As ERK biosensor, we used ERK-KTR-mRuby2 that is spectrally compatible with optoFGFR. ERK-KTR reversibly translocates from the nucleus to the cytosol upon ERK activation (Regot et al. 2014). We used a nuclear Histone 2B (H2B)-miRFP703 marker to identify and track single cells. After stably inserting these constructs into murine NIH3T3 fibroblasts, we used automated time-lapse microscopy to stimulate selected fields of view with defined blue light input patterns to activate optoFGFR. The corresponding ERK-KTR/H2B signals were recorded with a 1-minute temporal resolution. We observed that a 100 ms light pulse leads to reversible ERK-KTR translocation from the nucleus to the cytosol, indicative of transient ERK activation (Figure 1B, Appendix Movie S1). At the end of each experiment, we imaged the mCitrine signal to evaluate optoFGFR expression levels. We built a computer vision pipeline to automatically track each nucleus, compute ERK activity as the cytosolic/nuclear ratio of the ERK-KTR signals and correlate single-cell ERK responses with optoFGFR levels (Figure 1C). We then use this pipeline to evaluate the sensitivity and specificity of our system with dose response experiments using the FGFR inhibitor SU5402, the RAF inhibitor RAF709, the MEK inhibitor U0126 and the ERK inhibitor SCH772984 (Appendix Figure S1A).

To evaluate light-dependent optoFGFR activation dynamics, we engineered a mScarlet-tagged optoFGFR that is spectrally orthogonal to CRY2 absorption (Appendix Figure S1B). Total internal reflection (TIRF) microscopy visualized the formation of optoFGFR clusters in response to blue light-mediated dimerization in the plasma membrane (Appendix Figure S1B, blue arrows, Appendix Movie S2). Consistently with CRY2's dissociation half-life (Duan et al. 2017), these optoFGFR clusters appeared within 20 seconds after a blue light pulse and disappeared after ~ 5 minutes (Appendix Figure S1C). We assume that optoFGFR is active in its clustered form in which transphosphorylation occurs and inactive in its monomeric form due to tonic cytosolic phosphatase activity (Lemmon et al. 2016). As documented previously (Kim et al. 2014), light stimulation also triggered optoFGFR endocytosis (Appendix Figure S1B, red arrows).

Directly following light stimulation, we systematically observed a short ERK
inactivation period, that we refer to as "dip", lasting 2-3 minutes before activation of a
strong ERK activity (Appendix Figure S1D, green rectangle). This light-induced ERK
dip was insensitive to SCH772984-mediated ERK inhibition but could be suppressed
by Cyclosporin A-mediated calcineurin inhibition. Calcineurin is a Ca^{2+}-dependent
phosphatase that dephosphorylates Ser383 in Elk1 (Sugimoto et al. 1997). As ERK-
KTR contains an Elk-1 docking domain phosphorylated by ERK (Regot et al. 2014),
we hypothesized that it could be negatively affected by optoFGFR-evoked Ca^{2+} input
(Kim et al. 2014) (Appendix Figure S1E). Consistently, Ionomycin-evoked increase in
cytosolic Ca^{2+} induced a dip in absence of light stimulation (Appendix Figure S1F).
143

144
Figure 1: An optogenetic actuator-biosensor genetic circuit to study input-dependent ERK
dynamics. (A) Schematic representation of the optoFGFR system consisting of the optogenetic FGF
receptor (optoFGFR) tagged with mCitrine, the ERK biosensor (ERK-KTR) tagged with mRuby2 and a
nuclear marker (H2B) tagged with miRFP703. **(B)** Time lapse micrographs of ERK-KTR dynamics in
response to a 470 nm light pulse. Using a 20x air objective, ERK-KTR and H2B channels were acquired
every 1 minute and the optoFGFR channel was acquired once at the end of the experiment. Scale bar:
50 μm. **(C)** Image analysis pipeline developed to quantify single-cell ERK dynamics. Nuclear and
cytosolic ERK-KTR signals were segmented based on the H2B nuclear mask. Single-cell ERK activity
was then calculated as the cytosolic/nuclear ERK-KTR ratio. Single-cell optoFGFR intensity was
measured under the cytosolic ERK-KTR mask and used as a proxy for single-cell optoFGFR
expression.

Different optoFGFR inputs trigger transient, oscillatory and sustained ERK
dynamics

Next, we characterized optoFGFR-triggered ERK dynamics in response to a single
light pulse of different intensities and durations (Figure 2A). As ERK dynamics

depended on light power density, as well as pulse duration, we defined the light dose
(D, mJ/cm^2) as their product to quantify the total energy received per illuminated area.
To characterize ERK dynamics, we extracted the amplitude at the maximum of the
peak (maxPeak), and the full width at half maximum (FWHM) of the ERK trajectories
(Figure 2B). With increasing light doses, ERK peaks increased both in duration and
amplitude, until the latter reached saturation. Based on these observations, we
selected 180 mW/cm^2 and 100 ms (D = 18 mJ/cm^2) as the minimal light input to
generate an ERK transient of maximal amplitude. Using this light dose, we then
investigated ERK dynamics in response to multiple light pulses delivered at different
intervals (Figure 2C). All stimulation regimes led to identical maximal ERK amplitude
(Figure EV1A) and adaptation kinetics when optoFGFR input ceased (Figure EV1B).
Repeated light inputs applied at 10- or 20-minute intervals evoked population-
synchronous ERK transients. In contrast, repeated light inputs applied at higher
frequencies (2-minute intervals) led to sustained ERK dynamics. Given CRY2's 5-
minute dissociation half-life (Appendix Figure S1B-C) (Duan et al. 2017), this suggests
that light pulses delivered at a 2-minute interval reactivate optoFGFR faster than it
deactivates, leading to sustained optoFGFR activity. Hierarchical clustering of ERK
responses to sustained optoFGFR input highlighted the presence of sustained and
oscillatory single-cell ERK dynamics (Figure 2D). Classification of ERK trajectories
based on optoFGFR expression revealed that sustained/oscillatory ERK dynamics
correlated with high/low optoFGFR levels (Figure 2E, Appendix Movie S3). Oscillatory
ERK dynamics were also observed in optoFGFR high expressing cells in response to
low light input (Figure 2F). Thus, sustained optoFGFR input can trigger sustained or
oscillatory ERK dynamics depending on the input strength, a combination of light
energy and optoFGFR expression.

Figure 2: Different optoFGFR inputs trigger transient, oscillatory and sustained ERK dynamics.
(A) ERK responses to increasing light power densities and pulse durations of 470 nm transient light
input. The light dose "D" is calculated as the product of the power density and pulse duration. **(B)**
Quantification of the maxPeak (maximal ERK amplitude of the trajectory) and the FWHM (full width at
half maximum) of single-cell ERK responses shown in (A) (N_{min} = 40 cells per condition). **(C)** ERK
responses to 470 nm light pulses delivered every 20, 10, 5 and 2 minutes respectively (D = 18 mJ/cm^2).
(D) Hierarchical clustering (Euclidean distance and Ward D2 linkage) of trajectories from the 2-minute
interval stimulation shown in (C) (referred to as "sustained") (N = 60 cells). The number of clusters was
empirically defined to resolve the different ERK dynamics. The average ERK responses per cluster are
displayed on the right. **(E)** Separation of the trajectories shown in (D) in low and high optoFGFR cells,
based on the log10 intensity of optoFGFR-mCitrine. **(F)** ERK responses to increasing doses of
sustained optoFGFR input. Single-cell ERK trajectories were divided in low (top panel) and high (bottom
panel) optoFGFR expression.

ERK dynamics evoked by optoFGFR versus endogenous RTKs highlight different MAPK regulatory mechanisms

Because of the absence of an ectodomain, optoFGFR must be considered as a prototypic RTK that lacks some regulatory mechanisms inherent to the native FGFR. To evaluate if optoFGFR is relevant for studying the MAPK network, we compared ERK dynamics evoked by optoFGFR inputs versus stimulation of the endogenous FGFR or EGFR using increasing concentrations of basic FGF (bFGF) and EGF. All bFGF concentrations led to an ERK peak similar in amplitude to sustained optoFGFR input (Figure 3A, EV1C, compared to EV1A). However, FGFR inputs led to different ERK dynamics than optoFGFR: 1 ng/ml bFGF led to damped ERK oscillations followed by steady state sustained ERK activity, while 10 and 100 ng/ml bFGF concentrations led to a first ERK peak followed by a strong adaptation. The biphasic behavior induced by increasing bFGF concentrations was previously documented to emerge from the competition of bFGF for FGFR and heparan sulfate proteoglycan co-receptors (Kanodia et al. 2014; Blum et al. 2019). It is thus not surprising that optoFGFR, that lacks these extracellular interactions, produced different ERK dynamics than FGFR. All EGF concentrations led to an ERK peak similar in amplitude to optoFGFR and FGFR inputs (Figure 3B, EV1D). As for bFGF, 1 ng/ml EGF concentration evoked damped oscillatory ERK dynamics that decreased at higher EGF concentrations. However, EGFR inputs led to strong ERK adaptation, not observed in response to optoFGFR inputs, suggesting the existence of different regulatory mechanisms.

Both oscillatory and transient ERK dynamics can be explained by the presence of NFB (Kholodenko et al. 2010). Thus, we wondered if the different ERK dynamics induced by optoFGFR or EGFR input emerge from differences in downstream NFBs. We reasoned that if EGFR induces different NFBs than optoFGFR, pre-stimulating cells with EGF should activate these feedbacks, and affect subsequent optoFGFR-evoked ERK dynamics. To test this, we pre-stimulated cells with sustained EGFR input, subsequently applied sustained optoFGFR input, and evaluated ERK dynamics (Figure 3C). Pre-stimulation with 100 ng/ml EGF led to the characteristic adaptive ERK transient. Subsequent application of optoFGFR input yielded sustained ERK responses similar in amplitude and duration to non-pre-stimulated cells. However, EGF pre-stimulation led to a reduction of synchronous optoFGFR-evoked ERK oscillations in low optoFGFR expressing cells.

To provide intuition about the MAPK network circuitries leading to different ERK dynamics in response to optoFGFR and EGFR inputs, as well as the origin of the oscillatory behavior, we built a mathematical model consisting of the RAS GTPase and the three-tiered RAF/MEK/ERK network (Figure 3D, Appendix Table S1). We used ordinary differential equations with Michaelis-Menten kinetics (see Material and methods, Appendix Table S2 and S3). To account for the oscillatory ERK dynamics in response to EGFR and optoFGFR inputs, we included the well-documented ERK-RAF NFB (Kholodenko et al. 2010; Santos et al. 2007; Fritsche-Guenther et al. 2011; Blum et al. 2019). We also included a receptor level inactivation process for EGFR, but not for optoFGFR, to account for EGF-dependent regulatory mechanisms. We used a

Bayesian inference approach (Mikelson and Khammash 2020) to infer the model
parameters from averaged ERK trajectories in response to sustained low optoFGFR
input with or without sustained EGFR pre-stimulation (Figure 3E). After identification
of parameters that allowed the model to capture the training dataset (Figure 3F), we
simulated ERK dynamics evoked by low EGFR input (adaptative, oscillatory ERK
dynamics), high EGFR input (adaptative ERK dynamics without oscillation) and
sustained high optoFGFR input (sustained ERK dynamics) (Figure 3G). We observed
that our model with a NFB and EGFR inactivation was able to predict ERK dynamics
evoked by different EGFR and optoFGFR input strengths, while two simpler models
(one with only the EGFR inactivation reaction, but no NFB (Figure EV1E-G) and one
with only the NFB, but no EGFR inactivation (Figure EV1H-J) were not able to
reproduce experimentally observed ERK dynamics.

This suggested that oscillatory optoFGFR-evoked ERK dynamics emerge from a NFB
also present downstream of endogenous EGFR, while additional regulatory
mechanisms seem to be required for the strong ERK transient adaptation following
EGFR input. These mechanisms might consist of receptor-level regulations such as
endocytosis, which was recently shown to be an important regulator of the transient
adaptive EGF-triggered ERK dynamics in different cell systems (Kiyatkin et al. 2020;
Gerosa et al. 2020). While optoFGFR also gets endocytosed (Appendix Figure S1B,
(Kim et al. 2014)), it most likely is insensitive to inactivation by endosome acidification
since it lacks an ectodomain (Huotari and Helenius 2011). Additionally, light-mediated
optoFGFR dimerization might occur both at the plasma and endo-membranes,
allowing for reactivation of endocytosed optoFGFR. The hypothesis that a receptor
level mechanism is important for strong adaptation was further supported by inhibition
of optoFGFR with the FGFR kinase inhibitor (SU5402), which shifted ERK dynamics
from sustained to transient in a dose response-dependent manner (Figure EV1K).
Thus, these results suggest that optoFGFR lacks receptor-dependent regulatory
mechanisms but allows us to investigate the intracellular MAPK feedback structure
shaping ERK dynamics. In our model, we used the well-established ERK-RAF NFB.
However, several NFBs have been mapped in the MAPK signaling cascade, whose
role in shaping ERK dynamics is still unknown and which could also be responsible
for the observed oscillatory ERK dynamics.

Figure 3: ERK dynamics evoked by optoFGFR versus endogenous RTKs highlight different MAPK regulatory mechanisms. (A-B) Single-cell ERK trajectories under increasing concentrations of sustained **(A)** bFGF or **(B)** EGF input added at t = 5 minutes. **(C)** ERK responses of cells stimulated with sustained optoFGFR input (D = 18 mJ/cm^2) at t = 24 minutes without or with 100 ng/ml EGF sustained pre-stimulation at t = 5 minutes. Average ERK responses for optoFGFR high and low expression levels are shown (N = 20 cells for low and high optoFGFR, randomly selected out of at least 80 cells). **(D)** Mathematical model topology consisting of the RAS GTPase, the MAPK three-tiered (RAF, MEK, ERK) network and the ERK-KTR reporter. EGFR and optoFGFR inputs both activate the RAS/RAF/MEK/ERK cascade and the ERK-RAF NFB. EGFR activity is under receptor-dependent regulations. **(E)** Training dataset consisting of the average ERK responses evoked by sustained low optoFGFR input with or without pre-stimulation with 100 ng/ml sustained EGF. **(F)** Simulation of ERK responses from the training dataset, including the maximum a posteriori (MAP) estimate, the posterior envelope indicating the predictive density of our estimation, as well as an example trajectory. **(G)** Predictions of the model for ERK responses evoked by 1 ng/ml EGF, 100 ng/ml EGF and sustained high optoFGFR inputs. Note that for low EGFR input (1 ng/ml), the model predicts both adaptive and oscillatory ERK responses.

RNA interference screen reveals that ERK dynamics remain unaffected in response to perturbation of most MAPK signaling nodes

We then explored the network circuitry that shapes optoFGFR-evoked ERK dynamics with an RNA interference (RNAi) screen targeting 50 MAPK signaling nodes. We focused our screen on sustained optoFGFR input which captured the largest amount of information about ERK dynamics when compared to other stimulation schemes: it led to sustained and oscillatory ERK dynamics (Figure 2E,F) while recapitulating the rapid increase of ERK activity and adaptation observed with transient input (Figure EV1A,B). We used a bioinformatic approach to select 50 known interactors of the tripartite RAF/MEK/ERK cascade downstream of the FGFR receptor that were detected in a NIH3T3 proteome (Schwanhäusser et al. 2011) (Figure 4A, Appendix Table S4). We used the siPOOL technology to specifically knockdown (KD) these 50 MAPK signaling nodes while limiting off-target effects (Hannus et al. 2014). We first validated KD efficiency by quantifying transcript levels with different siPOOL concentrations targeting the ERK and MEK isoforms (Figure EV2A) and observed strong KD with 10 nM siRNA concentration. We then evaluated the effect of *ERK1* or *ERK2* KD on ERK dynamics. We observed only subtle phenotypes compared to the non-targeting siRNA (*CTRL*) used as negative control (Figure 4B), even though efficient KD was observed at protein level (Figure 4C). However, combined *ERK1/ERK2* KD strongly suppressed ERK dynamics indicating that the latter is not affected by the perturbation of individual ERK isoforms as previously reported (Fritsche-Guenther et al. 2011; Ornitz and Itoh 2015). Due to its strong phenotype, we used *ERK1/ERK2* KD as positive control throughout our screen.

We performed three replicates of the screen targeting the 50 nodes. Despite efficient KD quantified for different nodes (Figure EV2B), visual inspection of ERK trajectories only revealed subtle ERK dynamics phenotypes for a limited number of node perturbations (Figure EV2C,D). We used a feature-based approach to evaluate the effect of each perturbation on ERK dynamics. We focused our analysis on ERK responses evoked by high optoFGFR input to limit the single-cell heterogeneity due to optoFGFR expression variability. We quantified the average ERK activity before stimulation (baseline), the maximal ERK amplitude during stimulation (maxPeak), and the ERK amplitude at a fixed time point after response adaptation in the negative control (ERKpostStim). To evaluate these phenotypes, we z-scored the features associated to each perturbation to those of the negative control (Figure 4D, see Material and method for details). While many phenotypes were statistically significant, most of them remained mild as observed by visually inspection of the feature distributions (Figure EV3A). Apart from *ERK1+2* KD, only *GRB2, PTK2* and *ERK2* led to a reduction of ERK amplitude (maxPeak). KD of negative regulators such as *SPROUTY 2,3* and *4*, or phosphatases such as *PP2A* and several dual-specificity phosphatases (DUSPs*)* led to increased ERK amplitude. Increased basal ERK activity was observed for *RKIP, PP2A*, *DUSP4* and *DUSP6* KDs, indicating a function in regulating basal ERK levels. Prolonged ERK activity (ERKpostStim) was observed in

response to KD of *RKIP*, *PP2A*, *ERK2*, *DUSP1,2,3,4,6* and strikingly for *RSK2* KD (Figure EV3B), suggesting a role of these nodes in ERK adaptation.

Because both visual inspection of trajectories, as well as our feature-based approach might miss more subtle ERK dynamics phenotypes, we used CODEX (Jacques et al. 2021), a data-driven approach to identify patterns in single-cell time-series based on convolutional neural networks (CNNs) (Figure EV3C). We trained a CNN to classify ERK trajectories that originate from different siRNA perturbations and selected the ten perturbations for which the CNN classification accuracy was the highest (Appendix Table S4, "CODEX accuracy", see Material and methods for details). Projection of the CNN features in a t-distributed stochastic neighbor embedding (t-SNE) space revealed different clusters of ERK trajectories (Figure EV3D). Comparison of the ten trajectories with the highest classification confidence identified by CODEX to randomly selected ERK trajectories for low or high optoFGFR expression highlighted ERK phenotypes not accessible to visual inspection and the feature-based approach (Figure 4E). CODEX identified some of the perturbations that affect ERK amplitude, baseline or adaptation observed with the feature-based approach. However, it also highlighted perturbations affecting oscillatory ERK dynamics. *PP2A* KD led to sustained oscillatory behavior. *PLCG1* KD resulted in a first peak followed by damped oscillations, and absence of the dip. As phospholipase C mediates Ca^{2+} signaling in response to FGFR activation (Ornitz and Itoh 2015), this further validates the role of Ca^{2+} signaling in formation of the dip (Appendix Figure S1D-F). *RAPGEF1* KD led to oscillatory ERK responses of different amplitudes. *RSK2*, *ERK2* and *CRAF* KD displayed reduced oscillatory ERK behavior.

To validate the latter oscillatory ERK dynamics phenotypes, we evaluated the proportion of oscillatory trajectories (trajectories with at least 3 peaks) for each perturbation, both for high and low optoFGFR input (Figure 4F). This confirmed that *RSK2*, *CRAF* and *ERK2* KD led to decreased oscillatory ERK dynamics. We also observed that these perturbations reduced ERK oscillations in cells stimulated with 1 ng/ml EGF (Figure EV3E-G), suggesting a role of these nodes in the regulation of ERK oscillations in the context of a native RTK.

ERK2 and CRAF isoforms are implicated in the well-established ERK-RAF NFB, known to regulate ERK dynamics (Santos et al. 2007; Ryu et al. 2015; Blum et al. 2019), and to enable consistent ERK dynamics under MEK or ERK perturbations (Fritsche-Guenther et al. 2011; Sturm et al. 2010). *RSK2* encodes the p90 ribosomal S6 kinase 2 protein, an ERK substrate regulating survival and proliferation (Cargnello and Roux 2011; Yoo et al. 2015). RSK2 is also known to be involved in an ERK-induced NFB targeting SOS (Douville and Downward 1997; Saha et al. 2012; Lake et al. 2016), whose significance in the regulation of ERK dynamics has been less well studied. In addition to dampening ERK oscillations, *RSK2* KD also led to slower ERK adaptation when optoFGFR input ceased (Figure 4D, EV3A,B), suggesting an important role of this NFB in ERK dynamics regulation. Our results suggest that the ERK-RAF and ERK-RSK2-SOS NFBs simultaneously operate within the MAPK network to generate ERK oscillations and raise the question whether both NFBs contribute to the strong MAPK signaling robustness observed in our screen.

380
Figure 4: RNA interference screen reveals that ERK dynamics remain unaffected in response to
perturbation of most MAPK signaling nodes. (A) RNAi perturbation targets referred to by their
protein names. Nodes were spatially grouped based on the hierarchy of interactions within the MAPK
network and color-coded for their function. **(B)** ERK responses to sustained optoFGFR input (D = 18
mJ/cm^2) in cells transfected with 10 nM siRNA against *ERK1*, *ERK2* or a 5 nM combination of each
(*ERK1+2*). A non-targeting siRNA (*CTRL*) was used as control (N = 15 cells from low and high
optoFGFR levels). **(C)** Western blot analysis of cells transfected with 10 nM siRNA against *ERK1*, *ERK2*
or a 5 nM combination of each (*ERK1+2*). **(D)** Z-Score evaluation of the baseline, maxPeak and

ERKpostStim of single-cell ERK responses under sustained high optoFGFR input (D = 18 mJ/cm^2). The
z-score was calculated by comparing each RNAi perturbation to the *CTRL* KD (N$_{min}$ = 126 cells per
treatment, from 3 technical replicates). Non-significant (NS) results are in grey (see Figure EV3A for
statistical results). **(E)** Single-cell ERK trajectories (sustained optoFGFR input, D = 18 mJ/cm^2) for the
RNAi perturbations classified with the highest accuracy by CODEX. Top lines show single-cell ERK
trajectories for which CODEX had the highest classification confidence in the validation set (N = 10).
Bottom lines show single-cell ERK trajectories for low and high optoFGFR cells (N = 30 for each
condition, randomly selected out of at least 212 cells per perturbation from 3 technical replicates). For
easier visualization, the CTRL condition is shown twice. **(F)** Proportion of oscillating cells (trajectories
with at least 3 peaks) per RNAi perturbation for low and high optoFGFR expression (sustained
optoFGFR input, D = 18 mJ/cm^2, N$_{min}$ = 61 cells for low and 126 for high optoFGFR per perturbation
from 3 technical replicates). Perturbations were ordered based on the proportion of oscillating cells with
low optoFGFR expression. Statistical analysis was done using a pairwise t-test, comparing each
perturbation against the *CTRL* for each receptor level independently (*<0.05, **<0.005, ***<0.0005,
****<0.00005, FDR p-value correction method).

Direct optogenetic activation of RAS highlights different ERK dynamics phenotypes than optoFGFR input

To further explore the role of MAPK feedbacks in MAPK signaling robustness, we used
optoSOS (Johnson et al. 2017), an optogenetic actuator that activates RAS, and thus
bypasses the RSK2-mediated NFB regulation (Figure 5A). OptoSOS consists of a
membrane anchored light-activatable iLID domain, and an mCitrine-tagged SspB
domain fused to SOS's catalytic GEF domain. It was stably integrated into cells
expressing ERK-KTR and H2B. Because iLID displays faster dissociation rates than
CRY2 (t$_{1/2}$= 30 seconds for iLID versus ~ 5 minutes for CRY2 (Duan et al. 2017;
Benedetti et al. 2018)), optoSOS required repeated light pulses to prolong its
membrane recruitment and produce a robust ERK response (Figure 5B). Five
consecutive 100 ms light pulses at 6 W/cm^2 (D = 0.6 J/cm^2) applied at 20-second
intervals, provided the minimal light input to induce a saturated ERK amplitude (Figure
EV4A). Application of this light input at 2-minute intervals evoked sustained ERK
dynamics with small fluctuations at the same frequency as the light input pattern,
reflecting the fast optoSOS reversion to the dark state (Figure 5C). OptoSOS did not
induce ERK oscillations (Figure EV4B), even in cells with low optoSOS expression or
at lower light doses (Figure 5D). However, ERK amplitudes correlated with optoSOS
expression level, low optoSOS levels led to low ERK amplitudes, while high actuator
expression levels resulted in high ERK amplitudes. Using the minimal light input to
trigger saturating ERK amplitude, both optoSOS and optoFGFR led to steep ERK
activation and fast adaptation when light stimulation ceased (compare Figures 2C and
5C), as well as similar ERK amplitudes in cells expressing high actuator levels (Figure
5E). However, high optoSOS expression levels moderately increased ERK activity
baseline levels in comparison to optoFGFR (Figure EV4C), suggesting that this
system is leaky to some extent.
Using this specific light input, we performed siRNA screens targeting MAPK signaling
nodes downstream of optoSOS in triplicates (Figure EV4D,E). We extracted the
baseline, maxPeak, ERKpostStim features from optoSOS high expressing cells
(Figure EV4F) and z-scored feature values to the negative control (Figure 5F). We

observed more prominent ERK amplitude phenotypes in response to optoSOS input
than to optoFGFR input. Some of these phenotypes are shown in Figure 5G. Most
prominently, *CRAF*, *ERK2, DUSP4* KD led to a stronger reduction in ERK amplitude
than observed with optoFGFR input. *RSK2* KD also reduced ERK amplitude,
suggesting that it also regulates nodes downstream of RAS. However, *RSK2* KD did
not decrease ERK adaptation following optoSOS input removal (Figure EV4G),
suggesting that it is not involved in NFB regulation in this system. *PP2A* KD did not
induce increased ERK amplitude or baseline as observed in the optoFGFR system.
As for optoFGFR input, *DUSP6* KD increased basal ERK activity and decreased
adaptation (Figure EV4G). *DUSP22* KD led to increased amplitude, without affecting
ERK baseline and adaptation. *NF1* KD, which encodes a RAS-specific GAP, led to
increased ERK baseline and slower adaptation (Figure EV4G), without affecting ERK
amplitude. The NF1 baseline phenotype, that was not observed in the optoFGFR
system, might emerge from the optoSOS-mediated low levels of RAS activation due
to the optoSOS system's leakiness (Figure EV4C), that can then be amplified by loss
of NF1's RAS GAP activity. The finding that perturbation of specific nodes (e.g. ERK2
and CRAF) leads to more penetrant phenotypes in response to optoSOS versus
optoFGFR input suggested that the RAS/RAF/MEK/ERK part of the network is more
sensitive to perturbations than optoFGFR-triggered network, suggesting that the
RSK2 NFB that operates above RAS contributes to MAPK signaling robustness.

454
Figure 5: Direct optogenetic activation of RAS highlights different ERK dynamics phenotypes
than optoFGFR input. (A) Schematic representation of ERK signaling induced by optoSOS versus
optoFGFR input. **(B)** ERK dose responses under transient optoSOS input consisting of different
numbers of repeated 470 nm pulses (1x, 2x, 3x, 4x and 5x pulses applied at 20-second intervals, D =
0.6 J/cm^2). Repeated pulses are depicted as a single stimulation (blue bar). **(C)** ERK responses to
optoSOS inputs consisting of 5 repeated 470 nm light pulses delivered every 20, 10, 5 and 2 minutes
respectively (D = 0.6 J/cm^2). **(D)** ERK responses to increasing light doses of sustained optoSOS input
consisting of 2-minute interval input, each input made of 5 repeated light pulses. Cells were divided in
low and high optoSOS expression levels based on the log10 intensity of the optoSOS-mCitrine. **(E)**
Quantification of the maxPeak of single-cell ERK responses under sustained optoFGFR (Figure 2F, D
= 18 mJ/cm^2) and optoSOS (Figure 5D, D = 0.6 J/cm^2) input for low or high expression of each
optogenetic system (N = 40 cells per condition). Statistical analysis was done using a Wilcoxon test,
comparing each condition to each other (N$_{min}$ = 48 cells per condition, NS: non-significant, *<0.05,
<0.005, *<0.0005, ****<0.00005, FDR p-value correction method). **(F)** Z-Score evaluation of the

baseline, maxPeak and ERKpostStim of single-cell ERK responses under sustained high optoSOS
input ($D = 0.6$ J/cm^2). The z-score was calculated by comparing each RNAi perturbation to the *CTRL*
KD (N_{min} = 33 cells per treatment, from 3 technical replicates). Non-significant (NS) results are in grey
(see Figure EV4F for statistical results). **(G)** Single-cell ERK trajectories for low and high optoSOS cells
for selected RNAi perturbations (N = 40 randomly selected out of at least 193 trajectories from 3
technical replicates).

Perturbation of the RSK2-mediated NFB increases the efficiency of RAF, MEK and ERK targeting drugs

To further investigate the role of the RSK2-mediated NFB in MAPK signaling
robustness, we performed dose response experiments using different MAPK inhibitors
and compared ERK amplitudes evoked by optoFGFR (RSK2-feedback dependent)
versus optoSOS (RSK2-feedback independent) input, as well as optoFGFR input in
absence/presence of RSK2 perturbation. We used drugs targeting B/CRAF (RAF709),
MEK (U0126) and ERK (SCH772984). We evaluated the inhibition efficiency by
measuring ERK amplitude at a fixed time point, focusing on ERK responses evoked
by high optoFGFR or optoSOS inputs to limit the single-cell heterogeneity due to
expression variability of the optogenetic actuator. All inhibitors led to a stronger
reduction of ERK amplitude and EC$_{50}$ in response to optoSOS versus optoFGFR input
(Figure 6A-C, EV5A, Appendix Table S5). Visual evaluation of ERK amplitude
distributions (Figure 6B) and quantification of their standard deviations (Figure 6D)
revealed more compact ERK amplitude distributions in presence of increasing drug
concentrations in response to optoSOS versus optoFGFR input. This suggests a more
homogeneous drug inhibition in the cell population in response to optoSOS input. We
then performed the identical experiments in *CTRL* or *RSK2* KD cells in response to
optoFGFR input (Figure 6E-H, EV5B, Appendix Table S6). *RSK2* KD led to increased
inhibition of ERK amplitudes, decreased EC$_{50}$, and more compact ERK amplitude
distributions in response to increasing drug concentration than in *CTRL* KD cells.
Similar results were observed when the RSK2-mediated feedback was inhibited using
the RSK inhibitor SL0101 (Smith et al. 2005) (Figure EV5C-F, Appendix Table S7).
Thus, inhibition of the RSK2-mediated NFB sensitizes ERK responses to RAF, MEK
or ERK drug perturbations. Note that drug mediated ERK amplitude inhibition was
stronger in response to optoSOS input than to optoFGFR input with *RSK2* KD or RSK
inhibition, suggesting that additional mechanisms to the RSK2-mediated feedback
contribute to MAPK signaling robustness. However, our results suggest that
perturbation of the RSK2-mediated feedback can be exploited to enhance the
efficiency of MAPK-targeting drugs, reducing ERK amplitudes more homogeneously
across the cell population.

506
Figure 6: Perturbation of the RSK2-mediated NFB increases the efficiency of RAS, MEK and ERK
targeting drugs. (A) Schematic representation of the optoFGFR (RSK2-mediated feedback
dependent) and optoSOS (RSK2-mediated feedback independent) systems targeted with the B/CRAF
(RAF709), the MEK (U0126) or the ERK (SCH772984) inhibitor. **(B)** Single-cell ERK amplitudes from
sustained high optoFGFR input (D = 18 mJ/cm^2) or optoSOS input (D = 0.6 J/cm^2) under different
concentrations of the MAPK inhibitors, extracted at a fixed time point ($t_{fixed\ optoFGFR}$ = 15 minutes, t_{fixed}
$_{optoSOS}$ = 10 minutes, N = 200 cells with high optoFGFR or optoSOS expression per condition randomly
selected from 3 technical replicates). **(C)** A Hill function was fit to the normalized mean ERK activity as
shown in (B) (N_{min} = 200 cells per condition). Shaded area indicates the 95% CI and dashed lines the
EC$_{50}$. **(D)** Normalized standard deviation of ERK amplitudes shown in (B) (N_{min} = 200 cells per
condition). **(E)** Schematic representation of the optoFGFR system treated with *CTRL* KD (RSK2-
mediated feedback dependent) or *RSK2* KD (RSK2-mediated feedback independent) targeted with the
B/CRAF (RAF709), the MEK (U0126) or the ERK (SCH772984) inhibitor. **(F)** Single-cell ERK
amplitudes from sustained high optoFGFR input (D = 18 mJ/cm^2) under different concentrations of the
MAPK inhibitors, extracted at a fixed time point ($t_{fixed\ optoFGFR}$ = 15 minutes, N = 70 cells with high
optoFGFR expression per condition (apart from *RSK2* KD + 0 µM U0126 (32 cells), randomly selected
from 2 technical replicates for *RSK2* KD and 1 replicate for *CTRL* KD). **(G)** A Hill function was fit to the
normalized mean ERK activity as shown in (F) (N_{min} = 32 cells per perturbation). Shaded area indicates
the 95% CI and dashed lines the EC$_{50}$. **(H)** Normalized standard deviation of ERK amplitudes shown in
(F) (N_{min} = 32 cells per perturbation).

Targeting the RSK2-mediated feedback in an ErbB2 oncogenic signaling model increases MEK inhibition efficiency

The results above suggested an important role of the RSK2-mediated feedback in MAPK signaling robustness against node perturbation in response to optogenetic inputs in NIH3T3 cells. To test if this feedback also contributes to MAPK signaling robustness in a disease-relevant system, we evaluated its function in MCF10A cells, a breast epithelium model, using either wild-type (WT) or overexpressing ErbB2 (referred to as ErbB2^{over}) recapitulating the ErbB2 amplification observed in 20% of all breast cancers (Arteaga and Engelman 2014; Yarden and Pines 2012). We chose this specific model system because ErbB2 amplification leads to constitutive RTK input on the MAPK network, while retaining an intact downstream feedback structure (Figure 7A). This contrasts with other cancer model systems in which additional mutations might lead to RAS or RAF overactivation, and thus disrupt the feedback architecture. Further, previous work has highlighted the role of NFBs in ERK pulse formation in MCF10A cells (Kochańczyk et al. 2017), suggesting that EGFR and ErbB2 trigger a MAPK network with similar feedback circuitry as optoFGFR.

As described before (Albeck et al. 2013), WT cells displayed asynchronous low frequency ERK pulses in the absence of EGF, and high frequency ERK pulses in presence of EGF (Figure 7B). In marked contrast, ErbB2^{over} cells displayed high frequency ERK pulses, even in the absence of EGF (Figure 7C). To investigate the role of the RSK2-mediated feedback in MAPK signaling robustness, we performed a U0126 dose response in EGF-stimulated MCF10A WT cells and found that 3 µM U0126 decreased ERK amplitude without fully suppressing the response (Figure EV5G,H). As observed in response to optogenetic inputs, RSK inhibition with 50 µM SL0101 led to a mild reduction in ERK amplitude. However, in combination with 3 µM U0126, ERK amplitude was decreased to the level of unstimulated cells. Similar results were observed in ErbB2^{over} cells (Figure EV5I), suggesting that RSK2 perturbation increases the sensitivity of ERK responses to MEK inhibition.

As averaging ERK dynamics can hide asynchronous single-cell signaling activity, we further investigated the effect of these perturbations on single-cell trajectories using CODEX (Jacques et al. 2021) (see Material and methods for details). For WT cells, a tSNE projection of the CNN features built from single-cell ERK trajectories hinted that the CNN was able to construct features separating the treatments into well-defined clusters (Figure 7D, EV5J). Clustering of the CNN features confirmed the existence of discrete ERK dynamics clusters (Figure 7E) whose composition correlated with the treatments (Figure 7F). To characterize the dynamics captured by each cluster, we extracted the medoid trajectory and its 4 closest neighbors from each cluster (Figure 7G). This revealed that non-stimulated cells mostly display low frequency ERK activity pulses (cluster 4) or absence of pulses (cluster 5). Cells stimulated with EGF without inhibitor displayed ERK pulses of high amplitude (cluster 1). SL0101-treated cells displayed a sustained ERK activation at low amplitude (cluster 3). U0126-treated cells still displayed prominent ERK pulses but at a lower amplitude than EGF-treated cells in absence of drug (cluster 2). Finally, in cells treated with both U0126 and SL0101,

almost no ERK activity was observed (cluster 5). For ErbB2[over] cells, we observed that the CNN features were forming a more continuous space with less distinct clusters (Figure 7H,I, EV5K). A heterogeneous mix of ERK trajectory clusters was observed for the different treatments (Figure 7J,K). Untreated cells mostly displayed high frequency ERK pulses that were either sharp (cluster 3) or wider (cluster 2). SL0101-treated cells were almost equally shared between cluster 1 (relatively flat high amplitude ERK trajectories), cluster 2, cluster 4 (low amplitude ERK pulses) and cluster 5 (low baseline activity). U0126 led to a less heterogeneous mix mostly consisting of ERK trajectories from cluster 4 and 5. The U0126/SL0101 combination shifted most cells to cluster 5, indicating an efficient inhibition of ERK activity at a suboptimal U0126 concentration. Thus, RSK inhibition also sensitizes the MAPK network to U0126-mediated MEK inhibition both in MCF10A WT and ErbB2[over] cells.

Figure 7: Targeting the RSK2-mediated feedback in an ErbB2 oncogenic signaling model
increases MEK inhibition efficiency. (A) Schematic representation of MAPK signaling in response to
EGFR input in MCF10A WT cells or oncogenic ErbB2 input in ErbB2 overexpressing (ErbB2^{over}) cells.
(B-C) Single-cell ERK responses in MCF10A WT cells without or with stimulation with 10 ng/ml EGF at
t = 30 minutes **(B)** and in unstimulated MCF10A ErbB2^{over} cells **(C)**. **(D)** tSNE projection of CODEX's
CNN features from ERK trajectories of MCF10A WT cells without EGF stimulation, or with 10 ng/ml
EGF stimulation added at t = 30 min in absence of perturbation, with 50 µM SL0101, 3 µM U0126 or a
combination of both. **(E)** t-SNE projection of CODEX's CNN features shown in (D) colored by the CNN
feature clusters. Black diamonds indicate the position of the medoid and its 4 closest neighbor
trajectories for each cluster. **(F)** Distribution of the trajectories in the CNN features clusters per
treatment. Colors are as shown in (E). **(G)** Medoid trajectories and their 4 closest neighbors per cluster
highlighted in (E) (black diamonds). **(H)** tSNE projection of CODEX's CNN features from ERK
trajectories of non-stimulated ErbB2 overexpressing cells without perturbation, with 50 µM SL0101, 3
µM U0126 or a combination of both. **(I)** t-SNE projection of CODEX's CNN features shown in (H) colored
by the CNN feature clusters. Black diamonds indicate the position of the medoid and its 4 closest
neighbor trajectories for each cluster. **(J)** Distribution of the trajectories in the CNN features clusters per
treatment. Colors are as shown in (I). **(K)** Medoid trajectories and their 4 closest neighbors per cluster
highlighted in (I) (black diamonds).

Discussion

Optogenetic actuator-biosensor circuits allow for feedback structure mapping in the MAPK network

ERK dynamics is crucial for fate decisions. Yet, the topology of the network enabling the cells to sense different inputs and convert this information into finely tuned ERK dynamics remains poorly understood. We developed genetic circuits consisting of optogenetic actuators and an ERK biosensor (Figure 1A, 5A) that allow for a large-scale interrogation of single-cell ERK dynamics and investigated the effects of 50 RNAi perturbations targeting components of the MAPK signaling network (Figure 4A). In our optoFGFR screen, we only observed a small number of penetrant ERK dynamics phenotypes (Figure 4D-F), implying that the MAPK network can buffer against perturbations of most of its components. We cannot exclude that in some cases, even on the relatively short 72 hours timescale of the RNAi experiment, compensation by upregulation of specific nodes might occur. However, our data suggest that the MAPK network topology allows for MAPK signaling robustness – the production of consistent ERK outputs in presence of node perturbation. This might emerge from isoform redundancy for multiple nodes in the network, as observed for single or combined ERK isoforms perturbation (Figure 4B), but also for individual perturbation of RAS, RAF, MEK isoforms. Another mechanism might involve NFBs that have been shown to decrease the network sensitivity to node perturbation (Sturm et al. 2010; Fritsche-Guenther et al. 2011). Our screen suggested that RSK2, that mediates a NFB from ERK to SOS (Douville and Downward 1997; Saha et al. 2012), both regulates ERK dynamics (Figure 4D-F) and plays a role in MAPK signaling robustness (Figure 6E-H). In addition, our data suggest that the well-studied ERK-RAF NFB, which has been shown to buffer against MAPK node perturbations (Sturm et al. 2010; Fritsche-Guenther et al. 2011), also regulates ERK dynamics (Figure 4F). We therefore speculate that both feedbacks operate simultaneously in the MAPK network, and act at multiple levels within the cascade to warrant MAPK signaling robustness. Consistently with this hypothesis, we observed that the optoSOS-triggered network, which is not under the RSK2 NFB regulation, shows an increased sensitivity in ERK amplitude to perturbation of some nodes (Figure 5F,G). Indeed, ERK2 and CRAF perturbations, which led to loss of ERK oscillations, had relatively mild amplitude phenotypes in response to optoFGFR input, while both perturbations led to strong ERK amplitude phenotypes in response to optoSOS input. Because these phenotypes were not observed with other ERK and RAF isoforms, we propose that ERK2 and CRAF are the isoforms involved in the classic ERK-RAF NFB. Additional feedbacks have been reported within the MAPK network (Langlois et al. 1995; Lake et al. 2016; Kochańczyk et al. 2017), and even if they have not been highlighted in our screen, they might also regulate ERK dynamics.

While providing the experimental throughput to perturb and analyze ERK dynamics at scale, optoFGFR, that lacks an ectodomain, evoked different ERK dynamics than endogenous RTKs such as FGFR and EGFR (Figure 3A,B compared to Figure 2F).

These different ERK dynamics emerge likely because of receptor-level interactions
that involve competition of bFGF for FGFR and heparan sulfate proteoglycan co-
receptors (Kanodia et al. 2014; Blum et al. 2019) in the case of FGFR, or receptor
endocytosis in the case of EGFR (Kiyatkin et al. 2020; Gerosa et al. 2020). Our
combined modeling and experimental approach suggested that optoFGFR and EGFR
share similar downstream MAPK network circuitries and NFBs (Figure 3C-G).
OptoFGFR therefore provides a simplified system that allowed us to focus on
intracellular feedback structures, without confounding receptor level regulations. Our
Bayesian inference modeling approach, that is parameter agnostic, could provide
simple intuitions about the receptor-level and negative feedback structures that shape
ERK dynamics in response to optoFGFR and EGFR inputs. However, even if we had
access to many ERK dynamics phenotypes, our modeling approach did not allow us
to explore more sophisticated MAPK network topologies such as the presence of two
NFBs or multiple node isoforms. We interpreted our data using some of the feedback
structures that have been previously experimentally documented and modelled but
cannot formally exclude that the observed ERK dynamics emerge from different
network structures. In the future, information about the different nodes and their
dynamics might allow to further constrain the model topology and parameter space,
and hopefully address this limitation.

Additional novel insights into regulation of ERK dynamics

Our optoFGFR and optoSOS screens provided new system-wide insights into the
regulation of the MAPK network. Strikingly, the same perturbations induced different
ERK dynamics phenotypes in the optoFGFR and optoSOS screens. This might occur
because some regulators target the MAPK network at multiple levels, differently
affecting ERK responses triggered with optoFGFR or optoSOS inputs. Additionally, as
the two optogenetic systems are under the regulation of one versus two
simultaneously occurring NFBs, they might have different sensitivities to perturbations,
as discussed above.
With respect to the optoFGFR system, *GRB2* KD led to a reduction of ERK amplitude
(Figure 4D,E). GRB2 acts as the RTK-proximal adaptor to activate SOS (Chardin et
al. 1993; Belov and Mohammadi 2012). As GRB2 operates at the start of the cascade,
outside of most NFBs, heterogeneity in its expression levels might be less easily
buffered out. *PLCG1* KD increased damped oscillatory behavior (Figure 4E,F).
Phospholipase Cγ1 activates calcium signaling, which has itself been shown to
regulate RAS/MAPK signaling in a calcium spike frequency-dependent manner
(Kupzig et al. 2005; Cullen and Lockyer 2002). Further investigation will be required
to understand the significance of this crosstalk. *RKIP* KD resulted in higher ERK
baseline and slower ERK adaptation post stimulation, without affecting ERK amplitude
(Figure 4D). RKIP (RAF kinase inhibitor protein) prevents MEK phosphorylation by
CRAF (Yeung et al. 2000), suggesting that RKIP-dependent regulation is specifically
involved in keeping basal ERK activity low. With respect to phosphatases, none of
their perturbations led to a strong phenotype such as sustained ERK dynamics post

stimulation for example. The strongest phenotype was observed for *PP2A* KD that led
to increased ERK amplitude, baseline, and slower adaptation (Figure 4D, EV3A). This
might occur because the protein phosphatase 2A is an ubiquitous phosphatase that
acts at multiple levels by dephosphorylating SHC1, MEK1, MEK2, ERK1 and ERK2,
as well as a large number of other proteins (Junttila et al. 2008; Saraf et al. 2010). The
observation that in optoFGFR-low *PP2A* KD cells, ERK dynamics displayed increased
amplitude but still oscillated rather than exhibiting sustained behavior, suggests that
NFBs might buffer against the loss of phosphatase regulation to some extent.
Perturbation of the nuclear DUSPs, DUSP1,2,4, the atypical DUSP3 and most strongly
the cytosolic DUSP6 (Patterson et al. 2009) led to higher ERK baseline, reduced
adaptation, with only limited effects on amplitude (Figure 4D, EV3A). Consistently,
DUSP6 has previously been proposed to pre-emptively dephosphorylate MAPKs to
maintain low ERK activity baseline levels at resting state (Huang and Tan 2012). Our
results indicate that perturbation of single DUSPs might not be compensated by the
others, suggesting that individual DUSPs might regulate specific substrates within the
MAPK network. Except for *DUSP6*, KD of the different DUSPs did not significantly
affect oscillatory ERK behavior in optoFGFR-low cells (Figure 4F), suggesting that
they are not involved in the MAPK feedback circuitry that operates on timescales of
minutes.
The optoSOS screen revealed stronger ERK amplitude phenotypes, especially for
ERK2 and *CRAF* KD (Figure 5F versus 4D). Unlike for optoFGFR input, *RSK2* KD did
not result in slower ERK adaptation, suggesting that ERK responses triggered by the
optoSOS input are not regulated by the RSK2-mediated NFB. However, *RSK2* KD led
to a reduction of ERK amplitude, also observed to a lesser extent in response to
optoFGFR input, suggesting a role of RSK2 in ERK amplitude regulation downstream
of RAS. With respect to phosphatases, *PP2A* KD led to decreased amplitude, a
different phenotype than in response to optoFGFR input. This might occur because of
the broad specificity PP2A phosphatase, which might lead to different phospho-
proteomes in response to optoSOS versus optoFGFR input. Similar phenomena might
apply for most of the DUSPs.

The RSK2-mediated feedback can be targeted to potently inhibit oncogenic ErbB2 signaling

Our data suggest that the RSK2-mediated NFB is important for MAPK signaling
robustness downstream of our prototypic optoFGFR RTK (Figure 6). We found that
the RSK2-mediated NFB likely also operates downstream of EGFR and oncogenic
ErbB2 signaling in MCF10A cells (Figure 7). In response to EGF stimulation, or ErbB2
overexpression, a subset of RSK-inhibited cells displayed wider ERK pulses,
suggesting that the RSK2 NFB is also involved in ERK adaptation in this system
(Figure 7G cluster 3, Figure 7K cluster 1 and 2). Further, RSK inhibition led to a high
heterogeneity of ERK dynamics within the cell population especially visible in the case
of ErbB2 overexpressing cells (Figure 7J), which might result from the reduced ability
of the MAPK network to cope with nodes expression noise in absence of the RSK2

NFB. In EGF-treated cells, combination of RSK and suboptimal MEK inhibition led to strong and homogeneous ERK inhibition (Figure 7E-G, cluster 5). In the ErbB2 overexpressing cells, combined RSK/MEK inhibition shifted most of the cell population to flat, low amplitude ERK dynamics, enabling to further inhibit a large number of cells when compared to suboptimal MEK inhibition only (Figure 7I-K, cluster 5). These results suggest that pharmacological inhibition of the RSK2-mediated NFB can be used to reduce MAPK signaling robustness, sensitizing the network to MEK perturbation. Such non-trivial drug combinations might allow for homogeneous inhibition of ERK dynamics in most of the cells in a population. This homogeneous inhibition might mitigate the emergence of drug-tolerant persister cells from cell subpopulations that display residual ERK activity in response to inhibition of a single node. Our results imply that efficient pharmacological inhibition of the MAPK network requires precise understanding of its topology. The RSK2 NFB is an example of a druggable node that can be exploited to target MAPK signaling robustness.

Our scalable experimental pipeline provides new insight into the MAPK network wiring that produces ERK dynamics. However, our perturbation approach only highlighted very subtle ERK dynamics phenotypes, precluding a complete understanding of the MAPK network. We envision that this will require more precise knowledge about the dynamics of MAPK network nodes and their interactions in response to defined inputs and perturbations. Such data can now be produced at scale using optogenetic actuator/biosensor circuits as those we describe in this work. This information might allow for faithful parametrization of more complex models. With the increasing amount of optogenetic actuators and biosensors available, similar genetic circuits could also be designed to study the dynamics of other signaling pathways at scale.

Materials and methods

Cell culture and reagents

NIH3T3 cells were cultured in DMEM high glucose medium with 5% fetal bovine serum, 4 mM L-glutamine, 200 U/ml penicillin and $200\mu g$/ml streptomycin at 37°C with 5% CO_2. All imaging experiments with NIH3T3 were done in starving medium consisting of DMEM high glucose supplemented with 0.5% BSA (Sigma), 200 U/ml penicillin, 200 µg/ml streptomycin and 4 mM L-Glutamine. MCF10A human mammary cells were cultured in DMEM:F12 supplemented with 5% horse serum, 20 ng/ml recombinant human EGF (Peprotech), 10 µg/ml insulin (Sigma), 0.5 µg/ml hydrocortisone (Sigma), 200 U/ml penicillin and 200 µg/ml streptomycin. All imaging experiments with MCF10A were done in starving medium consisting in DMEM:F12 supplemented with 0.3% BSA, 0.5 µg/ml hydrocortisone, 200 U/ml penicillin and 200 µg/ml streptomycin. For growth factor stimulations, we used human EGF (AF-100, Peprotech) and human basic FGF (F0291, Sigma). Chemical perturbations were done with SU-5402 (SML0443, Sigma), RAF709 (HY-100510, Lucerna Chem), U0126 (S1102, Selleck chemicals, Lubio), SCH772984 (HY-50846, Lucerna-Chem), SL0101 (559285, Sigma), Cyclosporine A (10-1119, Lucerna-chem) and Ionomycin (sc-3592, Santa Cruz). Selection of the cells post transfection was done using Puromycin (P7255, Sigma), Blasticidin S HCl (5502, Tocris) and Hygromycin B (sc-29067, Lab Force).

Plasmids and stable cell line generation

The optoFGFR construct was a gift from Won Do Heo (Addgene plasmid # 59776) (Kim et al. 2014). It consists of the myristoylated FGFR1 cytoplasmic region fused with the PHR domain of the cryptochrome2 and tagged with mCitrine. It was cloned in a lentiviral backbone for stable cell line generation. A modified version of the optoFGFR tagged with the red fluorophore mScarlet (Bindels et al. 2017) was cloned in a PiggyBac plasmid pPBbSr2-MCS (blasticidin resistance), a gift from Kazuhiro Aoki. The optoSOS construct is a modified version of the tRFP-SSPB-SOScat-P2A-iLID-CAAX (Addgene plasmid #86439) (Johnson et al. 2017), in which we replaced the tRFP by mCitrine. The construct was cloned in the pPB3.0.Puro, an improved PiggyBac plasmid generated in our lab with puromycin resistance. The ERK-KTR-mRuby2 and ERK-KTR-mTurquoise2 reporters were generated by fusing the ERK Kinase Translocation Reporter (ERK-KTR) (Regot et al. 2014) with mRuby2 (Lam et al. 2012) or mTurquoise2 (Goedhart et al. 2012). The nuclear marker H2B-miRFP703 is a fusion of the human H2B clustered histone 11 (H2BC11) with the monomeric near-infrared fluorescent protein miRFP703 (Shcherbakova et al. 2016) (Addgene plasmid #80001). ERK-KTR-mRuby2, ERK-KTR-mTurquoise2 and H2B-miRFP703 were cloned in the PiggyBac plasmids pPB3.0.Hygro, pSB-HPB (gift of David Hacker, EPFL, (Balasubramanian et al. 2016)) and pPB3.0.Blast, respectively. All constructs in PiggyBac plasmids were co-transfected with the helper plasmid expressing the transposase (Yusa et al. 2011) for stable insertion using the jetPEI (Polyplus) transfection reagent for NIH3T3 cells or FuGene (Promega) transfection reagent for

MCF10A cells. After antibiotic selection, NIH3T3 cells were FACS-sorted to generate
stable cell lines homogeneously expressing the biosensors. In the case of MCF10A
cells, clones with uniform biosensor expression were isolated. To generate ErbB2
overexpressing MCF10A cells, lentiviral transduction using a pHAGE-ERBB2
construct (a gift from Gordon Mills & Kenneth Scott, Addgene plasmid #116734, (Ng
et al. 2018)) was performed in the presence of 8 µg/ml polybrene (TR1003, Sigma) in
cells already expressing H2B-miRFP703 and ERK-KTR-mTurquoise2. Cells were
further selected with 5 µg/ml puromycin.
805
Live imaging of ERK dynamics
NIH3T3 cells were seeded in 96 well 1.5 glass bottom plates (Cellvis) coated with 10
µg/ml Fibronectin (Huber lab) using 1.5×10^3 cells/well and incubated for 24 hours.
MCF10A cells were seeded in 24-well 1.5 glass bottom plates (Cellvis) coated with 5
µg/ml Fibronectin (Huber lab) at 1×10^5 cells/well and incubated for 48 hours. NIH3T3
cells were washed with PBS and incubated in starving medium for 4 hours in the dark
before starting the experiment. MCF10A cells were starved for 7 hours before starting
the experiments. In experiments involving drug perturbations, cells were incubated for
2 hours (or 1 hour in MCF10A experiments) with the inhibitor(s). Imaging was
performed with an epifluorescence Eclipse Ti inverted fluorescence microscope
(Nikon) using a Plan Apo air 20x (NA 0.8) objective. Nikon Perfect Focus System
(PFS) was used to keep cells in focus throughout the experiment. Illumination was
done with a SPECTRA X light engine (Lumencor) with the following filters (Chroma):
mTurquoise2: 440 nm LED, 470lp, 69308 CFP/YFP/mCherry-ET, CFP 458-482;
mCitrine: 508 nm LED, ET500/20x, 69308bs, ET535/30m; mRuby2 and mCherry: 555
nm LED, ET575/25x, 69008bs, 59022m, miRFP703: 640 nm LED, ET640/30x,
89100bs Sedat Quad, 84101m Quad. Images were acquired with an Andor Zyla 4.2
plus camera at a 16-bit depth. Image acquisition and optogenetic stimulation were
controlled with the NIS-Element JOBS module. For NIH3T3 experiments, ERK-KTR-
mRuby2 and H2B-miRFP703 were acquired at 1-minute interval and 470 nm light
inputs were delivered at specific frequencies and intensities (see below). MCF10A
image acquisition was performed at 5-minute time resolution. Growth factor
stimulations were done by manually pipetting EGF and bFGF during the experiment.
We used mCitrine intensity to quantify the expression level of the optogenetic
constructs. However, as mCitrine excitation leads to optoFGFR or optoSOS activation,
we acquired one frame with the ERK-KTR-mRuby2, the H2B-miRFP703 and the
mCitrine-tagged optoFGFR or optoSOS only at the end of each NIH3T3 experiments.
All experiments were carried on at 37°C with 5% CO_2.
834
Optogenetic stimulation
Light stimulations were delivered with a 470 nm LED light source that was hardware-
triggered by the camera to generate light pulses of reproducible duration. Light
stimulations of defined intensity and duration were programmed to be automatically
delivered at specific timepoints. To define the dose of light received by the cells, we
measured the 470 nm light intensity at the focal plane using an optical power meter

(X-Cite Power Meter, Lumen Dynamics Group) and converted this value to a power
density as
843

844
$$Light\ power\ density = Light\ intensity \times \frac{1}{\pi \times \left(\frac{FN}{2 \times Magnification}\right)^2} \left[\frac{mW}{cm^2}\right]$$

846 with FN = 18 mm. The obtained value was then multiplied by the duration of the pulse
847 to obtain the dose of light received by the cells for each light pulse.
848

849
$$Light\ dose\ (D) = Light\ power\ density \times Pulse\ duration = \left[\frac{mW \times s}{cm^2}\right] = \left[\frac{mJ}{cm^2}\right]$$

For stimulation of the optoFGFR cells, the 470 nm LED intensity was limited to a low
dose by combining a ZET470/10x filter and a ND filter 5% (Chroma). Transient
stimulations were done with a single pulse, while sustained stimulations were done
with single pulses delivered every 2 minutes. For stimulation of the optoSOS cells, we
used the 470 nm LED with a ET470/24x filter (no ND filter). Transient stimulations
were done with 5 pulses repeated at 20-second intervals, while sustained stimulations
were done using 5 pulses repeated at 20-second intervals, delivered every 2 minutes.
858

Figures	System	Power density	Pulse duration	Dose	Stimulation pattern
1B,C, Appendix S1A, Appendix S1D	optoFGFR	180 mW/cm^2	1 x 100 ms	18 mJ/cm^2	transient
Appendix S1B,C	optoFGFR (mScarlet)	> 180 mW/cm^2	1 x 100 ms	> 18 mJ/cm^2	transient
2A,B	optoFGFR	variable	variable	variable	transient
2C, EV1A,B	optoFGFR	180 mW/cm^2	1 x 100 ms	18 mJ/cm^2	variable
2D,E, 3C,E, EV1K, 4B,D-F, EV2C,D, EV3A-D, 5E, EV4C, 6B-D,F-H, EV5A,B,D-F	optoFGFR	180 mW/cm^2	1 x 100 ms	18 mJ/cm^2	sustained
2F	optoFGFR	variable	1 x 100 ms	variable	sustained
5B	optoSOS	6 W/cm^2	variable x 100 ms (20 sec interval)	0.6 J/cm^2	transient
EV4A	optoSOS	variable	variable x 100ms (20-sec interval)	variable	transient
5C	optoSOS	6 W/cm^2	5 x 100 ms (20-	0.6 J/cm^2	variable

			sec interval)		
5D	optoSOS	variable	5 x 100 ms (20-sec interval)	variable	sustained
5E-G, EV4B-G, 6B-D, EV5A	optoSOS	6 W/cm^2	5 x 100 ms (20-sec interval)	0.6 J/cm^2	sustained

TIRF imaging of optoFGFR dynamics

Cells were seeded at a density of 1 x 10^3 per well in 96 well 1.5 glass bottom plates
(Cellvis) coated with 10 µg/ml Fibronectin (Huber lab) and incubated for 24 hours at
37°C with 5% CO_2. Before imaging, cells were washed with PBS and incubated in
starving medium for 4 hours in the dark. Imaging was performed with an
epifluorescence Eclipse Ti inverted fluorescence microscope (Nikon) using a CFI
Apochromat TIRF 100x oil (NA 1.49). Images were acquired with an Andor Zyla 4.2
plus camera at a 16-bit depth. TIRF images were acquired with a 561 nm laser using
a ET575/25 filter in front of the ZT488/561rpc (Chroma) to prevent nonspecific
activation of the CRY2. MetaMorph software (Universal Imaging) was used for
acquisition. TIRF images of the optoFGFR-mScarlet were acquired at a 20-second
interval. Optogenetic stimulation was done using a 470 nm LED (SPECTRA X,
Lumencor) (Appendix Figure S1B). All experiments were carried on at 37°C with 5%
CO_2.

Image processing pipeline

Nuclear segmentation was done in CellProfiler 3.0 (McQuin et al. 2018) using a
threshold-based approach of the H2B channel. In the case of MCF10A cells, nuclear
segmentation was preceded by prediction of nuclear probability using a random forest
classifier based on different pixel features available in Ilastik software (Berg et al.
2019). To measure the ERK-KTR fluorescence in the cytosol, the nuclear mask was
first expanded by 2 pixels to exclude the blurred edges of the nucleus. The new mask
was then further expanded by 4 pixels in a threshold-based manner to obtain a "ring"
area corresponding to the cytoplasmic ERK-KTR. ERK activity was obtained by
calculating the ratio between the average cytosolic pixel intensity and the average
nuclear pixel intensity. Single-cell tracking was done on nuclear centroids with
MATLAB using µ-track 2.2.1 (Jaqaman et al. 2008). The final images containing the
ERK-KTR-mRuby2, H2B-miRFP703 and the optoFGFR-mCitrine (or optoSOS-
mCitrine) channels were processed using the same CellProfiler settings as the time
lapse images. Intensity of the mCitrine was extracted under the ERK-KTR cytoplasmic
mask and used to classify cells into low or high expressors in a threshold-based
manner. For optoFGFR-evoked ERK responses, the threshold was defined empirically
to separate oscillatory and non-oscillatory ERK responses (low < -1.75 (log10 mCitrine
intensity) < high). For optoSOS-evoked ERK responses, the threshold was defined
empirically to separate cells with low or high ERK response amplitudes (low < -1.25

(log10 mCitrine intensity) < high). The same thresholds were kept across experiments
to compare low and high expressors.
The optoFGFR-mScarlet dimers/oligomers were segmented using the pixel
classification module from Ilastik (Berg et al. 2019). OptoFGFR dimers, cell
background and trafficking vesicles were manually annotated on images before and
after the light stimulation. A probability map of the optoFGFR dimers classification was
exported as TIFF for each frame. We then computed the mean of pixel intensities from
the binarized mask obtained with Ilastik using Fiji (Appendix Figure S1C).
903
Quantification of ERK activity
We wrote a set of custom R scripts to automatically calculate the ERK-KTR
cytoplasmic/nuclear ratio as a proxy for ERK activity for each single-cell, link single-
cell ERK responses with the corresponding optoFGFR/optoSOS intensity value and
export the corresponding ERK single-cell trajectories. For NIH3T3 data, outliers in
ERK single-cell trajectories were removed using a clustering-based approach
(https://github.com/pertzlab/Outlier_app). Trajectories with an ERK activity higher than
0.8 or lower than 0.2 before stimulation, above 1.6 during the whole experiment or
displaying single time point spiking values were removed. For MCF10A data,
trajectories with an ERK activity above 2 or shorter than 90% of the total experiment
duration were removed. All the R codes used for further analysis are available as
supplementary information (see Data availability section). Hierarchical clustering
analysis of single-cell trajectories (Figure 2D, EV3F,G, EV4B) was done using Time
Course Inspector (Dobrzyński et al. 2019).
918
Modeling
The model for the EGF and light stimulated ERK cascade is a kinetic model,
representing the EGF receptor, the inter-cellular proteins (RAS, RAF, MEK, ERK) as
well as a negative feedback (NFB) from ERK to RAF and the inactivation of the EGF
receptor in the form of endocytosis (Figure 3D). We explicitly modelled the ERK-KTR
readout through nuclear and cytosolic KTR. The initial fraction of cytosolic KTR is
estimated from the data through the parameter ktr_{init}. The KTR readout $Y(t)$ was taken
to be the ratio of cytosolic KTR over nuclear KTR with additive Gaussian noise
927
928 $$Y(t) = \frac{KTR}{KTR^*} + \epsilon$$
929
930 $$\epsilon \sim Normal(0, \sigma^2)$$
931
where the variance of the measurement noise σ^2 was estimated from the data.
Appendix Table S1 shows all modelled species, their notation used for the equation,
as well as the initial values. We assume that in the beginning of the experiment, all
species are in the inactive form, reflecting the fact that the cells have been starved.
The total concentrations of all species have been normalized to 1. The model
equations are shown in Appendix Table S2. The phosphorylation events are modelled

with Michaelis-Menten kinetics. The NFB is modelled through the modeling species
NFB and its "active" version NFB^* which affects the dephosphorylation rate of RAF
linearly. The activation, endocytosis, and recycling of the EGF receptor is modelled
linearly. The model parameters are described in Appendix Table S3. For the modeling
of the two smaller models (without feedback (Figure EV1E) or without endocytosis
(Figure EV1H)), we set the corresponding parameters (k_{nfb} and $r_{2,3}$) to zero.
For the parameter inference, we used a Nested Sampling algorithm as described in
(Mikelson and Khammash 2020). The inference was performed on the ETH High-
performance Cluster Euler and was done using the parallel implementation on 48
cores. The algorithm was run for 24 hours or until the algorithm stopped because the
termination criterion Δ_{LFNS} (see (Mikelson and Khammash 2020) for details) was $-\infty$.
As prior distributions, we chose for all parameters non-informative log-uniform priors
between 10^{-5} and 10^5, except for ktr_{init} for which we chose a uniform prior on the
interval [0, 1] and for σ for which we chose a log-uniform prior between 10^{-5} and 1.
Predictive distributions can be found on Figure 3F,G, EV1F,G,I,J.
953
RNAi perturbation screen
We used Ingenuity Pathway Analysis (IPA, Qiagen) to select proteins directly
interacting with ERK, MEK, RAF, RAS and FGFR, that are known to be expressed in
NIH3T3 cells using a proteomics approach (Schwanhäusser et al. 2011; Jensen et al.
2009) (Appendix Table S4). We then imported this protein list in STRING (Jensen et
al. 2009) to generate an interaction network with a minimum interaction score of 0.4.
The final interactome was manually modified to display the protein names to facilitate
the readout (Figure 4A). We targeted these selected proteins with RNA interference,
using the siPOOL technology (one siPOOL containing a mix of 30 siRNAs targeting
the same gene (Hannus et al. 2014), sequences available in the Data availability
section). We arranged the siPOOLs in a 96 well plate format (in columns 2-5 and 8-
11, one well per siPOOL) with the non-targeting siRNA (*CTRL*) and the positive control
(mix of 5 nM siPOOL against *ERK1* and 5 nM siPOOL against *ERK2*) placed
alternately in columns 1, 6, 7 and 12. Cells were reverse transfected using RNAiMAX
(Thermofisher, 13778150) following the recommended siPOOL transfection protocol
(https://sitoolsbiotech.com/protocols.php). OptoFGFR-expressing cells were
transfected with 10 nM of siPOOL in a 96 well 1.5 glass bottom plate (Cellvis) coated
with 10 µg/ml Fibronectin (Huber Lab) at 0.3×10^3 cells/well density and incubated for
72 hours at 37°C and 5% CO_2. For the imaging, the 96 well plate was divided into 15
sub-experiments, each sub-experiment consisting of a negative control well, a positive
control well and 4 wells with different siPOOLs. We selected 2 FOVs per well and
programmed the microscope to run the 15 experiments sequentially, acquiring the
ERK-KTR-mRuby2 and the H2B-miRFP703 channels with a 1-minute interval,
stimulating the cells with sustained optoFGFR input (2-minute intervals, D = 18
mJ/cm^2), and acquiring a final frame with ERK-KTR-mRuby2, H2B-miRFP703 and
optoFGFR-mCitrine (Figure 4B,D-F, EV2C,D, EV3A-D, 6F-H, EV5B). For the optoSOS
system, we limited the perturbation screen to targets acting below RAS (Figure 5F,G,
EV4D-G). Stimulations were done with sustained optoSOS input (5 repeated pulses

at 2-minute intervals, D = 0.6 J/cm^2). For EGF experiments, cells were stimulated with

1 ng/ml EGF at t = 5 minutes (Figure EV3E-G).

Real-time qPCR

Cells were transfected with different concentrations of siPOOL in a 24 well plate at 5

x 10^3 cells/well density and incubated at 37°C with 5% CO_2 for 72 hours before RNA

isolation. Reverse transcription was done with the ProtoScript II reverse transcriptase

kit (Bioconcept, M0368L). Real-time qPCR reactions were run using the MESA Green

pPCR MasterMix Plus for SYBR Green assay (Eurogenetec, RT-SY2X-03+WOU) on

the Rotor-Gen Q device (Qiagen). Each sample was tested in triplicate. Expression

level of the gene of interest was calculated using the 2$^{-\Delta\Delta Ct}$ method with *GAPDH*

expression level as internal control (Figure EV2A). The following primers were used

for the RT-qPCR reaction (designed with the Real-time PCR (TaqMan) Primer and

Probes Design Tool from GenScript).

Target	Forward sequences	Reverse sequences
ERK1	5'-GGTTGTTCCCAAATGCTGACT-3'	5'-CAACTTCAATCCTCTTGTGAGGG-3'
ERK2	5'-TCCGCCATGAGAATGTTATAGGC-3'	5'-GGTGGTGTTGATAAGCAGATTGG-3'
MEK1	5'-AAGGTGGGGGAACTGAAGGAT-3'	5'-CGGATTGCGGGTTTGATCTC-3'
MEK2	5'-GTTACCGGCACTCACTATCAA C-3'	5'-CCTCCAGCCGCTTCCTTTG-3'
GAPDH	5'-ACCCAGAAGACTGTGGATGG-3'	5'-TCAGCTCAGGGATGACCTTG-3'

Immunoblotting

Cells were transfected with 10 nM siPOOL in 6 well plates at 6 x 10^4 cells/well density

and incubated at 37°C with 5% CO_2 for 72 hours. Cells were lysed in a buffer

containing 10 mM Tris HCl, 1 mM EDTA and 1% SDS. Protein concentration was

determined with the BCA^{TM} protein assay kit (Thermofisher, 23227). Home cast 10%

SDS gels or Novex 4%-20% 10 well Mini Gels (Thermofisher, XP04200) were used

for SDS page. Transfer was done using PVDF membranes and a Trans-Blot SD Semi-

Dry Electrophoretic Transfer Cell (Bio-Rad). Imaging was done with an Odyssey

Fluorescence scanner (Li-COR) (Figure 4C, EV2B). The following primary antibodies

were used: anti-total ERK (M7927, Sigma), anti-MEK1 (ab32091, Abcam), anti-MEK2

(ab32517, Abcam), anti-BRAF (sc-5284, Santa Cruz), anti-CRAF (9422S, Cell

Signaling Technology), anti-SOS1 (610096, Biosciences), anti-GRB2 (PA5-17692,

Invitrogen) and anti-RSK2 (sc-9986, Santa Cruz). Anti-GAPDH (sc-32233, Santa

Cruz) or anti-Actin (A2066, Merck) was used as protein of reference. For the

secondary antibodies, we used the IRDye 680LT donkey anti-mouse IgG (926-68022,

Li-COR), IRDye 800CW goat anti-mouse (926-32210, Li-COR) and IRDye 800CW

donkey anti-rabbit (926-32213, Li-COR). Protein quantification was done with the

Image Studio^{TM} Lite software.

1016

Time-series feature extraction

1017

We used custom scripts to extract features of ERK responses to transient optoFGFR
input (Figure 2B, EV1A,B), sustained GF input (Figure EV1C,D) and transient
optoSOS input (EV4A). The maximum peak (maxPeak) is the absolute value of the
highest ERK activity in the trajectory. To estimate the full width at half maximum
(FWHM), we first removed the baseline of the trajectories and increased their sampling
frequency by a factor 30 with spline interpolation. On the resulting trajectory, we
applied a "walk" procedure to quantify the FWHM. In this method, a pointer walks left
and right (*i.e.* opposite and along the direction of time respectively) from the maximum
point of the trajectory. The pointer stops whenever the half maximum value is crossed.
Both stops define a left and a right border, the time difference between these 2-border
time-points gives the FWHM. To avoid reporting aberrant FWHM values in cases
where a peak cannot be clearly defined, we excluded FWHM calculation for
trajectories where the fold change between the baseline (mean activity before
stimulation) and the maximum value of the trajectory was below a threshold manually
defined. ERKpostStim is the absolute value of ERK activity extracted 9 minutes after
the last stimulation pulse to evaluate ERK adaptation. Statistical analysis (Figure
EV1A,B) was done by comparing all conditions to the 20-minute interval stimulation
patterns with a Wilcoxon test using the FDR p-value correction (NS: non-significant,
*<0.01, **<0.001, ***<0.0001, ****<0.00001).

To evaluate ERK phenotypes under siRNA perturbations in response to sustained
optoFGFR or optoSOS input (Figure 4D, EV3A, 5F, EV4F), we extracted the baseline
(average ERK activity on 5 timepoints before stimulation), the maxPeak (maximum
ERK activity within a 10-minute time window following the start of the stimulation) and
the ERKpostStim (ERK activity at a fixed timepoint post-stimulation ($t_{optoFGFR}$ = 42 min
and $t_{optoSOS}$ = 40 min)) from 3 technical replicates. To avoid heterogeneity due to
differences in optogenetic expression, we focused our analysis on cells with high
optogenetic expression. The obtained baseline, maxPeak and ERKpostStim for each
siRNA perturbation was z-scored to the non-targeting siRNA (*CTRL*). Non-significant
results were manually set to grey. Statistical analysis was done by comparing each
perturbation to the control with a Wilcoxon test using the FDR p-value correction (NS:
non-significant, *<0.05, **<0.005, ***<0.0005, ****<0.00005).

For the comparison of both optogenetic systems (Figure 5E, EV4C), ERK baseline
was obtained by averaging ERK activity on 5 timepoints before stimulation and ERK
maxPeak was extracted within a 10-minute time window following the start of the
stimulation. Statistical analysis was done by comparing low and high expressing cells
within and across optogenetic systems with a Wilcoxon test using the FDR p-value
correction (NS: non-significant, *<0.05, **<0.005, ***<0.0005, ****<0.00005).

To quantify the efficiency of the three MAPK inhibitors on the reduction of ERK
amplitudes under sustained high optoFGFR or optoSOS input (Figure 6), extraction of
the maxPeak was limited by the fact that several concentrations led to a full
suppression of ERK amplitudes. Therefore, we extracted ERK amplitudes at a fixed
time point following the start of the stimulation ($t_{fixed\ optoFGFR}$ = 15 minutes, $t_{fixed\ optoSOS}$ =

10 minutes). The obtained ERK amplitudes were then plotted for each concentration
for a fixed number of cells randomly selected (Figure 6B,F, EV5D). To calculate the
EC_{50} of each drug, we normalized the data by setting the mean ERK responses of the
non-treated condition to 1 and the mean ERK responses of the maximum
concentration to 0. EC_{50} then was calculated by fitting a Hill function to the mean ERK
activity of each concentration (Figure 6C,G, EV5E, Appendix Table 5-7). The
heterogeneity of ERK amplitude at the fixed time point was evaluated by computing
the normalized standard deviation of the extracted ERK activity per condition (Figure
6D,H, EV5F).
1069
Identification of ERK dynamics phenotypes using CODEX
To investigate ERK dynamics phenotypes to siRNA perturbations, we first trained a
convolutional neural network (CNN) to classify input ERK trajectories into any of the
siRNA-perturbed conditions (Figure EV3C). For this purpose, we used a CNN
architecture composed of 4 1D-convolution layers with 20 kernels of size 5, followed
by a convolution layer with 20 kernels of size 3 and one layer of 10 kernels of size 3.
The responses are then pooled with global average pooling to generate a vector of 10
features that is passed to a (10,63) fully connected layer for classification. Each
convolutional layer is followed by ReLU and batch normalization. The CNN was trained
to minimize the cross-entropy loss, with L2 weight penalty of $1e^{-3}$.
To identify siRNA treatments that induced a distinctive phenotype, we selected the 10
conditions for which the CNN classification precision was the highest on the validation
set (Appendix Table S4, "CODEX accuracy"). To these 10 conditions, we also added
the negative control (non-targeting siRNA (*CTRL*)). We trained a second CNN, with
the same architecture and training parameters, but limited to recognizing the 11
selected treatments to obtain a clear embedding of these hits. With this new model,
we extracted the features used for the classification of the trajectories (i.e. the input
representation after the last convolution layer) and projected them with tSNE (Python's
sklearn implementation, perplexity of 100, learning rate of 600 and 2500 iterations)
(Figure EV3D). We selected 10 prototype curves for each treatment by taking the
trajectories for which the second CNN's classification confidence (*i.e.* the probability
for the actual class of the inputs) were the highest in the validation set (Figure 4E,
"CODEX").
To visualize the ERK dynamics landscape in MCF10A WT cells and in MCF10A cells
overexpressing ErbB2, we trained one CNN for each cell line. These CNNs were
trained to recognize the drug treatment applied on cells, using single-cell ERK traces
as input. The architecture of the CNNs is the same as described previously. The only
difference lies in the number of outputs in the final fully connected layer, which were
set to the number of drug treatments. Features used for the classification of the
trajectories were then projected with tSNE (Figure 7D,H, EV5J,K).
To identify clusters gathering similar ERK dynamics (Figure 7E,F,I,J), we clustered
trajectories based on their CNN features using a partition around medoids (PAM). This
iterative algorithm is similar to K-means clustering. PAM defines the cluster centers
(i.e. the medoids) as the observed data points which minimize the median distances

to all other points in its own cluster. This makes PAM more robust to outliers than K-means which uses the average coordinates of a cluster to define its center. Representative trajectories were obtained by taking the medoids of each cluster and their four closest neighbors (Figure 7G,K). Distances between points were defined with the Manhattan distance between the scaled CNN features (zero mean and unit variance). We manually verified that these clusters captured an actual trend by visualizing trajectories in each cluster with the interactive CODEX application.

Peak detection and classification of oscillatory trajectories

The number of ERK activity peaks was calculated with a custom algorithm that detects local maxima in time series. First, we applied a short median filter to smoothen the data with a window width of 3 time points. Then, we ran a long median filter to estimate the long-term bias with a window width of 15 time points. This bias was then subtracted from the smoothed time series and we only kept the positive values. If no point in this processed trajectory was exceeding a manual threshold of 0.075, all variations were considered as noise and no peak was extracted from the trajectory. The remaining trajectories were then rescaled to [0,1]. Finally, peaks were detected as points that exceeded a threshold which was manually set to 0.1. Peaks that were found before the first stimulation or after the last stimulation were filtered out.

The classification of trajectories into oscillatory and non-oscillatory behaviors was performed after the peak detection step. Cells were called oscillatory if at least 3 peaks were detected with the peak detection procedure (Figure 4F). Statistical analysis was done using a pairwise t-test comparing each perturbation to the control for high and low levels of optoFGFR independently, with FDR p-value correction ($*<0.05$, $**<0.005$, $***<0.0005$, $****<0.00005$).

Data availability

The datasets used in this study as well as all R codes used for further analysis are available at https://data.mendeley.com/datasets/st36dd7k23/1. Source code for the inference algorithm, model files and results are available at https://github.com/Mijan/LFNS_optoFGFR.

Acknowledgements

This work was supported by SystemsX.ch, Swiss Cancer League and Swiss National Science Foundation grants to Olivier Pertz, by the H2020-MSCA-IF, project No. 89631 - NOSCAR to Agne Frismantiene and by the European Union's Horizon 2020 and innovation program under grant agreement No. 730964 (TRANSVAC project) to Mustafa Khammash. We thank Won Do Heo for sharing the optoFGFR plasmid, Kazuhiro Aoki for sharing the pPBbSr2-MCS plasmid, and David Hacker for sharing the pSB-HPB plasmid. We thank the Microscopy Imaging Center of the University of Bern for its support.

Authors contribution

1148

O.P. and C.D. designed the study. C.D. developed the optogenetic systems and
imaging pipelines. CD performed the experiment and image analysis on NIH3T3. A.F
and P.A.G. performed the experiments and image analysis on MCF10A. M.D.
developed the processing pipelines. C.D processed the data. C.D., M.-A.J., A.F and
P.A.G. analyzed the data. M.-A.J. conducted the CNN analysis. J.M. performed
mathematical modeling. O.P and M.K. supervised the work. O.P., C.D. and J.M. wrote
the paper.
1156

1157 **Conflict of interest**

1158

1159 The authors declare having no conflict of interest.

References

Albeck, J.G., Mills, G.B. and Brugge, J.S. 2013. Frequency-modulated pulses of ERK activity transmit quantitative proliferation signals. *Molecular Cell* 49(2), pp. 249–261.

Arkun, Y. and Yasemi, M. 2018. Dynamics and control of the ERK signaling pathway: Sensitivity, bistability, and oscillations. *Plos One* 13(4), p. e0195513.

Arteaga, C.L. and Engelman, J.A. 2014. ERBB receptors: from oncogene discovery to basic science to mechanism-based cancer therapeutics. *Cancer Cell* 25(3), pp. 282–303.

Avraham, R. and Yarden, Y. 2011. Feedback regulation of EGFR signalling: decision making by early and delayed loops. *Nature Reviews. Molecular Cell Biology* 12(2), pp. 104–117.

Balasubramanian, S., Wurm, F.M. and Hacker, D.L. 2016. Multigene expression in stable CHO cell pools generated with the piggyBac transposon system. *Biotechnology Progress* 32(5), pp. 1308–1317.

Belov, A.A. and Mohammadi, M. 2012. Grb2, a double-edged sword of receptor tyrosine kinase signaling. *Science Signaling* 5(249), p. pe49.

Benedetti, L., Barentine, A.E.S., Messa, M., Wheeler, H., Bewersdorf, J. and De Camilli, P. 2018. Light-activated protein interaction with high spatial subcellular confinement. *Proceedings of the National Academy of Sciences of the United States of America* 115(10), pp. E2238–E2245.

Berg, S., Kutra, D., Kroeger, T., et al. 2019. ilastik: interactive machine learning for (bio)image analysis. *Nature Methods* 16(12), pp. 1226–1232.

Bindels, D.S., Haarbosch, L., van Weeren, L., et al. 2017. mScarlet: a bright monomeric red fluorescent protein for cellular imaging. *Nature Methods* 14(1), pp. 53–56.

Blum, Y., Mikelson, J., Dobrzyński, M., et al. 2019. Temporal perturbation of ERK dynamics reveals network architecture of FGF2/MAPK signaling. *Molecular Systems Biology* 15(11), p. e8947.

Blüthgen, N. and Legewie, S. 2013. Robustness of signal transduction pathways. *Cellular and Molecular Life Sciences* 70(13), pp. 2259–2269.

Cargnello, M. and Roux, P.P. 2011. Activation and function of the MAPKs and their substrates, the MAPK-activated protein kinases. *Microbiology and Molecular Biology Reviews* 75(1), pp. 50–83.

Chardin, P., Camonis, J.H., Gale, N.W., et al. 1993. Human Sos1: a guanine nucleotide exchange factor for Ras that binds to GRB2. *Science* 260(5112), pp. 1338–1343.

Cherfils, J. and Zeghouf, M. 2013. Regulation of small GTPases by GEFs, GAPs, and GDIs. *Physiological Reviews* 93(1), pp. 269–309.

Cullen, P.J. and Lockyer, P.J. 2002. Integration of calcium and Ras signalling. *Nature Reviews. Molecular Cell Biology* 3(5), pp. 339–348.

Dobrzyński, M., Jacques, M.-A. and Pertz, O. 2019. Mining single-cell time-series datasets with Time Course Inspector. *Bioinformatics*.

1212   Douville, E. and Downward, J. 1997. EGF induced SOS phosphorylation in PC12 cells involves P90
1213   RSK-2. *Oncogene* 15(4), pp. 373–383.

1214

1215   Duan, L., Hope, J., Ong, Q., et al. 2017. Understanding CRY2 interactions for optical control of
1216   intracellular signaling. *Nature Communications* 8(1), p. 547.

1217

1218   Erickson, K.E., Rukhlenko, O.S., Posner, R.G., Hlavacek, W.S. and Kholodenko, B.N. 2019. New
1219   insights into RAS biology reinvigorate interest in mathematical modeling of RAS signaling. *Seminars*
1220   *in Cancer Biology* 54, pp. 162–173.

1221

1222   Ferrell, J.E. and Bhatt, R.R. 1997. Mechanistic studies of the dual phosphorylation of mitogen-
1223   activated protein kinase. *The Journal of Biological Chemistry* 272(30), pp. 19008–19016.

1224

1225   Fritsche-Guenther, R., Witzel, F., Sieber, A., et al. 2011. Strong negative feedback from Erk to Raf
1226   confers robustness to MAPK signalling. *Molecular Systems Biology* 7, p. 489.

1227

1228   Gerosa, L., Chidley, C., Fröhlich, F., et al. 2020. Receptor-Driven ERK Pulses Reconfigure MAPK
1229   Signaling and Enable Persistence of Drug-Adapted BRAF-Mutant Melanoma Cells. *Cell Systems*
1230   11(5), p. 478–494.e9.

1231

1232   Goedhart, J., von Stetten, D., Noirclerc-Savoye, M., et al. 2012. Structure-guided evolution of cyan
1233   fluorescent proteins towards a quantum yield of 93%. *Nature Communications* 3, p. 751.

1234

1235   Hannus, M., Beitzinger, M., Engelmann, J.C., et al. 2014. siPools: highly complex but accurately
1236   defined siRNA pools eliminate off-target effects. *Nucleic Acids Research* 42(12), pp. 8049–8061.

1237

1238   Huang, C.-Y. and Tan, T.-H. 2012. DUSPs, to MAP kinases and beyond. *Cell & bioscience* 2(1), p.
1239   24.

1240

1241   Huotari, J. and Helenius, A. 2011. Endosome maturation. *The EMBO Journal* 30(17), pp. 3481–3500.

1242

1243   Jacques, M.A., Dobrzyński, M., Gagliardi, P.A., Sznitman, R. and Pertz, O. 2021. CODEX, a neural
1244   network approach to explore signaling dynamics landscapes. *Molecular Systems Biology* 17(4), p.
1245   e10026.

1246

1247   Jaqaman, K., Loerke, D., Mettlen, M., et al. 2008. Robust single-particle tracking in live-cell time-lapse
1248   sequences. *Nature Methods* 5(8), pp. 695–702.

1249

1250   Jensen, L.J., Kuhn, M., Stark, M., et al. 2009. STRING 8--a global view on proteins and their
1251   functional interactions in 630 organisms. *Nucleic Acids Research* 37(Database issue), pp. D412-6.

1252

1253   Johnson, H.E., Goyal, Y., Pannucci, N.L., Schüpbach, T., Shvartsman, S.Y. and Toettcher, J.E. 2017.
1254   The spatiotemporal limits of developmental erk signaling. *Developmental Cell* 40(2), pp. 185–192.

1255

1256   Junttila, M.R., Li, S.-P. and Westermarck, J. 2008. Phosphatase-mediated crosstalk between MAPK
1257   signaling pathways in the regulation of cell survival. *The FASEB Journal* 22(4), pp. 954–965.

1258

1259   Kanodia, J., Chai, D., Vollmer, J., et al. 2014. Deciphering the mechanism behind Fibroblast Growth
1260   Factor (FGF) induced biphasic signal-response profiles. *Cell Communication and Signaling* 12, p. 34.

1261

1262   Kholodenko, B.N. 2000. Negative feedback and ultrasensitivity can bring about oscillations in the
1263   mitogen-activted protein kinase cascades. *Eur. J. Biochem* 267, pp. 1583–1588.

1264

1265 Kholodenko, B.N., Hancock, J.F. and Kolch, W. 2010. Signalling ballet in space and time. *Nature*
1266 *Reviews. Molecular Cell Biology* 11(6), pp. 414–426.

1267

1268 Kim, N., Kim, J.M., Lee, M., Kim, C.Y., Chang, K.-Y. and Heo, W.D. 2014. Spatiotemporal control of
1269 fibroblast growth factor receptor signals by blue light. *Chemistry & Biology* 21(7), pp. 903–912.

1270

1271 Kiyatkin, A., van Alderwerelt van Rosenburgh, I.K., Klein, D.E. and Lemmon, M.A. 2020. Kinetics of
1272 receptor tyrosine kinase activation define ERK signaling dynamics. *Science Signaling* 13(645).

1273

1274 Kochańczyk, M., Kocieniewski, P., Kozłowska, E., et al. 2017. Relaxation oscillations and hierarchy of
1275 feedbacks in MAPK signaling. *Scientific Reports* 7, p. 38244.

1276

1277 Koseska, A. and Bastiaens, P.I.H. 2020. Processing temporal growth factor patterns by an epidermal
1278 growth factor receptor network dynamically established in space. *Annual Review of Cell and*
1279 *Developmental Biology* 36, pp. 359–383.

1280

1281 Kupzig, S., Walker, S.A. and Cullen, P.J. 2005. The frequencies of calcium oscillations are optimized
1282 for efficient calcium-mediated activation of Ras and the ERK/MAPK cascade. *Proceedings of the*
1283 *National Academy of Sciences of the United States of America* 102(21), pp. 7577–7582.

1284

1285 Lake, D., Corrêa, S.A.L. and Müller, J. 2016. Negative feedback regulation of the ERK1/2 MAPK
1286 pathway. *Cellular and Molecular Life Sciences* 73(23), pp. 4397–4413.

1287

1288 Lam, A.J., St-Pierre, F., Gong, Y., et al. 2012. Improving FRET dynamic range with bright green and
1289 red fluorescent proteins. *Nature Methods* 9(10), pp. 1005–1012.

1290

1291 Langlois, W.J., Sasaoka, T., Saltiel, A.R. and Olefsky, J.M. 1995. Negative feedback regulation and
1292 desensitization of insulin- and epidermal growth factor-stimulated p21ras activation. *The Journal of*
1293 *Biological Chemistry* 270(43), pp. 25320–25323.

1294

1295 Lavoie, H., Gagnon, J. and Therrien, M. 2020. ERK signalling: a master regulator of cell behaviour,
1296 life and fate. *Nature Reviews. Molecular Cell Biology* 21(10), pp. 607–632.

1297

1298 Lemmon, M.A., Freed, D.M., Schlessinger, J. and Kiyatkin, A. 2016. The dark side of cell signaling:
1299 positive roles for negative regulators. *Cell* 164(6), pp. 1172–1184.

1300

1301 McQuin, C., Goodman, A., Chernyshev, V., et al. 2018. CellProfiler 3.0: Next-generation image
1302 processing for biology. *PLoS Biology* 16(7), p. e2005970.

1303

1304 Mikelson, J. and Khammash, M. 2020. Likelihood-free nested sampling for parameter inference of
1305 biochemical reaction networks. *PLoS Computational Biology* 16(10), p. e1008264.

1306

1307 Nakakuki, T., Birtwistle, M.R., Saeki, Y., et al. 2010. Ligand-specific c-Fos expression emerges from
1308 the spatiotemporal control of ErbB network dynamics. *Cell* 141(5), pp. 884–896.

1309

1310 Ng, P.K.-S., Li, J., Jeong, K.J., et al. 2018. Systematic functional annotation of somatic mutations in
1311 cancer. *Cancer Cell* 33(3), p. 450–462.e10.

1312

1313 Ornitz, D.M. and Itoh, N. 2015. The Fibroblast Growth Factor signaling pathway. *Wiley*
1314 *interdisciplinary reviews. Developmental biology* 4(3), pp. 215–266.

1315

1316   Orton, R.J., Sturm, O.E., Vyshemirsky, V., Calder, M., Gilbert, D.R. and Kolch, W. 2005.
1317   Computational modelling of the receptor-tyrosine-kinase-activated MAPK pathway. *The Biochemical*
1318   *Journal* 392(Pt 2), pp. 249–261.
1319
1320   Patterson, K.I., Brummer, T., O'Brien, P.M. and Daly, R.J. 2009. Dual-specificity phosphatases:
1321   critical regulators with diverse cellular targets. *The Biochemical Journal* 418(3), pp. 475–489.
1322
1323   Purvis, J.E. and Lahav, G. 2013. Encoding and decoding cellular information through signaling
1324   dynamics. *Cell* 152(5), pp. 945–956.
1325
1326   Rauen, K.A. 2013. The RASopathies. *Annual Review of Genomics and Human Genetics* 14, pp. 355–
1327   369.
1328
1329   Regot, S., Hughey, J.J., Bajar, B.T., Carrasco, S. and Covert, M.W. 2014. High-sensitivity
1330   measurements of multiple kinase activities in live single cells. *Cell* 157(7), pp. 1724–1734.
1331
1332   Ryu, H., Chung, M., Dobrzyński, M., et al. 2015. Frequency modulation of ERK activation dynamics
1333   rewires cell fate. *Molecular Systems Biology* 11(11), p. 838.
1334
1335   Saha, M., Carriere, A., Cheerathodi, M., et al. 2012. RSK phosphorylates SOS1 creating 14-3-3-
1336   docking sites and negatively regulating MAPK activation. *The Biochemical Journal* 447(1), pp. 159–
1337   166.
1338
1339   Sakaue-Sawano, A., Kurokawa, H., Morimura, T., et al. 2008. Visualizing spatiotemporal dynamics of
1340   multicellular cell-cycle progression. *Cell* 132(3), pp. 487–498.
1341
1342   Samatar, A.A. and Poulikakos, P.I. 2014. Targeting RAS-ERK signalling in cancer: promises and
1343   challenges. *Nature Reviews. Drug Discovery* 13(12), pp. 928–942.
1344
1345   Santos, S.D.M., Verveer, P.J. and Bastiaens, P.I.H. 2007. Growth factor-induced MAPK network
1346   topology shapes Erk response determining PC-12 cell fate. *Nature Cell Biology* 9(3), pp. 324–330.
1347
1348   Saraf, A., Oberg, E.A. and Strack, S. 2010. Molecular determinants for PP2A substrate specificity:
1349   charged residues mediate dephosphorylation of tyrosine hydroxylase by the PP2A/B' regulatory
1350   subunit. *Biochemistry* 49(5), pp. 986–995.
1351
1352   Schlessinger, J. 2000. Cell signaling by receptor tyrosine kinases. *Cell* 103(2), pp. 211–225.
1353
1354   Schmick, M., Kraemer, A. and Bastiaens, P.I.H. 2015. Ras moves to stay in place. *Trends in Cell*
1355   *Biology* 25(4), pp. 190–197.
1356
1357   Schwanhäusser, B., Busse, D., Li, N., et al. 2011. Global quantification of mammalian gene
1358   expression control. *Nature* 473(7347), pp. 337–342.
1359
1360   Shankaran, H., Ippolito, D.L., Chrisler, W.B., et al. 2009. Rapid and sustained nuclear-cytoplasmic
1361   ERK oscillations induced by epidermal growth factor. *Molecular Systems Biology* 5, p. 332.
1362
1363   Shcherbakova, D.M., Baloban, M., Emelyanov, A.V., Brenowitz, M., Guo, P. and Verkhusha, V.V.
1364   2016. Bright monomeric near-infrared fluorescent proteins as tags and biosensors for multiscale
1365   imaging. *Nature Communications* 7, p. 12405.
1366
1367   Smith, J.A., Poteet-Smith, C.E., Xu, Y., Errington, T.M., Hecht, S.M. and Lannigan, D.A. 2005.

1368   Identification of the first specific inhibitor of p90 ribosomal S6 kinase (RSK) reveals an unexpected
1369   role for RSK in cancer cell proliferation. *Cancer Research* 65(3), pp. 1027–1034.
1370
1371   Sturm, O.E., Orton, R., Grindlay, J., et al. 2010. The mammalian MAPK/ERK pathway exhibits
1372   properties of a negative feedback amplifier. *Science Signaling* 3(153), p. ra90.
1373
1374   Sugimoto, T., Stewart, S. and Guan, K.L. 1997. The calcium/calmodulin-dependent protein
1375   phosphatase calcineurin is the major Elk-1 phosphatase. *The Journal of Biological Chemistry* 272(47),
1376   pp. 29415–29418.
1377
1378   Yarden, Y. and Pines, G. 2012. The ERBB network: at last, cancer therapy meets systems biology.
1379   *Nature Reviews. Cancer* 12(8), pp. 553–563.
1380
1381   Yeung, K., Janosch, P., McFerran, B., et al. 2000. Mechanism of suppression of the
1382   Raf/MEK/extracellular signal-regulated kinase pathway by the raf kinase inhibitor protein. *Molecular*
1383   *and Cellular Biology* 20(9), pp. 3079–3085.
1384
1385   Yoo, S.-M., Cho, S.J. and Cho, Y.-Y. 2015. Molecular targeting of erks/rsk2 signaling axis in cancer
1386   prevention. *Journal of cancer prevention* 20(3), pp. 165–171.
1387
1388   Yusa, K., Zhou, L., Li, M.A., Bradley, A. and Craig, N.L. 2011. A hyperactive piggyBac transposase for
1389   mammalian applications. *Proceedings of the National Academy of Sciences of the United States of*
1390   *America* 108(4), pp. 1531–1536.
1391