Lytic transglycosylases mitigate periplasmic crowding by degrading soluble cell wall turnover products

Anna Weaver[1,2], Laura Alvarez[3], Kelly Rosch[1], Asraa Ahmed[1,6] Garrett Wang[1], Michael VanNieuwenhze[4,5], Felipe Cava[3a], and Tobias Dörr[1,2,6a]

[1] Weill Institute for Cell and Molecular Biology, Cornell University, Ithaca, NY 14853, USA
[2] Department of Microbiology, Cornell University, Ithaca NY 14853, USA
[3] The Laboratory for Molecular Infection Medicine Sweden (MIMS), Department of Molecular Biology, Umeå University, 90187 Umeå, Sweden.
[4] Department of Molecular and Cellular Biochemistry, Indiana University, Bloomington, IN 47405, USA
[5] Department of Chemistry, Indiana University, Bloomington, IN 47405, USA
[6] Cornell Institute of Host-Microbe Interactions and Disease, Cornell University, Ithaca NY 14853, USA

[a] corresponding authors: tdoerr@cornell.edu, felipe.cava@umu.se

Abstract

The peptidoglycan cell wall is a predominant structure of bacteria, determining cell shape and supporting survival in diverse conditions. Peptidoglycan is dynamic and requires regulated synthesis of new material, remodeling, and turnover – or autolysis – of old material. Despite exploitation of peptidoglycan synthesis as an antibiotic target, we lack a fundamental understanding of how peptidoglycan synthesis and autolysis intersect to maintain the cell wall. Here, we uncover a critical physiological role for a widely misunderstood class of autolytic enzymes, lytic transglycosylases (LTGs). We demonstrate that LTG activity is essential to survival by contributing to periplasmic processes upstream and independent of peptidoglycan recycling. Defects accumulate in *Vibrio cholerae* LTG mutants due to generally inadequate LTG activity, rather than absence of specific enzymes, and essential LTG activities are likely independent of protein-protein interactions, as heterologous expression of a non-native LTG rescues growth of a conditionally LTG-null mutant. Lastly, we demonstrate that soluble,

uncrosslinked, endopeptidase-dependent peptidoglycan chains, also detected in the wild-

type, are enriched in LTG mutants, and that LTG mutants are hypersusceptible to the

production of diverse periplasmic polymers. Collectively, our results suggest that LTGs

prevent toxic crowding of the periplasm with synthesis-derived peptidoglycan polymers

and contrary to prevailing models, that this autolytic function can be temporally separate

from peptidoglycan synthesis.

Introduction

The bacterial cell wall is a nearly universal feature of the bacterial cell envelope. Made

primarily of the strong and elastic polymer peptidoglycan (PG), the cell wall preserves

bacterial shape while protecting the cell from its high internal turgor pressure and external

environmental challenges[1-5]. PG synthesis begins in the cytoplasm with the generation of

lipid II, consisting of the disaccharide [N-acetylmuramic acid (MurNAc)-N-

acetylglucosamine (GlcNAc)] that is modified with a pentapeptide side stem and attached

to the lipid carrier undecaprenol. Lipid II is flipped across the cytoplasmic membrane

where the cell wall is assembled in two reactions: First, lipid II is polymerized into longer

glycan strands by glycosyltransferases (GTases) followed by crosslinking of the

elongating PG strands via their peptide side stems by transpeptidases (TPases)[3,4,6-8].

Ultimately, the combined result of GTase and TPase activities is a covalently closed

mesh-like macromolecular network called the PG sacculus.

The strength of the PG sacculus is a double-edged sword. On the one hand, the covalent

network provides a mechanical structure strong enough to withstand the high cellular

turgor pressure and stresses of a changing environment. On the other hand, it acts as a

macromolecular cage that might inhibit cellular expansion and division as well as the

insertion of crucial trans-envelope machinery (i.e., flagella and pili).[9-14] Bacteria therefore

need to couple new PG synthesis with degradation of bonds within the PG network to

simultaneously maintain the integrity of the sacculus while also making space for the

insertion of new PG.[11,15,16]

PG degradation is accomplished by several divergent enzyme classes that are often

subsumed under the term "autolysins", i.e., enzymes that cleave various bonds within PG.

Endopeptidases (EPs), for example, cleave the peptide crosslinks and are particularly

integral to the "space-making" autolytic function that permits sacculus expansion during

cell elongation; without EP activity, PG synthesis results in a thicker cell wall or integrity

failure and lysis.[17,18] Another major class of autolysins, the lytic transglycosylases (LTGs),

cleave the glycosidic linkages between disaccharide subunits within PG strands. Their

biochemistry has been exquisitely well-studied and the diversity of their structures and

mechanisms of action well characterized.[19-22] Unlike the other autolysins, the primary

cleavage mechanism of LTGs is non-hydrolytic. Rather, LTGs perform an intramolecular

cyclization of MurNac residues to generate a unique and readily identifiable signature of

their activity, anhydro-MurNAc (anhMurNAc).[20,23,24] Early characterization of PG in the

bacterial sacculus suggested that every PG strand terminates in an anhMurNAc cap, long

implicating LTGs as potential "terminases" of GTase glycan elongation.[23,25] This was only

recently confirmed empirically with the novel discovery and characterization of MltG and

its functional analogs. [26-29] *E. coli* MltG associates with active PG synthetic complexes to

release new strands from the cytoplasmic membrane (to which they are initially tethered

via undecaprenyl pyrophosphate), presumably as they emerge from GTase activity;

consequently, MltG is a strong determinant of PG strand length.[26-29] Whether association

with active GTases is a conserved characteristic of MltG functional analogues (including

some hydrolytic glycosidases[30]) has not been established, so these enzymes may be

more broadly referred to as "PG release factors" since their cleavage of membrane-bound

PG may not directly influence PG elongation. Other physiological roles assigned to LTGs

include local PG editing for insertion of PG-spanning protein complexes[10,12,31] and PG

recycling. PG recycling, the re-incorporation of PG breakdown products into the

biosynthesis cycle, starts with re-uptake of PG turnover products by the importer AmpG.[32]

AmpG specifically imports LTG breakdown products (anhMurNAc-containing fragments)

and this process can theoretically be supported by any active LTG that produces

monomeric anhydromuropeptides. Despite being well-conserved throughout many

bacterial phyla, PG recycling is not an essential process under standard growth

conditions.[33] Another underappreciated function of LTGs has emerged through the work

of our group and others wherein certain individual or combinatorial LTG mutations,

including septal LTG RlpA, results in a daughter cell separation defect.[34-37] Intriguingly,

the roles of LTGs in all these functions – PG release from the membrane, PG recycling,

insertion of PG-spanning complexes, and daughter cell separation – do not appear to be

essential to bacterial growth in the contexts studied to date.

The apparent non-essentiality of lytic transglycosylase activity would seem contradictory

to this enzyme class's broad conservation across bacterial phyla (and chloroplasts) as

well as genetic and functional redundancy within many individual species – a trait indicative of an important, conserved function.[27] Yet answering the most general physiological questions about LTGs has been severely encumbered by LTG redundancy as rarely does a single LTG mutation yield a significant, readily investigable phenotype. We recently showed that collectively, LTG activity indeed seems to be essential for growth and division.[36] The actual physiological function for the majority of LTGs, however, and the reason for their collective essentiality, has remained elusive. Here, we sought to comprehensively illuminate physiological roles for LTGs by extensively characterizing mutants of the Gram-negative pathogen *Vibrio cholerae* that are defective for most or all of the species' eight currently annotated LTGs. We find that the vast majority of LTGs is dispensable for growth in laboratory media. Minimal LTG strains are defective for growth in low-salt media and hypersensitive to accumulation of periplasmic sugar polymers. Through analysis of PG turnover products, we show that soluble PG strands accumulate in the wild type and that this is exacerbated in ∆LTG mutants and alleviated by inactivating a major PG endopeptidase. Taken together, our data suggest that lytic transglycosylase activity downstream of PG synthesis mitigates toxic periplasmic accumulation of uncrosslinked, polymeric peptidoglycan turnover products released by endopeptidases.

Results

A single LTG is necessary and sufficient for V. cholerae growth

We recently reported that a ∆6 LTG mutant (*rlpA+ mltG+ ∆mltA ∆mltB ∆mltC ∆mltD ∆mltF ∆slt70*) was viable under standard laboratory conditions and exhibited only slight morphological defects, including an increase in cell length[36]. Depletion of RlpA from this

background resulted in a lethal chaining defect, suggesting that collectively, some degree

of LTG activity is essential for *V. cholerae* growth. Since we were previously unable to

delete MltG or observe its depletion from a Δ6 LTG background, we asked whether MltG

exhibited a synthetic-lethal relationship with the other LTGs using a quantitative

insertion/disruption assay.　Briefly, we conjugated a suicide vector targeting *mltG* (or

positive or negative control loci) into WT and Δ6 LTG and quantified viable recombinants.

Surprisingly, the WT and Δ6 LTG strains both tolerated inactivation of *mltG* by the suicide

vector so long as essential DNA synthesis genes downstream of *mltG* were expressed *in*

trans to ameliorate polar effects of *mltG* disruption (Figure 1A, Figure 1 – Figure

supplement 1AB). The resulting Δ6 LTG *mltG::kan* mutant was viable in LB but failed to

grow in low-salt LB (LB without added NaCl, hereafter designated LB0N) (Figure 1 –

Figure supplement 1C), which could explain why previous attempts to inactivate *mltG* in

this　background　using　SacB-based　allelic　exchange　　(requiring　selection　on

LB0N+sucrose)　were unsuccessful[36]. Consistent with these data, we were then able to

generate a clean, viable Δ7 LTG mutant (*rlpA⁺ mltG::stop ΔmltABCDF Δslt70*) using a

MqsR toxin-based allelic exchange system.[38]　The Δ7 mutant sacculus contained only a

tenth of the AnhMurNAc residues observed in the wild type sacculus (Figure 1 – Figure

supplement 1D, Supplemental File 1), suggesting that there is significantly reduced LTG

activity in the Δ7 LTG mutant. Additionally, by placing the native copy of *rlpA* under an

arabinose-inducible promoter, we were able to conditionally deplete RlpA by growing

these strains in the absence of arabinose to observe the effects of LTG insufficiency

(Figure 1B, Figure 1 – Figure supplement 2).　Compared to Δ6 LTG[36], RlpA depletion was

more severe in the ∆7 background, both by morphology and plating efficiency, consistent

with ∆7 exhibiting more limited LTG activity than ∆6.

Importantly, we confirmed that growth of ∆7 LTG depended on RlpA LTG activity, as

RlpA^{D145A}, a predicted active site mutant[35] (Figure 1 – Figure supplement 3AD) was

unable to promote growth despite being stably produced (Figure 1 – Figure supplement

4) and maintaining septal recruitment as indicated by the septal localization of an

RlpA^{D145A}-mCherry fusion (Figure 1C, Figure 1 – Figure supplement 4). Conversely, we

found that a truncated RlpA mutant lacking the conserved SPOR domain (which is

essential for septal localization in *P. aeruginosa*[35]) no longer localized to the division

septum in *V. cholerae* as an mCherry fusion (Figure 1 – Figure supplement 3C) yet still

fully complemented native RlpA depletion in both ∆6 and ∆7 (Figure 1C, Figure 1 – Figure

supplement 4), suggesting that RlpA LTG activity, but not septal localization, are essential

in these backgrounds. Taken together, these results demonstrate that at least during

growth in standard laboratory conditions, *V. cholerae* requires at least one active LTG of

the 8 currently annotated in its genome.

Only a subset of LTGs can independently fulfill all essential LTG functions

Since RlpA LTG activity, but not septal localization, was essential for ∆6 and ∆7 growth,

we asked whether growth required specialized LTG function, or just PG cleavage function

in general. To test this, we assessed the ability of other LTGs to complement ∆6 and ∆7

(Figure 1C, Figure 1 – Figure supplement 4). Only two native LTGs, MltD and Slt70, could

fully (MltD) or partially (Slt70) substitute for RlpA in the ∆7 LTG background,

demonstrating that RlpA, MltD, and (to a lesser degree) Slt70 are the only *V. cholerae*

LTGs capable of fulfilling all required LTG roles. Additional LTGs, MltA, MltB, and MltC,

were able to rescue RlpA depletion in the Δ6 LTG background (where MltG is present),

but not the Δ7 LTG background (where MltG is absent). These observations suggest that

LTGs perform at least two separable essential functions for viability, where MltG can

perform one function but requires MltA, MltB, or MltC to perform another, and vice versa,

as none of these LTGs are independently capable of supporting growth. Intriguingly, the

E. coli LTG MltE, for which *V. cholerae* has no known homologue, was also capable of

fully supporting *V. cholerae* growth as the sole functioning LTG, suggesting that LTG

essential functions may not depend on specific protein-protein interactions. Collectively,

these data suggest that LTGs are partially redundant; but some LTGs also exhibit varying

degrees of functional specificity.

LTGs are required during vegetative growth

We were surprised that our strains lacking major LTG activity (even Δ7) were viable. We

thus sought to characterize growth and morphology of these backgrounds in more detail.

While the Δ6 and Δ7 mutants grew at wild-type growth rates in LB medium when cultures

were started from a 100-fold dilution of saturated overnight cultures, we noticed a dilution-

dependent exacerbation of growth and morphology defects (Figure 2, Figure 2 – Figure

supplement 1, Figure 2 – Figure supplement 2). When Δ7 LTG cultures were highly diluted

from overnight cultures to increase the number of generations spent in exponential

growth, we started to observe a marked growth defect (Figure 2AB, Figure 2 – Figure

supplement 2A). Conversely, the *mltG::stop* single mutant and Δ6 LTG mutant grew at

wild-type growth rates independent of initial dilution factor (Figure 2B, Figure 2 – Figure

supplement 2A). These data suggest an essential or near-essential role for MltG during

sustained exponential growth, albeit only when other LTGs are also inactivated. We also

observed diverse aberrant morphologies within the LTG-deficient mutants (but not a

Δ*mltG*::stop single mutant) under these conditions (Figure 2C, Figure 2 – Figure

supplement 1). Interestingly, despite lacking most members of an entire class of PG

enzymes, Δ6 and Δ7 only exhibited only mild defects in length and width homeostasis

(Figure 2DE, Figure 2 – Figure supplement 2), which were only dilution-factor dependent

for cell length (suggesting a cumulative division defect). When we followed a time-course

after back-dilution from stationary phase, we observed morphological defects

accumulating in exponential phase and then largely disappearing in stationary phase

(Figure 2 – Figure supplement 3) in both Δ6 and Δ7. Collectively, these observations

suggest that LTG-deficient mutants suffer from cumulative damage during exponential

growth, which is partially alleviated in stationary phase.

LTG activity is required for survival in hypo-osmotic conditions

Our results detailed above suggested that the majority of LTGs are dispensable for

growth, catalyzing renewed interest in the question of what their physiological roles are.

To dissect potential roles for LTGs in cell envelope integrity maintenance, we subjected

the Δ6 and Δ7 mutants to growth in low osmolarity medium. Similar to the morphology

and growth defects in LB, both mutants were sensitive to low salt conditions in a dilution-

dependent manner (Figure 3A, Figure 3 – Figure supplement 1). Interestingly, the Δ7 LTG

mutant grew at wild-type rate during the initial growth period before a rapid decrease in

OD_{600}, indicative of lysis, suggesting that it is not the initial shock of changing osmotic

conditions that kills this mutant, but rather some cumulative damage during growth in low-

salt conditions (Figure 3A, Figure 3 – Figure supplement 1A). Importantly, this rules out

a simple cell envelope defect as a cause of LTG-deficient mutant osmo-sensitivity. We

then sought to dissect the contributions of individual LTGs to salt sensitivity. Other than

RlpA, which has an established role in LB0N growth[35,36] and is still present in the Δ6 and

Δ7 mutants, no other single LTG mutant exhibited a strong growth defect in LB0N (Figure

3 – Figure supplement 2A), all further suggesting that this defect is an accumulative

function of collective LTG insufficiency. Several LTGs (MltA, MltB, Slt70, and EcMltE)

were able to rescue Δ6 LTG growth in LB0N at higher dilutions (Figure 3 – Figure

supplement 2C), but none could rescue Δ7 growth in LB0N except MltG, which only

partially restored growth to Δ6 mutant levels (Figure 3 – Figure supplement 2D). This

suggests that no single LTG can fully support growth in LB0N, and that MltG, despite not

exhibiting a defect as a single mutant, is particularly important in this environment when

other LTGs are inactivated. Intriguingly, overexpression of *mltF* was toxic for both the Δ6

(LB0N only) and the Δ7 mutant (LB) (Figure 3 – Figure supplement 2BCD), preventing us

from assessing its potential for LTG-deficient mutant complementation.

LTG mutants are hypersensitive to accumulation of periplasmic polysaccharides

To determine the reason for ΔLTG defects in low salt media, we took advantage of the

spontaneous appearance of suppressors arising during Δ6 LTG growth in LB0N. (Figure

3B, Figure 3 – Figure supplement 1B).   Whole genome sequencing of three stable

suppressors identified three unique mutations: a deletion in *ptsH* (*vc0966*), a frameshift

mutation in *opgH* (*vc1287*), and a deletion mutation affecting both *opgH* and a

downstream gene (*vc1286*) (Figure 3C). *ptsH* encodes HPr, a key regulator of sugar

import via the phosphoenolpyruvate-carbohydrate phosphotransferase system (PTS).[39]

OpgH is critical to the synthesis of periplasmic glucans (OPGs), which accumulate under

conditions of low osmolarity and have been implicated in a variety of cell functions,

including steady-state maintenance of osmolarity in the periplasm.[40] We were particularly

intrigued by the role of periplasmic glucans and validated restoration of Δ6 LTG LB0N

growth using clean deletion mutants. Inactivation of *opgH* completely restored wild-type

growth of the Δ6 LTG mutant in LB0N, independent of initial dilution factor (Figure 3D,

Figure 3 – Figure supplement 1E). In contrast, inactivation of *opgH* was only able to

restore growth of the Δ7 LTG mutant in LB0N from a 10^{-2} inoculum (Figure 3D, Figure 3

– Figure supplement 1E), but not from greater dilutions, nor from a 10^{-5} inoculum in LB

(Figure 3 – Figure supplement 1CD), indicating that the Δ7 LTG mutant has additional

exponential growth-dependent defects unique from the Δ6 LTG mutant. Interestingly,

OpgH orthologues in *E. coli* and *Y. pseudotuberculosis* have also been ascribed a

moonlighting function, tying carbon availability with cell length by inhibiting FtsZ

filamentation when UDP-glucose levels are high in the cell[41,42]. The elongated cell

phenotype of the Δ6 LTG and Δ7 LTG mutants could in principle be indicative of a similar

activity by *V. cholerae* OpgH; however, deleting *opgH* from the Δ6 LTG mutant did not

restore wild-type cell length in Δ6 (Figure 3 – Figure supplement 3A). Additionally,

alignment of *V. cholerae* OpgH with *E. coli* and *Y. pseudotuberculosis* suggests that it

lacks much of the N-terminal domain that was shown to interact with FtsZ in *E. coli* (Figure

3 – Figure supplement 3B).[41]

We consequently considered a model where accumulation of periplasmic glucans is

detrimental to ΔLTG mutants, e.g., via periplasmic crowding or an increase in periplasmic

osmolarity. To test this model, we used an unrelated system to increase periplasmic

crowding. The *B. subtilis sacB* gene product, which is secreted and functions

extracellularly[43], isomerizes and polymerizes sucrose monomers into levan molecules up

to several kD[44], molecules much too large to escape the gram-negative periplasm through

outer membrane porins. This is often exploited in allelic exchange methods for mutant

generation as a means of counterselection,[45-47] where, for example, WT *V. cholerae* is

sensitive to *sacB* expression in LB0N+sucrose. We hypothesized that the ΔLTG mutants

might be hypersensitive to periplasmic levan synthesis even in standard LB salt

conditions (1% W/V). To test this, we engineered strains overexpressing *sacB*. Consistent

with the use of *sacB* as a counterselection method under low salt conditions, the wild type

is sensitive to *sacB* induction on LB0N (Figure 3 – Figure supplement 4), but not LB. In

contrast, we observed a *sacB*-dependent plating defect on sucrose in the Δ6 and Δ7 LTG

mutant on LB (Figure 3E, Figure 3 – Figure supplement 4). Interestingly, the *mltG::stop*

mutant also exhibited hypersensitivity to SacB activity (Figure 3E, Figure 3 – Figure

supplement 4), suggesting that accumulating MltG substrate is a particularly strong direct

or indirect contributor to this phenotype. Collectively, our data suggest that ΔLTG mutants

suffer from either hyperosmotic stress or excessive molecular crowding in the periplasm,

which can be exacerbated through the induction of long-chain polysaccharides.

PG transglycosylase activity causes Δ6 LTG mutant periplasmic stress

Based on our model that accumulation of polysaccharides in the periplasm is toxic, we

predicted that accumulation of uncrosslinked PG strands caused by uninterrupted GT

activity during TP inhibition, e.g. by β-lactams[48] (a process termed "futile cycling" when

coupled with LTG-mediated degradation[48]) should exacerbate LTG-deficient mutant

sickness, and this has indeed been shown with Slt70 in *E. coli*[48]. To test this hypothesis,

we assessed susceptibility of ΔLTG mutants to cell wall-acting antibiotics with varying

ability to induce futile cycling. In a disk diffusion assay, the Δ6 and Δ7 LTG mutants were

hypersensitive to inducers of futile cycling, i.e., general PBP inhibition by Penicillin G as

well as to inhibition of specific PBPs including PBP3 (aztreonam), PBP2 (mecillinam), and

PBP1b (cefsulodin) (Figure 4A). Conversely, both the Δ6 and the Δ7 mutant exhibited

wild-type sensitivity to moenomycin and fosfomycin, both of which inhibit cell wall

synthesis without inducing futile cycling. Thus, β-lactam sensitivity of LTG-deficient

mutants is not necessarily tied to simple inhibition of cell wall synthesis, but potentially

also to periplasmic crowding due to the accumulation of uncrosslinked PG strands.

Curiously, the Δ7 LTG mutant was hypersensitive to MreB inhibition by MP265. This

suggests that in the absence of other LTGs, MltG contributes to survival upon Rod system

insufficiency through an unknown mechanism.

To further dissect PG strand accumulation under these conditions, we visualized PG

synthesis and turnover during antibiotic treatment using the cell wall label BADA.[49,50] The

Δ6 and Δ7 LTG mutants were all readily labeled when grown with BADA (Figure 4 – Figure

supplement 1, Figure 4 – Figure supplement 2) prior to antibiotic addition. Upon treatment

with cell wall-targeting antibiotics, WT *V. cholerae* degrades its structural, rod-shaped

sacculus to ultimately yield stable, cell wall-deficient spheroplasts.[51,52] After up to 3 hours

of treatment with Penicillin G (100µg ml^{-1}, 20xMIC) and continued incubation with BADA,

the Δ6 and Δ7 LTG mutant spheroplasts accumulated strong periplasmic BADA signal

compared to the wild-type spheroplasts in which cell wall material is presumably

completely degraded and removed from the periplasm (Figure 4B, Figure 4 – Figure

supplement 1). Meanwhile, in the Δ6 and Δ7 LTG mutants, uncrosslinked PG likely

accumulates due to aberrant GT activity. Alternatively, degraded PG from the sacculus

may be sufficiently retained in the periplasm to be additionally labeled with BADA by L,

D-transpeptidases[53] in the Δ6 and Δ7 LTG but not in the WT. However, inhibition of cell

wall synthesis upstream of GT activity using fosfomycin (500µg ml^{-1}) (which is expected

to result in sacculus degradation without induction of futile cycling) did not result in

pronounced BADA accumulation in the periplasm, suggesting that the increasing BADA

signal observed during penicillin treatment does not solely reflect retention of labeled

debris from the degraded sacculus, but indeed is the result of ongoing cell wall synthesis

(Figure 4B, Figure 4 – Figure supplement 2). The diffuse nature of the BADA signal

additionally suggests that the PG in the Δ6 and Δ7 spheroplasts is solubilized debris

trapped in the periplasm, as opposed a thin, intact layer of peripheral PG. At earlier

stages of penicillin treatment, it also appeared that the Δ6 and Δ7 LTG mutants were

slower to degrade their rod-shaped poles (Figure 4 – Figure supplement 1).

PG recycling is not required during periplasmic stress

The roles for LTGs in PG recycling are well characterized. LTGs are required to digest

longer glycan strands down to single disaccharide subunits which can be imported into

the cytoplasm via AmpG, a permease that selectively recognizes the anhMurNAc residue

generated by LTG activity. Mutants lacking AmpG have been demonstrated to

accumulate extracellular monomeric disaccharide LTG-turnover products[54]. We therefore

asked whether a lack of recycling could account for some of our key phenotypes.

However, an Δ*ampG* mutant exhibited wild-type behavior for growth in LB0N, sacB

overexpression, beta-lactam resistance, and BADA staining after PenG exposure (Figure

4 – Figure supplement 3), demonstrating that lack of PG recycling does not promote the

LTG-deficient phenotypes observed here.

LTG insufficiency results in periplasmic PG strand accumulation

Why are LTG mutants sensitive to periplasmic accumulation of polymers? Accumulating

PG debris released during PG synthesis could, in principle, increase periplasmic

osmolarity and/or crowding, explaining defects associated with low salt conditions and

periplasmic polysaccharide accumulation. We therefore sought to quantify soluble,

periplasmic PG debris within crude cell lysates (excluding PG released into the growth

medium). Canonical PG architecture analysis relies on SDS-boiled sacculi isolated by

ultra-centrifugation, which permits sacculus characterization (Figure 1 – Figure

supplement 1D, Supplemental File 1), but ignores solubilized, uncrosslinked (freely-

diffusing) PG fragments associated with PG turnover processes such as those potentially

mediated by LTGs. We thus analyzed fragments that remained soluble after

sedimentation of purified sacculi, i.e., PG material that freely accumulates within the

periplasm (or inside the cell) but is not attached to the cell wall. Since entire periplasmic

PG strands cannot be easily resolved using subsequent LC-MS analysis, we also

digested the soluble PG fraction with muramidase (to generate smaller monomers

suitable for LC-MS detection) and then compared muramidase-treated vs. untreated

traces to determine soluble PG architecture (Figure 5A). We first analyzed soluble

products of LTG activity (Figure 5BC, Supplemental File 2). AnhMurNAc-tetrapeptide

(M4N) was abundant in the wild type during exponential phase, and predictably, M4N was

significantly depleted in the Δ7 LTG soluble muropeptide profile. Surprisingly, M4N was

enriched in the *mltG::kan* and Δ6 LTG mutants, suggesting that the absence of some

LTGs might cause upregulated activity of others.

Importantly, MurNAc-tetrapeptide species without anhMurNAc (M4) were detectable in

the WT during exponential phase, and significantly enriched in the Δ6/Δ7 LTG mutants

(Figure 5D, Supplemental File 2). By comparing the muramidase-treated vs. untreated

samples, we can infer the native state of these M4 and M4N (Figure 5ABD), i.e.,

determine whether these species occur predominantly as monomers or as parts of PG

polymers *in vivo*. Both species were significantly depleted in the muramidase-untreated

samples, suggesting that *in vivo*, they are part of polymeric, uncrosslinked peptidoglycan

strands. It is important to note that M4 monomers are not intermediates of cell wall

synthesis (which proceeds from a tripeptide directly to pentapeptide due to addition of

preformed D-Ala-D-Ala dipeptide).[55] This, in conjunction with a strong muramidase-

treatment-dependence of the abundance of M4 excludes a cytoplasmic origin of this

species.   It is also interesting to note that, unlike the WT, *mltG::kan*, and even Δ6 LTG

mutant, the Δ7 LTG appears to have a disproportionately large pool of polymeric M4

species (muramidase-dependent) compared to M4N species, suggesting that either the

soluble PG strands in the Δ7 LTG mutant are extremely long, or perhaps do not

ubiquitously terminate in anhMurNAc residues. The abundance of soluble PG strands

decreased in stationary phase (Figure 5CE, Supplemental File 2), which suggests that in

the LTG-deficient mutants, these strands are somehow cleared by RlpA, an unrecognized

LTG, or a cryptic PG hydrolase. Reduction of these uncrosslinked strands in stationary

phase is consistent with the alleviation of morphology defects observed as cells exit

exponential growth (Figure 2 – Figure supplement 3).

We next asked how these polymeric, uncrosslinked PG strands might be generated.

Since endopeptidases have been suggested to be essential for sacculus expansion

during cell elongation, we hypothesized that these soluble strands may reflect

endopeptidase activity, which would imply that EPs do not simply relax PG crosslinking

(as commonly assumed), but also might excise entire strands that accumulate - at least

transiently - in the periplasm. To test this hypothesis, we depleted the major

housekeeping EP ShyA[56] from a Δ6 LTG mutant, and found that the most abundant

monomer species, M4, was reduced in a ShyA-dependent manner in both the WT and

Δ6 LTG mutant (Figure 5F, Supplemental File 3). This strongly suggests that ShyA

produces the majority of the uncrosslinked strands in the periplasm. Curiously, depletion

of ShyA in the Δ6 LTG background resulted in large, irregular cells (Figure 5 – Figure

supplement 1), suggesting a cell envelope defect upon EP insufficiency. Additionally,

ShyA was required for Δ6 LTG mutant colony formation on LB plates, but Δ6 LTG liquid

cultures depleted for ShyA remained viable and could be rescued on plates restoring

shyA expression (Figure 5 – Figure supplement 1). Altogether, the data suggest that EPs

excise entire PG strands that accumulate in the periplasm until they are later cleared by

LTGs.

Discussion

Despite decades of work and a renewed research focus in the last few years, we still lack

a fundamental understanding of how bacteria harness both PG synthesis and constant

degradation to build and maintain an essential and dynamic wall structure that is able to

withstand a high internal turgor pressure. Even the basic physiological function of many

cell wall cleavage enzymes ("autolysins") remains unknown. The exact contributions of

LTGs to cell growth, for example, have remained elusive, mainly due to the high level of

apparent redundancy of these enzymes, which hampered classical genotype-phenotype

association analyses. While circumstantial evidence abounds, no collective physiological

characterization of essential LTG function has been conducted in any organism, and even

cases of demonstrated synthetic lethal relationships are rare and rely on indirect evidence

(inability to delete LTG genes).[34,57,58] The *V. cholerae* Δ6 and Δ7 LTG mutants described

here provide a vital platform for exploring both the collective and individual contributions

of lytic transglycosylases to bacterial viability.

Conserved LTGs are not functionally equivalent

In this study we confirm that LTG enzymatic activity is essential for growth and division.

We further demonstrate that some functional redundancy exists in *V. cholerae* between

diverse LTG families, as has been observed in other bacteria [12,20,59]. Members of LTG

families 1A (Slt70), 1D (MltD) and 6A (RlpA) can support growth independent of all other

natively encoded LTGs from families 1B, 1E, 2A, 3A, and 5A. Certain *V. cholerae* LTGs

additionally exhibit complementary genetic relationships in the absence of all other LTGs,

implying there are at least two essential roles for LTGs. For example, while neither MltG,

MltA, MltB, nor MltC can support growth on their own, strains expressing the MltG+MltA,

MltG+MltB, or MltG+MltC pairs are each viable. Surprisingly, and importantly, essential

LTG activities may not require conserved protein-protein interactions as suggested by the

viability of a *V. cholerae* mutant expressing non-native *E. coli* MltE as its sole LTG.

Scheurwater and Clarke unsuccessfully but informatively attempted to inactivate *mltF* in

an *E. coli* Δ6 LTG (Δ*mltABCDE* Δ*slt70*), suggesting that the essentiality of LTG activity is

conserved between *E. coli* and *V. cholerae,* and that MltF and MltG may not be

functionally redundant in *E. coli*.[58] In contrast, *mltF* inactivation is not synthetically lethal

with Δ*mltABCD* Δ*slt70* in *V. cholerae*. In fact, restoring MltF expression in the Δ6 and Δ7

LTG background is toxic, indicating that *V. cholerae* and *E. coli* exhibit different genetic

relationships between homologs of the same LTGs. This is perhaps most strongly

exemplified by the conservation of RlpA in both species, where *E. coli* RlpA has no

detectable LTG activity[35], yet *V. cholerae* RlpA can perform essential functions as the

sole LTG. Recently, the hydrolytic enzyme DigH in *E. coli* was shown to be functionally

similar to LTGs by contributing to the resolution of septal PG during daughter cell

separation[60]. The activity of DigH (of which *V. cholerae* does not possess a strong

homolog) may explain why *E. coli* does not appear to require RlpA LTG activity; this LTG-

like physiological role for a hydrolytic enzyme also suggests that we may need to look

beyond LTGs to fully address the roles for glycosidic bond cleavage within PG. Another

recent example of glycosidases serving as functional analogues includes MpgA from *S.*

aureus which shares enough homology with MltG to be identified bioinformatically as a

YceG-family LTG but harbors a single active site mutation to impart a muramidase

mechanism of PG cleavage.[30]

The functional redundancy of LTGs is also likely the root of an apparent paradox. Release

of new PG strands from undecaprenyl pyrophosphate should be an essential function, as

anchoring of PG to the inner membrane is toxic[61] and is likely necessary for lipid carrier

recycling. Yet MltG, the highly conserved "PG terminase," or more generally, "PG release

factor," that associates with PG synthetic complexes to release new strands, is not

essential. MltG can be inactivated in wild type *V. cholerae* (and *E. coli*) without significant

consequence, but our Δ7 LTG (*mltG*-null) mutant lost viability upon sustained growth in

exponential phase in LB. It is therefore likely that other LTGs can partially complement

for PG release activity. The dilution-dependent Δ7 LTG growth defect further suggests

that either RlpA has at least partial PG release activity and/or that during brief growth in

exponential phase, PG release activity is non-essential. MltG essentiality (or near-

essentiality) can thus perhaps only be appreciated in a multiple LTG background under

conditions where PG release activity is expected to be most important. This is also

evidenced by the increased sensitivity of the Δ7 LTG mutant to hypo-osmotic conditions

that cannot be rescued via the same mechanisms that rescue Δ6 LTG mutant, i.e.,

inactivation of OPG synthesis, as well as hypersensitivity to MreB-inhibitor MP265. All of

these defects point to a unique, MltG-dependent, conditionally essential LTG function.

While this study has focused on clearance of soluble PG debris, an activity that would

perhaps only be critical in organisms with a membrane-confined periplasmic space, PG

release factors analogous in function to MltG are broadly conserved even in Gram

positive organisms, additionally demonstrating a necessity for further investigation into

the role of PG release in bacterial cell wall homeostasis.

LTGs manage the periplasmic environment

By analyses of soluble PG, we discovered that even wild-type *V. cholerae* accumulates

loose strands of uncrosslinked PG of periplasmic origin (detected as muramidase-

dependent, soluble M4 products) and that production of these strands is dependent on

the EP ShyA. PG strands appear to accumulate to detrimental levels in the $\Delta 6$ and $\Delta 7$

mutants and we thus propose that processing soluble PG is a major collective function of

LTGs. PG strand accumulation induces hypersensitivity to other polymeric sugars in the

periplasm, such as OPGs, SacB-generated levans and β-lactam-induced futile cycling

products. Since these hypersensitivities are not shared by a $\Delta ampG$ mutant, previously

demonstrated to accumulate small PG monomers, this polysaccharide toxicity may arise

from the fact that these large polymers (PG, OPGs, or levans) cannot diffuse out through

OM porins, accumulating in a growth and/or substrate-dependent manner. This would

inappropriately increase the osmolarity and/or cause excessive crowding[5] of the

periplasm to interfere with normal growth-related processes, as exemplified by the mild

division defect of the *V. cholerae* LTG-deficient mutants.

While periplasmic crowding can clearly occur in Gram-negative bacteria due to their

diffusion-limiting OM, it is intriguing to consider implications for Gram-positives, where

LTGs are likewise conserved, and the cell wall may not provide a meaningful diffusion

barrier to long PG strands. We propose three possible (and testable) scenarios for the

role of LTGs in Gram-positive bacteria. For example, only PG release activity might be

essential in Gram-positives. It is also conceivable that LTGs are non-essential in Gram-

positive bacteria. Lastly, "periplasmic" crowding may in fact be detrimental in Gram-

positive bacteria: *B. subtilis* PG, for example, is very long chained and it is conceivable

that PG may constitute a reasonable diffusion barrier against such long strands.

Broader implications for LTGs in PG dynamics

Our data suggest that the most important roles for LTGs may not include "making space"

for new PG insertion, as has been assumed for fundamental models of PG synthesis.

More likely, LTGs clear the periplasm of most PG debris generated by endopeptidases

during normal growth by facilitating either recycling or release of small fragments into the

environment (Figure 6). As such, this collectively essential function of LTGs may not be

tied to a direct influence on PG synthesis. Importantly, uncrosslinked strands are not a

deviant feature unique to the LTG-deficient mutants nor β-lactam treatment (though they

become more apparent under these conditions), as they were also detectable in the wild

type during normal growth. Detection of EP-dependent, uncrosslinked PG strands

suggests that EP activity might be temporally separated from LTG-mediated turnover of

those PG strands.   This is seemingly contrary to pervasive models proposing a

synchronized synthetic/autolytic complex that digests old PG while simultaneously

building new PG. If such a complex exists, we propose that LTGs collectively maintain an

important role complementary to it, but not inevitably as part it.

Lastly, while high accumulation of PG strands is toxic, maintaining a small pool of

uncrosslinked PG material may actually serve a beneficial purpose to cells under some

conditions; bacteria might thus have an adaptive reason for avoiding tightly coordinating

LTG activity with synthesis. Inactivation of *sltY* in *E. coli,* for example, can promote β-

lactam resistance by upregulating L,D-crosslinking activity of L,D-transpeptidase LdtD

(YcbB), presumably due to the increased presence of uncrosslinked glycan strands under

these conditions.[62] It is possible that one function of L,D-TPs is to patch holes in the

sacculus via readily available soluble PG strands, constituting an innate "bike tire repair

kit" for the bacterial cell wall. Future studies of this LTG-mediated turnover of

uncrosslinked PG could be critical for understanding PG repair mechanisms, as well as

general periplasmic homeostasis, both of which have been historically elusive topics.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (*V. cholerae*)	*mltA*	UniProtKB	*vc_2312*	
gene (*V. cholerae*)	*mltB*	UniProtKB	*vc_1956*	
gene (*V. cholerae*)	*mltC*	UniProtKB	*vc_0450*	
gene (*V. cholerae*)	*mltD*	UniProtKB	*vc_2237*	

gene (*V. cholerae*)	*mltF*	UniProtKB	*vc_0866*	
gene (*V. cholerae*)	*mltG*	UniProtKB	*vc_2017*	
gene (*V. cholerae*)	*rlpA*	UniProtKB	*vc_0948*	
gene (*V. cholerae*)	*slt70*	UniProtKB	*vc_0700*	
gene (*V. cholerae*)	*shyA*	UniProtKB	*vc_a0079*	
gene (*V. cholerae*)	*opgH*	UniProtKB	*vc_1287*	
gene (*V. cholerae*)	*ampG*	UniProtKB	*vc_2300*	
gene (*V. cholerae*)	*lacZ*	Kegg2 Gene DB	*vc_2338*	
gene (*E. coli*)	*EcrlpA*	UniProtKB	*b0633*	
gene (*E. coli*)	*EcmltE*	UniProtKB	*b1193*	
gene (*P. aeruginosa*)	*ParlpA*	UniProtKB	*pa4000*	
gene (*B. subtilis*)	*sacB*	UniProtKB	*bsu34450*	

Strain, strain background (*V. cholerae*)	El Tor N16961; wild-type	PMID: 243277		
Recombinant DNA reagent	Δ*mltABCDF* Δ*slt70*; Δ6 LTG	PMID: 31286580		
Recombinant DNA reagent	Δ*mltABCDF* Δ*slt70* *mltG::stop;* Δ7 LTG	This Study		See Supplemental File 4; Request from Doerr lab
Recombinant DNA reagent	*mltG::stop*	This Study		See Supplemental File 4; Request from Doerr Lab
Recombinant DNA reagent	pCVD442	PMID: 15109831		
Recombinant DNA reagent	pTOX5	PMID: 31201277	Addgene #127450	
Recombinant DNA reagent	pAM224; pGP704-kanR	PMID: 2836362		
Recombinant DNA reagent	pAM299	PMID: 25631756		
Recombinant DNA reagent	pJL1	PMID: 24348240		
Antibody	anti-mCherry (rabbit polyclonal)	GeneTex	GTX59788	(1:5,000)
Antibody	Goat anti-rabbit IRDye 800CW secondary	Li-cor	926-32211	(1:16,000)

Antibody	anti-RpoA (mouse monoclonal)	BioLegend	663104	(1:10,000)
Antibody	Goat anti-mouse IRDye 800CW secondary	Li-cor	926-32210	(1:16,000)
Chemical compound, drug	Penicillin G potassium salt	Fisher Scientific	CAS: 113-98-4	
Chemical compound, drug	Aztreonam	Fisher Scientific	CAS: 78110-38-0	
Chemical compound, drug	Mecillinam	Sigma Aldrich	CAS: 32887-01-7	
Chemical compound, drug	Cefsulodin sodium salt hydrate	TCI Chemicals	CAS: 1426397-23-0	
Chemical compound, drug	S-(4-Chlorobenzyl) Isothiouronium chloride (MP265)	Chem Impex International	CAS: 544-47-8	
Chemical compound, drug	Phosphomycin disodium salt (Fosfomycin)	Sigma Aldrich	CAS: 26016-99-9	
Software, algorithm	FIJI	PMID: 22743772		
Software, algorithm	MicrobeJ	PMID: 27572972		
Software, algorithm	Oufti	PMID: 26538279		
Other	BADA	PMID: 28989665		

525
526

Bacterial strains and growth conditions

V. cholerae strains in this study are derivatives of *V. cholerae* WT El Tor strain N16961

[63]. Construction of plasmids and mutant *V. cholerae* strains is described in the next

section along with a table of strains and plasmids used in this work (Supplemental File

4).

Strains were grown at 30°C or 37°C in Luria-Bertani (LB-Miller, Fisher Bioreagents#

BP97235) with or without 1% NaCl, 1% sucrose, or 10% sucrose, or in M9 minimal media

+ 0.4% glucose (Cold Spring Harbor Protocols) where indicated in the figure legends.

Growth media were supplemented with kanamycin (50 µg ml^{-1}), ampicillin (25 µg ml^{-1}), or

chloramphenicol (5 µg ml^{-1}) in plates and overnight cultures when needed to maintain

plasmids or chromosomal integration of suicide vectors. Genes under P_{ara} and P_{tac}

regulation were induced with 0.4% L-arabinose or 200 µM isopropyl-β-D-1-

thiolgalactopyranoside (IPTG), respectively.

Construction of plasmids and strains

A summary of all strains, plasmids, and primers used in this study can be found in Tables

S9-S11. *E. coli* DH5α λ*pir* was used for general cloning, while *E. coli* SM10 or MFD λ*pir*

were used for conjugation into *V. cholerae*[64]. Plasmids were constructed using Gibson

assembly[65] with the exception of plasmids expressing *rlpAΔSPOR* or *rlpA^{D145A},* which

were generated by site directed mutagenesis of the parent wild-type sequence *rlpA*

plasmids. All Illumina whole genome sequencing and variant calling for Δ6 LTG and Δ7

LTG strain verification or suppressor identification was performed by the Microbial

Genome Sequencing Center (MiGS, Pittsburg, PA, USA).

Most chromosomal in-frame deletions (or premature stop codon mutants) were generated

using the pCVD442 *ampR/sacB* allelic exchange system[66]. 500bp regions flanking the

gene to be deleted were amplified from N16961 genomic DNA by PCR, cloned into

suicide vector pCVD442, and conjugated into *V. cholerae*. Conjugation was performed

by mixing and pelleting equal volumes of recipient *V. cholerae* and SM10 or MFD λ*pir*

donor LB overnight cultures, spotting the mixed pellet onto LB (+600 μM diaminopimelic

acid [DAP] for MFD λ*pir*) followed by incubation at 37°C for 3 hrs. The first round of

selection was performed on LB + streptomycin (200 μg ml^{-1}) + ampicillin (100 μg ml^{-1}) at

30°C followed by counterselection on salt-free LB + 10% sucrose + streptomycin at room

temperature. Inducers for conditionally essential genes were included in all media during

conjugation and selection. Addition of 0.2% glucose was required for maintenance in *E.*

coli of plasmids expressing *Vc mltB*. Deletions were verified by PCR.

Introduction of a premature stop codon to *mltG* at its native locus to yield Δ7 LTG as well

as in-frame deletions of *opgH* were constructed using the pTOX5 *cmR/msqR* allelic

exchange system[38]. Flanking regions were cloned into pTOX5 as described for

pCVD442. Conjugation was performed by mixing and pelleting equal parts of recipient

V. cholerae and donor MFD λ*pir,* and spotting onto LB + 1% glucose + 600μM DAP at

37°C for 5 hrs. The first round of selection was performed on LB + chloramphenicol (5 μg

μg ml^{-1}) + streptomycin + 1% glucose at 30°C. Chloramphenicol resistant colonies were

picked into a 96well plate containing 200µL LB + 1% glucose and incubated at 37°C

without agitation for 3 hrs, then counter selected on LB + 1% rhamnose at 30°C.

Mutations were verified by PCR.

Ectopic chromosomal expression from IPTG-inducible P_{tac} was achieved through use of

suicide vector pTD101, a pJL1[67] derivative carrying the P_{tac} promoter, a multiple cloning

site, and *lacIq* and integrates into the native *V. cholerae lacZ* (*vc2338*) locus. Single

genes of interest were amplified from N16961 genomic DNA, introducing a strong

consensus RBS (AGGAGA). Genes downstream of *mltG* (*vc2016, vc2015, vc2014*) were

amplified together maintaining their native organization, including 30bp upstream of

vc2016 to retain the native RBS. Selection for double crossover events was performed

as described for pCVD442. Due to hypersensitivity of the Δ7 LTG strain to β-lactams,

ampicillin was reduced to 25 µg ml^{-1} for the first selection. Due to the osmosensitivity of

the Δ7 LTG strain, pTD101 in this strain (and control strains) was maintained as an ampR

single crossover without counterselection.

Suicide vector pAM299[68] was used to place *rlpA* under P_{ara} control at its native locus for

RlpA depletion experiments. pAM299 was introduced via conjugation and selection for

single crossover events on LB + kanamycin (50 µg ml^{-1}) + streptomycin. IPTG-inducible

overexpression of *rlpA-mCherry* fusions for localization studies and *sacB* for levan toxicity

assays was achieved using pHL100[68] or its conjugatable derivative pHL100mob. The

sacB gene was amplified from pCVD442.

Gene insertion/disruption assay for gene essentiality

The suicide vector pAM224 was used to disrupt genes through single crossover integration events. 300bp internal regions in the first third of each respective *orf* (towards the 5' end) were cloned into pAM224 using methods described above. Quantitative conjugation was performed by washing, mixing, and pelleting 500 µL of recipient *V. cholerae* and donor MFD λ*pir*. Pellets were re-suspended in 50 µl of LB, spotted onto a 45 µm filter on LB + 600 µM DAP, and incubated at 37°C for 4 hrs. Cells were recovered from the filters into 1 ml of LB by vigorous vortexing and 20 µl of the suspension was reserved for 10-fold serial dilution and spotting onto LB + streptomycin (200 µg ml^{-1}) incubated at 30°C to calculate total CFU/ml for all viable *V. cholerae.* The remaining suspension was pelleted and plated on LB + kanamycin (50 µg ml^{-1}) + streptomycin incubated at 30°C. Viable CFU/ml were calculated, and kanamycin resistance was verified by patching 50 colonies back onto LB + streptomycin +/- kanamycin (all kanR colonies were patched if fewer than 50 colonies were recovered). Transformation efficiency was calculated as a ratio of kanR CFU to all strepR CFU.

Western Blot Analysis

Expression of translational mCherry fusions was induced in WT *V. cholerae* with 1 mM IPTG in LB and grown to OD600 ~ 0.6. Cells were harvested by centrifugation (9500xg, 15 min) at room temperature and resuspended in 1% SDS + 10 mM dithiothreitol (DTT) lysis buffer. Resuspended cells were incubated at 95°C for 3 min, then sonicated 4 x 5 s at 20% amplitude. Standard western blots against mCherry were performed using polyclonal mCherry antibody (Genetex #GTX59788) and detection by IRDye 800CW

secondary antibody (Li-cor #926-32211). After imaging for mCherry, the same blots were

then re-incubated with monoclonal RpoA antibody (BioLegend # 663104) detected by

IRDye 800CW secondary antibody on an Odyssey CLx imaging device (Li-cor).

Growth rate experiments

Saturated overnight cultures used for growth curve experiments were washed once and

resuspended in final growth media, normalizing to OD_{600} 2.0. Normalized cell

suspensions were serially diluted into 200 µL growth media and incubated in a Bioscreen

growth plate reader (Growth Curves America) at 37°C with random shaking at maximum

amplitude, and OD_{600} recorded at 5-min intervals. Calculations of doublings per hour

(DPH) were performed in R as previously described[59]. Briefly, logarithmic regressions

were fitted to sections of growth curves with >5 consecutive values corresponding to

OD_{600} 0.03-0.1 (for 10^{-2} diluted inocula) or OD_{600} 0.01-0.1 (for $<10^{-3}$ diluted inocula).

Logarithmic regressions of ≥3 replicates with fit value $r^2 > 0.95$ were used to estimate

mean doublings per hour for each strain in each growth condition. Growth rates were not

calculated for samples that did not reach OD_{600} 0.1, either through absence of detectable

growth or lysis prior to reaching OD_{600} 0.1. Growth rates were also not calculated for

replicates that were subsequently determined to be suppressors or grew due to

irreproducible adaptation. Means within strains between dilution factors and between

strains within dilution factors were compared using a two-way Anova and Tukey HSD

post-hoc test.

Morphology analysis by microscopy

Strains were grown as described in the figure legends and imaged without fixation on LB

0.8% agarose using a Leica DMi8 inverted microscope. Phase-contrast images were

analyzed using MicrobeJ. Default parameter settings were applied, and features (septa)

were defined as 25% constriction of cell width. Cell outlines were manually edited as

needed.

Sacculus composition analysis

PG composition from insoluble sacculi samples were analyzed as described previously

with some modifications[69,70]. Briefly, cells were harvested and resuspended and in boiled

5% SDS for 1 h. Sacculi were repeatedly washed by ultracentrifugation (110,000 rpm,

10 min, 20ºC) with MilliQ water until SDS was totally removed. Samples were treated with

20 µg Proteinase K (1 h, 37 °C) for removal of Braun's lipoprotein, and finally treated with

muramidase (100 µg/mL) for 16 hours at 37 ºC. Muramidase digestion was stopped by

boiling and coagulated proteins were removed by centrifugation (14,000 rpm, 15 min).

For sample reduction, the pH of the supernatants was adjusted to pH 8.5-9.0 with sodium

borate buffer and sodium borohydride was added to a final concentration of 10 mg/mL.

After incubating for 30 min at room temperature, the samples pH was adjusted to pH 3.5

with orthophosphoric acid.

UPLC analyses of muropeptides were performed on a Waters UPLC system (Waters

Corporation, USA) equipped with an ACQUITY UPLC BEH C18 Column, 130Å, 1.7 µm,

2.1 mm X 150 mm (Waters, USA) and a dual wavelength absorbance detector. Elution of

muropeptides was detected at 204 nm. Muropeptides were separated at 45°C using a

linear gradient from buffer A (formic acid 0.1% in water) to buffer B (formic acid 0.1% in

acetonitrile) in an 18-minute run, with a 0.25 ml/min flow.

Relative total PG amount was calculated by comparison of the total intensities of the

chromatograms (total area) from three biological replicas normalized to the same OD600

and extracted with the same volumes. Muropeptide identity was confirmed by MS/MS

analysis, using a Xevo G2-XS QTof system (Waters Corporation, USA) (*see next section*

for details). Quantification of muropeptides was based on their relative abundances

(relative area of the corresponding peak) normalized to their molar ratio. Analyses were

performed in biological triplicates and means were compared with unpaired t-tests.

Soluble peptidoglycan analysis

Sample preparation of soluble PG samples was performed as follows. Bacteria cultures

were harvested by centrifugation (4,000 rpm, 20 min, 4°C). Cell pellets were gently

resuspended and washed twice with ice-cold 0.9% NaCl solution. After pelleting the cells

again by centrifugation, they were resuspended in 1 ml water and boiled for 30 min.

Samples were centrifuged again to remove cell debris at 14,000 rpm for 15 min, and

soluble fractions were transferred to new tubes. Next, samples were filtered using 0.2 μm

pore size filters. Half of the sample was treated with muramidase (100 μg/mL) for 16 hours

at 37 ºC. Muramidase digestion was stopped by boiling and coagulated proteins were

removed by centrifugation (14,000 rpm, 15 min). Finally, sample pH was adjusted to pH

3.5 with orthophosphoric acid. When needed, samples were diluted or concentrated by

speed vacuum.

Soluble muropeptides were detected and characterized by MS/MS analysis, using a Xevo
G2-XS QTof system (Waters Corporation, USA) equipped with an ACQUITY UPLC BEH
C18 Column (130 Å, 1.7 µm, 2.1 mm x 150 mm (Waters, USA)). Muropeptides were
separated at 45 ºC using a linear gradient from buffer A (formic acid 0.1% in water) to
buffer B (formic acid 0.1% in acetonitrile) in an 18-min run, with a 0.25 ml/min flow. The
QTOF–MS instrument was operated in positive ionization mode. Detection of
muropeptides was performed by MS^E to allow for the acquisition of precursor and product
ion data simultaneously, using the following parameters: capillary voltage at 3.0 kV,
source temperature to 120 ºC, desolvation temperature to 350 ºC, sample cone voltage
to 40 V, cone gas flow 100 l/h, desolvation gas flow 500 l/h and collision energy (CE): low
CE: 6 eV and high CE ramp: 15-40 eV. Mass spectra were acquired at a speed of 0.25
s/scan. The scan was in a range of 100–2000 *m/z*. Data acquisition and processing was
performed using UNIFI software package (Waters Corp.).

An in-house compound library built in UNIFI was used for detection and identification of
muropeptides. Subsequent identification and confirmation of each muropeptide was
performed by comparison of the retention-times and mass spectrometric data to known
samples. Quantification was performed by integrating peak areas from extracted ion
chromatograms (EICs) of the corresponding *m/z* value of each muropeptide and
normalized to their molar ratio. Soluble muropeptide analyses were performed in
biological triplicates and means were compared with unpaired t-tests.

Antibiotic sensitivity

For zone of inhibition assays, a lawn of saturated overnight culture was spread on an LB

plate and allowed to dry for 15 minutes. Filter disks (6 mm) were dropped with 5µL of

antibiotic solutions (Table S4) onto the lawns and incubated at 37°C for 24 hrs before

measurements. Means were compared with a one-way ANOVA and Tukey post-hoc test.

Fluorescent D-amino acid PG labeling

Saturated overnight cultures were diluted 1:100 into LB with 100 µM BADA[49] at 37°C for

1.5 hrs before addition of antibiotic. Labeled samples were washed one time in LB before

imaging on LB 0.8% agarose pads using a Leica DMi8 inverted microscope set for 490

nm excitation.

Acknowledgements

We would like to acknowledge Dr. Stephen Zinder (Cornell University) for informative

comments based on his unique insight into bacterial physiology. We thank Dr. John

Helmann (Cornell University) and members of the Dörr lab for helpful comments on the

manuscript. Research on autolysins in the Dörr lab is funded by NIH R01 GM130971.

Research in the Cava lab is supported by MIMS, the Knut and Alice Wallenberg

Foundation (KAW), the Swedish Research Council and the Kempe Foundation. Research

in the VanNieuwenhze lab is supported by the NIH through R01 GM113172 and R35

GM136365.

References

1   Osawa, M. & Erickson, H. P. Turgor Pressure and Possible Constriction
Mechanisms in Bacterial Division. *Frontiers in Microbiology* **9**,
doi:10.3389/fmicb.2018.00111 (2018).
2   Cabeen, M. T. & Jacobs-Wagner, C. Bacterial cell shape. *Nature Reviews*
Microbiology **3**, 601-610, doi:10.1038/nrmicro1205 (2005).
3   Egan, A. J. F., Errington, J. & Vollmer, W. Regulation of peptidoglycan synthesis
and remodelling. *Nature Reviews Microbiology* **18**, 446-460, doi:10.1038/s41579-
020-0366-3 (2020).
4   Zhao, H., Patel, V., Helmann, J. D. & Dörr, T. Don't let sleeping dogmas lie: new
views of peptidoglycan synthesis and its regulation. *Mol Microbiol* **106**, 847-860,
doi:10.1111/mmi.13853 (2017).
5   Sochacki, K. A., Shkel, I. A., Record, M. T. & Weisshaar, J. C. Protein diffusion in
the periplasm of E. coli under osmotic stress. *Biophys J* **100**, 22-31,
doi:10.1016/j.bpj.2010.11.044 (2011).
6   Cho, H. *et al.* Bacterial cell wall biogenesis is mediated by SEDS and PBP
polymerase families functioning semi-autonomously. *Nat Microbiol* **1**, 16172,
doi:10.1038/nmicrobiol.2016.172 (2016).
7   Leclercq, S. *et al.* Interplay between Penicillin-binding proteins and SEDS proteins
promotes bacterial cell wall synthesis. *Scientific Reports* **7**, 43306,
doi:10.1038/srep43306 (2017).
8   Taguchi, A. *et al.* FtsW is a peptidoglycan polymerase that is functional only in
complex with its cognate penicillin-binding protein. *Nat Microbiol* **4**, 587-594,
doi:10.1038/s41564-018-0345-x (2019).
9   Nambu, T., Minamino, T., Macnab, R. M. & Kutsukake, K. Peptidoglycan-
Hydrolyzing Activity of the FlgJ Protein, Essential for Flagellar Rod Formation
in*Salmonella typhimurium*. *Journal of Bacteriology* **181**, 1555-1561,
doi:doi:10.1128/JB.181.5.1555-1561.1999 (1999).
10   Santin, Y. G. & Cascales, E. Domestication of a housekeeping transglycosylase
for assembly of a Type VI secretion system. *EMBO Rep* **18**, 138-149,
doi:10.15252/embr.201643206 (2017).
11   Höltje, J. V. Growth of the stress-bearing and shape-maintaining murein sacculus
of Escherichia coli. *Microbiol Mol Biol Rev* **62**, 181-203,
doi:10.1128/mmbr.62.1.181-203.1998 (1998).
12   Herlihey, F. A. & Clarke, A. J. in *Protein Reviews: Volume 17* (ed M. Zouhair
Atassi) 41-56 (Springer Singapore, 2017).
13   Viollier, P. H. & Shapiro, L. A lytic transglycosylase homologue, PleA, is required
for the assembly of pili and the flagellum at the Caulobacter crescentus cell pole.
Mol Microbiol **49**, 331-345, doi:10.1046/j.1365-2958.2003.03576.x (2003).
14   Uehara, T. & Park, J. T. Growth of Escherichia coli: significance of peptidoglycan
degradation during elongation and septation. *J Bacteriol* **190**, 3914-3922,
doi:10.1128/jb.00207-08 (2008).
15   Nguyen, L. T., Gumbart, J. C., Beeby, M. & Jensen, G. J. Coarse-grained
simulations of bacterial cell wall growth reveal that local coordination alone can be
sufficient to maintain rod shape. *Proceedings of the National Academy of Sciences*
112, E3689-E3698, doi:10.1073/pnas.1504281112 (2015).

16 Lee, T. K. & Huang, K. C. The role of hydrolases in bacterial cell-wall growth.
Current opinion in microbiology **16**, 760-766, doi:10.1016/j.mib.2013.08.005
(2013).
17 Murphy, S. G. *et al.* Class A Penicillin-Binding Protein-Mediated Cell Wall
Synthesis Promotes Structural Integrity during Peptidoglycan Endopeptidase
Insufficiency in Vibrio cholerae. *mBio* **12**, doi:10.1128/mBio.03596-20 (2021).
18 Singh, S. K., SaiSree, L., Amrutha, R. N. & Reddy, M. Three redundant murein
endopeptidases catalyse an essential cleavage step in peptidoglycan synthesis of
Escherichia coli K12. *Mol Microbiol* **86**, 1036-1051, doi:10.1111/mmi.12058
(2012).
19 Byun, B. *et al.* Mechanism of the Escherichia coli MltE lytic transglycosylase, the
cell-wall-penetrating enzyme for Type VI secretion system assembly. *Sci Rep* **8**,
4110, doi:10.1038/s41598-018-22527-y (2018).
20 Dik, D. A., Marous, D. R., Fisher, J. F. & Mobashery, S. Lytic transglycosylases:
concinnity in concision of the bacterial cell wall. *Crit Rev Biochem Mol Biol* **52**, 503-
542, doi:10.1080/10409238.2017.1337705 (2017).
21 Lee, M. *et al.* Exolytic and endolytic turnover of peptidoglycan by lytic
transglycosylase Slt of Pseudomonas aeruginosa. *Proceedings of the
National Academy of Sciences* **115**, 4393-4398, doi:10.1073/pnas.1801298115
(2018).
22 Vijayaraghavan, J. *et al.* Structural studies and molecular dynamics simulations
suggest a processive mechanism of exolytic lytic transglycosylase from
Campylobacter jejuni. *PLOS ONE* **13**, e0197136,
doi:10.1371/journal.pone.0197136 (2018).
23 Höltje, J. V., Mirelman, D., Sharon, N. & Schwarz, U. Novel type of murein
transglycosylase in Escherichia coli. *J Bacteriol* **124**, 1067-1076,
doi:10.1128/jb.124.3.1067-1076.1975 (1975).
24 Williams, A. H. *et al.* A step-by-step in crystallo guide to bond cleavage and 1,6-
anhydro-sugar product synthesis by a peptidoglycan-degrading lytic
transglycosylase. *J Biol Chem* **293**, 6000-6010, doi:10.1074/jbc.RA117.001095
(2018).
25 Kraft, A. R., Templin, M. F. & Höltje, J. V. Membrane-bound lytic
endotransglycosylase in Escherichia coli. *Journal of bacteriology* **180**, 3441-3447,
doi:10.1128/JB.180.13.3441-3447.1998 (1998).
26 Bohrhunter, J. L., Rohs, P. D. A., Torres, G., Yunck, R. & Bernhardt, T. G. MltG
activity antagonizes cell wall synthesis by both types of peptidoglycan polymerases
in Escherichia coli. *Molecular Microbiology* **115**, 1170-1180,
doi:https://doi.org/10.1111/mmi.14660 (2021).
27 Yunck, R., Cho, H. & Bernhardt, T. G. Identification of MltG as a potential terminase
for peptidoglycan polymerization in bacteria. *Molecular Microbiology* **99**, 700-718,
doi:10.1111/mmi.13258 (2016).
28 Sassine, J., Pazos, M., Breukink, E. & Vollmer, W. Lytic transglycosylase MltG
cleaves in nascent peptidoglycan and produces short glycan strands. *The Cell
Surface* **7**, 100053, doi:https://doi.org/10.1016/j.tcsw.2021.100053 (2021).
29 Tsui, H. C. *et al.* Suppression of a deletion mutation in the gene encoding essential
PBP2b reveals a new lytic transglycosylase involved in peripheral peptidoglycan

824   synthesis in Streptococcus pneumoniae D39. *Mol Microbiol* **100**, 1039-1065,
825   doi:10.1111/mmi.13366 (2016).

826   30   Taguchi, A., Page, J. E., Tsui, H.-C. T., Winkler, M. E. & Walker, S. Biochemical
827   reconstitution defines new functions for membrane-bound glycosidases in
828   assembly of the bacterial cell wall. *Proceedings of the National Academy of*
829   *Sciences* **118**, e2103740118, doi:10.1073/pnas.2103740118 (2021).

830   31   Koraimann, G. Lytic transglycosylases in macromolecular transport systems of
831   Gram-negative bacteria. *Cell Mol Life Sci* **60**, 2371-2388, doi:10.1007/s00018-
832   003-3056-1 (2003).

833   32   Jacobs, C., Huang, L. J., Bartowsky, E., Normark, S. & Park, J. T. Bacterial cell
834   wall recycling provides cytosolic muropeptides as effectors for beta-lactamase
835   induction. *Embo j* **13**, 4684-4694 (1994).

836   33   Dik, D. A., Fisher, J. F. & Mobashery, S. Cell-Wall Recycling of the Gram-Negative
837   Bacteria and the Nexus to Antibiotic Resistance. *Chem Rev* **118**, 5952-5984,
838   doi:10.1021/acs.chemrev.8b00277 (2018).

839   34   Heidrich, C., Ursinus, A., Berger, J., Schwarz, H. & Holtje, J. V. Effects of multiple
840   deletions of murein hydrolases on viability, septum cleavage, and sensitivity to
841   large toxic molecules in Escherichia coli. *J Bacteriol* **184**, 6093-6099,
842   doi:10.1128/jb.184.22.6093-6099.2002 (2002).

843   35   Jorgenson, M. A., Chen, Y., Yahashiri, A., Popham, D. L. & Weiss, D. S. The
844   bacterial septal ring protein RlpA is a lytic transglycosylase that contributes to rod
845   shape and daughter cell separation in Pseudomonas aeruginosa. *Mol Microbiol*
846   **93**, 113-128, doi:10.1111/mmi.12643 (2014).

847   36   Weaver, A. I. *et al.* Lytic transglycosylases RlpA and MltC assist in Vibrio cholerae
848   daughter cell separation. *Molecular Microbiology* **112**, 1100-1115,
849   doi:10.1111/mmi.14349 (2019).

850   37   Priyadarshini, R., Popham, D. L. & Young, K. D. Daughter cell separation by
851   penicillin-binding proteins and peptidoglycan amidases in Escherichia coli. *Journal*
852   *of bacteriology* **188**, 5345-5355, doi:10.1128/JB.00476-06 (2006).

853   38   Lazarus, J. E. *et al.* A New Suite of Allelic-Exchange Vectors for the Scarless
854   Modification of Proteobacterial Genomes. *Appl Environ Microbiol* **85**,
855   doi:10.1128/aem.00990-19 (2019).

856   39   Deutscher, J. *et al.* The bacterial phosphoenolpyruvate:carbohydrate
857   phosphotransferase system: regulation by protein phosphorylation and
858   phosphorylation-dependent protein-protein interactions. *Microbiol Mol Biol Rev* **78**,
859   231-256, doi:10.1128/mmbr.00001-14 (2014).

860   40   Bontemps-Gallo, S., Bohin, J. & Lacroix, J. Osmoregulated Periplasmic Glucans.
861   *EcoSal Plus*, doi:doi:10.1128/ecosalplus.ESP-0001-2017 (2017).

862   41   Hill, N. S., Buske, P. J., Shi, Y. & Levin, P. A. A Moonlighting Enzyme Links
863   Escherichia coli Cell Size with Central Metabolism. *PLOS Genetics* **9**, e1003663,
864   doi:10.1371/journal.pgen.1003663 (2013).

865   42   Quintard, K. *et al.* Evaluation of the Role of the opgGH Operon in Yersinia
866   pseudotuberculosis and Its Deletion during the Emergence of Yersinia pestis.
867   *Infection and immunity* **83**, 3638-3647, doi:10.1128/IAI.00482-15 (2015).

868   43   Pereira, Y., Petit-Glatron, M.-F. & Chambert, R. yveB, encoding endolevanase
869   LevB, is part of the sacB–yveB–yveA levansucrase tricistronic operon in Bacillus

subtilis. *Microbiology* **147**, 3413-3419, doi:https://doi.org/10.1099/00221287-147-
12-3413 (2001).

44   Tanaka, T., Oi, S. & Yamamoto, T. The molecular structure of low and high
molecular weight levans synthesized by levansucrase. *J Biochem* **87**, 297-303,
doi:10.1093/oxfordjournals.jbchem.a132737 (1980).

45   Steinmetz, M., Le Coq, D., Djemia, H. B. & Gay, P. [Genetic analysis of sacB, the
structural gene of a secreted enzyme, levansucrase of Bacillus subtilis Marburg].
Mol Gen Genet **191**, 138-144, doi:10.1007/bf00330901 (1983).

46   Gay, P., Le Coq, D., Steinmetz, M., Ferrari, E. & Hoch, J. A. Cloning structural
gene sacB, which codes for exoenzyme levansucrase of Bacillus subtilis:
expression of the gene in Escherichia coli. *J Bacteriol* **153**, 1424-1431,
doi:10.1128/jb.153.3.1424-1431.1983 (1983).

47   Blomfield, I. C., Vaughn, V., Rest, R. F. & Eisenstein, B. I. Allelic exchange in
Escherichia coli using the Bacillus subtilis sacB gene and a temperature-sensitive
pSC101 replicon. *Molecular Microbiology* **5**, 1447-1457,
doi:https://doi.org/10.1111/j.1365-2958.1991.tb00791.x (1991).

48   Cho, H., Uehara, T. & Bernhardt, T. G. Beta-lactam antibiotics induce a lethal
malfunctioning of the bacterial cell wall synthesis machinery. *Cell* **159**, 1300-1311,
doi:10.1016/j.cell.2014.11.017 (2014).

49   Hsu, Y. P. *et al.* Full color palette of fluorescent d-amino acids for in situ labeling
of bacterial cell walls. *Chem Sci* **8**, 6313-6321, doi:10.1039/c7sc01800b (2017).

50   Kuru, E. *et al.* In Situ probing of newly synthesized peptidoglycan in live bacteria
with fluorescent D-amino acids. *Angew Chem Int Ed Engl* **51**, 12519-12523,
doi:10.1002/anie.201206749 (2012).

51   Dörr, T., Davis, B. M. & Waldor, M. K. Endopeptidase-mediated beta lactam
tolerance. *PLoS Pathog* **11**, e1004850, doi:10.1371/journal.ppat.1004850 (2015).

52   Weaver, A. I. *et al.* Genetic Determinants of Penicillin Tolerance in Vibrio cholerae.
Antimicrob Agents Chemother **62**, doi:10.1128/aac.01326-18 (2018).

53   Kuru, E. *et al.* Fluorescent D-amino-acids reveal bi-cellular cell wall modifications
important for Bdellovibrio bacteriovorus predation. *Nature Microbiology* **2**, 1648-
1657, doi:10.1038/s41564-017-0029-y (2017).

54   Hernández, S. B., Dörr, T., Waldor, M. K. & Cava, F. Modulation of Peptidoglycan
Synthesis by Recycled Cell Wall Tetrapeptides. *Cell Reports* **31**, 107578,
doi:https://doi.org/10.1016/j.celrep.2020.107578 (2020).

55   Barreteau, H. *et al.* Cytoplasmic steps of peptidoglycan biosynthesis. *FEMS*
Microbiology Reviews **32**, 168-207, doi:10.1111/j.1574-6976.2008.00104.x
(2008).

56   Dorr, T., Cava, F., Lam, H., Davis, B. M. & Waldor, M. K. Substrate specificity of
an elongation-specific peptidoglycan endopeptidase and its implications for cell
wall architecture and growth of Vibrio cholerae. *Mol Microbiol* **89**, 949-962,
doi:10.1111/mmi.12323 (2013).

57   Chaput, C., Labigne, A. & Boneca, I. G. Characterization of Helicobacter pylori lytic
transglycosylases Slt and MltD. *J Bacteriol* **189**, 422-429, doi:10.1128/jb.01270-06
(2007).

58 Scheurwater, E. M. & Clarke, A. J. The C-terminal Domain of Escherichia coli YfhD
Functions as a Lytic Transglycosylase*. *Journal of Biological Chemistry* **283**, 8363-
8373, doi:https://doi.org/10.1074/jbc.M710135200 (2008).
59 Mueller, E. A., Egan, A. J. F., Breukink, E., Vollmer, W. & Levin, P. A. Plasticity of
Escherichia coli cell wall metabolism promotes fitness and antibiotic resistance
across environmental conditions. *eLife* **8**, e40754, doi:10.7554/eLife.40754 (2019).
60 Yakhnina, A. A. & Bernhardt, T. G. The Tol-Pal system is required for
peptidoglycan-cleaving enzymes to complete bacterial cell division. *Proceedings*
of the National Academy of Sciences **117**, 6777-6783,
doi:10.1073/pnas.1919267117 (2020).
61 Suzuki, M., Hara, H. & Matsumoto, K. Envelope disorder of Escherichia coli cells
lacking phosphatidylglycerol. *J Bacteriol* **184**, 5418-5425,
doi:10.1128/jb.184.19.5418-5425.2002 (2002).
62 Voedts, H. *et al.* Role of endopeptidases in peptidoglycan synthesis mediated by
alternative cross-linking enzymes in Escherichia coli. *bioRxiv*,
2021.2002.2012.430937, doi:10.1101/2021.02.12.430937 (2021).
63 Heidelberg, J. F. *et al.* DNA sequence of both chromosomes of the cholera
pathogen Vibrio cholerae. *Nature* **406**, 477-483, doi:10.1038/35020000 (2000).
64 Ferrières, L. *et al.* Silent mischief: bacteriophage Mu insertions contaminate
products of Escherichia coli random mutagenesis performed using suicidal
transposon delivery plasmids mobilized by broad-host-range RP4 conjugative
machinery. *J Bacteriol* **192**, 6418-6427, doi:10.1128/jb.00621-10 (2010).
65 Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred
kilobases. *Nat Methods* **6**, 343-345, doi:10.1038/nmeth.1318 (2009).
66 Donnenberg, M. S. & Kaper, J. B. Construction of an eae deletion mutant of
enteropathogenic Escherichia coli by using a positive-selection suicide vector.
Infection and immunity **59**, 4310-4317, doi:10.1128/iai.59.12.4310-4317.1991
(1991).
67 Miyata, S. T., Unterweger, D., Rudko, S. P. & Pukatzki, S. Dual expression profile
of type VI secretion system immunity genes protects pandemic Vibrio cholerae.
PLoS pathogens **9**, e1003752-e1003752, doi:10.1371/journal.ppat.1003752
(2013).
68 Möll, A. *et al.* A D, D-carboxypeptidase is required for Vibrio cholerae
halotolerance. *Environ Microbiol* **17**, 527-540, doi:10.1111/1462-2920.12779
(2015).
69 Alvarez, L., Hernandez, S. B., de Pedro, M. A. & Cava, F. in *Bacterial Cell Wall*
Homeostasis: Methods and Protocols (ed Hee-Jeon Hong) 11-27 (Springer New
York, 2016).
70 Desmarais, S. M., De Pedro, M. A., Cava, F. & Huang, K. C. Peptidoglycan at its
peaks: how chromatographic analyses can reveal bacterial cell wall structure and
assembly. *Mol Microbiol* **89**, 1-13, doi:10.1111/mmi.12266 (2013).
955
956
957

Figures

Figure 1. A single LTG is necessary and sufficient for *V. cholerae* growth and envelope homeostasis

(A) *Trans* expression of DNA synthesis genes *vc2016-2014* permitted pAM299 disruption of native *mltG* locus. *lacZ* and *rpoB* were targeted as positive and negative controls for disruption, respectively. TE = Transformation Efficiency. ND = below limit of detection. 3 biological replicates are shown. **(B)** RlpA was depleted from the WT, Δ6, and Δ7 LTG backgrounds by placing its native promoter under control of arabinose induction and growing from a 10^{-3} overnight culture dilution into 5mL LB +/- 0.4% arabinose (ara) at 37 °C with shaking for 3 hrs, back-diluting 10^{-3} into fresh media, and incubating for another 3 hrs. Cells were imaged on LB agarose pads. Scale bars = 5µm. Dotted line indicates 10^{-3} back-dilution. **(C)** Arabinose-dependent RlpA depletion in Δ6 and Δ7 LTG backgrounds was rescued with IPTG-inducible LTGs by growing cultures in LB +/- ara (0.4%) and +/- IPTG (200µM) in 96 well plates at 37 °C without shaking for 3hrs, back-diluting 10^{-3} into fresh media, incubating another 3 hrs, and spotting directly onto the same media +kan50. Plates were incubated at 30 °C for 24hrs before imaging. Complete plating efficiencies associated with panels (B) and (C) can be found in Figure 1 – Figure Supplement 2 and Figure 1 – Figure Supplement 4, respectively. Images are representative of 3 biological replicates.

Figure 1 – Figure Supplement 1. MltG is dispensable for growth of Δ6 LTG

(A) Transformation efficiencies represent the strep[R] and kan[R] CFU divided by total viable strep[R] CFU recovered from a single mating. Streptomycin[R] (Strep[R]) *V. cholerae* WT and Δ6 LTG strains were mated with a strep[S] *E. coli* MFD donor strain carrying the pAM299 suicide vector carrying kanamycin resistance (Kan[R]) and targeting positive control gene *lacZ*, negative control gene *rpoB*, *rlpA*, or *mltG*. **(B)** The genetic region downstream of *mltG* encodes essential DNA synthesis genes. **(C)** Mean growth curves for *mltG::kan* strains in WT $P_{tac}vc2016-2014$ and Δ6 LTG $P_{tac}vc2016-2014$ backgrounds diluted 1:100 from overnight cultures into LB or LB0N and grown at 37°C. **(D)** Relative molar abundance of all AnhMurNAc-containing muropeptide species in sacculus. Overnight cultures of WT, *mltG::kan,* Δ6 LTG and Δ6 LTG *mltG::kan* (Δ7 LTG) in *lacZ::P_{tac}vc2016-2014* backgrounds were diluted 1:100 into LB+200μM IPTG and samples collected for HPLC analysis at OD_{600} 0.3 (Exp, solid bars) and OD_{600} 1.2 (Stat, striped bars). Complete muropeptide profiles can be found in Supplemental File 1. Error bars represent standard deviation from three biological replicates. Mutants were compared to the WT within each growth phase with an unpaired t-test. ** = $p<0.01$, *** = $p<0.001$, **** = $p<0.0001$.

998 **Figure 1 – Figure Supplement 2. RlpA depletion in the Δ6 and Δ7 LTG backgrounds.**
999 Native RlpA was depleted from the WT, Δ6, and Δ7 LTG backgrounds as described in
1000 Figure 1B, **(A)** imaged on LB agarose pads and **(B)** spot plated in 10-fold serial dilutions
1001 (10^0-10^{-6}) onto LB + kan50 +/- 0.4% arabinose and incubated at 30 °C for 24 hrs before
1002 imaging.

Figure 1 – Figure Supplement 3. RlpA active site architecture and localization

(A) Clustal Omega alignment of a relevant portion of lyase domain from *V. cholerae* RlpA (*vc0948*), *P. aeruginosa* RlpA (*pa4000*), and *E. coli* RlpA (*b0633*). Two Asp residues are implicated in PaRlpA activity: PaRlpA D156 aligns with VcRlpA D133 and EcRlpA D135, and PaRlpA D168 aligns with VcRlpA D145 and EcRlpA S147. WT *V. cholerae* carrying **(B)** pHL100 *rlpA-mCherry* **(C)** *rlpAΔSPOR-mCherry* or **(D)** *rlpAD145A-mCherry* was grown in M9 + 0.2% glucose at 30 °C for 2 hrs, induced with 1mM IPTG, and imaged on M9 + 0.2% glucose agarose pads at OD_{600} ~0.2. Demographs were generated using Oufti. Panels B-D are representative of 2 biological replicates. Scale bars = 5µm. **(E)** mCherry (mC) fusion proteins were detected with mCherry antibody, with mCherry fused to a DsbA signal sequence serving as a soluble mCherry size standard. RpoA loading control

1015 detected with RpoA antibody. Raw and unedited blots can be found in Figure 1 – Source
1016 Data 1.

1017

1018 **Figure 1 – Figure Supplement 4. LTGs have variable ability to sustain growth**
1019 Native RlpA depletion in Δ6 and Δ7 LTG backgrounds was rescued by single LTGs as
1020 described in Figure 1B, spot plated in 10-fold serial dilutions (from 10^0-10^{-6}) at 0 hrs and
1021 6 hrs, then incubated at 30 °C for 24hrs before imaging. Images are representative of 3
1022 biological replicates.

1023

1024 **Figure 1 – Source Data 1. Raw and uncropped mCherry Western blots**

1025

Figure 2. LTG insufficiency causes cumulative growth and morphology defects

(A) Schema describing relationship between dilution factor (DF) of saturated cultures into fresh media and time spent in exponential growth (t_{exp}). **(B)** Mass doubling times (doublings per hour, DPH) from growth curves performed in LB inoculated with 10-fold serial dilutions of saturated overnight cultures. Values were calculated from growth curves shown in Figure 2 – Figure Supplement 2. Error bars represent standard deviation of the mean, n≥3. **(C)** Relative abundance of cell morphologies from cultures at OD_{600} 0.3 from panel C. n>500 cells. Definition criteria and images are shown in Figure 2 – Figure Supplement 1. Mean length **(D)** and width **(E)** as a function of dilution factor (DF) of rod cells from panel (C). n>500 rods.

A

Morphology	Features	Poles	Septa	Length
Spheroplast	Round	N/A	N/A	N/A
Branched	N/A	3+	0+	N/A
Rod	N/A	2	0	less than 2x the median length of cells with 0 septa
Filament	N/A	2	0	greater than 2x the median length of cells with 0 septa
Doublet	N/A	2	1	N/A
Chain	N/A	2	2+	N/A

B

1037
Figure 2 – Figure Supplement 1. Definitions of morphology defect categories
(A) Cell morphology classification criteria. **(B)** Representative micrographs of cell morphologies where each cell type depicted accounts for >1% of population as sampled from cultures grown in LB to OD_{600} 0.3 from 10-fold serial dilutions of overnight cultures. Scale bars = 5 µm.

1043

Figure 2 – Figure Supplement 2. Quantitative growth and morphology of LTG-deficient mutants in LB during extended exponential phase

(A) Growth curves were performed in LB at 37°C inoculated with 10-fold serial dilutions of saturated overnight cultures. Violin plots of raw **(B)** mean width or **(C)** length of single rods. n > 500 single rods. Error bars represent standard deviation of the mean, n≥3. DF = Dilution Factor

Figure 2 – Figure Supplement 3. Morphology of LTG-deficient mutants in LB is growth-phase dependent.

Saturated overnight cultures were diluted 1/1000 into LB at 37 °C and periodically sampled for **(A)** OD$_{600}$ and imaged on LB agarose pads for **(B)** mean width, **(C)** length, and **(D)** cell morphology as exhibited in **(E)** representative micrographs. Error bars in (B) and (C) represent standard deviation of the mean, n >1 00 single rods. n > 200 cells for (D). Scale bars = 5µm.

Figure 3. LTG mutants are hypersensitive to low osmolarity and accumulation of periplasmic polymers

(A) Mass doublings (doublings per hour, DPH) upon increasing dilutions (DF, dilution factor) during growth in LB0N. Values were calculated from growth curves in Figure 3 – Figure Supplement 1. Error bars represent standard deviation of the mean, n≥3. NG, no growth **(B)** Representative growth curve in low-salt LB (LB0N) showing late-growing spontaneous suppressor in Δ6 LTG **(C)** Whole genome sequencing of Δ6 LB0N suppressor mutations identify a premature stop resulting in a 6% 3' truncation of *opgH;* a deletion resulting in a 56% 3' truncation of *opgH* and 36% 3' truncation of *vc1286;* and a deletion of the 5' end of *ptsH*. **(D)** Validation of low osmolarity growth defect suppression by an *opgH* mutation. Shown are mass doublings (doublings per hour, DPH) upon increasing dilutions (DF, dilution factor) during growth in LB0N. Values were calculated from growth curves shown in Figure 3 – Figure Supplement 1 **(E)** Saturated overnight cultures harboring IPTG-inducible *sacB* were 10-fold serially diluted and plated on LB+kan50 +/- 200μM IPTG +/- 10% sucrose, incubated at 30 °C, and imaged 24 hrs before. Representative of 3 biological replicates. Empty vector and LB0N controls are shown in Figure 3 – Figure Supplement 4.

1076

Figure 3 – Figure Supplement 1. Suppression of growth defects of ∆LTG mutants in low salt LB

(A) Mean growth curves in LB0N at 37°C started from 10-fold dilutions of saturated overnight cultures used to generate growth rates in Figure 3A. Error bars = positive standard deviation of the mean, n > 3. Growth plot on the right is reproduced from Figure 3 for comparison (B) Suppressors of the ∆6 LTG LB0N growth defect were isolated on LB and then tested for growth in LB and LB0N from a 10^{-5} inoculum of saturated overnight (LB) culture. (C) Mass doubling times of ∆*opgH* mutants calculated from (D) growth curves of ∆*opgH* mutants in LB at 37°C inoculated with 10-fold serial dilutions of saturated overnight cultures. (E) Mean growth curves in LB0N at 37°C started from 10-fold dilutions

of saturated overnight cultures used to generate growth rates in Figure 3D. error bars =
standard deviation of the mean, n=3.

Figure 3 – Figure Supplement 2. Single LTG contributions to growth in LB and low salt LB.

Mass doubling times of strains diluted 10^{-3} or 10^{-5} into LB and LB0N (+ 200µM IPTG when induction was required) normalized to the appropriate WT controls under the same growth conditions. Solid line = WT mean, Dotted lines = +/- standard deviation of WT mean. **(A)** Single LTG mutant growth normalized to WT growth. **(B)** WT, **(C)** Δ6 LTG and **(D)** Δ7 LTG carrying chromosomal *lacZ::P~tac~ ltg* normalized to WT *lacZ::P~tac~* empty. Experiments in panels (c) and (d) were carried out simultaneously and therefore share the same control

1099 replicates. Error bars represent standard deviation of 3 biological replicates. NG = No
1100 Growth. Raw curves for (A-D) and uninduced data for (B-D) available in Figure 3 – Source
1101 Data 1.

1102

Figure 3 – Figure Supplement 3. Periplasmic glucans do not contribute to division
defects in ΔLTG mutants
(a) Δ*opgH* mutants in WT, Δ6, and Δ7 backgrounds were grown in LB at 37 °C to OD_{600}
0.3 from 10^{-3} inoculum of saturated overnight culture, imaged on LB agarose pads, and
single rod lengths analyzed in MicrobeJ. Dotted lines denote quartiles. Significance
determined by unpaired Mann-Whitney tests, n>70 single rods. **(b)** T-Coffee alignment of
predicted cytoplasmic N-terminal domains from *E. coli* OpgH *(b0149)*, *Y.*
pseudotuberculosis MdoH (*yptb2493*), and *V. cholerae* OpgH (*vc1287*).

1111

Figure 3 – Figure Supplement 4. SacB expression is toxic in ΔLTG mutants in a sucrose-dependent manner.

Saturated overnight cultures harboring **(a)** IPTG-inducible *sacB* or **(b)** empty vector were 10-fold serially diluted (10^0-10^{-6}) and plated on LB+kan50 and LB0N+kan50 +/- IPTG (200µM) + sucrose (0, 1,or 10% W/V) and incubated at 30 °C for 24 hrs before imaging. Representative of 3 biological replicates.

Figure 3 – Source Data 1. Raw growth curve data for single LTG mutants and LTG complementation in LTG-deficient mutants.

Presents raw OD_{600} measurements for growth curves performed in LB and low-salt LB (LB0N) with and without induction of *trans*-complementation by single LTGs in LTG-deficient mutant backgrounds, as well as growth curves in LB and LB0N of single LTG-deficient mutants. n≥3

Figure 4. LTG mutants are hypersensitive to antibiotics promoting periplasmic PG accumulation

(A) Sensitivity to Penicillin G (PenG), aztreonam (AZTM), mecillinam (Mec), cefsulodin (Cef), moenomycin (Moeno), MP265, and Fosfomycin (Fos) measured as Zone of Inhibition (ZOI) in a disk diffusion assay. ND = No ZOI around disk. Error bars = standard deviation. Significance determined by one-way Anova. ns = $p>0.05$, * = $p< 0.05$, ** = $p<0.01$, *** = $p<0.001$, **** = $p<0.0001$. Overnight cultures were diluted 1:100 into LB+BADA (100µM) and grown at 37°C to OD_{600} 0.5 before addition of **(B)** PenG (100µg ml^{-1}) or **(C)** Fosfomycin (500µg ml^{-1}). Resulting spheroplasts were washed and imaged after 3hrs of antibiotic exposure. Fluorescence was normalized to the same intensity threshold for visual comparison except where indicated (Exceptionally bright samples were normalized to a higher intensity threshold denoted by the multiplier). Representative of 2 biological replicates. Scale bar = 5µm

1140

Figure 4 – Figure Supplement 1. Periplasmic cell wall accumulation in response to Penicillin G.

Overnight cultures were diluted 1:100 into LB+BADA (100μM) and grown at 37°C to OD_{600}0.5 before addition of PenG (100μg ml^{-1}). Resulting spheroplasts were removed periodically, washed, and imaged on LB agarose. **(A)** Fluorescence was normalized to the same intensity threshold in Figure 4 – Figure Supplement 1 and Figure 4 – Figure Supplement 2 for comparison except where indicated (Exceptionally bright samples were normalized to a higher intensity threshold denoted by the multiplier). Representative of 2 biological replicates. Scale bar = 5μm **(B)** Mean fluorescence intensity (AU) for >100 cells

1150 or spheroplasts were measured in ImageJ. N/A = not applicable due to insufficient intact
1151 spheroplasts for measurement.
1152

Figure 4 – Figure Supplement 2. Lack of periplasmic cell wall accumulation in response to Fosfomycin.

Overnight cultures were diluted 1:100 into LB+BADA (100µM) and grown at 37°C to OD_{600} 0.5 before addition of Fosfomycin (500µg ml[-1]). Resulting spheroplasts were removed periodically, washed, and imaged on LB agarose. **(A)** Fluorescence was normalized to the same intensity threshold in Figure 4 – Figure Supplement 1 and Figure

1160 4 – Figure Supplement 2 for comparison. Representative of 2 biological replicates. Scale
1161 bar = 5µm. **(B)** Mean fluorescence intensity (AU) for >100 cells or spheroplasts were
1162 measured in ImageJ.

1163
Figure 4 – Figure Supplement 3. Defects of LTG-deficient mutants are independent
of PG recycling.
$\Delta ampG$ data for this figure was collected during the same experiments in Figures 2A and
3A-F, therefore WT data are reproduced from those figures and any statistical tests
shown here were performed on the complete data sets for those experiments (including
WT, $\Delta ampG$, and ΔLTG strains). **(A-B)** Mass doubling times from growth curves
performed in LB or LB0N inoculated with 10-fold serial dilutions (10^0-10^{-6}) of saturated
overnight cultures. Error bars represent standard deviation of the mean, n≥3. **(C)**
Saturated overnight cultures harboring IPTG-inducible *sacB* or empty vector were 10-fold
serially diluted (10 and plated on LB+kan50 and LB0N+kan50 +/- IPTG (200µM) +

sucrose (0, 1, or 10% W/V) and incubated at 30 °C for 24 hrs before imaging.
Representative of 3 biological replicates. **(D)** Sensitivity to Penicillin G (PenG), aztreonam
(AZTM), mecillinam (Mec), cefsulodin (Cef), moenomycin (Moeno), MP265, and
Fosfomycin (Fos) measured as Zone of Inhibition (ZOI) in a disk diffusion assay. ND =
No ZOI around disk. Error bars = standard deviation. Significance determined by one-
way Anova. ns = $p > 0.05$. Overnight cultures were diluted 1:100 into LB+BADA (100μM)
and grown at 37°C to OD_{600} 0.5 before addition of **(E)** PenG (100μg ml^{-1}) or **(F)** Fosfomycin
(500μg ml^{-1}). Samples were washed and imaged after 3hrs of antibiotic exposure.
Fluorescence was normalized to the same intensity threshold as Figures S12 and S13
for comparison. Representative of 2 biological replicates. Scale bar = 5μm. **(G-H)** Mean
fluorescence intensity (AU) for >100 BADA-labeled cells or spheroplasts were measured
in ImageJ.

1186
Figure 5. Periplasmic uncrosslinked PG strands accumulate in an endopeptidase-
dependent manner during normal growth
(A) Schema describing muramidase-treatment dependence of detection of monomeric or
polymeric PG fragments. M4N = anhMurNAc-tetrapeptide. M4 = reduced MurNAc-
tetrapeptide. **(B-E)** Overnight cultures of. WT and ΔLTG mutants were diluted 1:100 into

LB, grown at 37°C, and harvested at OD_{600} 0.3 (Exp, solid bars) and 1.2 (Stat, striped
bars) for soluble PG analysis by LC-MS. MS peak areas for M4N and M4 are shown here
and complete muropeptide profiles can be found in Supplemental File 2.   Means
compared to WT by unpaired t-tests, n=3. **(F)** WT and Δ6 LTG strains harboring a single
chromosomal copy of *shyA* under an IPTG-inducible promoter were grown from 10^{-2}
inocula for 3hrs (OD_{600} ~1.0) in LB with (ShyA+) or without (ShyA -) 200µM IPTG at 37
°C and harvested for soluble PG analysis by LC-MS. Complete muropeptide profiles can
be found in Supplemental File 3. Means compared by unpaired t-test, n=3. All error bars
= standard deviation. ns = p>0.01, ** = p<0.01, *** = p<0.001, **** = p<0.0001. ND = Not
detected in all replicates.

1202
Figure 5 – Figure Supplement 1. Endopeptidase depletion phenotypes in LTG-
deficient mutants.
WT and Δ6 LTG strains harboring a single chromosomal copy of *shyA* under an IPTG-
inducible promoter were grown from 10^{-2} inocula for 3hrs (OD_{600} ~1.0) in LB +/- IPTG
(200µM) at 37 °C. Samples were **(A)** imaged on LB + agarose pads and **(B)** spot plated

in 10-fold serial dilutions (10^0-10^{-6}) onto LB+/- IPTG (200μM), then incubated at 30 °C for
24hrs before imaging. Representative of 3 biological replicates.

1210
1211 Figure 6. Model for LTG-mediated removal of toxic PG debris
1212 **(A)** Endopeptidases (EPs, Yellow) excise PG strands (Red) from the sacculus (Blue),
1213 permitting sacculus expansion. **(B)** In wild-type cells, lytic transglycosylases (LTGs,
1214 Green) digest excised, uncrosslinked PG strands into smaller fragments that can be
1215 recycled by AmpG (Black) or released through porins (Violet) **(C)** In LTG deficient cells,
1216 excised PG debris crowds the periplasm and becomes toxic.

1217

1218 Supplemental File 1. Complete sacculus muropeptide composition profiles of LTG-
1219 deficient mutants
1220 Supports Figure 1 – Figure supplement 1D. Presents mean relative muropeptide
1221 abundance +/- standard deviation of the mean for all detectable muropeptide species
1222 from muramidase-treated sacculi isolated by ultracentrifugation from LTG-deficient
1223 mutant cultures during exponential and stationary growth phases. n = 3.

1224

1225 Supplemental File 2. Complete muropeptide detection profiles of soluble PG
1226 material from LTG-deficient mutants
1227 Supports Figure 5BCDE. Presents mean mass peak area +/- standard deviation of the
1228 mean for all soluble, detectable muropeptide species from muramidase treated and
1229 untreated lysates of LTG-deficient mutant cultures during exponential and stationary
1230 growth phases. n = 3.

1231

1232 Supplemental File 3. Complete muropeptide detection profiles of soluble PG
1233 material from ShyA-depleted mutants
1234 Supports Figure 5F. Presents mean mass peak area +/- standard deviation of the mean
1235 for all soluble, detectable muropeptide species from muramidase treated and untreated
1236 lysates of cultures grown with and without ShyA induction. n = 3

1237

1238 **Supplemental File 4. Bacterial strains, plasmids, and primers used in this study**