Title: spatialGE: Quantification and visualization of the tumor microenvironment heterogeneity

using spatial transcriptomics

Oscar E. Ospina

Department of Biostatistics and Bioinformatics, Moffitt Cancer Center, Tampa, FL 33612 USA

Christopher M. Wilson

Department of Biostatistics and Bioinformatics, Moffitt Cancer Center, Tampa, FL 33612, USA

Alex C. Soupir

Department of Biostatistics and Bioinformatics, Moffitt Cancer Center, Tampa, FL 33612, USA

Anders Berglund

Department of Biostatistics and Bioinformatics, Moffitt Cancer Center, Tampa, FL 33612, USA

Inna Smalley

Department of Tumor Biology, Moffitt Cancer Center, Tampa, FL 33612, USA

Kenneth Y. Tsai

Department of Anatomic Pathology, Moffitt Cancer Center, Tampa, FL 33612, USA

Brooke L. Fridley*

Department of Biostatistics and Bioinformatics, Moffitt Cancer Center, Tampa, FL 33612, USA

*Contact: Brooke.Fridley@Moffitt.org

Abstract

Summary: Spatially-resolved transcriptomics promises to increase our understanding of the

tumor microenvironment and improve cancer prognosis and therapies. Nonetheless, analytical

methods to explore associations between the spatial heterogeneity of the tumor and clinical data

are not available. Hence, we have developed spatialGE, a software that provides visualizations

and quantification of the tumor microenvironment heterogeneity through gene expression

surfaces, spatial heterogeneity statistics (SThet) that can be compared against clinical

information, spot-level cell deconvolution, and spatially-informed clustering (STclust), all using

a new data object to store data and resulting analyses simultaneously.

Availability and implementation: The R package and tutorial/vignette are available at

https://github.com/FridleyLab/spatialGE. A script to reproduce the analyses in this manuscript is

available in Supplementary information.

Contact: Fridley.Lab@Moffitt.org or Brooke.Fridley@Moffitt.org

Supplementary information: Available at Bioinformatics online.

42 1 Introduction

Spatially-resolved transcriptomics (ST) has allowed a better understanding of the tumor

microenvironment (TME), immune infiltration, and its relationship with immunotherapy

response (Tang, et al., 2016), as well as promising to improve the development of cancer

prognoses and therapies (Nederlof, et al., 2021). Many ST methods involve mRNA probe

hybridization in immobilized tissues on surfaces (Zhang, et al., 2021). Hence, gene expression

localization within the tissue architecture is preserved as opposed to other transcriptomic

approaches, such as bulk RNA (RNA-seq) or single-cell RNA sequencing (scRNAseq)

(Maniatis, et al., 2021; Yu, et al., 2021). Given the high-dimensional nature of ST experiments,

data structures are needed to accommodate the gene expression abundance, spatial locations, and

clinical information for samples taken from multiple subjects. Similarly, methods for ST

visualization and analysis are needed, with existing tools varying in flexibility and approaches

(Bergenstrahle, et al., 2020; Dries, et al., 2021; Hao, et al., 2021; Navarro, et al., 2017; Sun, et

al., 2020; Tan, et al., 2020; Zhao, et al., 2021).

To meet these unmet computational needs, we have developed a pipeline for the

processing and analysis of probe-based NGS ST experiments and exploration of the TME, which

we call spatialGE. Novel features of the software include the generation of transcriptomic

surfaces; quantification of spatial heterogeneity and the association with clinical information;

gene expression deconvolution; spatially-informed clustering (STclust); and a new data structure

to store multi-sample spatial data, subject-level metadata, and analytical results.

2 Implementation

Our new software takes gene expression and spatial location data as input, and optionally,

associated metadata (i.e., clinical information), and stores it in a new R object class referred to as

'STList'. Other existing data structures can store metadata, however the STList takes a single

table with just one row of information for each sample. This approach makes inputting data

easier for users and allows spatialGE to complete downstream analyses with few lines of code.

Users also have the possibility to provide Visium outputs directly from space ranger or Seurat

objects, as well as .dcc/.pkc files from GeoMx experiments. Additionally, results from spatialGE

analyses are saved in the STList making downstream statistical analysis and visualization of

results seamless. The software allows users to perform gene- or spot-wise filtering and generate

quality control visualizations, followed by library size normalization and logarithmic or voom

transformation (Law, et al., 2014). The relative gene expression at each spot can be visualized

using "quilt plots" (Fig. 1A). If the STList is created from space ranger outputs or Seurat objects,

histological images are imported automatically, allowing to plot gene expression and these

images side by side for comparative analysis. Higher resolution plots are achieved with

transcriptomic surfaces produced by spatial interpolation ("kriging"; Fig. 1B) (Diggle and

Ribeiro, 2007). Surfaces can also be generated for deconvoluted cell type scores (Fig. 1C), with

cell type compositions at each spot inferred via gene expression deconvolution in xCell (Aran, et

al., 2017). Lastly, tumor purity scores are generated using ESTIMATE (Yoshihara, et al., 2013),

followed by tumor/stroma classification using model-based clustering (Fig 1A-C) (Scrucca, et

al., 2016). See Supplementary methods for additional details.

spatialGE also provides users with spatial statistics to quantify TME heterogeneity

(Supplementary Table 1; Supplementary Fig. 1). The Moran's I (Moran, 1950) and Geary's C

(Geary, 1954) allow users to ascertain whether gene expression is uniform throughout the tissue.

The Getis-Ord G_i statistic measures the tendency of a gene to produce expression hot- or cold-

spots (Getis and Ord, 2010). Users can associate per-sample spatial heterogeneity statistics with

other measurable clinical outcomes present in the sample metadata (SThet; Fig 1D). Finally,

spatialGE includes a computationally efficient spatially-informed unsupervised clustering

method, referred to as STclust, to detect TME compartments or "niches" (Fig. 1E). First, the

approach computes principal components (PCs) from gene counts. Then, a distance matrix is

computed based on two scaled distance matrices: 1) transcriptomic autocorrelation between spots

using the PCs (D_1), and 2) spatial distances between spots (D_2). Next, the autocorrelation matrix

(D_1) is "shrunk" toward the spatial distances matrix (D_2) by calculating its weighted average as

$D = [(1 - w) * D_1] + (w * D_2)$. The user specifies the weight (w) to apply. Based on our

experience, a weight between 0.025 and 0.25 seems to best capture tissue heterogeneity

(Supplementary Figs. 2 and 3).

3 Discussion

The spatialGE package is a comprehensive analytical R package for the simultaneous analysis of

multiple tissues assayed with probe-based ST technologies (i.e., Visium, GeoMx) using the new

STList R object class. spatialGE is unique in implementing kriging to generate gene expression

surfaces at high resolution for multiple ST tissue sections simultaneously (for an alternative

method see (Zhao, et al., 2021)). Several ST clustering approaches are already available

(Supplementary Table 2) (Bergenstrahle, et al., 2020; Dries, et al., 2021; Tan, et al., 2020; Zhao,

et al., 2021), however our spatially-informed clustering approach, STclust, is computationally

efficient (Supplementary Fig. 4) and resembles tissue features by only weighting transcriptomic

similarities by their distances between spots. A comparison of Louvain clustering as

implemented in the widely used software Seurat to STclust, showed a Rand similarity index of

0.8 for k=4 (Supplementary Fig. 5), indicating good agreement of spot assignments between the

two methods. Another innovation of spatialGE is the estimation of quantifiable metrics (spatial

heterogeneity statistics, SThet) to capture transcriptomic complexity and the ability to compare

them against clinical outcomes.

Applying spatialGE to the Thrane et al. melanoma ST data, we observed agreement

between the transcriptomic surfaces (Supplementary Fig. 6) and tumor and stroma regions

observed in pathology images (see (Thrane, et al., 2018)). STclust provided TME compartments

(i.e., clusters; Supplementary Figs. 2 and 3) that resembled the features annotated in pathology

images (Thrane, et al., 2018). Finally, we observed an association between spatial heterogeneity

and patient survival, providing support to the hypothesis that heterogeneity is a predictor of

patient outcomes (Nederlof, et al., 2021).

In future versions, spatialGE will include analysis tools for different ST technologies

(e.g., Slide-seq). We are currently developing additional analytical methods and visualizations

that use histological annotation tools for non-gridded technologies (e.g, GeoMx), as well as a

web-based implementation of spatialGE. Another developing area is the integration of scRNA-

seq data with ST for spot-level cell phenotyping.

Acknowledgments

Funding: This work was supported by the National Institutes of Health (NIH) [T32-CA233399,

P30-CA076292, R00-CA226679]. This content is solely the responsibility of the authors and

does not necessarily represent the official view of the NIH. We thank the spatial research group

at the KTH Royal Institute of Technology – SciLifeLab (Stockholm, Sweden) for making their

melanoma data available.

Conflicts of interests: none declared.

References

Aran, D., Hu, Z. and Butte, A.J. xCell: digitally portraying the tissue cellular heterogeneity

landscape. *Genome Biol* 2017;18(1):220.

Bergenstrahle, J., Larsson, L. and Lundeberg, J. Seamless integration of image and molecular

analysis for spatial transcriptomics workflows. *BMC Genomics* 2020;21(1):482.

Diggle, P.J. and Ribeiro, P.J. Spatial prediction. In, *Model-based Geostatistics*. New York, NY:

Springer New York; 2007. p. 134-156.

Dries, R*., et al.* Giotto: a toolbox for integrative analysis and visualization of spatial expression

data. *Genome Biol* 2021;22(1):78.

Geary, R.C. The contiguity ratio and statistical mapping. *The incorporated statistician*

1954;5(3):115-146.

Getis, A. and Ord, J.K. The analysis of spatial association by use of distance statistics. In,

Perspectives on spatial data analysis. Springer; 2010. p. 127-145.

Hao, Y*., et al.* Integrated analysis of multimodal single-cell data. *Cell* 2021;184(13):3573-3587

e3529.

Law, C.W*., et al.* voom: Precision weights unlock linear model analysis tools for RNA-seq read

counts. *Genome Biol* 2014;15(2):R29.

Maniatis, S., Petrescu, J. and Phatnani, H. Spatially resolved transcriptomics and its applications

in cancer. *Curr Opin Genet Dev* 2021;66:70-77.

Moran, P.A. Notes on continuous stochastic phenomena. *Biometrika* 1950;37(1/2):17-23.

Navarro, J.F*., et al.* ST Pipeline: an automated pipeline for spatial mapping of unique transcripts.

Bioinformatics 2017;33(16):2591-2593.

Nederlof, I*., et al.* A High-Dimensional Window into the Micro-Environment of Triple Negative

Breast Cancer. *Cancers (Basel)* 2021;13(2).

Scrucca, L*., et al.* mclust 5: Clustering, Classification and Density Estimation Using Gaussian

Finite Mixture Models. *R Journal* 2016;8(1):289-317.

Sun, S., Zhu, J. and Zhou, X. Statistical analysis of spatial expression patterns for spatially

resolved transcriptomic studies. *Nat Methods* 2020;17(2):193-200.

Tan, X*., et al.* SpaCell: integrating tissue morphology and spatial gene expression to predict

disease cells. *Bioinformatics* 2020;36(7):2293-2294.

Tang, H., Qiao, J. and Fu, Y.X. Immunotherapy and tumor microenvironment. *Cancer Lett*

2016;370(1):85-90.

Thrane, K*., et al.* Spatially Resolved Transcriptomics Enables Dissection of Genetic

Heterogeneity in Stage III Cutaneous Malignant Melanoma. *Cancer Res* 2018;78(20):5970-5979.

Yoshihara, K*., et al.* Inferring tumour purity and stromal and immune cell admixture from

expression data. *Nat Commun* 2013;4:2612.

Yu, X*., et al.* Statistical and Bioinformatics Analysis of Data from Bulk and Single-Cell RNA

Sequencing Experiments. *Methods Mol Biol* 2021;2194:143-175.

Zhang, M*., et al.* Spatial molecular profiling: platforms, applications and analysis tools. *Brief*

Bioinform 2021;22(3).

Zhao, E*., et al.* Spatial transcriptomics at subspot resolution with BayesSpace. *Nat Biotechnol*

2021.

Figure captions

Graphical abstract. Overview of spatialGE features. **A.** The STList data object from spatialGE

can be creared from several sources, including comma- or tab-separated files containing gene

counts and spatial coordinates. The object can also be created directly from Visium outputs,

Seurat objects, or GeoMx outputs. **B.** Users can optionally provide a metadata file, containing

information associated with each sample (one row per sample, or per ROI if GeoMx data). **C.**

Methods for quality control of data are provided by spatialGE, including visualizations of counts

and genes per spot, as well as filtering of spots or genes within user-determined thresholds. **D.** A

novel method (STclust) performs spatially informed clustering of spots and tissue domain

identification. **E.** spatialGE provides different types of data visualization, including gene

expression at each spot ("quilt plots"), as well as adaptation of spatial interpolation ("kriging") to

spatial transcriptomics data (transcriptomic surface). **F.** spatialGE also leverages spatial statistics

(Moran's I, Geary's C, Getis-Ord Gi) to quantitatively describe heterogeneity within the tumor

microenvironment and to explore associations between spatial heterogeneity and clinical

oucomes. **G.** Gene expression deconvolution can also be applied to each spot to detect immune

cell types (xCell) and classification of spots as tumor or stroma (ESTIMATE).

Figure 1. Results from spatialGE analysis of a melanoma stage IIIc using spatial transcriptomics

(patient 1, sample 2 (Thrane, et al., 2018)). **A.** Quilt plot showing log-expression at each spot for

immune gene *CD74*, with squares and crosses indicating tumor and stromal spot classifications

respectively. **B.** Transcriptomic surface of *CD74* expression allows for a clearer visualization of

the tissue heterogeneity, with gray squares indicating tumor compartment. **C.** The inferred

abundance of B cells in the tissue can be visualized by the predicted surface. **D.** Scatter plots

showing the relationships between spatial statistics calculated for *CD74* and the survival of

patients in months from (Thrane, et al., 2018). The colored arrows provide a guide for the

interpretation of the spatial statistics. The dot with black border indicates the tissue section from

patient 1 featured in the other panels in Fig 1. **E.** Tumor/stroma assignments and spatially-

informed clusters with STclust (using a spatial weight of 0.025) are shown. The yellow (1) and

blue (2) clusters roughly represent tumor and stroma regions of the tissue, respectively. The red

cluster (3) likely correspond to spots showing immune activity given the topological match with

CD74 (Fig. 1B) and B cell scores (see Fig. 1**C**).

Supplementary Figure captions

Supplementary Fig. 1. Relationship between the spatial autocorrelation statistics and standard deviation of gene expression. The red dot represents *CD74*, and asterisk (*) indicates a significant relationship between the spatial statistic and the standard deviation.

Supplementary Fig. 2. Cluster assignment of the spots as generated with STclust and applying automatic dynamic tree cutting. Spatial weights (*w*) from 0.0 to 0.5 were evaluated. **A.** *w*=0; **B.** *w*=0.025; **C.** *w*=0.05; **D.** *w*=0.0.75; **E.** *w*=0.1; **F.** *w*=0.15; **G.** *w*=0.2; **H.** *w*=0.5.

Supplementary Fig. 3. Cluster assignment of the spots as generated with STClust using different values of k, k=2 to 5. **A.** k=2; **B.** k=3; **C.** k=4; **D.** k=5. Each page in this file shows clusters at different spatial weights from *w*=0.0 to *w*=0.5.

Supplementary Fig. 4. Execution time of unsupervised spatially-informed clustering with STclust (k=4) in relation to the number of spots from each of the tissue slices in (Thrane, et al., 2018). Labels accompanying points indicate the patient and replicate numbers.

Supplementary Fig. 5. Results Louvain clustering using Seurat and STclust with k=4 and spatial weights *w*=0 (**A**); *w*=0.025 (**B**); *w*=0.05 (**C**); *w*=0.075 (**D**); *w*=0.1 (**E**); *w*=0.15 (**F**); *w*=0.2; (**G**); and *w*=0.5 (**H**). Louvain clustering used a resolution parameter=0.8 (**I**). Clustering was applied to tissue slice 2 from patient 1 (Thrane, et al., 2018). Rand indexes (R) are shown for the comparison between STclust (**A-H**) and Louvain clustering (**I**). The red boxes indicate the two most concordant clustering solutions (R=0.81).

236 **Supplementary Fig. 6.** Transcriptomic surfaces resulting from spatial interpolation ("kriging")

237 of *PMEL* (**A**), *SPP1* (**B**), *CD74* (**C**), *IGLL5* (**D**), *SOX10* (**E**), and *SOX4* (**F**). The same genes are

238 shown for all the tissue slices in (Thrane, et al., 2018).

A *CD74*, patient 1 (sample 2)

B *CD74*, patient 1 (sample 2)
Morans I=0.18* Gearys C=0.81*
GetisOrd Gi=0.1591*

C B cells, patient 1 (sample 2)

+ stroma
■ tumor

log expr
0   1   2   3

pred_expr   □ tumor

pred_score   □ tumor
0.2   0.4   0.6

D *CD74*   Patients ● 1 ● 2 ● 3 ● 4

Survival (months)

Moran's I

Uniform
expression   Clustered
expression

Geary's C

Clustered
expression   Uniform
expression

Getis-ord Gi

More
"cold regions"   More
"hot regions"

E Cluster assignments

Clusters
1   2

+ stroma
■ tumor