Phosphate starvation response precedes drought response in field-grown plants

Yukari Nagatoshi[1], Kenta Ikazaki[2], Nobuyuki Mizuno[3†], Yasufumi Kobayashi[1], Kenichiro

Fujii[1], Eri Ogiso-Tanaka[4‡], Masao Ishimoto[4], Yasuo Yasui[3], Tetsuji Oya[2], Yasunari Fujita[1,5*]

[1] Biological Resources and Post-harvest Division, Japan International Research Center for

Agricultural Sciences (JIRCAS); Tsukuba, Ibaraki 305-8686, Japan.

[2] Crop, Livestock and Environment Division, Japan International Research Center for

Agricultural Sciences (JIRCAS); Tsukuba, Ibaraki 305-8686, Japan.

[3] Graduate School of Agriculture, Kyoto University; Kyoto, Kyoto 606-8502, Japan.

[4] Institute of Crop Science, National Agriculture and Food Research Organization (NARO);

Tsukuba, Ibaraki 305-8518, Japan.

[5] Graduate School of Life Environmental Science, University of Tsukuba; Tsukuba, Ibaraki

305-8572, Japan.

[†]Present address: Institute of Crop Science, NARO; Tsukuba, Ibaraki 305-8518, Japan.

[‡]Present address: Center for Molecular Biodiversity Research, National Museum of Nature and

Science; Tsukuba, Ibaraki 305-0005, Japan.

Yasunari Fujita[1,5*]

Biological Resources and Post-harvest Division, JIRCAS; 1-1 Ohwashi, Tsukuba, Ibaraki 305-

8686, Japan.

[*]Corresponding author. E-mail: yasuf@affrc.go.jp

Abstract:

Drought severely damages crop production, even under conditions so mild that the leaves show

no signs of wilting. As effective methods for analyzing the field drought response have not

been established, it is unclear how field-grown plants respond to mild drought. We show that

ridges are a useful experimental tool to mimic mild drought stress in the field. Mild drought

reduces inorganic phosphate levels in the leaves to activate the phosphate starvation response

(PSR) in field-grown soybean plants. PSR-related gene expression is mainly observed under

drought conditions that are too mild to activate abscisic acid-mediated gene expression. Thus,

our study provides insights into the molecular response to mild drought in field-grown plants

and into the link between nutritional and drought stress responses in plants.

One-Sentence Summary:

Mimicking mild drought using ridges reveals that the phosphate response is a sign of drought reducing growth in plants.

Introduction:

Drought is a major threat to global crop production. Although tremendous progress has been made in deciphering the molecular mechanism of the drought stress response in model plants grown under controlled conditions (*1-3*), little is known about this mechanism in crops grown in the field, where conditions fluctuate and are difficult to predict and control. As a major oil and protein source for food, feed, biodiesel, and industrial products, soybean (*Glycine max* L. Merr.) is the most economically important dicot crop in the world (*4*). However, since soybean is cultivated mostly under rainfed conditions, soybean production is severely constrained by abiotic stresses, particularly drought (*5*). Even mild drought stress that occurs without obvious outward signs of a stress response such as leaf wilting poses a major threat to soybean productivity (*6*). Given that strategies to increase arable land area and yield face major obstacles, identifying ways to reduce crop production losses due to mild drought stress is a key goal in attaining global food security. However, since it is challenging to induce drought conditions to conduct experiments in the field due to uncontrollable weather, effective methods for analyzing the field drought response have not been developed. Here, using soybean as a model field-grown crop, we established that ridges could be used to induce conditions that mimic mild drought stress. We then used this system to explore the mechanisms underlying the plant's response to such conditions.

Results and Discussion:

Each growing season for six consecutive years, we created flat and ridged plots (referred to as "flats" and "ridges", respectively) to conduct mild drought tests in the field. We monitored the volumetric water content (VWC) of the soil using soil moisture sensors (Fig. 1, A and B, and figs. S1 to S3). In 2016, the soil VWC in the flats of our experimental field fluctuated between 34% (–170 kPa) and 49% (–4 kPa) (Fig. 1C and fig. S4). Although the VWC was strongly affected by rainfall (Fig. 1C), the VWC in ridges (ridge height, 30 cm; Fig. 1, A and B) was consistently lower than that of the flats and fluctuated between 27% (–530 kPa) and 41% (–44 kPa) (Fig. 1C and fig. S4). We compared the growth of soybean grown on the flats and ridges. The aboveground biomass of the plants grown on ridges was clearly reduced compared with that grown on flats, even though leaf wilting was not observed in either group of plants (Fig.

1D and fig. S5). Thermal image analysis of the soybean plants showed that the leaf temperature

of the plants grown on ridges was higher than that of plants grown on flats (Fig. 1E), suggesting

that transpiration was lower in the plants grown on ridges, which may have reduced the

photosynthetic rate and biomass. Consequently, the yield of soybean grown on ridges was also

reduced compared with that of soybean grown on flats (Fig. 1, F and G, fig S6). A decrease in

transpiration and growth is a typical response of plants to water deficit, suggesting that growth

on ridges induces mild drought stress in field-grown plants.

To ascertain whether the reduced growth of the plants grown on ridges was indeed

due to water deficit or to changes in the nutrient composition of the soil, we measured the

contents of nitrogen (N), phosphorous (P), and potassium (K) in the flats and ridges. There was

no significant difference in the contents of these nutrients between the flats and ridges, both at

the beginning and the end of the soybean growing season (Fig. 1H). We thus next examined

whether watering the ridges would compensate for the negative effect of ridges on plant growth

(Fig. 2 and fig. S7). In rainfed plants, there was a noticeable difference in soybean growth on

flats and ridges, whereas in the irrigated plants, the difference was reduced or barely detectable

(Fig. 2 and fig. S7, B to E). Thus, the negative effect of ridges on plant growth was

complemented by irrigation, indicating that the reduced growth of the plants on ridges was

mainly caused by water deficit. While a water deficit was not adequately induced on ridges in

2017, the results of the remaining five years of field trials demonstrated that the growth

reduction of soybean on ridges compared to flats was a result of reduced VWC (Fig. 1 and figs.

S7 to S9). Together, these observations demonstrate that ridges are a valuable tool to induce

conditions that mimic mild drought stress.

To investigate how mild drought stress affects the expression of genes in field-grown

soybean plants, we analyzed the fully expanded second trifoliate leaves of 29-day-old pre-

flowering soybean plants in the 2015 field experiment by RNA sequencing (RNA-seq) (Fig. 3,

A and B, fig. S9). We identified 3,045 differentially expressed genes (DEGs) between the

soybean plants grown on flats and ridges (table S1), which had soil VWCs at sampling time of

41% (–26 kPa) and 34% (–123 kPa), respectively (figs. S4 and S9). We identified 990 and 400

DEGs that are up- and down-regulated ($|\log_2(FC)| \geq 1$, FPKM value > 0, $q < 0.05$), respectively,

under mild drought conditions (Fig. 3C and tables S2 and S3). Gene ontology (GO) analysis

(FDR < 0.05) revealed a total of 53 GO terms that were significantly enriched in up-regulated

DEGs, while no GO terms were significantly enriched in down-regulated DEGs (table S4).

Notably, 16 and 7 of the up-regulated DEGs were related to "cellular response to phosphate

starvation" and "phosphate ion homeostasis", respectively (fig. S10). Furthermore, 11.4% of

the up-regulated DEGs were significantly enriched in phosphate starvation response (PSR)

family genes, including those encoding signaling factors, phosphatases, and transporters, and in genes implicated in lipid metabolism, in *Arabidopsis thaliana* (*7*) and in soybean (*8*, *9*) (Fig. 3, D and E, and tables S5 and S6), indicating that a battery of PSR genes were up-regulated under mild drought conditions. Mild drought stress tended to be induced the expression of PSR marker genes in not only the second but also the first and third trifoliate leaves in the field-grown soybean plants (Fig. 3F).

To determine whether reduced P levels are indeed observed under mild drought conditions in the field, we determined the inorganic phosphate (Pi) concentration of the first to third trifoliate leaves. The Pi concentrations in leaves at each of these positions in soybean plants grown on ridges were significantly decreased compared with those grown on flats (Fig. 3G), indicating that mild drought stress decreases Pi levels in the leaves of field-grown soybean plants. Our collective observations (Fig. 1, C and E, figs. S7 to S9) suggest that reduced transpiration and water availability induced by mild drought inhibit Pi uptake by plants, leading to Pi deficiency conditions. Combined with a previous meta-analysis using various plants and soil types (*10,11*) and physiological experiments with soybean grown in pots (*12*), these observations support the notion that mild drought stress reduces Pi levels to induce the PSR in field-grown plants.

We next explored if the link between mild drought stress and the PSR identified in our field experiments could be reproduced in pot trials (Fig. 4). When the primary soybean leaves were fully expanded at 10 days after sowing in pots containing the same soil as that used in the field experiments, we started the drought stress treatments with five different levels of VWC, α to ε (increasing water deficit; Fig. 4A). The leaf surface temperatures of 26-day-old soybean plants visualized by infrared thermography paralleled the soil VWCs in the pots (Fig. 4, A and B), suggesting that the lower the VWC, the greater the reduction in transpiration under the experimental condition. We analyzed the gene expression profiles of the fully expanded first trifoliate leaves of 38-day-old seedlings grown in pots. The expression levels of PSR genes were largely inversely associated with VWC levels of α to δ, but not ε (Fig. 4C). Thus, the soil water status determines the degree of the PSR under mild drought stress conditions, supporting the findings of our field experiments. Interestingly, between pots δ and ε, even though the VWC decreased, PSR gene expression did not increase, but the expression of abscisic acid (ABA)-responsive genes, which are established markers of the response to severe drought stress (*13,14*), did increase (Fig. 4, C and D). Thus, the PSR, and not ABA, is an important indicator of mild drought stress in the field (Fig. 4E).

Here, we reveal a novel link between the PSR and drought response in plants. This insight was possible because the experiments were conducted in an actual field, where available

P is not abundant and P acquisition is dependent on the biotic and abiotic environment (*15*).
While ridges are widely used in crop production for various purposes, they have not been used
to explore the drought response in plants. Our research using ridges provides insight into the
link between plant nutritional and drought stress responses and paves the way for elucidating
the molecular mechanism of the drought response in field-grown plants. Since it is easy and
inexpensive to create ridges in experimental fields, this method of inducing mild drought stress
overcomes a major research barrier, and is expected to facilitate efforts around the world aimed
at deciphering the response of field-grown crops to various degrees of drought stress. Global
food security depends on producing crops that can maintain productivity even under mild to
moderate soil water deficits (*16*, *17*). The PSR could be used as an early marker of mild drought
stress, allowing irrigation to be optimized before the yield of field-grown crops is affected by
mild drought conditions.
153
Materials and Methods:
Field experiment
Soybean (*Glycine max* L. Merr.) cv. Williams 82, a sequenced model US cultivar (*18*), was
used in this study for all years other than 2019. Soybean cv. Enrei, a sequenced model Japanese
cultivar (*19*), was used only in the 2019 field experiment. Field experiments were conducted
at the experimental field at the Japan International Research Center for Agricultural Sciences
(JIRCAS, 36°03′14″ N, 140°04′46″ E, 24 m above sea level) located in Ibaraki, Japan, from
2015 to 2020. The soil type is Andosol (*20*) and the site has a typical temperate climate. In the
experimental field, no fertilizers have been applied. To mimic mild drought stress in the
experimental field, we created ridges with a tiller coupled to a rotary after plowing (fig. S1A).
In field trials conducted in the 2015–2018 growing seasons, the ridges were produced with a
TA6 or TA800N tiller (Kubota, Osaka, Japan) coupled to an RA3 rotary (Kubota) and were 30
cm high and 38 cm wide at the base. In the 2019 and 2020 growing seasons, since the same
size rotary was unavailable, ridges were created using a TA800N tiller (Kubota) coupled to a
SKB-E15C rotary (Kubota) and were 30 cm high and 63 cm wide at the base. Because the
thickness of A horizon, a topsoil characterized by organic matter accumulation, is greater than
30 cm in our experimental field, both the flat and ridged plots were composed of this topsoil.
In 2015, the experimental layout consisted of two areas with flat and ridged plots (fig.
S2A). In each area, three plots were used for the analysis. Soybean seeds were sown at 20-cm
intervals on August 3, and a total of 30 plants were grown per plot. For 5 days after sowing,
water was sprayed every day to prompt seed germination. In 2016, the experiment was laid out
in a randomized complete block design, including flat and ridged plots, with four blocks for

four replicates (fig. S2B). Soybean seeds were germinated on vermiculite in a temperature-
controlled phytotron at JIRCAS (27 ± 6 °C), as described previously (*21*), on July 7, and 30
healthy and uniform 6-day-old seedlings were planted per plot at 20-cm intervals on July 13.
In 2017, the experimental layout was divided into two areas, one irrigated and the other rainfed.
Each area was composed of randomized blocks, including flat and ridged plots, with four
replicates (fig. S3A). In the irrigated area, irrigation tubes (EVAFLOW, Mitsubishi Chemical
Agri Dream, Tokyo, Japan) were installed to supply water to the entire area of each plot.
Soybean seeds were sown at 20-cm intervals on July14, and a total of 15 plants were grown
per plot. To avoid drought stress for seedling establishment, all plots were surrounded by
irrigation tubes and irrigated a total of four times from the day before sowing to 7 days after
sowing. Subsequently, the irrigated area was irrigated 20 times during the period from 18 to 95
days after sowing. Too much irrigation may have increased soil moisture throughout the test
area and failed to induce adequate water deficit on ridges (fig. S8A). In 2018, the experimental
layout consisted of two areas, one irrigated and the other rainfed (fig. S2C). Each area included
randomized blocks, with flat and ridged plots, and with four replicates. In the irrigated area,
each plot was surrounded by irrigation tubes and irrigated 9 times during the period from 15 to
53 days after sowing (fig. S7A). Soybean seeds were sown at 20-cm intervals on July 19, and
a total of 25 plants were grown per plot. In 2019, the experiment was laid out in a randomized
complete block design, including flat and ridged plots, with four blocks for four replicates (fig.
S3B). Seeds were germinated on vermiculite in a temperature-controlled phytotron (27 ± 6°C)
on July 12, and 20 healthy and uniform 6-day-old seedlings were planted per plot at 20-cm
intervals. In 2020, the experimental layout consisted of two areas, one irrigated and the other
rainfed (fig. S3C). Each area included randomized blocks, with flat and ridged plots, and with
four replicates. In the irrigated area, each plot was surrounded by irrigation tubes and irrigated
only once, at 28 days after sowing (fig. S8C). Seeds were germinated on vermiculite in a
temperature-controlled phytotron (27 ± 6 °C) on July 30, and 25 healthy and uniform seedlings
were planted per plot at 20-cm intervals on August 6.
203
Pot experiment
Pot experiments were conducted in a greenhouse (20 ± 10°C) at JIRCAS to examine how the
soil volumetric water content (VWC) affects phosphate starvation response (PSR)-related gene
expression in soybean. Soybean seeds were germinated on moistened vermiculite in a
temperature-controlled phytotron (*21*) and then the five-day-old seedlings were transferred to
350 ml pots filled with A horizon collected from the JIRCAS experimental field. The seedlings
were grown under well-watered conditions (up to a VWC of 43%) for five days in the

greenhouse. When primary leaves were fully expanded at 10 days after sowing, we started to adjust the VWC to five different levels, α (42–57%), β (35–50%), γ (30–43%), δ (24–36%), and ε (20–29%), by measuring the weight of the pot, adding water every Monday, Wednesday, and Friday. The fully expanded first trifoliate leaves of 38-day-old seedlings were collected and immediately frozen in liquid nitrogen. Then, the total RNA from leaves was used for gene expression analysis.

Meteorological data

Meteorological data, such as amount of rainfall, were obtained from the Weather Data Acquisition System of Institute for Agro-Environmental Sciences, National Agriculture and Food Research Organization (http://www.naro.affrc.go.jp/org/niaes/aws/). The weather station is located at 36°01′ N, 140°07′ E, approximately 5 km from the JIRCAS experimental field.

Evaluation of growth and yield performance of soybean in the field

To investigate the effect of mild drought stress induced by ridges on soybean growth, we measured the aboveground biomass at three time points per growing season in 2015 and 2016. In 2015, a total of 5 plants (1 or 2 plants per plot), 7 plants (2 or 3 plants per plot), and 7 plants (2 or 3 plants per plot) were selected randomly to measure the aboveground biomass at 21, 30, and 44 days after sowing, respectively. In 2016, a total of 12 plants (3 plants per plot) were selected randomly for measurements of aboveground biomass at 20, 36, and 50 days after sowing. Dry weight of the aboveground biomass was measured after oven-drying at 65°C for more than 48 h.

We evaluated the yield of soybean plants in field trials during 2015–2020. In 2015, a total of 11 plants (3–4 plants per plot) grown on the flat plots and 12 plants (4 plants per plot) grown on the ridged plots were selected randomly and harvested on December 1. The number of seeds per plant and total air-dried seed weight per plant were measured. In 2016, a total of 20 plants (5 plants per plot) for each condition were selected and harvested on November 10. After oven-drying at 65°C for more than 48 h, we measured the following parameters for each plant: number of seeds, total seed weight, stem weight, and pod-shell weight. The harvest index was calculated as total seed weight / (total seed weight + stem weight + pod-shell weight). In 2017, a total of 24 plants (6 plants per plot) were selected randomly for each treatment and harvested on November 15. After oven-drying at 65°C for more than 48 h, we measured the following parameters for each plant: number of seeds, total seed weight, combined stem and pod-shell weight, and plant height. In 2018, a total of 48 plants (12 plants per plot) were selected randomly for each treatment and harvested on November 14. After oven-drying at

65°C for more than 48 h, we measured the following parameters for each plant: number of
seeds, total seed weight, stem and pod-shell weight, plant height, and total branch length. In
2019, a total of 40 plants (10 plants per plot) were selected randomly for each treatment and
harvested on November 13. Since soybean plants were damaged by a typhoon on September 8,
we measured only the total biomass, including seeds, stem, and pod shell as preliminary data.
In 2020, a total of 40 plants (10 plants per plot) were selected randomly for each treatment and
harvested on November 2. After oven-drying at 65°C for more than 48 h, we measured the
following parameters for each plant: total seed weight and combined stem and pod-shell weight.
254

255 **Image acquisition**
256 Thermal images of soybean plants were taken using a thermal imaging infrared camera (CPA-
257 T640A, CHINO, Tokyo, Japan) according to the manufacturer's protocol. Aerial images of the
258 experimental field were obtained using an unmanned aerial vehicle (Phantom 4, DJI, Shenzhen,
259 China) according to the manufacturer's protocol.
260

Measurements of soil water contents
Based on the preliminary investigation of soybean roots in the flat and ridged plots in this
experimental field of JIRCAS, we decided to measure the soil moisture content at 0–20 cm or
0–30 cm, starting from the top of the ridge or the flat, in this study. In 2015, the volumetric
water content (VWC, m^3 m^{-3}) at a depth of 0–20 cm was measured in each experimental plot
at intervals of about 3 days with a 20-cm-long time domain reflectometry (TDR) probe
(Hydrosense II, Campbell Scientific Inc., Logan, UT, USA). Then, *in situ* calibration was
conducted using the gravimetric method according to the manufacturer's protocol. The
tentative values of VWCs were measured using the TDR probe, and the soils at the measured
sites were collected from a depth of 0–20 cm with polyvinyl chloride pipes (4.3 cm in diameter
and 20 cm in length). Soil weights including water were determined. After pre-drying at 60°C
for more than 24 h, soil samples were removed from the pipes and oven-dried at 105°C for
more than 24 h to calculate the gravimetric water content (g g^{-1}). A total of eight points were
measured on any four days. Then, the gravimetric water content was converted into VWC by
multiplying it by the bulk density (BD, Mg m^{-3}), assuming that the water density is 1.0 Mg m^{-}
^{3}. Since Andosols generally have a high porosity (*22*) (about 70% in this study) and can be
easily compressed, it was difficult to collect an accurate volume of a soil from a depth of 0–20
cm using a single pipe in the field. Therefore, to determine the BD of soils from a depth of 0–
20 cm, we carefully collected soil samples from a depth of 10–15 cm using 100-ml soil
sampling cores (DIK-1801-11, Daiki Rika Kogyo, Konosu, Japan) at 12 points, and then

determined the dry wights of the 100-ml soil sample after oven-drying at 105°C for more than

48 h. The average BD of soils from a depth of 0–20 cm was 0.734 ± 0.017 (mean ± SD) Mg

m^{-3}. From the relationships between the tentative values of VWCs recorded by the TDR probe

(estimated value) and those determined *in situ* (true value), we obtained a calibration formula:

actual VWC = 0.9718 × tentative VWC + 12.554.

In 2016–2020, the VWC at a depth of 0–30 cm was automatically recorded in each plot

every day at 10:00 a.m. with a 30-cm-long TDR probe (CS616, Campbell Scientific, Inc.)

connected to a data logger (CR1000, Campbell Scientific, Inc.). Soil temperature at a depth of

12–18 cm was also recorded with a temperature probe (107, Campbell Scientific, Inc.) to

correct for the temperature dependence of the TDR probes, according to the manufacture's

protocol. The estimated VWCs recorded by the TDR probes were similarly calibrated using

the gravimetric method according to the manufacture's protocol. We selected eight TDR probes

for the calibration and inserted the probes into a plot with temperature probes in the

experimental field. The soil samples were collected from four points of the plot on any seven

days, yielding a total of 28 soil samples, with polyvinyl chloride pipes (4.3 cm in diameter and

30 cm in length), and then the soil weight was determined. The dry weights of the soil samples

were determined as described above. We measured the BD at a depth of 0–10 cm and 10–30

cm to determine the BD at a depth of 0–30 cm, because the BD differed at each depth under

our test conditions in the field. To determine the BD of soils at a depth of 0–30 cm, we carefully

collected soil samples from depths of 3–8 cm and 20–25 cm using the 100 ml soil sampling

cores (DIK-1801-11, Daiki Rika Kogyo) at 8 points each. The BD for depths of 0–30 cm was

determined to be 0.773 Mg m^{-3} by the combined weighted average of the values obtained for

depths of 0–10 cm and 10–30 cm. According to the manufacture's protocol, the output period

data of each TDR probe at the sampling times were corrected by temperature, and the tentative

VWCs were calculated. Using the averages of tentative values of VWCs obtained using the

eight probes and the corresponding actual VWCs determined *in situ* at each sampling point, we

generated a calibration formula: actual VWC = 1.0005 × tentative VWC + 0.1514.

Measurements of soil nutrients

Soil samples were taken from a depth of about 0–30 cm with a polyvinyl chloride cylinder or

soil sampler (DIK-1601, Daiki Rika Kogyo) in each plot on July 7 and November 17, 2016,

which corresponded to the start and end of the soybean cultivation period, respectively.

Samples were air-dried and passed through a 2–mm sieve. Then, the obtained fine earth was

subjected to soil chemical analysis. The oven-dried weight of the fine earth was measured to

obtain the moisture correction factors that were used to convert nutrient contents on air-dried

soil to a dry weight basis.

Total carbon and total nitrogen contents were determined using the dry combustion

method with an elemental analyzer (Sumigraph NC-220F, Sumika Chemical Analysis Service,

Tokyo, Japan). Exchangeable bases (K^+) extracted in 1 M ammonium acetate (pH 7) were

measured basically as described previously (23), except that inductively coupled plasma atomic

emission spectroscopy with a spectrometer (ICPE-9000, Shimadzu Corporation, Kyoto, Japan)

was used in this study. Available phosphorus was determined using the Bray-II method (24)

with a spectrophotometer (UV-1800, Shimadzu Corporation).

Soil water retention curve

Soil samples were collected from a depth of 10–15 cm using 100-ml soil sampling cores (DIK-

1801-11, Daiki Rika Kogyo) at six points in each plot. The soil water retention, a relationship

between the VWC (θ) and soil water potential (ψ), of each sample was measured with a

pressure plate apparatus at a water potential of between –3.9 kPa and –98 kPa, as described

previously (25), and with a dew point potentiometer (WP4, Decagon Devices, Pullman, WA,

USA) at values below –310 kPa, as described previously (26). The values of VWCs at a depth

of 0–20 cm and 0–30 cm were calculated using the BD values at each depth. The averaged

VWCs are shown at matric potential values of –3.9, –9.8, –31, –98, –619, and –1554 kPa (fig.

S4). Both regression curves at a depth of 0–20 cm and 0–30 cm were obtained by fitting a

hydraulic model (27) using SWRC Fit (28):

$$\theta = \theta_s \left[\frac{1}{\ln[e + (\psi/a)^n]} \right]^m$$

where θ_s is the saturated VWC; e is the Napier's constant; and a, n, and m are three different

soil parameters. The model showed a good fit (R^2=0.985, $p < 0.001$ for both curves).

Transcriptome analysis

A total of 10 plants (3 or 4 plants per plot) were randomly selected from each treatment group,

and the first to third trifoliate leaves were collected individually and immediately frozen in

liquid nitrogen in the field on September 1, 2015. The frozen second trifoliate leaves were

powdered using a multi-bead shocker (YASUI KIKAI, Osaka, Japan) and the other leaves were

powdered using a mortar and pestle. For RNA-seq analysis, total RNA was extracted from

powdered samples of fully expanded second trifoliate leaves using an RNeasy Plant Mini Kit

(Qiagen, Hilden, Germany) following the manufacturer's protocol. The extracted total RNA

samples from three independent plants per plot were mixed for one replicate, and in total three

biological replicates were performed in each treatment. The extracted RNA was used to construct paired-end libraries using a TruSeq RNA Library Preparation Kit v2 (Illumina, CA, USA). The libraries were sequenced on an Illumina HiSeq 4000 sequencer with a 100-bp paired-end protocol (Macrogen, Kyoto, Japan). For trimming low quality reads and adaptor sequences, a Trimmomatic v0.36 (*29*) with the option of SLIDINGWINDOW:4:20, MINLEN:40, LEADING:20, and TRAILING:20 was used. The trimmed reads were mapped against the reference sequence of Williams 82 (Wm82.a2.v1: https://phytozome.jgi.doe.gov/pz/portal.html) using STAR v2.5.1 (*30*) with the option of outSJfilterReads: Unique; outFilterMatchNminOverLread: 0.96; and outFilterScoreMinOverLread: 0.8. The FPKM (fragment per kilobase of exon per million mapped reads) of each gene was calculated using Cufflinks v2.2.1 (*31*). Differential expression analysis was performed by Cuffdiff embedded in Cufflinks v2.2.1 (*31*). The genes that showed q value < 0.05 were defined as differentially expressed genes (DEGs) between the samples grown on flat and ridged plots. Gene annotation such as homology to *Arabidopsis thaliana* genes and GO was obtained from gene annotation information in Wm82.a2.v1 (Gmax_275_Wm82.a2.v1.annotation_info.txt).

A heatmap of up- and down-regulated DEGs was constructed with the 'heatmap.2' function of the gplots package in R software v3.3.3 (*32*). The different types of representative phosphate starvation responsive genes were selected in the up-regulated genes based on a previous report (*7*). Using the clusterProfiler package in R (*33*), GO enrichment analysis was performed for the up- and down-regulated DEGs selected according to the following criteria: $|\log_2(\text{FC})| \geq 1$, FPKM value > 0, q < 0.05. A directed acyclic graph (DAG) was constructed using the clusterProfiler package in R. FDR < 0.05 was regarded as significantly enriched GO terms. The GO descriptions were obtained using AnnotationHub ("AH13355") in R (*34*). We assessed the significant enrichment of PSR marker genes (*7-9*) in the up- and down-regulated genes using the hypergeometric tests (phyper function in R).

RT-qPCR analyses

For RT-qPCR analysis, total RNA was extracted from soybean leaves using an RNeasy Plant Mini Kit (Qiagen) or RNAiso Plus (Takara Bio, Kusatsu, Japan) according to the manufacturer's protocol. The extracted total RNA was treated with RQ1 RNase-free DNase (Promega, Madison, WI, USA). Complementary DNA (cDNA) was synthesized using PrimeScript RT Master Mix (Takara Bio). RT-qPCR using GoTaq qPCR Master Mix (Promega) was performed by QuantStudio7 Flex (Applied Biosystems, Foster City, CA, USA). The relative amounts of target mRNAs were calculated using the relative standard curve

method and normalized against actin (*Glyma.15G050200*) as a reference gene. The primers used in this study are listed in Supplementary Table S7.

Measurements of inorganic phosphate contents

We examined the inorganic phosphate (Pi) contents in soybean leaves using a molybdate colorimetric assay (*35,36*). The frozen powdered leaf samples used for gene expression analysis were used to analyze the Pi content. A total of 20–45 mg of each sample was mixed with 1% acetic acid (20 μl/mg sample weight), vortexed for 15 min, and incubated at 42°C for 1 h in a heat block. Following centrifugation at 18,300 g for 5 min at room temperature, the supernatant was used for the Pi assay. Reaction solutions containing 140 μl of molybdate solution (a master mix of a 6:1 proportion of 0.42% (w/v) ammonium molybdate in 1 N H_2SO_4 and 10% ascorbic acid in water) and 60 μl of the supernatants were incubated in a 96-well plate at 42°C for 20 min using a thermal cycler. The absorbance of 100 μl of the reaction solutions at 820 nm was measured using a plate reader (ARVOX3, PerkinElmer, MA, USA). The amount of Pi in the solution was calculated using a calibration curve diluted based on Phosphorus Standard Solution (FUJIFILM Wako, Osaka, Japan).

Statistical analysis

Statistical tests in this study, except for the hypergeometric enrichment analysis of PSR (Fig. 3D) conducted by the R function "phyper", were performed in GraphPad Prism 9 (GraphPad software, La Jolla, CA, USA). An unpaired, two-tailed *t*-test or a one-way analysis of variance (ANOVA) with Tukey's test was used in most experiments. The data shown in Fig. 2 and Fig. S7 did not seem to be homoscedastic. Therefore, Brown-Forsythe and Welch ANOVA tests with a Dunnett's T3 multiple comparisons test were used.

References and Notes:

1. J-K. Zhu, Abiotic stress signaling and responses in plants. *Cell* **167**, 313−324 (2016). doi: 10.1016/j.cell.2016.08.029; pmid: 27716505

2. M. Dubois, D. Inzé, Plant growth under suboptimal water conditions: early responses and methods to study them. *J. Exp. Bot.* **71**, 1706−1722 (2020). doi: 10.1093/jxb/eraa037; pmid: 31967643

3. H. Claeys, D. Inzé, The agony of choice: how plants balance growth and survival under water-limiting conditions. *Plant Physiol.* **162**, 1768−1779 (2013). doi: 10.1104/pp.113.220921; pmid: 23766368

4. P. H. Graham, C. P. Vance, Legumes: importance and constraints to greater use. *Plant*

Physiol. **131**, 872−877 (2003). doi: 10.1104/pp.017004; pmid: 12644639

5. L. P. Manavalan, S. K. Guttikonda, L.-S. P. Tran, H. T. Nguyen, Physiological and molecular

approaches to improve drought resistance in soybean. *Plant Cell Physiol.* **50**, 1260−1276

(2009). doi: 10.1093/pcp/pcp082; pmid: 19546148

6. M. G. Gebre, H. J. Earl, Soil water deficit and fertilizer placement effects on root biomass

distribution, soil water extraction, water use, yield, and yield components of soybean

[*Glycine max* (L.) Merr.] grown in 1-m rooting columns. *Front. Plant Sci.* **12**, 581127 (2021).

doi: 10.3389/fpls.2021.581127; pmid: 33790918

7. G. Castrillo, P. J. P. L. Teixeira, S. H. Paredes, T. F. Law, L. de Lorenzo, M. E. Feltcher, O.

M. Finkel, N. W. Breakfield, P. Mieczkowski, C. D. Jones, J. Paz-Ares, J. L. Dangl, Root

microbiota drive direct integration of phosphate stress and immunity. *Nature* **543**, 513−518

(2017). doi: 10.1038/nature21417; pmid: 28297714

8. C. Li, S. Gui, T. Yang, T. Walk, X. Wang, H. Liao, Identification of soybean purple acid

phosphatase genes and their expression responses to phosphorus availability and symbiosis.

Ann. Bot. **109**, 275−285 (2012). doi: 10.1093/aob/mcr246; pmid: 21948626

9. Z. Yao, J. Tian, H. Liao, Comparative characterization of GmSPX members reveals that

GmSPX3 is involved in phosphate homeostasis in soybean. *Ann. Bot.* **114**, 477−488 (2014).

doi: 10.1093/aob/mcu147; pmid: 25074550

10. L. D. B. Suriyagoda, M. H. Ryan, M. Renton, H. Lambers, Plant responses to limited

moisture and phosphorus availability: a meta-analysis. *Adv. Agron.* **124***, 143–200 (2014).

doi: 10.1016/B978-0-12-800138-7.00004-8

11. M. He, F. A. Dijkstra, Drought effect on plant nitrogen and phosphorus: a meta-analysis.

New Phytol. **204**, 924−931 (2014). doi: 10.1111/nph.12952; pmid: 25130263

12. D. T. Krizek, A. Carmi, R. M. Mirecki, F. W. Snyder, J. A. Bunce, Comparative effects of

soil moisture stress and restricted root zone volume on morphogenetic and physiological

responses of soybean [*Glycine max* (L.) Merr.]. *J. Exp. Bot.* **36**, 25−38 (1985). doi:

10.1093/jxb/36.1.25

13. Y. Fujita, M. Fujita, K. Shinozaki, K. Yamaguchi-Shinozaki, ABA-mediated

transcriptional regulation in response to osmotic stress in plants. *J. Plant Res.* **124**, 509−525

(2011). doi: 10.1007/s10265-011-0412-3; pmid: 21416314

14. L. Song, S. C. Huang, A. Wise, R. Castanon, J. R. Nery, H. Chen, M. Watanabe, J. Thomas,

Z. Bar-Joseph, J. R. Ecker, A transcription factor hierarchy defines an environmental stress

response network. *Science*. **354**, aag1550 (2016). doi: 10.1126/science.aag1550. pmid:

27811239

15. J. Pang, M. H. Ryan, H. Lambers, K. H. M. Siddique, Phosphorus acquisition and utilisation

in crop legumes under global change. *Curr. Opin. Plant Biol.* **45**, 248−254 (2018). doi:

10.1016/j.pbi.2018.05.012; pmid: 29853281

16. F. Tardieu, Any trait or trait-related allele can confer drought tolerance: just design the right

drought scenario. *J. Exp. Bot.* **63**, 25−31 (2012). doi: 10.1093/jxb/err269; pmid: 21963615

17. F. Tardieu, T. Simonneau, B. Muller, The physiological basis of drought tolerance in crop

plants: A scenario-dependent probabilistic approach. *Annu. Rev. Plant Biol.* **69**, 733−759

(2018). doi: 10.1146/annurev-arplant-042817-040218; pmid: 29553801

18. J. Schmutz, S. B. Cannon, J. Schlueter, J. Ma, T. Mitros, W. Nelson, D. L. Hyten, Q. Song,

J. J. Thelen, J. Cheng, D. Xu, U. Hellsten, G. D. May, Y. Yu, T. Sakurai, T. Umezawa, M.

K. Bhattacharyya, D. Sandhu, B. Valliyodan, E. Lindquist, M. Peto, D. Grant, S. Shu, D.

Goodstein, K. Barry, M. Futrell-Griggs, B. Abernathy, J. Du, Z. Tian, L. Zhu, N. Gill, T.

Joshi, M. Libault, A. Sethuraman, X.-C. Zhang, K. Shinozaki, H. T. Nguyen, R. A. Wing,

P. Cregan, J. Specht, J. Grimwood, D. Rokhsar, G. Stacey, R. C. Shoemaker, S. A. Jackson,

Genome sequence of the palaeopolyploid soybean. *Nature* **463**, 178–183 (2010). doi:

10.1038/nature08670; pmid: 20075913

19. M. Shimomura, H. Kanamori, S. Komatsu, N. Namiki, Y. Mukai, K. Kurita, K. Kamatsuki,

H. Ikawa, R. Yano, M. Ishimoto, A. Kaga, Y. Katayose, The *Glycine max* cv. Enrei genome

for improvement of Japanese soybean cultivars. *Int. J. Genomic.* **2015**, 358127 (2015).

doi: 10.1155/2015/358127; pmid: 26199933

20. IUSS Working Group WRB, "World reference base for soil resources 2014, update 2015"

(World Soil Resources Rep. Vol. 106, Food and Agriculture Organization of the United

Nations, 2015; http://www.fao.org/3/i3794en/I3794en.pdf)

21. Y. Nagatoshi, Y. Fujita, Accelerating soybean breeding in a CO_2-supplemented growth

chamber. *Plant Cell Physiol.* **60**, 77–84 (2019). doi: 10.1093/pcp/pcy189; pmid: 30219921

22. P. Driessen, J. Deckers, O. Spaargaren, F. Nachtergaele, Eds., "Andosols" in *Lecture Notes*

on the Major Soils of the World (World Soil Resources Rep. Vol. 94, Food and Agriculture

Organization of the United Nations, 2001; http://www.fao.org/3/y1899e/y1899e.pdf).

23. L. P. Van Reeuwijk, "Procedures for Soil Analysis" (Tech. Paper Vol. 9, International Soil

Reference   and   Information   Centre   (ISRIC),   Wageningen,   2002;

https://www.isric.org/sites/default/files/ISRIC_TechPap09.pdf).

24. R. H. Bray, L. T. Kurtz, Determination of total, organic, and available forms of phosphorus

in soils. *Soil Sci.* **59**, 39–46 (1945). doi: 10.1097/00010694-194501000-00006

25. J. H. Dane, J. W. Hopmans, "Pressure plate extractor" in *Methods of Soil Analysis*, J. H.

Dane, G. C. Topp, Eds. (SSSA, Madison, 2002), Part 4: Physical methods, pp. 688–690.

26. B. R. Scanlon, B. J. Andraski, J. Bilskie, "*Dew point potentiameter*" in Methods of soil
analysis, Part 4: Physical methods, J. H. Dane, G. C. Topp, Eds. (SSSA, Madison, 2002)
pp. 662–665.

27. D. G. Fredlund, A. Xing, Equations for the soil-water characteristic curve. *Can. Geotech.*
J. **31**, 521–532 (1994). doi: doi.org/10.1139/t94-061

28. K. Seki, SWRC fit – a nonlinear fitting program with a water retention curve for soils
having unimodal and bimodal pore structure, *Hydrol. Earth Syst. Sci. Discuss.* **4**, 407–437
(2007). doi: 10.5194/hessd-4-407-2007

29. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114–2120 (2014). doi: 10.1093/bioinformatics/btu170; pmid:
24695404

30. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson,
T. R. Gingeras, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013). doi: 10.1093/bioinformatics/bts635; pmid: 23104886

31. C. Trapnell, B. A. Williams, G. Pertea, A. Mortazavi, G. Kwan, M. J. van Baren, S. L.
Salzberg, B. J. Wold, L. Pachter, Transcript assembly and quantification by RNA-Seq
reveals unannotated transcripts and isoform switching during cell differentiation. *Nat.*
Biotechnol. **28**, 511–515 (2010). doi: 10.1038/nbt.1621; pmid: 20436464

32. G. R. Warnes, B. Bolker, L. Bonebakker, R. Gentleman, W. Huber, A. Liaw, T. Lumley, M.
Maechler, A. Magnusson, S. Moeller, M. Schwartz, B. Venables, T. Galili, gplots: Various
R   programming   tools   for   plotting   data,   2016.   https://cran.r-
project.org/web/packages/gplots/index.html

33. G. Yu, L. G. Wang, Y. Han, Q. Y. He, clusterProfiler: an R package for comparing biological
themes among gene clusters. *OMICS* **5**, 284–287 (2012). doi: 10.1089/omi.2011.0118;
pmid: 22455463

34. M. Morgan, M. Carlson, D. Tenenbaum, S. Arora, AnnotationHub: Client to access
AnnotationHub   resources.   R   package   version   2.10.1.,   2017.   doi:
10.18129/B9.bioc.AnnotationHub

35. B. N. Ames, Assay of inorganic phosphate, total phosphate and phosphatases. *Methods*
Enzymol. **8**, 115–118 (1966). doi: 10.1016/0076-6879(66)08014-5

36. T. J. Chiou, K. Aung, S.-I. Lin, C. C. Wu, S.-F. Chiang, C.-L. Su, Regulation of phosphate
homeostasis by MicroRNA in *Arabidopsis*. *Plant Cell* **18**, 412–421 (2006). doi:
10.1105/tpc.105.038943; pmid: 16387831

Acknowledgments: We thank S. Shu, M. Toyoshima, K. Ozawa, K. Shimizu, N. Hisatomi, Y.
Saito, J. Baba, M. Ohsaki, Y. Shirai, Y. Nakamura, I. Gejima, H. Ohwada, N. Tayama, K.
Kawamura, T. Komatsu, and H. Ishiyama for technical assistance; and T. Taji, K. Nakashima,
M. Fujita, and T. Ogata for discussions.

Funding: This work was supported by the Grants-in-Aid for Scientific Research (KAKENHI)
by the Japan Society for the Promotion of Science (JSPS) (Grant Nos. JP18K05379 to YN;
JP21H02158 to YN, YF; JP16K07412, JP24510312 to YF), Cross-ministerial Moonshot
Agriculture, Forestry and Fisheries Research and Development Program by the Bio-oriented
Technology Research Advancement Institution (BRAIN) (Grant No. JPJ009237), the Science
and Technology Research Partnership for Sustainable Development (SATREPS) by the Japan
Science and Technology Agency (JST) and the Japan International Cooperation Agency
(JICA) (Grant No. JPMJSA1907), and the Ministry of Agriculture, Forestry and Fisheries
(MAFF) of Japan.

Author contributions: Y.N. and Y.F. conceived and designed the study and wrote the paper;
Y.N. performed most of the experiments and analyzed the data; K.I. performed the soil and
chemical analyses; Y.N., K.F., and Y.F. performed the field experiments, with support from
T.O. and Y.K.; Y.N., N.M., Y.K., E.O-T., M.I., and Y.Y. performed the transcriptome analysis.
All authors discussed and commented on the manuscript.

Competing interests: The authors declare no competing interests.

Data and materials availability: RNA-seq data are available from the DNA Data Bank of
Japan (www.ddbj.nig.ac.jp/) under accession number DRA012279. No restrictions are placed
on materials, such as materials transfer agreements. Details of all data, code, and materials used
in the analysis are available in the main text or the supplementary materials.

Supplementary Materials

Figs. S1 to S10

Tables S1 to S7

Fig. 1. Ridges can artificially induce mild drought stress, reducing soybean yield. (**A**) Photograph of the experimental field, including the VWC data collection station, watering tubes, and probes (yellow arrows). (**B**) Twelve-day-old soybean plants on flat and ridged plots (referred to as "flats" and "ridges", respectively). (**C**) Daily rainfall (dark green bars) and time-course of soil VWC ($n = 4$) in the flats (blue lines and points) and ridges (pink lines and points) over the study period in the 2016 field. (**D**) Aboveground biomass (dry weight, DW) per plant grown on the flats and ridges ($n = 12$). F and R denote flats and ridges, respectively. (**E**) Thermogram and the corresponding digital image of 9-week-old soybean plants grown on the flats and ridges in 2016. (**F**) Total seed weight per plant grown on the flats and ridges ($n = 20$). (**G**) Seeds from individual representative plants grown on the flats and ridges. Scale bars, 10 cm. (**H**) Nutrient contents (N, P, and K) of soil in the flats and ridges before (July) and after (November) soybean cultivation in 2016 ($n = 4$). $**P < 0.01$, $***P < 0.001$; Student's t-test; ns, no significant difference. Error bars in (C), (D), (F), and (H) denote SD.

Fig. 2. Irrigation compensates for the reduced growth of plants on ridges. (**A**) Aerial view of 7-week-old soybean plants in the 2018 test. F and R denote flats and ridges, respectively. Left half, irrigated area. Right half, rainfed area. (**B**) Representative view of soybean plants from the flats and ridges with and without irrigation. Scale bar, 10 cm. (**C**) Biological yield (total dry weight of seed, stem, and pod shells) per plant grown on the flats and ridges ($n = 48$). ***$P < 0.001$, Brown-Forsythe and Welch ANOVA tests with Dunnett's T3 multiple comparisons test; ns, no significant difference. Error bars denote SD.

Fig. 3. Mild drought induces PSR in field-grown soybean plants. (**A**) Positions of the second trifoliate leaves (dotted enclosure) used for RNA-seq analysis, in 30-day-old soybean plants grown on flats and ridges in 2015. Scale bars, 10 cm. (**B**) Schematic representation of a soybean plant. (**C**) Heat map of $\log_2$ fold change in expression of 990 up-regulated and 400 down-regulated genes ($|\log_2(FC)| \geq 1$, FPKM value > 0, $q < 0.05$) under mild drought conditions induced by ridges. Lower row indicates PSR genes. (**D**) Ratio of PSR marker genes to the total number of up- and down- regulated genes, respectively. (**E**) A list of representative up-regulated PSR marker genes under mild drought conditions. (**F**) PSR gene expression with quantitative RT-PCR in first to third trifoliate leaves of 29-day-old plants grown on ridges and flats in 2015 ($n = 9$ in second trifoliate leaves; $n = 10$ in first and third trifoliate leaves). F and R denote flats and ridges, respectively. (**G**) Pi contents of first to third trifoliate leaves in soybean plants grown on ridges and flats in the same samples as described in (F). $^{*}P < 0.05$, $^{**}P < 0.01$, and $^{***}P < 0.001$ [one-way analysis of variance (ANOVA) with Tukey's test], in ridged versus flat plots for each leaf position; ns, no significant difference. Error bars denote SD.

Fig. 4. Mild and severe water deficit induce the PSR and ABA response, respectively.
(**A**) Time-course analysis of VWCs in potted soybean plants with five different moisture contents ($n = 6$). The range of variation in VWCs during the period from day 7 after sowing (when plants were transplanted into pots) to day 38 (when sampling was done) is shown on the right. From 10 days after sowing, we started the drought stress treatments. (**B**) Thermogram and the corresponding digital image of 26-day-old soybean plants in the pots. (**C and D**) Relative expression of PSR (C) and ABA responsive (D) genes in the V1 leaves of the 38-day-old plants shown in (B), as determined by quantitative RT-PCR ($n = 6$). $^*P < 0.05$, $^{**}P < 0.01$, and $^{***}P < 0.001$ [one-way analysis of variance (ANOVA) with Tukey's test], in β–ε versus α. Error bars denote SD. (**E**) Proposed model of the plant's response in terms of gene expression to increasing levels of water deficit stress.