The transcriptional corepressor CTBP-1 acts with the SOX family

transcription factor EGL-13 to maintain AIA interneuron cell identity in *C.*

elegans

Josh Saul[1], Takashi Hirose[1,#a], H. Robert Horvitz[1,*]

Affiliations

[1] Howard Hughes Medical Institute, Department of Biology, Massachusetts

Institute of Technology, Cambridge, MA, USA

[#a] Current Address: Sysmex Corporation, 4-4-4 Takatsukadai, Nishi-ku,

Kobe 651-2271, Japan

*Corresponding author

email: horvitz@mit.edu (HRH)

Draft: 10/6/2021

Abstract

Cell identity is characterized by a distinct combination of gene expression, cell morphology and cellular function established as progenitor cells divide and differentiate. Following establishment, cell identities can be unstable and require active and continuous maintenance throughout the remaining life of a cell. Mechanisms underlying the maintenance of cell identities are incompletely understood. Here we show that the gene *ctbp-1,* which encodes the transcriptional corepressor C-terminal binding protein-1 (CTBP-1), is essential for the maintenance of the identities of the two AIA interneurons in the nematode *Caenorhabditis elegans*. *ctbp-1* is not required for the establishment of the AIA cell fate but rather functions cell-autonomously and can act in later larval stage and adult worms to maintain proper AIA gene expression, morphology and function. From a screen for suppressors of the *ctbp-1* mutant phenotype, we identified the gene *egl-13,* which encodes a SOX family transcription factor. We found that *egl-13* regulates AIA function and aspects of AIA gene expression, but not AIA morphology. We conclude that the CTBP-1 protein maintains AIA cell identity in part by utilizing EGL-13 to repress transcriptional activity in the AIAs. More generally, we propose that transcriptional corepressors like CTBP-1 might be critical factors in the maintenance of cell identities, harnessing the DNA-binding specificity of transcription factors like EGL-13 to selectively regulate gene expression in a cell-specific manner.

Introduction

Over the course of animal development, complex networks of transcription factors act and interact to drive the division and differentiation of progenitor cells towards terminal cell identities [1–8]. These networks of transcriptional activity often culminate in the activation of master transcriptional regulators that are responsible for directing the differentiation of a diverse range of cell and tissue types [4,9–12]. Examples of such master transcriptional regulators include the mammalian bHLH transcription factor MyoD, which specifies skeletal muscle cells [13–15]; the *Drosophila* Pax-family transcription factor Eyeless, which drives differentiation of the fly eye [16–20]; and the *C. elegans* GATA transcription factor ELT-2, essential for development of the worm intestine [21–24]. Many such master transcriptional regulators are not only required to establish the identities of specific cell types but are subsequently continuously required to maintain those identities for the remaining life of the cell [4,23,25–29]. Defects in the maintenance of cell identities can manifest as late-onset misregulated gene expression, altered morphology or disrupted cellular function, and often become progressively worse as the cell ages [25,28,30–32].

Previous studies of the nematode *Caenorhabditis elegans* have identified a class of master transcriptional regulators, termed terminal selectors [4,12,33–37]. Terminal selectors drive the expression of whole batteries of gene activity that ultimately define the unique features of many different cell types [10–12]. Individual terminal selectors have been shown to contribute to the establishment and maintenance of multiple distinct *C. elegans* cell types and to drive the

expression of many cell-type specific genes [7,37–41]. However, it has been

unclear how individual terminal selectors can drive the expression of cell-type

specific genes in only the appropriate cell types rather than in all cells in which

they act [42–44]. Recent work has shown that terminal selectors appear to

broadly activate the expression of many genes, including cell-type specific

genes, in all cells in which they function [42,45]. Piecemeal assemblies of

transcription factors are then responsible for pruning this broad expression to

restrict expression of cell-type specific genes to the appropriate cell types

[42,45]. This restriction of the activation of gene expression by terminal selectors

appears to be an essential aspect of proper cell-identity maintenance

[28,42,45,46]. However, it is not known how the myriad of transcription factors

utilized to restrict terminal selector gene activation are coordinated and

controlled.

Here we report the discovery that the *C. elegans* gene *ctbp-1,* which

encodes the sole worm ortholog of the C-terminal Binding Protein (CtBP) family

of transcriptional corepressors [47–55], functions to maintain the cell identity of

the two AIA interneurons. We demonstrate that CTBP-1 functions with the SOX-

family transcription factor EGL-13 [56,57] to maintain multiple aspects of the AIA

cell identity and propose that CTBP-1 does so in part by utilizing EGL-13 to

repress transcriptional activity in the AIAs.

Results

Mutations in *ctbp-1* cause *ceh-28* reporter misexpression in the AIA neurons

In previous studies, we screened for and characterized mutations that prevent the programmed cell death of the sister cell of the *C. elegans* M4 neuron [58,59]. For these screens, we used the normally M4-specific GFP transcriptional reporter P_{ceh-28}::*gfp* and identified isolates with an undead M4 sister cell, which expresses characteristics normally expressed by the M4 cell, on the basis of ectopic GFP expression. In addition to mutants with an undead M4 sister cell, we isolated 18 mutant strains that express P_{ceh-28}::*gfp* in a manner uncharacteristic of M4 or its undead sister. These mutants express P_{ceh-28}::*gfp* in a bilaterally symmetric pair of cells located near the posterior of the *C. elegans* head, far from both M4 and the single M4 sister cell (Fig. 1A).

These mutations define a single complementation group and all 18 mutant strains have mutations in the transcriptional corepressor gene *ctbp-1* (Figs. 1C; S1A-B; S2A). These *ctbp-1* alleles include three splice-site mutations and nine nonsense mutations (such as the mutation *n4784,* an early nonsense mutation and one of many presumptive null alleles of the gene). The mutant phenotype is recessive, and a transgenic construct carrying a wild-type copy of *ctbp-1* expressed under its native promoter fully rescued the GFP misexpression caused by *n4784* (Figs. 1A; S2A). *tm5512*, a 632 bp deletion spanning the transcription start site and first two exons of the *ctbp-1a* isoform and a presumptive null allele of this gene [60], likewise caused P_{ceh-28}::*gfp*

misexpression in two cells in the posterior region of the head (Fig. S1C-D),

similar to our *ctbp-1* isolates. These findings demonstrate that loss of *ctbp-1*

function is responsible for P_{ceh-28}::*gfp* misexpression.

To determine the identity of the cells misexpressing the normally M4-

specific marker P_{ceh-28}::*gfp*, we examined reporters for cells in the vicinity of the

observed misexpression in *ctbp-1* mutants. The AIA-neuron reporter *nIs843[P_{gcy-}*

$_{28.d}$::mCherry] showed complete overlap with misexpressed P_{ceh-28}::*gfp*, indicating

that the cells misexpressing the M4 reporter are the two bilaterally symmetric and

embryonically-generated AIA interneurons (Fig. 1B).

The penetrance of *ceh-28* reporter misexpression in the AIA neurons

increases with age

While characterizing *ctbp-1* mutants, we noticed that fewer young worms

misexpress P_{ceh-28}::*gfp* in the AIAs than do older worms (Fig. 1D). To investigate

the temporal aspect of this phenotype, we scored *ctbp-1* mutants for P_{ceh-28}::*gfp*

misexpression throughout the four worm larval stages (L1-L4) and into the first

day of adulthood ("early" and "day 1" adults). *ctbp-1* mutants rarely misexpressed

P_{ceh-28}::*gfp* at early larval stages, but displayed an increasing penetrance, though

invariant expressivity, of this defect as worms transitioned through larval

development, such that by the last larval stage (L4) nearly all worms exhibited

reporter misexpression specifically and solely in the AIAs (Fig. 1E). A similar

stage-dependent increase in reporter expression in *ctbp-1* mutants occurred in

mutants carrying a second independently-generated *ceh-28* reporter, *nIs348[P_{ceh-}*

$_{28}$::mCherry] (Fig. S1E). These results demonstrate that ctbp-1 function prevents

an age-dependent misexpression of the M4-specific gene ceh-28 in the unrelated

AIA neurons.

We next asked in what cells and at what stages ctbp-1 functions to

suppress P_{ceh-28}::gfp expression in the AIAs. We generated a transgenic

construct that expresses wild-type ctbp-1 specifically in the AIAs, nIs743[P_{gcy-}

$_{28.d}$::ctbp-1(+)] (hereafter referred to as nIs743[P_{AIA}::ctbp-1(+)]). We found that

AIA-specific restoration of ctbp-1 was sufficient to suppress P_{ceh-28}::gfp

misexpression in an otherwise ctbp-1 mutant background (Figs. 1F; S2A),

demonstrating that ctbp-1 is able to act cell-autonomously to regulate ceh-28

expression in the AIA neurons.

To determine if ctbp-1 can act in older animals to suppress AIA gene

misexpression, we generated a transgenic construct that drives expression of

wild-type ctbp-1 throughout the worm in response to a short heat shock,

nEx2351[$P_{hsp-16.2}$::ctbp-1(+);$P_{hsp-16.41}$::ctbp-1(+)] (hereafter referred to as

nEx2351[P_{hsp}::ctbp-1(+)]). We found that heat shock during the L4 larval stage

was sufficient to suppress P_{ceh-28}::gfp misexpression in adult ctbp-1 mutant AIAs

(Figs. 1G-H; S2B), demonstrating that ctbp-1 can act in L4-to-young adult stage

worms to regulate AIA gene expression.

From these data we conclude that ctbp-1 is able to act cell-autonomously

and in L4-to-young adult worms to prevent expression of at least one non-AIA

gene in the AIA neurons.

151   ***ctbp-1* mutant AIAs are not transdifferentiating into an M4-like cell identity**

152   We asked if P_{ceh-28}::*gfp* misexpression in the AIAs of *ctbp-1* mutants might

153   be a consequence of the AIAs transdifferentiating into an M4-like cell identity. We

154   scored *ctbp-1* mutants for cell-type markers expressed in, although not

155   necessarily unique to, either M4 or the AIA neurons (Figs. 2A-B; S3A-B). We

156   found that *ctbp-1* mutant AIAs expressed all five of five AIA markers tested and

157   did not express any of four other (non-*ceh-28)* M4 markers tested. Of particular

158   note, *ctbp-1* mutant AIAs did not misexpress either of the two tested M4 genes

159   known to be directly regulated by *ceh-28* (i.e. *dbl-1* and *egl-17)*, indicating that

160   the *ceh-28* misexpression in mutant AIAs does not activate the *ceh-28* regulatory

161   pathway [61,62]. We conclude that *ctbp-1* mutant AIAs are not transdifferentiated

162   into M4-like cells and instead seem to retain much of their AIA identity while

163   gaining at least one M4 characteristic (i.e. *ceh-28* expression) later in life.

164

165   ***ctbp-1* mutants display an increasingly severe disruption of AIA**

166   **morphology**

167   Because of the time-dependency of the defect of *ctbp-1* mutants in AIA

168   cell identity, we hypothesized that *ctbp-1* might act to maintain the AIA cell

169   identity. To test this hypothesis, we examined morphological and functional

170   aspects of AIA identity at both early and late larval stages. To assay AIA

171   morphology, we generated a transgenic construct driving expression of GFP

172   throughout the AIA cell (*nIs840[P_{gcy-28.d}::gfp]*). We crossed this construct into

173   *ctbp-1* mutant worms and visualized AIA morphology in L1 and L4 larvae as well

as in day 1 adults (Fig. 3A). We found that L1 *ctbp-1* mutant AIAs appeared

grossly wild-type in morphology (Fig. 3A). However, L4 and adult *ctbp-1* mutant

AIAs had ectopic neurite branches that extended from both the anterior and

posterior ends of the AIA cell body (Fig. 3A). The penetrance of these ectopic

branches increased progressively in later larval stage and adult mutants (Fig. 3B-

C). Older *ctbp-1* mutant AIAs also appeared to have an elongated cell body

compared to wild-type AIAs. Quantification of this defect revealed that L4 and

adult mutant AIA cell bodies, but not those of L1s, were significantly longer than

their wild-type counterparts (Fig. 3D). To assess if this increase in AIA length was

a consequence of an increase in AIA size, we measured the maximum area of

the AIA cell body from cross-sections of these cells. We found that the maximum

area of the AIA cell body did not significantly differ between wild-type and mutant

AIAs at any stage (Fig. S4A), indicating that mutant AIAs were misshapen but not

enlarged. To confirm that we were not biased by an awareness of genotype while

measuring AIA lengths, we blinded the wild-type and *ctbp-1* AIA images used for

length measurements and scored the blinded images as either "normal" or

"elongated" (Fig. S4B). Again, at the L1 larval stage both wild-type and *ctbp-1*

mutant AIAs appeared overwhelmingly "normal," whereas at both the L4 larval

stage and in day 1 adults *ctbp-1* mutant AIAs were scored as "elongated" at a

consistently higher rate than their wild-type counterparts. Collectively, these

results demonstrate that *ctbp-1* mutant AIAs display abnormal morphology and

that the severity of the observed morphological defects in *ctbp-1* mutants

increases from L1 to L4 to adulthood. Furthermore, the relative lack of AIA

morphological defects in L1 *ctbp-1* mutants suggests that *ctbp-1* is not required

for the establishment of proper AIA morphology but instead acts to maintain AIA

morphology over time.

We next asked if *ctbp-1* acts cell-autonomously and at later stages to

regulate AIA morphology as it does for AIA gene expression. We visualized *ctbp-*

1 mutant AIAs carrying the AIA-specific *ctbp-1(+)* rescue construct

nIs743[P$_{AIA}$::ctbp-1(+)] (Fig. 3E). We found that AIA-specific restoration of *ctbp-1*

in mutant worms rescued all AIA morphological defects to near-wild-type levels at

all stages tested, indicating that *ctbp-1* can act cell-autonomously to regulate AIA

morphology (Fig. 3F-H). Next, we visualized *ctbp-1* mutant worms carrying the

heat shock-inducible *ctbp-1(+)* rescue construct *nEx2351[P$_{hsp}$::ctbp-1(+)]*. We

found that heat shock at the L4 stage did not restore *ctbp-1* mutant AIA

morphology in day 1 adults back to wild-type, nor was there any significant

difference in the frequency of morphological defects between heat-shocked and

non-heat-shocked adults carrying the rescue construct (Fig. 3I-L). We speculate

that the lack of restoration of morphology in late-stage worms might be a

consequence of the defects being irreversible, and that *ctbp-1* might be

continuously required to prevent such defects from occurring. From these data

we conclude that *ctbp-1* can act cell-autonomously, and possibly continuously, to

maintain aspects of AIA morphology in a manner similar to AIA gene expression.

***ctbp-1* mutants display a progressive decline of AIA function**

The AIA interneurons integrate sensory information from a number of

sensory neurons, resulting in modulation of the movement of the worm in

response to environmental stimuli [63–65]. The AIAs function in response to

volatile odors and play an important role in learning associated with the sensation

of volatile odors or salts [63,66]. We asked if *ctbp-1* mutants are abnormal in a

behavior known to require the AIAs – adaptation to the volatile odor 2-butanone

[66] – reasoning that if AIA function is disrupted in *ctbp-1* mutants, there should

be a reduction of adaptation (and thus greater attraction) to butanone in *ctbp-1*

worms relative to wild-type worms.

Consistent with previous studies [66], we found that worms that had been

briefly starved with 90 minutes of food deprivation and had no prior experience

with butanone (so-called "naïve" worms) were generally attracted to the odor,

while worms that were briefly starved in the presence of butanone ("conditioned"

worms) adapted to the odor and exhibited mild repulsion to it (Fig. 4A-E). We

next compared wild-type and *ctbp-1* mutant worms for their ability to adapt to

butanone. We found that while L1 *ctbp-1* worms showed an ability to adapt to

butanone roughly similar to that of their wild-type counterparts, conditioned L4

ctbp-1 mutants displayed a significant decrease in repulsion from butanone

relative to wild-type L4 animals, indicating a decrease in their ability to adapt to

the odor (Fig. 4B-E). As a control, we assayed a strain carrying a transgenic

construct that genetically ablates the AIA neurons, JN580. As expected, JN580

worms displayed decreased butanone adaptation at both the L1 and L4 larval

stages. Thus, *ctbp-1* mutant worms displayed a defect in butanone adaptation

similar to that of an AIA-ablated strain and did so only at a later larval stage,

suggesting a potential loss of AIA function in L4 *ctbp-1* mutants. However, while

ctbp-1 mutant L4s exhibited weaker butanone adaptation than their wild-type

counterparts, this defect was not as severe as that of JN580 L4s, indicating that

ctbp-1 mutant AIAs might retain some function. Additionally, the lack of a

butanone adaptation defect in L1 *ctbp-1* mutants similar to that of L1 JN580

worms further suggests that loss of *ctbp-1* does not disrupt early AIA function

and shows that *ctbp-1* is not required for the establishment of functional AIA

neurons.

We next asked if *ctbp-1* can act cell-autonomously in the AIAs and in older

worms to regulate butanone adaptation. We assayed *ctbp-1* mutants carrying the

AIA-specific rescue construct *nIs743[P_{AIA}::ctbp-1(+)]* for butanone adaptation

(Fig. 4B-E) and found that AIA-specific restoration of *ctbp-1* rescued butanone

adaption of conditioned *ctbp-1* mutant L4s to near wild-type levels (Fig. 4E). We

conclude that the butanone adaptation defect of *ctbp-1* mutants is caused by a

disruption of AIA function and that *ctbp-1* can act cell-autonomously to regulate

this AIA function. Next, we assayed *ctbp-1* mutants carrying the heat shock-

inducible *ctbp-1(+)* rescue construct *nEx2351[P_{hsp}::ctbp-1(+)]* for butanone

adaptation. We found that restoration of *ctbp-1* by heat shock at the L4 larval

stage rescued the butanone adaptation defect in day 1 adults, indicating that

ctbp-1 can act in L4-to-day 1 adult worms to maintain proper AIA function after

the initial establishment of the AIA cell identity (Fig. 4F-G). Taken together, these

data establish that loss of *ctbp-1* disrupts the function of the AIA neurons and

that *ctbp-1* can act cell-autonomously and in L4-to-day 1 adult worms to maintain

AIA function.

While conducting these assays, we observed that naïve *ctbp-1* mutant

worms displayed a mildly weaker attraction to butanone than did their wild-type

counterparts at both the L1 and L4 larval stages (Fig. 4B,D). AIA-specific rescue

of *ctbp-1* did not rescue this mild chemotaxis defect – naïve *ctbp-1* mutants

carrying the $P_{AIA}::ctbp-1(+)$ construct still displayed weaker butanone attraction

than wild-type worms (Fig. 4B,D). We suggest that this defect in attraction to

butanone is not a consequence of dysfunction of the AIAs but rather of some

other cell(s) involved in butanone chemotaxis. Consistent with this hypothesis,

we found that *ctbp-1* mutants were defective in chemotaxis to the volatile odor

isoamyl alcohol but were not defective in the response to diacetyl (Fig. S5A-B)

while AIA-ablated strain JN580 animals were not defective in either response

(Fig. S5A-B). These observations indicate that *ctbp-1* mutant worms have a

broader defect in chemotaxis caused by the disruption of the function of cells

other than the AIAs. Because our primary focus has been on how *ctbp-1*

functions to maintaining the AIA cell identity, we did not attempt to identify the

other cells with functions perturbed by the loss of *ctbp-1*.

***ctbp-1* mutant AIAs have additional defects in gene expression**

To better characterize the genetic changes occurring in mutant AIAs, we

performed a single, exploratory single-cell RNA-Sequencing (scRNA-Seq)

experiment comparing wild-type and *ctbp-1* mutant worms. We sequenced RNA

from the neurons of wild-type and *ctbp-1* L4 worms and processed the resulting

data using the 10X CellRanger pipeline to identify presumptive AIA neurons

based on the expression of several AIA markers (*gcy-28*, *ins-1*, *cho-1*) shown

above to be expressed in both wild-type and *ctbp-1* mutant AIAs (Fig. 2B).

Confirming that these data captured changes in the AIA transcriptional profiles,

we found that *ctbp-1* mutant AIAs showed high levels of expression of *ceh-28,*

while wild-type AIAs showed no detectable *ceh-28* expression (Fig. S6).

We analyzed AIA transcriptional profiles to identify genes that appeared to

be either expressed in *ctbp-1* mutant AIAs and not expressed in wild-type AIAs

(similar to *ceh-28*) or expressed in wild-type AIAs but not expressed in *ctbp-1*

AIAs. To confirm candidate genes, we crossed existing reporters for those genes

to *ctbp-1* mutants or, in cases for which reporters were not readily available,

generated our own transgenic constructs. We identified and confirmed one gene

that, similar to *ceh-28*, was not expressed in wild-type AIAs but was

misexpressed in *ctbp-1* mutant AIAs: *acbp-6*, which is predicted to encode an

acyl-Coenzyme A binding protein [67] (Fig. 5A; S6). We also identified and

confirmed two genes expressed in wild-type AIAs but not expressed in *ctbp-1*

mutant AIAs: *sra-11*, which encodes a transmembrane serpentine receptor [68];

and *glr-2,* which encodes a glutamate receptor [69] (Fig. 5C,E; S6). We

visualized the *acbp-6* reporter *nEx3081[P$_{acbp-6}$::gfp]*, the *sra-11* reporter

otIs123[P$_{sra-11}$::gfp] and the *glr-2* reporter *ivEx138[P$_{glr-2}$::gfp]* in wild-type and

ctbp-1 L4 worms and confirmed that *acbp-6* was absent in wild-type AIAs but

misexpressed in *ctbp-1* mutants (Fig. 5A-B), while both *sra-11* and *glr-2* were

consistently expressed in wild-type AIAs but not expressed in the AIAs of *ctbp-1*

mutants (Fig. 5C-F). We also visualized these reporters in L1 wild-type and *ctbp-*

1 worms and found that both $P_{acbp-6}::gfp$ and $P_{sra-11}::gfp$ displayed a time-

dependence to their expression similar to that of $P_{ceh-28}::gfp$ − $P_{acbp-6}::gfp$ was

rarely detectible in the AIAs of either wild-type of *ctbp-1* AIAs at the L1 stage but

was consistently expressed in *ctbp-1* mutant L4 AIAs (Fig. 5A-B), while P_{sra-}

$_{11}::gfp$ was rarely detectible in the AIAs of either wild-type or *ctbp-1* mutant L1

worms but was expressed in the AIAs of most wild-type worms by the L4 stage

while remaining off in the AIAs of most L4 *ctbp-1* mutants (Fig. 5C-D). These

observations suggest that, like *ceh-28* expression, *acbp-6* and *sra-11* expression

is regulated by *ctbp-1* primarily in the AIAs of late-stage larvae and adults. By

contrast, *glr-2* was expressed in wild-type but not *ctbp-1* AIAs in both L1 and L4

larvae (Fig. 5E-F).

These data demonstrate that mutant AIAs fail to turn on and/or maintain

the expression of genes characteristic of the adult AIA neuron (*sra-11* and *glr-2*)

while misexpressing at least two genes uncharacteristic of AIA (*ceh-28* and *acbp-*

6). That the majority of these abnormalities in AIA gene expression occurred long

after the AIAs are generated during embryogenesis further supports the

conclusion that *ctbp-1* does not act to establish the AIA cell identity.

Collectively, our findings concerning AIA gene expression, morphology

and function demonstrate that *ctbp-1* acts to maintain the AIA cell identity, plays

little to no role in the initial establishment of the AIA cell fate, and can act cell-

autonomously and in older worms to maintain these aspects of the AIA identity.

egl-13 mutations suppress the _ctbp-1_ mutant phenotype

To investigate how _ctbp-1_ acts to maintain AIA cell identity, we performed

a mutagenesis screen for suppression of P_{ceh-28}::_gfp_ misexpression in the AIAs of

L4 _ctbp-1_ mutants (Fig. 6A). Using a combination of Hawaiian SNP mapping [70]

and whole-genome sequencing, we identified the gene _egl-13_, which encodes a

SOX family transcription factor, as a suppressor of _ctbp-1_. _egl-13_ has been

shown to act in the establishment of the BAG and URX cell fates and in vulval

development of _C. elegans_ [56,71], and its mammalian orthologs SOX5 and

SOX6 act in neural fate determination [72,73]. We isolated three alleles of _egl-13_

as _ctbp-1_ suppressors: _n5937_, a mutation of the splice acceptor site at the

beginning of the 6th exon of the _egl-13a_ isoform resulting in a frameshift and early

stop; _n6013_, a Q381ochre nonsense mutation towards the end of the _egl-13_

transcript; and _n6313_, a 436-nucleotide deletion spanning the 7th and 8th exons of

the _egl-13a_ isoform (Figs. 6B; S7A-B). We generated and introduced a

transgenic construct carrying a wild-type copy of _egl-13_ under its native promoter

into these mutant strains and found that this construct was capable of rescuing

the suppression of P_{ceh-28}::_gfp_ misexpression by all three _egl-13_ alleles,

demonstrating that loss of _egl-13_ function suppresses this aspect of the _ctbp-1_

mutant phenotype and suggesting that these alleles are likely loss-of-function

alleles of _egl-13_ (Fig. S7C).

We assayed the loss-of-function allele of _egl-13_ with the highest

penetrance of suppression, _n5937_, for its ability to suppress P_{ceh-28}::_gfp_

357   misexpression over the course of larval development and into adulthood of *ctbp-*

358   *1* mutant worms (Fig. 6C). *egl-13(n5937)* strongly suppressed *ctbp-1* at all

359   stages, resulting in little to no misexpression of P_{ceh-28}*::gfp* in the AIAs of *egl-13*

360   *ctbp-1* double mutants at any larval stage or in day 1 adults.

361   To determine if, like *ctbp-1*, *egl-13* can act cell-autonomously in the AIAs,

we generated a transgenic construct that drives expression of a wild-type copy of

egl-13 in the AIAs (*nEx3055[$P_{gcy-28.d}$::egl-13(+)]*). Introduction of this construct to

egl-13(n5937) ctbp-1 double mutants rescued the *egl-13* suppression of P_{ceh-}

$_{28}$*::gfp* misexpression in the AIAs, indicating that *egl-13* can function cell-

autonomously (Fig. S8A-B). These results suggest that, in the absence of *ctbp-1*

function, ectopic *egl-13* activity drives *ceh-28* misexpression, and thus that *ctbp-1*

likely normally acts to repress *egl-13* activity in the AIAs.

369   We next asked if *egl-13(n5937)* could suppress the AIA morphological and

functional defects of *ctbp-1* mutants. To both test suppression of AIA

morphological defects and confirm the presence of the AIA neurons in *egl-13*

ctbp-1 double mutants, we crossed the AIA morphology reporter *nIs840[P_{gcy-}*

$_{28.d}$*::gfp]* into *egl-13 ctbp-1* double mutants and scored AIA morphology in L1, L4

and day 1 adult worms (Fig. 6D-G). *egl-13 ctbp-1* double mutant AIAs displayed

a mild (though significant) reduction in the penetrance of ectopic anterior neurites

only in adult worms, no significant change in the frequency of posterior neurites

at any stage, and a slight increase in AIA cell body length of L4 worms (though

the difference was no longer significant in adults). These data demonstrate that

loss of *egl-13* has little consistent effect on the AIA morphological defects caused

by a loss of *ctbp-1* activity, suggesting that *ctbp-1* maintains AIA morphology

primarily through *egl-13*-independent pathways.

We next assayed the ability of *egl-13(n5937)* to suppress AIA functional

defects. We tested *egl-13 ctbp-1* double mutants for butanone adaptation and

found that, at the L1 larval stage, this double mutant strain displayed a detectable

response to butanone similar to *ctbp-1* single mutants (Fig. 6H-I). By contrast, at

the L4 larval stage, mutation of *egl-13* strongly suppressed the *ctbp-1* mutant

defect in butanone adaptation, causing near wild-type levels of repulsion in

conditioned worms (Fig. 6J-K). These results indicate that loss of *egl-13* activity

suppressed AIA functional defects of *ctbp-1* mutant worms and suggest that

ctbp-1 maintains at least this aspect of AIA cellular function primarily through an

egl-13-dependent pathway.

From these data we conclude that *ctbp-1* maintains AIA function and at

least some aspects of AIA gene expression by antagonizing *egl-13* function and

that *ctbp-1* likely also acts primarily through one or more *egl-13*-independent

pathways to maintain AIA cellular morphology.

egl-13 regulates AIA function through control of ceh-28 expression

We next asked if mutation of *egl-13* could suppress other *ctbp-1* mutant

AIA gene expression defects besides that of *ceh-28*. We crossed in *acbp-6*, *sra-*

11 and *glr-2* reporters to *egl-13 ctbp-1* double mutants and visualized reporter

expression at the L1 and L4 larval stages. We found that mutation of *egl-13*

suppressed P_{acbp-6}::*gfp* misexpression in the AIAs (Fig. 7A,D, compare to Fig. 5),

just as *egl-13* mutation suppressed P_{ceh-28}*::gfp* misexpression. By contrast,

mutation of *egl-13* had no effect on the loss of P_{sra-11}*::gfp* or P_{glr-2}*::gfp* expression

in *ctbp-1* mutants (Fig. 7B-D). We speculated that EGL-13 might directly regulate

expression of *ceh-28* or *acbp-6*. To test this hypothesis, we examined the *ceh-28*

and *acbp-6* promoter regions for possible EGL-13 binding sites. We failed to

identify any promising candidates, suggesting that regulation of these genes by

EGL-13 is likely indirect. These results demonstrate that some, though not all, of

the AIA gene expression defects seen in *ctbp-1* mutants are regulated through

egl-13.

As *egl-13* is required for misexpression of both *ceh-28* and *acbp-6* as well

as for disruption of AIA function in *ctbp-1* mutants, we hypothesized that

misexpressed *ceh-28* or *acbp-6* might be contributing to the observed AIA

functional defect in *ctbp-1* mutants. If so, we expected that mutations that

eliminated the functions of these ectopically expressed genes should restore AIA

function in *ctbp-1* mutants. To test this hypothesis, we crossed mutant alleles of

ceh-28 (cu11) or *acbp-6 (tm2995)* (both deletion alleles spanning greater than

half their respective genes) to *ctbp-1(n4784)* mutants and assayed the resulting

double mutants for butanone adaptation in L1 and L4 worms. We found that

acbp-6; ctbp-1 double mutants were nearly identical to both naïve and

conditioned *ctbp-1* single mutants at both the L1 and L4 larval stages (Fig. 7E-H),

indicating that misexpressed *acbp-6* is likely not responsible for the observed AIA

functional defect. Conditioned *ctbp-1 ceh-28* double mutants appeared similar to

both the wild type and *ctbp-1* single mutants at the L1 stage (Fig. 7I-J). These

double mutants display an minor, though not statistically significant, difference in

adaptation at the L4 larval stage when compared to either wild-type or *ctbp-1*

animals (Fig. 7K-L). We speculate that overexpression of *ceh-28,* caused by a

loss of *ctbp-1* and likely driven by ectopic *egl-13* activity, might partially account

for the defect in butanone adaptation seen in L4 and day 1 adult *ctbp-1* mutants

and that removal of *egl-13* in part restores AIA function by eliminating *ceh-28*

misexpression. We propose that *ctbp-1* functions to maintain aspects of the AIA

cell identity by preventing *egl-13* from promoting *ceh-28* expression and that *ceh-*

28 misexpression can perturb proper AIA function. These results also indicate

that *ceh-28* overexpression alone is not solely responsible for the observed AIA

functional defect, suggesting that the regulation of other, as-of-yet unidentified

genes controlled by *ctbp-1* (and potentially *egl-13*) also contribute to the

maintenance of the AIA cell identity.

Discussion

We have shown that the *ctbp-1* transcriptional corepressor gene is

required to maintain AIA cell identity and that *ctbp-1* negatively and selectively

regulates the function of the *egl-13* transcription factor gene. We suggest that

the CTBP-1 protein functions as a transcriptional corepressor to selectively

regulate the transcriptional output (either directly or indirectly) of the EGL-13

protein. c*tbp-1* mutant AIAs undergo a progressive decline in their initially wild-

type gene-expression pattern, morphology and function. *ctbp-1* can act cell-

autonomously and is able to act in older animals to maintain these aspects of the

AIA identity. We conclude that CTBP-1 functions to maintain AIA cell identity and

speculate that other transcriptional corepressors similarly function in the

maintenance of specific cell identities and do so by silencing undesired gene

expression through repression of transcriptional activators, such as EGL-13.

Such a mechanism could explain how the breadth of transcriptional activation by

terminal selectors can be fine-tuned in a coordinated fashion to fit the

requirements of specific cell types, with selective transcriptional silencing

providing a crucial aspect of proper cell-identity maintenance.

CTBP-1 might physically interact with EGL-13 to maintain the AIA cell

identity

The mammalian CtBPs (CtBP1 and CtBP2) bind PXDLS-like motifs on a

number of diverse transcription factors to target specific genetic loci for silencing

[49–51]. The mammalian ortholog of EGL-13, SOX6, interacts with the

mammalian ortholog of CTBP-1, CtBP2, through a PLNLS motif located in SOX6

to repress *Fgf-3* expression in the developing mouse auditory otic vesicle [74].

This motif is 100% conserved in *C. elegans* EGL-13. We speculate that CTBP-1

interacts with EGL-13 through its PLNLS motif to regulate EGL-13 activity as part

of AIA cell-identity maintenance. Specifically, we propose (Fig. 8) that CTBP-1

physically interacts with EGL-13 to target specific genetic loci for silencing as an

aspect of normal AIA cell-identity maintenance. Following the establishment of

the AIA cell fate, for which CTBP-1 is not required, CTBP-1 binds EGL-13,

recruiting CTBP-1 to EGL-13 DNA binding sites. CTBP-1 then silences

surrounding genetic loci, resulting in the repression of specific target genes (Fig.

8B). This repression is necessary for proper maintenance of the AIA cell identity,

and when disrupted, as in *ctbp-1* mutants, CTBP-1 binding partners, such as

EGL-13, inappropriately act as transcriptional activators in the AIAs, resulting in

disruption of AIA gene expression, morphology and function (Fig. 8C). In the

absence of such CTBP-1 interactors, as in *egl-13 ctbp-1* double mutants,

aberrant transcription is not activated and some of the defects in AIA

maintenance are avoided (Fig. 8D).

CTBP-1 likely utilizes additional transcription factors besides EGL-13 to

maintain the AIA cell identity

Our understanding of how CTBP-1 acts to maintain the AIA cell identity is

incomplete. While we have identified a few genes with expression that changes

in the absence of *ctbp-1* (*ceh-28*, *acbp-6*, *sra-11*, *glr-2*), none of these genes

seems to individually account for the full range of AIA defects seen in older *ctbp-*

1 mutants. We speculate that there are many more unidentified transcriptional

changes occurring in *ctbp-1* mutant AIAs that contribute to the observed AIA

morphological and functional defects.

Our observations suggest that EGL-13 is not the sole transcription factor

through which CTBP-1 functions to maintain AIA cell identity – neither AIA

morphological defects nor some AIA gene-expression defects (i.e. *sra-11* and *glr-*

2 expression) in *ctbp-1* mutants were suppressed in *egl-13 ctbp-1* double

mutants (Figs. 6D-G; 7A-B). We propose that CTBP-1 maintains different

aspects of AIA cell identity through interactions with multiple different

transcription factors. Given CTBP-1's known function as a transcriptional

corepressor and EGL-13's observed role in driving gene misexpression in the

absence of *ctbp-1*, we speculate that CTBP-1 likely utilizes not just EGL-13 but

also other transcription factors (possibly through interacting with PXDLS-like

motifs located in those transcription factors) to target multiple specific DNA

sequences for transcriptional silencing, effectively turning these transcription

factors into transcriptional repressors. When *ctbp-1* is absent, these unregulated

transcription factors can aberrantly function as transcriptional activators, resulting

in either the direct or indirect expression of genes which can in turn lead to

defects in other aspects of cell identity. Such a mechanism for the selective and

continuous silencing of multiple genetic loci in cell-type specific contexts by a

transcriptional corepressor like CTBP-1 might explain how the broad activating

activities of terminal selectors are restricted in the context of maintaining the

identities of distinct cell types.

CTBP-1 likely maintains the identities of other cells besides that of the AIAs

Others have previously reported a near pan-neuronal expression pattern

of *ctbp-1* in *C. elegans* [53], suggesting that *ctbp-1* might be acting in more cells

than just the AIAs to maintain cell identities. Why then have we thus far only

been able to identify defects in the maintenance of the AIA identity in *ctbp-1*

mutants? We speculate that, like the relatively subtle defects we have observed

in AIA gene expression, morphology and function, *ctbp-1* mutant defects in the

maintenance of other cell identities might be similarly subtle and easily missed if

not specifically sought. In addition, the AIAs might be particularly susceptible to

perturbations of maintenance of their identity, with defects manifesting either

earlier in the life of the cell or in more distinct ways (e.g. more gene

misexpression).

Both our findings and the work of others [54,55] provide further support for

the hypothesis that *ctbp-1* maintains other cell identities besides that of the AIAs.

We observed that *ctbp-1* mutants have an additional AIA-independent

chemotaxis defect (Figs. 4B-E; S5A-B), suggesting that other cells, likely neurons

that sense and/or execute responses to volatile odors, are also dysfunctional*.

Additionally, others have shown that, in *ctbp-1* mutants another pair of *C.*

elegans neurons, the SMDDs, display late-onset morphological abnormalities

coupled with a defect in *C. elegans* foraging behavior associated with these cells

[54,55], indicating that CTBP-1 might act to maintain SMDD cell identity as well.

The broad expression of *ctbp-1* throughout much of the *C. elegans* nervous

system is also consistent with the hypothesis that *ctbp-1* functions broadly to

maintain multiple neuronal cell identities [53].

Transcriptional corepressors might function broadly in the maintenance of

cell identities

The neuron-specific expression of *ctbp-1* [53] suggests that CTBP-1 likely

does not function in maintaining the identities of non-neuronal cells. How might

non-neuronal cell identities be maintained? We speculate that transcriptional

corepressors function in maintaining cell identities in both neuronal and non-

neuronal cells. There are known tissue-specific activities of other corepressor

complexes, such as those of NCoR1 in mediating the downstream effects of

hormone sensation in the mammalian liver [75,76] or of Transducin-Like

Enhancer of Split (TLE) in regulating gene expression and chromatin state in the

developing mouse heart and kidney [77–79]. We propose that, by analogy to

CTBP-1, distinct transcriptional corepressors might specialize in the maintenance

of a wide range of cell identities in distinct tissue types throughout metazoa.

Materials and Methods

C. elegans strains and transgenes

All *C. elegans* strains were grown on Nematode Growth Medium (NGM) plates

seeded with *E. coli* OP50 as described previously [80]. We used the N2 Bristol

strain as wild type. Worms were grown at 20°C unless otherwise indicated.

Standard molecular biology and microinjection methods, as previously described

[81], were used to generate transgenic worms.

The following strains were used in this study:

Strain	Genotype
N2	wild type
MT15670	*nIs175[P$_{ceh-28}$::gfp]*
MT15672	*nIs177[P$_{ceh-28}$::gfp]*
MT15677	*nIs175; ctbp-1(n4778)*
MT16225	*nIs175; ctbp-1(n4784)*
MT15688	*nIs175; ctbp-1(n4789)*
MT15801	*nIs175; ctbp-1(n4800)*
MT15805	*nIs175; ctbp-1(n4804)*
MT15806	*nIs175; ctbp-1(n4805)*
MT15809	*nIs175; ctbp-1(n4808)*
MT15811	*nIs175; ctbp-1(n4810)*
MT15813	*nIs175; ctbp-1(n4813)*
MT15820	*nIs175; ctbp-1(n4819)*
MT15824	*nIs175; ctbp-1(n4823)*
MT15825	*nIs175; ctbp-1(n4824)*
MT15841	*nIs177; ctbp-1(n4840)*
MT15850	*nIs177; ctbp-1(n4849)*
MT15853	*nIs177; ctbp-1(n4852)*
MT15862	*nIs177; ctbp-1(n4861)*
MT15865	*nIs177; ctbp-1(n4864)*
MT15866	*nIs177; ctbp-1(n4865)*
MT26446	*nIs175; ctbp-1(tm5512)*
MT15918	*nIs175* introgressed into CB4856 "Hawaiian" background
MT16295	*nIs177* introgressed into CB4856 background
MT26522	*nIs175; ctbp-1(n4784)* introgressed into CB4856 background
MT23360	*nIs175; ctbp-1(n4784); nEx2346[ctbp-1(+)]*

MT23361	*nIs175; ctbp-1(n4784); nEx2347[ctbp-1(+)]*
MT23714	*nIs175; ctbp-1(n4784); nIs743[P$_{gcy-28.d}$::ctbp-1(+)]*
MT25271	*nIs843[P$_{gcy-28.d}$::mCherry]*
MT26437	*nIs175; ctbp-1(n4784); nIs843*
MT23365	*nIs175; ctbp-1(n4784); nEx2351[P$_{hsp-16.2}$::ctbp-1(+);P$_{hsp-16.41}$::ctbp-1(+)]*
MT18778	*nIs348[P$_{ceh-28}$::mCherry]; lin-15AB(n765ts)*
MT20844	*nIs348[P$_{ceh-28}$::mCherry]; ctbp-1(n4784)*
NH2466	*ayIs4[P$_{egl-17}$::gfp]; dpy-20(e1282ts)*
MT26417	*ayIs4; nIs348; ctbp-1(n4784)*
BW1946	*ctIs43[P$_{dbl-1}$::gfp] unc-42(e270)*
MT23726	*nIs348; ctIs43 unc-42(e270); ctbp-1(n4784)*
MT20852	*nIs491[P$_{ser-7.b}$::mCherry]*
MT23427	*nIs491; ctbp-1(n4784)*
NY2080	*ynIs80[P$_{flp-21}$::gfp]*
MT23718	*nIs348; ctbp-1(n4784); ynIs80*
OH10237	*otIs326[P$_{ins-1}$::gfp]*
MT26422	*ctbp-1(n4784); otIs326*
JN1716	*peIs1716[P$_{ins-1s}$::gfp;P$_{ttx-3}$::mCherry]*
MT23717	*nIs348; ctbp-1(n4784); peIs1716*
OH11030	*otIs317[P$_{mgl-1}$::mCherry]; otIs379[P$_{cho-1}$::gfp]*
MT26421	*nIs348; ctbp-1(n4784); otIs317; otIs379*
MT26420	*ctbp-1(n4784); otIs317*
MT25268	*nIs840[P$_{gcy-28.d}$::gfp]*
MT25270	*nIs842[P$_{gcy-28.d}$::gfp]*
MT26412	*nIs348; ctbp-1(n4784); nIs840*
MT26438	*nIs348; ctbp-1(n4784); nIs743; nIs840*
MT26439	*nIs348; ctbp-1(n4784); nIs840; nEx2351*
JN580	*peIs580[P$_{ins-1s}$::casp1;P$_{ins-1s}$::venus;P$_{unc-122}$::gfp]*
MT23746	*nIs175; egl-13(n5937) ctbp-1(n4784)*
MT24129	*nIs175; egl-13(n6013) ctbp-1(n4784)*
MT25352	*nIs175; egl-13(n6313) ctbp-1(n4784)*
MT26486	*nIs175; egl-13(n5937) ctbp-1(n4784); nEx3062[egl-13(+)]*
MT26487	*nIs175; egl-13(n5937) ctbp-1(n4784); nEx3063[egl-13(+)]*
MT26549	*nIs175; egl-13(n6013) ctbp-1(n4784); nEx3080[egl-13(+)]*
MT26523	*nIs175; egl-13(n6313) ctbp-1(n4784); nEx3074[egl-13(+)]*
MT26548	*nIs175; egl-13(n6313) ctbp-1(n4784); nEx3079[egl-13(+)]*
MT26481	*nIs175; egl-13(n5937) ctbp-1(n4784); nEx3055[P$_{gcy-28.d}$::egl-13(+)]*
MT26441	*nIs175; egl-13(n5937) ctbp-1(n4784); nIs840*
MT26415	*evIs111[P$_{rgef-1}$::gfp]; nIs843*
MT26416	*ctbp-1(n4784); evIs111; nIs843*
MT26444	*otIs123[P$_{sra-11}$::gfp]; nIs843*
MT26445	*nIs348; ctbp-1(n4784); otIs123*
MT26524	*nIs348; egl-13(n5937) ctbp-1(n4784); otIs123*
MT26504	*nIs843; ivEx138[P$_{glr-2}$::gfp]*

MT26505	*nIs348; ctbp-1(n4784); ivEx138*
MT26550	*nIs348; egl-13(n5937) ctbp-1(n4784); ivEx138*
MT26581	*nIs843; nEx3081[P$_{acbp-6}$::gfp]*
MT26551	*nIs348; ctbp-1(n4784); nEx3081*
MT26582	*nIs348; egl-13(n5937) ctbp-1(n4784); nEx3081*
MT26605	*acbp-6(tm2995); nIs175; ctbp-1(n4784)*
MT23725	*nIs175; ctbp-1(n4784) ceh-28(cu11)*

Plasmid construction

The *nIs175[P$_{ceh-28}$::gfp], nIs177[P$_{ceh-28}$::gfp]* and *nIs348[P$_{ceh-28}$::mCherry]*

transgenes have been previously described [59]. *nIs743[P$_{gcy-28.d}$::ctbp-1(+)]*

contains 3.0 kb of the 5' promoter of *gcy-28.d* fused to the *ctbp-1a* coding region

inserted into plasmid pPD49.26. *nIs840[P$_{gcy-28.d}$::gfp]* contains 3.0 kb of the 5'

promoter of *gcy-28.d* inserted into pPD95.77. *nIs843[P$_{gcy-28.d}$::mCherry]* contains

3.0 kb of the 5' promoter of *gcy-28.d* inserted into pPD122.56 containing

mCherry. *nEx2351[P$_{hsp-16.2}$::ctbp-1(+);P$_{hsp-16.41}$::ctbp-1(+)]* contains *ctbp-1a*

cDNA, isolated by RT-PCR, inserted into pPD49.73 and pPD49.83.

nEx3055[P$_{gcy-28.d}$::egl-13(+)] contains 3.0 kb of the 5' promoter of *gcy-28.d* fused

to the *egl-13* coding region inserted into pPD49.26. *nEx3081[P$_{acbp-6}$::gfp]*

contains 2.0 kb of the 5' promoter of *acbp-6* inserted into pPD122.56. Plasmid

construction was performed using Infusion cloning enzymes (Takara Bio,

Mountain View, CA).

Mutagenesis screens

ctbp-1 mutants were isolated from genetic screens for mutations that cause the

survival of the M4 sister cell as scored by extra GFP-positive cells carrying the

M4-cell-specific markers *nIs175[P$_{ceh-28}$::gfp]* or *nIs177[P$_{ceh-28}$::gfp]* [58,59]. *egl-13*

mutants were isolated from genetic screens for mutations that suppress *nls175*

misexpression in the AIAs of *ctbp-1(n4784)* mutants while retaining GFP

expression in the M4 neuron. For both screens, mutagenesis was performed with

ethyl methanesulfonate (EMS) as previously described [80]. Mutagenized P_0

animals were allowed to propagate, and their F_2 progeny were synchronized by

hypochlorite treatment and screened at the L4 stage for extra GFP-positive cells

(*ctbp-1* screens) or fewer GFP-positive cells (suppressor screens) on a

dissecting microscope equipped to examine fluorescence. From both screens,

mutant alleles were grouped into functional groups by complementation testing

when possible. Mutants were mapped using SNP mapping [70] by crossing

mutants to strains containing *nls175*, *nls177*, or *nls175;ctbp-1(n4784)*

introgressed into the Hawaiian strain CB4856. Whole-genome sequencing was

performed on mutants and a combination of functional groupings and mapping

data suggested genes with mutations that were likely causal for the mutant

phenotypes. Rescue of mutant phenotypes with wild-type *ctbp-1(+)* and *egl-13(+)*

constructs as well as the mutant phenotype of a separately isolated deletion

allele of *ctbp-1, tm5512,* confirmed the identities of the causal mutations.

Microscopy

All images were obtained using an LSM 800 confocal microscope (Zeiss) and

ZEN software. Images were processed and prepared for publication using FIJI

software and Adobe Illustrator.

Heat-shock assays

Rescue of AIA defects in day 1 adult worms was assayed using the

nEx2351[P_{hsp-16.2}::ctbp-1;P_{hsp-16.41}::ctbp-1] transgene. Worms were synchronized

and grown at 20°C. Subsets of L1 and L4 worms carrying *nEx2351* were

removed from this population for scoring at the appropriate stages. At the L4

stage, half of the worms were heat-shocked at 34°C for 30 minutes and returned

to 20°C for 24 hours while the other half remained at 20°C throughout. After 24

hours, heat-shocked and non-heat-shocked worms carrying *nEx2351* were

scored.

Single-cell RNA-sequencing

Dissociation of animals into cell suspensions. Single-cell suspensions were

generated as described [82–84] with minor modifications. Briefly, synchronized

populations of worms were grown on NGM plates seeded with OP50 to the L4

larval stage. Worms were harvested from these plates, washed three times with

M9 buffer and treated with SDS-DTT (200 mM DTT, 0.25% SDS, 20 mM HEPES,

3% sucrose, pH 8.0) for two to three minutes. Worms were washed five times

with 1x PBS and treated with pronase (15 mg/mL) for 20-23 minutes. During the

pronase treatment, worm suspensions were pipetted with a P200 pipette rapidly

for four sets of 80 repetitions. The pronase treatment was stopped by the

addition of L-15-10 media (90% L-15 media, 10% FBS). The suspension was

then passed through a 35 μm nylon filter into a collection tube, washed once with

1x PBS, and prepared for FACS.

FACS of fluorescently-labeled neurons. FACS was performed using a BD

FACSAria III cell sorter running BD FACS Diva software. DAPI was added to

samples at a final concentration of 1 μg/mL to label dead and dying cells. GFP-

positive, DAPI-negative neurons were sorted from the single-cell suspension into

1x PBS containing 1% FBS. Non-fluorescent and single-color controls were used

to set gating parameters. Cells were then concentrated and processed for single-

cell sequencing.

Single-cell sequencing. Samples were processed for single-cell sequencing

using the 10X Genomics Chromium 3'mRNA-sequencing platform. Libraries were

prepared using the Chromium Next GEM Single Cell 3' Kit v3.1 according to the

manufacturer's protocol. The libraries were sequenced using an Illumina

NextSeq 500 with 75 bp paired end reads.

Single-cell RNA-sequencing data processing. Data processing was performed

using 10X Genomics' CellRanger software (v4.0.0). Reads were mapped to the

C. elegans reference genome from Wormbase, version WBcel235. For

visualization and analysis of data, we used 10X Genomics' Loupe Browser

(v4.2.0). AIAs were identified by expression of multiple AIA markers confirmed to

be expressed in both wild-type and *ctbp-1* mutant AIAs (i.e. *gcy-28, ins-1, cho-1*;

Fig. 2B). Candidate genes for misexpression (either ectopic or missing) in mutant

AIAs were identified and tested as described in the text.

Morphology scoring

We assayed AIA morphology by visualizing and imaging AIAs expressing *nIs840*

using an LSM 800 confocal microscope (Zeiss) and a 63x objective. AIA cell

body length and area were quantified using FIJI software.

Image blinding and scoring

A subset of 60 wild-type and 60 *ctbp-1* mutant images per stage (randomly

chosen from the existing images taken to measure AIA cell body length) were

selected and the genotype of each was blinded. Blinded images were then

scored as either "Normal" or "Elongated" in appearance in batches of 40 images

(20 each of wild-type and *ctbp-1* mutant, randomly assorted), repeated three

times per stage. Scored images were then matched back to their genotypes and

percentage of AIAs scored as "Elongated" per genotype was calculated and

graphed.

Behavioral Assays

Butanone adaptation. Assay conditions were adapted from Cho et al., 2016

[66]. Staged worms were washed off non-crowded NGM plates seeded with *E.*

coli OP50 with S basal. Worms were washed two times with S basal and split

evenly into the "naïve" and "conditioned" populations. Naïve worms were

incubated in 1 mL S basal for 90 minutes. Conditioned worms were incubated in

1 mL S basal with 2-butanone diluted to a final concentration of 120 µM for 90

minutes. During conditioning, unseeded NGM plates were spotted with two 1 µL

drops of 10% ethanol ("control") and two 1 µL drops of 2-butanone diluted in 10%

ethanol at 1:1000 ("odor") as well as four 1 µL drops of 1 M NaN_3 at the same

loci. After conditioning, both populations were washed three more times in S

basal and placed at the center of the unseeded NGM plates. Worms were

allowed to chemotax for two hours. Plates were moved to 4°C for 30-60 minutes

to stop the assay and then scored. Worms that had left the origin were scored as

chemotaxing to the odor spots ("#odor") or control spots ("#control"), and a

chemotaxis index was determined as (#odor - #control) / (#odor + #control).

Assays were repeated on at least three separate days with one to three plates

per strain ran in parallel on any given day based on the number of appropriately-

staged worms available. Plates in which fewer than 50 worms left the origin were

not scored.

Chemotaxis assays. L4 worms were washed off non-crowded NGM plates

seeded with *E. coli* OP50 with S basal. Worms were washed three times with S

basal. Unseeded NGM plates were spotted with two 1 µL drops of 100% ethanol

("control") and two 1 µL drops of diacetyl diluted in 100% ethanol at 1:1000 or

two 1 µL drops of isoamyl alcohol diluted in 100% ethanol at 1:100 ("odor") as

well as four 1 µL drops of 1 M NaN_3 at the same loci. Worms were placed at the

center of the unseeded NGM plates. Worms were allowed to chemotax for two

hours. Plates were moved to 4°C for 30-60 minutes to stop the assay and then

scored. Worms that had left the origin were scored, and a chemotaxis index was

determined as above. Assays were repeated on at least three separate days.

Plates in which fewer than 40 worms left the origin were not scored.

Statistical analyses

Unpaired t-tests were used for the comparisons of AIA gene expression and AIA

morphological features. One-way ANOVA tests with Tukey's correction were

used for comparisons of AIA morphological features and for behavioral assays.

Statistical tests were performed using GraphPad Prism software (GraphPad

Prism version 6.0h, RRID: SCR_002798).

Accession Number

The GEO accession number for the RNA-Seq dataset in this paper is

GSE179484.

Acknowledgments

We thank N. An, R. Droste, S. Mitani, and the *Caenorhabditis* Genetics Center (CGC), which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440), for strains and reagents. We thank C. Diehl, E. Lee, C. Pender, S. Sando, V. Dwivedi, K. Burkhart, S. Wong, C. Fincher, C. Cho, G. Johnson, D. Lee, D. Ghosh, A. Amon, P. Reddien, and Horvitz laboratory members for discussion and advice.

Funding

National Institutes of Health (GM024663)

- Josh Saul
- H. Robert Horvitz

National Institutes of Health (T32GM007287)

- Josh Saul

Howard Hughes Medical Institute

- Josh Saul
- Takashi Hirose
- H. Robert Horvitz

Friends of the McGovern Institute Fellowship (2733360)

- Josh Saul

Author Contributions

H.R.H. supervised the project. T.H. initiated the project. J.S. and T.H. designed

and performed experiments, generated reagents, and analyzed data. All authors

contributed to interpretation of data. J.S. wrote the original manuscript draft. All

authors contributed to review and editing of the manuscript.

Declaration of Interests

The authors declare no competing interests.

References

References

1.   Davidson EH, Cameron RA, Ransick A. Specification of cell fate in the sea

urchin embryo: summary and some proposed mechanisms. Dev Camb Engl.

1998 Sep;125(17):3269–90.

2.   Davidson EH. A Genomic Regulatory Network for Development. Science.

2002 Mar 1;295(5560):1669–78.

3.   Levine M, Davidson EH. Gene regulatory networks for development. Proc

Natl Acad Sci. 2005 Apr 5;102(14):4936–42.

4.   Hobert O. A map of terminal regulators of neuronal identity in

Caenorhabditis elegans. Wiley Interdiscip Rev Dev Biol. 5(4):474–98.

5.   Hsieh J, Zhao X. Genetics and Epigenetics in Adult Neurogenesis. Cold

Spring Harb Perspect Biol [Internet]. 2016 Jun [cited 2021 Mar 18];8(6).

Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4888816/

6.   Homem CCF, Repic M, Knoblich JA. Proliferation control in neural stem and

progenitor cells. Nat Rev Neurosci. 2015 Nov;16(11):647–59.

7.   Altun-Gultekin Z, Andachi Y, Tsalik EL, Pilgrim D, Kohara Y, Hobert O. A

regulatory cascade of three homeobox genes, ceh-10, ttx-3 and ceh-23,

controls cell fate specification of a defined interneuron class in C. elegans.

Development. 2001 Jun 1;128(11):1951–69.

8.   Zeng H, Sanes JR. Neuronal cell-type classification: challenges,

opportunities and the path forward. Nat Rev Neurosci. 2017 Sep;18(9):530–

46.

9.   Baker NE. Master regulatory genes; telling them what to do. BioEssays.

2001;23(9):763–6.

10. Deneris ES, Hobert O. Maintenance of postmitotic neuronal cell identity. Nat

Neurosci. 2014 Jul;17(7):899–907.

11. Hobert O, Carrera I, Stefanakis N. The molecular and gene regulatory

signature of a neuron. Trends Neurosci. 2010 Oct 1;33(10):435–45.

12. Masoudi N, Tavazoie S, Glenwinkel L, Ryu L, Kim K, Hobert O.

Unconventional function of an Achaete-Scute homolog as a terminal

selector of nociceptive neuron identity. PLOS Biol. 2018 Apr

19;16(4):e2004979.

13. Weintraub H, Dwarki VJ, Verma I, Davis R, Hollenberg S, Snider L, et al.

Muscle-specific transcriptional activation by MyoD. Genes Dev. 1991 Aug

1;5(8):1377–86.

14. Lassar AB. Finding MyoD and lessons learned along the way. Semin Cell

Dev Biol. 2017 Dec;72:3–9.

15. Wardle FC. Master control: transcriptional regulation of mammalian Myod. J

Muscle Res Cell Motil. 2019 Jun;40(2):211–26.

16. Halder G, Callaerts P, Gehring WJ. Induction of ectopic eyes by targeted

expression of the eyeless gene in Drosophila. Science. 1995 Mar

24;267(5205):1788–92.

17. Gehring WJ. The master control gene for morphogenesis and evolution of

the eye. Genes Cells. 1996;1(1):11–5.

18. Shen W, Mardon G. Ectopic eye development in Drosophila induced by

directed dachshund expression. Dev Camb Engl. 1997 Jan;124(1):45–52.

19. Treisman JE. Retinal differentiation in Drosophila. WIREs Dev Biol.

2013;2(4):545–57.

20. Lima Cunha D, Arno G, Corton M, Moosajee M. The Spectrum of PAX6

Mutations and Genotype-Phenotype Correlations in the Eye. Genes. 2019

Dec;10(12):1050.

21. Fukushige T, Hawkins MG, McGhee JD. The GATA-factor elt-2 is essential

for formation of the Caenorhabditis elegans intestine. Dev Biol. 1998 Jun

15;198(2):286–302.

22. Fukushige T, Hendzel MJ, Bazett-Jones DP, McGhee JD. Direct

visualization of the elt-2 gut-specific GATA factor binding to a target

promoter inside the living Caenorhabditis elegans embryo. Proc Natl Acad

Sci. 1999 Oct 12;96(21):11883–8.

23. McGhee JD, Fukushige T, Krause MW, Minnema SE, Goszczynski B,

Gaudet J, et al. ELT-2 Is the Predominant Transcription Factor Controlling

Differentiation and Function of the C. elegans Intestine, from Embryo to

Adult. Dev Biol. 2009 Mar 15;327(2):551–65.

24. Block DH, Shapira M. GATA transcription factors as tissue-specific master

regulators for induced responses. Worm. 2015 Oct 2;4(4):e1118607.

25. Matson CK, Murphy MW, Sarver AL, Griswold MD, Bardwell VJ, Zarkower

D. DMRT1 prevents female reprogramming in the postnatal mammalian

testis. Nature. 2011 Aug 4;476(7358):101–4.

26. Mall M, Kareta MS, Chanda S, Ahlenius H, Perotti N, Zhou B, et al. Myt1l

safeguards neuronal identity by actively repressing many non-neuronal

fates. Nature [Internet]. 2017 Apr 5 [cited 2017 Apr 7];advance online

publication. Available from:

http://www.nature.com/nature/journal/vaop/ncurrent/full/nature21722.html

27. Simon HH, Thuret S, Alberi L. Midbrain dopaminergic neurons: control of

their cell fate by the engrailed transcription factors. Cell Tissue Res. 2004

Sep 1;318(1):53–61.

28. Vissers JHA, Froldi F, Schröder J, Papenfuss AT, Cheng LY, Harvey KF.

The Scalloped and Nerfin-1 Transcription Factors Cooperate to Maintain

Neuronal Cell Fate. Cell Rep. 2018 Nov 6;25(6):1561-1576.e7.

29. Hsiao H-Y, Jukam D, Johnston R, Desplan C. The neuronal transcription

factor erect wing regulates specification and maintenance of Drosophila R8

photoreceptor subtypes. Dev Biol. 2013 Sep 15;381(2):482–90.

30. Riddle MR, Weintraub A, Nguyen KCQ, Hall DH, Rothman JH.

Transdifferentiation and remodeling of post-embryonic C. elegans cells by a

single transcription factor. Development. 2013 Dec 15;140(24):4844–9.

31. O'Meara MM, Zhang F, Hobert O. Maintenance of Neuronal Laterality in

Caenorhabditis elegans Through MYST Histone Acetyltransferase Complex

Components LSY-12, LSY-13 and LIN-49. Genetics. 2010 Dec

1;186(4):1497–502.

32. Xu J, Hao X, Yin M-X, Lu Y, Jin Y, Xu J, et al. Prevention of medulla neuron

dedifferentiation by Nerfin-1 requires inhibition of Notch activity.

Development. 2017 Apr 15;144(8):1510–7.

33. Hobert O. Regulatory logic of neuronal diversity: Terminal selector genes

and selector motifs. Proc Natl Acad Sci. 2008 Dec 23;105(51):20067–71.

34. Hobert O. Regulation of Terminal Differentiation Programs in the Nervous

System. Annu Rev Cell Dev Biol. 2011;27(1):681–96.

35. Hobert O, Kratsios P. Neuronal identity control by terminal selectors in

worms, flies, and chordates. Curr Opin Neurobiol. 2019 Jun 1;56:97–105.

36. Hobert O. Terminal Selectors of Neuronal Identity. Curr Top Dev Biol.

2016;116:455–75.

37. Zhang F, Bhattacharya A, Nelson JC, Abe N, Gordon P, Lloret-Fernandez C,

et al. The LIM and POU homeobox genes ttx-3 and unc-86 act as terminal

selectors in distinct cholinergic and serotonergic neuron types.

Development. 2014 Jan 15;141(2):422–35.

38. Serrano-Saiz E, Poole RJ, Felton T, Zhang F, De La Cruz ED, Hobert O.

Modular Control of Glutamatergic Neuronal Identity in C. elegans by Distinct

Homeodomain Proteins. Cell. 2013 Oct 24;155(3):659–73.

39. Duggan A, Ma C, Chalfie M. Regulation of touch receptor differentiation by

the Caenorhabditis elegans mec-3 and unc-86 genes. Dev Camb Engl. 1998

Oct;125(20):4107–19.

40. Kim J, Yeon J, Choi S-K, Huh YH, Fang Z, Park SJ, et al. The Evolutionarily

Conserved LIM Homeodomain Protein LIM-4/LHX6 Specifies the Terminal

845   Identity of a Cholinergic and Peptidergic C. elegans Sensory/Inter/Motor

846   Neuron-Type. PLOS Genet. 2015 Aug 25;11(8):e1005480.

847 41. Alqadah A, Hsieh Y-W, Vidal B, Chang C, Hobert O, Chuang C-F.

848   Postmitotic diversification of olfactory neuron types is mediated by

849   differential activities of the HMG-box transcription factor SOX-2. EMBO J.

850   2015 Oct 14;34(20):2574–89.

851 42. Kerk SY, Kratsios P, Hart M, Mourao R, Hobert O. Diversification of

852   C. elegans Motor Neuron Identity via Selective Effector Gene Repression.

853   Neuron. 2017 Jan 4;93(1):80–98.

854 43. Zhou HM, Walthall WW. UNC-55, an Orphan Nuclear Hormone Receptor,

855   Orchestrates Synaptic Specificity among Two Classes of Motor Neurons in

856   Caenorhabditis elegans. J Neurosci. 1998 Dec 15;18(24):10438–44.

857 44. Winnier AR, Meir JY-J, Ross JM, Tavernarakis N, Driscoll M, Ishihara T, et

858   al. UNC-4/UNC-37-dependent repression of motor neuron-specific genes

859   controls synaptic choice in Caenorhabditis elegans. Genes Dev. 1999 Nov

860   1;13(21):2774–86.

861 45. Yu B, Wang X, Wei S, Fu T, Dzakah EE, Waqas A, et al. Convergent

862   Transcriptional Programs Regulate cAMP Levels in C. elegans GABAergic

863   Motor Neurons. Dev Cell. 2017 Oct 23;43(2):212-226.e7.

864 46. Wyler SC, Spencer WC, Green NH, Rood BD, Crawford L, Craige C, et al.

865   Pet-1 Switches Transcriptional Targets Postnatally to Regulate Maturation of

866   Serotonin Neuron Excitability. J Neurosci. 2016 Feb 3;36(5):1758–74.

47. Turner J, Crossley M. The CtBP family: enigmatic and enzymatic

transcriptional co-repressors. BioEssays. 2001 Aug 1;23(8):683–90.

48. Chinnadurai G. CtBP, an Unconventional Transcriptional Corepressor in

Development and Oncogenesis. Mol Cell. 2002 Feb;9(2):213–24.

49. Chinnadurai G. CtBP family proteins: More than transcriptional

corepressors. BioEssays. 2003 Jan 1;25(1):9–12.

50. Shi Y, Sawada J, Sui G, Affar EB, Whetstine JR, Lan F, et al. Coordinated

histone modifications mediated by a CtBP co-repressor complex. Nature.

2003 Apr 17;422(6933):735–8.

51. Stankiewicz TR, Gray JJ, Winter AN, Linseman DA. C-terminal binding

proteins: central players in development and disease. Biomol Concepts.

2014;5(6):489–511.

52. Nicholas HR, Lowry JA, Wu T, Crossley M. The Caenorhabditis elegans

Protein CTBP-1 Defines a New Group of THAP Domain-Containing CtBP

Corepressors. J Mol Biol. 2008 Jan 4;375(1):1–11.

53. Reid A, Yücel D, Wood M, Llamosas E, Kant S, Crossley M, et al. The

transcriptional repressor CTBP-1 functions in the nervous system of

Caenorhabditis elegans to regulate lifespan. Exp Gerontol. 2014

Dec;60:153–65.

54. Reid A, Sherry TJ, Yücel D, Llamosas E, Nicholas HR. The C-terminal

binding protein (CTBP-1) regulates dorsal SMD axonal morphology in

Caenorhabditis elegans. Neuroscience. 2015 Dec 17;311:216–30.

55. Sherry T, Handley A, Nicholas HR, Pocock R. Harmonization of L1CAM

expression facilitates axon outgrowth and guidance of a motor neuron.

Development [Internet]. 2020 Jan 1 [cited 2020 Oct 8]; Available from:

https://dev.biologists.org/content/early/2020/10/07/dev.193805

56. Petersen JG, Romanos TR, Juozaityte V, Riveiro AR, Hums I, Traunmüller

L, et al. EGL-13/SoxD Specifies Distinct O2 and CO2 Sensory Neuron Fates

in Caenorhabditis elegans. PLOS Genet. 2013 May 9;9(5):e1003511.

57. Cinar HN, Richards KL, Oommen KS, Newman AP. The EGL-13 SOX

domain transcription factor affects the uterine pi cell lineages in

Caenorhabditis elegans. Genetics. 2003 Nov;165(3):1623–8.

58. Hirose T, Horvitz HR. An Sp1 transcription factor coordinates caspase-

dependent and -independent apoptotic pathways. Nature. 2013

Aug;500(7462):354–8.

59. Hirose T, Galvin BD, Horvitz HR. Six and Eya promote apoptosis through

direct transcriptional activation of the proapoptotic BH3-only gene egl-1 in

Caenorhabditis elegans. Proc Natl Acad Sci. 2010 Aug 31;107(35):15479–

84.

60. C. elegans Deletion Mutant Consortium. large-scale screening for targeted

knockouts in the Caenorhabditis elegans genome. G3 Bethesda Md. 2012

Nov;2(11):1415–25.

61. Ramakrishnan K, Okkema PG. Regulation of C. elegans Neuronal

Differentiation by the ZEB-Family Factor ZAG-1 and the NK-2

Homeodomain Factor CEH-28. PLOS ONE. 2014 Dec 4;9(12):e113893.

62. Ramakrishnan K, Ray P, Okkema PG. CEH-28 activates dbl-1 expression

and TGF-β signaling in the C. elegans M4 neuron. Dev Biol. 2014 Jun

15;390(2):149–59.

63. Tomioka M, Adachi T, Suzuki H, Kunitomo H, Schafer WR, Iino Y. The

Insulin/PI 3-Kinase Pathway Regulates Salt Chemotaxis Learning in

Caenorhabditis elegans. Neuron. 2006 Sep 7;51(5):613–25.

64. Iino Y, Yoshida K. Parallel use of two behavioral mechanisms for

chemotaxis in Caenorhabditis elegans. J Neurosci Off J Soc Neurosci. 2009

Apr 29;29(17):5370–80.

65. Shinkai Y, Yamamoto Y, Fujiwara M, Tabata T, Murayama T, Hirotsu T, et

al. Behavioral Choice between Conflicting Alternatives Is Regulated by a

Receptor Guanylyl Cyclase, GCY-28, and a Receptor Tyrosine Kinase,

SCD-2, in AIA Interneurons of Caenorhabditis elegans. J Neurosci. 2011

Feb 23;31(8):3007–15.

66. Cho CE, Brueggemann C, L'Etoile ND, Bargmann CI. Parallel encoding of

sensory history and behavioral preference during Caenorhabditis elegans

olfactory learning. eLife. 2016 Jul 6;5:e14000.

67. Shaye DD, Greenwald I. OrthoList: A Compendium of C. elegans Genes

with Human Orthologs. PLOS ONE. 2011 May 25;6(5):e20085.

68. Troemel ER, Chou JH, Dwyer ND, Colbert HA, Bargmann CI. Divergent

seven transmembrane receptors are candidate chemosensory receptors in

C. elegans. Cell. 1995 Oct 20;83(2):207–18.

69. Brockie PJ, Madsen DM, Zheng Y, Mellem J, Maricq AV. Differential

Expression of Glutamate Receptor Subunits in the Nervous System of

Caenorhabditis elegans and Their Regulation by the Homeodomain Protein

UNC-42. J Neurosci. 2001 Mar 1;21(5):1510–22.

70. Davis MW, Hammarlund M, Harrach T, Hullett P, Olsen S, Jorgensen EM.

Rapid single nucleotide polymorphism mapping in C. elegans. BMC

Genomics. 2005;6:118.

71. Feng G, Yi P, Yang Y, Chai Y, Tian D, Zhu Z, et al. Developmental stage-

dependent transcriptional regulatory pathways control neuroblast lineage

progression. Development. 2013 Sep 15;140(18):3838–47.

72. Ji EH, Kim J. SoxD Transcription Factors: Multifaceted Players of Neural

Development. Int J Stem Cells. 2016 May 30;9(1):3–8.

73. Saleem M, Barturen-Larrea P, Gomez JA. Emerging roles of Sox6 in the

renal and cardiovascular system. Physiol Rep. 2020 Nov;8(22):e14604.

74. Murakami A, Ishida S, Thurlow J, Revest J-M, Dickson C. SOX6 binds

CtBP2 to repress transcription from the Fgf-3 promoter. Nucleic Acids Res.

2001 Aug 15;29(16):3347–55.

75. Feng X, Jiang Y, Meltzer P, Yen PM. Transgenic Targeting of a Dominant

Negative Corepressor to Liver Blocks Basal Repression by Thyroid

Hormone Receptor and Increases Cell Proliferation*. J Biol Chem. 2001 Jan

1;276(18):15066–72.

76. Mottis A, Mouchiroud L, Auwerx J. Emerging roles of the corepressors

NCoR1 and SMRT in homeostasis. Genes Dev. 2013 Apr 15;27(8):819–35.

77. Sharma M, Brantley JG, Vassmer D, Chaturvedi G, Baas J, Vanden Heuvel

GB. The homeodomain protein Cux1 interacts with Grg4 to repress p27kip1

expression during kidney development. Gene. 2009 Jun 15;439(1):87–94.

78. Kaltenbrun E, Greco TM, Slagle CE, Kennedy LM, Li T, Cristea IM, et al. A

Gro/TLE-NuRD Corepressor Complex Facilitates Tbx20-Dependent

Transcriptional Repression. J Proteome Res. 2013 Dec 6;12(12):5395–409.

79. Agarwal M, Kumar P, Mathew SJ. The Groucho/Transducin-like enhancer of

split protein family in animal development. IUBMB Life. 2015 Jul

1;67(7):472–81.

80. Brenner S. The Genetics of CAENORHABDITIS ELEGANS. Genetics. 1974

May;77(1):71–94.

81. Mello CC, Kramer JM, Stinchcomb D, Ambros V. Efficient gene transfer in

C.elegans: extrachromosomal maintenance and integration of transforming

sequences. EMBO J. 1991 Dec;10(12):3959–70.

82. Taylor SR, Santpere G, Reilly M, Glenwinkel L, Poff A, McWhirter R, et al.

Expression profiling of the mature C. elegans nervous system by single-cell

RNA-Sequencing. bioRxiv. 2019 Aug 17;737577.

83. Kaletsky R, Lakhina V, Arey R, Williams A, Landis J, Ashraf J, et al. The C.

elegans adult neuronal IIS/FOXO transcriptome reveals adult phenotype

regulators. Nature. 2016 Jan;529(7584):92–6.

84. Zhang S. Cell isolation and culture. WormBook. 2013 Feb 21;1–39.

Figures

Figure 1

 Figure 1. *ctbp-1* mutants misexpress *P_ceh-28_::gfp* in the AIA neurons

(A) Expression of the M4-specific marker $nIs175[P_{ceh-28}::gfp]$ in the wild type (left

panel), a *ctbp-1(n4784)* mutant (middle panel), and a *ctbp-1* mutant carrying an

extrachromosomal array expressing wild-type *ctbp-1* under its native promoter

(*nEx2347*) (right panel). Arrow, M4 neuron. Circle, AIAs. Scale bar, 10 μm.

(B) A *ctbp-1(n4784)* mutant expressing *nIs175* (left panel) and the AIA marker

$nIs843[P_{gcy-28.d}::mCherry]$ (middle panel). Merge, right panel. Arrow, M4 neuron.

Circle, AIAs. Scale bar, 10 μm.

(C) Gene diagram of the *ctbp-1a* isoform. Arrows (above), point mutations. Line

(below), deletion. Scale bar (bottom right), 1 kb. Additional *ctbp-1* alleles are

shown in Fig. S1B.

(D) *nIs175* expression in wild-type (top) and *ctbp-1(n4784)* (bottom) worms at the

L1 larval stage (left) and L4 larval stage (right). Arrow, M4 neuron. Circle, AIAs.

Scale bar, 10 μm.

(E) Percentage of wild-type, *ctbp-1(n4784)*, and *ctbp-1(n4808)* worms expressing

nIs175 in the AIA neurons over time. Time points correspond to the L1, L2, L3,

and L4 larval stages, early adult, and day 1 adult worms (indicated below X axis).

Mean ± SEM. $n \geq 60$ worms scored per strain per stage, 4 biological replicates.

(F) Expression of *nIs175* in *ctbp-1* mutants containing a transgene driving

expression of wild-type *ctbp-1* under an AIA-specific promoter ($nIs743[P_{gcy-}$

$_{28.d}::ctbp-1(+)]$) in L1 and L4 larval worms. Arrow, M4 neuron. Circle, AIAs. Scale

bar, 10 μm.

(G) Schematic for the heat shock experiment shown in Fig. 1H.

(H) *nIs175* expression in *ctbp-1(n4784)* mutants carrying the heat shock-

inducible transgene *nEx2351[P$_{hsp-16.2}$::ctbp-1(+);P$_{hsp-16.41}$::ctbp-1(+)]*. Arrow, M4

neuron. Circle, AIAs. Scale bar, 10 μm.

All strains shown contain the transgene *nIs175[P$_{ceh-28}$::gfp]*.

Images are oriented such that left corresponds to anterior, top to dorsal.

1010

Figure 2

1011

Figure 2. *ctbp-1* mutant AIAs retain multiple aspects of their AIA gene

expression profile

(A-B) Expression of (A) M4 markers *egl-17*, *dbl-1*, *ser-7.b* and *flp-21* and (B) AIA

markers *gcy-28.d*, *ins-1*, *ttx-3*, *cho-1* and *mgl-1* in wild-type (left image) and *ctbp-*

1(n4784) (right image) L4 larval worms. Arrow, M4 neuron. Circles, AIAs. Scale

bar, 5 μm.

Images are oriented such that left corresponds to anterior, top to dorsal.

1019

Figure 3

1020

Figure 3. Loss of *ctbp-1* results in a progressive decline in AIA morphology

(A) Three representative images of an AIA neuron in wild-type (left) and *ctbp-1(n4784)* (right) worms at L1 (top), L4 (middle) and day 1 adult (bottom) stages. Arrows, examples of ectopic neurites protruding from the AIA cell body. Scale bar, 5 μm.

(B-C) Percentage of AIAs in wild-type and *ctbp-1* worms at the L1, L4 and day 1 adult stages with an ectopic neurite protruding from the (B) anterior or (C) posterior of the AIA cell body. Mean ± SEM. *n* = 60 AIAs scored per strain per

stage, 4 biological replicates. ns, not significant, **p<0.01, ****p<0.0001,

unpaired t-test.

(D) Quantification of AIA cell body length in wild-type and *ctbp-1* worms at the

L1, L4 and day 1 adult stages. Mean ± SEM. *n* = 30 AIAs scored per strain per

stage. ns, not significant, *p<0.05, ****p<0.0001, unpaired t-test.

(E) Three representative images of an AIA neuron in *ctbp-1; nIs743[P_{gcy-}*

_{28.d}::ctbp-1(+)] worms at L1 (top), L4 (middle) and day 1 adult (bottom) stages.

Arrows, examples of ectopic neurites protruding from the AIA cell body. Scale

bar, 5 μm.

(F-G) Percentage of AIAs in wild-type, *ctbp-1* and *ctbp-1; nIs743* worms at the

L1, L4 and day 1 adult stages with an ectopic neurite protruding from the (F)

anterior or (G) posterior of the AIA cell body. Mean ± SEM. *n* = 30 AIAs scored

per strain per stage, 3 biological replicates. ns, not significant, **p<0.01,

***p<0.001, ****p<0.0001, one-way ANOVA with Tukey's correction.

(H) Quantification of AIA cell body length in wild-type, *ctbp-1* and *ctbp-1; nIs743*

worms at the L1, L4 and day 1 adult stages. Mean ± SEM. *n* ≥ 30 AIAs scored

per strain per stage. ns, not significant, **p<0.01, ****p<0.0001, one-way ANOVA

with Tukey's correction.

(I) Three representative images of an AIA neuron in *ctbp-1; nEx2351[P_{hsp-}*

_{16.2}::ctbp-1(+);P_{hsp-16.41}::ctbp-1(+)] worms at L4 (top), day 1 adult with heat shock

(HS) (middle) and day 1 adult without heat shock (bottom). Arrows, examples of

ectopic neurites protruding from the AIA cell body. Scale bar, 5 μm.

(J-K) Percentage of AIAs in wild-type, *ctbp-1* and *ctbp-1; nEx2351* worms at L4

and day 1 adult (with or without heat shock) stages with an ectopic neurite

protruding from the (J) anterior or (K) posterior of the AIA cell body*.* Mean ±

SEM. *n* = 30 AIAs scored per strain per stage, 3 biological replicates. ns, not

significant, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, one-way ANOVA with

Tukey's correction.

(L) Quantification of AIA cell body length in wild-type, *ctbp-1* and *ctbp-1;*

nEx2351 worms at L4 and day 1 adult (with or without heat shock) stages*.* Mean

± SEM. *n* ≥ 30 AIAs scored per strain per stage. ns, not significant, *p<0.05,

****p<0.0001, one-way ANOVA with Tukey's correction.

The *ctbp-1* allele used for all panels of this figure was *n4784*.

All strains contain *nIs840[P$_{gcy-28.d}$::gfp],* and all strains other than "Wild type"

contain *nIs348[P$_{ceh-28}$::mCherry]* (not shown in images).

Images are oriented such that left corresponds to anterior, top to dorsal.

1065

Figure 4

Figure 4. Loss of *ctbp-1* results in a disruption of AIA function in L4-to-day

1 adult worms

(A) Schematic of the butanone adaptation assay. L1 or L4 worms from

synchronized populations were washed off plates with S Basal, washed with S

Basal, split into naïve and conditioned populations, incubated in S Basal with or

without 2-butanone for 1.5 hours, washed again with S Basal, allowed to

chemotax for two hours on unseeded plates containing two 1 μl spots of 10%

ethanol (blue dots) and 2-butanone diluted in 10% ethanol (orange dots), and

then scored.

(B-E) Chemotaxis indices of (B,D) naïve or (C,E) conditioned wild-type (N2 and

nIs175), AIA-ablated (JN580), *nIs175*; *ctbp-1(n4784)*, and *nIs175*; *ctbp-1* mutants

containing a transgene driving expression of wild-type *ctbp-1* under an AIA-

specific promoter (*nIs743[P$_{gcy-28.d}$::ctbp-1(+)]*) at the (B-C) L1 or (D-E) L4 larval

stage. Mean ± SEM. *n* ≥ 6 assays per condition, ≥ 50 worms per assay. ns, not

significant, **p<0.01, ***p<0.001, ****p<0.0001, one-way ANOVA with Tukey's

correction.

(F-G) Chemotaxis indices of (F) naïve or (G) conditioned *nIs175*, *nIs175*; *ctbp-1*,

and *nIs175*; *ctbp-1* mutants carrying the heat-shock-inducible transgene

nEx2351[P$_{hsp-16.2}$::ctbp-1(+);P$_{hsp-16.41}$::ctbp-1(+)] with or without heat shock (HS)

at the day 1 adult stage. Mean ± SEM. *n* ≥ 5 assays per condition, ≥ 50 worms

per assay. ns, not significant, **p<0.01, ****p<0.0001, one-way ANOVA with

Tukey's correction.

The *ctbp-1* allele used for all panels of this figure was *n4784*.

1090

Figure 5

Figure 5. Loss of *ctbp-1* results in a disruption to normal AIA gene expression

(A,C,E) (A) *nEx3081[P$_{acbp-6}$::gfp]*, (C) *otIs123[P$_{sra-11}$::gfp],* or (E) *ivEx138[P$_{glr-2}$::gfp]* expression in wild-type (top) and *ctbp-1(n4784)* (bottom) worms at the L1 larval stage (left) and L4 larval stage (right). Wild-type strains contain *nIs843[P$_{gcy-28.d}$::mCherry]. ctbp-1* mutant strains contain *nIs348[P$_{ceh-28}$::mCherry].*   Arrow, M4 neuron. Circle, AIAs. Scale bar, 10 µm.

(B,D,F) Percentage of wild-type and *ctbp-1(n4784)* expressing (B) *P$_{acbp-6}$::gfp*, (D) *P$_{sra-11}$::gfp,* or (F) *P$_{glr-2}$::gfp* in the AIA neurons at L1 and L4 larval stages. Wild-type strains contain *nIs843[P$_{gcy-28.d}$::mCherry]. ctbp-1* mutant strains contain *nIs348[P$_{ceh-28}$::mCherry].* Mean ± SEM. *n* ≥ 50 worms per strain per stage, 3 biological replicates. ns, not significant, ****p<0.0001, unpaired t-test.

Figure 6

Figure 6. A suppressor screen reveals *egl-13* as a *ctbp-1* genetic interactor

(A) Schematic of *ctbp-1* suppressor screen design. *ctbp-1* mutant worms carrying

nIs175[P_ceh-28_::gfp] were mutagenized with ethyl methanesulfonate (EMS), and

their F2 progeny were screened for continued *nls175* expression in M4 and loss

of expression in the AIA neurons (red circle).

(B) Gene diagram of the *egl-13a* isoform. Arrows (above), point mutations. Line

(below), deletion. Scale bar (bottom right), 1 kb.

(C) Percentage of wild-type, *ctbp-1* and *egl-13(n5937)* worms expressing *nls175*

in the AIA neurons over time. Time points correspond to the L1, L2, L3, L4 larval

stages, and day 1 adult worms (indicated below X axis). All strains contain

nls175[P$_{ceh-28}$::gfp]. Mean ± SEM. *n* ≥ 100 worms per strain per stage, 3

biological replicates.

(D) Three representative images of an AIA neuron in *egl-13 ctbp-1* worms at L1

(top), L4 (middle) and day 1 adult (bottom) stages. Arrows, examples of ectopic

neurites protruding from the AIA cell body. Image oriented such that left

corresponds to anterior, top to dorsal. Scale bar, 5 μm.

(E-F) Percentage of AIAs in wild-type, *ctbp-1* and *egl-13 ctbp-1* worms at the L1,

L4 and day 1 adult stages with an ectopic neurite protruding from the (E) anterior

or (F) posterior of the AIA cell body. Mean ± SEM. *n* = 30 AIAs scored per strain

per stage, 3 biological replicates. ns, not significant, *p<0.05, **p<0.01,

***p<0.001, ****p<0.0001, one-way ANOVA with Tukey's correction.

(G) Quantification of AIA cell body length in wild-type, *ctbp-1* and *egl-13 ctbp-1*

worms at the L1, L4 and day 1 adult stages. Mean ± SEM. *n* ≥ 30 AIAs scored

per strain per stage. ns, not significant, *p<0.05, ****p<0.0001, one-way ANOVA

with Tukey's correction.

(H-K) Chemotaxis indices of (H,J) naïve or (I,K) conditioned wild-type, *ctbp-1* and

egl-13 ctbp-1 worms at the (H-I) L1 or (J-K) L4 larval stage. Mean ± SEM. *n* ≥ 5

assays per condition, ≥ 50 worms per assay. ns, not significant, *p<0.05,

**p<0.01, ****p<0.0001, one-way ANOVA with Tukey's correction.

The *ctbp-1* allele used for all panels of this figure was *n4784*. The *egl-13* allele

used for all panels of this figure was *n5937*.

All strains in (D-G) contain *nIs840[P$_{gcy-28.d}$::gfp]* and all strains in (D-G) other than

"Wild type" contain *nIs348[P$_{ceh-28}$::mCherry]* (not shown in images).

All strains in (C, H-K) contain *nIs175[P$_{ceh-28}$::gfp].*

1140

Figure 7

Figure 7. EGL-13 disrupts AIA function partially through driving misexpression of *ceh-28* in *ctbp-1* mutants

(A-C) Expression of markers for AIA misexpressed genes (A) *nEx3081[P$_{acbp-}$*

$_6$::gfp], (B) *otIs123[P$_{sra-11}$::gfp]*, or (C) *ivEx138[P$_{glr-2}$::gfp]* in *egl-13(n5937) ctbp-*

1(n4784) double mutants at the (top) L1 and (bottom) L4 larval stages. Arrow, M4

neuron. Circle, AIAs. Scale bar, 10 μm.

(D) Percentage of *egl-13(n5937) ctbp-1(n4784)* double mutants expressing the

indicated reporter in the AIA neurons at the L1 and L4 larval stages. Mean ±

SEM. *n* ≥ 50 worms scored per strain, 3 biological replicates.

(E-H) Chemotaxis indices of (E,G) naïve or (F,H) conditioned wild-type (*nIs175*),

nIs175; ctbp-1(n4784), and *acbp-6(tm2995)*; *nIs175; ctbp-1* mutants at the (E-F)

L1 or (G-H) L4 larval stage. Mean ± SEM. *n* ≥ 6 assays per condition, ≥ 50

worms per assay. ns, not significant, *p<0.05, ***p<0.001, ****p<0.0001, one-way

ANOVA with Tukey's correction.

(I-L) Chemotaxis indices of (I,K) naïve or (J,L) conditioned wild-type (*nIs175*),

nIs175; ctbp-1(n4784), and *nIs175; ctbp-1 ceh-28(cu11)* mutants at the (I-J) L1

or (K-L) L4 larval stage. Mean ± SEM. *n* ≥ 6 assays per condition, ≥ 50 worms

per assay. ns, not significant, *p<0.05, ***p<0.001, one-way ANOVA with Tukey's

correction.

The *ctbp-1* allele used for all panels of this figure was *n4784.*

All strains in Fig. 7A-D contain *nIs348[P$_{ceh-28}$::mCherry]*.

Images are oriented such that left corresponds to anterior, top to dorsal.

1164

Figure 8

Figure 8. Model for the maintenance of the AIA cell identity by *ctbp-1*

(A) The genetic pathway in which *ctbp-1* promotes AIA morphology and *glr-2* and *sra-11* expression. *ctbp-1* also inhibits *egl-13*, thereby repressing expression of *ceh-28* and *acbp-6* in the AIAs and promoting proper AIA function

(B-D) Model for how CTBP-1 maintains the AIA cell identity.

(B) We propose that CTBP-1 acts in the maintenance but not establishment of

the AIA cell identity, and does so by targeting specific genetic loci for regulation

through physical interaction with transcription factors such as EGL-13.

(C) In the absence of CTBP-1, EGL-13 and other CTBP-1 targets drive

expression at multiple genetic loci, resulting in changes to the gene expression,

morphology and function (as assessed by butanone adaptation) of the AIAs.

(D) When EGL-13 activity is also removed, gene expression and cellular function

are no longer perturbed, while normal morphology is not restored, resulting in a

"Partially Maintained AIA."

1180

Figure S1

A

Allele	Nucleotide Change	Amino Acid Change	Codon Position
n4778	83 bp deletion, frameshift	-	389-417
n4784	CGA to TGA	R ➔ Opal	27
n4789	AGg to AGa	splice site donor	370
n4800	706 bp deletion, frameshift	-	166-216
n4804	GCA to GTA	A ➔ V	261
n4805	CCA to TCA	P ➔ S	29
n4808	CAC to TAC	H ➔ Y	60
n4810	CGA to TGA	R ➔ Opal	55
n4813	TCA to TAA	S ➔ Ochre	113
n4819	CAG to TAG	Q ➔ Amber	208
n4823	AGg to AGa	splice site donor	486
n4824	CAA to TAA	Q ➔ Ochre	100
n4840	TGG to TGA	W ➔ Opal	470
n4849	CGA to TGA	R ➔ Opal	324
n4852	GGg to GGa	splice site donor	265
n4861	GAA to TAA	E ➔ Ochre	220
n4864	CAA to TAA	Q ➔ Ochre	312
n4865	GGA to AGA	G ➔ R	332

B

C

D

nIs175; ctbp-1(tm5512)

Stage	% AIA expressing P_{ceh-28}::gfp	n
L1	1	100
L4	98	100

E

Figure S1. Additional *ctbp-1* mutant alleles cause misexpression of P_{ceh-}

$_{28}$::*gfp* in the AIA neurons

(A) Table of *ctbp-1* mutant alleles we showed to result in *nIs175[P_{ceh-28}::gfp]* or

nIs177[P_{ceh-28}::gfp] misexpression in the AIA neurons. Specific nucleotide

changes are denoted in red. Codon positions correspond to the *ctbp-1a* isoform.

(B) Gene diagram of the *ctbp-1a* isoform showing all 18 *ctbp-1* alleles isolated in

this study. Arrows, point mutations. Lines, deletions. Scale bar (bottom right), 1

kb.

(C) Expression of *nIs175* in *ctbp-1(tm5512)* L1 and L4 mutant worms. Arrow, M4

neuron. Circle, AIAs. Scale bar, 10 µm.

(D) Percentage of *ctbp-1(tm5512)* worms expressing *nIs175* in the AIA neurons

at the L1 and L4 larval stages.

(E) *nIs348[P_{ceh-28}::mCherry]* expression in wild-type (top) and *ctbp-1(n4784)*

(bottom) worms at the L1 larval stage (left) and L4 larval stage (right). Arrow, M4

neuron. Circle, AIAs. Scale bar, 10 µm.

All strains in Fig. S1A-C contain the either *nIs175[P_{ceh-28}::gfp]* or *nIs177[P_{ceh-}*

$_{28}$::gfp].

Images are oriented such that left corresponds to anterior, top to dorsal.

1200

Figure S2

A

Genotype	% AIA expressing $P_{ceh\text{-}28}$::gfp	n
Wild type	0	100
ctbp-1(n4784)	98	100
ctbp-1(n4808)	98	100
ctbp-1(n4784); nEx2346[ctbp-1(+)]	12	100
ctbp-1(n4784); nEx2347[ctbp-1(+)]	4	100
ctbp-1(n4784); nIs743[P_{AIA}::ctbp-1(+)]	0	100

B

ctbp-1(n4784); nEx2351[P_{hsp}::ctbp-1(+)]

Stage	% AIA expressing $P_{ceh\text{-}28}$::gfp	n
L1	8	100
L4	99	100
Day 1 Adult (- HS)	100	100
Day 1 Adult (+ HS)	8	100

1201

Figure S2. Quantification of *ctbp-1* strains misexpressing $P_{ceh\text{-}28}$::gfp

(A) Percentage of worms of indicated genotypes expressing *nIs175[$P_{ceh\text{-}28}$::gfp]*

in the AIA neurons at the L4 larval stage.

(B) Percentage of *ctbp-1(n4784); nEx2351[$P_{hsp\text{-}16.2}$::ctbp-1(+);$P_{hsp\text{-}16.41}$::ctbp-1(+)]*

worms expressing *nIs175* at the L1, L4 and day 1 adult stages. Day 1 adults

shown ± heat shock (HS) at the L4 stage.

1208

Figure S3

A

M4 markers

Reporter	Wild type			ctbp-1(n4784)		
	% expressing in M4	% expressing in AIAs	*n*	% expressing in M4	% expressing in AIAs	*n*
ayIs4	100	0	50	100	0	50
ctIs43	100	0	50	100	0	50
nIs491	100	0	50	100	0	50
ynIs80	100	0	50	100	0	50

B

AIA markers

Reporter	Wild type			ctbp-1(n4784)		
	% expressing in M4	% expressing in AIAs	*n*	% expressing in M4	% expressing in AIAs	*n*
nIs843	0	100	50	0	100	50
otIs326	0	100	50	0	100	50
peIs1716	0	100	50	0	100	50
otIs379	0	100	50	0	100	50
otIs317	0	100	50	0	100	50

1209

1210 **Figure S3. Quantification of M4 and AIA marker expression**

1211 (A-B) Quantification of wild-type and *ctbp-1(n4784)* L4 worms expressing the

1212 indicated (A) M4 or (B) AIA markers from Fig. 2A-B in the M4 and AIA neurons.

1213

Figure S4

1214

Figure S4. Loss of *ctbp-1* results in a disruption of AIA morphology but not

AIA size

(A) Quantification of maximum AIA cell body area in wild-type and *ctbp-1(n4784)*

worms at L1, L4 and day 1 adult stages*.* Both strains contain *nIs840[P$_{gcy-28.d}$::gfp]*

and the *ctbp-1* strain contains *nIs348[P$_{ceh-28}$::mCherry]*. Mean ± SEM. *n* ≥ 30

AIAs scored per strain per stage. ns, not significant, unpaired t-test.

(B) Scoring of wild-type and *ctbp-1(n4784)* AIA images at the L1, L4 and day 1

adult stages. A random subset of AIA images used for length measurements in

Fig. 3D, 3H and 3L were blinded and scored as having either "Normal" or

"Elongated" AIA cell bodies. *n* ≥ 20 AIAs scored per strain per stage, 3 replicates.

ns, not significant, *p<0.05, **p<0.01, unpaired t-test.

1226

    Figure S5

1227

1228   **Figure S5. *ctbp-1* mutants display a non-AIA-dependent chemotaxis defect**

1229   (A-B) Chemotaxis indices of wild-type (N2), AIA-ablated (JN580) and *nIs175*;

1230   *ctbp-1(n4784)* mutants at the L4 larval stage to (A) diacetyl or (B) isoamyl alcohol

1231   diluted in pure ethanol. Mean ± SEM. *n* ≥ 3 assays per condition, ≥ 40 worms per

1232   assay. ns, not significant, *p<0.05, one-way ANOVA with Tukey's correction.

1233

Figure S6

Average Expression (A.U.)		
gene	Wild-type	*ctbp-1*
acbp-6	0.572	13.667
ceh-28	0	4.349
glr-2	0.208	0
sra-11	0.052	0

1234

1235   **Figure S6. Gene expression in wild-type and *ctbp-1* AIAs**

1236   Average expression level of confirmed scRNA-Seq hits in the AIAs of L4 wild-

1237   type and *ctbp-1* mutant animals. A.U., arbitrary expression units.

1238

Figure S7

A

Allele	Nucleotide Change	Amino Acid Change	Codon Position
n5937	gAT to aAT	splice site acceptor	158
n6013	CAA to TAA	Q → Ochre	381
n6313	436 bp deletion	-	205-334

Figure S7. Characterization of *egl-13* alleles isolated as *ctbp-1* suppressors

(A) Table of *egl-13* mutant alleles isolated in this study as suppressors of *ctbp-1*-mediated *nIs175[P_{ceh-28}::gfp]* misexpression in the AIA neurons. Specific nucleotide changes are denoted in red. Codon positions correspond to *egl-13a* isoform.

(B) Percentage of wild-type, *ctbp-1(n4784)* and *egl-13 ctbp-1* worms expressing *nIs175* in the AIA neurons at the L4 larval stage. Mean ± SEM. *n* ≥ 100 worms scored per strain, 3 biological replicates. ****p<0.0001, one-way ANOVA with Tukey's correction.

(C) Percentage of *ctbp-1(n4784), egl-13 ctbp-1* and *egl-13 ctbp-1* worms carrying

transgenic constructs expressing wild-type *egl-13* under its native promoter

expressing *nIs175* in the AIA neurons at the L4 larval stage. Mean ± SEM. *n* ≥ 50

worms scored per strain, 3 biological replicates. ****p<0.0001, one-way ANOVA

with Tukey's correction.

All strains in Fig. S7B-C contain *nIs175[P$_{ceh-28}$::gfp]*.

1255

Figure S8

1256

Figure S8. EGL-13 functions cell-autonomously to regulate AIA gene

expression

(A) Expression of *nIs175[P$_{ceh-28}$::gfp]* in *egl-13(n5937) ctbp-1(n4784)* (left panel),

and *egl-13 ctbp-1* mutants carrying an extrachromosomal array expressing wild-

type *egl-13* under the AIA-specific promoter *gcy-28.d* (*nEx3055*, right panel) in L4

worms. Arrow, M4 neuron. Circle, AIAs. Scale bar, 10 µm.

(B) Percentage of *ctbp-1(n4784)*, *egl-13(n5937) ctbp-1* and *egl-13 ctbp-1;*

nEx3055 worms expressing *nIs175* in the AIA neurons at the L4 larval stage. All

strains contain *nIs175[P$_{ceh-28}$::gfp].* Mean ± SEM. *n* = 100 worms scored per

strain, 3 biological replicates. ***p<0.001, ****p<0.0001, one-way ANOVA with

Tukey's correction.

The alleles used for all panels of this figure were *ctbp-1(n4784)* and *egl-*

13(n5937).

All strains contain *nIs175[P$_{ceh-28}$::gfp].*

Images are oriented such that left corresponds to anterior, top to dorsal.