CRISPR-Cas Controls Cryptic Prophages

Sooyeon Song[1,2,3*], Ekaterina Semenova[4], Konstantin Severinov[4], Laura Fernández-García[1], Michael J. Benedik[5], Toshinari Maeda[6], and Thomas K. Wood[1*]

[1]Department of Chemical Engineering, Pennsylvania State University, University Park, Pennsylvania, 16802-4400, USA

Departments of Animal Science[2] and Agricultural Convergence Technology[3], Jeonbuk National University, 587 Baekje-Daero, Deojin-Gu, Jeonju-Si, Jellabuk-Do, 54896, South Korea

[4]Waksman Institute of Microbiology, Rutgers, The State University of New Jersey, Piscataway, NJ 08854

[5]Hamad bin Khalifa University, Education City, Doha, Qatar

[6]Department of Biological Functions Engineering, Kyushu Institute of Technology, Kitakyushu, Japan 808-0196

*For correspondence. E-mail songsy@jbnu.ac.kr, tuw14@psu.edu

Tel. (+)1 814-863-4811; Fax (1) 814-865-7846

Running title: **CRISPR-Cas tames cryptic prophages**.

Keywords: CRISPR-Cas, persisters, cryptic prophage

SUMMARY

The bacterial archetypal adaptive immune system, CRISPR-Cas, is thought to be repressed in the best-studied bacterium, *Escherichia coli* K-12. We show here that the *E. coli* CRISPR-Cas system is active and serves to inhibit its nine defective (i.e., cryptic) prophages. Specifically, compared to the wild-type strain, reducing the amounts of specific interfering RNAs (crRNA) decreases growth by 40%, increases cell death by 700%, and prevents persister cell resuscitation. Similar results were obtained by inactivating CRISPR-Cas by deleting the entire 13 spacer region (CRISPR array); hence, CRISPR-Cas serves to inhibit the remaining deleterious effects of these cryptic prophages, most likely through CRISPR array-derived crRNA binding to cryptic prophage mRNA, rather than through cleavage of cryptic prophage DNA; i.e., self-targeting. Consistently, four of the 13 *E. coli* spacers contain complementary regions to the mRNA sequences of six cryptic prophages, and inactivation of CRISPR-Cas increases the level of mRNA for lysis protein YdfD of cryptic prophage Qin and lysis protein RzoD of cryptic prophage DLP-12. Also, lysis is clearly seen via transmission electron microscopy when the whole CRISPR-Cas array is deleted, and eliminating spacer #12, which encodes crRNA with complementary regions for Rac, Qin, and CP4-57 cryptic prophages, also results in growth inhibition and cell lysis. Therefore, we report the novel results that (i) CRISPR-Cas is active in *E. coli* and (ii) CRISPR-Cas is used to tame cryptic prophages, likely through RNAi; i.e., unlike with active lysogens, active CRISPR-Cas and cryptic prophages may stably co-exist.

INTRODUCTION

Along with restriction/modification (Vasu and Nagaraja, 2013) and toxin/antitoxin (TA) systems (Pecota and Wood, 1996), prokaryotes utilize clustered, regularly-interspaced, short palindromic repeats (CRISPR) and CRISPR-associated (Cas) (Makarova et al., 2020) proteins to combat phages. These systems are interrelated in that some Cas proteins and TA systems have a common ancestor; for example, *Sulfolobus solfataricus* Cas2 is structurally similar to antitoxin GhoS (an RNase) of the *Escherichia coli* GhoT/GhoS TA system (Wang et al., 2012). In addition, in a manner similar to our discovery (Pecota and Wood, 1996) that toxins of host TA systems inhibit phage by degrading mRNA when the phage shuts down transcription (e.g., Hok/Sok inhibits T4 phage), some Cas proteins induce host dormancy rather than degrading phage DNA to inhibit phage propagation (Meeske et al., 2019). Also, TA systems have been found to stabilize CRISPR-Cas systems by making them addictive to the host (Li et al., 2021).

Although CRISPR-Cas systems exclude both external lytic and temperate (lysogenic) phages (Goldberg et al., 2018), CRISPR-Cas systems of lysogens that target their own integrated prophages decrease long-term fitness and either the cell dies or the prophage is lost (Edgar and Qimron, 2010; Goldberg *et al.*, 2018). Also, the class I-E (Makarova *et al.*, 2020) CRISPR–Cas system of *E. coli* is not related to immunity for external phages (Touchon et al., 2011) and appears to be inactive in the wild-type strain at standard laboratory conditions (Shmakov et al., 2017), due to repression by H-NS, although it is functional when induced (Pougach et al., 2010a). To date, the relationship of CRISPR-Cas to cryptic prophages; i.e., those phage remnants that are unable to form lytic particles, has not been investigated.

Up to 50% of bacterial genomes may contain stably-integrated phage DNA (Howard-Varona et al., 2017), and for *E. coli,* we discovered that its nine cryptic prophages are not extraneous DNA but instead encode genes for proteins that increase resistance to sub-lethal concentrations of quinolone and β-lactam antibiotics as well as protect the cell from osmotic, oxidative, and acid stresses (by deleting 166 kb) (Wang et al., 2010). Although these cryptic prophages do not enhance the formation of persister cells, a subpopulation of cells that weather extreme stress by entering into a dormant state (Wood and Song, 2020), these phage remnants facilitate the resuscitation of persisters via nutrient sensing (Song et al., 2021).

Therefore, the bacterial cell can co-opt the genome of its former parasite to both combat stress (Wang *et al.*, 2010) as well as to revive from dormancy (Song *et al.*, 2021).

Given these active roles of cryptic prophages in the stress response (Wang *et al.*, 2010) and persister cell resuscitation (Song *et al.*, 2021), we hypothesized here that the native *E. coli* CRISPR-Cas system plays an active role in the regulation of cryptic prophages. We find that the CRISPR-Cas system is required for inhibiting the expression of deleterious cryptic prophage genes, since, if CRISPR-Cas is inactivated by preventing crRNA production, cells die due to activation of the cryptic prophage lysis proteins YdfD of Qin and RzoD from DLP-12. Hence, we discovered CRISPR-Cas is active in *E. coli* and serves to regulate its former phage foe.

RESULTS

Deletions of CRISPR-Cas components reduce growth. We assayed the importance of CRISPR-Cas in *E. coli* K-12 by testing the individual deletions of the CRISPR-Cas genes on growth in M9 glucose medium (*cas1, cas2, cas3, casA, casB, casC, casD,* and *casE*) and found only the *cas2* deletion, with its kanamycin cassette insertion, had an effect: *cas2* causes 40% slower growth in rich medium compared to the wild-type strain (specific growth rate of 0.79 ± 0.21/h vs. 1.3 ± 0.11/h, respectively). Similarly, in minimal glucose (0.4 wt%) medium, deletion of *cas2* also reduces growth by 33% (0.42 ± 0.02/h vs. 0.62 ± 0.02/h, respectively) and reduced the yield in stationary phase (**Fig. S1A**). We hypothesized that the *cas2* deletion/kanamycin insertion caused a polar effect by reducing spacer production given it is directly upstream of the spacer region (Pougach *et al.*, 2010a) (**Fig. S2**). Fittingly, inactivation of CRISPR-Cas by eliminating the CRISPR arrays in the isogenic host BW25113 (henceforth Δspacer) also reduced growth by 32% in rich medium and eliminating kanamycin resistance in the *cas2* deletion strain to prevent the polar mutation affecting the spacer region restored nearly wild-type growth (**Fig. S1A**) as well as reduced toxicity (**Fig. S1B**) relative to the *cas2* Ωkan^R strain. Therefore, the *E. coli* CRISPR-Cas represses some processes that inhibit growth via crRNA, and by this criterion, is active, providing a clear advantage to the host.

CRISPR-Cas increases single-cell resuscitation. Since deletions of CRISPR-Cas components decrease growth, we tested for their effect on persister cell resuscitation using single-cell microscopy. Persister cell resuscitation is germane in that the dormant cells are highly stressed and have limited resources for their revival via activation of hibernating ribosomes (Kim et al., 2018; Song and Wood, 2020; Yamasaki et al., 2020); for example, we have shown that inhibiting ATP synthesis leads to a 5,000-fold increase in persister cell formation (Kwan et al., 2013). Since we discovered facile means for converting the whole population of cells into persister cells (Kim *et al.*, 2018; Kwan *et al.*, 2013; Yamasaki *et al.*, 2020) that has been used by at least 17 independent labs to date with various bacterial species (Song and Wood, 2021), these stressed cells are an excellent model for testing the effects of CRISPR-Cas on *E. coli* physiology.

Here, we found that deletion of *cas2* reduces persister cell resuscitation by 31-fold (**Fig. 1A, Table S1**). Persister cell resuscitation was similarly affected (15-fold reduction) in the Δspacer strain (**Fig. 1A, Table S1**). Since the *cas2* mutant grows more slowly than the wild-type in rich medium, as a control, we tested an *E. coli* mutant that grows 22% slower than the wild-type, *ssrA*, to confirm that the slower growth does not affect persister resuscitation and found that the *ssrA* mutant resuscitates at nearly the identical rate as the wild-type strain (**Fig. S3**). Hence, CRISPR-Cas is active in *E. coli* and plays key roles in its growth and recovery from extreme stress.

CRISPR-Cas prevents cell death by preventing cell lysis. To explore how CRISPR-Cas increases growth (including persister cell resuscitation), we checked for death among resuscitating persister cells of the *cas2* deletion strain using the Live/Dead stain. We found that inactivating CRISPR-Cas leads to a 7-fold increase in death among resuscitating cells (**Fig. 1B, Table S2**). In addition, there were 34-fold more cells termed "ghosts" (Cheng et al., 2014) that lack cytosolic material (**Fig. S4A**), are likely dead, and have intact membranes, so these cells are not stained by the propidium iodide dye. Corroborating these results, there was 11-fold and 5-fold more death for stationary- and exponential-phase cells, respectively, when CRISPR-Cas was inactivated via *cas2* (**Fig. S5**, **Table S2**). Moreover, for the Δspacer strain, there was 81-fold more cell death in the for stationary-phase cells (**Fig. S5**) and 7-fold more death for resuscitating persister cells

(**Fig. 1B, Table S2**). Critically, for the Δspacer strain, waking persister cells clearly show lysis (**Fig. S4A**), and transmission electron microscopy images confirm this lysis with cytosolic materials seen next to lysed cells (**Fig. S4B**). These results indicate inactivating CRISPR-Cas leads to cell death and that the mechanism for cell death is via lysis.

CRISPR-Cas reduces cryptic prophage lysis gene mRNA. We hypothesized that since CRISPR-Cas systems inhibit some phage before they become lysogens (Edgar and Qimron, 2010; Goldberg *et al.*, 2018), the *E. coli* system may be preventing cell death by repressing certain cryptic prophage genes. To test this hypothesis, we first examined the *E. coli* CRISPR-Cas system for spacers related to the nine cryptic prophages. *E. coli* K-12 CRISPR array located next to the *cas* operon contains 13 spacers (Militello and Lazatin, 2017; Pougach *et al.*, 2010a), each containing 32 or 33 nt (Brouns et al., 2008). Between the 14 29-nt repeat sequences (5'-GTGTTCCCCGCATCAGCGGGGATAAACCG), we found that four of the 13 spacers contain 11 to 16 nt of perfect matches to six of the nine cryptic prophages (CPS-53, CP4-6, Rac, Qin, CP4-57, and e14) (**Fig. 2A**) based on binding to cryptic prophage mRNA (**Fig. S6**). Critically, spacer 12 encodes crRNA with the potential to bind mRNA from Rac (*stfR*), Qin (*stfQ*), and CP4-57 (*alpA*) (**Fig. S6**). In general, spacer lengths vary from 21 to 72 nt (Barrangou and Marraffini, 2014) with perfect complementarity of 6 to 12 nt (Shmakov *et al.*, 2017). Together, these results suggest CRISPR-Cas potentially regulates the *E. coli* cryptic prophages. Note that the second CRISPR array in *E. coli* K-12 in between genes *ygcE* and *queE* lacks spacers with sequences that matched the cryptic prophages.

The presence of the cryptic prophage-related targeting spacers suggests that *E. coli* CRISPR-Cas may be preventing expression of cryptic prophage genes and, given the cell phenotype, is suggestive of prophage-encoded lethal genes; hence, we checked the mRNA levels of the cryptic prophage lysis genes encoded by these seven cryptic prophages with spacer matches. Specifically, transcription of *ydfD* (Qin), *hokD* (Qin), *ypjF* (CP4-57), *essD* (DLP-12), and *rzoD* (DLP-12) was checked via quantitative reverse transcription polymerase chain reaction (qRT-PCR). We found that inactivation of CRISPR-Cas via the *cas2* deletion results in a 35-fold increase in *ydfD* mRNA and a 39-fold increase in *rzoD* mRNA in resuscitating persister cells (**Fig. 1C**). RzoD is a putative DLP-12 lysis lipoprotein that we previously

120   showed was toxic through its interaction with toxin Hha (García Contreras et al., 2008) of the Hha/TomB

TA system (Marimon et al., 2016). YdfD of Qin cryptic prophage has been shown to lyse cells when

induced (Masuda et al., 2016). To investigate if YdfD and RzoD are toxic, we induced production of both

proteins and found they are toxic (**Fig. S7**). Moreover, deleting *cas2* to inactivate CRISPR-Cas in the same

host that lacks all nine cryptic prophages (delta9) (Wang *et al.*, 2010) has no effect on growth (**Fig. S8**) and

125   does not cause lysis (**Fig. S5**). Hence, these results show CRISPR-Cas represses at least two *E. coli* cryptic

prophage proteins RzoD and YdfD that can reduce cell growth.

Excision of DLP-12 is not regulated by CRISPR-Cas. Since inactivation of CRISPR-Cas leads to

derepression of the DLP-12 *rzoD* lysis gene, we checked for increased DLP-12 excision with the *cas2*

deletion strain using qPCR. We found the *cas2* deletion has little impact on DLP-12 excision (**Table S3**).

130   Since the DLP-12 toxin gene *essD* is not derepressed, the effect is not due to excision of the prophage. We

also tested the effect of CRISPR-Cas on two other cryptic prophages with significant excision (CP4-57 and

e14) (Wang *et al.*, 2010), and also found no effect of deleting *cas2* (**Table S3**). Hence, CRISPR-Cas does

not affect cryptic prophage excision.

CRISPR-Cas regulates lytic gene mRNA levels. Since CRISPR-Cas does not affect cryptic prophage

135   excision and is unlikely to degrade cryptic prophage DNA as this would be lethal (Edgar and Qimron, 2010;

Goldberg *et al.*, 2018), we reasoned that CRISPR-Cas, via crRNA, must be interfering with expression of

the mRNA of the lytic genes to prevent production of the lytic proteins. To test this hypothesis, we

investigated whether *rzoD* and *ydfD* transcript levels are increased in the *cas2* and Δspacer strain relative

to the wild-type after inhibiting transcription via rifampicin. If CRISPR-Cas prevents lytic gene expression

140   by inhibiting lytic transcripts via crRNA, then inactivating CRISPR-Cas should lead to increased *rzoD* and

ydfD mRNA. In accordance with this hypothesis, *rzoD* mRNA was increased 320% in the Δspacer mutant

and 50% in the *cas2* mutant. Moreover, as expected, inactivating CasE increased cell lysis 49-fold

compared to the wild-type strain (**Fig. S5, Table S2)**, likely by preventing CasE from properly forming

crRNA from pre-crRNA; hence, there is less crRNA to inhibit the cryptic prophage mRNA from toxic

145   genes. Similarly, preventing spacer RNA formation by inactivating Cas2 led to a 10-fold increase in CRISPR array RNA levels in resuscitating persister cells as assayed by qRT-PCR, likely due to a lack of processing of the CRISPR array into crRNA (**Table S3B**). Finally, deleting only spacer #12 (which encodes crRNA targeting three separate regions in the cryptic prophage mRNA of CP4-57, Rac, and Qin, **Fig. S6**) in a markerless fashion inhibits cell growth in the stationary phase (**Fig. S8**) due to cell lysis (**Fig. S5, Table**

150   **S2**) to roughly the same extent as deleting all 13 spacers. In contrast, deleting spacer #3, which does not encode crRNA with matches to cryptic prophage mRNA, has no effect on growth (**Fig. S8**). Therefore, CRISPR-Cas prevents cryptic prophage lytic gene expression by interfering with the transcripts from these genes (**Fig. 2B**), specifically through spacer #12.

DISCUSSION

155   Our results reveal a new (non-canonical) role for CRISPR-Cas systems: regulation of phage fossils. The evidence for this includes that inactivating CRISPR-Cas by deleting *cas2* (i) reduces growth by 40%, (ii) nearly eliminates resuscitation from the persister state (**Fig. 1**), (iii) causes ghost cell formation and cell death (**Fig. S4**), (iv) increases mRNA levels of the cryptic prophage lysis genes *ydfD* and *rzoD,* and (v) increases CRISPR array levels by preventing processing into crRNA. Corroborating these results, the *cas2*

160   deletion has no effect in a strain which lacks cryptic prophage genes and eliminating the CRISPR array with prophage-matching spacers in the wild-type strain also reduced the growth rate, eliminated persister cell resuscitation, resulted in cell death via lysis, and activated *rzoD*. Furthermore, deleting only prophage-matching spacer #12 similarly inhibited growth and caused cell lysis. Given that Cas2 is not required for CRISPR interference, it is likely that the phenotypes produced upon deleting *cas2* are the result of a polar

165   mutation (**Fig. S2**) that inactivates CRISPR-Cas by altering production/processing of the spacers.

        Since there is no change in excision of DLP-12 upon inactivating CRISPR-Cas, our results suggest the mechanism for regulating the lysis genes *ydfD* and *rzoD* of the cryptic prophages is via CRISPR-Cas RNA binding to the cryptic prophage mRNA (analogous to RNA interference as seen previously in procaryotic systems with CRISPR-Cas (Mohanraju et al., 2022)) rather than cleaving excised DNA or directly cleaving

170 the mRNA. This mechanism is predicted based on bioinformatics analysis of 230 phages, that CRISPR-Cas functions in *E. coli* via endogenous gene expression rather than acting as an immune system (Bozic et al., 2019). In support of this, the excision rates of the cryptic prophages are so low that there may be insufficient excised DNA to cleave (1 excision in 10^6 cells for CP4-57) (Wang *et al.*, 2010), and a class I-F CRISPR-Cas system in *P. aeruginosa* uses crRNA to regulate endogenous LasR mRNA to reduce the

175 pro-inflammatory response (Li et al., 2016). Moreover, *E. coli* crRNA have been shown to be activated during infection in mice (Li *et al.*, 2016), and the *Salmonella* spp. system is identical to the *E. coli* one (e.g., the repeats are the same), but the spacers are different, which suggests it, too, is for endogenous gene expression. Critically, it has been demonstrated *in vitro* that that the *E. coli* Cascade binds ssRNA complementary to crRNA (Jore et al., 2011). Therefore, there is ample precedence for our proposed RNA

180 interference mechanism.

This new function for CRISPR-Cas acting on cryptic prophages is likely general and may explain why many species appear to have inactive CRISPR-Cas systems as was previously thought for *E. coli* (Shmakov *et al.*, 2017); i.e., instead of protecting cells from external phages, CRISPR-Cas systems may also control resident cryptic prophages which are prevalent. As additional evidence, we found matches in CRISPR-Cas

185 spacers not only for *E. coli* K-12 (**Fig. 2A**) but also for *E. coli* O157:H7, *Salmonella* spp., and *K. pneumoniae* (**Fig. S9**). Critically, our results provide the first example where it is beneficial for the host to have an active CRISPR-Cas system that targets *inactive* integrated phages (i.e., cryptic prophages) since previous reports show targeting *active* temperate phages is deleterious; i.e., either the cell dies or the phage is lost (Edgar and Qimron, 2010; Goldberg *et al.*, 2018; Rollie et al., 2020). Since *E. coli* cryptic prophages

190 like *rac* have been present in its genome for 4.5 million years (Perna et al., 2001), the active *E. coli* K-12 CRISPR-Cas system is stable with the cryptic prophages; in fact, there has been little change in the *E. coli* spacers for at least 42,000 years (Savitskaya et al., 2017).

Our results also indicate that, although the cryptic prophages are stable and the cell makes use of the genetic tools encoded by its former foe to combat myriad stresses (Wang *et al.*, 2010) and to sense nutrients

195 prior to exiting the persister state (Song *et al.*, 2021), the source of these tools must be elegantly regulated

by CRISPR-Cas since they often harbor deleterious membrane lysis proteins like YdfD and RzoD. Similarly, host Rho has been found recently to silence cryptic prophage toxin/antitoxin systems through transcription termination (Hafeezunnisa et al., 2021), and H-NS silences cryptic prophages through 65 binding sites (Ishihama and Shimada, 2021). Therefore, phages may be captured by the host, but they must be tamed, and this now includes repression by CRISPR-Cas. These results also suggest a role for CRISPR-Cas in gene regulation beyond self-defense.

MATERIALS AND METHODS

Bacteria and growth conditions. Bacteria (**Table S4**) were cultured routinely in lysogeny broth (Bertani, 1951) at 37°C, while M9 glucose (0.4 wt%) (Rodriguez and Tait, 1983) was used to assay the growth rate and to resuscitate persister cells. pCA24N-based plasmids (Kitagawa et al., 2005b) were retained in overnight cultures via chloramphenicol (30 μg/mL), and kanamycin (50 μg/mL) was used for deletion mutants, where applicable. The BW25113 *cas2* and spacer deletion were confirmed by PCR (primers shown in **Table S5**). Spacers 3 and 12 were deleted from BW25113, and *cas2* was deleted from the strain that lacks cryptic prophages (Δ9) using the method of Datsenko and Wanner (Datsenko and Wanner, 2000) using plasmids pKD4 with the primers in **Table S5**; kanamycin resistance was removed using pCP20 (Datsenko and Wanner, 2000).

Persister cells. Exponentially-growing cells (turbidity of 0.8 at 600 nm) were converted nearly completely to persister cells (Kim *et al.*, 2018; Kwan *et al.*, 2013) by adding rifampicin (100 μg/mL) for 30 min to stop transcription, centrifuging, and adding LB with ampicillin (100 μg/mL) for 3 h to lyse non-persister cells. To remove ampicillin, cells were washed twice with 0.85% NaCl then re-suspended in 0.85% NaCl. Persister concentrations were enumerated via a drop assay (Donegan et al., 1991).

Single-cell persister resuscitation. Persister cells (5 μL) were added to 1.5% agarose gel pads containing M9 glucose (0.4 wt%) medium (Rodriguez and Tait, 1983), and single-cell resuscitation was visualized at 37°C via a light microscope (Zeiss Axio Scope.A1, bl_ph channel at 1000 ms exposure). For each condition, at least two independent cultures were used with 150 to 300 individual cells assessed per culture.

Membrane integrity assay. To determine membrane integrity, the persister cells were analyzed with the LIVE/DEAD BacLight Bacterial Viability Kit (Molecular Probes, Inc., Eugene, OR, catalog number L7012). The fluorescence signal was analyzed via a Zeiss Axioscope.A1 using excitation at 485 nm and emission at 530 nm for green fluorescence and using excitation at 485 nm and emission at 630 nm for red fluorescence.

qRT-PCR. To quantify transcription from the cryptic prophage lytic genes, RNA was isolated from persister cells that were resuscitated by adding M9 glucose (0.4 wt%) medium for 10 min and from exponential cells grown to a turbidity of 0.8. For quantifying transcription from the CRISPR array, RNA was isolated from persister cells that were resuscitated by adding M9 glucose (0.4 wt%) medium for 10 min. Samples were cooled rapidly using ethanol/dry ice in the presence of RNA Later. RNA was isolated using the High Pure RNA Isolation Kit (Roche). The following qRT-PCR thermocycling protocol was used with the iTaq™ universal SYBR® Green One-Step kit (Bio-Rad): 95 ºC for 5 min; 40 cycles of 95 ºC for 15 s, 60 ºC for 1 min for two replicate reactions for each sample/primer pair. The annealing temperature was 60ºC for all primers (**Table S5**).

qPCR. To quantify prophage excision and the levels of DNA flanking the CRISPR-Cas cleavage sites, total DNA (100 ng) was isolated from exponentially-growing and persister resuscitating cells using an UltraClean Microbial DNA Isolation Kit (Mo Bio Laboratories). Excised cryptic prophage was quantified using primers for each prophage excisionase (**Table S5**) that only yield a PCR product upon prophage excision, and the relative amount of each target gene was determined using reference gene *purM*. The level of cryptic prophage flanking the CRISPR-Cas cleave site was quantified using primers that flank each site (**Table S5**). The qPCR reaction performed using CFX96 Real Time System. The reaction and analysis was conducted using the StepOne Real-Time PCR System (Bio-Rad).

Transmission electron microscopy. For transmission electron microscopy (TEM), the samples were prepared from persister cells that were resuscitated by adding M9 glucose (0.4%) medium for 10 min then washed with 0.85% NaCl. The samples were fixed with buffer (2.5% glutaraldehyde in 0.1M cacodylate buffer, pH 7.4). The negative staining was performed with 2% uranyl acetate for 1 hr, then dehydrated. The

sectioned specimens were stained again with uranyl acetate and lead citrate after dehydration and embedded

with resin. The TEM image was observed using a Hitachi (H-7605) instrument.

ACKNOWLEDGEMENTS

250   This work was supported by funds derived from the Biotechnology Endowed Professorship at the

Pennsylvania State University for TKW and from the National Research Foundation of Korea (NRF) grant

from the Korean Government (NRF-2020R1F1A1072397) for SYS. We appreciate the feedback of Dr.

Joseph Wade on the PAM sequences and of Joy Muthami. The authors have no conflicts of interest.

REFERENCES

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K.A., Tomita, M., Wanner, B.L., and Mori, H. (2006). Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol. Syst. Biol. *2*, 2006 0008.

Barrangou, R., and Marraffini, Luciano A. (2014). CRISPR-Cas Systems: Prokaryotes Upgrade to Adaptive Immunity. Molecular Cell *54*, 234-244.

Bertani, G. (1951). Studies on Lysogenesis .1. The Mode of Phage Liberation by Lysogenic *Escherichia-Coli*. J Bacteriol *62*, 293-300.

Bozic, B., Repac, J., and Djordjevic, M. (2019). Endogenous Gene Regulation as a Predicted Main Function of Type I-E CRISPR/Cas System in E. coli. Molecules *24*, 784.

Brouns, S.J.J., Jore, M.M., Lundgren, M., Westra, E.R., Slijkhuis, R.J.H., Snijders, A.P.L., Dickman, M.J., Makarova, K.S., Koonin, E.V., and van der Oost, J. (2008). Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes. Science *321*, 960-964.

Cheng, H.-Y., Soo, V.W.C., Islam, S., McAnulty, M.J., Benedik, M.J., and Wood, T.K. (2014). Toxin GhoT of the GhoT/GhoS toxin/antitoxin system damages the cell membrane to reduce adenosine triphosphate and to reduce growth under stress. Environmental Microbiology *16*, 1741-1754.

Datsenko, K.A., Pougach, K., Tikhonov, A., Wanner, B.L., Severinov, K., and Semenova, E. (2012). Molecular memory of prior infections activates the CRISPR/Cas adaptive bacterial immunity system. Nature Communications *3*, 945.

Datsenko, K.A., and Wanner, B.L. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. Proc. Natl. Acad. Sci. USA *97*, 6640-6645.

Donegan, K., Matyac, C., Seidler, R., and Porteous, A. (1991). Evaluation of methods for sampling, recovery, and enumeration of bacteria applied to the phylloplane. Appl Environ Microbiol *57*, 51-56.

Edgar, R., and Qimron, U. (2010). The *Escherichia coli* CRISPR System Protects from λ Lysogenization, Lysogens, and Prophage Induction. Journal of Bacteriology *192*, 6291-6294.

García Contreras, R., Zhang, X.-S., Kim, Y., and Wood, T.K. (2008). Protein Translation and Cell Death: The Role of Rare tRNAs in Biofilm Formation and in Activating Dormant Phage Killer Genes. PLoS ONE *3*, e2394.

Goldberg, G.W., McMillan, E.A., Varble, A., Modell, J.W., Samai, P., Jiang, W., and Marraffini, L.A. (2018). Incomplete prophage tolerance by type III-A CRISPR-Cas systems reduces the fitness of lysogenic hosts. Nature Communications *9*, 61.

Hafeezunnisa, M., Chhakchhuak, P.I.R., Krishnakumar, J., and Sen, R. (2021). Rho-dependent transcription termination regulates the toxin–antitoxin modules of cryptic prophages to silence their expression in *Escherichia coli*. FEBS Lett *15*, 2057-2067.

Howard-Varona, C., Hargreaves, K.R., Abedon, S.T., and Sullivan, M.B. (2017). Lysogeny in nature: mechanisms, impact and ecology of temperate phages. The ISME Journal *11*, 1511-1520.

Ishihama, A., and Shimada, T. (2021). Hierarchy of transcription factor network in *Escherichia coli* K-12: H-NS-mediated silencing and Anti-silencing by global regulators. FEMS Microbiology Reviews.

Jore, M.M., Lundgren, M., van Duijn, E., Bultema, J.B., Westra, E.R., Waghmare, S.P., Wiedenheft, B., Pul, Ü., Wurm, R., Wagner, R., et al. (2011). Structural basis for CRISPR RNA-guided DNA recognition by Cascade. Nature Structural & Molecular Biology *18*, 529-536.

Kim, J.-S., Yamasaki, R., Song, S., Zhang, W., and Wood, T.K. (2018). Single Cell Observations Show Persister Cells Wake Based on Ribosome Content. Environ Microbiol *20*, 2085-2098.

Kitagawa, M., Ara, T., Arifuzzaman, M., Ioka-Nakamichi, T., Inamoto, E., Toyonaga, H., and Mori, H. (2005a). Complete set of ORF clones of *Escherichia coli* ASKA library (a complete set of *E. coli* K-12 ORF archive): unique resources for biological research. DNA Res. *12*, 291-299.

Kitagawa, M., Ara, T., Arifuzzaman, M., Ioka-Nakamichi, T., Inamoto, E., Toyonaga, H., and Mori, H. (2005b). Complete set of ORF clones of *Escherichia coli* ASKA library (A Complete Set of *E. coli* K-12 ORF Archive): Unique Resources for Biological Research. DNA Res *12*, 291-299.

Kwan, B.W., Valenta, J.A., Benedik, M.J., and Wood, T.K. (2013). Arrested protein synthesis increases persister-like cell formation. Antimicrob Agents Chemother *57*, 1468-1473.

Li, M., Gong, L., Cheng, F., Yu, H., Zhao, D., Wang, R., Wang, T., Zhang, S., Zhou, J., Shmakov, S.A., et al. (2021). Toxin-antitoxin RNA pairs safeguard CRISPR-Cas systems. Science *372*, eabe5601.

Li, R., Fang, L., Tan, S., Yu, M., Li, X., He, S., Wei, Y., Li, G., Jiang, J., and Wu, M. (2016). Type I CRISPR-Cas targets endogenous genes and regulates virulence to evade mammalian host immunity. Cell Res *26*, 1273-1287.

Makarova, K.S., Wolf, Y.I., Iranzo, J., Shmakov, S.A., Alkhnbashi, O.S., Brouns, S.J.J., Charpentier, E., Cheng, D., Haft, D.H., Horvath, P., et al. (2020). Evolutionary classification of CRISPR–Cas systems: a burst of class 2 and derived variants. Nature Reviews Microbiology *18*, 67-83.

Marimon, O., Teixeira, J.M.C., Cordeiro, T.N., Soo, V.W.C., Wood, T.L., Mayzel, M., Amata, I., García, J., Morera, A., Gay, M., et al. (2016). An oxygen-sensitive toxin–antitoxin system. Nature Communications *7*, 13634.

Masuda, H., Awano, N., and Inouye, M. (2016). ydfD encodes a novel lytic protein in *Escherichia coli*. FEMS Microbiology Letters *363*.

Meeske, A.J., Nakandakari-Higa, S., and Marraffini, L.A. (2019). Cas13-induced cellular dormancy prevents the rise of CRISPR-resistant bacteriophage. Nature *570*, 241-245.

Militello, K.T., and Lazatin, J.C. (2017). Discovery of *Escherichia coli* CRISPR sequences in an undergraduate laboratory. Biochem Molec Biol Edu *45*, 262-269.

Mohanraju, P., Saha, C., van Baarlen, P., Louwen, R., Staals, R.H.J., and van der Oost, J. (2022). Alternative functions of CRISPR–Cas systems in the evolutionary arms race. Nature Reviews Microbiology.

Pecota, D.C., and Wood, T.K. (1996). Exclusion of T4 Phage by the *hok/sok* Killer Locus from Plasmid R1. J Bacteriol *178*, 2044-2050.

Perna, N.T., Plunkett, G., Burland, V., Mau, B., Glasner, J.D., Rose, D.J., Mayhew, G.F., Evans, P.S., Gregor, J., Kirkpatrick, H.A., et al. (2001). Genome sequence of enterohaemorrhagic *Escherichia coli* O157: H7. Nature *409*, 529-533.

Pfaffl, M.W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res. *29*.

Pougach, K., Semenova, E., Bogdanova, E., Datsenko, K.A., Djordjevic, M., Wanner, B.L., and Severinov, K. (2010a). Transcription, processing and function of CRISPR cassettes in *Escherichia coli*. Mol Microbiol *77*, 1367-1379.

Pougach, K., Semenova, E., Bogdanova, E., Datsenko, K.A., Djordjevic, M., Wanner, B.L., and Severinov, K. (2010b). Transcription, processing and function of CRISPR cassettes in *Escherichia coli*. Molecular Microbiology *77*, 1367-1379.

Rodriguez, R.L., and Tait, R.C. (1983). Recombinant DNA Techniques: An Introduction (Benjamin/Cummings Publishing).

Rollie, C., Chevallereau, A., Watson, B.N.J., Chyou, T.-y., Fradet, O., McLeod, I., Fineran, P.C., Brown, C.M., Gandon, S., and Westra, E.R. (2020). Targeting of temperate phages drives loss of type I CRISPR–Cas systems. Nature.

Savitskaya, E., Lopatina, A., Medvedeva, S., Kapustin, M., Shmakov, S., Tikhonov, A., Artamonova, II, Logacheva, M., and Severinov, K. (2017). Dynamics of *Escherichia coli* type I-E CRISPR spacers over 42 000 years. Molecular ecology *26*, 2019-2026.

Shmakov, S.A., Sitnik, V., Makarova, K.S., Wolf, Y.I., Severinov, K.V., and Koonin, E.V. (2017). The CRISPR Spacer Space Is Dominated by Sequences from Species-Specific Mobilomes. mBio *8*, e01397-01317.

Song, S., Kim, J.-S., Yamasaki, R., Oh, S., Benedik, M.J., and Wood, T.K. (2021). *Escherichia coli* cryptic prophages sense nutrients to influence persister cell resuscitation. Environ Microbiol *23*, 7245-7254.

Song, S., and Wood, T.K. (2020). Persister Cells Resuscitate via Ribosome Modification by 23S rRNA Pseudouridine Synthase RluD. Environ. Microbiol. *22*, 850–857.

Song, S., and Wood, T.K. (2021). Are We Really Studying Persister Cells? Environmental Microbiology Reports *13*, 3–7.

Touchon, M., Charpentier, S., Clermont, O., Rocha, E.P.C., Denamur, E., and Branger, C. (2011). CRISPR Distribution within the *Escherichia coli* Species Is Not Suggestive of Immunity-Associated Diversifying Selection. Journal of Bacteriology *193*, 2460-2467.

Vasu, K., and Nagaraja, V. (2013). Diverse Functions of Restriction-Modification Systems in Addition to Cellular Defense. Microbiology and Molecular Biology Reviews *77*, 53-72.

Wang, X., Kim, Y., Ma, Q., Hong, S.H., Pokusaeva, K., Sturino, J.M., and Wood, T.K. (2010). Cryptic prophages help bacteria cope with adverse environments. Nat Commun *1*, 147.

Wang, X., Kim, Y., and Wood, T.K. (2009). Control and Benefits of CP4-57 Prophage Excision in *Escherichia coli* Biofilms. ISME J *3*, 1164-1179.

Wang, X., Lord, D.M., Cheng, H.-Y., Osbourne, D.O., Hong, S.H., Sanchez-Torres, V., Quiroga, C., Zheng, K., Herrmann, T., Peti, W., et al. (2012). A Novel Type V TA System Where mRNA for Toxin GhoT is Cleaved by Antitoxin GhoS. Nat. Chem. Biol. *8*, 855-861.

Wood, T.K., and Song, S. (2020). Forming and waking dormant cells: The ppGpp ribosome dimerization persister model. Biofilm *2*, 100018.

Yamasaki, R., Song, S., Benedik, M.J., and Wood, T.K. (2020). Persister Cells Resuscitate Using Membrane Sensors that Activate Chemotaxis, Lower cAMP Levels, and Revive Ribosomes. iScience *23*, 100792.

Fig. 1. Inactivating CRISPR-Cas eliminates persister cell resuscitation by activating cryptic prophage lytic proteins, causing cell death. (A) Single cell persister resuscitation for wild-type BW25113, the *cas2* mutant, and the Δspacer mutant after 6 hours on 0.4 wt% glucose minimal medium. Black arrows indicate cells that resuscitate, and the scale bar indicates 10 μm. Cells were observed using light microscopy (Zeiss Axio Scope.A1). Representative results from two independent cultures are shown, and tabulated cell numbers are in **Table S1**. **(B)** LIVE/DEAD staining of resuscitating persister cells shows the *cas2* and Δspacer mutations cause cell death. DF is dark field, SYTO9 is a membrane permeable stain for nucleic acids (green), and PI is propidium iodide, which is a membrane impermeable stain for the nucleic acids of dead cells (red). Tabulated cell numbers are in **Table S2**. **(C)** The *cas2* mutation derepresses cryptic prophage lysis genes *ydfD* (in resuscitating persister cells) and *rzoD* (in both resuscitating persister cells and exponentially-growing cells). Lytic genes from five cryptic prophages were checked by qRT-PCR: *ydfD* (Qin), *hokD* (Qin), *ypjF* (CP4-57), *essD* (DLP-12), and *rzoD* (DLP-12). Error bar indicates standard deviations.

Fig. 2. *E. coli* CRISPR-Cas spacer sequences and lytic gene inhibition mechanism. (A) The 14 repeat (R, hexagon) and 13 spacer (squares) sequences of the CRISPR-Cas system (from the *iap* to *cas2* part of the *E. coli* genome) showing the cryptic prophage spacer matches (red text) and prophage DNA protospacer sequences (blue text) which includes matches to seven of the nine cryptic prophages (DLP-12, CP4-57, CPS-53, CP4-6, rac, Qin, and e14). Matches indicate mRNA binding to spacer sequences (**Fig. S6**). Pink highlight and pink dashed lines indicate the spacer positions relative to the cryptic prophage genes. The potential PAM sequences are shown in **Fig. S4**. **(B)** Schematic for the mechanism by which CRISPR-Cas controls cryptic phage lytic genes.

Supporting Information

Sooyeon Song[1,2,3*], **Ekaterina Semenova**[4]**, Konstantin Severinov**[4]**, Laura Fernández-García**[1]**, Michael J. Benedik**[5]**, Toshinari Maeda**[6]**, and Thomas K. Wood**[1*]

[1]Department of Chemical Engineering, Pennsylvania State University,
University Park, Pennsylvania, 16802-4400, USA

Departments of Animal Science[2] and Agricultural Convergence Technology[3], Jeonbuk National
University, 587 Baekje-Daero, Deojin-Gu, Jeonju-Si, Jellabuk-Do, 54896, South Korea

[4]Waksman Institute of Microbiology, Rutgers, The State University of New Jersey, Piscataway, NJ 08854

[5]Hamad bin Khalifa University, Education City, Doha, Qatar

[6]Department of Biological Functions Engineering, Kyushu Institute of Technology, Kitakyushu, Japan
808-0196

*For correspondence. E-mail songsy@jbnu.ac.kr, tuw14@psu.edu

Tel. (+)1 814-863-4811; Fax (1) 814-865-7846

Table S1. Inactivating CRISPR-Cas eliminates persister cell resuscitation on glucose agarose gel pads. Single persister cells were observed using light microscopy (Zeiss Axio Scope.A1). The total number and waking number of persister cells are shown after 6 hours on 0.4 wt% glucose minimal medium. Fold-change in waking is relative to BW25113. These results are the combined observations from two independent experiments (independent culture results separated by "/"), and standard deviations are shown. Sample microscope images are shown in **Fig. 1.**

	Total cells	Waking cells	% waking	Fold-change
BW25113	504/382	199/248	52 ± 18	1
$\Delta cas2$	516/677	6/15	1.7 ± 0.7	-30.9
Δspacer	323/174	10/7	3.6 ± 0.7	-14.6

Table S2. Inactivating CRISPR-Cas causes cell death and ghost cell formation. For resuscitated cells, persister cells were washed with PBS twice, resuscitated by M9 0.4% glucose for 10 min and stained with LIVE/DEAD reagents. Exponential cells were grown to a turbidity of 0.8 (at 600 nm) and stationary cells were grown to turbidity of 2.0. The ghost cells in the persister population were visualized using a Zeiss Axioscope.A1microscope. The results are the combined observations from two independent experiments (independent culture results separated by "/"). The microscope images are shown in **Fig. 1AB** and **Fig. S3**.

	Strains	Total cells	Dead cells	Ghost cells	% Ghost cells	fold-change ghost cells	% dead	fold-change dead cells
Resuscitated	**BW25113**	52/115	2/7	0/1	0.6	1	5.4	1
	Δ*cas2*	142/106	54/38	38/13	20.6	34	37.1	6.9
	Δ**spacer**	161/221	76/76	-	-	-	37.8	7.0
Exponential	**BW25113**	153/147	0/2	-	-	-	0.7	1
	Δ*cas2*	59/40	½	-	-	-	3.4	4.9
	Δ**spacer**	95/146	1/5				2.1	3.15
Stationary	**BW25113**	492/123	1/0				0.1	1
	Δ*cas2*	1136/1736	14/18	-	-	-	1.13	11.2
	Δ**spacer**	200/113	19/8				8.26	81.3
	Δ*casE*	208/175	9/10	-	-	-	5.02	49.4
	Δ9 Δ*cas2*	833/1541	10/13	-	-	-	1.02	
	Δ**spacer3**	136/297	8/73	-	-	-	15.23	
	Δ**spacer12**	513/396	213/92	-	-	-	32.38	

Table S3. CRISPR-Cas does not affect cryptic prophage excision in stationary cells but produces the CRISPR array. (A) Fold changes in excision are relative to the wild-type strain as determined by qPCR. **(B)** Fold changes in the CRISPR array are relative to the wild-type strain as determined by qRT-PCR (the Δspacer strain was used as a negative control). Cycle numbers (C_t) are indicated for each sample including that for the target genes as well as that of the house-keeping gene, *purM*, which was used to normalize the data. *cas2* served to inactivate crRNA production due to a polar mutation. Fold changes in the transcription of various targets (i.e., cryptic prophage CP4-57, e14, DLP-12, Spacer3, and Spacer12) with and without $cas2^+$ ($cas2^+/cas2^-$) were calculated as described earlier (Pfaffl, 2001):

$$2\wedge\text{-}(C_{t\ target\_cas2+}\text{-}C_{t\ purM\_cas2+})/2\wedge\text{-}(C_{t\ target\_cas2-}\text{-}C_{t\ purM\_cas2-})$$

(A) qPCR

Gene	*purM*		CP4-57		e14		DLP-12	
Strain	WT	*cas2*	WT	*cas2*	WT	*cas2*	WT	*cas2*
C_T	9.30 ± 0.13	9.70 ± 0.18	30.45 ± 1.73	30.37 ± 0.68	16.30 ± 0.06	17.45 ± 0.51	30.58 ± 0.49	30.05 ± 0.5
ΔC_T			21.16 ± 1.73	20.67 ± 0.70	7.00 ± 0.15	8.16 ± 0.54	21.29 ± 0.51	20.35 ± 0.53
$\Delta\Delta C_T$				-0.48 ± 0.7		1.16 ±0.54		-0.94 ±0.53
fold				1.40		-2.23		1.92

(B) qRT-PCR

Gene	*purM*			Spacer3			Spacer12		
Strain	WT	*cas2*	spacer	WT	*cas2*	spacer	WT	*cas2*	spacer
C_T	25.3 ± 0.5	22.2 ± 0.5	22.0 ± 0.2	30.68 ± 1.73	24.35 ± 0.93	-	33.60 ± 1.20	27.12 ± 1.22	-
ΔC_T				5.37 ± 1.08	2.19 ± 1.1	-	8.28 ± 1.31	4.96 ± 1.32	-
$\Delta\Delta C_T$					-3.20 ± 1.51	-		-3.33 ±1.86	-
fold					9.11	-∞		10.02	-∞

Table S4. *E. coli* bacterial strains and plasmids utilized.

Strains and Plasmids	Features	dSource
Strains		
BW25113	*rrnB3 ΔlacZ4787 hsdR514 Δ(araBAD)567 Δ(rhaBAD)568 rph-1*	(Baba et al., 2006)
BW25113 Δ*cas1*	Δ*cas1*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*cas2*	Δ*cas2*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*cas2*	Δ*cas2*, ΔKm^R	this work
BW25113 Δ*cas3*	Δ*cas3*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*casA*	Δ*casA*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*casB*	Δ*casB*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*casC*	Δ*casC*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*casD*	Δ*casD*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*casE*	Δ*casE*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δspacer (also called BW39292)	Δ(13 spacer region), ΩKm^R	(Pougach et al., 2010b)
BW40114	*placuv5::cas3 paraB8p::casABCDE12spacers*	(Datsenko et al., 2012)
BW25113 Δspacer3	Δ(spacer region 3), ΔKm^R	this work
BW25113 Δspacer12	Δ(13 spacer region 12), ΔKm^R	this work
BW40114	BW25113 *lac*UV5::*cas3 araB*8p::*casABCDE12*	(Datsenko *et al.*, 2012)
BW15113 Δ9	BW15113 lacking all 9 cryptic prophages	(Wang *et al.*, 2010)
BW15113 Δ9 *cas2*	BW15113 Δ9 *cas2*, ΔKm^R	this work
ssrA	*ssrA*, ΩKm^R	(Wang et al., 2009)
Plasmids		
pCA24N	Cm^R; *lacI^q*	(Kitagawa et al., 2005a)
pCA24N_*cas2*	Cm^R; *lacI^q*, P$_{T5-lac}$::*cas2$^+$*	(Kitagawa *et al.*, 2005a)
pKD4	FRT::Kan^R::FRT, Amp^R	(Datsenko and Wanner, 2000)
pCP20	FLP synthase, Amp^R	(Datsenko and Wanner, 2000)

Table S5. Primers used in this study for qRT-PCR and qPCR. * indicates excision primers.

Gene	Sequence (5' to 3')
qRT-PCR	
purM	**F:** CGGTGTTGATATTGACGCGGG
	R: CAGCACGGGTTCACGATATTTTG
*ydfD (*Qin*)*	**F:** TCAGCATTTGTGCTTGTTCTG
	R: CTGCCGGGATTTCGATATTA
*hokD (*Qin*)*	**F:** GCCCTGATCGTCATCTGTTT
	R: AGCTGTGAAGACAGCGACCT
ypjF (CP4-57*)*	**F:** TATGGCCTGACACTGAACGA
	R: GTTGTTGCCACGAAAATCCT
rzoD (DLP-12)	**F:** TGCTCTGCGTGATGATGTTG
	R: TCTCTCTGAGGGTGAAATAATCC
essD (DLP-12)	**F:** CTCACAGTGGGCAGCAATAG
	R: TATTCACCTCTCGCAGCCTT
Spacer3	**F:** GATAAACCGCTTTCGCAGAC
	R: AGAAATTCCAGACCCGATCC
Spacer12	**F:** AACGCCTGAACCGAGAGTT
	R: CTGGGAGTTCTACCGCAGAG
qPCR	
*CP4-57	**F:** AAGCATGTAGTACCGAGGATGTAGG
	R: TATGTCTCCTCACCGTCTGGTCGG
*e14	**F:** GTGCAAACATCGGTGACGAA
	R: TTCAGCAGCTTAGCGCCTTC
*DLP-12	**F:** CAAAAGCCATTGACTCAGCAAGG
	R: CGGATAAGACGGGCATAAATGA
purM 400	**F:** CTGATTGCACTCGGTTCCAG
	R: CGTTTTCACCGTTGGCATTG
Spacer 3-1	**F:** CATGCAATTACAACATCAGGGTA
	R: GCCATTGGTAAAACCTTCCA
Spacer 3-2	**F:** AAATTCCGAAAAAGCTCCTGA
	R: TGCTTTCAAGATTATGGCGTA
Spacer 6	**F:** CACGAAAGCCAGCCTATTCC
	R: CCGCTGTTTCTTTCTCCAGG
Spacer 8	**F:** GCCAGCATAATGAGATCGGC
	R: TTTTACCCAAACTCAGCGCG
Spacer 11	**F:** TGTGCAGTTGTACCAGTGGA
	R: CAACCCAGCAAAGTTTCGGA
Spacer 12-1	**F:** TGTATGTGCCCCGGTGTTAT
	R: GCAGATGAAGGCGCATTACA
Spacer 12-2	**F:** AGCTTTACACCTCGGCTCAT
	R: CCGGAACTCTTGTGTTGGTG
Spacer 12-3	**F:** TCTACAGGGAAAGGACGACC
	R: CTCTGCAACCAAAGTGAACCA
Spacer 13	**F:** ACAACCACTATCGCCCCTTT
	R: GGTAAGGCTGCATTGGGAAG
Deletions	
cas2	**F:** ACCTTAATGTAACATTTCCTTATTATTAAAGATCAGCTAATTCTTTGTTTGTGTAGGCTGGAGCTGCTTC
	R: CATTCCGCTTCCTGTTTCACTGGGAGATGCAGGCCATCGGAGTAGCTGAACATATGAATATCCTCCTTAG
Spacer 3	**F:** GCTGGCGCGGGGAACTCGCGACCGCTCAGAAATTCCAGACCCGATCCAAAGTGTAGGCTGGAGCTGCTTC
	R: AAAGGTGATGCCGAACACGCTGAGTTCCCCGCGCCAGCGGGGATAAACCGCATATGAATATCCTCCTTAG
Spacer 12	**F:** CAGAGGCGGGGGAACTCCAAGTGATATCCATCATCGCATCCAGTGCGCCCGTGTAGGCTGGAGCTGCTTC
	R: GGTTTGCAACGCCTGAACCGAGAGTTCCCCGCGCCAGCAGGGATAAACC

GCATATGAATATCCTCCTTAG

Verification for *cas2* and Spacer

cas2	**F:** GAAAATGTACCTCCGCGCTT	
	R: CCATCCAAATCTACCGGGGT	
Spacer	**F:** GGATCGGGTCTGGAATTTCT	
	R: GGGGAACTCGTAGTCCATCA	

(A)

(B)

Supplementary Figure 1. Removing the kanamycin marker from the *E. coli Δcas2 ΩkanR* strain to create the *Δcas2 ΔkanR* strain **(A)** increased growth in M9 glucose medium at 37°C and **(B)** reduced toxicity via the LIVE/DEAD assay for stationary cells showing the polar mutation due to the kanamycin marker in the *Δcas2 ΔkanR* strain inhibits spacer formation, which results in reduced growth and increased toxicity.

BW25113 (K12)

Δcas2 (K12)

Δspacer (K12)

Supplementary Figure 2. Schematics of the CRISPR-Cas strains used in this study.

BW25113   ***ΔssrA***

Supplementary Figure 3. As a control for growth rate and resuscitation, the *ssrA* mutation was checked and found to have no effect on persister resuscitation on LB plates although it grows 22% more slowly than the wild-type strain (1.18 ± 0.07/h vs. 1.49 ± 0.03/h, respectively). The number of *ssrA* persisters that resuscitated is 101 ± 2 vs. 100.0 ± 0.1 for the wild-type. Plates were incubated for 24 h at 37°C.

Supplementary Figure 4. CRISPR-Cas prevents cell lysis induced by cryptic phages. (A)
Single cell persister waking of BW25113 Δ*cas2* and the Δspacer mutant on M9 0.4% glucose agar
plates incubated at 37°C for 4 hours. The scale bar indicates 10μm. (**B**) TEM image for persister
waking of BW25113 Δ*cas2* and the Δspacer mutant. Persister cells were resuscitated by M9 0.4%
glucose for 10 min. One representative image is shown. The scale bar indicates 1 μm.

Supplementary Figure 5. LIVE/DEAD staining of stationary (turbidity 2.0) and exponential (turbidity 0.8) cells shows the *cas2*, Δspacer (all 13 spacers removed), and mutations cause cell death. DF is dark field, SYTO9 is a membrane permeable stain for nucleic acids (green), and PI is propidium iodide, which is a membrane impermeable stain for the nucleic acids of dead cells (red). Tabulated cell numbers are shown in **Table S2**.

yfdR gene (Start and stop codon)(CPS-53, encodes 5'-deoxynucleotidase)

ATGTCATTTATTAAAACTTTTTCCGGGAAGCATTTTTATTATGACAAGATAAATAAAGACGACATCGTGATTAACGA
TATCGCGGTTTCCCTTTCAAATATCTGCCGCTTTGCCGGTCATCTTTCTCACTTCTACAGTGTCGCCCAACATGCGG
                                                          Spacer 8 3'-ACCCGCCGA
TGCTTTGCAGCCAGCTGGTGCCGCAGGAATTTGCTTTTGAAGCATTAATGCATGATGCAACAGAAGCGTATTGCCAG
|||||||||||||||||
ACGGAACGTCGGTCGAGGTCGTC-5'
GACATCCCCGCCACCACTGAAACGCCTTCTTCCTGACTATAAACGGATGGAAGAAAAAATAGATGCAGTAATCCGTGA
GAAATACGGGTTACCTCCTGTTATGAGCACGCCAGTGAAATATGCCGATCTCATTATGCTGGCAACCGAACGCCGTG
ATCTCGGGCTTGATGATGGCTCTTTCTGGCCTGTACTGGAAGGCATCCCGGCAACAGAGATGTTCAACGTGATTCCA
CTGGCACCGGGTCATGCCTACGGGATGTTTATGGAACGTTTTAACGATTTATCGGAGTTACGCAAATGCGCATGA

perR gene (Start and stop codon)(CP4-6, encodes transcriptional regulator)

ATGAAGCTCTTAGCAAAAGCACCTCTAAATCTGTTACGCGCCTTTGAAGCGGCTGGTCGTACCGGAGCTTTTGCGTT
GGCCGCCTCTGAGTTGGAACTGTCACCCAGTGCGATCAGCCATGCCATCCGCAAACTGGAAAACTTGCTTGATGTAC
GCCTTTTTCAGCGCAGTACGCGAGAAATTACGCTGACGAAAGAAGGTGAAATACTGCTTGAGCACATACAGCGGGGA
TTTAACGAATTACAGCAGGGGTTGGCATTAGTGACGGCTGATGAATCCAGGCCTTTGCGCCTTCATACTGCACCGAG
TTTTGCCCATCAATGGCTTTTACCACGTCTTGGTAAGTTCATACGTGAGAACCCAAGCATCGATCTGCGACTTTCAG
CCAGCACAGAATATGCACGTTTTGAACAGGATGATTTTGATCTCGATATAGTCTACGGTGAACCCCGCCCATCACCC
TATGAGAAGATCCCGCTTGCTGTTGAAGAACTTACGCCACTGTGTTCTCCCCAACTGGCTGAGCGGCTAAAGAAACC
AGAGGATCTCTATGCGTTGACATTGATTCAGTGCGATGTGCAGTTGTACCAGTGGAAAGGATGGTTTGAGGCGAATA
AGATGACGCCACCCAATAATTATGGCCTTCGATTTGACCGTAGCTTTATGGCGATTGCTGCCGCAGTTGACGGCTTG
GGAGTTGTGCTTGAATCAAAACTACTGGCTGAACGTGAAATCGCTAGCGGTAAACTGGTATGCCCACTTGTAAACAG
TACCAGCGAAATACATTACATCGGTCATTATTTGGTTTTCCCCCAGCACCAACACATGCATTCTGCGCTTGATGTGT
                                                    // // / / //
                                    Spacer 11 3'-AGCCAAGTCCGCA
TCAAAACCTGGCTACTGAATGAACTCAATCTGGGTAAAATTCGTTGA
| |||||||||||| |
ACGTTTGGACCGATGGCCC-5'

stfR gene (Start and stop codon)(Rac, side tail fiber protein)

ATGGCAGTAAAGATTTCAGGTGTACTGAAAGACGGCACAGGAAAACCGGTACAGAACTGCACAATCCAGCTGAAAGC
AAAACGTAACAGCACCACGGTGGTGGTGAACACGCTGGCCTCAGAAAATCCGGATGAAGCCGGGCGTTACAGCATGG
ACGTTGAGTACGGTCAGTACAGCGTTATTCTGTTGGTGGAAGGATTCCCGCCGTCACATGCCGGGACCATTACCGTG
TATGAAGATTCTCAACCCGGTACGCTGAATGATTTTCTCGGTGCCATGACGGAGGATGATGCCCGTCCGGAGGCACT
GCGCCGTTTTGAACTGATGGTGGAAGAGGTGGCGCGTAACGCGTCCGCGGTGGCACAGAACACGGCAGCCGCGAAGA
AGTCAGCCAGTGATGCCAGCACATCAGCCCGTGAGGCGGCAACCCATGCGGCTGATGCTGCGGACTCAGCACGCGCA
GCCAGCACGTCAGCCGGACAGGCCGCGTCGTCGGCTCAGTCAGCGTCTTCCAGCGCAGGAACGGCATCAACAAAGGC
CACTGAAGCATCAAAAAGTGCTGCCGCTGCAGAGTCCTCAAAAAGCGCGGCGGCCACCAGTGCCGGTGCGGCGAAAA
CGTCAGAAACGAATGCTTCAGCGTCACTACAATCAGCAGCCACATCTGCATCCACCGCGACCACGAAGGCATCAGAA
GCTGCGACCTCGGCCCGGGATGCGGCGGCCTCAAAAGAAGCGGCAAAATCATCAGAAACGAACGCATCATCAAGCGC
CAGTAGTGCAGCTTCCTCGGCAACGGCGGCAGGAAATTCCGCGAAGGCGGCAAAAACGTCCGAGACGAACGCCAGGT
CTTCTGAAACGGCAGCGGGACAGAGCGCCTCGGCTGCGGCAGGCTCAAAAACAGCGGCTGCGTCGTCTGCCAGTGCA
GCGTCAACAAGTGCCGGGCAGGCCTCAGCCAGTGCCACCGCCGCCGGAAAATCGGCAGAAAGCGCCGCATCGTCTGC
TTCAACAGCCACAACGAAGGCTGGCGAAGCCACTGAACAGGCCAGCGCAGCAGCGAGGTCTGCTTCCGCAGCGAAGA
CATCCGAAACGAACGCGGAAAGCGTCGGAAACAAGCGCAGAATCCTCAAAAACGGCTGCCGCATCGTCAGCCAGTTCG
GCGGCGTCATCGGCATCATCGGCGTCTGCTTCAAAAGATGAGGCGACCAGACAAGCGCTCAGCAGCGAAGAGCAGCGC
CACGACGGCATCCACGAAGGCGACAGAGGCTGCTGGCAGTGCGACGGCGGCAGCTCAGAGCAAAAGTACGGCGGAAT
CCGCGGCAACGCGCGCCGAGACAGCAGCTAAACGGGCAGAGGATATTGCATCCGCCGTGGCGCTTGAGGATGCAAGT
ACGACGAAAAAAGGGGATAGTACAGCTCAGCAGTGCGACCAACAGTACGTCTGAAACGCTGGCGGCAACGCCAAAGGC
AGTAAAAATCAGCCTATGACAATGCAGAGAAACGTCTGCAGAAAGACCAGAACGGCGCTGATATACCCGATAAGGGAT
GCTTCCTGAACAACATTAACGCGGTCAGTAAAACAGACTTTGCTGATAAGCGTGGTATGCGTTATGTGCGGGTTAAC
GCTCCTGCAGGTGCAACATCTGGAAAATATTACCCTGTTGTTGTTATGCGTTCTGCTGGCTCAGTAAGCGAACTGGC
ATCAAGAGTCATTATCACCACGGCAACGCGAACCGCAGGCGATCCGATGAATAACTGCGAGTTTAACGGATTTGTTA
TGCCTGGTGGCTGGACTGACAGGGGGCGTTATGCTTATGGCATGTTCTGGCAATATCAAAACAATGAACGAGCCATT
CACTCAATAATGATGAGTAATAAAGGGCGATGATTTGCGCTCTGTGTTCTATGTTGATGGCGCTGCTTTCCCTGTTTT
TGCGTTTATTGAAGATGGCCTGTCAATATCCGCACCTGGTGCTGATCTCGTTGTTAATGATACGACCTATAAGTTTG
GGGCAACAAATCCGGCGACTGAATGTATCGCGGCGGACGTTATCCTTGATTTTAAGAGTGGGCGTGGTTTTTATGAG
TCTCATTCGTTAATCGTTAACGATAACTTGTCGTGCAAAAAACTTTTTGCCACAGACGAAATTGTAGCGCGTGGTGG
TAATCAGATTCGAATGATAGGTGGGGAGTATGGTGCATTATGGCGTAATGATGGCGCTAAAACTTACCTGCTGCTTA
CCAATCAAGGTGATGTTTATGGTGGCTGGAATACATTAAGACCGTTTGCTATTGATAACGCAACCGGCGAACTGGTT
ATTGGAACCAAACTGTCCGCAAGTCTGAACGGTAATGCATTAACAGCAACAAAGCTGCAAACGCCAAGACGGGTTTC
TGGTGTTGAGTTTGATGGTTCCAAAGATATTACTTTAACCGCCGCGCATGTGGCTGCTTTTGCCAGAAGGGCAACGG
ATACATATGCCGATGCGGATGGTGGCGTTCCATGGAATGCCGAATCTGGCGCTTACAATGTCACCCGCTCTGGCGAC
AGCTATATTCTGGTTAACTTCTATACCGGAGTCGGAAGTTGCCGGACCCTGCAGATGAAGGCGCATTACAGAAATGG
TGGTCTGTTCTACCGTTCTTCAAGAGACGGTTATGGTTTTGAGGAAGACTGGGCAGAAGTTTATACCTCGAAAAATC
TTCCACCAGAAAGCTACCCAGTCGGCGCACCAATCCCGTGGCCATCAGATACCGTTCCGTCTGGTTATGCCCTGATG
CAGGGGCAGGCTTTTGACAAATCTGCTTACCCGAAACTTGCAGCCGCTTATCCGTCAGGCGTGATCCCTGATATGCG
                                       |||||||||||| || | ||
                          Spacer 12 3'-GTCGCAGTCCGCACTTTAGAGTGGCAGC
AACG-5'
| ||
TGGCTGGACGATTAAGGGCAAACCTGCCAGTGGTCGGGCCGTATTGTCTCAGGAACAGGACGGCATTAAATCGCATA
CCCACAGCGCCAGCGCATCCAGTACAGATTTGGGGACGAAAACCACATCGTCGTTTGATTACGGCACTAAATCCACG
AATAACACCGGGGCACATACACACAGTGTGAGCGGCTCTACAAACTCGGCTGGAGCACACACACACTCACTAGCCAA
CGTGAACACGGCTAGTGCTAACTCCGGTGCTGGTAGTGCATCAACAAGATTGTCTGTTGTGCATAATCAAAACTATG
CAACATCATCTGCTGGCGCACATACCCACTCACTGTCCGGCACTGCTGCAAGCGCAGGTGCACACGCGCATACTGTC
GGTATTGGTGCTCATACGCACTCCGTTGCGATTGGTTCACATGGACACACCATCACCGTTAACGCTGCTGGTAACGC
GGAAAACACCGTCAAAAACATCGCATTTAACTATATTGTGAGGCTTGCATAA

stfQ gene (`Start` and `stop` codon) (**Qin**, side tail fiber protein)

```
GTGAATATAACCGCGCTGACAGACAATACGCAGGGGGCAGCAGGTCTTGAGTTATACGAGGTGTATAACAACGGATA
TCCAACAGCGTATGGAAATATCATTCACCTGAAAGGGATGACAGCCGTTGGCGAAGGTGAGTTACTCATCGGCTGGA
GTGGTACAAGCGGTGCTCATGCTCCGGCATTTATTCGTTCACGACGGGATACGACCGACGCAAACTGGTCGCCGTGG
GCGCAGCTTTACACCTCGGCTCATCCTCCTGCAGAGTTTTATCCAGTCGGTGCACCAATCCCGTGGCCATCAGATAC
CGTTCCGTCTGGTTATGCCCTGATGCAGGGGCAGACTTTTGACAAATCTGCATACCCGAAACTTGCAGTTGCTTATC
                                                      Spacer 12 3'-GTC
CGTCAGGCGTGATCCCTGATATGCGTGGCTGGACGATTAAGGGCAAGCCCGCCAGTGGTCGGGCCGTATTATCTCAG
|||||||||||||   || |    ||| ||
GCAGTCCGCACTTTAGAGTGGCAGCAACG-5'
GAACAGGACGGCATTAAATCGCACACCCACAGCGCCAGCGCATCCAGTACGGATTTGGGGACGGAAACCACATCGTC
GTTTGATTACGGAACCAAATCCACGAATAACACCGGGGCGCATACCCATAGTATTAGCGGGACCGCAAATAGTGCCG
GTGCGCACCAACACAAGAGTTCCGGTGCATTTGGTGGCACGAACACGAGCATTTTCCCTAATGGTTATACCGCGATT
TCAAATCTAAGCGCGGGGATTATGAGCACAACAAGCGGTAGTGGCCAGACTCGTAATGCAGGGAAGACATCATCAGA
TGGTGCTCATACCCACTCGCTGTCCGGCACTGCTGCAAGCGCAGGCGCGCATGCACATACTGTCGGTATTGGTGCTC
ATACGCACTCCGTTGCGATTGGTTCACATGGACACACCATCACCGTTAACGCTGCTGGTAACGCGGAAAACACCGTC
AAAAACATCGCATTTAACTATATTGTGAGGCTTGCATAA
```

alpA gene (`Start` and `stop` codon) (**CP4-57**, transcriptional regulator)

```
ATGAACAATCGATCGGCCGTTAGAATACTACGGTTACCAGCGGTTATCCAAAAAACAGGTATGGCACGGGCCACCAT
CTATGACTGGTTGAACCCCAAATCACCACGATACGATGCCACCTTTCCCAAAAAGCGAATGCTCGGCGTGAAATCTG
                                             | |    ||||||||||||
                                      Spacer 12 3'-GTCGCAGTCCGCACTTTAGAG
TGGCAGCAACG-5'
 ||| |
TCGGATGGATTGAGGCCGAGATTGATGAGTGGTTATCACAACGCTGTAAACTTATTTGA
```

tfaP gene (`Start` and `stop` codon)(**e14**, tail fiber assembly)

```
ATGAAAATATACTGTTGCTTAAATACCGTTGGTTTTTTTATGGATGGCTGTGGCGTCATTCCGCCAGATTCTAAAGA
AATAACGGCAGAACACTGGCAGTCATTATTAAAATCTCAAGCTGAAGGAGGCGTGATCGATTTTTCTGTTTTTCCTC
CTTCTATTAAAGAGGTTATCCGTACTCATGATGATGAAGTCGCAGATGCGAACTTTCAAAAGCAGATGCTTATCTCT
GATGCAACTGATTTTATCAATAGCAGACAGTGGCAGGGTAAGGCTGCATTGGGAAGACTTAAAGAAGATGAGCTGAA
ACAATATAATTTGTGGCTGGATTATCTGGAAGCACTGGAACTGGTTGATACATCCAGTGCGCCAGATATTGAATGGC
                                  | ||  |    |    |||||||||||||
                      Spacer 13 3'-GTTCACTATAGGTAGTAGCGTAGGTCACGCGG-5'
CTACGCCTCCGGCAGTTCAGGCCAGATGA
```

Supplementary Figure 6. Predicted base-pairing interactions between each of the CRISPR spacers (i.e., spacers 8, 11, 12, and 13) and a protospacer within the cryptic phage mRNA (CPS-53, CP4-6, Rac, Qin, CP4-57, and e14). Note, some spacers match more than one region within a particular cryptic prophage. The complementary nt between the spacers and cryptic prophage protospacers is indicated by blue text and vertical lines.

Supplementary Figure 7. Toxicity of YdfD and RzoD. Growth curve of *E. coli* BW25113 harboring empty plasmid pCA24N (blue line), plasmid pCA24N encoding *ydfD* (orange line), and encoding plasmid pCA24N encoding for *rzoD*. Red arrow indicates addition of 1 mM of IPTG produce the toxins. Average of two independent cultures shown.

Supplementary Figure 8. Deleting the DNA encoding spacer #12, which encodes crRNA targeting three separate regions in cryptic prophage mRNA (mRNA of CP4-57, Rac, and Qin), inhibits cell growth (**A**) due to cell lysis (**B**). In contrast, deleting *cas2* in Δ9 and deleting spacer 3, which encodes crRNA that lacks matches in the mRNA of all 9 cryptic prophages, has no effect on growth and cell lysis. Δ9 is BW25113 that lacks all cryptic prophages and Kan[R] is kanamycin resistance.

D

Spacer sequence

Cryptic prophage Gene

S. enterica subsp. *enterica* serovar *Typhimurium* str. LT2

Supplementary Figure 9. *E. coli* O157:H7 and *Salmonella* spp. CRISPR-Cas spacer sequences related to cryptic prophages. **(A)** Repeat (R, blue hexagon) and spacer (rectangle) sequences of each CRISPR-Cas system of *E. coli* O157:H7, *Salmonella* spp., and *K. pneumoniae* indicating matches with their cryptic prophages. **(A)** *E. coli* O157:H7, **(B)** *S. enterica* subsp. *enterica* serovar *Typhi* strain TY2, **(C, D, E)** three CRISPR-Cas systems of *S. enterica* subsp. *enterica* serovar *Typhimurium* str. LT2, and **(F)** *K. pneumoniae*. Pink highlight and dashed lines indicate the spacer positions relative to the cryptic prophage genes.