CRISPR-Cas Controls Cryptic Prophages

Sooyeon Song[1,2,3*], Ekaterina Semenova[4], Konstantin Severinov[4], Laura Fernández-García[1], Michael J. Benedik[5], Toshinari Maeda[6] and Thomas K. Wood[1*]

[1]Department of Chemical Engineering, Pennsylvania State University, University Park, Pennsylvania, 16802-4400, USA

[2]Department of Animal Science, Jeonbuk National University, 587 Baekje-Daero, Deojin-Gu, Jeonju-Si, Jellabuk-Do, 54896, South Korea

[3]Agricultural Convergence Technology, Jeonbuk National University, 587 Baekje-Daero, Deojin-Gu, Jeonju-Si, Jellabuk-Do, 54896, South Korea

[4]Waksman Institute of Microbiology, Rutgers, The State University of New Jersey, Piscataway, NJ 08854

[5]Hamad bin Khalifa University, Education City, Doha, Qatar

[6]Department of Biological Functions Engineering, Kyushu Institute of Technology, Kitakyushu, Japan 808-0196

* Correspondence: songsy@jbnu.ac.kr, tuw14@psu.edu; Tel (+)1 814-863-4811; Fax (1) 814-865-7846

Abstract: The bacterial archetypal adaptive immune system, CRISPR-Cas, is thought to be repressed in the best-studied bacterium, *Escherichia coli* K-12. We show here that the *E. coli* CRISPR-Cas system is active and serves to inhibit its nine defective (i.e., cryptic) prophages. Specifically, compared to the wild-type strain, reducing the amounts of specific interfering RNAs (crRNA) decreases growth by 40%, increases cell death by 700%, and prevents persister cell resuscitation. Similar results were obtained by inactivating CRISPR-Cas by deleting the entire 13 spacer region (CRISPR array); hence, CRISPR-Cas serves to inhibit the remaining deleterious effects of these cryptic prophages, most likely through CRISPR array-derived crRNA binding to cryptic prophage mRNA, rather than through cleavage of cryptic prophage DNA; i.e., self-targeting. Consistently, four of the 13 *E. coli* spacers contain complementary regions to the mRNA sequences of seven cryptic prophages, and inactivation of CRISPR-Cas increases the level of mRNA for lysis protein YdfD of cryptic prophage Qin and lysis protein RzoD of cryptic prophage DLP-12. Also, lysis is clearly seen via transmission electron microscopy when the whole CRISPR-Cas array is deleted, and eliminating spacer #12, which encodes crRNA with complementary regions for DLP-12 (including *rzoD*), Rac, Qin (including *ydfD*), and CP4-57 cryptic prophages, also results in growth inhibition and cell lysis. Therefore, we report the novel results that (i) CRISPR-Cas is active in *E. coli* and (ii) CRISPR-Cas is used to tame cryptic prophages, likely through RNAi; i.e., unlike with active lysogens, active CRISPR-Cas and cryptic prophages may stably co-exist.

Keywords: CRISPR-Cas; persisters; cryptic prophage

1. INTRODUCTION

Along with restriction/modification (Vasu and Nagaraja, 2013) and toxin/antitoxin (TA) systems (Pecota and Wood, 1996), prokaryotes utilize clustered, regularly-interspaced, short palindromic repeats (CRISPR) and CRISPR-associated (Cas) (Makarova et al., 2020) proteins to combat phages. These systems are interrelated in that some Cas proteins and TA systems have a common ancestor; for example, *Sulfolobus solfataricus* Cas2 is structurally similar to antitoxin GhoS (an RNase) of the *Escherichia coli* GhoT/GhoS TA system (Wang et al., 2012). In addition, in a manner similar to our discovery (Pecota and Wood, 1996) that toxins of host TA systems inhibit phage by degrading mRNA when the phage shuts down transcription (e.g., Hok/Sok inhibits T4 phage), some Cas proteins induce host dormancy rather than degrading phage DNA to inhibit phage propagation (Meeske et al., 2019). Also, TA systems have been found to stabilize CRISPR-Cas systems by making them addictive to the host (Li et al., 2021).

Although CRISPR-Cas systems exclude both external lytic and temperate (lysogenic) phages (Goldberg et al., 2018), CRISPR-Cas systems of lysogens that target their own integrated prophages decrease long-term fitness and either the cell dies or the prophage is

lost (Edgar and Qimron, 2010; Goldberg *et al.*, 2018). Also, the class I-E (Makarova *et al.*, 2020) CRISPR–Cas system of *E. coli* is not related to immunity for external phages (Touchon et al., 2011) and appears to be inactive in the wild-type strain at standard laboratory conditions (Shmakov et al., 2017), due to repression by H-NS, although it is functional when induced (Pougach et al., 2010). To date, the relationship of CRISPR-Cas to cryptic prophages; i.e., those phage remnants that are unable to form lytic particles, has not been investigated.

Up to 50% of bacterial genomes may contain stably-integrated phage DNA (Howard-Varona et al., 2017), and for *E. coli,* we discovered that its nine cryptic prophages are not extraneous DNA but instead encode genes for proteins that increase resistance to sub-lethal concentrations of quinolone and β-lactam antibiotics as well as protect the cell from osmotic, oxidative, and acid stresses (by deleting 166 kb) (Wang et al., 2010). Although these cryptic prophages do not enhance the formation of persister cells, a subpopulation of cells that weather extreme stress by entering into a dormant state (Wood and Song, 2020), these phage remnants facilitate the resuscitation of persisters via nutrient sensing (Song et al., 2021). Therefore, the bacterial cell can co-opt the genome of its former parasite to both combat stress (Wang *et al.*, 2010) as well as to revive from dormancy (Song *et al.*, 2021).

Given these active roles of cryptic prophages in the stress response (Wang *et al.*, 2010) and persister cell resuscitation (Song *et al.*, 2021), we hypothesized here that the native *E. coli* CRISPR-Cas system plays an active role in the regulation of cryptic prophages. We find that the CRISPR-Cas system is required for inhibiting the expression of deleterious cryptic prophage genes, since, if CRISPR-Cas is inactivated by preventing crRNA production, cells die due to activation of the cryptic prophage lysis proteins YdfD of Qin and RzoD from DLP-12. Hence, we discovered CRISPR-Cas is active in *E. coli* and serves to regulate its former phage foe*.

2. RESULTS

Deletions of CRISPR-Cas components reduce growth. We assayed the importance of CRISPR-Cas in *E. coli* K-12 by testing the individual deletions of the CRISPR-Cas genes on growth in M9 glucose medium (*cas1, cas2, cas3, casA, casB, casC, casD,* and *casE*) (**Fig. S1**) and found only the *cas2* deletion, with its kanamycin cassette insertion, had an effect: *cas2* causes 40% slower growth in rich medium compared to the wild-type strain (specific growth rate of 0.79 ± 0.21/h vs. 1.3 ± 0.11/h, respectively). Similarly, in minimal glucose (0.4 wt%) medium, deletion of *cas2* also reduces growth by 33% (0.42 ± 0.02/h vs. 0.62 ± 0.02/h, respectively) and reduced the yield in stationary phase (**Fig. S2A**). We hypothesized that the *cas2* deletion/kanamycin insertion caused a polar effect by reducing spacer production given it is directly upstream of the spacer region (Pougach *et al.*, 2010) (**Fig. S3**). Fittingly, inactivation of CRISPR-Cas by eliminating the CRISPR arrays in the isogenic host BW25113 (henceforth Δspacer) also reduced growth by 32% in rich medium and eliminating kanamycin resistance in the *cas2* deletion strain to prevent the polar mutation affecting the spacer region restored nearly wild-type growth (**Fig. S2A**) as well as reduced toxicity (**Fig. S2B**) relative to the *cas2* Ωkan^R strain. Therefore, the *E. coli* CRISPR-Cas represses some processes that inhibit growth via crRNA, and by this criterion, is active, providing a clear advantage to the host.

CRISPR-Cas increases single-cell resuscitation. Since deletions of CRISPR-Cas components decrease growth, we tested for their effect on persister cell resuscitation using single-cell microscopy. Persister cell resuscitation is germane in that the dormant cells are highly stressed and have limited resources for their revival via activation of hibernating ribosomes (Kim et al., 2018; Song and Wood, 2020; Yamasaki et al., 2020); for example, we have shown that inhibiting ATP synthesis leads to a 5,000-fold increase in persister cell formation (Kwan et al., 2013). Since we discovered facile means for converting the whole population of cells into persister cells (Kim *et al.*, 2018; Kwan *et al.*, 2013; Yamasaki *et al.*, 2020) that has been used by at least 17 independent labs to date with various bacterial

species (Song and Wood, 2021), these stressed cells are an excellent model for testing the effects of CRISPR-Cas on *E. coli* physiology.

Here, we found that deletion of *cas2* reduces persister cell resuscitation by 31-fold (**Fig. 1A, Table S1**). Persister cell resuscitation was similarly affected (15-fold reduction) in the Δspacer strain (**Fig. 1A, Table S1**). Since the *cas2* mutant grows more slowly than the wild-type in rich medium, as a control, we tested an *E. coli* mutant that grows 22% slower than the wild-type, *ssrA*, to confirm that the slower growth does not affect persister resuscitation and found that the *ssrA* mutant resuscitates at nearly the identical rate as the wild-type strain (**Fig. S4**). Hence, CRISPR-Cas is active in *E. coli* and plays key roles in its growth and recovery from extreme stress.

CRISPR-Cas prevents cell death by preventing cell lysis. To explore how deletion of CRISPR-Cas decreases growth (including persister cell resuscitation), we checked for death among resuscitating persister cells of the *cas2* deletion strain using the Live/Dead stain. We found that inactivating CRISPR-Cas leads to a 7-fold increase in death among resuscitating cells (**Fig. 1B, Table S2**). In addition, there were 34-fold more cells termed "ghosts" (Cheng et al., 2014) that lack cytosolic material (**Fig. S5A**), are likely dead, and have intact membranes, so these cells are not stained by the propidium iodide dye. Corroborating these results, there was 11-fold and 5-fold more death for stationary- and exponential-phase cells, respectively, when CRISPR-Cas was inactivated via *cas2* (**Fig. S6, Table S2**). Moreover, for the Δspacer strain, there was 81-fold more cell death in the for stationary-phase cells (**Fig. S6**) and 7-fold more death for resuscitating persister cells (**Fig. 1B, Table S2**). Critically, for the Δspacer strain, waking persister cells clearly shows lysis (**Fig. S5A**), and transmission electron microscopy images confirm this lysis with cytosolic materials seen next to lysed cells (**Fig. S5B**). Unfortunately, this toxicity could not be reversed by CRISPR-Cas induction after placing the complete CRISPR-Cas system on the chromosome under inducible control (BW40114), since this led to more rapid growth when CRISPR-Cas was not induced as a result of the metabolic burden of this complete system; hence, inducing CRISPR-Cas reduced growth (**Fig. S7**). These results indicate inactivating CRISPR-Cas leads to cell death and that the mechanism for cell death is via lysis.

Figure 1. Inactivating CRISPR-Cas eliminates persister cell resuscitation by activating cryptic prophage lytic proteins, causing cell death. (A) Single cell persister resuscitation for wild-type BW25113, the *cas2* mutant, and the Δspacer mutant after 6 hours on 0.4 wt% glucose minimal medium. Black arrows indicate cells that resuscitate, and the scale bar indicates 10 μm. Cells were observed using light microscopy (Zeiss Axio Scope.A1). Representative results from two independent cultures are shown, and tabulated cell numbers are in Table S1. (B) LIVE/DEAD staining of resuscitating persister cells shows the *cas2* and Δspacer mutations cause cell death. DF is dark field, SYTO9 is a membrane permeable stain for nucleic acids (green), and PI is propidium iodide, which is a membrane impermeable stain for the nucleic acids of dead cells (red). Representative results from two independent cultures are shown, and tabulated cell numbers are in Table S2. (C) The *cas2* mutation derepresses cryptic prophage lysis genes *ydfD* (in resuscitating persister cells) and *rzoD* (in both resuscitating persister cells and exponentially-growing cells). Five lytic genes from three cryptic prophages were checked by qRT-PCR: *ydfD* (Qin), *hokD* (Qin), *ypjF* (CP4-57), *essD* (DLP-12), and *rzoD* (DLP-12). Error bars indicate standard deviations from two independent cultures.

CRISPR-Cas reduces cryptic prophage lysis gene mRNA. We hypothesized that since CRISPR-Cas systems inhibit some phage before they become lysogens (Edgar and Qimron, 2010; Goldberg *et al.*, 2018), the *E. coli* system may be preventing cell death by repressing certain cryptic prophage genes. To test this hypothesis, we first examined the *E. coli* CRISPR-Cas system for spacers related to the nine cryptic prophages. *E. coli* K-12 CRISPR array located next to the *cas* operon contains 13 spacers (Militello and Lazatin, 2017; Pougach *et al.*, 2010), each containing 32 or 33 nt (Brouns et al., 2008). Between the

14 29-nt repeat sequences (5'-GTGTTCCCCGCATCAGCGGGGAfTAAACCG), we found 154
that four of the 13 spacers contain 7 to 16 nt of perfect matches to seven of the nine cryptic 155
prophages (DLP-12, CPS-53, CP4-6, Rac, Qin, CP4-57, and e14) (**Fig. 2A**) based on base- 156
pairing matches with cryptic prophage mRNA (**Fig. S8**), indicating putative binding to 157
cryptic prophage mRNA. Critically, spacer 12 encodes crRNA with the potential to bind 158
mRNA from DLP-12 (*rzoD*, *ybcN*), Rac (*stfR*), Qin (*stfQ*), and CP4-57 (*alpA*) (**Fig. S8**). In 159
general, spacer lengths vary from 21 to 72 nt (Barrangou and Marraffini, 2014) with perfect 160
complementarity of 6 to 12 nt (Shmakov *et al.*, 2017). Together, these results suggest 161
CRISPR-Cas potentially regulates the *E. coli* cryptic prophages. Note that the second 162
CRISPR array in *E. coli* K-12 in between genes *ygcE* and *queE* lacks spacers with sequences 163
that matched the cryptic prophages. 164

The presence of the cryptic prophage-related targeting spacers suggests that *E.* 165
coli CRISPR-Cas may be preventing expression of cryptic prophage genes and, given the 166
cell phenotype, is suggestive of prophage-encoded lethal genes; hence, we checked the 167
mRNA levels of the cryptic prophage lysis genes encoded by these seven cryptic pro- 168
phages with spacer matches. Specifically, transcription of *ydfD* (Qin), *hokD* (Qin), *ypjF* 169
(CP4-57), *essD* (DLP-12), and *rzoD* (DLP-12) was checked via quantitative reverse tran- 170
scription polymerase chain reaction (qRT-PCR). We found that inactivation of CRISPR- 171
Cas via the *cas2* deletion results in a 35-fold increase in *ydfD* mRNA and a 39-fold increase 172
in *rzoD* mRNA in resuscitating persister cells (**Fig. 1C**). RzoD is a putative DLP-12 lysis 173
lipoprotein that we previously showed was toxic through its interaction with toxin Hha 174
(García Contreras et al., 2008) of the Hha/TomB TA system (Marimon et al., 2016). YdfD 175
of Qin cryptic prophage has been shown to lyse cells when induced (Masuda et al., 2016). 176
To confirm that YdfD and RzoD are toxic, we induced production of both proteins and 177
found they are indeed toxic (**Fig. S9**). Moreover, deleting *cas2* to inactivate CRISPR-Cas in 178
the same host that lacks all nine cryptic prophages (delta9) (Wang *et al.*, 2010) has no effect 179
on growth (**Fig. S10**) and does not cause lysis (**Fig. S6**). Hence, these results show CRISPR- 180
Cas represses at least two *E. coli* cryptic prophage proteins RzoD and YdfD that can reduce 181
cell growth. 182

Figure 2. *E. coli* **CRISPR-Cas spacer sequences and lytic gene inhibition mechanism. (A)** The 14 repeat (R, hexagon) and 13 spacer (squares) sequences of the CRISPR-Cas system (from the *iap* to *cas2* part of the *E. coli* genome) showing the cryptic prophage spacer matches (red text) and prophage DNA protospacer sequences (blue text) which includes matches to seven of the nine cryptic prophages (CPS-53, CP4-6, Rac, Qin, CP4-57, DLP-12, and e14). Matches indicate mRNA binding to spacer sequences (**Fig. S8**). Pink highlight and pink dashed lines indicate the spacer positions relative to the cryptic prophage genes. **(B)** Schematic for a hypothetical mechanism by which CRISPR-Cas controls cryptic phage lytic genes.

Excision of DLP-12 is not regulated by CRISPR-Cas. Since inactivation of CRISPR-Cas leads to derepression of the DLP-12 *rzoD* lysis gene, we checked for increased DLP-12 excision with the *cas2* deletion strain using qPCR. We found the *cas2* deletion has little impact on DLP-12 excision (**Table S3**). Since the DLP-12 toxin gene *essD* is not derepressed, the effect is not due to excision of the prophage. We also tested the effect of

CRISPR-Cas on two other cryptic prophages with significant excision (CP4-57 and e14) (Wang *et al.*, 2010), and also found no effect of deleting *cas2* (**Table S3**). Hence, CRISPR-Cas does not affect cryptic prophage excision.

CRISPR-Cas regulates lytic gene mRNA levels. Since CRISPR-Cas does not affect cryptic prophage excision and is unlikely to degrade cryptic prophage DNA as this would be lethal (Edgar and Qimron, 2010; Goldberg *et al.*, 2018), we reasoned that CRISPR-Cas, via crRNA, must be interfering with expression of the mRNA of the lytic genes to prevent production of the lytic proteins. To test this hypothesis, we investigated whether *rzoD* and *ydfD* transcript levels are increased in the *cas2* and Δspacer strain relative to the wild-type after inhibiting transcription via rifampicin. If CRISPR-Cas prevents lytic gene expression by inhibiting lytic transcripts via crRNA, then inactivating CRISPR-Cas should lead to increased *rzoD* and *ydfD* mRNA. In accordance with this hypothesis, *rzoD* mRNA was increased 320% in the Δspacer mutant and 50% in the *cas2* mutant. Moreover, as expected, inactivating CasE increased cell lysis 49-fold compared to the wild-type strain (**Fig. S6, Table S2)**, likely by preventing CasE from properly forming crRNA from pre-crRNA; hence, there is less crRNA to inhibit the cryptic prophage mRNA from toxic genes. Similarly, preventing spacer RNA formation by inactivating Cas2 led to a 10-fold increase in CRISPR array RNA levels in resuscitating persister cells as assayed by qRT-PCR, likely due to a lack of processing of the CRISPR array into crRNA (**Table S3B**). Finally, deleting only spacer #12 (which encodes crRNA targeting three separate regions in the cryptic prophage mRNA of CP4-57, Rac, and Qin, **Fig. S8**), in a markerless fashion, inhibits cell growth in the stationary phase (**Fig. S10**) due to cell lysis (**Fig. S6, Table S2**) to roughly the same extent as deleting all 13 spacers. In contrast, deleting spacer #3 in a markerless fashion, which does not encode crRNA with matches to cryptic prophage mRNA, has little effect on overall growth (**Fig. S10**), but did increase cell death (**Fig. S6, Table S2**), which indicates it does serve some role. Therefore, CRISPR-Cas prevents cryptic prophage lytic gene expression likely by interfering with the transcripts from these genes (**Fig. 2B**), specifically through spacer #12.

3. DISCUSSION

Our results reveal a new (non-canonical) role for CRISPR-Cas systems: regulation of phage fossils. The evidence for this includes that inactivating CRISPR-Cas by deleting *cas2* (i) reduces growth by 40%, (ii) nearly eliminates resuscitation from the persister state (**Fig. 1**), (iii) causes ghost cell formation and cell death (**Fig. S5**), (iv) increases mRNA levels of the cryptic prophage lysis genes *ydfD* and *rzoD*, and (v) increases CRISPR array levels by preventing processing into crRNA. Corroborating these results, the *cas2* deletion has no effect in a strain which lacks cryptic prophage genes and eliminating the CRISPR array with prophage-matching spacers in the wild-type strain also reduced the growth rate, eliminated persister cell resuscitation, resulted in cell death via lysis, and activated *rzoD*. Furthermore, deleting only prophage-matching spacer #12 similarly inhibited growth and caused cell lysis. Given that Cas2 is not required for CRISPR interference, it is likely that the phenotypes produced upon deleting *cas2* are the result of a polar mutation (**Fig. S3**) that inactivates CRISPR-Cas by altering production/processing of the spacers.

Since there is no change in excision of DLP-12 upon inactivating CRISPR-Cas, our results suggest the mechanism for regulating the lysis genes *ydfD* and *rzoD* of the cryptic prophages is via CRISPR-Cas RNA binding to the cryptic prophage mRNA (analogous to RNA interference as seen previously in procaryotic systems with CRISPR-Cas (Mohanraju et al., 2022)) rather than cleaving excised DNA or directly cleaving the mRNA. Our proposed RNAi mechanism has precedent since a previous bioinformatics analysis of 230 phages suggested CRISPR-Cas functions in *E. coli* via endogenous gene expression rather than acting as an immune system (Bozic et al., 2019). In support of this, the excision rates of the cryptic prophages are so low that there may be insufficient excised DNA to cleave (1 excision in 10^6 cells for CP4-57) (Wang *et al.*, 2010), and a class I-F CRISPR-Cas system in *P. aeruginosa* uses crRNA to regulate endogenous LasR mRNA to reduce the pro-inflammatory response (Li et al., 2016). Moreover, *E. coli* crRNA have been shown to be

activated during infection in mice (Li *et al.*, 2016), and the *Salmonella* spp. system is identical to the *E. coli* one (e.g., the repeats are the same), but the spacers are different, which suggests it, too, is for endogenous gene expression. Critically, it has been demonstrated *in vitro* that that the *E. coli* Cascade binds ssRNA complementary to crRNA (Jore et al., 2011). Therefore, there is ample precedence for our proposed RNA interference mechanism.

Although we identified spacer #12 sequences that probably bind and interfere with the cryptic prophages Qin 12 (including *ydfD*) and DLP-12 (including *rzoD*) mRNA, there may be another level of regulation that remains to be identified. For example, we show a spacer #12 region that would bind *alpA* mRNA in cryptic prophage CP4-57, and AlpA is a regulator that impacts persister resuscitation by sensing phosphate nutrients (Song et al., 2021).

This new function for CRISPR-Cas acting on cryptic prophages is likely general and may explain why many species appear to have inactive CRISPR-Cas systems as was previously thought for *E. coli* (Shmakov *et al.*, 2017); i.e., instead of protecting cells from external phages, CRISPR-Cas systems may also control resident cryptic prophages which are prevalent. As additional evidence, we found matches in CRISPR-Cas spacers not only for *E. coli* K-12 (**Fig. 2A**) but also for *E. coli* O157:H7, *Salmonella* spp., and *K. pneumoniae* (**Fig. S11**). Critically, our results provide the first example where it is beneficial for the host to have an active CRISPR-Cas system that targets *inactive* integrated phages (i.e., cryptic prophages) since previous reports show targeting *active* temperate phages is deleterious; i.e., either the cell dies or the phage is lost (Edgar and Qimron, 2010; Goldberg *et al.*, 2018; Rollie et al., 2020). Since *E. coli* cryptic prophages like *rac* have been present in its genome for 4.5 million years (Perna et al., 2001), the active *E. coli* K-12 CRISPR-Cas system is stable with the cryptic prophages; in fact, there has been little change in the *E. coli* spacers for at least 42,000 years (Savitskaya et al., 2017).

Our results also indicate that, although the cryptic prophages are stable and the cell makes use of the genetic tools encoded by its former foe to combat myriad stresses (Wang *et al.*, 2010) and to sense nutrients prior to exiting the persister state (Song *et al.*, 2021), the source of these tools must be elegantly regulated by CRISPR-Cas since they often harbor deleterious membrane lysis proteins like YdfD and RzoD. Similarly, host Rho has been found recently to silence cryptic prophage toxin/antitoxin systems through transcription termination (Hafeezunnisa et al., 2021), and H-NS silences cryptic prophages through 65 binding sites (Ishihama and Shimada, 2021). Therefore, phages may be captured by the host, but they must be tamed, and this now includes repression by CRISPR-Cas. These results also suggest a role for CRISPR-Cas in gene regulation beyond self-defense.

4. MATERIALS AND METHODS

Bacteria and growth conditions. Bacteria (**Table S4**) were cultured routinely in lysogeny broth (Bertani, 1951) at 37°C, while M9 glucose (0.4 wt%) (Rodriguez and Tait, 1983) was used to assay the growth rate and to resuscitate persister cells. pCA24N-based plasmids (Kitagawa et al., 2005) were retained in overnight cultures via chloramphenicol (30 μg/mL), and kanamycin (50 μg/mL) was used for deletion mutants, where applicable. The BW25113 *cas2* and spacer deletion were confirmed by PCR (primers shown in **Table S5**). Spacers 3 and 12 were deleted from BW25113, and *cas2* was deleted from the strain that lacks cryptic prophages (Δ9) using the method of Datsenko and Wanner (Datsenko and Wanner, 2000) using plasmids pKD4 with the primers in **Table S5**; kanamycin resistance was removed using pCP20 (Datsenko and Wanner, 2000).

Persister cells. Exponentially-growing cells (turbidity of 0.8 at 600 nm) were converted nearly completely to persister cells (Kim *et al.*, 2018; Kwan *et al.*, 2013) by adding rifampicin (100 μg/mL) for 30 min to stop transcription, centrifuging, and adding LB with ampicillin (100 μg/mL) for 3 h to lyse non-persister cells. To remove ampicillin, cells were washed twice with 0.85% NaCl then re-suspended in 0.85% NaCl. Persister concentrations were enumerated via a drop assay (Donegan et al., 1991).

Single-cell persister resuscitation. Persister cells (5 μL) were added to 1.5% agarose gel pads containing M9 glucose (0.4 wt%) medium (Rodriguez and Tait, 1983), and single-

cell resuscitation was visualized at 37°C via a light microscope (Zeiss Axio Scope.A1, bl_ph channel at 1000 ms exposure). For each condition, at least two independent cultures were used with 150 to 300 individual cells assessed per culture.

Membrane integrity assay. To determine membrane integrity, the persister cells were analyzed with the LIVE/DEAD BacLight Bacterial Viability Kit (Molecular Probes, Inc., Eugene, OR, catalog number L7012). The fluorescence signal was analyzed via a Zeiss Axioscope.A1 using excitation at 485 nm and emission at 530 nm for green fluorescence and using excitation at 485 nm and emission at 630 nm for red fluorescence.

qRT-PCR. To quantify transcription from the cryptic prophage lytic genes, RNA was isolated from persister cells that were resuscitated by adding M9 glucose (0.4 wt%) medium for 10 min and from exponential cells grown to a turbidity of 0.8. For quantifying transcription from the CRISPR array, RNA was isolated from persister cells that were resuscitated by adding M9 glucose (0.4 wt%) medium for 10 min. Samples were cooled rapidly using ethanol/dry ice in the presence of RNA Later. RNA was isolated using the High Pure RNA Isolation Kit (Roche). The following qRT-PCR thermocycling protocol was used with the iTaq™ universal SYBR® Green One-Step kit (Bio-Rad): 95 ºC for 5 min; 40 cycles of 95 ºC for 15 s, 60 ºC for 1 min for two replicate reactions for each sample/primer pair. The annealing temperature was 60ºC for all primers (**Table S5**).

qPCR. To quantify prophage excision and the levels of DNA flanking the CRISPR-Cas cleavage sites, total DNA (100 ng) was isolated from exponentially-growing and persister resuscitating cells using an UltraClean Microbial DNA Isolation Kit (Mo Bio Laboratories). Excised cryptic prophage was quantified using primers for each prophage excisionase (**Table S5**) that only yield a PCR product upon prophage excision, and the relative amount of each target gene was determined using reference gene *purM*. The level of cryptic prophage flanking the CRISPR-Cas cleave site was quantified using primers that flank each site (**Table S5**). The qPCR reaction performed using CFX96 Real Time System. The reaction and analysis was conducted using the StepOne Real-Time PCR System (Bio-Rad).

Transmission electron microscopy. For transmission electron microscopy (TEM), the samples were prepared from persister cells that were resuscitated by adding M9 glucose (0.4%) medium for 10 min then washed with 0.85% NaCl. The samples were fixed with buffer (2.5% glutaraldehyde in 0.1M cacodylate buffer, pH 7.4). The negative staining was performed with 2% uranyl acetate for 1 hr, then dehydrated. The sectioned specimens were stained again with uranyl acetate and lead citrate after dehydration and embedded with resin. The TEM image was observed using a Hitachi (H-7605) instrument.

ACKNOWLEDGEMENTS: This work was supported by funds derived from the Biotechnology Endowed Professorship at the Pennsylvania State University for TKW and from the National Research Foundation of Korea (NRF) grant from the Korean Government (NRF-2020R1F1A1072397) for SYS. We appreciate the feedback of Joy Muthami. The authors have no conflicts of interest.

REFERENCES

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K.A., Tomita, M., Wanner, B.L., and Mori, H. (2006). Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol. Syst. Biol. *2*, 2006 0008.

Barrangou, R., and Marraffini, Luciano A. (2014). CRISPR-Cas Systems: Prokaryotes Upgrade to Adaptive Immunity. Molecular Cell *54*, 234-244.

Bertani, G. (1951). Studies on Lysogenesis .1. The Mode of Phage Liberation by Lysogenic *Escherichia-Coli*. J Bacteriol *62*, 293-300.

Bozic, B., Repac, J., and Djordjevic, M. (2019). Endogenous Gene Regulation as a Predicted Main Function of Type I-E CRISPR/Cas System in E. coli. Molecules *24*, 784.

Brouns, S.J.J., Jore, M.M., Lundgren, M., Westra, E.R., Slijkhuis, R.J.H., Snijders, A.P.L., Dickman, M.J., Makarova, K.S., Koonin, E.V., and van der Oost, J. (2008). Small CRISPR RNAs Guide Antiviral Defense in Prokaryotes. Science *321*, 960-964.

Cheng, H.-Y., Soo, V.W.C., Islam, S., McAnulty, M.J., Benedik, M.J., and Wood, T.K. (2014). Toxin GhoT of the GhoT/GhoS toxin/antitoxin system damages the cell membrane to reduce adenosine triphosphate and to reduce growth under stress. Environmental Microbiology *16*, 1741-1754.

Datsenko, K.A., Pougach, K., Tikhonov, A., Wanner, B.L., Severinov, K., and Semenova, E. (2012). Molecular memory of prior infections activates the CRISPR/Cas adaptive bacterial immunity system. Nature Communications *3*, 945.

Datsenko, K.A., and Wanner, B.L. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. Proc. Natl. Acad. Sci. USA *97*, 6640-6645.

Donegan, K., Matyac, C., Seidler, R., and Porteous, A. (1991). Evaluation of methods for sampling, recovery, and enumeration of bacteria applied to the phylloplane. Appl Environ Microbiol *57*, 51-56.

Edgar, R., and Qimron, U. (2010). The *Escherichia coli* CRISPR System Protects from λ Lysogenization, Lysogens, and Prophage Induction. Journal of Bacteriology *192*, 6291-6294.

García Contreras, R., Zhang, X.-S., Kim, Y., and Wood, T.K. (2008). Protein Translation and Cell Death: The Role of Rare tRNAs in Biofilm Formation and in Activating Dormant Phage Killer Genes. PLoS ONE *3*, e2394.

Goldberg, G.W., McMillan, E.A., Varble, A., Modell, J.W., Samai, P., Jiang, W., and Marraffini, L.A. (2018). Incomplete prophage tolerance by type III-A CRISPR-Cas systems reduces the fitness of lysogenic hosts. Nature Communications *9*, 61.

Hafeezunnisa, M., Chhakchhuak, P.I.R., Krishnakumar, J., and Sen, R. (2021). Rho-dependent transcription termination regulates the toxin–antitoxin modules of cryptic prophages to silence their expression in *Escherichia coli*. FEBS Lett *15*, 2057-2067.

Howard-Varona, C., Hargreaves, K.R., Abedon, S.T., and Sullivan, M.B. (2017). Lysogeny in nature: mechanisms, impact and ecology of temperate phages. The ISME Journal *11*, 1511-1520.

Ishihama, A., and Shimada, T. (2021). Hierarchy of transcription factor network in *Escherichia coli* K-12: H-NS-mediated silencing and Anti-silencing by global regulators. FEMS Microbiology Reviews.

Jore, M.M., Lundgren, M., van Duijn, E., Bultema, J.B., Westra, E.R., Waghmare, S.P., Wiedenheft, B., Pul, Ü., Wurm, R., Wagner, R., et al. (2011). Structural basis for CRISPR RNA-guided DNA recognition by Cascade. Nature Structural & Molecular Biology *18*, 529-536.

Kim, J.-S., Yamasaki, R., Song, S., Zhang, W., and Wood, T.K. (2018). Single Cell Observations Show Persister Cells Wake Based on Ribosome Content. Environ Microbiol *20*, 2085-2098.

Kitagawa, M., Ara, T., Arifuzzaman, M., Ioka-Nakamichi, T., Inamoto, E., Toyonaga, H., and Mori, H. (2005). Complete set of ORF clones of *Escherichia coli* ASKA library (A Complete Set of *E. coli* K-12 ORF Archive): Unique Resources for Biological Research. DNA Res *12*, 291-299.

Kwan, B.W., Valenta, J.A., Benedik, M.J., and Wood, T.K. (2013). Arrested protein synthesis increases persister-like cell formation. Antimicrob Agents Chemother *57*, 1468-1473.

Li, M., Gong, L., Cheng, F., Yu, H., Zhao, D., Wang, R., Wang, T., Zhang, S., Zhou, J., Shmakov, S.A., et al. (2021). Toxin-antitoxin RNA pairs safeguard CRISPR-Cas systems. Science *372*, eabe5601.

Li, R., Fang, L., Tan, S., Yu, M., Li, X., He, S., Wei, Y., Li, G., Jiang, J., and Wu, M. (2016). Type I CRISPR-Cas targets endogenous genes and regulates virulence to evade mammalian host immunity. Cell Res *26*, 1273-1287.

Makarova, K.S., Wolf, Y.I., Iranzo, J., Shmakov, S.A., Alkhnbashi, O.S., Brouns, S.J.J., Charpentier, E., Cheng, D., Haft, D.H., Horvath, P., et al. (2020). Evolutionary classification of CRISPR–Cas systems: a burst of class 2 and derived variants. Nature Reviews Microbiology *18*, 67-83.

Marimon, O., Teixeira, J.M.C., Cordeiro, T.N., Soo, V.W.C., Wood, T.L., Mayzel, M., Amata, I., García, J., Morera, A., Gay, M., et al. (2016). An oxygen-sensitive toxin–antitoxin system. Nature Communications *7*, 13634.

Masuda, H., Awano, N., and Inouye, M. (2016). ydfD encodes a novel lytic protein in *Escherichia coli*. FEMS Microbiology Letters *363*.

Meeske, A.J., Nakandakari-Higa, S., and Marraffini, L.A. (2019). Cas13-induced cellular dormancy prevents the rise of CRISPR-resistant bacteriophage. Nature *570*, 241-245.

Militello, K.T., and Lazatin, J.C. (2017). Discovery of *Escherichia coli* CRISPR sequences in an undergraduate laboratory. Biochem Molec Biol Edu *45*, 262-269.

Mohanraju, P., Saha, C., van Baarlen, P., Louwen, R., Staals, R.H.J., and van der Oost, J. (2022). Alternative functions of CRISPR–Cas systems in the evolutionary arms race. Nature Reviews Microbiology.

Pecota, D.C., and Wood, T.K. (1996). Exclusion of T4 Phage by the *hok/sok* Killer Locus from Plasmid R1. J Bacteriol *178*, 2044-2050.

Perna, N.T., Plunkett, G., Burland, V., Mau, B., Glasner, J.D., Rose, D.J., Mayhew, G.F., Evans, P.S., Gregor, J., Kirkpatrick, H.A., et al. (2001). Genome sequence of enterohaemorrhagic *Escherichia coli* O157: H7. Nature *409*, 529-533.

Pfaffl, M.W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res. *29*.

Pougach, K., Semenova, E., Bogdanova, E., Datsenko, K.A., Djordjevic, M., Wanner, B.L., and Severinov, K. (2010). Transcription, processing and function of CRISPR cassettes in *Escherichia coli*. Mol Microbiol *77*, 1367-1379.

Rodriguez, R.L., and Tait, R.C. (1983). Recombinant DNA Techniques: An Introduction (Benjamin/Cummings Publishing).

Rollie, C., Chevallereau, A., Watson, B.N.J., Chyou, T.-y., Fradet, O., McLeod, I., Fineran, P.C., Brown, C.M., Gandon, S., and Westra, E.R. (2020). Targeting of temperate phages drives loss of type I CRISPR–Cas systems. Nature.

Savitskaya, E., Lopatina, A., Medvedeva, S., Kapustin, M., Shmakov, S., Tikhonov, A., Artamonova, II, Logacheva, M., and Severinov, K. (2017). Dynamics of *Escherichia coli* type I-E CRISPR spacers over 42 000 years. Molecular ecology *26*, 2019-2026.

Shmakov, S.A., Sitnik, V., Makarova, K.S., Wolf, Y.I., Severinov, K.V., and Koonin, E.V. (2017). The CRISPR Spacer Space Is Dominated by Sequences from Species-Specific Mobilomes. mBio *8*, e01397-01317.

Song, S., Kim, J.-S., Yamasaki, R., Oh, S., Benedik, M.J., and Wood, T.K. (2021). *Escherichia coli* cryptic prophages sense nutrients to influence persister cell resuscitation. Environ Microbiol *23*, 7245-7254.

Song, S., and Wood, T.K. (2020). Persister Cells Resuscitate via Ribosome Modification by 23S rRNA Pseudouridine Synthase RluD. Environ. Microbiol. *22*, 850–857.

Song, S., and Wood, T.K. (2021). Are We Really Studying Persister Cells? Environmental Microbiology Reports *13*, 3–7.

Touchon, M., Charpentier, S., Clermont, O., Rocha, E.P.C., Denamur, E., and Branger, C. (2011). CRISPR Distribution within the *Escherichia coli* Species Is Not Suggestive of Immunity-Associated Diversifying Selection. Journal of Bacteriology *193*, 2460-2467.

Vasu, K., and Nagaraja, V. (2013). Diverse Functions of Restriction-Modification Systems in Addition to Cellular Defense. Microbiology and Molecular Biology Reviews *77*, 53-72.

Wang, X., Kim, Y., Ma, Q., Hong, S.H., Pokusaeva, K., Sturino, J.M., and Wood, T.K. (2010). Cryptic prophages help bacteria cope with adverse environments. Nat Commun *1*, 147.

Wang, X., Kim, Y., and Wood, T.K. (2009). Control and Benefits of CP4-57 Prophage Excision in *Escherichia coli* Biofilms. ISME J *3*, 1164-1179.

Wang, X., Lord, D.M., Cheng, H.-Y., Osbourne, D.O., Hong, S.H., Sanchez-Torres, V., Quiroga, C., Zheng, K., Herrmann, T., Peti, W., et al. (2012). A Novel Type V TA System Where mRNA for Toxin GhoT is Cleaved by Antitoxin GhoS. Nat. Chem. Biol. *8*, 855-861.

Wood, T.K., and Song, S. (2020). Forming and waking dormant cells: The ppGpp ribosome dimerization persister model. Biofilm *2*, 100018.

Yamasaki, R., Song, S., Benedik, M.J., and Wood, T.K. (2020). Persister Cells Resuscitate Using Membrane Sensors that Activate Chemotaxis, Lower cAMP Levels, and Revive Ribosomes. iScience *23*, 100792.

Supporting Information

Sooyeon Song[1,2,3*]**, Ekaterina Semenova**[4]**, Konstantin Severinov**[4]**, Laura Fernández-García**[1]**, Michael J. Benedik**[5]**, Toshinari Maeda**[6]**, and Thomas K. Wood**[1*]

[1]Department of Chemical Engineering, Pennsylvania State University,
University Park, Pennsylvania, 16802-4400, USA

Departments of Animal Science[2] and Agricultural Convergence Technology[3], Jeonbuk National University, 587 Baekje-Daero, Deojin-Gu, Jeonju-Si, Jellabuk-Do, 54896, South Korea

[4]Waksman Institute of Microbiology, Rutgers, The State University of New Jersey, Piscataway, NJ 08854

[5]Hamad bin Khalifa University, Education City, Doha, Qatar

[6]Department of Biological Functions Engineering, Kyushu Institute of Technology, Kitakyushu, Japan
808-0196

*For correspondence. E-mail songsy@jbnu.ac.kr, tuw14@psu.edu

Tel. (+)1 814-863-4811; Fax (1) 814-865-7846

Table S1. Inactivating CRISPR-Cas eliminates persister cell resuscitation on glucose agarose gel pads. Single persister cells were observed using light microscopy (Zeiss Axio Scope.A1). The total number and waking number of persister cells are shown after 6 hours on 0.4 wt% glucose minimal medium. Fold-change in waking is relative to BW25113. These results are the combined observations from two independent experiments (independent culture results separated by "/"), and standard deviations are shown. Sample microscope images are shown in **Fig. 1**.

	Total cells	**Waking cells**	**% waking**	**Fold-change**
BW25113	504/382	199/248	52 ± 18	1
$\Delta cas2$	516/677	6/15	1.7 ± 0.7	-30.9
Δspacer	323/174	10/7	3.6 ± 0.7	-14.6

Table S2. Inactivating CRISPR-Cas causes cell death and ghost cell formation. For resuscitated cells, persister cells were washed with PBS twice, resuscitated by M9 0.4% glucose for 10 min and stained with LIVE/DEAD reagents. Exponential cells were grown to a turbidity of 0.8 (at 600 nm) and stationary cells were grown to turbidity of 2.0. The ghost cells in the persister population were visualized using a Zeiss Axioscope.A1microscope. The results are the combined observations from two independent experiments (independent culture results separated by "/"). The microscope images are shown in **Fig. 1AB** and **Fig. S6**.

	Strains	Total cells	Dead cells	Ghost cells	% Ghost cells	fold-change ghost cells	% dead	fold-change dead cells
Resuscitated	**BW25113**	52/115	2/7	0/1	0.6	1	5.4	1
	Δ*cas2*	142/106	54/38	38/13	20.6	34	37.1	6.9
	Δspacer	161/221	76/76	-	-	-	37.8	7.0
Exponential	**BW25113**	153/147	0/2	-	-	-	0.7	1
	Δ*cas2*	59/40	½	-	-	-	3.4	4.9
	Δspacer	95/146	1/5				2.1	3.15
Stationary	**BW25113**	492/123	1/0				0.1	1
	Δ*cas2*	1136/1736	14/18	-	-	-	1.13	11.2
	Δspacer	200/113	19/8				8.26	81.3
	Δ*casE*	208/175	9/10	-	-	-	5.02	49.4
	Δ9 Δ*cas2*	833/1541	10/13	-	-	-	1.02	
	Δspacer3	136/297	8/73	-	-	-	15.23	
	Δspacer12	513/396	213/92	-	-	-	32.38	

Table S3. CRISPR-Cas does not affect cryptic prophage excision in stationary cells but produces the CRISPR array. (**A**) Fold changes in excision are relative to the wild-type strain as determined by qPCR. (**B**) Fold changes in the CRISPR array are relative to the wild-type strain as determined by qRT-PCR (the Δspacer strain was used as a negative control). Cycle numbers (C_t) are indicated for each sample including that for the target genes as well as that of the house-keeping gene, *purM*, which was used to normalize the data. *cas2* served to inactivate crRNA production due to a polar mutation. Fold changes in the transcription of various targets (i.e., cryptic prophage CP4-57, e14, DLP-12, Spacer3, and Spacer12) with and without $cas2^+$ ($cas2^+/cas2^-$) were calculated as described earlier (Pfaffl, 2001):

$$2^{\wedge}\text{-}(C_{t\ target\_cas2+}\text{-}C_{t\ purM\_cas2+})/2^{\wedge}\text{-}(C_{t\ target\_cas2-}\text{-}C_{t\ purM\_cas2-})$$

(A) qPCR

Gene	*purM*		CP4-57		e14		DLP-12	
Strain	WT	*cas2*	WT	*cas2*	WT	*cas2*	WT	*cas2*
C_T	9.30 ± 0.13	9.70 ± 0.18	30.45 ± 1.73	30.37 ± 0.68	16.30 ± 0.06	17.45 ± 0.51	30.58 ± 0.49	30.05 ± 0.5
ΔC_T			21.16 ± 1.73	20.67 ± 0.70	7.00 ± 0.15	8.16 ± 0.54	21.29 ± 0.51	20.35 ± 0.53
$\Delta\Delta C_T$			-0.48 ± 0.7		1.16 ±0.54		-0.94 ±0.53	
fold			1.40		-2.23		1.92	

(B) qRT-PCR

Gene	*purM*			Spacer3			Spacer12		
Strain	WT	*cas2*	spacer	WT	*cas2*	spacer	WT	*cas2*	spacer
C_T	25.3 ± 0.5	22.2 ± 0.5	22.0 ± 0.2	30.68 ± 1.73	24.35 ± 0.93	-	33.60 ± 1.20	27.12 ± 1.22	-
ΔC_T				5.37 ± 1.08	2.19 ± 1.1	-	8.28 ± 1.31	4.96 ± 1.32	-
$\Delta\Delta C_T$				-3.20 ± 1.51		-	-3.33 ±1.86		-
fold				9.11	-∞		10.02	-∞	

Table S4. *E. coli* **bacterial strains and plasmids utilized.**

Strains and Plasmids	Features	Source
Strains		
BW25113	*rrnB3 ΔlacZ4787 hsdR514 Δ(araBAD)567 Δ(rha-BAD)568 rph-1*	(Baba et al., 2006)
BW25113 Δ*cas1*	Δ*cas1*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*cas2*	Δ*cas2*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*cas2* ΔKm^R	Δ*cas2*, ΔKm^R	this work
BW25113 Δ*cas3*	Δ*cas3*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*casA*	Δ*casA*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*casB*	Δ*casB*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*casC*	Δ*casC*, ΩMr.	(Baba *et al.*, 2006)
BW25113 Δ*casD*	Δ*casD*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δ*casE*	Δ*casE*, ΩKm^R	(Baba *et al.*, 2006)
BW25113 Δspacer (also called BW39292)	Δ(13 spacer region), ΩKm^R	(Pougach et al., 2010)
BW40114	*placuv5::cas3 paraB8p::casABCDE12spacers*	(Datsenko et al., 2012)
BW25113 Δspacer3	Δ(spacer region 3), ΔKm^R	this work
BW25113 Δspacer12	Δ(spacer region 12), ΔKm^R	this work
BW40114	BW25113 *lac*UV5::*cas3 araB*8p::*casABCDE12*	(Datsenko *et al.*, 2012)
BW15113 Δ9	BW15113 lacking all 9 cryptic prophages	(Wang *et al.*, 2010)
BW15113 Δ9 *cas2*	BW15113 Δ9 *cas2*, ΔKm^R	this work
ssrA	*ssrA*, ΩKm^R	(Wang et al., 2009)
Plasmids		
pCA24N	Cm^R; *lacI^q*	(Kitagawa et al., 2005)
pCA24N_*cas2*	Cm^R; *lacI^q*, P$_{T5-lac}$::*cas2$^+$*	(Kitagawa et al., 2005)
pKD4	FRT::Kan^R::FRT, Amp^R	(Datsenko and Wanner, 2000)
pCP20	FLP synthase, Amp^R	(Datsenko and Wanner, 2000)

Table S5. Primers used in this study for qRT-PCR and qPCR. * indicates excision primers.

Gene	Sequence (5' to 3')
qRT-PCR	
purM	**F:** CGGTGTTGATATTGACGCGGG
	R: CAGCACGGGTTCACGATATTTTG
*ydfD (*Qin*)*	**F:** TCAGCATTTGTGCTTGTTCTG
	R: CTGCCGGGATTTCGATATTA
*hokD (*Qin*)*	**F:** GCCCTGATCGTCATCTGTTT
	R: AGCTGTGAAGACAGCGACCT
*ypjF (*CP4-57*)*	**F:** TATGGCCTGACACTGAACGA
	R: GTTGTTGCCACGAAAATCCT
rzoD (DLP-12)	**F:** TGCTCTGCGTGATGATGTTG
	R: TCTCTCTGAGGGTGAAATAATCC
essD (DLP-12)	**F:** CTCACAGTGGGCAGCAATAG
	R: TATTCACCTCTCGCAGCCTT
Spacer3	**F:** GATAAACCGCTTTCGCAGAC
	R: AGAAATTCCAGACCCGATCC
Spacer12	**F:** AACGCCTGAACCGAGAGTT
	R: CTGGGAGTTCTACCGCAGAG
qPCR	
*CP4-57	**F:** AAGCATGTAGTACCGAGGATGTAGG
	R: TATGTCTCCTCACCGTCTGGTCGG
*e14	**F:** GTGCAAACATCGGTGACGAA
	R: TTCAGCAGCTTAGCGCCTTC
*DLP-12	**F:** CAAAAGCCATTGACTCAGCAAGG
	R: CGGATAAGACGGGCATAAATGA
purM 400	**F:** CTGATTGCACTCGGTTCCAG
	R: CGTTTTCACCGTTGGCATTG
Spacer 3-1	**F:** CATGCAATTACAACATCAGGGTA
	R: GCCATTGGTAAAACCTTCCA
Spacer 3-2	**F:** AAATTCCGAAAAAGCTCCTGA
	R: TGCTTTCAAGATTATGGCGTA
Spacer 6	**F:** CACGAAAGCCAGCCTATTCC
	R: CCGCTGTTTCTTTCTCCAGG
Spacer 8	**F:** GCCAGCATAATGAGATCGGC
	R: TTTTACCCAAACTCAGCGCG
Spacer 11	**F:** TGTGCAGTTGTACCAGTGGA
	R: CAACCCAGCAAAGTTTCGGA
Spacer 12-1	**F:** TGTATGTGCCCCGGTGTTAT
	R: GCAGATGAAGGCGCATTACA
Spacer 12-2	**F:** AGCTTTACACCTCGGCTCAT
	R: CCGGAACTCTTGTGTTGGTG
Spacer 12-3	**F:** TCTACAGGGAAAGGACGACC
	R: CTCTGCAACCAAAGTGAACCA
Spacer 13	**F:** ACAACCACTATCGCCCCTTT
	R:GGTAAGGCTGCATTGGGAAG
Deletions	

cas2	**F**:	ACCTTAATGTAACATTTCCTTATTATTAAAGATCAGCTAATTCTTT-GTTTGT
		GTAGGCTGGAGCTGCTTC
	R: CATTCCGCTTCCTGTTTCACTGGGAGATGCAGGCCATCGGAGTAGCTGA	
	ACATATGAATATCCTCCTTAG	
Spacer 3	**F**: GCTGGCGCGGGGAACTCGCGACCGCTCAGAAATTCCAGACCCGATCCAA	
	AGTGTAGGCTGGAGCTGCTTC	
	R: AAAGGTGATGCCGAACACGCTGAGTTCCCCGCGCCAGCGGGGATAAACC	
	GCATATGAATATCCTCCTTAG	
Spacer 12	**F**: CAGAGGCGGGGGAACTCCAAGTGATATCCATCATCGCATCCAGTGCGCC	
	CGTGTAGGCTGGAGCTGCTTC	
	R: GGTTTGCAACGCCTGAACCGAGAGTTCCCCGCGCCAGCAGGGATAAACC	
	GCATATGAATATCCTCCTTAG	

Verification for *cas2* and Spacer

cas2	**F:** GAAAATGTACCTCCGCGCTT	
	R: CCATCCAAATCTACCGGGGT	
Spacer	**F:** GGATCGGGTCTGGAATTTCT	
	R: GGGGAACTCGTAGTCCATCA	

Supplementary Figure 1. Growth curves for △*cas1*, △*cas2*, *cas3*, △*casA*, △*casB*, △*casC*, △*casD*, and △*casE* in M9 glucose medium at 37ºC. Turbidity at 600 nm shown, and error bars indicate standard deviations from two independent cultures.

(A)

(B)

Supplementary Figure 2. Removing the kanamycin marker from the *E. coli Δcas2 ΩkanR* strain to create the *Δcas2 ΔkanR* strain **(A)** increased growth in M9 glucose medium at 37°C (turbidity at 600 nm shown and error bars indicate standard deviations from two independent cultures) and **(B)** reduced toxicity via the LIVE/DEAD assay for stationary cells showing the polar mutation due to the kanamycin marker in the *Δcas2 ΩkanR* strain inhibits spacer formation, which results in reduced growth and increased toxicity. Representative images shown from two independent cultures.

BW25113

Δ*cas2*

Δspacer

BW40114

Supplementary Figure 3. Schematics of the CRISPR-Cas strains used in this study. All the strains are derivatives of isogenic host, BW25113 (see **Table S4**).

BW25113　　　　　　　**Δ*ssrA***

Supplementary Figure 4. As a control for growth rate and resuscitation, the *ssrA* mutation was checked and found to have no effect on persister resuscitation on LB plates although it grows 22% more slowly than the wild-type strain (1.18 ± 0.07/h vs. 1.49 ± 0.03/h, respectively). The number of *ssrA* persisters that resuscitated is 101 ± 2 vs. 100.0 ± 0.1 for the wild-type. Plates were incubated for 24 h at 37°C, and representative images shown from two independent cultures.

Supplementary Figure 5. CRISPR-Cas prevents cell lysis induced by cryptic phages. (**A**) Single cell persister waking of BW25113 Δ*cas2* and the Δspacer mutant on M9 0.4% glucose agar plates incubated at 37°C for 4 hours. The scale bar indicates 10µm. (**B**) TEM image for persister waking of BW25113 Δ*cas2* and the Δspacer mutant. Persister cells were resuscitated by M9 0.4% glucose for 10 min. One representative image from two independent cultures is shown. The scale bar indicates 1 µm.

Supplementary Figure 6. LIVE/DEAD staining of stationary (turbidity 2.0) and exponential (turbidity 0.8) cells shows the *cas2*, Δspacer (all 13 spacers removed), and mutations cause cell death. DF is dark field, SYTO9 is a membrane permeable stain for nucleic acids (green), and PI is propidium iodide, which is a membrane impermeable stain for the nucleic acids of dead cells (red). Tabulated cell numbers are shown in **Table S2**, and representative images from two independent cultures shown.

Supplementary Figure 7. Growth curves upon inducing the full CRISPR-Cas system in *E. coli* BW40114 (via IPTG and arabinose) at 37°C in LB and M9 glucose media (turbidity at 600 nm shown). The growth is reduced due to the metabolic burden of producing the complete CRISPR-Cas system. Error bars indicate standard deviations from two independent cultures.

yfdR gene (`Start` and `stop` codon)(**CPS-53**, encodes 5'-deoxynucleotidase)

`ATG`TCATTTATTAAAACTTTTTCCGGGAAGCATTTTTATTATGACAAGATAAATAAAGACGACATCGTGATTAACGA
TATCGCGGTTTCCCTTTCAAATATCTGCCGCTTTGCCGGTCATCTTTCTCACTTCTACAGTGTCGCCCAACATGCGG
                                                            `Spacer 8` 3'-ACCCGCCGA
TGCTTTGCAGCCAGCTGGTGCCGCAGGAATTTGCTTTTGAAGCATTAATGCATGATGCAACAGAAGCGTATTGCCAG
|||||||||||||||||
`ACGGAACGTCGGTCGA`GGTCGTC-5'
GACATCCCCGCACCACTGAAACGCCTTCTTCCTGACTATAAACGGATGGAAGAAAAAATAGATGCAGTAATCCGTGA
GAAATACGGGTTACCTCCTGTTATGAGCACGCCAGTGAAATATGCCGATCTCATTATGCTGGCAACCGAACGCCGTG
ATCTCGGGCTTGATGATGGCTCTTTCTGGCCTGTACTGGAAGGCATCCCGGCAACAGAGATGTTCAACGTGATTCCA
CTGGCACCGGGTCATGCCTACGGGATGTTTATGGAACGTTTTAACGATTTATCGGAGTTACGCAAATGCGCA`TGA`

perR gene (`Start` and `stop` codon)(**CP4-6**, encodes transcriptional regulator)

`ATG`AAGCTCTTAGCAAAAGCACCTCTAAATCTGTTACGCGCCTTTGAAGCGGCTGGTCGTACCGGAGCTTTTGCGTT
GGCCGCCTCTGAGTTGGAACTGTCACCCAGTGCGATCAGCCATGCCATCCGCAAACTGGAAAACTTGCTTGATGTAC
GCCTTTTTCAGCGCAGTACGCGAGAAATTACGCTGACGAAAGAAGGTGAAATACTGCTTGAGCACATACAGCGGGGA
TTTAACGAATTACAGCAGGGGTTGGCATTAGTGACGGCTGATGAATCCAGGCCTTTGCGCCTTCATACTGCACCGAG
TTTTGCCCATCAATGGCTTTTTACCACGTCTTGGTAAGTTCATACGTGAGAACCCAAGCATCGATCTGCGACTTTCAG
CCAGCACAGAATATGCACGTTTTGAACAGGATGATTTTGATCTCGATATAGTCTACGGTGAACCCCGCCCATCACCC
TATGAGAAGATCCCGCTTGCTGTTGAAGAACTTACGCCACTGTGTTCTCCCCAACTGGCTGAGCGGCTAAAGAAACC
AGAGGATCTCTATGCGTTGACATTGATTCAGTGCGATGTTGCAGTTGTACCGGTGGAAAGGATGGTTTGAGGCGAATA
AGATGACGCCACCCAATAATTATGGCCTTCGATTTGACCGTAGCTTTATGGCGATTGCTGCCGCAGTTGACGGCTTG
GGAGTTGTGCTTGAATCAAAACTACTGGCTGAACGTGAAATCGCTAGCGGTAAACTGGTATGCCCACTTGTAAACAG
TACCAGCGAAATACATTACATCGGTCATTATTTGGTTTTCCCCCAGCACCAACACATGCATTCTGCGCTTGATGTGT
                                                          // // / / //
                                              `Spacer 11` 3'-AGCCAAGTCCGCA
TCAAAACCTGGCTACTGAATGAACTCAATCTGGGTAAAATTCGT`TGA`
| ||||||||||| |
ACG`TTTGGACCGATG`GCCC-5'

stfR gene (`Start` and `stop` codon)(**Rac**, side tail fiber protein)

`ATG`GCAGTAAAGATTTCAGGTGTACTGAAAGACGGCACAGGAAAACCGGTACAGAACTGCACAATCCAGCTGAAAGC
AAAACGTAACAGCACCACGGTGGTGGTGAACACGCTGGCCTCAGAAAATCCGGATGAAGCCGGGCGTTACAGCATGG
ACGTTGAGTACGGTCAGTACAGCGTTATTCTGTTGGTGGAAGGATTCCCGCCGTCACATGCCGGGACCATTACCGTG
TATGAAGATTCTCAACCCGGTACGCTGAATGATTTTCTCGGTGCCATGACGGAGGATGATGCCCGTCCGGAGGCACT
GCGCCGTTTTGAACTGATGGTGGAAGAGGTGGCGCGTAACGCGTCCGCGGTGGCACAGAACACGGCAGCCGCGAAGA
AGTCAGCCAGTGATGCCAGCACATCAGCCCGTGAGGCGGCAACCCATGCGGCTGATGCTGCGGACTCAGCACGCGCA
GCCAGCACGTCAGCCGGACAGGCCGCGTCGTCGGCTCAGTCAGCGTCTTCCAGCGCAGGAACGGCATCAACAAAGGC
CACTGAAGCATCAAAAAGTGCTGCCGCTGCAGAGTCCTCAAAAAGCGCGGCGGCCACCAGTGCCGGTGCGGCGAAAA
CGTCAGAAACGAATGCTTCAGCGTCACTACAATCAGCAGCCACATCTGCATCCACCGCGACCACGAAGGCATCAGAA
GCTGCGACCTCGGCCCGGGATGCGGCGGCCTCAAAAGAAGCGGCAAAATCATCAGAAACGAACGCATCATCAAGCGC
CAGTAGTGCAGCTTCCTCGGCAACGGCGGGCAGGAAATTCCGCGAAGGCGGCAAAAACGTCCGAGACGAACGCCAGGT
CTTCTGAAACGGCAGCGGGACAGAGCGCCTCGGCTGCGGCAGGCTCAAAAACAGCGGCTGCGTCGTCTGCCAGTGCA
GCGTCAACAAGTGCCGGGCAGGCCTCAGCCAGTGCCACCGCCGCCGGAAAATCGGCAGAAAGCGCCGCATCGTCTGC
TTCAACAGCCACAACGAAGGCTGGCGAAGCCACTGAACAGGCCAGCGCAGCAGCGAGGTCTGCTTCCGCAGCGAAGA
CATCCGAAACGAACGCGAAAGCGTCGGAAACAAGCGCAGAATCCTCAAAAACGGCTGCCGCATCGTCAGCCAGTTCG
GCGGCGTCATCGGCATCATCGGCGTCTGCTTCAAAAGATGAGGCGGACCAGACAAGCGTCAGCAGCGAAGAGCAGCGC
CACGACGGCATCCACGAAGGCGACAGAGGCTGCTGGCAGTGCGACGGCGGCAGCTCAGAGCAAAAGTACGGCGGAAT
CCGCGGCAACGCGCGCCGAGACAGCAGCTAAACGGGCAGAGGATATTGCATCCGCCGTGGCGCTTGAGGATGCAAGT
ACGACGAAAAAGGGGATAGTACAGCTCAGCAGTGCGACCAACAGTACGTCTGAAACGCTGGCGGCAACGCCAAAGGC
AGTAAAATCAGCCTATGACAATGCAGAGAAACGTCTGCAGAAAGACCAGAACGGCGCTGATATACCCGATAAGGGAT
GCTTCCTGAACAACATTAACGCGGTCAGTAAAACAGACTTTGCTGATAAGCGTGGTATGCGTTATGTGCGGGTTAAC
GCTCCTGCAGGTGCAACATCTGGAAAATATTACCCTGTTGTTGTTATGCGTTCTGCTGGCTCAGTAAGCGAACTGGC
ATCAAGAGTCATTATCACCACGGCAACGCGAACCGCAGGCGATCCGATGAATAACTGCGAGTTTAACGGATTTGTTA
TGCCTGGTGGCTGGACTGACAGGGGGCGTTATGCTTATGGCATGTTCTGGCAATATCAAAACAATGAACGAGCCATT
CACTCAATAATGATGAGTAATAAGGGCGATGATTTGCGCTCTGTGTTCTATGTTGATGGCGCTGCTTTCCCTGTTTT
TGCGTTTATTGAAGATGGCCTGTCAATATCCGCACCTGGTGCTGATCTCGTTGTTAATGATACGACCTATAAGTTTG
GGGCAACAAATCCGGCGACTGAATGTATCGCGGCGGACGTTATCCTTGATTTTAAGAGTGGGCGTGGTTTTTATGAG
TCTCATTCGTTAATCGTTAACGATAACTTGTCGTGCAAAAAACTTTTTGCCACAGACGAAATTGTAGCGCGTGGTGG
TAATCAGATTCGAATGATAGGTGGGGAGTATGGTGCATTATGGCGTAATGATGGCGCTAAAACTTACCTGCTGCTTA
CCAATCAAGGTGATGTTTATGGTGGCTGGAATACATTAAGACCGTTTGCTATTGATAACGCAACCGGCGAACTGGTT
ATTGGAACCAAACTGTCCGCAAGTCTGAACGGTAATGCATTAACAGCAACAAAGCTGCAAACGCCAAGACGGGTTTC
TGGTGTTGAGTTTGATGGTTCCAAAGATATTACTTTAACCGCCGCGCATGTGGCTGGCTGCTTTTGCCAGAAGGGCAACGG
ATACATATGCCGATGCGGATGGTGGCGTTCCATGGAATGCCGAATCTGGCGCTTACAATGTCACCCGCTCTGGCGAC
AGCTATATTCTGGTTAACTTCTATACCGGAGTCGGAAGTTGCCGGACCCTGCAGATGAAGGCGCATTACAGAAATGG
TGGTCTGTTCTACCGTTCTTCAAGAGACGGTTATGGTTTTGAGGAAGACTGGGCAGAAGTTTATACCTCGAAAAATC
TTCCACCAGAAAGCTACCCAGTCGGCGCACCAATCCCGTGGCCATCAGATACCGTTCCGTCTGGTTATGCCCTGATG
CAGGGGCAGGCTTTTGACAAATCTGCTTACCCGAAACTTGCAGCCGCTTATC`CGTCAGGCGTGAT`CCCTGATATGCG
                                                     |||||||||||| ||| ||
                                         `Spacer 12` 3'-GTC`GCAGTCCGCACT`TTAGAGTGGCAGC
AACG-5'
| ||
TGGCTGGACGATTAAGGGCAAACCTGCCAGTGGTCGGGCCGTATTGTCTCAGGAACAGGACGGCATTAAATCGCATA
CCCACAGCGCCAGCGCATCCAGTACAGATTTGGGGACGAAAACCACATCGTCGTTTGATTACGGCACTAAATCCACG
AATAACACCGGGGCACATACACACAGTGTGAGCGGCTCTACAAACTCGGCTGGAGCACACACACACTCACTAGCCAA
CGTGAACACGGCTAGTGCTAACTCCGGTGCTGGTAGTGCATCAACAAGATTGTCTGTTGTGCATAATCAAAACTATG
CAACATCATCTGCTGGCGCACATACCCACTCACTGTCCGGCACTGCTGCAAGCGCAGGTGCACACGCGCATACTGTC
GGTATTGGTGCTCATACGCACTCCGTTGCGATTGGTTCACATGGACACACCATCACCGTTAACGCTGCTGGTAACGC
GGAAAACACCGTCAAAAACATCGCATTTAACTATATTGTGAGGCTTGCA`TAA`

stfQ gene (<mark>Start</mark> and <mark>stop</mark> codon) (**Qin**, side tail fiber protein)

<mark>GTG</mark>AATATAACCGCGCTGACAGACAATACGCAGGGGGCAGCAGGTCTTGAGTTATACGAGGTGTATAACAACGGATA
TCCAACAGCGTATGGAAATATCATTCACCTGAAAGGGATGACAGCCGTTGGCGAAGGTGAGTTACTCATCGGCTGGA
GTGGTACAAGCGGTGCTCATGCTCCGGCATTTATTCGTTCACGACGGGATACGACCGACGCAAACTGGTCGCCGTGG
GCGCAGCTTTACACCTCGGCTCATCCTCCTGCAGAGTTTTATCCAGTCGGTGCACCAATCCCGTGGCCATCAGATAC
CGTTCCGTCTGGTTATGCCCTGATGCAGGGGCAGACTTTTGACAAATCTGCATACCCGAAACTTGCAGTTGCTTATC
                                                    <mark>Spacer 12</mark> 3'– GTC
CGTCAGGCGTGATCCCTGATATGCGTGGCTGGACGATTAAGGGCAAGCCCGCCAGTGGTCGGGCCGTATTATCTCAG
    |||||||||||| || | ||| ||
<mark>GCAGTCCGCACT</mark>TTAGAGTGGCAGCAACG–5'
GAACAGGACGGCATTAAATCGCACACCCACAGCGCCAGCGCATCCAGTACGGATTTGGGGACGGAAACCACATCGTC
GTTTGATTACGGAACCAAATCCACGAATAACACCGGGGCGCATACCCATAGTATTAGCGGGACCGCAAATAGTGCCG
GTGCGCACCAACACAAGAGTTCCGGTGCATTTGGTGGCACGAACACGAGCATTTTCCCTAATGGTTATACCGCGATT
TCAAATCTAAGCGCGGGGATTATGAGCACAACAAGCGGTAGTGGCCAGACTCGTAATGCAGGGAAGACATCATCAGA
TGGTGCTCATACCCACTCGCTGTCCGGCACTGCTGCAAGCGCAGGCGCGCATGCACATACTGTCGGTATTGGTGCTC
ATACGCACTCCGTTGCGATTGGTTCACATGGACACACCATCACCGTTAACGCTGCTGGTAACGCGGAAAACACCGTC
AAAAACATCGCATTTAACTATATTGTGAGGCTTGCA<mark>TAA</mark>

alpA gene (`Start` and `stop` codon) (**CP4-57**, transcriptional regulator)

```
ATGAACAATCGATCGGCCGTTAGAATACTACGGTTACCAGCGGTTATCCAAAAAACAGGTATGGCACGGGCCACCAT
CTATGACTGGTTGAACCCCAAATCACCACGATACGATGCCCACCTTTCCCAAAAAGCGAATGCTCGGCGTGAAATCTG
                                                      |  |  ||||||||||||
                               Spacer 12 3'-GTCGCAGTCCGCACTTTAGAG
TGGCAGCAACG-5'
 |  |  |
TCGGATGGATTGAGGCCGAGATTGATGAGTGGTTATCACAACGCTGTAAACTTATTTGA
```

ybcN gene (`Start` and `stop` codon) (**DLP-12**, DNA base-flipping protein)

```
ATGCGCACATACAATCCAAACTCTCTTCTCCCTTCACAGATGCAGAAATGCACCTGCAATTCTTTGCATCTAGCGTT
TGACCTCTGCGGAGGGGAAGCGTGAACCTCTCACAAGACGGCATCAAATTACATCGCGGCAACTTCACCGCTATCGG
                      |||||  |   |       |
           Spacer 12 3'-GTCGCAGTCCGCACTTTAGAGTGGCAGCAACG-5'
TCGGCAGATCCAGCCTTATCTGGAGGAGGGCAAATGCTTTCGCATGGTGCTTAAACCGTGGCGTGAGAAACGCAGTC
TTTCCCAGAATGCACTCAGCCACATGTGGTACAGCGAAATCAGTGAATACCTCATCAGCAGGGGTAAAACGTTCGCC
ACTCCAGCTTGGGTAAAAGATGCTCTCAAACACACATATCTCGGTTATGAAACCAAAGACCTGGTTGATGTCGTAAC
CGGTGATATCACCACTATCCAGTCGTTACGCCATACCTCCGATCTTGATACCGGAGAGATGTATGTCTTCCTGTGTA
AGGTTGAAGCCTGGGCGATGAATATTGGTTGCCACCTGACTATTCCACAGAGCTGCGAGTTCCAGCTGCTGCGCGAC
AAGCAGGAGGCGTAATGGCTACACCGCTTATTCGTGTC
```

rzoD gene (`Start` and `stop` codon) (**DLP-12**, putative prophage lysis lipoprotein)

```
ATGCGAAAGCTGAAAATGATGCTCTGCGTGATGATGTTGCCGCTGGTCGTCGTCGGTTGCACATCAAAGCAGTCTGT
                            |  ||    ||  |||||| |
            Spacer 12 3'-GTCGCAGTCCGCACTTTAGAGTGGCAGCAACG-5'          3'-G
CAGTCAGTGCGTGAAGCCACCACGGCCTCCGGCGTGGATAATGCAGCCTCCCCCCGACTGGCAGACACCGCTGAACG
       ||||||||  |  |  ||  |  |
TCGCAGTCCGCACTTTAGAGTGGCAGCAACG-5'  Spacer 12
GGATTATTTCACCCTCAGAGAGAGGCTGA
```

ydfD gene (`Start` and `stop` codon) (**Qin**, lysis protein)

```
ATGAATTCAGCATTTGTGCTTGTTCTGACAGTTTTTCTTGTTTCCGGAGAGCCAGTTGATATTGCAGTCAGTGTTCA
                                                   |||||   |  |
                            Spacer 12 3'-GTCGCAGTCCGCACT
TTAGAGTGGCAGCAACG-5'
 |   ||   |   |   |
CAGGACAATGCAGGAGTGTATGACTGCAGCAACCGAACAGAAAATTCCCGGTAACTGTTACCCGGTCGATAAAGTTA
TTCACCAGGATAATATCGAAATCCCGGCAGGTCTTTAA
  |    |   |   ||||||  |   |        |
  3'-GTCGCAGTCCGCACTTTAGAGTGGCAGCAACG-5'  Spacer 12
```

tfaP gene (`Start` and `stop` codon) (**e14**, tail fiber assembly)

```
ATGAAAATATACTGTTGCTTAAATACCGTTGGTTTTTTTATGGATGGCTGTGGCGTCATTCCGCCAGATTCTAAAGA
AATAACGGCAGAACACTGGCAGTCATTATTAAAATCTCAAGCTGAAGGAGGCGTGATCGATTTTTCTGTTTTTCCTC
CTTCTATTAAAGAGGTTATCCGTACTCATGATGATGAAGTCGCAGATGCGAACTTTCAAAAGCAGATGCTTATCTCT
GATGCAACTGATTTTATCAATAGCAGACAGTGGCAGGGTAAGGCTGCATTGGGAAGACTTAAAGAAGATGAGCTGAA
ACAATATAATTTGTGGCTGGATTATCTGGAAGCACTGGAACTGGTTGATACATCCAGTGCGCCAGATATTGAATGGC
                        |  ||  |   |      ||||||||||||
            Spacer 13 3'-GTTCACTATAGGTAGTAGCGTAGGTCACGCGG-5'
CTACGCCTCCGGCAGTTCAGGCCAGATGA
```

Supplementary Figure 8. Predicted base-pairing interactions between each of the CRISPR spacers (i.e., spacers 8, 11, 12, and 13) and a protospacer within the cryptic phage mRNA (CPS-53, CP4-6, Rac, Qin, CP4-57, DLP-12, and e14). Note, some spacers match more than one region within a particular cryptic prophage. The complementary nt between the spacers and cryptic prophage protospacers is indicated by yellow highlight and vertical lines. Note that the spacer sequences are written 3' to 5' to aid an understanding of the putative RNAi matching.

Supplementary Figure 9. Toxicity of YdfD and RzoD. Growth curve at 37°C in LB (turbidity at 600 nm shown) of *E. coli* BW25113 harboring empty plasmid pCA24N (blue line), plasmid pCA24N encoding *ydfD* (orange line), and encoding plasmid pCA24N encoding for *rzoD*. Red arrow indicates addition of 1 mM of IPTG produce the toxins. Average of two independent cultures shown, and error bars indicate standard deviations.

Supplementary Figure 10. Deleting the DNA encoding spacer #12, which encodes crRNA targeting four separate regions in cryptic prophage mRNA (mRNA of Rac, Qin, CP4-57, and DLP-12), leads to lower cell yield. As seen **Table S2** and **Fig. S6**, this is associated with increased cell lysis. In contrast, deleting *cas2* in Δ9 and deleting spacer 3, which encodes crRNA that lacks matches in the mRNA of all 9 cryptic prophages has little effect on growth and cell lysis. Δ9 is BW25113 that lacks all cryptic prophages and Kan[R] is kanamycin resistance. Growth (turbidity at 600 nm) at 37°C in LB shown, and error bars indicate standard deviations from two independent cultures.

D

S. enterica subsp. *enterica* serovar *Typhimurium* str. LT2

E

S. enterica subsp. *enterica* serovar *Typhimurium* str. LT2

Spacer sequence

CGGTTTATCCCGTGGGCGGGGGGGAACAC / A**TCTTCATATTGCG**TGACGCTGCCGATGAACG
TCTTTATCAGCTAACCATTTCCAGAACTGTC
TATA**TATGAATTAAT**TTTTGGGCATAACCTG
TGCCCGTTCTGCCTCTTCGCACTCGATCAA
TGGGTAATGGGCTACCTGAACTTCACATATCC
ATTAAGCGGCAAAAGTTTGGGTTAATTGGACA
CGTATTCGTCACACAGCCCGTCCAGAAATGA
TAACGAACTGAATAAAATGTCAGAAAGTGACG
GCAGCTTAGCGACGAGAATTAAAACCGAACTCAC
TGCCA**GTGACTACGAGAGCGT**CGCTATCGGG
ACCGATAAACACCCGCATAGCCTCTTTCGTTT
TGCTCAATAACGTCGTAAATAGGGTAAGCTGG
TATTTCGCCTTCGGCACTGACGTCACCGTCAA
GTGCCGTTCGTTGCCGGTATAGACCAGCGTCA
ATCGAATCGAAACCCCAGCC**CACAGAAATAAT**T
GCTCATGTCAAACGCCATCAGCGTTCCGGCAT
AATCGCCAGCCTCGGAAATATTCCATCCTCCG
AGGAACTAAAACAGCCTGACCGTTGAGGATCG
ACCGGACAAATCTTTTTTCCTGTTCCTGTT
GGGCCACTATGAACGGATCGGCGCTGATGCCGG
GGTAAAGCCACACCATTTTTATTGGACCTGC
GTGGTGGCCTCAAATAAATTCGACGCGCTGGAG
TCGACGTGGACGAGGAGTACTCAACCGCTGC
AGCCGCACATGGCCCACCGGCACCACCCGATC
AAATGACCAAATCAGAAATCTTCACCAAAAGCC
TAAT**GGCCACAGTAAG**TCAAACGGTTCTGGA
GAGTCCGGGGGGTATATAGTTATTTAATGAGC
TTGGGCGTCGGTTTTTCAGGTTTAGGTCCGG
TCAACTGTCAGTTCGTCGTTAGCCAGTAATTC
CTGAAAAACGCATGGAGTCCGGTATAAAACAGTC
GATGTAACTGATAGCG**GAAATATATTGGGGATAA**
GAAAC**GTAAACAGGGTAA**AGATACAACTCTGCA

Cryptic prophage Gene

ATATGAATTAAT
PHAGE_Entero_HK630
pseudogene / *pagK*

ATGAATTAATTT
PHAGE_Entero_HK630
RBS / sopE

TTGCGTGACGCTG
PHAGE_Burkho_phiE255
Baseplate protein / *Protein D* / Hypothetical protein

TC__CATATTGCGTGAC_C
PHAGE_Burkho_phiE255
Baseplate protein / *Protein D* / Hypothetical protein

TCTTCATATTGCG
PHAGE_Burkho_phiE255
Phage tail protein I / *Baseplate protein*

TG_CTACAGAAGCGT
PHAGE_Burkho_phiE255
RBS / Hypothetical protein / *glycosyltransferase*

CACAGAAATAAT
PHAGE_Entero_HK630
rarD / eamA / *rarD*

GGCCACAGTAAG_C
PHAGE_Burkho_phiE255
RBS / Hypothetical protein / *RBS*

TTGG_C__CGGTTTTTTCAGGTT
PHAGE_Cronob_vB_CsaM_GAP32
yljA / clpA

G__ATATATTGGGAT_A
PHAGE_Salmon_118970_sal3
RBS / NTPase

GTAAACAGGGTAA
PHAGE_Salmon_Fels_1
oligopeptidase B

Supplementary Figure 11. *E. coli* **O157:H7 and *Salmonella* spp. CRISPR-Cas spacer sequences related to cryptic prophages. (A)** Repeat (R, blue hexagon) and spacer (rectangle) sequences of each CRISPR-Cas system of *E. coli* O157:H7, *Salmonella* spp., and *K. pneumoniae* indicating matches with their cryptic prophages. (**A**) *E. coli* O157:H7, (**B**) *S. enterica* subsp. *enterica* serovar *Typhi* strain TY2, (**C, D, E**) three CRISPR-Cas systems of *S. enterica* subsp. *enterica* serovar *Typhimurium* str. LT2, and (**F**) *K. pneumoniae*. Pink highlight and dashed lines indicate the spacer positions relative to the cryptic prophage genes.