1

PLB-985 neutrophil-like cells as a model to study *Aspergillus fumigatus* pathogenesis

Muhammad Rafiq[a,b,#], Flora Rivieccio[a,b,#], Ann-Kathrin Zimmermann[a,b,#], Corissa Visser[a,b], Alexander Bruch[c],

Thomas Krüger[a], Katherine González Rojas[a,b], Olaf Kniemeyer[a,b], Matthew G. Blango[c], Axel A. Brakhage[a,b]*

[a] Department of Molecular and Applied Microbiology, Leibniz Institute for Natural Product Research and

Infection Biology - Hans Knöll Institute (Leibniz-HKI), Jena, Germany.

[b] Department of Microbiology and Molecular Biology, Institute of Microbiology, Friedrich Schiller University,

Jena, Germany.

[c] Junior Research Group RNA Biology of Fungal Infections, Leibniz Institute for Natural Product Research and

Infection Biology - Hans Knöll Institute (Leibniz-HKI), Jena, Germany.

Running Title: Neutrophil-like model of *A. fumigatus* infection

Key Words: *Aspergillus fumigatus*, PLB-985, phagocytosis, fungal pathogens, extracellular vesicles, HL-60

*Correspondence: E-mail axel.brakhage@leibniz-hki.de

Tel. +49 (0)3641-532 1001.

# These authors contributed equally to the manuscript and were listed alphabetically

ABSTRACT

Fungal infections remain a major global concern. Emerging fungal pathogens and increasing rates of resistance mean that additional research efforts and resources must be allocated to advancing our understanding of fungal pathogenesis and developing new therapeutic interventions. Neutrophilic granulocytes are a major cell type involved in protection against the important fungal pathogen *Aspergillus fumigatus*, where they employ numerous defense mechanisms, including production of antimicrobial extracellular vesicles. A major draw-back to work with neutrophils is the lack of a suitable cell line system for the study of fungal pathogenesis. To address this problem, we assessed the feasibility of using differentiated PLB-985 neutrophil-like cells as an *in vitro* model to study *A. fumigatus* infection. We find that dimethylformamide-differentiated PLB-985 cells provide a useful recapitulation of many aspects of *A. fumigatus* interactions with primary human polymorphonuclear leukocytes. We show that differentiated PLB-985 cells phagocytose fungal conidia and acidify conidia-containing phagolysosomes similar to primary neutrophils, release neutrophil extracellular traps, and also produce antifungal extracellular vesicles in response to infection. In addition, we provide an improved method for the isolation of extracellular vesicles produced during infection by employing a size-exclusion chromatography-based approach. Advanced LC-MS/MS proteomics revealed an enrichment of extracellular vesicle marker proteins and a decrease of cytoplasmic proteins in extracellular vesicles isolated using this improved method. Ultimately, we find that differentiated PLB-985 cells can serve as a genetically tractable model to study many aspects of *A. fumigatus* pathogenesis.

IMPORTANCE

Polymorphonuclear leukocytes are an important defense against human fungal pathogens, yet our model systems to study this group of cells remains very limited in scope. In this study, we established that differentiated PLB-985 cells can serve as a model to recapitulate several important aspects of human polymorphonuclear leukocyte interactions with the important human fungal pathogen *Aspergillus fumigatus*. The proposed addition of a cultured neutrophil-like cell line to the experimental toolbox to study fungal pathogenesis will allow for a more mechanistic description of neutrophil antifungal biology. In addition, the easier handling of the cell line compared to primary human neutrophils allowed us to use PLB-985 cells to provide an improved method for isolation of neutrophil-derived extracellular vesicles using size-exclusion chromatography. Together, these results provide significant tools and a baseline knowledge for the future study of neutrophil-derived extracellular vesicles in the laboratory.

INTRODUCTION

Fungal infections remain a tremendous source of global morbidity and mortality. More than 1 billion individuals are affected by fungal infections per year, with invasive infections killing numbers comparable to other leading bacterial pathogens (>1.5 million per year; gaffi.org; (1)). Deadly invasive infections are caused by a relatively small number of fungi, with most of these attributed to members of the genera *Candida*, *Pneumocystis*, *Cryptococcus*, and *Aspergillus* (2). *Aspergillus fumigatus* is the major cause of aspergillosis and is particularly dangerous to immunocompromised individuals suffering from neutropenia (3). There is also emerging evidence to suggest that invasive aspergillosis may contribute to COVID-19-related deaths (4, 5), but challenges in safely obtaining bronchoalveolar lavage samples from these patients have often made confirmatory diagnosis difficult. Despite the obvious importance of *A. fumigatus* in the clinic, our understanding of this important pathogen remains lacking in many aspects, in part due to a lack of tractable experimental systems in the laboratory.

Mammals are continuously challenged by fungal pathogens. In fact, asexual spores of *A. fumigatus*, termed conidia, are thought to be inhaled by humans on a scale of hundreds per day (6). For most fungi, the body temperature of mammals is too high to allow for growth, but for organisms like *A. fumigatus* that thrive in compost piles at high temperatures, the human host provides fertile ground in the absence of a functional immune system (7). However, humans do have multiple additional defenses, including a mucociliary escalator to remove particles from the lungs, a robust epithelium, resident alveolar macrophages that eliminate the majority of the remaining fungal conidia, and infiltrating polymorphonuclear leukocytes (PMNs) that aid in clearance of conidia and destruction of fungal hyphae, among others (3, 8). Neutrophils play an essential role in antifungal defense, due to their importance in killing fungal hyphae. This is well-illustrated by the high susceptibility of neutropenic patients to *A. fumigatus* infections in the clinics (reviewed in (3, 9)).

Studies in primary human neutrophils have revealed the capacity of these cells to phagocytose conidia and release granules, neutrophil extracellular traps (NETs), and extracellular vesicles in response to invading pathogens (10-12). Phagocytosis occurs in conjunction with recognition of pathogen-associated molecular patterns by host pathogen recognition receptors. Studies in zebrafish and mice have shown that these internalized conidia can even be passed from neutrophils to macrophages for destruction of the fungus (13), implying complex intracellular trafficking. In neutrophils and macrophages however, internalized wild-type *A. fumigatus* conidia are capable of stalling phagolysosomal acidification to facilitate outgrowth.

In addition to phagocytosis, NETs are an important mechanism of defense against *A. fumigatus* that are produced in response to fungal recognition in a CD11b-dependent manner (11, 14). NET production is most abundant against hyphae, but NETs are also sometimes produced in response to resting and swollen conidia (15). In vivo, NETs were shown to be present in mouse lungs during infection but were generally dispensable for fungal

clearance (15, 16), suggesting that alternative measures are required to eliminate fungal hyphae that escaped
phagocytosis as conidia.

Neutrophilic granules and extracellular vesicles are two heterogeneous subcellular populations that also
contribute widely to the antimicrobial response. Proteins like lactoferrin associated with granules are known to
inhibit *A. fumigatus* conidial germination due to iron sequestration (17) and the abundant granule protein
myeloperoxidase inhibits fungal hyphal growth (18). Extracellular vesicles were also shown to provide an
antifungal mechanism against *A. fumigatus* hyphae and conidia (10). Antifungal extracellular vesicles produced by
neutrophils in response to infection are capable of associating with the fungal cell wall and are in some cases
internalized to deliver an unknown antifungal cargo. Intriguingly, the antifungal effect of these extracellular
vesicles appeared to be tailored to the infecting pathogen, as spontaneously released vesicles and vesicles
released in response to infection with a knockout strain deficient in production of the conidial pigment
dihydroxynaphthalene (DHN)-melanin were not antifungal against wild-type fungus (10).

The dissection of the molecular mechanisms behind these differences in antifungal potential are difficult
to discern in the absence of a tractable model system for the genetic manipulation of neutrophils. Despite the
importance of neutrophils in antifungal defense, the majority of studies are still performed in primary human
PMNs isolated from venous blood, a resource that can only be used for small-scale studies and is unfortunately
highly variable between blood donors. A laborious isolation procedure and limited half-life of about 19 hours *ex*
vivo (19) also make more mechanistic studies difficult. For all these reasons, establishment of a cell culture line to
study *A. fumigatus* pathogenesis in neutrophils or neutrophil-like cells would be highly advantageous. In recent
years some options have emerged to investigate specific aspects of neutrophil biology in culture (20). In particular,
the myeloid cell line PLB-985, a derivative of HL-60 granulocytic cells, has recently emerged as a tractable model
to study NET formation in response to bacterial infection after differentiation into neutrophil-like cells (21, 22). Of
note, the related HL-60 cell line has also been used previously to deliver azole drugs to *A. fumigatus*-infected mice
(23), but not to our knowledge in studies of *A. fumigatus* pathogenesis.

In this study we hypothesized that differentiated human PLB-985 neutrophil-like cells could serve as
model system to study the host-pathogenesis of *A. fumigatus* in a blood donor-independent manner. We tested
this hypothesis by defining the phagocytosis and intracellular trafficking of *A. fumigatus* conidia by PLB-985 cells,
the production of neutrophil extracellular traps, and the production of extracellular vesicles produced in response
to fungal infection. We find numerous similarities between this cell line and primary neutrophils that suggest
differentiated PLB-985 cells can serve as a tractable *in vitro* model system for further study of some aspects of *A.*
fumigatus pathogenesis.

RESULTS

Differentiated PLB-985 neutrophil-like cells phagocytose opsonized *A. fumigatus* conidia.

We set out to assess the feasibility of using *N,N*-dimethylformamide (DMF)-differentiated PLB-985 (dPLB) cells as a model for neutrophil phagocytosis, NET production, and extracellular vesicles release in response to *A. fumigatus* infection. Upon infection, neutrophils are known to rapidly phagocytose and process *A. fumigatus* conidia. We therefore challenged dPLB cells with opsonized *A. fumigatus* resting conidia and assessed various parameters of infection.

First, we determined the phagocytic capacity of PLB-985 cells using confocal laser scanning microscopy following incubation of cells with fluorescein isothiocyanate (FITC)-labelled, opsonized conidia (green). We observed that dPLBs were capable of phagocytosis, as evidenced by a phagosomal membrane surrounding the conidia (**Fig. 1A**). A similar result was confirmed for primary human PMNs with this experimental setup (**Fig. S1A**). Interestingly, dPLB cells only infrequently phagocytosed unopsonized conidia, whereas primary human neutrophils indiscriminately phagocytosed both unopsonized and opsonized conidia (data not shown). To better quantify phagocytosis, we analyzed 5,000 host cells post-infection by single-cell analysis using imaging flow cytometry. For this we counterstained with calcofluor white (CFW; blue) to elucidate non-phagocytosed conidia (**Fig. 1B and Fig. S1B to C**). Quantification of imaging flow cytometry data revealed that approximately 22% of wild-type and 26% of non-pigmented Δ*pksP* conidia were phagocytosed by dPLBs after 2 hours, while undifferentiated PLB-985 cells exhibited limited phagocytosis of both wild-type and non-pigmented conidia (**Fig. 1C**). As expected, these phagocytosis percentages were lower than those of primary human PMNs, which phagocytosed 62% of wild-type and 54% of Δ*pksP* conidia after 2 hours and 90% of wild-type and 85% of Δ*pksP* conidia after 4 hours of infection (**Fig. S1D**). We also measured the induction of cell damage following infection using lactate dehydrogenase release assays and found levels below the limit of detection for dPLB cells after 2 and 4 hours of incubation (data not shown). To determine if infection also resulted in an activated immune response, we measured proinflammatory cytokine levels by enzyme-linked immunosorbent assay (ELISA). Co-incubation of dPLB cells with wild-type spores resulted in production of significantly increased levels of interleukin (IL)-8, but not IL-1β after 24 hours of co-incubation, comparable to primary neutrophils at the corresponding earlier time points (**Fig. 1D and Fig. S2**). Together, these results suggested that DMF-differentiated PLB-985 cells could serve as a suitable model for *A. fumigatus* pathogenesis and warranted further investigation.

Internalized *A. fumigatus* conidia are processed inside dPLB phagolysosomes.

Following internalization of conidia by professional phagocytes like macrophages and neutrophils, conidia-containing phagosomes fuse with lysosomes to acidify the compartment and aid in fungal killing. Numerous

additional enzymes are activated upon phagolysosomal acidification to degrade and digest the internalized fungal conidia. The acidification and maturation of phagolysosomes are regulated by protein complexes assembled on the phagolysosomal membrane, including lysosomal-associated membrane protein (LAMP)-1, 2, and 3; several vacuolar ATPases (V-ATPases); Ras-related protein (RAB) 5 and 7; Flotillin 1 and 2; and numerous others (24). DHN-melanin on the surface of conidia can interfere with these processes in alveolar macrophages, monocytes, and primary neutrophils to delay fungal processing and phagolysosomal acidification (25). To test whether dPLB cells process conidia similarly to primary human PMNs, we first stained the cells with LAMP-1, a general endocytic marker on the membrane of phagolysosomes. LAMP-1 showed a clear signal around phagolysosomes of infected cells (**Fig. 2A**), and the intensity of the signal did not differ between wild-type and *ΔpksP* conidia (**Fig. 2B**). Furthermore, loading of dPLB cells with Lysotracker, a weak base that becomes fluorescent under acidic conditions, showed that the *ΔpksP* conidia-containing phagosomes were significantly more acidified than those containing wild-type conidia, revealing that DHN-melanin can also block the acidification process in the dPLB model as well (**Fig. 2C to D**).

The interference of the acidification pathway in alveolar macrophages by fungal DHN-melanin occurs through the inhibition of V-ATPase assembly (26, 27). This multiprotein complex plays a major role in lowering the pH from 6 to <4.5 by pumping H^+ ions across the phagolysosomal membrane. Staining of the V-ATPase V1 subunit in dPLB cells revealed that the percentage of recruitment to conidia-containing phagolysosomes was fairly similar for wild-type and *ΔpksP* conidia (**Fig. S3A to B**), suggesting that other proton pumps may contribute to acidification of these phagolysosomes in neutrophil-like cells. An additional defense mechanism for intra-phagosomal degradation of pathogens occurs via reactive oxygen species (ROS) production. As expected, we observed dPLB cells to be capable of producing ROS after staining with CellROX Orange (**Fig. S3C**), consistent with the literature (28, 29).

A. fumigatus triggers formation of NETs from dPLB cells.

One defense mechanism employed by neutrophilic granulocytes to fight against pathogens is the formation of NETs. These structures consist of condensed chromatin and various enzymes that are released into the extracellular space and typically correlate with death of the cell (30). NETs contain several anti-fungal proteins, which are responsible for the fungistatic effect against *A. fumigatus* hyphae (15). To test if dPLBs are also able to form NETs in response to *A. fumigatus* and serve as a model for this pathway, we co-incubated dPLBs cells with hyphae and stained for nucleic acid with 4',6-diamidino-2-phenylindole (DAPI; **Fig. 3A**). As a positive control, we induced NET formation using phorbol myristate acetate, a known trigger of NET formation in PLB-985 cells (**Fig. 3B**; (21, 31)). We observed the presence of histone H3 embedded in the DNA fibers and an association of NETs

with fungal hyphae (**Fig. 3B**). Taken together these results showed that *A. fumigatus* is able to trigger NET

formation in dPLB cells and that these NETs are specifically directed against the hyphae.

dPLB cells produce extracellular vesicles in response to infection.

Recently, a new defense mechanism from neutrophilic granulocytes was discovered, the production of

antimicrobial extracellular vesicles (32, 33). These small lipid-enclosed nanoparticles are released from primary

PMNs after contact with microorganisms and, in the case of *A. fumigatus* conidia, can inhibit fungal growth after

coincubation (10). This effect is likely due in part to their protein cargo, which consists of antimicrobial peptides

such as myeloperoxidase, azurocidin, and cathepsin G. To assess if dPLB cells can produce extracellular vesicles

spontaneously or in response to *A. fumigatus*, we incubated cells with or without conidia for 2 and 4 hours. Next,

we isolated extracellular vesicles using two different methods; first using a previously described differential

centrifugation-based approach (DC; (10)) that enriches for medium-sized extracellular vesicles and a second

approach that relies on size-exclusion chromatography (SEC) to purify a more selective population of smaller

extracellular vesicles. Using these methods, dPLB cells were observed to actively secrete extracellular vesicles over

time in a manner comparable to primary human PMNs (**Fig. 4A to B).** Using nanoparticle tracking analysis, we

observed the median size of particles to be around 200 nm, similar to extracellular vesicles derived from primary

neutrophils, a feature that was comparable between infection-derived and spontaneously released extracellular

vesicles (**Fig. 4C to D**).

We next compared the protein content of dPLB-derived extracellular vesicles isolated, at the 2 hours

timepoint, using each isolation method with and without fungal infection by LC-MS/MS-based proteomics

analysis. We were able to identify 1,984 unique proteins across all four samples (**Dataset S1**). The majority of

identified proteins (737 proteins) were found in all four samples (**Fig. 5A**). We expected that isolation using size-

exclusion chromatography would improve the quality of the isolated particles as has been shown previously (34),

and this was in fact the case. We observed an increase in extracellular vesicle markers like the tetraspanins CD63

and CD81, and tumor susceptibility gene 101 (TSG101), and a decrease in cytoplasmic proteins like calnexin (CANX;

Table 1).

Extracellular vesicles produced by primary PMNs in response to *A. fumigatus* infection are known to be

enriched for antimicrobial cargo proteins like azurocidin, cathepsin G, and defensin (10). In line, with the data

from primary neutrophils we found that dPLBs also contained many of the same proteins in both infection-derived

and spontaneously released extracellular vesicles (**Table 1**), as evidenced by an UpSetR plot showing overlapping

proteins cohorts (**Fig. S4;** (10)), although with some exceptions. For example, dPLBs appeared to lack defensin,

neutrophil elastase, and some histone proteins previously observed. Ultimately, improvements in LC-MS/MS

technology and the advantage of higher input amounts of extracellular vesicle protein using large amounts of
dPLB cells in culture resulted in significantly more proteins detected in the dataset provided here compared to
efforts using primary neutrophils (10), proving that dPLBs offer a scalable system for the elucidation of novel
mechanisms of neutrophil extracellular vesicle biology.

dPLB extracellular vesicles produced against *A. fumigatus* limit fungal growth.

The most compelling feature of infection-derived extracellular vesicles of human neutrophils is likely their
antifungal capacity, as we previously reported (10). We set out to determine if the extracellular vesicles produced
by dPLB cells in response to *A. fumigatus* opsonized conidia are antifungal to a mitochondrial-GFP reporter strain
used previously as a marker of fungal viability (35). First, we incubated the reporter strain with dPLB cells and
primary human PMNs and assessed the capacity of the cells to control infection. PMNs were capable of minimizing
fungal outgrowth after 22 hours, whereas dPLBs did not completely contain fungal growth (**Fig. 6A**). We then
assessed the antifungal capacity of the dPLB extracellular vesicles. For this experiment, extracellular vesicles from
equal numbers of dPLBs and primary human PMNs were isolated using differential centrifugation and found to be
antifungal against the mitochondrial-GFP reporter strain. This was evidenced by fragmentation of fungal
mitochondria after administration of extracellular vesicles to germinating conidia, which were allowed to
germinate for 6 hours prior to overnight incubation with extracellular vesicles, 3 mM H_2O_2 as a positive control,
or left untreated as a negative control (**Fig. 6B**). Although, the mitochondrial reporter clearly indicated
fragmentation, the extracellular vesicles from dPLB did appear to be in some cases less effective in limiting the
length development of the hyphae than those from primary PMNs. It is important to note that dPLBs produced
slightly fewer extracellular vesicles, which could potentially explain the decreased antifungal activity in these
assays (**Fig. 6B**). Additional representative images of dPLB-derived extracellular vesicles are shown to indicate the
spectrum of phenotypes observed for each experimental condition (**Fig. S5**). Collectively, these results suggest
that dPLB infection-derived extracellular vesicles can limit *A. fumigatus* hyphal growth, similar in fashion to
primary human PMNs (10).

MATERIALS AND METHODS

Fungal strains cultivation and opsonization.

Cultivation of *A. fumigatus* strains CEA10 (Fungal Genetics Stock Center; A1163), CEA17 Δ*akuB^{ku80}* Δ*pksP* (36), and
AfS35/pJW103 (35) was performed on malt agar plates (Sigma-Aldrich) supplemented to a final concentration of
2% (wt/vol) agar for 5 days at 37 °C. Conidia were harvested in sterile, ultrafiltrated water, filtered through a 30-
μm pore filter (MACS, Miltenyi Biotec). Prior to confrontation with PLB-985 cells, conidia were opsonized with

normal human serum (Merck Millipore). Briefly, 900 µl of spore suspension was mixed with 100 µl of normal

human serum and incubated in a thermomixer at 37 °C for 30 min shaking at 500 rpm. The spore suspension was

washed three times by collecting spores *via* centrifugation at 1800 x *g* for 1 min at 4 °C and resuspending in fresh

PBS 1X. Following washing, spores were enumerated in a Thoma chamber in preparation for infection assays.

Cell culture cultivation, differentiation, and infection.

PLB-985 cells were maintained in Roswell Park Memorial Institute medium 1640 (RPMI; Gibco) supplemented with

10% (vol/vol) fetal bovine serum (FBS; HyClone, GE Life science), 100 U/ml penicillin-streptomycin antibiotic

solution (Lonza) and 2 mM UltraGlutamine (alanyl L-glutamine; Gibco). For differentiation into a neutrophilic

granulocyte cell type, 4×10^5 cells/ml were resuspended in supplemented RPMI medium containing 2.5% (vol/vol)

FBS and 0.5% DMF (Sigma-Aldrich) for 4 days at 37 °C with 5% (vol/vol) CO_2. On day 4, fresh differentiation medium

was added, and the cells were incubated for 3 additional days. On day 7, cells were collected, centrifuged at 300

x *g* for 5 min and then seeded in wells for experiments at the appropriate concentrations. The medium used for

all the assays was composed of RPMI supplemented with 1% (vol/vol) exosome-depleted FBS (Life Technologies

GmbH), 2 mM UltraGlutamine, and 0.5% (vol/vol) DMF.

Isolation of human PMNs. Fresh venous blood was drawn from adult healthy volunteers, aged 20 to 35 years,

after obtaining written consent. This study was approved by the Jena University Hospital Institutional Review

Board (approval number 5074-02/17) in agreement with the Declaration of Helsinki. Blood was collected in EDTA

BD Vacutainer tubes (BD Biosciences) and mixed with PBS 1X. 6 ml of this solution was carefully layered onto

(avoiding any mixing) 5 ml (1:1 vol/vol) of PolymorphPrep solution (Progen) and centrifuged 50 min at 500 x *g* for

gradient separation. After centrifugation the neutrophil layer was collected, mixed with an equal volume of PBS

0.5% and re-centrifuged for 15 min at 400 x *g*. To limit contamination of red blood cells, the pellet was lysed with

the ACK lysis buffer (Gibco) and centrifuged as before. The PMNs were finally re-suspended in RPMI medium

without phenol red (Thermo Fischer scientific) and after addition of Trypan blue counted with a Luna automated

cell counter (Logos biosystem). For each experiment the viability of neutrophils was ≥ 95%.

Determination of cell viability.

To assess cell viability after confrontation with *A. fumigatus* conidia, the release of LDH was determined using the

CyQUANT™ LDH cytotoxicity assay kit (Thermo Fisher Scientific) according to the manufacturer´s instruction. Cells

treated with *A. fumigatus* conidia with a multiplicity of infection (MOI) of 5 for 2 hours were tested for their LDH

activity, which was compared to the spontaneous LDH activity and the maximum LDH activity. Release of LDH was

below the limit of detection for all three independent biological replicates performed.

Phagocytosis assays.

To assess phagocytic ability, we used a combination of imaging flow cytometry and confocal fluorescence

microscopy. For both methods the conidia were first stained with fluorescein isothiocyanate (FITC) and then after

confrontation of dPLBs or primary neutrophils, counterstained with calcofluor white (CFW; Sigma -Aldrich). The

FITC solution was obtained by dissolving FITC powder (Sigma-Aldrich) in 5 mL of 0.1 M sodium carbonate (Na_2CO_3)

followed by filtration through a 0.22-μm pore size filter (Carl Roth). Afterwards 1 ml of this solution was mixed

with 500 μl of spore suspension and incubated for 30 min at 37 °C while shaking at 1000 rpm in the dark. The

spores were then pelleted and washed three times with PBS 1X with 0.001% (vol/vol) Tween 20 (Carl Roth). During

the last washing step Tween was removed to avoid residual detergent in the samples (25, 37). Before counting,

the conidia were opsonized following the protocol described above. They were then added to the cells and co-

incubated for 2 and 4 hours at 37 °C with 5% (vol/vol) CO_2. At the end of each time point CFW was added to a final

concentration of 1 μg/ml and incubated for 1 min. Afterwards the cell suspension was transferred to

microcentrifuge tubes, centrifuged at 600 x g for 2 min at 4 °C and washed with PBS 1X twice. After discarding the

supernatant, the pellet was fixed with 150 μL of 3.7% (vol/vol) formaldehyde in PBS at room temperature for 15

min and subsequently washed again as described above. For imaging flow cytometry measurements, the cells

were resuspended in 150 μL of PBS 1X and analyzed immediately or stored at 4 °C for no more than 24 hours. Four

independent experiments were performed and for each replicate 5000 cells were analyzed using the ImageStream

X Mark II (Luminex). A 488 nm laser was used to detect FITC staining and a 405 nm laser for the CFW. The laser

voltage was adjusted depending on the sample. For wild-type conidia, we used 10 mW of 488 nm laser and 2 mW

for the 405 nm laser, and for Δ$pksP$ conidia 1 and 2 mW of the respective lasers. The fluorescence intensities of

the samples were compensated and analyzed with the IDEAS software (Luminex). An example of the gating

strategy used for analysis can be found in the supplement files (**Fig. S1B to C**). For the fluorescence microscopy

analysis, cells were seeded in 8-well μ-slides (Ibidi) and visualized using a Zeiss LSM 780 confocal microscope (Carl

Zeiss) from at least three biological replicates as described above.

Immunofluorescence assays.

For immunofluorescence, cells were allowed to adhere in a 24-well plate with poly-L-lysine-coated glass coverslips

for 1 hour (Merck). After infection with labelled *A. fumigatus* spores and incubation for the noted times, cells were

fixed using 3.7% (vol/vol) formaldehyde in PBS for 10 min, rinsed three times with PBS 1X, permeabilized for 15

min using 0.1% (vol/vol) Triton X-100 in PBS 1X or 0.1% (wt/vol) saponin in PBS 1X, and then blocked for 30 min

with 2% (wt/vol) bovine serum albumin (BSA). After permeabilization, washed cells were incubated with primary

rabbit anti-Lamp-1 antibody (Abcam 24170, 1:100 dilution), anti-V-ATPase V1 subunit antibody (Abcam 73404,

1:100 dilution), or anti-Histone H3 (Cell signaling DIH2, 1:200 dilution) antibody in 1% (wt/vol) BSA in PBS 1X,

followed by incubation with secondary goat anti-rabbit IgG antibody DyLight 633 (Thermo Fisher Scientific). The

glass cover slips were mounted onto microscopy slides and visualized using a Zeiss LSM 780 confocal microscope

(Carl Zeiss). LAMP-1 recruitment was quantified by comparing the positive signal from stained phagolysosomes to

non-stained phagolysosomes.

Reactive oxygen species measurements.

ROS production was detected in dPLB cells by seeding cells in 8-well μ-slides (Ibidi) coated with poly-L-lysine

followed by infection with *A. fumigatus conidia* at an MOI of 5 for either 2 or 4 hours. CellROX® Orange Reagent

(Thermo Fisher Scientific) was added 30 minutes prior the end of the infection time points, and ROS production

was imaged using a Zeiss LSM 780 confocal microscope (Carl-Zeiss) for fluorescent intensity.

Cytokine measurement.

For the detection of IL-8 and IL-1β, $2x10^5$ dPLB cells or primary neutrophils were seeded in 24-well plates. After

addition of conidia at a MOI of 5, plates were incubated at 37 °C with 5% (vol/vol) CO_2 for 2, 4, or 6 hours. An

additional 24-hour timepoint was included for dPLB cells. As a positive control, cells were treated with 5 μg/mL of

lipopolysaccharide (LPS; Sigma-Aldrich L4516). At the appropriate end point, samples were collected, centrifuged

at 300 x *g* for 5 min to remove cell debris, and frozen at -20°C and analyzed within 3 days. Cytokines were

measured using human ELISA Max deluxe kits (BioLegend) according to the manufacturer´s instruction.

Acidification assays.

Acidification assays were performed as mentioned previously (27). Briefly, the cells were incubated with 50 nM

lysotracker DND-99 (Thermo Fisher Scientific) for 1 hour prior to infection. After that, FITC-labeled conidia were

added to the cells for 2 or 4 hours and imaged using Zeiss LSM 780 confocal microscope (Carl Zeiss). For

quantification, 200 conidia-containing phagolysosomes were counted and evaluated for acidification. The values

represent the mean ± SD of three separate experiments.

Extracellular vesicle isolation.

After infection with *A. fumigatus* the supernatant of dPLB cells was collected, and extracellular vesicles were

isolated using two different methods: a differential centrifugation-based approach or a size exclusion

chromatography-based approach. The first method was described previously for the isolation of neutrophil-

derived extracellular vesicles (10, 32). In both approaches samples were centrifuged at 3000 x *g* for 15 min at 4 °C

and then filtered through 5-μm pore size filters (Carl Roth). In the first method, samples were then centrifuged at

19,500 x *g* for 20 min at 4 °C to collect extracellular vesicles. For the second approach, the clarified filtrate was

concentrated using Amicon Ultra-15 centrifugal filters (Merck) with a molecular mass cut-off (MWCO) of 100 kDa

for 10 min at 4 °C and 3220 *x g* and finally loaded on size-exclusion chromatography qEV 70 nm columns (Izon).

After discarding the 3 mL void volume, 1.5 mL of extracellular vesicle sample were collected and measured. When

necessary, extracellular vesicles were further concentrated using 10 kDa cutoff Amicon Ultra-0.5 ml filters (Merck).

Extracellular vesicles were used fresh for most of the downstream experiments except for proteomic analysis

where they were frozen at -20 °C.

Nanoparticle tracking analysis.

Particle concentration and size distribution were analyzed using a nanoparticle tracking analysis (NTA) NS300

device with a 650 nm laser (Malvern Instruments Ltd). Fresh samples were measured with a constant flow rate of

20 and a temperature set to 25 °C. Five 60-second videos were recorded with a camera level of 11. Videos were

then analyzed with the NTA 3.2.16 software using a detection threshold of 4.

Protein preparation for LC-MS/MS.

After isolation as described above, extracellular vesicles were delipidated as described previously (10) using the

protein precipitation protocol of Wessel and Flügge (38). Delipidated extracellular vesicles were resolubilized in

100 μl of 50 mM triethyl ammonium bicarbonate (TEAB) in 1:1 (vol/vol) trifluoroethanol/water. For reduction of

cysteine thiols, the solution was mixed with 10 mM Tris(2-carboxyethyl) phosphine and alkylated with 12.5 mM

chloroacetamide at 70 °C for 30 min in the dark. Proteins were digested for 18 hours at 37 °C with trypsin/LysC

mix (Promega) at a protein-to-protease ratio of 25:1. Tryptic peptides were first completely evaporated using a

vacuum concentrator (Eppendorf) and then resolubilized in 0.05% (vol/vol) trifluoroacetic acid (TFA) in 2:98

(vol/vol) acetonitrile/water. Finally, the samples were filtered through Ultrafree-MC Hydrophilic PTFE membrane

(0.2-μm pore size) spin filters (Millipore) and stored at -20 °C until measurement. Each sample was measured in

triplicate (3 analytical replicates of 3 biological replicates) as follows.

LC-MS/MS analysis.

LC-MS/MS analysis was performed on an Ultimate 3000 nano RSLC system connected to a QExactive HF mass
spectrometer (both Thermo Fisher Scientific). Peptide trapping for 5 min on an Acclaim Pep Map 100 column (2
cm x 75 µm, 3 µm) at 5 µL/min was followed by separation on an analytical Acclaim Pep Map RSLC nano column
(50 cm x 75 µm, 2µm). Mobile phase gradient elution of eluent A (0.1% (vol/vol) formic acid in water) mixed with
eluent B (0.1% (vol/vol) formic acid in 90/10 acetonitrile/water) was performed as follows: 0 min at 4% B, 30 min
at 12% B, 75 min at 30% B, 85 min at 50% B, 90-95 min at 96% B, 95.1-120 min at 4% B.

Positively charged ions were generated at spray voltage of 2.2 kV using a stainless-steel emitter attached
to the Nanospray Flex Ion Source (Thermo Fisher Scientific). The quadrupole/orbitrap instrument was operated in
Full MS / data dependent MS2 (Top15) mode. Precursor ions were monitored at m/z 300-1500 at a resolution of
120,000 FWHM (full width at half maximum) using a maximum injection time (ITmax) of 120 ms and an AGC
(automatic gain control) target of $3x10^6$. Precursor ions with a charge state of z=2-5 were filtered at an isolation
width of m/z 1.6 amu for further HCD fragmentation at 27% normalized collision energy (NCE). MS2 ions were
scanned at 15,000 FWHM (ITmax=100 ms, AGC= $2x10^5$). Dynamic exclusion of precursor ions was set to 25 s and
the underfill ratio was set to 1.0%. The LC-MS/MS instrument was controlled by Chromeleon 7.2, QExactive HF
Tune 2.8 and Xcalibur 4.0 software.

Database search and data analysis
Tandem   mass   spectra   were   searched   against   the   UniProt   database   of   *Homo   sapiens*
(https://www.uniprot.org/proteomes/UP000005640;   2021/05/17)   and   *Neosartorya   fumigata*
(https://www.uniprot.org/proteomes/UP000002530; 2021/05/17) using Proteome Discoverer (PD) 2.4 (Thermo)
and the algorithms of Mascot 2.4.1 (Matrix Science), Sequest HT (version of PD2.2), MS Amanda 2.0, and MS
Fragger 3.2. Two missed cleavages were allowed for the tryptic digestion. The precursor mass tolerance was set
to 10 ppm and the fragment mass tolerance was set to 0.02 Da. Modifications were defined as dynamic Met
oxidation, and protein N-term acetylation with and without methionine-loss as well as static Cys
carbamidomethylation. A strict false discovery rate (FDR) < 1% (peptide and protein level) were required for
positive protein hits. If only 1 peptide per protein has been identified the hit was accepted if the Mascot score
was >30 or the MS Amanda score was >300 or the Sequest score was >4 or the MS Fragger score was >8. The
Percolator node of PD2.4 and a reverse decoy database was used for qvalue validation of spectral matches. Only
rank 1 proteins and peptides of the top scored proteins were counted. Label-free protein quantification was based
on the Minora algorithm of PD2.2 using a signal-to-noise ratio >5. Imputation of missing quan values was applied
by setting the abundance to 75% of the lowest abundance identified for each sample. Normalization was based
on a replicate median total peptide sum approach, which was calculated based on the sum of all identified peptide

abundance values per replicate sample. The sums of each of the three replicates from the four sample groups

were used to calculate median values. Normalization factors were calculated by dividing median values of the

respective sample group by the abundance sum of each sample. Normalization factors were multiplied with single

protein abundance values of each replicate/sample. The p-values are based on a Student`s t-test. Ratio-adjusted

p-values were calculated by dividing p-values with the log4ratio of the protein abundance levels. Significant

differences in protein abundance were defined when the following three requirements were reached: At least a

4-fold change in abundance (up and down), a ratio-adjusted p-value <0.05, and at least identified in 2 of 3

replicates of the sample group with the highest abundance. Intersection plots were created using the UpSetR

package (39) and only include proteins that were detected in at least two replicates of a given sample. Volcano

plots were created using ggplot2 in R using the replicate median total peptide sum normalized (RMN) data for all

proteins detected in **Dataset S1**.

Fungal mitochondrial reporter inhibition assay

To characterize growth inhibition and killing of hyphae by fungal-induced extracellular vesicles an *A. fumigatus*

strain expressing a mitochondrial GFP reporter (AfS35/pJW103; (35)) has been used as described previously (10).

Extracellular vesicles produced by $1x10^7$ dPLB cells or PMNs were isolated and co-incubated with the reporter

strain. After 16 hours of incubation samples were stained with CFW and images were taken using Zeiss LSM 780

confocal microscope (Carl Zeiss).

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE

partner repository (40) with the dataset identifier PXD027032.

Statistical Analysis. Data were plotted and statistically analyzed using GraphPad Prism software 5.0 (GraphPad

Software) unless otherwise noted. The Student's t test was used for significance testing when comparing two

groups. Differences between the groups were considered significant at a P value of <0.05. Throughout the article,

significance is denoted as follows: *, P = <0.05; **, P = <0.01, ***, P = <0.001; ns, nonsignificant.

Ethics Statement. Peripheral human blood was collected from healthy volunteers only after written informed

consent was provided. The study was conducted in accordance with the Declaration of Helsinki and approved by

the Ethics Committee of the University Hospital Jena (approval number 5074-02/17).

DISCUSSION

Neutrophils are critical players in the immune response to fungal pathogens stemming from their numerous antimicrobial capacities (10, 41). To study neutrophil functions, numerous cell lines that model these abilities *in vitro* have been considered with limited success. The most promising remains the HL-60 leukemia cell line and PLB-985 sublineage, which have been used as models for aspergillosis and bacterial killing (21-23, 28). In the present study, we analyze different aspects of the interaction between DMF-differentiated PLB-985 myeloid cells and the fungus *A. fumigatus* for the first time, making frequent comparisons with data from primary neutrophils.

Primary neutrophils are capable of phagocytosing and processing *A. fumigatus* conidia, features that were maintained in dPLB cells. As expected for cultured cells, the observed phagocytosis rate of *A. fumigatus* conidia by dPLB cells was lower than that of primary human PMNs. It is possible that additional differentiation methods will ultimately reveal higher rates of phagocytosis, as PLB-985 cells differentiated with DMF are known to exhibit lower phagocytic activity than cells differentiated with dimethyl sulfoxide (42). In terms of cytokine release, we observed increased release of IL-8 from dPLBs after infection with *A. fumigatus* opsonized conidia, but not IL-1β, consistent with our observations in human PMNs. These results are also in agreement with those collected from primary PMNs co-incubated with L-ficolin opsonized conidia for 8 hours, where secretion of IL-8 was also much higher than IL-1β (43).

We next stained dPLB cells for the endosomal marker LAMP-1 and measured the acidification of the phagolysosomes via Lysotracker to assess intracellular processing of fungal conidia. These experiments confirmed that around 20% of wild-type conidia are completely internalized inside the phagosomes and that lysosomal fusion occurs as a next step of conidial processing. Interestingly, and as already demonstrated for primary monocytes and neutrophils (25), melanized conidia showed less acidification around the phagolysosomal membrane than the mutant lacking DHN-melanin. This effect in monocytes is caused by inhibition of V-ATPase assembly. Surprisingly, staining for the V1 subunit of the V-ATPase did not reveal a significant difference between the wild-type and Δ*pksP* strains in dPLB cells as has been observed in other cell systems. These results suggest that neutrophils potentially employ additional mechanisms to acidify phagolysosomes in neutrophilic granulocytes after *A. fumigatus* infection or that the methods used were not sufficiently sensitive to detect slight differences in acidification. Together, these results suggest that dPLBs can serve as an intriguing model to study various aspects of neutrophil phagocytosis and intracellular processing in a more tractable *in vitro* model system.

A conidium that escapes phagocytosis can sometimes germinate into a hypha, a morphotype that is much
more difficult to eliminate and requires the antifungal activity of neutrophils. One mechanism that aids in control
of *A. fumigatus* hyphae is the release of fungistatic NETs composed of DNA and antimicrobial proteins. By staining
for extracellular DNA and NET constituent histone H3, we could clearly show that dPLB cells also produce NETs in
response to *A. fumigatus* hyphae, comparable to primary human PMNs and in agreement with other studies
challenging dPLBs with bacterial pathogens (21, 22).

Production of extracellular vesicles from primary neutrophils co-incubated with *Aspergillus* conidia or
bacteria like *Staphylococcus aureus* is an important defense strategy against microorganisms (10, 32). We found
that dPLB cells were also able to generate comparable populations of extracellular vesicles to PMNs upon contact
with opsonized conidia. Accumulation of extracellular vesicles of approximately 200 nm in diameter was observed
over time, independently of the isolation method we applied. Proteomic analysis revealed that by using a size-
exclusion chromatography-based isolation approach, the extracellular vesicles population could be enriched for
extracellular vesicle marker proteins like CD63, CD81, and TSG101. We observed that the samples obtained by
this method had a higher abundance of proteins and showed more differences between the spontaneously
released and infection-derived extracellular vesicles. In particular, infection-derived extracellular vesicles were
enriched in immunoglobulins and complement factors when compared to spontaneously released vesicles. These
serum proteins may be introduced to the infection system during opsonization of the fungal conidia despite
thorough washing. It remains unclear whether these immunoglobulins and complement factors play an important
role in extracellular vesicle biology during *A. fumigatus* infection. As a hint for such a role, the association of
immunoglobulins and complement proteins was shown previously to be important in systemic lupus
erythematosus (SLE) disease, where extracellular vesicles with immunoglobulin cargo could act as immune
complexes to mediate inflammation (44, 45).

In the proteomic analysis of extracellular vesicles derived from primary neutrophils infected with *A.*
fumigatus, several antimicrobial peptides, such as cathepsin G and azurocidin, were detected (10). Co-incubation
of these extracellular vesicles from primary neutrophils with fungal hyphae arrested growth, which was confirmed
in this study. Using dPLB cells, we identified extracellular vesicles containing cathepsin G and azurocidin; however,
the distribution remained relatively equal between spontaneously released and infection-derived extracellular
vesicles. We suspect that this is due to the differentiation procedure of dPLB cells with DMF, which provides a
mild inflammatory stimulus. Nevertheless, we observed mitochondrial damage in hyphae treated with infection-
derived but not spontaneously released extracellular vesicles. We can postulate several reasons for this result.
First, it is possible that immunoglobulins are playing a role in targeting extracellular vesicles to fungal hyphae. In
this case, the protein cargo may be the same, but targeting would be inefficient in spontaneously released

extracellular vesicles. Of note, opsonization of the bacterial pathogen *Staphylococcus aureus* was not found to influence the antimicrobial capacity of neutrophil-derived extracellular vesicles (32). A second option is that additional extracellular vesicle cargo molecules such as RNA or lipids may play a role in the antifungal activity. It has been shown in many instances that extracellular vesicles can contain small RNAs that exert distinct functions during infection, especially in regard to plant fungal pathogens (46, 47). We believe that a combination of factors is likely involved in the antifungal effect of extracellular vesicles.

Part of the explanation may also come from similarities between neutrophilic granules and extracellular vesicles. Granules and extracellular vesicles are distinct cellular features, but do share some biogenesis pathways and cargo molecules. For example, the azurophilic granules contain myeloperoxidase to aid in pathogen killing in the endocytic pathway via fusion of the granules to pathogen-containing phagosomes (48), but myeloperoxidase is also released in a CD63-negative population of extracellular vesicles known as microvesicles (49). Based on the proteomics analysis of this study, as well as our previous work (10), it seems possible that microvesicle-cargo proteins like myeloperoxidase contribute at least in part to the antifungal activity. However, the presence of myeloperoxidase in spontaneously released extracellular vesicles and the lack of susceptibility of patients lacking myeloperoxidase to fungal infections (50) hint that other factors are also involved. Besides, the shared marker protein CD63 nicely highlights another link between granules and extracellular vesicles, the shared biogenesis from multivesicular bodies of azurophilic granules and the extracellular vesicle subset called exosomes (51).

In conclusion our results suggest that DMF-differentiated PLB-985 cells can be used as a model to study aspects of the interaction of the human pathogenic fungus *A. fumigatus* with neutrophilic granulocytes. Although they will never substitute for all experiments with neutrophils, we do believe that this system will serve as a useful tool for the genetic dissection of *A. fumigatus* pathogenesis in the future.

ACKNOWLEDGEMENTS

We would like to thank Johannes Wagener for providing the *A. fumigatus* mitochondrial GFP reporter strain. We also thank Natascha Wilker for excellent technical assistance. The work presented here was generously supported by the Deutsche Forschungsgemeinschaft (DFG)-funded Collaborative Research Center/Transregio FungiNet 124 'Pathogenic fungi and their human host: Networks of Interaction' (210879364, project A1 and Z2), the DFG-funded ANR project (316898429, project AfuInf), and by the Federal Ministry for Education and Research (BMBF: https://www.bmbf.de/), Germany, Project FKZ 01KI2012 "RFIN – RNA-Biologie von Pilzinfektionen." The authors declare no conflicts of interest.

REFERENCES

1. Brown GD, Denning DW, Gow NA, Levitz SM, Netea MG, White TC. 2012. Hidden killers: human fungal infections. Sci Transl Med 4:165rv13.

2. Schmiedel Y, Zimmerli S. 2016. Common invasive fungal diseases: an overview of invasive candidiasis, aspergillosis, cryptococcosis, and Pneumocystis pneumonia. Swiss Med Wkly 146:w14281.

3. Latge JP, Chamilos G. 2019. *Aspergillus fumigatus* and Aspergillosis in 2019. Clin Microbiol Rev 33.

4. Nasir N, Farooqi J, Mahmood SF, Jabeen K. 2020. COVID-19-associated pulmonary aspergillosis (CAPA) in patients admitted with severe COVID-19 pneumonia: An observational study from Pakistan. Mycoses 63:766-770.

5. Salmanton-Garcia J, Sprute R, Stemler J, Bartoletti M, Dupont D, Valerio M, Garcia-Vidal C, Falces-Romero I, Machado M, de la Villa S, Schroeder M, Hoyo I, Hanses F, Ferreira-Paim K, Giacobbe DR, Meis JF, Gangneux JP, Rodriguez-Guardado A, Antinori S, Sal E, Malaj X, Seidel D, Cornely OA, Koehler P, FungiScope European Confederation of Medical Mycology/The International Society for H, Animal Mycology Working G. 2021. COVID-19-Associated Pulmonary Aspergillosis, March-August 2020. Emerg Infect Dis 27:1077-1086.

6. Rivera A, Hohl TM, Collins N, Leiner I, Gallegos A, Saijo S, Coward JW, Iwakura Y, Pamer EG. 2011. Dectin-1 diversifies *Aspergillus fumigatus*-specific T cell responses by inhibiting T helper type 1 CD4 T cell differentiation. J Exp Med 208:369-81.

7. Brakhage AA, Langfelder K. 2002. Menacing mold: the molecular biology of *Aspergillus fumigatus*. Annu Rev Microbiol 56:433-55.

8. Drummond RA, Gaffen SL, Hise AG, Brown GD. 2014. Innate Defense against Fungal Pathogens. Cold Spring Harb Perspect Med 5.

9. Mircescu MM, Lipuma L, van Rooijen N, Pamer EG, Hohl TM. 2009. Essential role for neutrophils but not alveolar macrophages at early time points following *Aspergillus fumigatus* infection. J Infect Dis 200:647-56.

10. Shopova IA, Belyaev I, Dasari P, Jahreis S, Stroe MC, Cseresnyes Z, Zimmermann AK, Medyukhina A, Svensson CM, Kruger T, Szeifert V, Nietzsche S, Conrad T, Blango MG, Kniemeyer O, von Lilienfeld-Toal M, Zipfel PF, Ligeti E, Figge MT, Brakhage AA. 2020. Human Neutrophils Produce Antifungal Extracellular Vesicles against *Aspergillus fumigatus*. mBio 11.

11. McCormick A, Heesemann L, Wagener J, Marcos V, Hartl D, Loeffler J, Heesemann J, Ebel F. 2010. NETs formed by human neutrophils inhibit growth of the pathogenic mold *Aspergillus fumigatus*. Microbes Infect 12:928-36.

12. Brogan TD. 1964. Effect of Complement and of the Carbohydrate Components of Sputum on Phagocytosis by Human Polymorphonuclear Leucocytes. Immunology 7:626-38.

13. Pazhakh V, Ellett F, Croker BA, O'Donnell JA, Pase L, Schulze KE, Greulich RS, Gupta A, Reyes-Aldasoro CC, Andrianopoulos A, Lieschke GJ. 2019. beta-glucan-dependent shuttling of conidia from neutrophils to macrophages occurs during fungal infection establishment. PLoS Biol 17:e3000113.

14. Silva JC, Rodrigues NC, Thompson-Souza GA, Muniz VS, Neves JS, Figueiredo RT. 2020. Mac-1 triggers neutrophil DNA extracellular trap formation to *Aspergillus fumigatus* independently of PAD4 histone citrullination. J Leukoc Biol 107:69-83.

15. Bruns S, Kniemeyer O, Hasenberg M, Aimanianda V, Nietzsche S, Thywissen A, Jeron A, Latge JP, Brakhage AA, Gunzer M. 2010. Production of extracellular traps against *Aspergillus fumigatus* in vitro and in infected lung tissue is dependent on invading neutrophils and influenced by hydrophobin RodA. PLoS Pathog 6:e1000873.

16. Clark HL, Abbondante S, Minns MS, Greenberg EN, Sun Y, Pearlman E. 2018. Protein Deiminase 4 and CR3 Regulate *Aspergillus fumigatus* and beta-Glucan-Induced Neutrophil Extracellular Trap Formation, but Hyphal Killing Is Dependent Only on CR3. Front Immunol 9:1182.

17. Zarember KA, Sugui JA, Chang YC, Kwon-Chung KJ, Gallin JI. 2007. Human polymorphonuclear leukocytes
inhibit *Aspergillus fumigatus* conidial growth by lactoferrin-mediated iron depletion. J Immunol 178:6367-
73.
18. Gazendam RP, van Hamme JL, Tool AT, Hoogenboezem M, van den Berg JM, Prins JM, Vitkov L, van de
Veerdonk FL, van den Berg TK, Roos D, Kuijpers TW. 2016. Human Neutrophils Use Different Mechanisms
To Kill *Aspergillus fumigatus* Conidia and Hyphae: Evidence from Phagocyte Defects. J Immunol 196:1272-
83.
19. Lahoz-Beneytez J, Elemans M, Zhang Y, Ahmed R, Salam A, Block M, Niederalt C, Asquith B, Macallan D.
2016. Human neutrophil kinetics: modeling of stable isotope labeling data supports short blood neutrophil
half-lives. Blood 127:3431-8.
20. Blanter M, Gouwy M, Struyf S. 2021. Studying Neutrophil Function in vitro: Cell Models and Environmental
Factors. J Inflamm Res 14:141-162.
21. Marin-Esteban V, Turbica I, Dufour G, Semiramoth N, Gleizes A, Gorges R, Beau I, Servin AL, Lievin-Le Moal
V, Sandre C, Chollet-Martin S. 2012. Afa/Dr diffusely adhering *Escherichia coli* strain C1845 induces
neutrophil extracellular traps that kill bacteria and damage human enterocyte-like cells. Infect Immun
80:1891-9.
22. Amulic B, Knackstedt SL, Abu Abed U, Deigendesch N, Harbort CJ, Caffrey BE, Brinkmann V, Heppner FL,
Hinds PW, Zychlinsky A. 2017. Cell-Cycle Proteins Control Production of Neutrophil Extracellular Traps.
Developmental Cell 43:449-462.e5.
23. Baistrocchi SR, Lee MJ, Lehoux M, Ralph B, Snarr BD, Robitaille R, Sheppard DC. 2017. Posaconazole-
Loaded Leukocytes as a Novel Treatment Strategy Targeting Invasive Pulmonary Aspergillosis. J Infect Dis
215:1734-1741.
24. Goyette G, Boulais J, Carruthers NJ, Landry CR, Jutras I, Duclos S, Dermine JF, Michnick SW, LaBoissiere S,
Lajoie G, Barreiro L, Thibault P, Desjardins M. 2012. Proteomic characterization of phagosomal membrane
microdomains during phagolysosome biogenesis and evolution. Mol Cell Proteomics 11:1365-77.
25. Thywissen A, Heinekamp T, Dahse HM, Schmaler-Ripcke J, Nietzsche S, Zipfel PF, Brakhage AA. 2011.
Conidial Dihydroxynaphthalene Melanin of the Human Pathogenic Fungus *Aspergillus fumigatus*
Interferes with the Host Endocytosis Pathway. Front Microbiol 2:96.
26. Heinekamp T, Schmidt H, Lapp K, Pahtz V, Shopova I, Koster-Eiserfunke N, Kruger T, Kniemeyer O,
Brakhage AA. 2015. Interference of *Aspergillus fumigatus* with the immune response. Semin
Immunopathol 37:141-52.
27. Schmidt F, Thywissen A, Goldmann M, Cunha C, Cseresnyes Z, Schmidt H, Rafiq M, Galiani S, Graler MH,
Chamilos G, Lacerda JF, Campos A, Jr., Eggeling C, Figge MT, Heinekamp T, Filler SG, Carvalho A, Brakhage
AA. 2020. Flotillin-Dependent Membrane Microdomains Are Required for Functional Phagolysosomes
against Fungal Infections. Cell Rep 32:108017.
28. Pedruzzi E, Fay M, Elbim C, Gaudry M, Gougerot-Pocidalo MA. 2002. Differentiation of PLB-985 myeloid
cells into mature neutrophils, shown by degranulation of terminally differentiated compartments in
response to N-formyl peptide and priming of superoxide anion production by granulocyte-macrophage
colony-stimulating factor. Br J Haematol 117:719-26.
29. Volk AP, Barber BM, Goss KL, Ruff JG, Heise CK, Hook JS, Moreland JG. 2011. Priming of neutrophils and
differentiated PLB-985 cells by pathophysiological concentrations of TNF-alpha is partially oxygen
dependent. J Innate Immun 3:298-314.
30. Branzk N, Papayannopoulos V. 2013. Molecular mechanisms regulating NETosis in infection and disease.
Semin Immunopathol 35:513-30.
31. Xie K, Varatnitskaya M, Maghnouj A, Bader V, Winklhofer KF, Hahn S, Leichert LI. 2020. Activation leads to
a significant shift in the intracellular redox homeostasis of neutrophil-like cells. Redox Biol 28:101344.

32. Timar CI, Lorincz AM, Csepanyi-Komi R, Valyi-Nagy A, Nagy G, Buzas EI, Ivanyi Z, Kittel A, Powell DW,
McLeish KR, Ligeti E. 2013. Antibacterial effect of microvesicles released from human neutrophilic
granulocytes. Blood 121:510-8.
33. Brakhage AA, K. ZA, Rivieccio F, Visser C, Blango MG. 2021. Host-derived extracellular vesicles for
antimicrobial defense. microLife doi:10.1093/femsml/uqab003/6228840.
34. Lane RE, Korbie D, Trau M, Hill MM. 2019. Optimizing Size Exclusion Chromatography for Extracellular
Vesicle Enrichment and Proteomic Analysis from Clinically Relevant Samples. Proteomics 19:e1800156.
35. Ruf D, Brantl V, Wagener J. 2018. Mitochondrial Fragmentation in *Aspergillus fumigatus* as Early Marker
of Granulocyte Killing Activity. Front Cell Infect Microbiol 8:128.
36. Hillmann F, Novohradska S, Mattern DJ, Forberger T, Heinekamp T, Westermann M, Winckler T, Brakhage
AA. 2015. Virulence determinants of the human pathogenic fungus *Aspergillus fumigatus* protect against
soil amoeba predation. Environ Microbiol 17:2858-69.
37. Kraibooj K, Schoeler H, Svensson CM, Brakhage AA, Figge MT. 2015. Automated quantification of the
phagocytosis of *Aspergillus fumigatus* conidia by a novel image analysis algorithm. Front Microbiol 6:549.
38. Wessel D, Flugge UI. 1984. A method for the quantitative recovery of protein in dilute solution in the
presence of detergents and lipids. Anal Biochem 138:141-3.
39. Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. 2014. UpSet: Visualization of Intersecting Sets.
IEEE Trans Vis Comput Graph 20:1983-92.
40. Perez-Riverol Y, Xu QW, Wang R, Uszkoreit J, Griss J, Sanchez A, Reisinger F, Csordas A, Ternent T, Del-
Toro N, Dianes JA, Eisenacher M, Hermjakob H, Vizcaino JA. 2016. PRIDE Inspector Toolsuite: Moving
Toward a Universal Visualization Tool for Proteomics Data Standard Formats and Quality Assessment of
ProteomeXchange Datasets. Mol Cell Proteomics 15:305-17.
41. Feldmesser M. 2006. Role of neutrophils in invasive aspergillosis. Infect Immun 74:6514-6.
42. Pivot-Pajot C, Chouinard FC, El Azreq MA, Harbour D, Bourgoin SG. 2010. Characterisation of
degranulation and phagocytic capacity of a human neutrophilic cellular model, PLB-985 cells.
Immunobiology 215:38-52.
43. Bidula S, Sexton DW, Schelenz S. 2016. Serum opsonin ficolin-A enhances host-fungal interactions and
modulates cytokine expression from human monocyte-derived macrophages and neutrophils following
Aspergillus fumigatus challenge. Med Microbiol Immunol 205:133-42.
44. Buzas EI, Toth EA, Sodar BW, Szabo-Taylor KE. 2018. Molecular interactions at the surface of extracellular
vesicles. Semin Immunopathol 40:453-464.
45. Perez-Hernandez J, Redon J, Cortes R. 2017. Extracellular Vesicles as Therapeutic Agents in Systemic Lupus
Erythematosus. Int J Mol Sci 18.
46. Cai Q, Qiao L, Wang M, He B, Lin FM, Palmquist J, Huang SD, Jin H. 2018. Plants send small RNAs in
extracellular vesicles to fungal pathogen to silence virulence genes. Science 360:1126-1129.
47. He B, Cai Q, Qiao L, Huang C-Y, Wang S, Miao W, Ha T, Wang Y, Jin H. 2021. RNA-binding proteins
contribute to small RNA loading in plant extracellular vesicles. Nature Plants doi:10.1038/s41477-021-
00863-8.
48. Klebanoff SJ, Kettle AJ, Rosen H, Winterbourn CC, Nauseef WM. 2013. Myeloperoxidase: a front-line
defender against phagocytosed microorganisms. J Leukoc Biol 93:185-98.
49. Hess C, Sadallah S, Hefti A, Landmann R, Schifferli JA. 1999. Ectosomes released by human neutrophils are
specialized functional units. J Immunol 163:4564-73.
50. Desai JV, Lionakis MS. 2018. The role of neutrophils in host defense against invasive fungal infections. Curr
Clin Microbiol Rep 5:181-189.
51. Edgar JR. 2016. Q&A: What are exosomes, exactly? BMC Biol 14:46.

FIGURES AND FIGURE LEGENDS

FIG 1. Phagocytosis of *A. fumigatus* conidia by PLB-985 cells.

(A). Confocal microscopy images of wild-type conidia (FITC-stained; green) phagocytosed by dPLB cells after 4 hours co-incubation. Membranes were stained with Cell Mask (red). Images are representative of three independent biological experiments. Scale bars are 10 μm. **(B)** Representative images of wild-type FITC-stained conidia phagocytosed by dPLB cells after 4 hours co-infection obtained by imaging flow cytometry (top two rows). External conidia are visualized by counter staining with CFW (purple), as seen in the bottom row. Images are representative of three independent experiments. **(C)** Quantification of PLB-985 cell phagocytosis using the IDEAs software on 5,000 cells per condition with either wild-type or Δ*pksP* conidia from four independent experiments

(D) ELISA detection of IL-8 cytokine released from infected dPLBs and primary human PMNs at different time

points. Lipopolysaccharide (LPS) was included for comparison of a bacterial stimulus. Data are presented as mean

±SD, from six biological replicates; * = p < 0.05.

FIG 2. Processing of *A. fumigatus* conidia inside phagolysosomes.

(A) dPLB cells were stained with LAMP-1 (red) after infection with wild-type or Δ*pksP c*onidia (FITC labeled; green)

for 4 h. Images are representative of three biological replicates. **(B)** Quantification of LAMP-1 co-localization with

CFW-labelled conidia in phagolysosomes and **(C)** Percentage of acidified conidia in dPLB cells after 2- and 4-hours

post-infection with wild-type and Δ*pksP c*onidia. Data are presented as mean ±SD; *** = p < 0.001. **(D)**

Colocalization of conidia (CFW; blue) in acidified compartments labeled with Lysotracker (red). Images are

representative of three biological replicates. Scale bars are 10 μm.

FIG 3. Production of NETs in response to *A. fumigatus* infection.

(A) Confocal scanning laser microscopy of nucleic acid stained with DAPI (blue) released from dPLB cells stained

with Cell Mask (red) after challenge with fungal hyphae containing a mitochondrial GFP reporter (*A. fumigatus*

strain AfS35 / pJW103 expressing a mitochondrial GFP reporter; green). Phorbol myristate acetate (PMA) was

used as a positive control. Data representative of three biological replicates. **(B)** Confocal micrographs of NET

markers, Histone H3 (red) and nucleic acid stained with DAPI (blue), produced by dPLB cells after contact with *A.*

fumigatus hyphae (FITC-labeled; green). Images are representative of three biological replicates. Scale bars are 10

μm.

FIG 4. dPLB cells produce extracellular vesicles comparable in size to primary human PMNs.

Extracellular vesicles released spontaneously or produced in response to infection with opsonized *A. fumigatus*

conidia were isolated using a differential centrifugation-based approach (DC) or a size-exclusion chromatography-

based approach (SEC) and quantified using nanoparticle tracking analysis. Extracellular vesicles were quantified

from **(A)** dPLB cells or **(B)** primary human PMNs at 0, 2, and 4 hours post infection and show the average of at

least six biological replicates and three biological replicates, respectively. Representative size histograms from five

biological replicates are shown for extracellular vesicles derived from **(C)** dPLB cells and **(D)** primary human PMNs.

FIG 5. SEC enriches for extracellular vesicle populations.

LC-MS/MS proteomics analysis was performed on extracellular vesicles isolated from dPLB cells using a differential

centrifugation-based approach (DC) or a size-exclusion chromatography-based approach (SEC) in the presence or

absence of infection with opsonized *A. fumigatus* conidia. **(A)** Proteins identified in spontaneously released

extracellular vesicles (sEVs) and infection-derived extracellular vesicles (iEVs) from at least two replicates of a

given sample were intersected using UpSetR. The red bar indicates proteins that were found in all four samples.

Volcano plots show the $\log_2$ ratio of infection-derived extracellular vesicles (iEVs) versus spontaneously released

EVs (sEVs) for **(B)** DC-based isolation and **(C)** SEC-based isolation. Input data included values from all replicates

using the RMN data included in **Dataset S1**. Plots were created using ggplot2 in R. Proteomics data is from three

analytical replicates of three independent biological replicates. Orange circles represent proteins with greater

than 2-fold change and $p = < 0.05$. Selected proteins are named for clarity.

FIG 6. Infection-derived extracellular vesicles from dPLBs are antifungal to *A. fumigatus* hyphae.

(A) Conidia from *A. fumigatus* strain AfS35 containing plasmid pJW103 expressing a mitochondrial GFP reporter (green) were opsonized and co-incubated with freshly harvested human PMNs or dPLBs. After overnight incubation (22 hours) samples were stained with CFW for 10 min and images were taken using a Zeiss LSM 780 confocal microscope. Scale bars are 20 μm. **(B)** The *A. fumigatus* strain AfS35 expressing a mitochondrial GFP reporter (green) was first grown for 6 hours and then stained with calcofluor white (blue) and incubated overnight with spontaneously released extracellular vesicles or infection-derived extracellular vesicles isolated from primary human PMNs or dPLBs. As a control, untreated hyphae and hyphae treated with 3 mM H_2O_2 to induce cell death

are included. An intact mitochondrial network is shown by a filamentous network, whereas a disrupted network

is shown by fragmentation or the lack of green signal. Scale bars are 5 μm.

TABLES

Table 1. RMN-fold change of selected proteins identified in extracellular vesicles (EVs) from dPLB cells.

Abbreviations: DC, differential centrifugation-based approach; SEC, size-exclusion chromatography-based

approach.

Gene	Accession	Infection-derived EVs (DC) / Spontaneous EVs (DC)	Infection-derived EVs (SEC) / Spontaneous EVs (SEC)	Infection-derived EVs (SEC) / Infection-derived EV (DCs)	Spontaneous EVs (SEC) / Spontaneous EVs (DC)
EV Markers					
CD63	F8VWK8	1.274	1.424	1.478	1.323
TSG101	F5H442	1.215	1.630	1.554	1.158
CD81	E9PRJ8	1.150	1.413	2.055	1.673
Non-EV Markers					
CANX (Calnexin)	P27824	-1.031	-1.082	-1.468	-1.398
MPO (Myeloperoxidase)	P05164	-1.024	1.026	2.063	1.963
CTSG (Cathepsin G)	P08311	1.045	1.042	1.117	1.120
AZU1 (Azurocidin)	P20160	-1.141	1.029	1.521	1.295

SUPPLEMENTAL FIGURES

FIG S1. Primary human neutrophils phagocytose wild-type conidia.

(A) Confocal microscopy images of wild-type conidia (CFW-stained; false-colored green) phagocytosed by primary human PMNs after 2 hours co-incubation. Membranes were stained with Cell Mask (red) to indicate cell membranes. Images are representative of at least two independent biological experiments. Scale bars are 10 µm. **(B and C)** indicate the gating strategy for determining phagocytosis rates of dPLB cells and PMNs using imaging flow cytometry. **(B)** Single cell populations were selected based on their area and aspect ratio in the bright field channel. **(C)** Phagocytosed conidia were gated based on their higher FITC fluorescence intensity and low CFW fluorescence intensity. Cells without conidia had low FITC fluorescence intensity. The data subject to analysis

764   included only cells in focus and were pre-gated during data acquisition based on their high root mean square
765   gradient in the bright field channel. **(D)** Quantification of phagocytosis by primary human PMNs using the IDEAs
766   software on 5,000 cells per condition with either wild-type or Δ*pksP* conidia from four independent experiments.
767

768
769   **FIG S2. Cytokine production by dPLB cells and PMNs.**
770   ELISA detection of the IL-1β cytokine released from infected dPLBs or primary human PMNs at different time
771   points. Lipopolysaccharide (LPS) was included as a comparison to a bacterial stimulus. Data are presented as mean
772   ±SD, from six biological replicates; * = p < 0.05.
773
774

775
FIG S3. V-ATPase V1 localization to phagolysosomal membranes and ROS production by dPLB cells.
(A) Immunofluorescence staining for V-ATPase V1 (red) in dPLB cells after infection with wild-type or Δ*pksP c*onidia
(FITC labeled; green) for 4 hours. Images are representative of three biological replicates. Scale bars are 10 μm.
(B) Quantification of V-ATPase V1 positive phagolysosomes after 4 hours of infection with wild-type or Δ*pksP*
conidia. Graphs are from three biological replicates. **(C)** Confocal microscopy images of dPLB cells infected with

wild-type conidia (CFW-stained; blue), left uninfected, or treated with PMA as a positive control and stained for
detection of reactive oxygen species using CellROX Orange stain.
783

784
FIG S4. Intersection of proteomics data with previously reported neutrophil-derived EV proteomes.
LC-MS/MS proteomics analysis was performed on extracellular vesicles isolated from dPLB cells using a
centrifugation-based approach (DC) or a size-exclusion chromatography-based approach (SEC) in the presence or
absence of infection with opsonized *A. fumigatus* conidia and compared to data from (10). Proteins that were
identified in this study were compared to proteins identified by a similar approach using label-free quantification
from Shopova et al. (10). Proteins had to be found in at least two replicate of a given sample to be included in the
UpSetR analysis. The red bar indicates proteins that were found in all six samples.
792
793

795
FIG S5. Representative images of dPLB-derived extracellular vesicles effects on *A. fumigatus* hyphae.
A. fumigatus strain AfS35 contains plasmid pJW103, which expresses a mitochondrial GFP reporter (green). The

strain was grown for 6 hours and then stained with calcofluor white (blue) and incubated overnight with

spontaneously released extracellular vesicles or infection-derived extracellular vesicles from dPLBs. Additional

representative images are included to show extent of variability that occurs in regard to extracellular vesicle killing

experiments shown in **Fig. 6**. As a control, untreated hyphae and hyphae treated with 3 mM H_2O_2 to induce cell

death are included. An intact mitochondrial network is shown by a filamentous network, whereas a disrupted

network is shown by fragmentation or the lack of green signal. Scale bars are 5 µm.

SUPPLEMENTAL TABLES

Table S1. Proteins identified only in infection-derived extracellular vesicles (≥2 replicates) by both isolation

methods. Abbreviations: PSMs, peptide spectrum matches; AAs, amino acids; Cov%, percent coverage of

mapped peptides.

Gene	Accession	Description	(PSMs/AAs)*(Cov%/100)
IGKC	P01834	Immunoglobulin kappa constant	3.45421
APOA1	P02647	Apolipoprotein A-I	1.73397
IGHM	P01871	Immunoglobulin heavy constant mu	1.26349
IGLC2	P0DOY2	Immunoglobulin lambda constant 2	0.82453
IGHG1	P01857	Immunoglobulin heavy constant gamma 1	0.51688
HBB	P68871	Hemoglobin subunit beta	0.43599
IGHG2	A0A286YEY4	Immunoglobulin heavy constant gamma 2 (Fragment)	0.27205
APOB	P04114	Apolipoprotein B-100	0.26471
IGHA1	P01876	Immunoglobulin heavy constant alpha 1	0.16677
C1QB	P02746	Complement C1q subcomponent subunit B	0.16126
NPTN	Q9Y639	Neuroplastin	0.12879
APCS	P02743	Serum amyloid P-component	0.12233
C1QC	P02747	Complement C1q subcomponent subunit C	0.07286
APOH	P02749	Beta-2-glycoprotein 1	0.06858
APOD	P05090	Apolipoprotein D	0.06349
EMB	D6RDX7	Embigin	0.04667
CFH	P08603	Complement factor H	0.02729
IGKV2-40	A0A087WW87	Immunoglobulin kappa variable 2-40	0.02545
DNAJA1	P31689	DnaJ homolog subfamily A member 1	0.01884
HPX	P02790	Hemopexin	0.01541
C4B	F5GXS0	C4a anaphylatoxin	0.00783
FN1	P02751	Fibronectin	0.00560
C3	P01024	Complement C3	0.00394
C8A	P07357	Complement component C8 alpha chain	0.00394
AHSG	P02765	Alpha-2-HS-glycoprotein	0.00229
C1R	B4DPQ0	Complement subcomponent C1r	0.00118

813 **SUPPLEMENTAL DATASETS**

815 **Dataset S1. Proteomics data of extracellular vesicles produced in response to *A. fumigatus* infection by dPLB**
816 **cells.**