Microbiome structure of a wild *Drosophila* community along

tropical elevational gradients and comparison to laboratory lines

Joel J. Brown[* 1, 2, 3]**, Anna Jandová**[3]**, Christopher T. Jeffs**[4]**, Megan Higgie**[5]**, Eva Nováková**[2, 6]**,**

Owen T. Lewis[4] **& Jan Hrček**[2, 3]

[1] Cornell University, Department of Entomology, Ithaca, New York, USA (current address)

[2] University of South Bohemia, Faculty of Science, České Budějovice, Czech Republic

[3] Institute of Entomology, Biology Centre of the Czech Academy of Sciences, České Budějovice, Czech Republic

[4] Department of Zoology, University of Oxford, Oxford, U.K.

[5] College of Science & Engineering, James Cook University, Townsville, QLD, Australia

[6] Institute of Parasitology, Biology Centre of the Czech Academy of Sciences, České Budějovice, Czech Republic

[*] Corresponding author: joeljbrown15@gmail.com

Statement of authorship: JJB, CTJ, MH, OTL, JH designed the research. CTJ and OTL

executed the fieldwork. JJB, AJ, EN performed the molecular work. JJB analysed the data. JJB

wrote the first version of the manuscript. All authors contributed to the improvement of the

manuscript.

Conflict of interest: The authors declare no conflict of interest.

Running title: Microbiome along elevation and wild vs lab

Keywords: Bacteria; community; *Drosophila;* ecology; elevation gradient; field and

laboratory; metabarcoding; microbiome; symbiosis

Abstract

Variation along environmental gradients in host-associated microbial communities is not

well understood, compared to free-living microbial communities. Because elevational

gradients may serve as natural proxies for climate change, understanding patterns along

these gradients can inform our understanding of the threats hosts and their symbiotic

microbes face in a warming world. In this study, we analysed bacterial microbiomes from

pupae and adults of four *Drosophila* species native to Australian tropical rainforests. We

sampled wild individuals at high and low elevation along two mountain gradients, to

determine natural diversity patterns. Further, we sampled laboratory-reared individuals

from isofemale lines established from the same localities, to see if any natural patterns are

retained in the lab. In both environments, we controlled for diet to help elucidate other

deterministic patterns of microbiome composition. We found small but significant

differences in *Drosophila* bacterial community composition across elevation, with some

notable taxonomic differences between different *Drosophila* species and sites. Further, we

found that field-collected fly pupae had significantly richer microbiomes than laboratory-

reared pupae. We also found similar microbiome composition in both types of provided

diet, suggesting that the significant differences found among *Drosophila* microbiomes are

the product of surrounding environments with different bacterial species pools, possibly

bound to elevational differences in temperature. Our results suggest that comparative

studies between lab and field specimens help reveal the true variability in microbiome

communities that can exist within a single species.

Importance

Bacteria form microbial communities inside most higher-level organisms, but we know little about how the microbiome varies along environmental gradients and between natural host populations and laboratory colonies. To explore such effects on insect-associated microbiomes, we studied gut microbiome in four *Drosophila* species over two mountain gradients in tropical Australia. We also compared these data to individuals kept in the laboratory to understand how different settings changed microbiome communities. We found that field sampled individuals had significantly higher microbiome diversity than those from the lab. In wild *Drosophila* populations, elevation explains a small but significant amount of the variation in their microbial communities. Our study highlights the importance of environmental bacterial sources for *Drosophila* microbiome composition across elevational gradients, and shows how comparative studies help reveal the true flexibility in microbiome communities that can exist within a species.

Introduction

Patterns of diversity over environmental gradients like latitude, elevation, or environmental degradation have long been of interest in community ecology and are of renewed interest for studying the potential consequences of climate change[1–5]. Most studies have focused on animals and plants to investigate these patterns, but bacterial communities are receiving increased attention. Some studies suggest free-living bacteria do not follow the same broad biogeographic patterns as plants and animals[6–8]. Fierer et al.[1] showed that soil bacteria did not change significantly in diversity when sampled across an elevational gradient, in contrast to trends documented in most other taxa. Subsequent studies have found

inconsistent patterns in bacterial communities sampled from streams and soils across

elevational gradients, with differences usually being attributed to changes in pH and C:N

ratio [2,8–10].

Many insects maintain intimate communities of symbiotic microbes (their 'microbiome').

Insect microbiomes can play important roles in host health, digestion, thermal regulation,

and protection against natural enemies (reviewed in[11–13]). In turn, many factors can

influence insect microbiome composition, some host-dependent (e.g. diet, insect species

identity, ontogeny, and parent-to-offspring transmission), others host-independent (e.g.

abiotic factors like local environment and temperature)[14–22]. Symbioses between insects and

bacteria have been particularly well investigated[23], notably because insect microbiome

communities tend to be less complex than those of vertebrates[24]. However, in contrast to

environmental microbial communities, the effect of elevational change on insect-associated

microbiomes has yet to be investigated in-depth. The most conspicuous aspect of a change

in elevation is a difference in mean temperature, creating different environments that can

be used as a proxy for climate change scenarios[25,26]. Elevational differences in temperature

mean we would expect to see differences in microbiome composition related to

temperature-dependent development in both insects[27–31] and bacteria[32–34]. Thus, at

different elevations, and in climate change scenarios, insect-associated microbiomes could

have different composition[20] with potentially important consequences on microbiome and

host functionality.

Drosophila spp. are established models for studying insect-associated microbiomes[35–40]

because they are cosmopolitan, occurring in a wide variety of habitats, and easy to maintain

in laboratory cultures. *Drosophila*-associated microbiomes have important functional

impacts on many aspects of their ecology including thermal tolerance[41], development[42],

ability to recognise kin[43], and immunity[38,44]. The microbiomes are of moderate-to-low

diversity, making them relatively simple to characterise. Additionally, some *Drosophila*

species/populations possess intracellular bacterial symbionts (*Wolbachia* and *Spiroplasma*)

that can influence host immunity and protect against natural enemies, including pathogenic

fungi, nematodes, and parasitoids[45–49]. This, combined with the well-studied nature of

Drosophila, makes them ideal candidates for investigating insect-associated microbiomes

over elevational gradients.

Here we examine the effects of elevation change on insect microbiome composition by

focusing on the underlying biotic and abiotic factors, including elevation, host species

identity, and site location. We sampled wild populations of four focal species of frugivorous

Drosophila from two mountain gradients in Queensland, Australia: *Drosophila rubida, D.*

pseudoananassae, D. pallidifrons, and *D. sulfurigaster.* These species occur throughout

north Queensland along multiple altitudinal gradients in the Wet Tropics. We chose these

four species because they are abundant and occur in sympatry across the full elevational

gradient at our focal sites[50]. We hypothesised that variation in microbiome composition

between high and low elevation populations will reflect temperature differences at these

sites. To control for diet in the field we exclusively sampled pupae from banana-baited

bottle traps (see Jeffs et al.[50]). The sampling approach guarantees that each analysed

individual originated from an egg laid in our bottle traps and therefore fed solely on yeasted

banana. To reinforce our investigation, we analysed lab-reared flies of the same species

collected from the same field sites to test if their microbiomes retained any natural

differences when reared in the laboratory on a standard, yeast-based diet. We expected *a*

priori to find high among-individual variation and hypothesised that species identity,

elevation, and environment of origin (i.e. lab vs field) would be the primary causes of

difference in host microbiome composition[35,39,40,51].

2. Material & Methods

2.1 Study Sites

The Australian Wet Tropics World Heritage Area is a 450 km long, narrow section of

rainforest along Queensland's northeast coast between Cooktown and Townsville (15-19'S,

145-146.30'E). Samples were collected from two altitudinal gradients: Paluma Range Road

(within Paluma Range National Park 19°00'S, 146°14'E) and Kirrama Range Road (within

Girramay National Park 18°12'S, 145°50'E). The Paluma gradient ranges from 59 m to 916 m

above sea level (a.s.l.) and the Kirrama gradient ranges from 92 m to 770 m a.s.l[50]. We chose

sites at high and low elevations (Paluma: 880m, 70m; Kirrama: 730m, 70m) to capture a

~5°C temperature range (mean temperatures 21°C at high elevation, 26°C at low

elevation[50]. Temperature was recorded by multiple dataloggers suspended next to bottle

traps at each site, which took a reading every hour. Our previous study on these gradients

determined ~5°C temperature range and found substantial differences in *Drosophila*

community composition at high and low elevation, with some species not present at the low

and high ends of the gradient and others changing in abundance[50].

2.2 Sample Collection and Selection

To ensure a comparable dataset, we selected stratified subsets of samples from the field and laboratory. We primarily collected pupae because they are more easily standardized than larvae and allow us to account for discrepancies in development rate between species. We additionally collected a small number of adult flies to compare microbiome composition between pupae and adults. Based on the results of Jeffs et al.[50], which used pupae samples to identify the natural *Drosophila*-parasitoid food web with COI metabarcoding and Multiplex PCR methods, we selected 214 field samples of the four most abundant *Drosophila* species that occurred at all elevations along both transects: *Drosophila rubida, D. pseudoananassae, D. pallidifrons,* and *D. sulfurigaster* (Table 1). Eight *D. rubida* pupae were parasitised by parasitoid wasps, enabling us to test if there are any changes in microbiome richness or unique microbial taxa associated with a developing parasitoid. We subsequently sampled 70 pupae and 70 adults from isofemale (IF) laboratory lines (2-4 lines per species) of these four elevationally ubiquitous species (20 pupae and 20 adults from *D. sulfurigaster, D. rubida,* and *D. pseudoananassae*, and 10 pupae and 10 adults from *D. pallidifrons*) to investigate if suspected natural patterns (site- and species-specific influence) were retained in lab-reared flies. Additionally, we took 10 samples of the food source provided to lab-reared *Drosophila* and 20 samples of the banana bait we used in our field sampling, to compare *Drosophila* microbiome samples to a dietary reference and determine how congruent the microbiome communities were between the food source and the insect hosts. Table 1 presents a detailed breakdown of all samples used in this study.

The *Drosophila* pupae field samples were collected from banana-baited bottle traps placed at low and high elevation sites along both altitudinal gradients. Each bottle trap had a piece

of cardboard to assist *Drosophila* larvae in pupation. Bottle traps were left exposed for

either 11-12, 14-15, or 24 days, to capture the natural variation in community colonisation

and ontogenetic development in different *Drosophila* species[50]. On the day of sampling,

these cards were removed and sealed in ziplock bags. The individuals we collected as pupae

thus only fed on banana bait. Pupae from each card were sampled by placing the card on a

white plastic dinner plate and adding distilled water, using a small paintbrush to remove all

pupae. Each pupa was placed into an individual well in 96-well PCR plates and preserved in

100% ethanol. Adults were aspirated from bottle traps hung at James Cook University,

Townsville (JCU), 2 days after provision of fresh banana bait and placed into individual vials

in 100% ethanol. JCU became a supplementary sampling site after Kirrama became

inaccessible due to heavy rainfall and landslides. Laboratory-reared pupae were collected

with forceps from standard fly food. Adults were collected with an aspirator and sexed, then

placed in individual vials with 100% ethanol. Laboratory isofemale lines were established

from the same populations sampled in the field (i.e., they were collected at the same sites

and shipped live to the lab in Czech Republic in 2017-18, after collection of field samples in

2016). Isofemale lines were kept in the lab on a standard *Drosophila* diet medium (corn

flour, sugar, agar, yeast, and methyl-4-hydroxybenzoate) for between 18-30 months by the

time of sampling.

Sample DNA was individually extracted using single column GeneAid Blood and Tissue kits

according to manufacturer instructions, with one extraction negative control accompanying

every 29 samples. For confirmation of *Drosophila* species identification we utilised a

multiplex PCR approach[50]. The PCRs were based on the ITS2 and COI regions using custom

specific forward and reverse primers. We combined primers into multiplex PCR reactions

based on difference in length of product. Gel electrophoresis of PCR products showed

product length, allowing us to identify species. When multiplex PCR was inconclusive, we

used Sanger sequencing with custom Diptera specific primers to identify the *Drosophila*

species. Full details of these processes and primer sequences can be found in the Supporting

Information of [50].

2.3 Library Preparation & Sequencing

After extraction and identification, all samples were moved to 96-well plates in a

randomised order in preparation for bacterial sequencing. DNA templates were stored at -

75ºC. These templates were used for amplification of ~400 bp of the V4/V5 hypervariable

region of the 16S rRNA gene according to Earth Microbiome Project standards (EMP;

http://www.earthmicrobiome.org/protocols-and-standards/16s/). Briefly, sample

multiplexing was based on the EMP-proposed double barcoding strategy using the EMP-

recommended modifications (12 bp Golay barcodes included on the forward primer 515F

(5Ⓐ-GTGYCAGCMGCCGCGGTAA), and additional 5 bp barcodes on the reverse primer 926R

(5Ⓐ-CCGYCAATTYMTTTRAGTTT)[52,53]. We also added a custom 18S rRNA gene blocking primer

(named 926X - 5Ⓐ GTGCCCTTCCGTCAATTCCT-C3 3') to counteract the low specificity of EMP

primers towards the 16S rRNA gene[22]. PCRs were carried out in triplicate and successful

amplification was confirmed with gel electrophoresis. Combined triplicates for each sample

were purified with AMPure XP (Beckman Coulter) magnetic beads and equimolarly pooled

to a single library (based on DNA concentration measured using a Synergy H1 (BioTek)

spectrophotometer). The library was purified using Pippin Prep (Sage Science) from all

fragments outside the 300-1100 bp range. To control for contamination, PCR biases and

confirm barcoding success, we included four negative controls from the extraction

procedure (ENC), eight negative controls from the PCR process (NC), and eight positive

controls (PC) of mock microbial communities. PCs were supplied commercially and

comprised 4 samples of gDNA templates with equal abundance of 10 bacterial species

(ATCC® MSA-1000™) and 4 samples with staggered abundance for the same bacteria (ATCC®

MSA-1001™). Altogether the library comprised PCR products from four 96-well plates, each

containing one ENC, two NCs, and two different PCs. The library was sequenced in a single

run of Illumina MiSeq platform using v3 chemistry with 2 x 300 bp output (Norwegian High

Throughput Sequencing Centre, Department of Medical Genetics, Oslo University Hospital).

2.4 Data Processing

The sequencing process returned 15,893,914 reads. These raw reads were quality checked

(FastQC[54]) and trimmed using USEARCH v9.2.64[55], to keep the quality score above Q20. We

trimmed the primers, demultiplexed and merged the reads which resulted in a final

amplicon length of 357 bp. We then clustered the reads at 100% identity for a

representative set of sequences and used the USEARCH global alignment option at both

99% and 97% identity[55] for *de novo* OTU assignment. We subsequently used the BLAST

algorithm[56] on the representative sequences, matching them against the SILVA 132

database[57] for taxonomic identification, producing a dataset of 1132 OTUs at 97% identity

and 1118 at 99% identity. We used the 97% identity OTU table as the primary dataset and

the 99% identity table as a supplemental dataset to confirm that the patterns we found

were not a product of identity threshold (Fig. S1). Raw sequence data is available on NCBI

Raw Sequence Archive (accession no.: PRJNA849960).

We recovered a mean of 16,898 reads per sample and a median of 14,751 reads (not

including negative controls). From our positive controls, we recovered microbiome profiles

that matched the expected community composition in each of the 'staggered' and 'even'

mock communities. The two low abundant species from the staggered templates (present at

0.04%) were successfully recovered from all four staggered mock samples. In the even

mocks, there was consistent overrepresentation of *Clostridium beijerinckii* and *Escherichia*

coli (1.4x - 4.7x expected), leading to reductions in other taxa. Overall, the positive controls

in this sequencing run matched our previous sequence outcomes[22,58].

Any chloroplast, mitochondrial, or eukaryotic OTUs were identified in the OTU table and

excluded. Potential bacterial contaminants were systematically evaluated using the R

package 'decontam' (V1.5.0[59]). The 'decontam' package uses the prevalence or frequency of

OTUs detected in negative controls to remove suspected contaminants. We used the

prevalence method, which compares the presence/absence of each sequence in true

positive samples compared to the prevalence in negative controls. We used a strict

threshold of 0.5, meaning any OTU with a higher proportion of reads in negative controls

than test samples was excluded as a contaminant. The prevalence of each OTU is shown in

Fig. S2. 43 OTUs were eliminated from the dataset via this process. Singletons were also

excluded, as was any OTU that made up less than one percent of the total reads for each

sample, which collectively removed 837 OTUs. These procedures resulted in a dataset of

110 OTUs and 360 samples.

2.5 Statistical Analyses

Sample analysis was carried out using the packages 'vegan'[60] and 'phyloseq'[61] in R[62]. To

measure alpha diversity we calculated the Shannon diversity index, using paired Wilcoxon

and paired ANOVA tests to compare values between sets of samples. We calculated Bray-

Curtis dissimilarity as a quantitative measure of beta diversity and used these values to

create non-metric multidimensional scaling ordinations (NMDS), to simultaneously evaluate

the roles of elevation, site, *Drosophila* species, life stage, environment of origin, parasitism,

trap identity, duration of trap exposure (and isofemale line and generation number for

laboratory samples). To support these ordinations statistically, we calculated PERMANOVA

tests on Bray-Curtis dissimilarity values. We applied the Benjamini-Hochberg (B-H)

correction on PERMANOVA p-values to control for multiple comparisons and false discovery

rate. We also evaluated statistical significance of differences in dispersion among groups

using multivariate homogeneity analysis (function "betadisper" in 'vegan'[60]) and ran each of

these tests with 999 permutations. In addition, we used DeSEQ differential abundance

analysis from the R package 'microbiomeSeq'[63] to compare the relative abundances of

bacterial taxa across different sample sets, e.g., differential abundance between *Drosophila*

species or between sites.

To systematically order our analyses, we first tested every variable (environment of origin,

elevation, gradient identity, species identity, developmental stage, trap identity, trap

duration of exposure, generation number, isofemale line) with all samples to determine the

relative importance of all studied factors (results shown below). We subsequently separated

the data by environment of origin, i.e., we tested lab *Drosophila* samples separately and

field *Drosophila* samples separately. Splitting the dataset in this way meant some variables

were not relevant to field samples, and vice versa. For example, when analysing field-only

samples we did not include isofemale line or generation number because these only

pertained to lab samples. Similarly, when focusing on lab-only samples, we did not include

trap identity, trap exposure time, or parasitoid status, because none were parasitised, and

traps were only a component of the field study. Furthermore, trap identity and site were

correlated variables because each trap was only used at one site. We tested the food/bait

samples separately because they were not focal samples and were used to support

conclusions about *Drosophila* microbiomes. The only samples positive for parasitoid

detection were from *Drosophila rubida* pupae, thus we only compared these samples to

other *D. rubida* pupae, and not as part of the core analyses.

3. Results

We first tested which of the studied factors influenced microbiome composition in all core

field and lab samples combined, using PERMANOVA analysis on Bray-Curtis values. The

dominant explanatory variable was environment of origin (i.e., whether a sample came from

the lab or field; NMDS mean stress ≈ 0.15, PERMANOVA, R^2 = 0.150, Benjamini-Hochberg

corrected p ≤ 0.001; Beta-dispersion F = 126.8, p ≤ 0.001; Table S1, Fig. 1). The R^2 values (see

Table S1) indicated that environment of origin was clearly an overarching explanatory

variable, thus we decided to further analyse the field and lab samples separately to

establish the important deterministic factors within each environment. The main trend in

our results was a significant reduction in microbiome richness in lab-reared flies of all

species based on a paired Wilcoxon rank sum test between Shannon index values for lab

and field samples (B-H corrected p ≤ 0.001; Fig. 2). This significant trend held when pupae or

adult *Drosophila* were analysed separately (Fig. S3, S4, S5).

3.1 Microbiome patterns from field samples

When focusing on core field samples, the main factors explaining variation in microbiome structure were trap identity (PERMANOVA R^2 = 0.13, B-H corrected p ≤ 0.001; Beta-dispersion p ≤ 0.001) and the interaction between site & trap (PERMANOVA R^2 = 0.13, B-H corrected p ≤ 0.001; Beta-dispersion p ≤ 0.001; Table S2), suggesting that local environmental differences in site location, as well as trap identity, have a significant effect on *Drosophila* microbiome composition (Fig. 3). Elevation alone explained a small, but still significant, proportion of the variation observed (R^2 = 0.03). The banana-baited bottle traps were left exposed in the field for different durations (between 11 and 24 days) to ensure we characterized the *Drosophila* community fully. PERMANOVA results suggest that the length of exposure had significant influence on microbiome composition but of lower importance compared to the other factors we identified based on the amount of variation explained (R^2 = 0.03). Based on the PERMANOVA analysis, field site was a significant variable but only explained 7% of the variation in diversity (reflected in the minimal differences in average Shannon index value in Fig. S6). We also found evidence of *Drosophila* species-specific differences in microbiome diversity, based on paired Wilcoxon tests between Shannon index values for different species (p ≤ 0.001; Fig. 4). DeSEQ analysis of differential abundance indicated some bacterial genera were significantly more abundant in some *Drosophila* species but not others, most prominently *Kozakia* and *Corynebacterium* (Table S3). The most abundant bacterial genera (Fig. 5) were evenly distributed throughout all four *Drosophila* species sampled here, including *Acinetobacter,* which was the most dominant genus overall.

3.2 Microbiome patterns from laboratory samples

The main factors explaining variation in microbiome structure from the lab-reared samples

were *Drosophila* species identity (PERMANOVA R^2 = 0.11, B-H corrected p ≤ 0.001, Beta-

dispersion p ≤ 0.001), isofemale line (PERMANOVA R^2 = 0.13, B-H corrected p ≤ 0.001, Beta-

dispersion p ≤ 0.001), and the number of generations an isofemale line had been in the lab

(PERMANOVA R^2 = 0.12, B-H corrected p ≤ 0.001, Beta-dispersion p = 0.007; Table S4),

suggesting that the duration a lineage of flies had been in the lab environment was

important for explaining microbiome composition. The amount of variation explained by

species-specific differences can largely be attributed to the presence of *Wolbachia* in *D.*

pseudoananassae (Fig. 4 & Fig. S7), because the alpha diversity differences between species

were not significant (Shannon index values, Fig. 6). All four *Drosophila* species contained

high proportions of *Acetobacter*, *Lactobacillus* and *Gluconobacter*, but only *D.*

pseudoananassae contained *Wolbachia*. The interaction between isofemale line and

generation was also highly significant and explained a high proportion of variation relative

to other factors, which was expected because IF lines were established at different times,

and therefore had been in the lab for varying numbers of generations at the time of

sampling. Life stage (i.e., the difference between pupae and adults) was not one of the most

important explanatory factors based on the amount of variation explained (PERMANOVA R^2

= 0.06). There was no detectable effect of elevation (PERMANOVA R^2 = 0.01), i.e., the

elevation of the field site where an isofemale line was collected did not have a significant

effect on the microbiome structure in the lab.

3.3 Comparisons between field and laboratory samples and microbiome of the food source

There was no significant difference in microbiome composition of laboratory food samples

from Czech Republic and the banana bait that we used in the field in Australia (Paired

Wilcoxon test on Shannon index values, p = 0.09; Fig. S8). Banana bait and lab fly food both

contained high relative proportions of *Acetobacter* and *Lactobacillus,* with some lab food

samples containing *Gluconobacter* (Fig. S9). Despite the physical differences in food sources,

the most abundant bacterial genera were the same, and the food source microbiome

diversity was low. In lab-reared flies, these three bacterial genera dominated the

microbiomes of pupae and adults (Fig. S7). In the field, however, *Acetobacter* and

Lactobacillus were not the most abundant genera of either pupae or adult gut microbiomes.

Acetobacter and *Lactobacillus* had the greatest difference in relative abundance between

field and lab samples (summary statistics in Table 2, Fig. S10). Whilst still present in field

samples, the relative abundance of *Acetobacter* and *Lactobacillus* was proportionally lower

in their taxa-rich microbiomes (Fig. 1).

3.4 Robustness of results

There was no detectable difference in microbiome diversity of parasitised pupae of *D.*

rubida, compared to unparasitized pupae (Paired Wilcoxon test on Shannon index values, p

= 0.25; Fig. S11), nor were there bacterial genera unique to parasitised samples. We also

found that the 99% identity OTU table produced qualitatively the same results as the 97%

identity OTU table. For example, in an NMDS on all samples the dominant explanatory

variable was still environment of origin (NMDS mean stress ≈ 0.17, PERMANOVA, R^2 = 0.17,

B-H corrected p ≤ 0.001; Beta-dispersion p ≤ 0.001; Fig. S1). Removing *Wolbachia* from the

dataset also did not qualitatively change the outcome of statistical tests, e.g., the main

factors explaining microbiome structure in lab-reared samples were *Drosophila* species

identity (PERMANOVA R^2 = 0.11, B-H corrected p ≤ 0.001, Beta-dispersion p ≤ 0.001),

isofemale line (PERMANOVA R^2 = 0.17, B-H corrected p ≤ 0.001, Beta-dispersion p ≤ 0.001),

and the number of generations an isofemale line had been in the lab (PERMANOVA R^2 =

0.12, Benjamini-Hochberg corrected p ≤ 0.001, Beta-dispersion p ≤ 0.001).

4. Discussion

Our results revealed small but significant variation in microbiome structure between

Drosophila populations from high and low elevation across both gradients. We expected to

find greater differences across elevation, because there is well-documented evidence of

both insects and bacteria developing differently according to differences in temperature of

>5°C[29–31,33,34,64]. This finding could be a result of the *Drosophila* species sampled here being

ubiquitous across elevation without forming sufficiently distinct populations at high and low

elevation sites, or because the ~5°C temperature shift between our sites was not strong

enough to drastically alter microbiome composition. Standardising diet may have

homogenised *Drosophila* microbiomes to some extent, but the overall lack of similarity in

microbiome composition between food microbiome and field-caught pupae suggests that

diet was not a dominant variable in structuring *Drosophila* microbiomes.

The most pronounced differences in microbiome composition were between individuals

raised in the laboratory and those raised in the field. Interestingly, the two food sources

(banana bait in the field and yeast-based *Drosophila* medium in the lab) had very similar

microbiome profiles, suggesting that dietary factors were not primarily responsible for the

observed differences between environments. The bacterial community from lab food

matches well with the microbiomes found within lab-reared pupae and adult flies. This was

expected, because it reflects a well-established pathway of insect microbiome colonisation -

Drosophila ingest food and acquire bacteria associated with that food source[36]. Yet in the

field, *Drosophila* microbiomes do not correspond well with the bacterial communities found

on banana bait samples. The observed pattern can be explained by significant differences in

microbiome colonisation from environmental bacterial species pools[65,66]. The flies sampled

from the lab come from a highly regulated environment, with a specific and consistent food

source provided into heat-sterilised glass vials, so the only 'available' bacteria for colonising

their microbiomes come from the diet, surrounding lab environment, and vertically

inherited endosymbionts (e.g., *Wolbachia* in *D. pseudoananassae*). In contrast, the bacterial

species pool in Australian tropical rainforest comprises much greater diversity and

abundance of different taxa, creating a greater variety of possible microbiome composition

within *Drosophila* hosts. This diversity of taxa creates more room for ecological drift,

dispersal, and selection to act on microbiome communities, in turn creating greater among-

individual and between-species variation in wild flies. Consistently higher diversity in wild

Drosophila microbiomes suggests that microbiomes are predominantly colonised from the

wider environment and dependent on local species pool diversity. For instance, bottle traps

were visited by other organisms - e.g., staphylinid beetles, neriid flies, Lepidoptera - which

could also have been a source of bacteria colonising the microbiome of the *Drosophila*

sampled in this study.

There was consistently low microbiome diversity found in both lab-reared pupae and lab-

reared adults, suggesting that low diversity within pupae is an accurate representation of

lab-reared microbiomes. This result was surprising because we anticipated some stage-

specific microbiome community patterns, given that *Drosophila* are holometabolous and thus undergo substantial gut remodelling during complete metamorphosis[67]. The consistency across life stages from lab-reared individuals provides further evidence for the depauperate nature of the lab microbial environment. In contrast to the lab, the field-caught adults of *D. rubida* lacked congruence with the field-caught pupae, and both stages lack similarity with the banana bait (Fig. S12, S13). The lack of geographic variation in microbiome composition suggests that different sites are unlikely to fully explain the discrepancy. With adult flies we cannot rule out that they might have fed on a substance other than our yeasted banana bait prior arriving at our bottle traps, making dietary variation a parsimonious explanatory factor in life-stage differences[36,68]. It is also possible that the high diversity we found in pupal samples reflects their metabolic activity (despite their lack of feeding), given that complete metamorphosis is an intense period of organismal change[69]. The substantial differences in microbiomes between lab and field specimens suggests that future studies should be cautious and specific with the types of microbiome-related questions investigating laboratory flies. Interpreting microbiome community composition from lab-kept specimens, particularly those from cultures maintained across multiple generations, is unlikely to yield data entirely representative of natural microbiomes[70,71], except perhaps in scenarios where microbiome communities from lab-reared specimens are a subset of the more diverse microbiomes found in the field (as our results show in Figure 5).

Previous studies on *Drosophila* have demonstrated high intra- and interspecific variation in microbiome community composition from both wild-caught and lab-reared flies[39,40,70,72,73], which our results corroborate. Controlling for diet in both scenarios allowed us to recognise

this species-specificity more accurately. We found a significant effect of species identity

(11% variation explained) but it did not explain as much variation as the results from Adair

et al.[40] who found species identity explaining 42% and 70% variation in two different sets of

Drosophila spp. This discrepancy could be a product of the species themselves (i.e., we used

a different set of *Drosophila* species) or the number of species studied (we studied four

species here; Adair et al.[40] eighteen), but the evidence from both studies suggests that

species-specificity is maintained in the lab, albeit not in a consistent manner. Statistically

significant differences in microbiome alpha diversity measures between different *Drosophila*

species in the field were not reflected in the lab. From the field, the only significant

difference in microbiome richness was between *D. pseudoananassae* and *D. rubida*, and this

was non-significant in the lab. Moreover, there was a highly significant difference in richness

between *D. rubida* and *D. sulfurigaster* in the lab, but not in the field. Some of our

significant PERMANOVA results were accompanied by significant Beta-dispersion, which

suggests that there was some heterogeneous dispersion in the variables we measured. This

was expected based on the high variation typically exhibited by microbiome community

data and typically found when sampling wild individuals. Even with greater and more even

sample size, we would still expect to find a large, significant difference in community

richness and composition between field and lab *Drosophila* microbiomes.

In other insect species, host transmission of extracellular symbionts (like those in the gut)

have been hypothesised to result in long-term associations between insect and microbe[74–

76]. The long-term laboratory survival of our four *Drosophila* species (minimum 24

generations) with radically different microbiome composition (compared to their

counterparts from the field) suggests that their symbiotic relationship with bacteria is not

highly specialised[77], corroborating previously published findings. For example, Wong et al.[37]

found no consistent evidence for a core microbiome in multiple *Drosophila* species and

Storelli et al.[78] described axenic *Drosophila* returning to normal growth and development in

the presence of a single bacteria species. Furthermore, Coon et al.[79,80] showed that any

bacterium was sufficient to facilitate mosquito larval development, whereas axenic

mosquito larvae do not develop past the first instar. Thus, our results provide further

evidence of satisfactory microbiome function being provided by a limited number of

bacterial species, which poses questions about the advantages of diverse microbiomes

within wild insects and the links between microbiome community diversity and host fitness.

Overall, we found significant differences in microbiome diversity of field-caught and lab-

reared *Drosophila*, which were consistent across species and life stage. We hypothesise that

these differences in diversity are the products of environments with markedly different

bacterial species pools. To elucidate functional conclusions from insect-microbiome

analyses, more in-depth molecular analysis (e.g., metagenomics, transcriptomics) is

required. We suggest that comparative studies between lab and field specimens help reveal

the true variability in microbiome communities that can exist within a single species.

Acknowledgements

We thank Hana Konvičková and Jan Zima Jr. for their assistance in the molecular lab. We

acknowledge funding support from the Czech Science Foundation grant no. 17-27184Y to

JH. OTL was further supported by the UK National Environment Research Council

(NE/N010221/1). JH was additionally supported by the Czech Ministry of Education, Youth

and Sports grant ERC CZ LL2001. Computational resources were supplied by the project "e-

Infrastruktura CZ" (e-INFRA LM2018140) provided within the program Projects of Large

Research, Development, and Innovations Infrastructures. Fieldwork was conducted under

permit WITK16977516 from Queensland's Department of Environment and Heritage

Protection.

References

1 Fierer N, McCain CM, Meir P, Zimmermann M, Rapp JM, Silman MR, *et al.* Microbes do
not follow the elevational diversity patterns of plants and animals. *Ecology* 2011;**92**:797–
804. https://doi.org/10.1890/10-1170.1.
2 Wang J, Soininen J, Zhang Y, Wang B, Yang X, Shen J. Contrasting patterns in elevational
diversity between microorganisms and macroorganisms. *J Biogeogr* 2011;**38**:595–603.
https://doi.org/10.1111/j.1365-2699.2010.02423.x.
3 Pärtel M, Bennett JA, Zobel M. Macroecology of biodiversity: disentangling local and
regional effects. *New Phytol* 2016;**211**:404–10. https://doi.org/10.1111/nph.13943.
4 Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, *et al.* A communal
catalogue reveals Earth's multiscale microbial diversity. *Nature* 2017;**551**:457–63.
https://doi.org/10.1038/nature24621.
5 Alruiz JM, Peralta-Maraver I, Bozinovic F, Santos M, Rezende EL. Thermal tolerance in
Drosophila: Repercussions for distribution, community coexistence and responses to
climate change. *J Anim Ecol* 2022;**91**:667. https://doi.org/10.1111/1365-2656.13653.
6 Fierer N, Jackson RB. The diversity and biogeography of soil bacterial communities. *Proc*
Natl Acad Sci 2006;**103**:626–31. https://doi.org/10.1073/pnas.0507535103.
7 Lauber CL, Hamady M, Knight R, Fierer N. Pyrosequencing-based assessment of soil pH as
a predictor of soil bacterial community structure at the continental scale. *Appl Environ*
Microbiol 2009;**75**:5111–20. https://doi.org/10.1128/AEM.00335-09.
8 Meyer KM, Memiaghe H, Korte L, Kenfack D, Alonso A, Bohannan BJM. Why do microbes
exhibit weak biogeographic patterns? *ISME J* 2018;**12**:1404–13.
https://doi.org/10.1038/s41396-018-0103-3.

9 Shen C, Ni Y, Liang W, Wang J, Chu H. Distinct soil bacterial communities along a small-
scale elevational gradient in alpine tundra. *Front Microbiol* 2015;**6**:.
https://doi.org/10.3389/fmicb.2015.00582.
10 Siles JA, Margesin R. Abundance and diversity of bacterial, archaeal, and fungal
communities along an altitudinal gradient in alpine forest soils: what are the driving
factors? *Microb Ecol* 2016;**72**:207–20. https://doi.org/10.1007/s00248-016-0748-2.
11 McLean AHC, Parker BJ, Hrček J, Henry LM, Godfray HCJ. Insect symbionts in food webs.
Phil Trans R Soc B 2016;**371**:20150325. https://doi.org/10.1098/rstb.2015.0325.
12 Corbin C, Heyworth ER, Ferrari J, Hurst GDD. Heritable symbionts in a world of varying
temperature. *Heredity* 2017;**118**:10–20. https://doi.org/10.1038/hdy.2016.71.
13 Brown JJ, Mihaljevic JR, Des Marteaux L, Hrček J. Metacommunity theory for transmission
of heritable symbionts within insect communities. *Ecol Evol* 2020;**10**:1703–21.
https://doi.org/10.1002/ece3.5754.
14 Colman DR, Toolson EC, Takacs-Vesbach CD. Do diet and taxonomy influence insect gut
bacterial communities? *Mol Ecol* 2012;**21**:5124–37. https://doi.org/10.1111/j.1365-
294X.2012.05752.x.
15 Yun J-H, Roh SW, Whon TW, Jung M-J, Kim M-S, Park D-S, *et al.* Insect gut bacterial
diversity determined by environmental habitat, diet, developmental stage, and
phylogeny of host. *Appl Env Microbiol* 2014;**80**:5254–64.
https://doi.org/10.1128/AEM.01226-14.
16 Xie J, Winter C, Winter L, Mateos M. Rapid spread of the defensive endosymbiont
Spiroplasma in *Drosophila hydei* under high parasitoid wasp pressure. *FEMS Microbiol
Ecol* 2015;**91**:1–11. https://doi.org/10.1093/femsec/fiu017.
17 Martinson VG, Carpinteyro-Ponce J, Moran NA, Markow TA. A distinctive and host-
restricted gut microbiota in populations of a cactophilic *Drosophila* species. *Appl Environ
Microbiol* 2017;**83**:. https://doi.org/10.1128/AEM.01551-17.
18 Nováková E, Woodhams DC, Rodríguez-Ruano SM, Brucker RM, Leff JW, Maharaj A, *et al.*
Mosquito microbiome dynamics, a background for prevalence and seasonality of West
Nile virus. *Front Microbiol* 2017;**8**:526. https://doi.org/10.3389/fmicb.2017.00526.
19 Bing X, Gerlach J, Loeb G, Buchon N. Nutrient-dependent impact of microbes on
Drosophila suzukii development. *MBio* 2018;**9**:. https://doi.org/10.1128/mBio.02199-17.
20 Moghadam NN, Thorshauge PM, Kristensen TN, de Jonge N, Bahrndorff S, Kjeldal H, *et al.*
Strong responses of *Drosophila melanogaster* microbiota to developmental temperature.
Fly (Austin) 2018;**12**:1–12. https://doi.org/10.1080/19336934.2017.1394558.
21 Park R, Dzialo MC, Spaepen S, Nsabimana D, Gielens K, Devriese H, *et al.* Microbial
communities of the house fly *Musca domestica* vary with geographical location and
habitat. *Microbiome* 2019;**7**:147. https://doi.org/10.1186/s40168-019-0748-9.
22 Brown JJ, Rodríguez-Ruano SM, Poosakkannu A, Batani G, Schmidt JO, Roachell W, *et al.*
Ontogeny, species identity, and environment dominate microbiome dynamics in wild
populations of kissing bugs (Triatominae). *Microbiome* 2020;**8**:146.
https://doi.org/10.1186/s40168-020-00921-x.
23 Douglas AE. How multi-partner endosymbioses function. *Nat Rev Microbiol* 2016;**14**:731–
43. https://doi.org/10.1038/nrmicro.2016.151.
24 Woodhams DC, Bletz MC, Becker CG, Bender HA, Buitrago-Rosas D, Diebboll H, *et al.*
Host-associated microbiomes are predicted by immune system complexity and climate.
Genome Biol 2020;**21**:23. https://doi.org/10.1186/s13059-019-1908-8.

25 Wadgymar SM, Mactavish RM, Anderson JT. Transgenerational and within-generation
plasticity in response to climate change: insights from a manipulative field experiment
across an elevational gradient. *Am Nat* 2018;**192**:698–714.
https://doi.org/10.1086/700097.
26 Nottingham AT, Whitaker J, Ostle NJ, Bardgett RD, McNamara NP, Fierer N, *et al.*
Microbial responses to warming enhance soil carbon loss following translocation across a
tropical forest elevation gradient. *Ecol Lett* 2019;**22**:1889–99.
https://doi.org/10.1111/ele.13379.
27 Economos AC, Lints FA. Growth rate and life span in *Drosophila*. III. Effect of body size and
development temperature on the biphasic relationship between growth rate and life
span. *Mech Ageing Dev* 1984;**27**:153–60. https://doi.org/10.1016/0047-6374(84)90040-X.
28 James AC, Azevedo RBR, Partridge L. Genetic and environmental responses to
temperature of *Drosophila melanogaster* from a latitudinal cline. *Genetics* 1997;**146**:881–
90.
29 Kinjo H, Kunimi Y, Nakai M. Effects of temperature on the reproduction and development
of *Drosophila suzukii* (Diptera: Drosophilidae). *Appl Entomol Zool* 2014;**49**:297–304.
https://doi.org/10.1007/s13355-014-0249-z.
30 Tochen S, Dalton DT, Wiman N, Hamm C, Shearer PW, Walton VM. Temperature-related
development and population parameters for *Drosophila suzukii* (Diptera: Drosophilidae)
on cherry and blueberry. *Environ Entomol* 2014;**43**:501–10.
https://doi.org/10.1603/EN13200.
31 Brankatschk M, Gutmann T, Knittelfelder O, Palladini A, Prince E, Grzybek M, *et al.* A
temperature-dependent switch in feeding preference improves *Drosophila* development
and survival in the cold. *Dev Cell* 2018;**46**:781-793.e4.
https://doi.org/10.1016/j.devcel.2018.05.028.
32 Ratkowsky DA, Olley J, McMeekin TA, Ball A. Relationship between temperature and
growth rate of bacterial cultures. *J Bacteriol* 1982;**149**:1–5.
33 Pettersson M, Bååth E. Temperature-dependent changes in the soil bacterial community
in limed and unlimed soil. *FEMS Microbiol Ecol* 2003;**45**:13–21.
https://doi.org/10.1016/S0168-6496(03)00106-5.
34 Tsuji S, Ushio M, Sakurai S, Minamoto T, Yamanaka H. Water temperature-dependent
degradation of environmental DNA and its relation to bacterial abundance. *PLoS ONE*
2017;**12**:. https://doi.org/10.1371/journal.pone.0176608.
35 Chandler JA, Lang JM, Bhatnagar S, Eisen JA, Kopp A. Bacterial communities of diverse
Drosophila species: ecological context of a host–microbe model system. *PLOS Genet*
2011;**7**:e1002272. https://doi.org/10.1371/journal.pgen.1002272.
36 Blum JE, Fischer CN, Miles J, Handelsman J. Frequent replenishment sustains the
beneficial microbiome of *Drosophila melanogaster*. *MBio* 2013;**4**:e00860-13.
https://doi.org/10.1128/mBio.00860-13.
37 Wong AC-N, Chaston JM, Douglas AE. The inconstant gut microbiota of *Drosophila* species
revealed by 16S rRNA gene analysis. *ISME J* 2013;**7**:1922–32.
https://doi.org/10.1038/ismej.2013.86.
38 Chaplinska M, Gerritsma S, Dini-Andreote F, Salles JF, Wertheim B. Bacterial communities
differ among *Drosophila melanogaster* populations and affect host resistance against
parasitoids. *PLOS ONE* 2016;**11**:e0167726.
https://doi.org/10.1371/journal.pone.0167726.

39 Adair KL, Wilson M, Bost A, Douglas AE. Microbial community assembly in wild
populations of the fruit fly *Drosophila melanogaster*. *ISME J* 2018:1.
https://doi.org/10.1038/s41396-017-0020-x.
40 Adair KL, Bost A, Bueno E, Kaunisto S, Kortet R, Peters-Schulze G, *et al.* Host determinants
of among-species variation in microbiome composition in drosophilid flies. *ISME J*
2020;**14**:217–29. https://doi.org/10.1038/s41396-019-0532-7.
41 Henry Y, Colinet H. Microbiota disruption leads to reduced cold tolerance in *Drosophila*
flies. *Sci Nat* 2018;**105**:59. https://doi.org/10.1007/s00114-018-1584-7.
42 Elgart M, Stern S, Salton O, Gnainsky Y, Heifetz Y, Soen Y. Impact of gut microbiota on the
fly's germ line. *Nat Commun* 2016;**7**:. https://doi.org/10.1038/ncomms11280.
43 Lizé A, McKay R, Lewis Z. Kin recognition in *Drosophila*: the importance of ecology and gut
microbiota. *ISME J* 2014;**8**:469–77. https://doi.org/10.1038/ismej.2013.157.
44 Sansone CL, Cohen J, Yasunaga A, Xu J, Osborn G, Subramanian H, *et al.* Microbiota-
dependent priming of antiviral intestinal immunity in *Drosophila*. *Cell Host Microbe*
2015;**18**:571–81. https://doi.org/10.1016/j.chom.2015.10.010.
45 Chrostek E, Marialva MSP, Esteves SS, Weinert LA, Martinez J, Jiggins FM, *et al.*
Wolbachia variants induce differential protection to viruses in *Drosophila melanogaster*:
A phenotypic and phylogenomic analysis. *PLOS Genet* 2013;**9**:e1003896.
https://doi.org/10.1371/journal.pgen.1003896.
46 Hamilton PT, Perlman SJ. Host defense via symbiosis in *Drosophila*. *PLOS Pathog*
2013;**9**:e1003808. https://doi.org/10.1371/journal.ppat.1003808.
47 Haselkorn TS, Cockburn SN, Hamilton PT, Perlman SJ, Jaenike J. Infectious adaptation:
potential host range of a defensive endosymbiont in *Drosophila*. *Evolution* 2013;**67**:934–
45. https://doi.org/10.1111/evo.12020.
48 Xie J, Butler S, Sanchez G, Mateos M. Male killing *Spiroplasma* protects *Drosophila*
melanogaster against two parasitoid wasps. *Heredity* 2014;**112**:399–408.
https://doi.org/10.1038/hdy.2013.118.
49 Yadav S, Frazer J, Banga A, Pruitt K, Harsh S, Jaenike J, *et al.* Endosymbiont-based
immunity in *Drosophila melanogaster* against parasitic nematode infection. *PLOS ONE*
2018;**13**:e0192183. https://doi.org/10.1371/journal.pone.0192183.
50 Jeffs CT, Terry JCD, Higgie M, Jandová A, Konvičková H, Brown JJ, *et al.* Molecular analyses
reveal consistent food web structure with elevation in rainforest *Drosophila* – parasitoid
communities. *Ecography* 2021;**44**:403–13. https://doi.org/10.1111/ecog.05390.
51 Staubach F, Baines JF, Künzel S, Bik EM, Petrov DA. Host species and environmental
effects on bacterial communities associated with *Drosophila* in the laboratory and in the
natural environment. *PLoS ONE* 2013;**8**:. https://doi.org/10.1371/journal.pone.0070749.
52 Quince C, Lanzen A, Davenport RJ, Turnbaugh PJ. Removing noise from pyrosequenced
amplicons. *BMC Bioinformatics* 2011;**12**:38. https://doi.org/10.1186/1471-2105-12-38.
53 Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA
primers for marine microbiomes with mock communities, time series and global field
samples. *Environ Microbiol* 2016;**18**:1403–14. https://doi.org/10.1111/1462-2920.13023.
54 Andrews S. FASTQC. A quality control tool for high throughput sequence data 2010.
55 Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat*
Methods 2013;**10**:996–8. https://doi.org/10.1038/nmeth.2604.
56 Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, *et al.* BLAST+:
architecture and applications. *BMC Bioinformatics* 2009;**10**:421.
https://doi.org/10.1186/1471-2105-10-421.

57 Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, *et al.* The SILVA ribosomal RNA
gene database project: improved data processing and web-based tools. *Nucleic Acids Res*
2013;**41**:D590–6. https://doi.org/10.1093/nar/gks1219.

58 Rodríguez-Ruano SM, Škochová V, Rego ROM, Schmidt JO, Roachell W, Hypša V, *et al.*
Microbiomes of North American Triatominae: the grounds for Chagas disease
epidemiology. *Front Microbiol* 2018;**9**:1167. https://doi.org/10.3389/fmicb.2018.01167.

59 Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. Simple statistical
identification and removal of contaminant sequences in marker-gene and metagenomics
data. *Microbiome* 2018;**6**:226. https://doi.org/10.1186/s40168-018-0605-2.

60 Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, *et al.* vegan:
Community Ecology Package 2019.

61 McMurdie PJ, Holmes S. phyloseq: An R package for reproducible interactive analysis and
graphics of microbiome census data. *PLOS ONE* 2013;**8**:e61217.
https://doi.org/10.1371/journal.pone.0061217.

62 R Core Team. R: A language and environment for statistical computing. 2021.

63 Ssekagiri A. microbiomeSeq: Microbial community analysis in an environmental context.
2021.

64 Cooper RO, Vavra JM, Cressler CE. Targeted manipulation of abundant and rare taxa in
the *Daphnia magna* Microbiota with antibiotics impacts host fitness differentially.
MSystems 2021;**6**:. https://doi.org/10.1128/mSystems.00916-20.

65 Cornell HV, Harrison SP. What are species pools and when are they important? *Annu Rev*
Ecol Evol Syst 2014;**45**:45–67. https://doi.org/10.1146/annurev-ecolsys-120213-091759.

66 Kohl KD. Ecological and evolutionary mechanisms underlying patterns of phylosymbiosis
in host-associated microbial communities. *Philos Trans R Soc B Biol Sci*
2020;**375**:20190251. https://doi.org/10.1098/rstb.2019.0251.

67 Hammer TJ, Moran NA. Links between metamorphosis and symbiosis in holometabolous
insects. *Philos Trans R Soc B Biol Sci* 2019;**374**:20190068.
https://doi.org/10.1098/rstb.2019.0068.

68 Obadia B, Keebaugh ES, Yamada R, Ludington WB, Ja WW. Diet influences host–
microbiota associations in *Drosophila*. *Proc Natl Acad Sci* 2018;**115**:E4547–8.
https://doi.org/10.1073/pnas.1804948115.

69 Merkey AB, Wong CK, Hoshizaki DK, Gibbs AG. Energetics of metamorphosis in *Drosophila*
melanogaster. *J Insect Physiol* 2011;**57**:1437–45.
https://doi.org/10.1016/j.jinsphys.2011.07.013.

70 Bost A, Martinson VG, Franzenburg S, Adair KL, Albasi A, Wells MT, *et al.* Functional
variation in the gut microbiome of wild *Drosophila* populations. *Mol Ecol* 2018;**27**:2834–
45. https://doi.org/10.1111/mec.14728.

71 Dada N, Benedict AC, López F, Lol JC, Sheth M, Dzuris N, *et al.* Comprehensive
characterization of internal and cuticle surface microbiota of laboratory-reared F1
Anopheles albimanus originating from different sites. *Malar J* 2021;**20**:414.
https://doi.org/10.1186/s12936-021-03934-5.

72 Bost A, Franzenburg S, Adair KL, Martinson VG, Loeb G, Douglas AE. How gut
transcriptional function of *Drosophila melanogaster* varies with the presence and
composition of the gut microbiota. *Mol Ecol* 2018;**27**:1848–59.
https://doi.org/10.1111/mec.14413.

73 Solomon GM, Dodangoda H, McCarthy-Walker T, Ntim-Gyakari R, Newell PD. The
microbiota of *Drosophila suzukii* influences the larval development of *Drosophila*
melanogaster. *PeerJ* 2019;**7**:e8097. https://doi.org/10.7717/peerj.8097.
74 Sanders JG, Powell S, Kronauer DJC, Vasconcelos HL, Frederickson ME, Pierce NE. Stability
and phylogenetic correlation in gut microbiota: lessons from ants and apes. *Mol Ecol*
2014;**23**:1268–83. https://doi.org/10.1111/mec.12611.
75 Kwong WK, Medina LA, Koch H, Sing K-W, Soh EJY, Ascher JS, *et al.* Dynamic microbiome
evolution in social bees. *Sci Adv* 2017;**3**:e1600513.
https://doi.org/10.1126/sciadv.1600513.
76 Sinotte VM, Renelies-Hamilton J, Taylor BA, Ellegaard KM, Sapountzis P, Vasseur-Cognet
M, *et al.* Synergies between division of labor and gut microbiomes of social insects. *Front*
Ecol Evol 2020;**7**:. https://doi.org/10.3389/fevo.2019.00503.
77 Leftwich PT, Clarke NVE, Hutchings MI, Chapman T. Gut microbiomes and reproductive
isolation in *Drosophila*. *Proc Natl Acad Sci* 2017;**114**:12767–72.
78 Storelli G, Defaye A, Erkosar B, Hols P, Royet J, Leulier F. *Lactobacillus plantarum*
Promotes *Drosophila* Systemic Growth by Modulating Hormonal Signals through TOR-
Dependent Nutrient Sensing. *Cell Metab* 2011;**14**:403–14.
https://doi.org/10.1016/j.cmet.2011.07.012.
79 Coon KL, Vogel KJ, Brown MR, Strand MR. Mosquitoes rely on their gut microbiota for
development. *Mol Ecol* 2014;**23**:2727–39. https://doi.org/10.1111/mec.12771.
80 Coon KL, Brown MR, Strand MR. Mosquitoes host communities of bacteria that are
essential for development but vary greatly between local habitats. *Mol Ecol*
2016;**25**:5806–26. https://doi.org/10.1111/mec.13877.
726
727

Table 1: Breakdown of the sample set used in this study. PL = Paluma Low, PH = Paluma High, KL = Kirrama Low, KH = Kirrama High, JCU = James Cook University campus in Townsville.

Species	Stage	Environment of origin	Sites	Number of samples	Core or Additional
D. rubida	pupae	field	PL, PH, KL, KH	71	Core
D. rubida (parasitised)	pupae	field	PL, PH, KL, KH	8	Core
D. rubida	adult	field	JCU	14	Core
D. pseudoananassae	pupae	field	PL, PH, KL, KH	48	Core
D. pallidifrons	pupae	field	PL, PH, KL, KH	39	Core
D. sulfurigaster	pupae	field	PL, PH, KL, KH	10	Core
D. rubida	pupae	lab	PL, PH, KL, KH	20	Core
D. rubida	adult	lab	PL, PH, KL, KH	20	Core
D. pseudoananassae	pupae	lab	PL, KL, KH	20	Core

D. pseudoananassae	adult	lab	PL, KL, KH	20	Core
D. pallidifrons	pupae	lab	PH, KH	10	Core
D. pallidifrons	adult	lab	PH, KH	10	Core
D. sulfurigaster	pupae	lab	PL, PH, KL, KH	20	Core
D. sulfurigaster	adult	lab	PL, PH, KL, KH	20	Core
banana bait	na	field	PL, PH, JCU	20	Additional
lab fly food	na	lab	na	10	Additional

Figure 1: NMDS analysis of microbiome communities from all samples in this study, from

the lab (blue) and the field (red). Samples of *Drosophila rubida, D. pseudoananassae, D.*

*pallidifron*s*, D. sulfurigaster*, and food bait are indicated by different shapes.

Figure 2: Comparison of Shannon index values for all samples in this study, split by

environment of origin. Lab samples are shown in blue and field samples in red. The triple

asterisk (***) shows high significance (p > 0.001) of a paired Wilcoxon test.

Figure 3: PCoA analysis of microbiome communities from the core field samples. Samples

of *Drosophila rubida, D. pseudoananassae, D. pallidifron*s, and *D. sulfurigaster* are indicated

by different colours, and the different sites of collection are indicated by shape. K = Kirrama,

P = Paluma, JCU = James Cook University, L = Low [elevation], H = High [elevation]. Samples

of *D. rubida* collected at JCU were all adults.

Figure 4: Comparison of Shannon index values from each species of *Drosophila* from (A)

field and (B) lab environments. Each colour represents a different species. Statistical

comparisons come from paired Wilcoxon tests. Three asterisks (***) denotes a highly

significant result (p ≤ 0.001). Two asterisks (**) indicates a result of moderate significance

(between p < 0.1 and p ≤ 0.001). One asterisk (*) denotes a marginally significant result (p <

0.1). NS. = not significant.

Figure 5: The most abundant bacterial genera from all samples in this study. Different

colours mark different bacterial genera. <2.5% is a conglomerate category of low abundance

taxa that made up less than 2.5% of the median number of reads. Each individual column

represents an individual sample. Relative abundance is on the y axis.

Figure 6: NMDS analysis of microbiome communities from core lab samples. Samples of

*Drosophila rubida, D. pseudoananassae, D. pallidifron*s, and *D. sulfurigaster* are indicated by

different colours and shapes.

Table 2: Summary output of DeSEQ differential abundance analysis showing the top 5 bacterial genera by origin of sample. Positive test statistic values indicate a genus significantly more abundant in lab samples and negative test statistics indicate a genus significantly more abundant in field samples.					
Genus	baseMean	log2foldchange	lfcSE	Test statistic	Adjusted p value
Acetobacter	436.99	3.59	0.38	9.46	≤ 0.001
Lactobacillus	183.75	3.16	0.38	8.34	≤ 0.001
Acinetobacter	195.62	-7.91	0.33	-23.70	≤ 0.001
Klebsiella	105.22	-7.01	0.37	-18.99	≤ 0.001
Chishuiella	59.69	-6.19	0.41	-14.97	≤ 0.001

Dominant bacterial genera