Recent SIV-Macaque Trials:

A Reassessment of Immune Priming and Varying

Infectability in Repeated Low-dose Challenge Studies

Short title: Repeated Low-dose SIV Challenges in Macaques

John L. Spouge[1]*

[1]National Center for Biotechnology Information, National Library of Medicine, National

Institutes of Health, Bethesda, Maryland 20894, USA

*Corresponding author

E-mail: spouge@nih.gov

Abstract

Nowadays, most preclinical HIV treatment trials use a protocol of administering repeated low-

dose challenges (RLCs) of simian immunodeficiency virus (SIV) to macaques. Statistical

analyses of treatment efficacy under the RLC protocol need to consider two confounding

hypotheses, both pertinent biologically to HIV: (1) the non-infecting challenges may immunize

animals against SIV; and (2) the animals may vary in intrinsic infectability ("frailty"). To

explore the two hypotheses, a previous study (Regoes 2012) assembled a database from 7 articles

with SIV-macaque treatment trials. With two explicable exceptions, Regoes concluded that the

control data did not support either confounding hypothesis. Recent SIV-macaque trials present

opportunities to evaluate the conclusions' robustness. Accordingly, the present article assembles

from 24 articles an updated database containing net survival curves from both control and

treatment arms in SIV-macaque treatment trials. Broad patterns of statistical significance (at

$p<0.05$, uncorrected for multiple testing) made it difficult to dismiss the confounding hypotheses

completely in the controls. Although statistical analysis has focused on defense against variable

frailty, only one set of controls showed significant variable frailty, whereas many sets showed

significant immunization. As trials progressed, changes in the probability of infection per

challenge were significant in 8/28 trials (1/3 trials using oral challenges; 2/4 trials using vaginal

challenges; and 5/21 trials using rectal challenges). The results suggest the possibility that

vaginal challenges may immunize animals faster than rectal challenges, and they also bear on

previous conclusions that repeated exposure to HIV without treatment may have no effect on

infectability or may even reduce it.

Author Summary

Many preclinical trials of HIV treatments rely on repeatedly administering low-dose SIV challenges to macaques until infection occurs. The repeated low-dose protocol reuses macaques and is more sensitive to subtle therapeutic efficacies than a protocol administering a single large dose to each macaque. The animal reuse raises some pertinent biological questions, notably: (1) do macaques have intrinsically variable infectabilities? and (2) do the repeated SIV challenges immunize macaques against infection? A 2012 study collected a database of eight macaque trials, concluding that variable infectability and immunization were at most sporadic and readily explicable. I expanded the 2012 database to twenty-eight trials, discovering that the conclusions were not robust. Although only 1/28 SIV-macaque trials showed variable infectability, 7/28 showed immunization, with few ready explanations. Statistical analysis of SIV-macaque trials has focused on the confounding effects of variable infectability to the neglect of immunization, so the expanded database provides a rich empirical resource. The trials have general medical importance because they provide a model for analyzing animal trials of infectious disease therapies and other sparse trials, e.g., for breast cancer. My findings also indirectly suggest that repeated human exposure to HIV inconsistently immunizes and can foster either immune priming or tolerance.

55 Introduction

Preclinical experiments play a fundamental role in determining treatments for an infectious

disease. In a modern re-interpretation, some of Koch's postulates specify reproducing the disease

in an animal model and then reisolating the agent from an animal with signs resembling the

human disease [1]. Historically, experimenters therefore tested vaccines or preventive treatment

strategies against human-immunodeficiency virus (HIV) preclinically in non-human primates.

They found that HIV infects in the chimpanzee animal model, although it does not consistently

produce an AIDS-like disease.

Typically, preclinical HIV trials have control and treatment arms, and they infer therapeutic

protection if treated animals are less easily infected than controls. In the 1980s, preclinical HIV

trials were extremely sparing because of the cost of chimpanzees, e.g., one trial used 6

chimpanzees in a control arm and 2 chimpanzees in a treatment arm [2]. Such trials ensured

infection of controls with a large viral challenge [3, 4]. The trials then applied a statistical test,

e.g., the Fisher's exact test [5], to the resulting 2x2 tables of counts of animals in control/treated

vs infected/uninfected groups. Unfortunately, with so few counts, the single high-dose protocol

provided scant statistical power for detecting subtle protection against infection. Accordingly,

experimenters relied heavily on surrogates of protection to detect a treatment effect, e.g., an

increase in virus-specific immune responses from a vaccine treatment, or a decrease in the

steady-state viral load (a.k.a. the viral set-point). The surrogates are chosen to correlate with

protection, although they are not necessarily causal.

Since 2007, chimpanzees are an endangered species no longer bred for research [6], requiring

other animal models for HIV infection and AIDS. Mice can receive transplanted components of

a human immune system [7], and when applicable, the humanized mouse model is cost-effective.

The mouse model has, however, many serious limitations in cell number and volume, life span,

integrated operation of an immune system, etc. [8]. By contrast, over 40 different simian

immunodeficiency viruses (SIVs) infect primate species without such limitations [9], and some

do produce AIDS-like illnesses. Presently, the rhesus macaque and its SIV provide the most

common preclinical primate model of AIDS [10-33].

Although rhesus macaques provide less expensive preclinical data than chimpanzees, their

cost is still severe. Excluding the historical controls used in recent trials [26, 29, 32], macaque

trials have limited themselves on average to about 13 contemporary controls. Under such

limitations, if the trials used the single high-dose protocol, they would remain seriously

underpowered for detecting a subtle treatment effect [34].

Accordingly, preclinical HIV treatment trials have adopted a protocol using repeated low-

dose challenges (RLCs) [10-33]. As Table 1 of [18] helpfully illustrates, typically each animal is

challenged repeatedly at regular intervals with the same low viral dose, until either it is infected

or the experiment finishes. Experimenters usually challenge weekly, because the anesthetized

animal can then receive the current challenge after tests for infection from the previous week are

initiated. To limit the number of challenges required to infect controls, the challenge dose often

approximates the animal ID_{50} [4]. Some experiments challenge beyond infection (e.g., [10]), but

post-infection challenges are irrelevant to present purposes, and we ignore them.

The RLC protocol has several advantages [35], both biological and statistical [36]. First,

RLCs mimic human exposure to HIV, making the animal studies conceptually closer to

subsequent human studies. Second, they can in principle detect immunization of controls

statistically during repeated challenge, as follows. RLCs yield net survival curves for each

control or treatment arm (e.g., Figure 3 of [18]), where the curves count the uninfected animals

before and after each challenge. Non-infecting challenges might gradually immunize animals,

but statistical analysis of the fraction of controls infected after each challenge can detect, e.g.,

systematic decreases in the fraction over time. In effect, the statistical analysis defines

"infectability" as a probability of infection, with "immunization" as the corresponding change in

infectability. To paraphrase [35], the statistical definition of immunization presumes neither

knowledge nor measurement of the immune surrogates of protection, but instead specifies what

matters epidemiologically. In contrast, biological studies examining laboratory surrogates of

protection (e.g., [2]) implicitly rely on a strong but unproved assumption, that the laboratory

surrogates are causally linked to immune protection [37]. Though limited in other ways, a

statistical definition of immunization deftly avoids implicit assumptions about causality.

Third, RLCs can also expose intrinsic differences in infectability between subpopulations of

animals, their "frailties" in the terminology of survival analysis [35]. Immune surrogates have

demonstrated some specific subpopulation differences, e.g., the peak plasma RNA in

SIVsmE660 infection is about 10-fold reduced in rhesus macaques with the *Mamu-A*01* class I

allele [19]. On one hand, therefore, a study examining immune surrogates can draw specific

biological conclusions. On the other hand, a statistical analysis of infectability has the

complementary strength of generality, because it may detect immunization or variable frailty

regardless of specific biological mechanisms like alleles or restriction factors [38].

Unfortunately, because of a phenomenon known as "heterogeneity's ruse" [39] (see also the

Discussion and the SI), statistical analysis of a net survival curve does not dependably

distinguish variable frailty from immunization. Further biological analysis is often still required

to disentangle their effects [35].

The RLC protocol has already demonstrated substantial advantages over the single high-dose

protocol [36], but some advantages may remain unrealized. Presently, statistical analyses of

many SIV-macaque trials use a standard survival analysis with the logrank test [22-24, 26, 29,

32]. The standard logrank test requires a large-sample Gaussian approximation, whereas the

sparsity of SIV-macaque data suggests the need for an exact test. The development of an exact

test is eased if immunization, variable frailty, or other confounding effects do not occur in SIV-

macaque trials, because then every challenge in an RLC protocol presents the same hazard of

infection to every control animal [40, 41]. In 2012, therefore, Regoes collected a database of 7

articles with SIV-macaque treatment trials, concluding that most did not display immunization or

variable frailty [35].

In 2015, Nolen et al [42] pointed out that a Fisher exact test is valid in a single high-dose

challenge study, but not in a RLC study if it is applied to the total challenge counts in different

arms of the trial. They therefore suggested exact permutation tests based on exchangeability of

control and treatment animals under the null hypothesis of treatment having no effect. After a

careful theoretical analysis, they applied the permutation tests, demonstrating that large-sample

p-value approximations could yield misleading scientific conclusions in small trials. In passing,

however, observe that the single Ellenberger (2006) trial [17] mentioned explicitly in Section 5

of Nolen et al [42] was mildly atypical (see comments in the Results), and that the only

confounding hypotheses Nolen et al [42] simulated involved variable frailty. These two

observations partially motivated the present article by suggesting that statistical analysis might

profit from a comprehensive database of SIV-macaque trials.

Although the present article notes some technical points, it does not improve substantially on

Regoes' statistical analysis of immunization and variable frailty [35]. New SIV-macaque trials

contain many pertinent data, however. The present article therefore assembles a comprehensive

updated database from 24 articles containing the net survival curves for control and treatment

arms in SIV-macaque treatment trials. It has the purview of applying Regoes' statistical analysis

to a larger database to examine the robustness of his scientific conclusions.

The RLC protocol has regularities not available in clinical trials, notably compulsory follow

up and a possibility that between observations, all hazards to controls may be roughly the same,

i.e., all probabilities of infection per challenge are roughly equal. The database may therefore

provide useful biological examples for applications with data more irregular than SIV-macaque

treatment trials but still requiring the development of small-sample statistical tests, e.g., small

studies of breast cancer therapies [43]. As Regoes also pointed out [35], the database also

provides an animal model for exploring repeated low-dose exposure to HIV in humans without

treatment.

The present article is structured as follows. The Materials and Methods section gives the

criteria for trial data to enter our SIV-Macaque database of control results. The human effort in

assembly of the database and its subsequent manipulation was demanding, so the Materials and

Methods section discusses at length the various representations of data in the original articles,

during digitization of survival curves, and for the survival analysis [44]. It then gives Regoes'

models and two additional full models (Geometric priming and Delta frailty), promoting

continuity with Regoes' statistical analysis [35] by using his term "priming models" for

immunization. The Results section presents three tables pertinent to our SIV-Macaque database:

(1) the metadata; (2) the count of controls uninfected after each trial's maximum number of

challenges, with estimates of the infection probability per challenge under the assumption that

the probability is constant; and (3) p-values pertinent to the constancy of infection probabilities

per challenge. The three tables are drawn from an aggregate table in the SI. Finally, the

Discussion section explores the implications of the results.

Results

The comprehensive Excel file in the SI yielded all rounded numbers in Tables 1-3.

Table 1 presents the metadata of the SIV-Macaque database. Each row corresponds to a

single control arm and its trial. The columns indicate: (1) the citation in the present article; (2) its

first author; (3) its year of publication; (4) its ID in PubMed; (5) the source within it for each set

of controls (a citation may contain more than one trial and therefore more than one source of

data); (6) the virus used to challenge; (7) the species of macaque challenged; (8) the challenge

route; and (9) the number of weeks between the regular challenges. In the final 3 columns,

uncommon values are highlighted in yellow, e.g., the Ellenberger (2006) trial [12] used SHIV-

SF162P3 to challenge pigtailed macaques rectally every week, the Wilson (2009) trial [16] used

SIVsmE660 to challenge rhesus macaques rectally every 3 weeks, etc.

The first 5 columns of Table 1 appear in all tables, so the reader can recall references and still

find them immediately and unambiguously in the literature. In all tables, the rows are ordered on

the PubMed ID in Column 4; and the heavy borders enclose the Regoes' dataset [35].

Table 1: **Metadata for the SIV-Macaque database**

Cite	First Author	Year	PMID	Source	Virus	Macaque	Route	Weeks
[10]	Ma	2004	15564513	Table-1	SIVmac251	Rhesus	Vaginal	1
[11]	Van Rompay	2005	15671796	Figure-4A	SIVmac251	Rhesus	Oral	1
[12]	Ellenberger	2006	16725169	Figure-1	SHIV-SF162P3	Pigtailed	Rectal	1
[13]	Wilson	2006	16731926	Table-2	SIVmac239	Rhesus	Rectal	1
[14]	Garcia-Lerma	2008	18254653	Figure-2	SHIV-SF162P3	Rhesus	Rectal	1
[15]	Hansen	2009	19219024	Figure-4c	SIVmac239	Rhesus	Rectal	1
[16]	Wilson	2009	19403685	Table-2	SIVsmE660	Rhesus	Rectal	3
[17]	Hessell	2009	19525965	Figure-2	SHIV-SF162P3	Rhesus	Vaginal	0.5
[18]	Reynolds	2010	20592091	Table-1	SIVsmE660	Rhesus	Rectal	1
[19]	Letvin	2011	21543722	Figure-1A	SIVmac251	Rhesus	Rectal	1
[19]	Letvin	2011	21543722	Figure-1B	SIVsmE660	Rhesus	Rectal	1
[19]	Letvin	2011	21543722	Figure-1C	SIVsmE660	Rhesus	Rectal	1
[20]	Hansen	2011	21562493	Figure-S6	SIVmac239	Rhesus	Rectal	1
[21]	Barouch	2012	22217938	Figure-1e	SIVmac251	Rhesus	Rectal	1
[21]	Barouch	2012	22217938	Figure-3e	SIVmac251	Rhesus	Rectal	1
[22]	Pegu	2013	23175374	Figure-1B	SIVmac251	Rhesus	Rectal	1
[23]	Strbo	2013	23401588	Figure-3AB	SIVmac251	Rhesus	Rectal	1
[23]	Strbo	2013	23401588	Figure-3C	SIVmac251	Rhesus	Rectal	1
[24]	Roederer	2014	24352234	Figure-1a	SIVsmE660	Rhesus	Rectal	1
[25]	Radzio	2016	26743846	Figure-1B	SHIV-SF162P3	Pigtailed	Vaginal	1
[26]	Vaccari	2016	27239761	Figure-1bc	SIVmac251	Rhesus	Rectal	1
[27]	Jensen	2016	27252535	Figure-1C	SIVmac251	Rhesus	Oral	1
[28]	Butler	2016	27355414	Figure-1A	SHIV-SF162P3	Pigtailed	Vaginal	1
[29]	Gordon	2016	27591322	Figure-4E	SIVmac251	Rhesus	Rectal	1
[30]	Xu	2017	28302457	Figure-2c	SIVmac251	Rhesus	Rectal	1
[31]	Eudailey	2018	29359183	Figure-7A	SHIV-1157ipd3N4	Rhesus	Oral	1
[32]	Vaccari	2018	29785023	Figure-1b	SIVmac251	Rhesus	Rectal	1
[33]	Om	2020	32881968	Figure-1B	SHIV-1157ipd3N4	Rhesus	Rectal	1

Table 2 focuses on results pertinent to the Constant Hazard reduced model, which assumes

that every challenge has a constant probability of infection. Columns after Column 5 indicate: (6)

the number n of control animals; (7) the number of uninfected controls remaining after RLCs,

with non-0 values highlighted in yellow; (8) the number m of RLCs that uninfected controls

survived; (9) the MLE $\hat{p}$ of the probability p of infection in the reduced model; and the (10)

lower and (11) upper endpoint of the corresponding 95% confidence interval for p. Thus, Table

2 indicates for macaque trials the potential sparsity of the control arm, controls remaining

uninfected after RLCs, and an approximation to the infection probability p of each challenge.

For the Hansen (2009) trial [15], Regoes [35] gives the control number as $n=15$, whereas the

legend of Figure-4c, the source, gives $n=16$, yielding minor numerical discrepancies of no

scientific substance. In the Hansen (2011) trial [20], the single uninfected control eventually

showed signs of infection, but long after the 1-week window for establishing infection during the

trial. Its designation as "uninfected" is therefore somewhere arbitrary but at least consistent with

the intent of the RLC protocol. Some controls were historical compared to treated animals [26,

29, 32]. Both the Vaccari (2016) trial [26] and the Gordon (2016) trial [29] used 23 concurrent and

24 historical controls. The two control datasets yielded the same net survival curve. To make the

presentation consistent, the statistical analysis appears twice in all tables, once for each trial. The

Vaccari (2018) trial [32] used 6 concurrent and 47=23+24 historical controls.

Table 2: Constant Hazard reduced model for controls in the SIV-Macaque database

Cite	First Author	Year	PMID	Source	n	Not inf	m	p_hat	p_lo	p_hi
[10]	Ma	2004	15564513	Table-1	8	0	5	0.364	0.186	0.572
[11]	Van Rompay	2005	15671796	Figure-4A	5	0	25	0.312	0.125	0.555
[12]	Ellenberger	2006	16725169	Figure-1	14	1	13	0.206	0.119	0.317
[13]	Wilson	2006	16731926	Table-2	8	0	8	0.216	0.105	0.365
[14]	Garcia-Lerma	2008	18254653	Figure-2	18	1	14	0.205	0.128	0.300
[15]	Hansen	2009	19219024	Figure-4c	15	0	12	0.242	0.147	0.358
[16]	Wilson	2009	19403685	Table-2	8	2	5	0.250	0.108	0.443
[17]	Hessell	2009	19525965	Figure-2	4	0	40	0.400	0.146	0.700
[18]	Reynolds	2010	20592091	Table-1	8	1	10	0.318	0.151	0.525
[19]	Letvin	2011	21543722	Figure-1A	20	0	5	0.500	0.349	0.651
[19]	Letvin	2011	21543722	Figure-1B	25	3	12	0.200	0.133	0.281
[19]	Letvin	2011	21543722	Figure-1C	20	5	12	0.124	0.073	0.190
[20]	Hansen	2011	21562493	Figure-S6	28	1	25	0.156	0.107	0.215
[21]	Barouch	2012	22217938	Figure-1e	8	0	6	0.727	0.435	0.924
[21]	Barouch	2012	22217938	Figure-3e	8	0	6	0.533	0.291	0.765
[22]	Pegu	2013	23175374	Figure-1B	11	0	5	0.333	0.189	0.502
[23]	Strbo	2013	23401588	Figure-3AB	12	0	10	0.480	0.293	0.671
[23]	Strbo	2013	23401588	Figure-3C	12	0	6	0.324	0.189	0.483
[24]	Roederer	2014	24352234	Figure-1a	20	0	12	0.339	0.227	0.465
[25]	Radzio	2016	26743846	Figure-1B	5	0	16	0.313	0.125	0.555
[26]	Vaccari	2016	27239761	Figure-1bc	47	0	10	0.367	0.287	0.453
[27]	Jensen	2016	27252535	Figure-1C	6	0	9	0.240	0.103	0.428
[28]	Butler	2016	27355414	Figure-1A	11	5	20	0.045	0.018	0.089
[29]	Gordon	2016	27591322	Figure-4E	47	0	10	0.367	0.287	0.453
[30]	Xu	2017	28302457	Figure-2c	6	2	5	0.235	0.080	0.465
[31]	Eudailey	2018	29359183	Figure-7A	7	1	8	0.176	0.074	0.326
[32]	Vaccari	2018	29785023	Figure-1b	53	0	10	0.414	0.331	0.500
[33]	Om	2020	32881968	Figure-1B	16	7	3	0.237	0.122	0.386

Table 3 focuses on various full models nesting the Constant Hazard reduced model, and it

displays the p-values (uncorrected for multiple testing) from the corresponding LRTs. Columns

after the 5-th indicate the p-values for the following full models: (6) Arithmetic Priming (AP);

(7) Geometric Priming (GP); (8) Step Priming with l_{step}=1 (SP1); (9) Step Priming with l_{step}=2

(SP2); (10) Step Priming with l_{step}=3 (SP3); (11) Beta Frailty (BF); and (12) Delta Frailty (DF).

In the final 7 columns, any p-value less than 0.05 is highlighted in yellow.

Table 3 had 8/28 rows corresponding to trials displaying deviation from a constant infection

probability per challenge at statistical significance $p<0.05$, uncorrected for multiple testing. The

8 trials showing deviation included 1/3 trials using oral challenges, 2/4 trials using vaginal

challenges, and 5/21 using rectal challenges. For each full model of immune priming with a

significant LRT, the second of its two MLE parameters (given in the SI) indicates whether the

infection probability per challenge increased or decreased (tolerance or priming) as the trial

progressed. The direction of change, increase or decrease, was different in different trials.

225   **Table 3: Likelihood ratio test (LRT) p-values for the SIV-Macaque database**

Cite	First Author	Year	PMID	Source	AP	GP	SP1	SP2	SP3	BF	DF
[10]	Ma	2004	15564513	Table-1	0.084	0.060	0.395	0.054	0.250	1.000	1.000
[11]	Van Rompay	2005	15671796	Figure-4A	0.390	0.229	0.099	0.045	0.097	0.110	1.000
[12]	Ellenberger	2006	16725169	Figure-1	0.446	0.406	0.492	0.098	0.131	0.380	0.600
[13]	Wilson	2006	16731926	Table-2	0.017	0.024	0.035	0.036	0.428	1.000	1.000
[14]	Garcia-Lerma	2008	18254653	Figure-2	0.425	0.371	0.837	0.176	0.155	0.321	0.665
[15]	Hansen	2009	19219024	Figure-4c	0.856	0.834	0.799	0.307	0.992	1.000	1.000
[16]	Wilson	2009	19403685	Table-2	1.000	1.000	0.297	0.807	0.768	1.000	0.825
[17]	Hessell	2009	19525965	Figure-2	0.048	0.103	0.016	0.749	0.158	1.000	1.000
[18]	Reynolds	2010	20592091	Table-1	0.020	0.028	0.170	0.036	0.124	0.043	0.023
[19]	Letvin	2011	21543722	Figure-1A	0.810	0.769	0.204	0.489	0.677	0.696	1.000
[19]	Letvin	2011	21543722	Figure-1B	0.101	0.076	0.234	0.027	0.033	0.086	0.119
[19]	Letvin	2011	21543722	Figure-1C	0.214	0.222	0.705	0.407	0.706	0.238	0.217
[20]	Hansen	2011	21562493	Figure-S6	0.269	0.264	0.371	0.319	0.868	0.275	0.320
[21]	Barouch	2012	22217938	Figure-1e	0.910	0.898	0.785	0.412	1.000	1.000	1.000
[21]	Barouch	2012	22217938	Figure-3e	0.504	0.454	0.782	0.601	0.251	1.000	1.000
[22]	Pegu	2013	23175374	Figure-1B	0.006	0.007	0.025	0.022	0.183	1.000	1.000
[23]	Strbo	2013	23401588	Figure-3AB	0.877	0.869	0.847	0.568	0.584	1.000	1.000
[23]	Strbo	2013	23401588	Figure-3C	0.003	0.003	0.020	0.012	0.058	1.000	1.000
[24]	Roederer	2014	24352234	Figure-1a	0.966	0.962	0.898	0.652	0.506	1.000	1.000
[25]	Radzio	2016	26743846	Figure-1B	0.096	0.089	0.030	0.889	0.361	1.000	1.000
[26]	Vaccari	2016	27239761	Figure-1bc	0.267	0.268	0.388	0.258	0.449	1.000	1.000
[27]	Jensen	2016	27252535	Figure-1C	0.655	0.557	0.549	0.569	0.910	1.000	1.000
[28]	Butler	2016	27355414	Figure-1A	0.052	0.133	0.315	0.993	0.600	0.224	0.099
[29]	Gordon	2016	27591322	Figure-4E	0.267	0.268	0.388	0.258	0.449	1.000	1.000
[30]	Xu	2017	28302457	Figure-2c	0.181	0.257	0.488	0.442	0.121	0.305	0.222
[31]	Eudailey	2018	29359183	Figure-7A	0.476	0.386	0.417	0.912	0.364	1.000	1.000
[32]	Vaccari	2018	29785023	Figure-1b	0.178	0.180	0.282	0.246	0.267	1.000	1.000
[33]	Om	2020	32881968	Figure-1B	0.738	0.757	0.538	0.906	1.000	1.000	1.000

226

227

228   **Discussion**

All p-values below are uncorrected for multiple testing.

In the tables, the rows inside the thick borders correspond to Regoes' database [35]. The

rows in Table 2 differ from Regoes' Table 2 in: (1) the subdivision of data from the Letvin

(2011) trial [19]; and (2) the number of controls in the Hansen (2009) trial [15]. Table 3 and

Regoes' Tables 3 and 4 have analogous differences. Otherwise, the present analysis agrees

broadly within numerical error on its overlap with Regoes' [35]. Regoes' Tables 3 and 4

displayed only a few anomalous rows with $p<0.05$, consistent with his conclusion that a constant

hazard of infection per challenge could explain most of his data [35]. The anomaly in the Wilson

(2006) trial was ascribed to possible misidentification of infecting challenges; the anomaly in the

Letvin (2011) Figure-1B trial, to the known biological effect of TRIM5 alleles on SIV infection

(e.g., [52]).

The trials in Table 3 outside Regoes' Tables 3 and 4 continue to display a broad pattern of

anomalies with $p<0.05$, however, making the anomalies harder to dismiss. Perhaps some

anomalies still reflect misidentification of the infecting challenges, but the misidentification then

becomes a persistent feature of the SIV-macaque trials. Other anomalies may depend on trials

using oral or vaginal challenge, which Regoes [35] excluded. The remaining anomalies may

reflect unknown and therefore interesting biological effects. The persistent pattern of anomalies

in Table 3 made Regoes' extensive analysis of the power of the LRTs against alternatives

unnecessary. By themselves, the anomalies clearly discourage any simplifying assumption of a

constant hazard of infection per challenge in SIV-macaque trials, particularly for oral or vaginal

challenges, particularly without specific evidence that supports the assumption's use as an

approximation.

The SIV-Macaque database and Table 3 noticeably enrich the empirical basis for the

statistical analysis of sparse animal trials. Presently, permutation tests provide the leading

candidates for exact tests. The power analysis of the permutation tests in [42], e.g., focused on

confounding hypotheses of variable frailty, namely, the two frailty full models in the present

article. The frailty models accurately reflect experimenters' concerns about the confounding

effects of, e.g., sex [33] or the *Mamu-A*01* class I allele [19]. Table 3 suggests, however, that a

statistical exploration of the two confounding hypotheses should probably be more concerned

about immunization than variable frailty in [42].

The pattern of anomalies in Table 3 also has a notable feature. Except for the Hessell (2009)

trial [17] and the Radzio (2016) trial [25], both of which challenged vaginally, whenever a full

model in Table 3 displayed $p<0.05$, the full model SP2 for step priming after two challenges in

the same row also displayed $p<0.05$. Under a weekly challenge, the full model SP2 could be

interpreted as changes in control immunity occurring after 2 weeks, e.g., the recommended

period for full immunity after COVID-19 vaccination. The immune changes were inconsistent:

neither immune priming nor immune tolerance occurred consistently (see end of the Results and

the SI). For the full model SP2, e.g., 3/6 changes indicated increased probability of infection per

challenge (tolerance), rather than priming. Typically, however, net survival curves alone cannot

distinguish variable frailty from immunization ("heterogeneity's ruse" [39]). The inconsistent

immune interpretation of the SP2 full model does, however, bear on previous biological

conclusions that repeated exposure without treatment may have no effect [35] or may even

reduce [53] human infectability in HIV acquisition.

The exceptions above, the Hessell (2009) trial [17] and the Radzio (2016) trial [25], used

vaginal challenge. Neither had full model SP2 displaying $p<0.05$, but instead had full model SP1

for step priming after one challenge displaying $p<0.05$, suggesting that immune priming in

vaginal challenge may occur faster than in rectal challenge.

The priming models do have some mild technical superiorities over frailty models, namely:

(1) in the context of net survival curves, priming models are more general than frailty models, at

least in one sense (see the SI); (2) they have strictly concave log-likelihoods (see the SI), slightly

simplifying numerical computations; and (3) they nest the reduced model in the interior of their

parameter space (see the Materials and Methods section), thereby satisfying the usual hypotheses

for theorems about the LRT [50]. Table 3 shows that for most trials, however, if one full model

achieves statistical significance at $p<0.05$, several others often follow. Typically, the underlying

SIV-macaque data are sparse, and the net survival curves are correspondingly nondescript. If

several models show significance, then, no specific biological interpretation can be considered

pre-eminent without further investigation.

If some controls remain uninfected after RLCs, e.g., it suggests the confounding hypothesis

that the controls may have an unusually low infectability. Fortunately, hyper-challenges with

increased viral doses after the RLCs can remove the doubt about low infectability [16, 31, 33],

although its cause, immunization or variable frailty, would remain unclear.

To summarize, then, the present article updated Regoes' database [35] with the SIV-Macaque

database in the SI. The updated database displays several trials where the controls appear not to

have a constant hazard of infection, and where full models of immune priming (or tolerance) or

variable frailty fit control data significantly better at uncorrected $p<0.05$ than the reduced model

positing a constant probability of infection per challenge. Given sparse animal data, exact tests

have advantages over large-sample tests like the standard logrank test that use Gaussian

approximations. Unfortunately, the present results show that any statistical test under

consideration should also be shown to be robust against confounding hypotheses about control

animals [54], notably both repeated low-dose immunization and variations in frailty.

Biologically, the data from SIV-macaque trials are consistent with the RLC protocol changing

the infectability of animals after about two weeks in rectal or oral challenge, and possibly even

earlier in vaginal challenge. Explanations other than immunization remain very possible,

however, particularly because changes in infectability were sometimes consistent with immune

priming, and sometimes with immune tolerance. Nonetheless, the results have some bearing on

previous biological conclusions that repeated exposure to HIV without treatment may have no

effect [35] or may even reduce human infectability [53].

Materials and Methods

The SIV-Macaque database. Our meta-analysis [45] permitted an article on an SIV-

macaque trial to contribute data to the SIV-Macaque database if it had: (1) at least 5 control

animals; (2) at least 3 repeats of the same challenge; (3) a regular challenge schedule; (4)

infection assigned to a specific challenge; and (5) publicly available data. Regoes [35] used more

stringent criteria than ours, including a restriction to rectal challenges, so his database has 7

articles [12-16, 19, 20] and is a subset of ours. Our SIV-Macaque database includes 2 articles

predating his; 2 articles contemporary with his; and 13 articles postdating his.

Table 1 in the Results section displays metadata for our SIV-Macaque database. To favor

completeness, our database included two borderline cases: (1) an article exploring the RLC

protocol with control animals only [10]; and (2) an article using (almost regular) twice-weekly

challenges [17]. The present article followed Regoes [35] in including the Wilson (2009) trial

[16], which followed its RLCs with viral hyper-challenges at greatly increased doses to test the

infectability of animals uninfected after Challenge 5. As in [35], our SIV-Macaque database

includes data corresponding to the RLCs, truncating the challenge data after the RLCs and before

any hyper-challenges. The Eudailey (2018) trial [31] also used hyper-challenges after Challenge

8. The Om (2020) trial [33] increased the challenge after Challenge 3, possibly just readjusting

the challenge ensure infection of controls, rather than testing their infectability with hyper-

challenges.

A reanalysis in the Letvin (2011) trial [19] regrouped some of its data to refine biological

conclusions about the *Mamu-A*01* allele. Similarly, the Om (2020) trial [33] regrouped their

data by sex to refine biological conclusions about vaccine protection by sex. The present article

avoided analyzing data twice, so it ignored the regroupings.

Each article yielded challenge data for its trial arms. If the data were figures with net survival

curves, the Windows® 10 Snip & Sketch tool captured the curves with a screen shot. Where

possible, two people independently digitized each net survival curve with the WebPlotDigitizer

(version 4.4) at https://apps.automeris.io/wpd/, later reconciling any discrepancies. No other data

format required digitization.

Some articles contained more than one set of control data. Within the articles, however,

every arm of every trial mapped to precisely one set of control data. All trial data in an article

were grouped by their control arm, and each group was stored as a DataFrame in a separate file

of comma-separated values (CSV), following the precepts of tidy data [46]. Each filename

corresponded to a composite key combining the article ID in PubMed and the source graphic(s)

or table(s) containing its "Control", anchoring all data unambiguously into the public literature.

In the Results section, the columns "PMID" and "Source" in each table implicitly display the

composite key for the relevant control.

The next sections detail the various representations used to assemble, retrieve, and analyze

the SIV-Macaque database. Flexible representation was indispensable in reducing human effort.

Data Representation by Individuals. To focus the following discourse, unless stated otherwise,

consider only the control arm of an RLC protocol in a specific trial. Following Regoes' notation

[35] where feasible, let $n = n_1$ count the controls, initially uninfected and indexed by

$j \in \{1,...,n\}$. Each Animal j receives serial challenges at regular intervals, e.g., weekly. The

challenges are numbered consecutively $t \in \{1,2,3,...\}$. Let Challenge c_j be the final challenge

to Animal j, with $m = \max_j c_j$ the largest challenge index among the controls, so $c_j \leq m$. To

account for frequent right-censoring in general survival analysis [44], Regoes [35] carefully

permitted the possibility that the final Challenge c_j does not infect Animal j, even if $c_j < m$.

On one hand, if Animal j becomes infected, Criterion (4) above specifies that the index c_j

of the infecting challenge is known. Some trials continue to challenge infected animals, but for

present purposes, we ignore such challenges and consider c_j as the final challenge to Animal j.

On the other hand, the final Challenge c_j may not infect, so some animals may remain

uninfected at the end of the trial.

Consider, therefore, the final status $(\mathbf{c}, \mathbf{i}) = \{(c_j, i_j) : j = 1,...,n\}$ of every Animal j, where i_j

indicates whether Challenge c_j infected Animal j: $i_j = 1$ indicates infection, and $i_j = 0$

otherwise. For a generic animal, drop the subscript j, so (c, i) denotes its final status. The final

status $(\mathbf{c}, \mathbf{i})$ is particularly useful if an analysis uses multiple covariates (e.g., alleles) to

distinguish individual animals, a rare circumstance indeed in sparse animal trials. We therefore

summarize the final status $(\mathbf{c}, \mathbf{i})$ with the statistics relevant to discrete-time hazards in analysis of

net survival curves [44].

Hazards in Discrete Time. Given sampled Animals $j \in \{1, ..., n\}$, assume a probability model

where the corresponding components (c_j, i_j) of the final status $(\mathbf{c}, \mathbf{i})$ are mutually independent.

The hazard h_t gives the probability of infection of a randomly sampled control at Challenge t,

conditional on previous challenges not having infected it. The Iverson bracket $[A]$ provides a

standard notation for the indicator of the event A: $[A] = 1$ if the event A occurs and $[A] = 0$

otherwise [47]. The probability that a generic control has final status $(c, 0)$ is

$(1 - h_1)...(1 - h_{c-1})(1 - h_c)$; final status $(c, 1)$, $(1 - h_1)...(1 - h_{c-1})h_c$. In either case, the natural

logarithm of the probability that a particular control has final status (c, i) is

$$\ln L = \sum_{t=1}^{m} \left\{ [t = c][i = 1]\left(\ln h_t - \ln(1 - h_t)\right) + [t \le c]\ln(1 - h_t) \right\}, \qquad (1)$$

proved as follows. The terms on the right side are 0 for any Challenge t after Challenge c, so

the sum on the right may be broken into terms for $t = 1, ..., c - 1$ and a term for $t = c$. Enumeration

of two cases $[i = 0]$ and $[i = 1]$ then shows that Eq (1) gives the natural logarithm of the stated

probability.

Let $n_t = \sum_{j=1}^{n} [t \le c_j]$ count the controls remaining in the trial at least to Challenge t;

$d_t = \sum_{j=1}^{n} [t = c_j][i_j = 1]$, the controls infected by Challenge t. The sum over all controls for

log-likelihood for the final status $(\mathbf{c}, \mathbf{i})$ in Eq (1) can therefore be rewritten as

$$\ln L = \sum_{j=1}^{n} \sum_{t=1}^{m} \left\{ \left[t = c_j \right] \left[i_j = 1 \right] \left(\ln h_t - \ln \left(1 - h_t \right) \right) + \left[t \le c_j \right] \ln \left(1 - h_t \right) \right\}$$

380
$$= \sum_{t=1}^{m} \left\{ d_t \left(\ln h_t - \ln \left(1 - h_t \right) \right) + n_t \ln \left(1 - h_t \right) \right\} \qquad , \qquad (2)$$

$$= \sum_{t=1}^{m} d_t \ln h_t + \sum_{t=1}^{T} \left(n_t - d_t \right) \ln \left(1 - h_t \right)$$

the standard expression from Kaplan-Meier survival analysis [44], put into the present context.

The representation by deaths $\left\{ \left(n_t, d_t \right) : t = 1,...,m \right\}$ quickly yields a net survival curve (e.g.,

Figure 3 of [18] or Fig 1 below). On one hand, Kaplan-Meier survival analysis derives its

generality from handling right-censoring due to incomplete patient follow-up. On the other hand,

the RLC protocol in animal trials typically makes follow-up compulsory, so every uninfected

control receives its next scheduled challenge. Challenge t then infects $d_t = n_t - n_{t+1}$ of the n_t

controls uninfected before Challenge t ($t = 1,...,m$), where n_{m+1} counts the uninfected controls at

the end of the trial (possibly, $n_{m+1} = 0$). At the end of a typical trial, each of the n_{m+1} uninfected

controls have therefore survived m challenges. Assume below that animal trials have

compulsory follow up within each arm, yielding several compact alternatives to the

representation by deaths $\left\{ \left(n_t, d_t \right) : t = 1,...,m \right\}$.

Alternative Data Representations (see Fig 1). The representation by survivors $\left(n_1, n_2,..., n_{m+1} \right)$

satisfies $n_1 \ge n_2 \ge ... \ge n_{m+1}$, and it permits challenge data from each arm in a trial to occupy one

column in a DataFrame. The SIV-Macaque database in the SI is a directory of CSV files, each a

DataFrame for one trial. Most trials enforce the same maximum m of challenges for each arm,

so most of the DataFrames are well-formed, with m rows in every column and 0s padding an

arm after infection of all animals in it. In fact, contrary to an assertion of plurality elsewhere

[42], only one trial in the SIV-Macaque database, Ellenberger (2006) [12], overtly permitted m

to vary with the arm. In the one trial [12], challenge was discontinued before complete infection

of controls and before the end of the trial. The corresponding DataFrame therefore has NaNs

padding the control arm, instead of 0s.

405   **Fig 1. A hypothetical net survival curve for a control arm with 8 animals**

Fig 1 plots percent uninfected animals against the number of challenges for a control arm with 8 animals. Typical

net survival curves use percentages to facilitate comparison of results from trial arms with differing numbers of

animals. At each Challenge $t = 1,...,m$, the open point (t, n_t) gives the percentage of the originally uninfected

animals receiving Challenge t; the closed point $(t, n_t - d_t)$ drops by d_t due to animals infected by Challenge t

(if not obscured by an open point when $d_t = 0$). By convention, n_{m+1} counts the uninfected controls at the end of

the trial, e.g., $n_{m+1} = 1$ in Fig 1. Fig 1 also shows that $m = 7$, so with $n = 8$ animals in the arm, the representation by

deaths $\{(n_t, d_t) : t = 1,...,m\}$ is { (8,4), (4,2), (2,0), (2,1), (1,0), (1,0), (1,0) }. The representation by survivors

$(n_1, n_2,..., n_{m+1})$ uses only the open circles and n_{m+1}: (8,4,2,2,1,1,1,1). The minimal representation { (

$\tau(i) - 1, n_{\tau(i)}$): $i = 1,...,k$ } uses only the closed points and (m, n_{m+1}): { (0,8), (1,4), (2,2), (4,1), (7,1) }.

A minimal representation reduces the human effort in digitizing the net survival curves from

SIV-macaque trials. In the representation by survivors, omit challenges where no infections

occurred to yield $\left(n_{\tau(1)}, n_{\tau(2)}, \ldots, n_{\tau(k-1)}, n_{\tau(k)} \right)$, where $n_1 = n_{\tau(1)} > n_{\tau(2)} > \ldots > n_{\tau(k-1)} \geq n_{\tau(k)} = n_{m+1}$.

Formally, start with $\tau(1) = 1$ and define $\tau(i+1) = \min\left\{ t > \tau(i) : n_{\tau(i)} > n_t \text{ for } t \leq m \right\}$ recursively

until the set is empty, say, after $\tau(k-1)$ has been defined. Then, define $\tau(k) = m+1$. The

minimal representation $\left\{ \left(\tau(i) - 1, n_{\tau(i)} \right) : i = 1, \ldots, k \right\}$ compacts the representation by survivors

without loss and greatly speeds digitization of net survival curves. Most experiments show the

net survival curve for each arm as percent survival, as in Fig 1. Accordingly, we digitized only

the points corresponding to the minimal representation, multiplied the percentages by $n = n_1$, and

then rounded to integers to yield the minimal representation. If two people digitize and then

disagree, the minimal representation facilitates a rapid reconciliation of their results.

Finally, a few trials (e.g., [13, 16]) use a representation $\left(t_j : j = 1, \ldots, n \right)$ by individual

infecting challenges [35], where Challenge t_j infects Animal j. Retain our convention that

$t_j = m+1$ if Animal j remains uninfected throughout the trial. Then, the data may be converted

to $n_t = \#\left\{ t_j \geq t : j = 1, 2, \ldots, n \right\}$ for $t = 1, \ldots, m$, where "#" counts the number of elements in the set.

Constant Hazard reduced model. The following nests a reduced model H_0, the Constant

Hazard model, within several different full models, to detect deviation from H_0 with Likelihood

Ratio tests (LRTs). The Constant Hazard model (Regoes' Geometric infection model [35]) posits

that all challenges yield mutually independent results and have a constant probability p of

infecting a control. Every animal is equally infectable, so under the reduced model H_0, the

hazard per challenge is the constant

$$H_0 : h_t = p. \tag{3}$$

438   The corresponding log-likelihood

$$\ln L_0 = \sum_{t=1}^{m} d_t \ln p + \sum_{t=1}^{m} (n_t - d_t) \ln (1 - p) \tag{4}$$

440   yields the maximum likelihood estimator (MLE)

$$\hat{p} = \sum_{t=1}^{m} d_t \, / \sum_{t=1}^{m} n_t, \tag{5}$$

a convenient initialization for maximizing likelihoods of the full models. Even if the reduced

model is rejected, Eq (5) is still the fraction of all challenges that infected and provides a

practical summary of the average potency of a challenge.

For brevity, to truncate a real number x to a probability $U(x)$ between 0 and 1, define the

unit function $U(x) = x$ for $0 \le x \le 1$, with $U(x) = 0$ for $x < 0$ and $U(x) = 1$ for $x > 1$. Below,

we characterize several full models $H_\bullet$, where "$\bullet$" is an arbitrary subscript, simply by

specifying their hazard h_t. If required, Eq (2) then implicitly gives the corresponding likelihoods

$L_\bullet$ and log-likelihoods $\ell_\bullet = \ln L_\bullet$. Our full models include all of Regoes' full models [35].

The first elaboration of the Constant Hazard model that Regoes [35] addressed was immune

priming: even if the animals had the same infectability, their infectability might evolve during

the trial because the RLCs provided immunization against the virus, either in the direction of

immune priming or immune tolerance. In mathematical models, negative immune priming

corresponds to immune tolerance.

Arithmetic priming full model (AP). Regoes [35] examined a full model equivalent to

$$H_{AP} : h_t = U\left(p + (t-1)\varepsilon\right). \tag{6}$$

H_{AP} reduces to H_0 if $\varepsilon = 0$. Priming corresponds to $\varepsilon < 0$; tolerance, to $\varepsilon > 0$.

Geometric priming full model (GP). Arithmetic priming has a hard boundary at 0 from the unit

function. A geometric sequence avoids it:

$$H_{GP} : h_t = U\left(pr^{t-1}\right). \tag{7}$$

H_{GP} reduces to H_0 if $r = 1$. Priming corresponds to $r < 1$; tolerance, to $r > 1$.

Step priming full model (SP1, SP2, SP3). Following Regoes [35] (see also [48]), consider full

models where the hazard is piecewise constant, changing from p to p_2 after Challenge l_{step}=1,2,

or 3.

$$H_{SP} : h_t = \begin{cases} p, & t \le l_{step} \\ p_2, & t > l_{step} \end{cases}. \tag{8}$$

H_{SP} reduces to H_0 if $p_2 = p$. Priming corresponds to $p_2 < p$; tolerance, to $p_2 > p$.

The log-likelihoods for priming models are all strictly concave in their two variables (p, ε),

(p, r), or (p, p_2) (see the SI), so hill-climbing methods always determine the MLE as the

unique global maximum. Regoes [35] also addressed frailty models without priming, where

animals might have different intrinsic infectabilities that remained constant during challenge

[49]. The log-likelihoods for the two frailty models below are not concave in general (see the SI),

mildly complicating numerical computation. In addition, the two frailty models nest the reduced

model on the boundary of their parameter space, violating the hypotheses of standard statistical

theorems [50]. The present article has the purview of examining the robustness of Regoes'

scientific conclusions against expanding his database, making technically arduous statistical

improvements peripheral.

Beta Frailty full model (BF). The Beta distribution with parameters (a,b) has probability

density function

$$f_{BF}(x) = \frac{(a+b-1)!}{(a-1)!(b-1)!} x^{a-1}(1-x)^{b-1}, \tag{9}$$

480   where factorials replace the cumbrous Gamma function notation, even if a and b are not

481   integers. The prefactor is the reciprocal of the Beta function

$$B(a,b) = \frac{(a-1)!(b-1)!}{(a+b-1)!} = \int_0^1 x^{a-1}(1-x)^{b-1}\,dx, \tag{10}$$

483   yielding the required normalization $\int_0^1 f_{BF}(x)\,dx = 1$.

484   Consider a frailty model where each challenge to Animal j has a probability x of infection,

485   the infectability x being intrinsic to Animal j with the probability density function in Eq (9).

486   The corresponding hazard, the probability of infection on Challenge t conditional on previous

487   non-infection, is

$$H_{BF}: h_t = \frac{\int_0^1 x(1-x)^{t-1} f_{BF}(x)\,dp}{\int_0^1 (1-x)^{t-1} f_{BF}(x)\,dp} = \frac{\dfrac{a!(b+t-2)!}{(a+b+t-1)!}}{\dfrac{(a-1)!(b+t-2)!}{(a+b+t-2)!}} = \frac{a}{a+b+t-1}, \tag{11}$$

489   equivalent to animal frailty with a beta distribution (see the SI). Hence, the log-likelihood

$$\ln L_{BF} = \sum_{t=1}^m d_t \ln \frac{a}{a+b+t-1} + \sum_{t=1}^m (n_t - d_t)\ln \frac{b+t-1}{a+b+t-1}. \tag{12}$$

To nest H_0 within H_{BF}, Regoes [35] noted that the beta distribution in Eq (9) can be

parametrized by its mean $p = a/(a+b)$ and variance $v = ab/(a+b)^2/(a+b+1)$. For $v > 0$, let

$s = p(1-p)/v - 1$ so $a = sp$, $b = s(1-p)$. H_{BF} reduces to H_0 if $v \downarrow 0$ with p fixed, i.e.,

$a,b \to \infty$ with $p = a/(a+b)$ fixed.

Delta Frailty full model (DF) [34, 49]: In the context of a treatment trial, Hudgens et al [34]

posit a frailty model for controls with parameters (θ, p). In their model, each control is

independently susceptible with probability $1-\theta$. Only susceptible controls can be infected, and

each challenge infects susceptible controls with probability p. The analog to Eq (9) is

$$f_{DF}(x) = (1-\theta)\delta(x-p) + \theta\delta(x),\qquad\qquad(13)$$

where $\delta(x)$ symbolizes the Dirac delta, which satisfies $\int_0^1 h(x)\delta(x)dx = h(0)$ for functions h.

Thus,

$$H_{DF} : h_t = \frac{\int_0^1 x(1-x)^{t-1} f_{DF}(x)dx}{\int_0^1 (1-x)^{t-1} f_{DF}(x)dx} = \frac{(1-\theta)p(1-p)^{t-1}}{\theta + (1-\theta)(1-p)^{t-1}}.\qquad\qquad(14)$$

H_{DF} reduces to H_0 if $\theta = 0$.

Model Parameter Estimation. Animal data are sparse, so if the reduced model holds, we base a

confidence interval for p on likelihood ratios [35, 51], rather than moments.

$$CI_{1-\alpha}(p) = \left\{ x : 2\left[\ell_0(\hat{p}) - \ell_0(x) \right] \le \chi^2_{1,1-\alpha} \right\},\qquad\qquad(15)$$

where α is the desired significance level and $\chi^2_{1,1-\alpha}$ is the $(1-\alpha)$-percentile of the chi-squared

distribution with one degree of freedom, i.e., for a random variate X with the distribution,

$P\left(X \le \chi^2_{1,1-\alpha}\right) = 1-\alpha$. We take $\alpha = 0.05$ to yield 95% confidence intervals. To facilitate

simulation of actual trials, the Excel file in the SI contains the MLE for each full model, along

with the corresponding Fisher information matrix, if calculable numerically.

Acknowledgments

I acknowledge the invaluable aid of Ms. Chelsea Zebaze in digitizing data from animal trials and

several useful conversations with Dr DoHwan Park and Dr Junyong Park.

References

1.   Veazey RS, Lackner AA. Nonhuman Primate Models and Understanding the Pathogenesis of HIV Infection and AIDS. ILAR Journal. 2017;58(2):160-71.

2.   Arthur LO, Bess JW, Jr., Waters DJ, Pyle SW, Kelliher JC, Nara PL, et al. Challenge of chimpanzees (Pan troglodytes) immunized with human immunodeficiency virus envelope glycoprotein gp120. J Virol. 1989;63(12):5046-53.

3.   Spouge JL. Statistical-Analysis of Sparse Infection Data and Its Implications For Retroviral Treatment Trials in Primates. Proc Natl Acad Sci U S A. 1992;89(16):7581-5.

4.   Roederer M. Parsimonious Determination of the Optimal Infectious Dose of a Pathogen for Nonhuman Primate Models. PLoS Pathog. 2015;11(8):e1005100.

5.   Siegel S, Castellan NJ. Nonparametric Statistics for the Behavioral Sciences. 2 ed. New York: MacGraw-Hill; 1988.

6.   Cohen J. NIH to End Chimp Breeding for Research. Science. 2007;316(5829):1265.

7.   Marx JL. Progress reported on mouse models for AIDS. Science. 1988;242(4886):1638.

8.   Marsden MD. Benefits and limitations of humanized mice in HIV persistence studies. Retrovirology. 2020;17(1):7.

9.   Keele BF, Jones JH, Terio KA, Estes JD, Rudicell RS, Wilson ML, et al. Increased mortality and AIDS-like immunopathology in wild chimpanzees infected with SIVcpz. Nature. 2009;460(7254):515-9.

10.   Ma Z-M, Abel K, Rourke T, Wang Y, Miller CJ. A Period of Transient Viremia and Occult Infection Precedes Persistent Viremia and Antiviral Immune Responses during Multiple Low-Dose Intravaginal Simian Immunodeficiency Virus Inoculations. 2004;78(24):14048-52.

11.   Van Rompay KKA, Abel K, Lawson JR, Singh RP, Schmidt KA, Evans T, et al. Attenuated Poxvirus-Based Simian Immunodeficiency Virus (SIV) Vaccines Given in Infancy Partially Protect Infant and Juvenile Macaques Against Repeated Oral Challenge With Virulent SIV. JAIDS Journal of Acquired Immune Deficiency Syndromes. 2005;38(2).

12.   Ellenberger D, Otten RA, Li B, Aidoo M, Rodriguez IV, Sariol CA, et al. HIV-1 DNA/MVA vaccination reduces the per exposure probability of infection during repeated mucosal SHIV challenges. Virology. 2006;352(1):216-25.

13.   Wilson NA, Reed J, Napoe GS, Piaskowski S, Szymanski A, Furlott J, et al. Vaccine-induced cellular immune responses reduce plasma viral concentrations after repeated low-dose challenge with pathogenic simian immunodeficiency virus SIVmac239. J Virol. 2006;80(12):5875-85.

14.   García-Lerma JG, Otten RA, Qari SH, Jackson E, Cong ME, Masciotra S, et al. Prevention of rectal SHIV transmission in macaques by daily or intermittent prophylaxis with emtricitabine and tenofovir. PLoS Med. 2008;5(2):e28.

15.   Hansen SG, Vieville C, Whizin N, Coyne-Johnson L, Siess DC, Drummond DD, et al. Effector memory T cell responses are associated with protection of rhesus monkeys from mucosal simian immunodeficiency virus challenge. Nat Med. 2009;15(3):293-9.

16.   Wilson NA, Keele BF, Reed JS, Piaskowski SM, MacNair CE, Bett AJ, et al. Vaccine-induced cellular responses control simian immunodeficiency virus replication after heterologous challenge. J Virol. 2009;83(13):6508-21.

17.   Hessell AJ, Poignard P, Hunter M, Hangartner L, Tehrani DM, Bleeker WK, et al. Effective, low-
titer antibody protection against low-dose repeated mucosal SHIV challenge in macaques. Nat Med.
2009;15(8):951-4.
18.   Reynolds MR, Weiler AM, Piaskowski SM, Kolar HL, Hessell AJ, Weiker M, et al. Macaques
vaccinated with simian immunodeficiency virus SIVmac239Delta nef delay acquisition and control
replication after repeated low-dose heterologous SIV challenge. J Virol. 2010;84(18):9190-9.
19.   Letvin NL, Rao SS, Montefiori DC, Seaman MS, Sun Y, Lim SY, et al. Immune and Genetic
Correlates of Vaccine Protection Against Mucosal Infection by SIV in Monkeys. Sci Transl Med.
2011;3(81):81ra36.
20.   Hansen SG, Ford JC, Lewis MS, Ventura AB, Hughes CM, Coyne-Johnson L, et al. Profound
early control of highly pathogenic SIV by an effector memory T-cell vaccine. Nature.
2011;473(7348):523-7.
21.   Barouch DH, Liu J, Li H, Maxfield LF, Abbink P, Lynch DM, et al. Vaccine protection against
acquisition of neutralization-resistant SIV challenges in rhesus monkeys. Nature. 2012;482(7383):89-93.
22.   Pegu P, Vaccari M, Gordon S, Keele BF, Doster M, Guan Y, et al. Antibodies with high avidity
to the gp120 envelope protein in protection from simian immunodeficiency virus SIV(mac251)
acquisition in an immunization regimen that mimics the RV-144 Thai trial. J Virol. 2013;87(3):1708-19.
23.   Strbo N, Vaccari M, Pahwa S, Kolber MA, Doster MN, Fisher E, et al. Cutting edge: novel
vaccination modality provides significant protection against mucosal infection by highly pathogenic
simian immunodeficiency virus. J Immunol. 2013;190(6):2495-9.
24.   Roederer M, Keele BF, Schmidt SD, Mason RD, Welles HC, Fischer W, et al. Immunological
and virological mechanisms of vaccine-mediated protection against SIV and HIV. Nature.
2014;505(7484):502-8.
25.   Radzio J, Henning T, Jenkins L, Ellis S, Farshy C, Phillips C, et al. Combination Emtricitabine
and Tenofovir Disoproxil Fumarate Prevents Vaginal Simian/Human Immunodeficiency Virus Infection
in Macaques Harboring Chlamydia trachomatis and Trichomonas vaginalis. J Infect Dis.
2016;213(10):1541-5.
26.   Vaccari M, Gordon SN, Fourati S, Schifanella L, Liyanage NP, Cameron M, et al. Adjuvant-
dependent innate and adaptive immune signatures of risk of SIVmac251 acquisition. Nat Med.
2016;22(7):762-70.
27.   Jensen K, Nabi R, Van Rompay KKA, Robichaux S, Lifson JD, Piatak M, et al. Vaccine-Elicited
Mucosal and Systemic Antibody Responses Are Associated with Reduced Simian Immunodeficiency
Viremia in Infant Rhesus Macaques. Journal of Virology. 2016;90(16):7285-302.
28.   Butler K, Ritter JM, Ellis S, Morris MR, Hanson DL, McNicholl JM, et al. A Depot
Medroxyprogesterone Acetate Dose That Models Human Use and Its Effect on Vaginal SHIV
Acquisition Risk. J Acquir Immune Defic Syndr. 2016;72(4):363-71.
29.   Gordon SN, Liyanage NP, Doster MN, Vaccari M, Vargas-Inchaustegui DA, Pegu P, et al.
Boosting of ALVAC-SIV Vaccine-Primed Macaques with the CD4-SIVgp120 Fusion Protein Elicits
Antibodies to V2 Associated with a Decreased Risk of SIVmac251 Acquisition. J Immunol.
2016;197(7):2726-37.
30.   Xu H, Andersson AM, Ragonnaud E, Boilesen D, Tolver A, Jensen BAH, et al. Mucosal
Vaccination with Heterologous Viral Vectored Vaccine Targeting Subdominant SIV Accessory Antigens
Strongly Inhibits Early Viral Replication. EBioMedicine. 2017;18:204-15.
31.   Eudailey JA, Dennis ML, Parker ME, Phillips BL, Huffman TN, Bay CP, et al. Maternal HIV-1
Env Vaccination for Systemic and Breast Milk Immunity To Prevent Oral SHIV Acquisition in Infant
Macaques. mSphere. 2018;3(1).
32.   Vaccari M, Fourati S, Gordon SN, Brown DR, Bissa M, Schifanella L, et al. HIV vaccine
candidate activation of hypoxia and the inflammasome in CD14(+) monocytes is associated with a
decreased risk of SIV(mac251) acquisition. Nat Med. 2018;24(6):847-56.

33.   Om K, Paquin-Proulx D, Montero M, Peachman K, Shen XY, Wieczorek L, et al. Adjuvanted
HIV-1 vaccine promotes antibody-dependent phagocytic responses and protects against heterologous
SHIV challenge. Plos Pathogens. 2020;16(9).
34.   Hudgens MG, Gilbert PB, Mascola JR, Wu CD, Barouch DH, Self SG. Power to detect the
effects of HIV vaccination in repeated low-dose challenge experiments. J Infect Dis. 2009;200(4):609-13.
35.   Regoes RR. The role of exposure history on HIV acquisition: insights from repeated low-dose
challenge studies. PLoS Comput Biol. 2012;8(11):e1002767.
36.   Regoes RR, Longini IM, Feinberg MB, Staprans SI. Preclinical assessment of HIV vaccines and
microbicides by repeated low-dose virus challenges. PLoS Medicine. 2005;2(8):798-807.
37.   Shedlock DJ, Silvestri G, Weiner DB. Monkeying around with HIV vaccines: using rhesus
macaques to define 'gatekeepers' for clinical trials. Nature Reviews Immunology. 2009;9(10):717-28.
38.   Thippeshappa R, Kimata JT, Kaushal D. Toward a Macaque Model of HIV-1 Infection:
Roadblocks, Progress, and Future Strategies. 2020;11(882).
39.   Vaupel JW, Yashin AI. Heterogeneity's ruses: some surprising effects of selection on population
dynamics. Am Stat. 1985;39(3):176-85.
40.   Hudgens MG, Gilbert PB. Assessing vaccine effects in repeated low-dose challenge experiments.
Biometrics. 2009;65(4):1223-32.
41.   Kang C, Huang Y, Miller CJ. A discrete-time survival model with random effects for designing
and analyzing repeated low-dose challenge experiments. Biostatistics. 2015;16(2):295-310.
42.   Nolen T, Hudgens M, Senb P, Koch G. Analysis of Repeated Low-Dose Challenge Studies. Stat
Med. 2015;34.
43.   Heinze G, Gnant M, Schemper M. Exact log-rank tests for unequal follow-up. Biometrics.
2003;59(4):1151-7.
44.   Kaplan EL, Meier P. Nonparametric-Estimation from Incomplete Observations. J Am Stat Assoc.
1958;53(282):457-81.
45.   Nakagawa S, Noble DWA, Senior AM, Lagisz M. Meta-evaluation of meta-analysis: ten
appraisal questions for biologists. BMC Biology. 2017;15(1):18.
46.   Wickham H. Tidy Data. Journal of Statistical Software. 2014;59(10):1-23.
47.   Knuth DE. 2 Notes on Notation. American Mathematical Monthly. 1992;99(5):403-22.
48.   Matthews DE, Farewell VT. On testing for a constant hazard against a change-point alternative.
Biometrics. 1982;38(2):463-8.
49.   Longini IM, Halloran ME. A frailty mixture model for estimating vaccine efficacy. Journal of the
Royal Statistical Society Series C-Applied Statistics. 1996;45(2):165-73.
50.   Self SG, Liang KY. Asymptotic Properties of Maximum-Likelihood Estimators and Likelihood
Ratio Tests under Nonstandard Conditions. J Am Stat Assoc. 1987;82(398):605-10.
51.   Cox DR. Analysis of Binary Data. London: Methuen; 1970.
52.   Wu F, Kirmaier A, White E, Ourmanov I, Whitted S, Matsuda K, et al. TRIM5α Resistance
Escape Mutations in the Capsid Are Transferable between Simian Immunodeficiency Virus Strains.
Journal of virology. 2016;90(24):11087-95.
53.   Stranford SA, Skurnick J, Louria D, Osmond D, Chang SY, Sninsky J, et al. Lack of infection in
HIV-exposed individuals is associated with a strong CD8(+) cell noncytotoxic anti-HIV response. Proc
Natl Acad Sci U S A. 1999;96(3):1030-5.
54.   MacKinnon DP, Krull JL, Lockwood CM. Equivalence of the mediation, confounding and
suppression effect. Prev Sci. 2000;1(4):173-81.

Supporting information

Supplementary Information.zip. This is a ZIP file of the following items.

README.txt. This is an index for the Supporting Information.

Supplementary_Methods.docx. This contains additional methods and a legend for
Supplementary_Table.xlsx.

Supplementary_Table.xlsx. This table includes Tables 1-3 and other information in
floating-point precision.

SIV-Macaque_Net_Survival_Curves/. This is a directory containing the CSV files,
DataFrames containing the data from the net survival curves in the SIV-Macaque
database.

666