A new protocol for multiple muscle mapping using nTMS

Fang Jin, Sjoerd Bruijn[*], Andreas Daffertshofer

Faculty of Behavioural and Movement Sciences, Department of Human Movement Sciences, Vrije Universiteit, Institute of Brain and Behavior Amsterdam and Amsterdam Movement Sciences, Van der Boechorststraat 9, 1081 BT Amsterdam, The Netherlands

[*] Corresponding author: s.m.bruijn@vu.nl

Keywords: nTMS, motor mapping, multiple muscles, cortical representation, active area

Abstract

Background: Single-pulse transcranial magnetic stimulation is a safe and non-invasive tool for investigating cortical representation of muscles in the primary motor cortex. While non-navigated TMS has been successfully applied to simultaneously induce motor-evoked potentials (MEPs) in multiple muscles, a more rigorous assessment of the corresponding cortical representation can greatly benefit from navigated transcranial magnetic stimulation (nTMS).

Objective: We designed a protocol to map the entire precentral gyrus using neural navigation while recording responses of eight muscles simultaneously. Here, we evaluated the feasibility, validity, and reliability of this protocol.

Method: Twenty participants underwent conventional (i.e., muscle-based, grid-constrained) and gyrus-based nTMS mapping. For both protocols, we investigated three different stimulation intensities during two consecutive sessions.

Results: The gyrus-based nTMS mapping was received well by all participants and was less time consuming than the grid-constrained standard. On average, MEP amplitudes, latencies, and centre-of-gravity and size of the active areas largely agreed across protocols supporting validity. Intraclass coefficients between sessions unscored the reliability of our protocol.

Conclusion: We designed an nTMS protocol for the simultaneous mapping multiple muscles on the cortex. The protocol takes only about ten minutes per participant when including as many as eight muscles. Our assessments revealed that the cortical representation of multiple muscles can be determined with high validity and reliability.

Introduction

Single-pulse transcranial magnetic stimulation (TMS) is a non-invasive and painless technique that allows for monitoring neurophysiological alterations of the human motor cortex [1, 2]. When a TMS coil discharges at a suitable intensity, the time-varying magnetic field will induce transient currents causing depolarisation of axons of nerve cells [3]. This will elicit a motor-evoked potential (MEP), which can be recorded in contralateral target muscles. Amplitudes and latencies of these MEPs can reveal the excitability and conduction times of the cortical-spinal tract. Both have been conceived as valid parameters of TMS motor mapping [4]. Neuroscientists and physicians alike utilised TMS motor mapping to evaluate motor cortical plasticity [5], to plan brain tumour surgery [6] and to follow the recovery after stroke [7]. There is ample evidence that the maximum MEP elicited using TMS is closer to the site found using direct cortical stimulation – the gold standard in motor mapping – than that found using magnetoencephalography (MEG) [8] and functional magnetic resonance imaging (fMRI) [9].

In traditional TMS motor mapping procedures, a grid is printed on a cap worn by the subject. This grid serves to manoeuvre the coil and stimulate at the adequate position [10, 11]. Navigated TMS (nTMS) has been developed to substitute the cap, yet the grid-based positions remain. The grid can be displayed on an anatomical scan (typically an MRI) allowing to position the coil according to neuroanatomy. This reduces the error from relative movements of cap and head [12]. That advance has been improved by also accounting for the orientation of the coil, next to the mere positioning [13]. The most recent addition in nTMS systems has been positioning with instantly assessed electromyography (EMG), to display MEP parameters (almost) online on the stimulated sites of the cortex. By this, one can, e.g., a create map of MEP amplitudes for a certain muscle during an experiment (rather than during post-processing) [14].

These advances in technology have led to improvements in experimental protocols. As said, for TMS without neural navigation, a grid on a cap serves to guide coil positioning [10]. In nTMS, the grid can be readily repositioned around the so-called hotspot of a target muscle, i.e. the centre of the area of interest for subsequent assessments, e.g., with different stimulation intensities [15, 16] quantified relative to the muscle-specific resting motor threshold (RMT) [17]. Littmann and co-workers [18] designed an optimal grid by increasing the number of points over the precentral gyrus and reducing the number of points far away from it. Despite these improvements, however, it remains a challenge to move the coil between closely neighbouring points. Van De Ruit, Perenboom, Grey [19] compared grid-based procedures to a protocol in which stimulations were applied at pseudorandom positions. Not only did this alternative approach reduce the time needed to map a single muscle, it turned out to be as reliable as the conventional procedures; see also Cavaleri, Schabrun, Chipchase [20].

Identifying the cortical map of multiple muscles in one experimental run will be difficult – if at all

possible – without neural navigation [10]. Yet, studies exploiting nTMS for that sake are few and far

between – see [21] for an exception where up to four muscles were targeted. The focus on single

muscle mapping arguably stems from the fact that – in most nTMS protocols – stimulation intensities

at a specific percentage of a single muscle's RMT are deemed important. Detecting RMTs for several

muscles is laborious because it often involves offline EMG analyses. Quite recently, however, it has

been shown that the RMT of a small hand muscle (there, the first dorsal interosseous, FDI) may be

similar to the RMT of all upper extremity muscles [16].

All these developments led us to design a new nTMS-based protocol for multiple muscles on the pre-

central gyrus (primary motor cortex). Here, we illustrate its feasibility, validity, and reliability. As will

be shown, our mapping protocol significantly reduces operation times and drastically simplifies pro-

cedures. By the same token, it comes with proper validity and reliability. The protocol can provide

information about the cortical representations of multiple muscles and the degree to which they over-

lap. For this, we also submit a new way to define and measure the so-called active area. Through our

research, we anticipate changing the nTMS paradigm in research and clinic: from grid- to gyrus-based

and from single to multiple muscle mapping.

Materials and methods

Participants

Twenty healthy, right-handed volunteers (average age: 29.55 ± 7.49, eight females) participated in the

study. Prior to the experiment, all participants were screened for contraindications of MRI and TMS

through questionnaires [22]. All of them provided signed informed consent. The Edinburgh Handed-

ness Inventory was used to determine hand dominance [23]. The study had been approved by the

medical ethics committee of Amsterdam University Medical Center (VUmc, 018.213 -

NL65023.029.18).

Materials

Our set-up consisted of three devices: a TMS system, an EMG amplifier, and a neural navigator. Single-

pulse TMS was delivered by a Magstim 200^2 stimulator (Magstim Company Ltd., Whitland, Dyfed, UK)

using a figure-of-eight coil with 70 mm windings. Eight bipolar EMG signals were recorded using a 16-

channel EMG amplifier (Porti, TMSi, Oldenzaal, the Netherlands) and continuously sampled at a rate

of 2 kHz. The EMG recordings were triggered by the TMS to allow for online EMG-assessments using

a custom-made Labview-programme with embedded Matlab functions (designed at our department

using Labview 2016, National Instruments, Austin, TX, and Matlab 2018b, The MathWorks, Natick,

MA). In brief, upon receiving a trigger, peak-to-peak amplitudes and latencies of MEPs were estimated

from all EMG signals during the following 500 ms. These parameters, as well as the original EMG sig-

nals (duration = 500 ms), were sent to the neural navigation software (Neural Navigator, Brain Science

Tools, De Bilt, The Netherlands, www.brainsciencetools.com) for online monitoring and storage. The

neural navigation software also stored the position and orientation of the coil with respect to the head.

Prior to running the TMS protocol, we acquired the participants' anatomical T1-weighted MRI (3 Tesla

Philips Achieva System, Philips, Best, The Netherlands; matrix size 256 × 256 × 211, voxel size

1.0 × 1.0 × 1.0 mm^3, TR/TE 6.40/2.94 ms). The MRI grey matter was segmented using SPM (SPM12,

https://www.fil.ion.ucl.ac.uk/spm/software/spm12/).

We considered the first dorsal interosseous (FDI), abductor digiti minimi (ADM), abductor pollicis

brevis (APB), flexor pollicis brevis (FPB), extensor digitorum communis (EDC), flexor digitorum super-

ficialis (FDS), extensor carpi radialis (ECR) and flexor carpi radialis (FCR) muscles, which were measured

using bipolar electrodes (Blue Sensor N-00-S, Ambu, Ballerup, Denmark), placed after cleaning the skin

with alcohol; cf. Figure 1. The ground electrode was attached to the ulnar styloid process. We moni-

tored and kept the electrode impedance below 5 kΩ. During the experiment, the orientation of the

TMS coil was held 45 degrees to the sagittal plane, tangential to the scalp. By this, we meant to induce

currents in the cortex along the posterior-to-anterior direction. To control the TMS output, we modi-

fied the employed Matlab-toolbox (https://github.com/armanabraham/Rapid2) when adjusting in-

tensities and intervals between stimulations.

Figure 1.   Placement of bipolar electrodes for the first dorsal interosseous (FDI), abductor digiti minimi (ADM), abductor pollicis brevis (APB), flexor pollicis brevis (FPB), extensor digitorum communis (EDC), flexor digitorum super-ficialis (FDS), extensor carpi radialis (ECR) and flexor carpi radialis (FCR) muscles.

Experimental procedures

Participants were comfortably seated in an armchair, relaxing muscles of hands and arms. The exper-

iment consisted of two identical sessions, which were separated by one hour to test for test-retest

reliability of our outcomes – electrodes were kept fixed to minimise placement errors. The interval of

one hour only was set to prevent drying of the conductive electrolyte gel. Each session contained three

parts: (1) Testing the RMTs for FDI, EDC and FCR; there, we stimulated thirty points near the omega-

shaped area of the precentral gyrus to identify the respective hotspots defined as the coordinates for

which the largest peak-to-peak amplitudes in the corresponding EMG signals could be observed. This

served to determine the RMTs for the three target muscles following [24] (i.e. the stimulator output

at which peak-to-peak amplitudes were higher than 50µV in five out of ten stimuli). (2) We conducted

two sets of stimulations to map the representations of all eight muscles using three intensities each,

namely 105% RMT of FDI, EDC and FCR, respectively. For the first set, we adopted the conventional

grid-based method [25] using a square grid (5 cm × 5 cm) with either the FDI-, EDC- or FCR-hotspot as

the centre and applied 80 stimuli. This was immediately followed by a pseudorandom positioning over

the whole gyrus (40 stimuli), yielding a total of 80+40=120 stimuli – in the following we consider the

total of 120 stimuli for our gyrus-based approach while the grid mapping contained only the first 80

stimuli. (3) We analysed the last 40 stimulations and estimated the hotspots of the other five muscles

(ADM, APB, FPB, FDS and ECR) and determined their RMTs.

Offline data processing

During the measurement, peak-to-peak amplitudes and latencies of the MEPs were estimated. The

MEP was defined as the range between the minimum and maximum peak of the EMG signal, and the

latency as the onset of the MEP signal (https://github.com/marlow17/surfaceanalysis). The onset was

defined as the point in time at which the signal exceeded mean ± 1.96×sd of the signal baseline

(100 ms prior to up to the moment of stimulus).

For every muscle and stimulation, we determined whether a MEP was elicited; see *Supplementary*

Material for details. Whenever MEPs were present, the corresponding parameters were included

when computing the mean MEP amplitude and latency, the centre-of-gravity of the stimulation sur-

face ($CoG_{x,y,z}$) and the size of the active area, i.e., the size of that surface. We defined the centre-of-

gravity as follows [26]:

149
$$CoG_x = \frac{\sum MEP_i \cdot X_i}{\sum MEP_i}, \; CoG_y = \frac{\sum MEP_i \cdot Y_i}{\sum MEP_i}, \text{ and } CoG_z = \frac{\sum MEP_i \cdot Z_i}{\sum MEP_i}.$$

150   Here, MEP_i represents the peak-to-peak amplitude for stimulation i at position (X_i, Y_i, Z_i). The defi-

151   nition of the active area is more involved, but we provide more details in the *Supplementary Material*.

152   All the analyses were performed separately for the 3 × 2 × 2 cases: three different muscle-specific in-

153   tensities, for grid- and gyrus-based mapping, and for both sessions. The different steps are illustrated

154   in Figure 2.

Figure 2. The schematic illustrates the procedures of the experiment and data analysis. (A) The gyrus-based method is illustrated by the points of stimulation projected on subjects' pial MRI surface in the neural navigation system in the above picture. The two points highlighted in the lower panel inlet represent the points of the FDI and FCR representation, respectively, corresponding to the motor evoked potentials (MEPs) shown in column (B): top = FDI, bottom = FCR. The MEPs were defined as the peak-to-peak value of the EMG signal under the provision that it exceeded 50µV and the latency was between 15 to 30 ms. (C) The cortical representations of FDI and FCR, i.e., the muscle-specific points shown in column (A) with proper MEPs.

Statistics

All statistical analyses were conducted in Matlab. First, we calculated the average values of RMTs over

all the subjects and sessions of the eight muscles. Possible differences of RMTs across the eight mus-

cles were assessed using a two-way repeated ANOVA with *Muscle* and *Session* as factors, with the

consecutive post-hoc assessment after Bonferroni correction. To evaluate the validity of the gyrus-

based protocol as compared to the grid-based one, we used Bland-Altman plots analysing the ampli-

tude, latency, $CoG_{x,y,z}$, and the size of the active area. We did this only for the parameters of the

muscles that were targeted with the specific intensity, i.e., we compared the grid-based and the gyrus-

based protocol to map the cortical representations of FDI at 105% RMT of FDI, etc. To test the relia-

bility of our gyrus-based protocol, we used intraclass correlation coefficients (ICC) between Session 1

and Session 2. These were calculated for the MEP amplitude and latency, the $CoG_{x,y,z}$, and the size of

the active area. Effects of stimulation intensity were investigated by estimating the ICCs for all MEP

parameters. For the sake of legibility, however, we defined distinct classes of reliability by discretising

the ICC-values as follows: Excellent: 0.8 ≤ ICC; Good: 0.65 ≤ ICC < 0.8; Moderate: 0.5 ≤ ICC < 0.65; Poor:

ICC<0.5 [20]. We also performed a two-way repeated measures ANOVA with the factors of *Intensity*

and *Session* on all the parameters (again with Bonferroni correction for multiple comparisons). Prior

to hypothesis testing, sphericity was verified via Mauchly's test and, if applicable a Greenhouse-

Geisser correction was performed. The significance threshold was set to α = .05.

Results

All $N = 20$ participants completed the experimental procedure without adverse reactions. Overall, of

20 subjects × 8 muscles × 3 intensities × 2 sessions=960 cortical representations, only 1.77% did not

elicit proper MEPs (11 of 320 stimulations in the intensity of 105% RMT of FDI, 5 of 320 stimulations

in the 105% RMT of EDC and 1 of 320 stimulations in the intensity of 105% RMT of FCR; see Table S1

in the *Supplementary Material* for more details). For five subjects we could not detect MEPs for ADM

when using the second intensity in both sessions.

Resting motor thresholds

For all the participants and sessions, the average RMTs of the eight muscles were: FDI (44.90±1.46%),

ADM (47.90±1.64%), APB (46.15±1.45%), FPB (46.78±1.73%), EDC (45.28±1.50%), FDS (47.75±1.51%),

ECR (46.55±1.50%) and FCR (48.00±1.52%). The repeated ANOVA revealed a significant effect of *Mus-*

cle (F(7,133) = 3.63, p = .001). There was no significant effect of *Session* (F(1,19) = 0.85, p = .369) or

interaction between *Muscle* and *Session* (F(7,133) = 1.69, p = .118). Post-hoc pairwise comparisons in-

dicated significant differences between EDC-FCR (p = .020) and FDI-FCR (p = .005) in RMTs.

Validity of the gyrus-based mapping

In the validation study, we performed grid- and gyrus-based protocols to map the cortical represen-

tations of FDI (intensity of 105% RMT of FDI), EDC (intensity of 105% RMT of EDC) and FCR (intensity

of 105% RMT of FCR) in Session 1; cf. Figure 3 and Figure S1 in the *Supplementary Material*. The dif-

ferences in $CoG_{x,y,z}$ between grid- and gyrus-based mapping was ±5mm for most subjects in all three

intensities, without systematic effect for the $CoG_{x,y,z}$ location. The amplitude differences between

the two protocols were usually in the region of ±100μV, while the latency differences were typically

in the range of ±1.5ms for all the subjects. The size of the active area showed larger differences be-

tween grid- and gyrus-based mapping, with the analysis of the slope indicating that the differences in

the size of the active area between the two methods increased when the active area increased (test

for slopes, FDI t(18) = 2.91, p = .010, EDC t(20) = 4.00, p = .0008, and FCR t(20) = 3.89, p = .001); see

Figure S1 in the *Supplementary Material* for more details.

Figure 3. Comparisons of the grid- and the gyrus-based protocols. The panels show the Bland-Altman plots for the parameters of $CoG_{x,y,z}$ for the three intensities. The coloured points represent the difference between the results of the gyrus- and grid-based protocol.

199   *Between session reliability*

200   Table 1 provides an overview of the ICCs of the gyrus-based mapping between Session 1 and 2 of the

201   MEP amplitude and latency, as well as of the $CoG_{x,y,z}$ and the size of the active area for the stimulation

202   intensity based on EDC-RMT. ICC values for the mappings based on the other muscles RMT are shown

203   in Table S2 and Table S3 in the *Supplementary Material*.

Table 1. Intraclass correlation coefficients (ICCs) indicating the reliability of the estimated MEP amplitude and latency and $CoG_{x,y,z}$ and the size of the active area at stimulation intensity of 105% RMT of EDC.

	FDI	ADM	APB	FPB	EDC	FDS	ECR	FCR
Amplitude	0.93	0.83	0.78	0.87	0.84	0.84	0.98	0.77
Latency	0.97	0.91	0.65	0.59	0.80	0.82	0.92	0.84
CoG_x	0.85	0.84	0.86	0.86	0.88	0.86	0.87	0.85
CoG_y	0.70	0.79	0.78	0.82	0.76	0.76	0.74	0.74
CoG_z	0.87	0.88	0.88	0.91	0.89	0.92	0.87	0.88
Active area	0.63	0.64	0.65	0.75	0.43	0.58	0.77	0.48

205   In Figure 4, we illustrate the cortical representations of the eight muscles in Figure 4 based on EDC-

206   RMT. Please note that Figures S2 and S3 in the *Supplementary Materials* show the cortical represen-

207   tations based on FDI-RMT and FCR-RMT.

Figure 4.   Cortical representation of the eight muscles in a representative subject for the intensity of 105% RMT of EDC. The cortical representation from Session 1 is depicted on the left panel, whereas the cortical representation from Session 2 is depicted on the right panel. MEP amplitudes colour-code the active area.

Reliability of gyrus-based mapping at different intensities

The reliability of gyrus-based mapping at different stimulation intensities is summarised in Table 2.

The MEP-responses (across parameters) showed an excellent agreement in intensity 105% RMT of

EDC, and the ICCs of latency performed almost excellent consistent for a stimulation intensity of 105%

RMT of EDC and FCR. The ICCs for in $CoG_{x,y,z}$ at all the intensities were in the range of good to excel-

lent.

Table 2.   The number of muscles in the four levels of ICCs cross the three intensities. ICC was interpreted as follows: <0.5 = poor, 0.5-0.64 = moderate, 0.65-0.79= good and >=0.8 was excellent.

		Excellent	Good	Moderate	Poor
Amp.	105 % RMT of FDI	4	4	0	0
	105 % RMT of EDC	6	2	0	0
	105 % RMT of FCR	3	3	1	1
Latency	105 % RMT of FDI	3	3	1	1
	105 % RMT of EDC	6	1	1	0
	105 % RMT of FCR	7	0	0	1

CoG_x	105 % RMT of FDI	8	0	0	0
	105 % RMT of EDC	8	0	0	0
	105 % RMT of FCR	8	0	0	0
CoG_y	105 % RMT of FDI	8	0	0	0
	105 % RMT of EDC	1	7	0	0
	105 % RMT of FCR	0	8	0	0
CoG_z	105 % RMT of FDI	8	0	0	0
	105 % RMT of EDC	8	0	0	0
	105 % RMT of FCR	8	0	0	0
Active area	105 % RMT of FDI	0	1	6	1
	105 % RMT of EDC	0	3	3	2
	105 % RMT of FCR	0	3	3	2

We did not find any significant effects of *Intensity* or *Session* (nor of their interaction) on $CoG_{x,y,z}$ (see

also Table S4 in the *Supplementary Material*). However, the size of the active area for each muscle

differed significantly across intensities, without significant effects of *Session* or *Intensity × Session* in-

teraction. The post-hoc pairwise comparisons revealed that the active area tested in the intensity of

105% RMT of FCR was markedly larger than in the other intensities; cf. Table 3. For several muscles,

the amplitude and latency varied significantly across the intensities (see Table S4 in the *Supplementary*

Material for details).

Table 3. The statistical results of the size of the active area: the average values, the outcomes of the repeated ANOVA (for the factors of *Intensity*, *Session* and *Intensity×Session*) and the pairwise comparisons in the intensity 105% RMT of FDI ~ 105% RMT of EDC, 105% RMT of FDI ~ 105% RMT of FCR and 105% RMT of EDC ~ 105% RMT of FCR.

	Average of active area (mm³)			Intensity		Session		Intensity ×Session		p-value of pairwise comparisons		
	105% RMT of FDI	105% RMT of EDC	105% RMT of FCR	F(2,34)	p-value	F(1,17)	p-value	F(2,34)	p-value	105% RMT FDI ~ EDC	105% RMT FDI ~ FCR	105% RMT EDC~ FCR
FDI	619.61±65.32	711.47±62.73	966.43±77.85	8.798	**0.0008**	0.181	0.676	0.385	0.684	0.750	**0.006**	**0.023**
ADM	439.54±80.66	495.22±73.36	688.80±86.01	4.735	**0.015**	0.781	0.389	0.057	0.945	1.000	0.061	0.131
APB	576.17±91.58	619.43±68.58	885.56±66.29	6.515	**0.004**	0.011	0.918	0.748	0.481	1.000	**0.024**	**0.027**
FPB	580.83±98.33	661.24±79.00	878.70±69.46	6.177	**0.005**	0.299	0.592	0.287	0.752	0.859	0.859	0.062
EDC	652.80±94.43	694.80±59.15	933.34±89.09	6.391	**0.004**	0.861	0.366	0.039	0.962	1.000	**0.029**	**0.024**
FDS	565.54±83.40	583.56±70.26	834.56±76.11	5.642	**0.008**	4.145	0.058	0.513	0.603	1.000	**0.026**	**0.045**
ECR	572.26±95.14	567.46±73.18	871.06±82.29	8.216	**0.001**	0.365	0.554	0.023	0.978	1.000	**0.009**	**0.005**
FCR	531.05±89.14	530.23±76.39	791.67±78.86	5.201	**0.011**	1.745	0.204	0.518	0.601	1.000	**0.028**	0.062

Discussion

We designed a protocol to investigate the cortical representation of multiple muscles using navigated

single-pulse TMS. Conventional grid-based mapping and our gyrus-based mapping were performed

using three stimulation intensities in two consecutive sessions. We determined RMTs of eight muscles

as well as MEP amplitudes and latencies, the centre-of-gravity and the size of the corresponding active

areas. We found that our new protocol is as valid and as reliable as the commonly applied grid-based

approach but appears much more feasible. Our protocol reduces assessment times and simplifies ex-

perimental procedures. In our experiment, it took only about 10 minutes to map eight muscles simul-
taneously (5 second intervals × 120 points / 60), while for the conventional grid-based approach with
serial muscle assessment, at least 80 minutes would be spent. Experimenting over such a long period
is prone to navigational errors and aggravates participants' fatigue. Moreover, by construction sepa-
rate measurements will not provide reliable insight about overlapping cortical muscle representations.

One of the problems in designing multiple muscles mapping is that the RMT of a single muscle is con-
sidered a reference when setting the stimulation intensity. We found that the RMTs of the hand and
forearm muscles considered are indeed marginally different (EDC-FCR and FDI-FCR). The average
RMTs indicated that the difference between RMTs is small (not more than 3.1% of stimulator output).
Intensities of 105%, 110% to 120% [27] RMT have been widely used in motor mapping [8, 27, 28],
suggesting that the here-observed difference is acceptable if not negligible. Hence, forearm and hand
muscles might be pooled in a group of muscles with "similar RMTs" and may indeed be evaluated at
the same intensity.

The validity testing clearly revealed that our gyrus-based mapping agrees with the grid-based standard,
consistent with a previous study on pseudorandom stimulus positioning [20, 29]. The variation of
$CoG_{x,y,z}$ values between the two protocols was restricted to a range of about -4mm to 4mm. The
differences in size of the active area between protocols are more considerable, and the between-
subjects variability is clearly noteworthy. One explanation for this is that any outlier stimulation site
may strongly affect the estimate of the active area. Moreover, the individual MRIs profoundly differed
in size. While the first calls for more statistical evaluation and (spatial) outlier detection, the latter will
soon be addressed by projecting the individual MRI to, e.g., the MNI template [13].

The reliability of amplitude and latency were excellent or good for all the muscles and all the param-
eters analysed in APB and ECR achieved excellent to good ICCs, indicating the required reliability of
our approach. The ICC values of the size of the active area in FDI, ADM, EDC and FCR were moderate
to poor. This may be related to the factors. First, in some participants, the RMTs of these muscles were
higher than the intensities we set. That is, the intensity of stimulation was too low to activate the
corresponding cortical areas. Second, one must realise that even a figure-of-eight coil comes with a
widespread focus. Hence, TMS activates not isolated but also nearby areas. Yet, the estimated centres-
of-gravity appeared very consistent and should be considered reliable parameters in motor mapping,
in line with previous studies [20, 21].

A gyrus-based protocol with nTMS for multiple muscles allows for not just assessing multiple muscles
in parallel, especially when stepping away from mere "active point" drawing [30] to estimating muscle-

specific active areas in detail. Isolated muscles are not mapped to isolated cortical areas, but one de-
termines the cortical representation of a set of muscles. Results show that these cortical representa-
tion manifest overlaps that diverges across the muscles [31]; cf. Figure 4. This comes particularly to
the fore when analysing cortical surfaces in 3D. There are several methods to determine mapping
areas using nTMS [19, 32] that stand out for their computational ease. However, by ignoring the 3D
representation of surface folds, one runs the risk of missing important information about active areas
in the gyri vis-à-vis the sulci. We used a 3D-alternative (as explained in the *Supplementary Material*)
that overcomes these potential shortcomings. This also provides a welcome visualisation of the active
area on the cortex (for single or multiples muscles), which might serve as a convenient tool. In future
research, we will exploit its capacity to unravel the nature of the overlap region of active areas of
multiple muscles.

We selected three intensities when eliciting MEPs, namely 105% RMT of FDI as a representative of the
hand muscle, EDC as extensor and FCR for flexor muscles. Despite the small variation in RMTs of all
muscles, one may still consider this sub-optimal. Image-based RMT prediction may be an excellent
way to address this issue. This should involve modelling the electric field distributions of TMS, which
may also serve to quantify overlapping cortical representation with even higher precision than cur-
rently possible.

Conclusion

We designed a time-efficient protocol for identifying the cortical representation of multiple muscles
in parallel. Our procedure takes only about ten minutes per subject when including eight muscles. The
results of the active area's centre-of-gravity at three distinct stimulation intensities can be considered
very reliable. All the here-considered outcome parameters confirm the validity of the procedure when
compared to a conventional, grid-based approach with muscle-specific resting motor threshold-scaled
stimulation intensities.

Acknowledgements

The authors are thankful to Moira van Leeuwen, who provided assistance during the measurements.
Sjoerd Bruijn was funded by a grant from the Netherlands Organization for Scientific Research
(016.Vidi.178.014), https://www.nwo.nl/en/.

References

[1]   Barker AT, Jalinous R, Freeston IL. Non-invasive magnetic stimulation of human motor cortex. The Lancet 1985;325(8437):1106-7.

[2]   Schambra H, Sawaki L, Cohen L. Modulation of excitability of human motor cortex (M1) by 1 Hz transcranial magnetic stimulation of the contralateral M1. Clinical Neurophysiology 2003;114(1):130-3.

[3]   Rossini PM, Burke D, Chen R, Cohen L, Daskalakis Z, Di Iorio R, et al. Non-invasive electrical and magnetic stimulation of the brain, spinal cord, roots and peripheral nerves: basic principles and procedures for routine clinical and research application. An updated report from an IFCN Committee. Clinical Neurophysiology 2015;126(6):1071-107.

[4]   Rossini PM, Barker A, Berardelli A, Caramia M, Caruso G, Cracco R, et al. Non-invasive electrical and magnetic stimulation of the brain, spinal cord and roots: basic principles and procedures for routine clinical application. Report of an IFCN committee. Electroencephalography and clinical neurophysiology 1994;91(2):79-92.

[5]   Siebner H, Rothwell J. Transcranial magnetic stimulation: new insights into representational cortical plasticity. Experimental brain research 2003;148(1):1-16.

[6]   Krieg SM, Shiban E, Buchmann N, Gempt J, Foerschler A, Meyer B, et al. Utility of presurgical navigated transcranial magnetic brain stimulation for the resection of tumors in eloquent motor areas. Journal of neurosurgery 2012;116(5):994-1001.

[7]   Mark V, Taub E, Morris D. Neuroplasticity and constraint-induced movement therapy. Europa medicophysica 2006;42(3):269.

[8]   Tarapore PE, Tate MC, Findlay AM, Honma SM, Mizuiri D, Berger MS, et al. Preoperative multimodal motor mapping: a comparison of magnetoencephalography imaging, navigated transcranial magnetic stimulation, and direct cortical stimulation. Journal of neurosurgery 2012;117(2):354-62.

[9]   Forster M-T, Hattingen E, Senft C, Gasser T, Seifert V, Szelényi A. Navigated transcranial magnetic stimulation and functional magnetic resonance imaging: advanced adjuncts in preoperative planning for central region tumors. Neurosurgery 2011;68(5):1317-25.

[10] Melgari J-M, Pasqualetti P, Pauri F, Rossini PM. Muscles in "concert": study of primary motor cortex upper limb functional topography. PloS one 2008;3(8):e3069.

[11] Wassermann EM, McShane LM, Hallett M, Cohen LG. Noninvasive mapping of muscle representations in human motor cortex. Electroencephalography and Clinical Neurophysiology/Evoked Potentials Section 1992;85(1):1-8.

[12] Gugino LD, Romero JR, Aglio L, Titone D, Ramirez M, Pascual-Leone A, et al. Transcranial magnetic stimulation coregistered with MRI: a comparison of a guided versus blind stimulation technique and its effect on evoked compound muscle action potentials. Clinical neurophysiology 2001;112(10):1781-92.

[13] Kraus D, Gharabaghi A. Projecting navigated TMS sites on the gyral anatomy decreases inter-subject variability of cortical motor maps. Brain stimulation 2015;8(4):831-7.

[14] Vink JJ, Petrov PI, Mandija S, Dijkhuizen RM, Neggers SF. Outcome of TMS-based motor mapping depends on TMS current direction. bioRxiv 2018:371997.

[15] Julkunen P, Säisänen L, Danner N, Niskanen E, Hukkanen T, Mervaala E, et al. Comparison of navigated and non-navigated transcranial magnetic stimulation for motor cortex mapping, motor threshold and motor evoked potentials. Neuroimage 2009;44(3):790-5.

[16] Krieg SM, Lioumis P, Mäkelä JP, Wilenius J, Karhu J, Hannula H, et al. Protocol for motor and language mapping by navigated TMS in patients and healthy volunteers; workshop report. Acta neurochirurgica 2017;159(7):1187-95.

[17] Kleim JA, Kleim ED, Cramer SC. Systematic assessment of training-induced changes in corticospinal output to hand using frameless stereotaxic transcranial magnetic stimulation. Nature Protocols 2007;2(7):1675.

[18] Littmann AE, McHenry CL, Shields RK. Variability of motor cortical excitability using a novel mapping pro-
cedure. Journal of neuroscience methods 2013;214(2):137-43.

[19] Van De Ruit M, Perenboom MJ, Grey MJ. TMS brain mapping in less than two minutes. Brain stimulation
2015;8(2):231-9.

[20] Cavaleri R, Schabrun SM, Chipchase LS. The reliability and validity of rapid transcranial magnetic stimulation
mapping. Brain stimulation 2018.

[21] Weiss C, Nettekoven C, Rehme AK, Neuschmelting V, Eisenbeis A, Goldbrunner R, et al. Mapping the hand,
foot and face representations in the primary motor cortex—retest reliability of neuronavigated TMS versus
functional MRI. Neuroimage 2013;66:531-42.

[22] Rossi S, Hallett M, Rossini PM, Pascual-Leone A. Screening questionnaire before TMS: an update. 2011.

[23] Oldfield RC. The assessment and analysis of handedness: the Edinburgh inventory. Neuropsychologia
1971;9(1):97-113.

[24] Groppa S, Oliviero A, Eisen A, Quartarone A, Cohen L, Mall V, et al. A practical guide to diagnostic transcra-
nial magnetic stimulation: report of an IFCN committee. Clinical Neurophysiology 2012;123(5):858-82.

[25] Malcolm M, Triggs W, Light K, Shechtman O, Khandekar G, Rothi LG. Reliability of motor cortex transcranial
magnetic stimulation in four muscle representations. Clinical neurophysiology 2006;117(5):1037-46.

[26] Opitz A, Zafar N, Bockermann V, Rohde V, Paulus W. Validating computationally predicted TMS stimulation
areas using direct electrical stimulation in patients with brain tumors near precentral regions. NeuroImage:
Clinical 2014;4:500-7.

[27] Akiyama T, Ohira T, Kawase T, Kato T. TMS orientation for NIRS-functional motor mapping. Brain topogra-
phy 2006;19(1-2):1-9.

[28] Bohning D, He L, George M, Epstein C. Deconvolution of transcranial magnetic stimulation (TMS) maps.
Journal of neural transmission 2001;108(1):35-52.

[29] Jonker ZD, van der Vliet R, Hauwert CM, Gaiser C, Tulen JH, van der Geest JN, et al. TMS motor mapping:
Comparing the absolute reliability of digital reconstruction methods to the golden standard. Brain stimula-
tion 2019;12(2):309-13.

[30] Seynaeve L, Haeck T, Gramer M, Maes F, De Vleeschouwer S, Van Paesschen W. Optimised preoperative
motor cortex mapping in brain tumors using advanced processing of transcranial magnetic stimulation data.
NeuroImage: Clinical 2019;21:101657.

[31] Mathew J, Kübler A, Bauer R, Gharabaghi A. Probing corticospinal recruitment patterns and functional syn-
ergies with transcranial magnetic stimulation. Frontiers in cellular neuroscience 2016;10:175.

[32] Julkunen P. Methods for estimating cortical motor representation size and location in navigated transcra-
nial magnetic stimulation. Journal of neuroscience methods 2014;232:125-33.

368