Antibody Evolution after SARS-CoV-2 mRNA Vaccination

Alice Cho[1,*], Frauke Muecksch[2,*], Dennis Schaefer-Babajew[1,*], Zijun Wang[1,*], Shlomo Finkin[1,*], Christian Gaebler[1], Victor Ramos[1], Melissa Cipolla[1], Marianna Agudelo[1], Eva Bednarski[2], Justin DaSilva[2], Irina Shimeliovich[1], Juan Dizon[1], Mridushi Daga[1], Katrina Millard[1], Martina Turroja[1], Fabian Schmidt[2], Fengwen Zhang[2], Tarek Ben Tanfous[1], Mila Jankovic[1], Thiago Y. Oliveria[1], Anna Gazumyan[1], Marina Caskey[1], Paul D. Bieniasz[2,3], Theodora Hatziioannou[2], and Michel C. Nussenzweig[1,3].

[1]Laboratory of Molecular Immunology, The Rockefeller University, New York, NY 10065, USA

[2]Laboratory of Retrovirology, The Rockefeller University, New York, NY 10065, USA

[3]Howard Hughes Medical Institute

*equal contribution

Address correspondence to: Paul D. Bieniasz pbieniasz@rockefeller.edu; Marina Caskey
mcaskey@rockefeller.edu; Theodora Hatziioannou thatziio@rockefeller.edu; or Michel C.
Nussenzweig nussen@rockefeller.edu

Summary

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection produces B-cell responses that continue to evolve for at least one year. During that time, memory B cells express increasingly broad and potent antibodies that are resistant to mutations found in variants of concern[1]. As a result, vaccination of coronavirus disease 2019 (COVID-19) convalescent individuals with currently available mRNA vaccines produces high levels of plasma neutralizing activity against all variants tested[1,2]. Here, we examine memory B cell evolution 5 months after vaccination with either Moderna (mRNA-1273) or Pfizer-BioNTech (BNT162b2) mRNA vaccines in a cohort of SARS-CoV-2 naïve individuals. Between prime and boost, memory B cells produce antibodies that evolve increased neutralizing activity, but there is no further increase in potency or breadth thereafter. Instead, memory B cells that emerge 5 months after vaccination of naïve individuals express antibodies that are equivalent to those that dominate the initial response. We conclude that memory antibodies selected over time by natural infection have greater potency and breadth than antibodies elicited by vaccination. These results suggest that boosting vaccinated individuals with currently available mRNA vaccines would produce a quantitative increase in plasma neutralizing activity but not the qualitative advantage against variants obtained by vaccinating convalescent individuals.

Between January 21 and June 23, 2021, we recruited 32 volunteers with no history of prior SARS-CoV-2 infection receiving either Moderna (mRNA-1273; n=8) or Pfizer-BioNTech (BNT162b2; n=24) mRNA vaccines for sequential blood donation. Matched samples were obtained an average of 2.5 and 5 weeks after the prime and boost, respectively, and an additional

set of matched samples an average of 2 and 5 months after the boost. The volunteers ranged in

age from 23-78 years (median 34.5 years), 53% were male and 47% female (for details see

Methods and Supplementary Tables 1 and 2).

Plasma binding and neutralization assays

Plasma IgM, IgG, and IgA responses to SARS-CoV-2 receptor binding domain (RBD) were

measured by enzyme linked immunosorbent assay (ELISA)[3]. As reported by others[2,4-6] there was

a significant increase in IgG reactivity to RBD between prime and boost (p<0.0001, Fig. 1a).

IgM and IgA titers were lower than IgG titers and remained low after the second vaccine dose

(Extended data Fig. 1a and b). The magnitude of the response was inversely correlated with age

after the prime (r=-0.54, p=0.005), but the difference was no longer significant after the boost

(Fig. 1b). Between 2 and 5 months after the boost, anti-RBD titers of all isotypes decreased

significantly. IgG titers decreased by an average of 5.2-fold (range: 2.1- to 10.2-fold) and the

loss of activity was directly correlated to the time after vaccination (p<0.0001, Fig. 1c and d and

Extended data Fig. 1c and d).

Neutralizing activity was measured using HIV-1 pseudotyped with the SARS-CoV-2 spike[1,3,7,8].

Naïve individuals showed variable responses to the initial vaccine dose with a geometric mean

half-maximal neutralizing titer (NT_{50}) of 171 (Fig. 1e and Supplementary Table 2). The

magnitude of the neutralizing responses to the initial vaccine dose in naïve volunteers was

inversely correlated with age (r=-0.39, p=0.05, Fig. 1f). Both binding and neutralizing responses

to the second vaccine dose were correlated to the prime (r=0.46, p=0.02, Extended data. Fig. 1e;

r=0.54, p=0.003, Extended data Fig. 1f) and produced a nearly 15-fold increase in the geometric

mean neutralizing response that was similar in males and females and eliminated the age-related

difference in neutralizing activity (Extended data 2g and Fig. 1f). After the boost naïve vaccinees

had 6.2-fold higher neutralizing titers than a cohort of infected individuals measured 1.3 months

after symptom onset[3] (p<0.0001, Fig. 1e). Neutralizing responses were directly correlated to IgG

anti-RBD titers (r=0.96, p<0.0001, Fig. 1g). Thus, the data obtained from this cohort agree with

prior observations showing a significant increase in plasma neutralizing activity that are

correlated with improved vaccine efficacy in naïve individuals that receive the second dose of

mRNA vaccine[2,6,9,10].

The 15 individuals assayed 5 months after vaccination had an average 4.7-fold decrease in

geometric mean neutralizing activity from their 2-month measurement (p=0.04, Fig. 1h), with a

range of 1.4- to 27-fold (Fig. 1i). Neutralizing activity was inversely correlated with the time

from vaccination (r=-0.75, p<0.0001, Fig. 1j), and directly correlated to IgG anti-RBD binding

titers when assessed 5 months after vaccination (Extended data. Fig. 1h).

We and others showed that the neutralizing responses elicited by mRNA vaccination are more

potent against the original Wuhan Hu-1 strain than for some of the currently circulating variants

of concern[2,11-13]. To confirm these observations, we measured the neutralizing activity of the

paired plasmas from naive individuals 2 and 5 months after the second vaccine dose against

B.1.1.7 (alpha variant), B.1.351 (beta variant), B.1.526 (first isolated in New York City), P.1

(gamma variant) and B.1.617.2 (delta variant). Consistent with previous reports[12,14-16] the

neutralizing activity against the variants was lower than against the original Wuhan Hu-1 strain

(Fig. 1k, Supplementary Table 3). Initial geometric mean neutralizing titers at 2 months against

B.1.351, B.1.1.7, B.1.526, P.1 and B.1.617.2 were 5.7, 1.8, 1.1, 1.4 and 2.7-fold lower than

against Wuhan-Hu respectively (Fig. 1k). In the months following vaccination there was a

decrease in neutralizing activity against all variants that paralleled the drop in activity against

Wuhan Hu-1 (R683G) with geometric mean neutralizing titers for WT, B.1.351, B.1.1.7,

B.1.526, P.1 and B.1.617.2 decreasing by 2.9-, 1.8-, 2.3-, 2.9-, 2.4- and 2.6-fold, respectively

(Fig. 1k and Supplementary Table 3). Nevertheless, vaccine-induced neutralizing activity at 5

months against all variants exceeds Wuhan-Hu plasma neutralization of convalescent individuals

after 6.2 months.

Monoclonal Antibodies

Circulating antibodies produced by plasma cells can prevent infection if present at sufficiently

high concentrations at the time of exposure. In contrast, the memory B cell compartment

contains long lived antigen-specific B cells that mediate rapid recall responses that contribute to

long term protection[17]. To examine the nature of the memory compartment elicited by one or two

mRNA vaccine doses and its evolution after 5 months we used flow cytometry to enumerate B

cells expressing receptors that bind to Wuhan Hu-1 (wild type) and the B.1.351

K417N/E484K/N501Y (KEN) variant RBDs (Fig. 2a and b, and Extended data Fig. 2). Wuhan-

Hu RBD-specific memory B cells developed after the prime in all volunteers examined and their

numbers increased for up to 5 months after vaccination (Fig. 2a). Memory B cells binding to the

B.1.351 RBD were detectable but in lower numbers than wild type RBD-binding B cells in all

samples examined (Fig. 2b). Whereas IgG memory cells increased after the boost, IgM-

expressing memory B cells that made up 23% of the memory compartment after the prime were

nearly absent after boosting (Fig. 2c). Finally, circulating RBD-specific plasmablasts were

readily detected after the prime but were infrequent after the boost (Fig. 2d, and Extended data

Fig. 2d).

The memory compartment continues to evolve up to one year after natural infection with

selective enrichment of cells producing broad and potent neutralizing antibodies[1]. To determine

how the memory compartment evolves after vaccination, we obtained 1524 paired antibody

sequences from 6 individuals sampled at 2.5 weeks after prime and 5 weeks or 5 months after

boost, and an additional 804 paired antibody sequences from 5 individuals sampled after 2- or 5-

months after boost (Fig. 2e and f, Extended Data Fig 3, Supplementary Table 4). As expected

IGHV3-30 and *IGHV3-53* were over-represented after the first and second vaccine dose and

remained over-represented 5 months after vaccination[18-20] (Extended data Fig. 4).

All individuals examined showed expanded clones of memory B cells that expressed closely

related IGHV and IGHL genes (Fig. 2e and f, Extended data Fig. 4). Paired prime and boost

samples showed expanded clones of memory B cells some of which were shared across

plasmablast, IgM and IgG prime, and IgG boost memory cells (Extended data Fig. 3 and 5).

Thus, the cell fate decision controlling the germinal center versus plasmablast decision is not

entirely affinity dependent since cells with the same initial affinity can enter both

compartments[21].

The relative fraction of memory cells found in expanded clones varied between prime and boost

and between individuals (Fig. 2e). However, new clones that develop after the boost represent a

greater fraction (80%) of the total clones than the conserved clones in all individuals (Fig. 2e).

Finally, memory B cells emerging after the boost showed significantly higher levels of somatic

mutations than plasmablasts or memory B cells isolated after the prime (Extended data Fig. 3b).

After 5 months, there was an overall decrease in the percentage of clones in the RBD-binding

memory compartment (Fig. 2g). Nevertheless, clones of memory B cells continued to evolve for

up to 5 months in vaccinated individuals as evidenced by the dominance of newly emerging

clones (86%, Fig. 2e and f) and the significant increase in somatic mutation between the time

points (p<0.0001, Fig. 2h, Extended data Fig. 3c). The number of mutations in antibody genes

was comparable between vaccinated and convalescent individuals after 5 months (Fig. 2h,

Extended data Fig. 3c). In conclusion the memory B cell compartment continues to evolve for up

to 5 months after mRNA vaccination.

Neutralizing Activity of Monoclonal Antibodies

We performed ELISAs to confirm that the antibodies isolated from memory B cells bind to RBD

(Extended data Fig. 6). 403 antibodies were tested by ELISA including: 86 isolated after the first

vaccine dose; 92 isolated after the second vaccine dose; 111 isolated 2 months after the second

vaccine dose and 114 isolated from individuals that had been fully vaccinated 5 months earlier.

Among the 403 antibodies tested 381 (95%) bound to the Wuhan Hu-1 RBD indicating that the

method used to isolate RBD-specific memory B cells was highly efficient (Supplementary Table

5-7). The geometric mean ELISA half-maximal concentration (EC_{50}) of the antibodies obtained

after prime, boost, 2 months and 5 months was 3.9, 2.6, 3.2 and 2.8 ng/ml respectively,

suggestive of no change in binding over time after vaccination (Extended data Fig. 6 and

Supplementary Table 5). Binding among all antibodies did not improve between prime and

boost, or 2 and 5 months (Extended data Fig. 6). However, a slight improvement was observed

after 5 months among clones (Extended data Fig.6).

381 RBD-binding antibodies were tested for neutralizing activity using HIV-1 pseudotyped with

the SARS-CoV-2 spike[3,8]. The geometric mean half-maximal inhibitory concentration (IC_{50}) of

the RBD-specific memory antibodies improved from 380 ng/ml to 170 ng/ml between the first

and second vaccine dose (p=0.005, Fig. 3a). The improvement was reflected in all clones (IC_{50}

370 vs. 182 ng/ml, p=0.03 Fig. 3b), newly arising clones (IC_{50} 430 vs. 165 ng/ml, p=0.02 Fig.

3c), single antibodies (IC_{50} 357 vs. 131 ng/ml, Fig. 3d) and conserved clones (IC_{50} 314 vs. 208

ng/ml, Fig. 3e, Supplementary Table 6). The increase in neutralizing activity between the first

and second vaccine dose was associated with a decrease in the percentage of non-neutralizing

antibodies (defined as IC50 >1000ng/ml) and increased representation of neutralizing antibodies

(p= 0.03, Fig. 3a). In conclusion, memory B cells recruited after the boost account for most of

the improvement in neutralizing activity in this compartment between the 2 vaccine doses. Thus,

in addition to the quantitative improvement in serum neutralizing activity there is a qualitative

improvement in the memory compartment after boosting.

There was no further improvement in neutralizing activity of the monoclonal antibodies obtained

between 2 and 5 months after vaccination (IC_{50} 140 vs. 144 ng/ml, Fig. 3f). This was true when

considering all B cell clones, newly arising clones or single memory B cells separately (Fig. 3g-j,

Supplementary table 7). In contrast, memory antibodies obtained from convalescent individuals

showed improved neutralizing activity between 1.3[3] and 6.2 months[7] with IC_{50} of 171 ng/ml to

116 ng/ml (Fig. 3f), which improved further after 1 year[1]. This improvement was due to

increased neutralizing activity among persisting clones (p=0.003, Fig. 3h).

Affinity, Epitopes and Neutralization Breadth

To examine affinity maturation after vaccination, we performed biolayer interferometry (BLI)

experiments using the Wuhan Hu-1 RBD[3]. 60 randomly selected antibodies were assayed from

the prime-boost cohort, 30 obtained after the prime and 30 after the boost, with similar overall

geometric mean IC_{50}s ranging from <10 ng/ml to >1000 ng/ml (Extended data Fig. 7a). There

was no significant difference in affinity between the antibodies obtained after the prime and

boost, and no correlation between affinity and neutralizing activity (Fig. 4a and Extended Data

Fig. 7 b and c). Similar experiments were performed on 31 and 30 antibodies obtained from the

2- and 5-month time points from vaccinated individuals, respectively. Despite similar levels of

neutralizing activity (Extended data Fig. 7d), there was an average 2.6-fold increase in the

affinity of the antibodies obtained between 2 and 5 months after vaccination (p=0.02, Fig. 4b and

Extended Data Fig. 7e and f).

We also compared the affinities of pairs of antibodies obtained from conserved clones between 2

and 5 months after vaccination. Conserved clones obtained at 2 and 5 months from vaccinated

individuals showed an average 1.9-fold increase in affinity (p=0.03, Fig. 4c). In contrast, a

comparable group of conserved clonal antibodies obtained from convalescent individuals 1.3 and

6.2 months after infection showed an average 24-fold increase in affinity (p=0.002, Fig. 4c).

In addition to the increase in potency, the neutralizing breath of memory antibodies obtained

from conserved clones from convalescent individuals increases with time after infection[1,7,22]. To

determine whether there is a similar increase in breadth with time after vaccination, we selected

20 random antibodies from the prime or boost, with representative levels of activity against the

original Wuhan Hu-1 strain, and measured their neutralization potency against a panel of

psudotypes encoding RBD mutations associated with circulating variants of concern (Extended

data. Fig. 8). There was little change in breadth between prime and boost, with only a small

increase in resilience to K417N and A475V substitutions (Extended data Fig. 8, Supplementary

Table 8).

In addition, we assayed 10 conserved pairs of neutralizing antibodies obtained 2 and 5 months

after vaccination against the same RBD mutant pseudotypes (Fig. 4d and Supplementary Table

9). They were compared to 6 previously reported[22], plus 10 additional pairs of antibodies

obtained from convalescent individuals at 1.3- and 6.2-month time points (Fig. 4e and

Supplementary Table 9). Whereas only 4 of 10 of the vaccine antibodies showed improved

breadth, 15 of the 16 convalescent pairs developed increased breadth. Moreover, only 1 of the 10

vaccine antibody pairs showed improved potency against pseudotypes carrying B.1.617.2 (delta

variant)-specific RBD amino acid substitutions (L452R/T478K), while 11 out of 16 convalescent

antibody pairs showed improved activity against this virus (Fig. 4d and e). We conclude that

antibody evolution differs in convalescent and vaccinated individuals in that there is less affinity

maturation and little increase in breadth between 2 and 5 months after mRNA vaccination.

Circulating antibodies are produced by plasma cells with variable longevity[23-26]. SARS-CoV-2

infection or mRNA vaccination produces an early peak antibody response that decreases by 5-

10-fold after 5 months[7,27-31]. Notably, peak neutralization titres after vaccination exceed those of

COVID-19 recovered individuals. Nevertheless, neutralizing potency against variants is

significantly lower than against Wuhan Hu-1, with up to 5-10-fold reduced activity against the

B.1.351 variant[5,6,12,13,32]. Taken together with the overall decay in neutralizing activity there can

be 1-2 orders of magnitude decrease in serum neutralizing activity after 5 or 6 months against

variants when compared to the peak of neutralizing activity against Wuhan Hu-1. Thus, antibody

mediated protection against variants is expected to wane significantly over a period of months,

consistent with reports of reinfections in convalescent individuals and breakthrough infection by

variants in fully vaccinated individuals[33-35].

In contrast to circulating antibodies, memory B cells are responsible for rapid recall responses[36-

39], and this compartment is relatively stable over the first 6 months after natural infection[7,40] or

mRNA vaccination. In both cases memory B cells continue to evolve for up to 5 months as

evidenced by increasing levels of somatic mutation and emergence of newly expanded clones.

The memory response would be expected to protect individuals that suffer breakthrough

infection from developing serious disease. However, memory B cell evolution differs in

important ways between infection and mRNA vaccination. Both natural infection and mRNA

vaccination produce memory antibodies that evolve increased affinity, but the increase in affinity

is more modest after vaccination. This difference is consistent with the observation that vaccine-

elicited memory antibodies fail to show the increased neutralizing breadth that developed after

natural infection[1,7].

There are innumerable differences between natural infection and mRNA vaccination that could

account for the differences in antibody evolution over time. These include but are not limited to:

1. Route of antigen delivery, respiratory tract vs. intra-muscular injection[41,42]; 2. The physical

nature of the antigen, intact virus vs. S protein[43]; 3. Antigen persistence, weeks in the case of

natural infection[7] vs. hours to days for mRNA[44]. Each of these could impact on B cell evolution

and selection directly, and indirectly through differential T cell recruitment.

The increase in potency and breadth in the memory compartment that develops after natural

infection accounts for the exceptional responses to Wuhan Hu-1 and its variants that

convalescent individuals develop when boosted with mRNA vaccines[1,5]. The expanded memory

B cell compartment in mRNA vaccinees should also produce high titers of neutralizing

antibodies but with decreased breath in comparison to natural infection when vaccinees are

boosted or when they are re-exposed to the virus[45].Thus, boosting vaccinated individuals with

currently available mRNA vaccines will produce strong responses that mirror their initial vaccine

responses to Wuhan-Hu with similarly decreased coverage against variants. Finally, timing a

boost for optimal responses will depend on whether the objective is to prevent viral acquisition

or disease[46]. Given the rapid emergence of variants, in the former, boosting would be needed on

a far shorter time scale than the latter. The optimal timing for boosting to prevent serious disease

will depend on the stability and further evolution of the memory B cell compartment.

Fig. 1: Plasma ELISAs and neutralizing activity.

a, Graph shows area under the curve (AUC, Y-axis) for plasma IgG antibody binding to SARS-CoV-2 RBD after prime and boost for paired samples. (n=26) **b**, Graph shows plasma IgG antibody binding (AUC, Y-axis) plotted against age (X-axis) after prime (black) and boost (orange). **c,** Graph shows AUC (Y-axis) for plasma IgG antibody binding to SARS-CoV-2 RBD for paired samples obtained 2 and 5 months after the boost. (n=15) **d,** Graph shows AUC values from **a**, and **b,** (Y-axis) plotted against time after vaccination (X-axis). **e,** NT50 in convalescent individuals 1.3m after infection[3] and in vaccinated individuals after 1 dose (prime) or 2 doses (boost) of an mRNA vaccine. **f**, NT50 values (Y-axis) vs. age (years X-axis) in individuals receiving 1 dose (prime, black) or two doses (boost, orange) of an mRNA vaccine. **g**, NT50 values (Y-axis) vs. IgG

antibody binding (AUC, X-axis) after boost in individuals receiving two doses of an mRNA

vaccine. **h**, NT50 values in convalescent individuals 1.3m[3] and 6.2m[7] after infection and in

vaccinated individuals 2- [12] and 5-months (m) after receiving 2 doses of an mRNA vaccine. **i,**

NT50 values in vaccinated individuals 2-[12] and 5-months after receiving 2 doses of an mRNA

vaccine. Lines connect paired longitudinal samples from the same individual. **j**, Graph shows

NT50 values (Y-axis) vs. days (X-axis) after boost in individuals receiving two doses of an mRNA

vaccine. **k**, Plasma neutralizing activity against indicated SARS-CoV-2 variants of concern (n=15

paired samples at 2- and 5-months after full vaccination). Refer to Methods for a list of all

substitutions/deletions/insertions in the spike variants. All experiments were performed at least in

duplicate. NT50 values for Moderna mRNA-1273 and Pfizer-BioNTech BNT162b2 in **d, e, h-j**

are shown in black and red, respectively. Red bars and values in **e, h,** and **k** represent geometric

mean NT50 values. Statistical significance in **a, c, i** and **k** was determined by Wilcoxon test, in **b,**

d, f, g and **j** by spearman correlation test and in **e** and **h** by Kruskal-Wallis test with subsequent

Dunn's multiple comparisons.

Fig. 2: Anti-SARS-CoV-2 RBD B cells after vaccination. a-d, Graphs summarizing **a,** the

number of Wuhan-Hu RBD (WT)-specific memory B cells per 10 million B cells for vaccinees

after prime or boost (n=26) or at follow-up of 2-[12] and 5-months after full vaccination (n=9),

including 3 individuals who were sampled at prime, boost, and again 5 months post-vaccination

(illustrated by red dots). **b,** the number of antigen-specific memory B cells cross-reactive with both

WT and K417N/E484K/N501Y (KEN) RBD mutant per 10 million B cells after prime or boost

(n=26), **c,** the frequency of IgM, IgG, or IgA isotype expression by Wuhan-Hu RBD-specific

memory B cells after prime or boost (n=10), and **d,** number of Wuhan-Hu RBD-binding

plasmablasts per 10 million B cells (n=26) after prime or boost. Red numbers indicate geometric

means. Gating strategy is in Extended Data Fig. 2. **e-f,** Pie charts show the distribution of IgG

antibody sequences obtained from memory B cells from 11 individuals after **e,** prime or boost or

5-months, and **f,** 2- or 5- months. The number inside the circle indicates the number of sequences

analyzed for the individual denoted above the circle, with Pfizer vaccinees indicated by (P) and

Moderna by (M). Invidiuals in red text indicate those that were sampled at prime, boost, and again

5 months post-vaccination. Pie slice size is proportional to the number of clonally related

sequences. The black outline and associated numbers indicate the percentage of clonally expanded

sequences detected at each time point. Colored slices indicate persisting clones (same IGHV and

IGLV genes, with highly similar CDR3s) found at more than one timepoint within the same

individual. Grey slices indicate clones unique to the timepoint. White slices indicate repeating

sequences isolated only once per time point. **g,** Graph shows the relative percentage of expanded

clonal sequences at each time point in **e** and **f**. The red numbers indicate the geometric means. **h,**

Number of nucleotide mutations in the IGVH and IGVL combined (also Supplementary Table 4)

in the antibodies illustrated in **e** and **f**, compared to the number of mutations obtained after 1.3[3] or

6.2[7] months after infection (illustrated by grey dots)**.** Red horizontal bars and numbers indicate

mean number of nucleotide mutations at each time point. Statistical significance in **a-d** and **g** was

determined using Wilcoxon matched-pairs signed rank test. Statistical significance in **h** was

determined by Kruskal Wallis test with subsequent Dunn's multiple comparisons.

Fig. 3: Anti-SARS-CoV-2 RBD monoclonal antibodies. a-j, Graphs show anti-SARS-CoV-2 neutralizing activity of monoclonal antibodies measured by a SARS-CoV-2 pseudotype virus neutralization assay using wild-type (Wuhan Hu-1[47]) SARS-CoV-2 pseudovirus[3,8]. **a-e,** Half-maximal inhibitory concentration (IC_{50}) values for all antibodies (**a**), all clones (**b**), non-persisting clones (**c**), singlets (**d**) and shared clones (**e**) isolated after prime or boost. **f-j,** Half-maximal inhibitory concentration (IC_{50}) values for all antibodies (**f**), all clones (**g**), shared clones (**h**), non-persisting clones (**i**) and singlets (**j**) isolated from COVID-19 convalescent individuals 1.3[3] and 6.2[7] months after infection or from vaccinated individuals 2m[12] or 5m after vaccination. Each dot represents one antibody. Pie charts illustrate the fraction of non-neutralizing (IC50 > 1000 ng/ml) antibodies (grey slices), inner circle shows the number of antibodies tested per group. Horizontal bars and red numbers indicate geometric mean values. Statistical significance in **a-d, f, g, i, j** was

337   determined by Mann-Whitney test and in **e** and **h** by Wilcoxon test. Statistical significance for ring

338   plots was determined using Fisher's exact test. All experiments were performed at least twice.

a (Prime/Boost Vaccinated): p = 0.47; geometric means 36.4, 20.0
b (2m/5m Vaccinated): p = 0.02; geometric means 4.9, 1.9
c shared clones — Convalescent 1.3m/6.2m: p = 0.002; geometric means 8.16, 0.34. Vaccinated 2m/5m: p = 0.03; geometric means 3.26, 1.74

d Vaccination persisting clones 2m/5m

	wt	R683G	R346S	K417N	N440K	A475V	E484K R683G	Q493R	N501Y	KEN R683G	L452R E484Q R683G	L452R T478K R683G
C1332	5	4	8	7	6	5	5	6	5	4	4	5
C1207	103	68	121	86	125	19	61	58	26	3	61	61
C1189	113	103	129	6	112	78	5	80	77	73	8	76
C1213	101	101	14	178	164	148	52	132	133	48	100	100
C1344	77	13	77	96	66	100	93	50	82	78	78	83
C1335	48	79	73	54	68	48	86	64	50	61	141	105
C1333	86	99	97	87	89	84	95	109	93	88	92	114
C1334	65	107	173	88	109	82	99	95	66	79	112	111
C1211	115	113	165	109	188	142	122	105	126	125	130	128
C1302	85	21	70	588	64	109	77	97	102	529	78	107

%IC50 of shared clone isolated 2m after vaccination

e Convalescent infection persisting clones 1.3m/6m

	wt	R683G	R346S	K417N	N440K	A475V	E484K R683G	Q493R	N501Y	KEN R683G	L452R E484Q R683G	L452R T478K R683G
C542	2	4	2	1	2	2	1	2	1	13	1	2
C099	3	1	4	2	3	2	1	5	8	0.4	2	1
C513	16	5	19	9	10	4	15	15	15	5	4	2
C573	1	5	100	1	1	1	2	1	2	1	2	4
C043	18	2.9	17	51	13	4	1	0.3	14	2	1	3
C512	10	2	100	2	25	7	0.3	9	53	0.2	1	2
C055	2	1	2	3	2	1	42	1	1	30	1	100
C080	70	69	3	11	4	35	10	68	71	20	1	1
C549	29	6	62	38	46	44	4	1	15	34	0.1	1
C511	83	45	101	34	71	3	24	10	1	0.2	25	23
C057	47	43	38	35	30	30	89	4	22	100	0.1	31
C095	88	19	121	15	44	57	100	0.3	81	100	100	0.2
C502	154	61	169	136	97	124	100	1	126	100	100	1
C572	148	28	96	60	127	138	100	100	401	144	248	184
C051	295	49	525	293	412	241	52	1	419	100	7	34
C052	48	112	37	50	36	35	192	457	27	100	100	41

%IC50 of shared clone isolated 1.3m after infection

Fig. 4: Affinity and Breadth. a-c, Graphs show antibody K_{DS} for Wuhan-Hu RBD measured by BLI. **a,** antibodies isolated from vaccinees after prime (red, n=30) or boost (green, n=30). **b**, same as **a,** but from vaccinees after 2- (red, n=31) and 5-months (green, n=30). **c**, Clonal-paired antibodies isolated from convalescents 1.3[3]- and 6[7]-months after infection (n=15) or vaccinated individuals 2- and 5-months after full vaccination (n=16). Black horizontal bars and numbers indicate geometric mean values. Statistical significance was determined using two-tailed unpaired

t-test (**a, b**) or two tailed Kruskal Wallis test with subsequent Dunn's multiple comparisons (**c**).

BLI traces can be found in Extended Data Fig 7. **d-e,** Heat-maps show inhibitory concentrations

of antibodies isolated 5m after vaccination (**d**) or 6.2 months[7] after infection (**e**) normalized to their

shared clone isolated 2m after vaccination (**d**) or 1.3 months[3] after infection (**e**), expressed as

%IC50, against indicated mutant SARS-CoV-2 pseudoviruses (Supplementary Table 9).

Antibodies with improved (<30%) IC50 compared to their clonal relative isolated at an earlier

timepoint are colored in shades of green with most improved antibodies in darkest green.

Antibodies with worse (>300%) IC50 than their clonal relative isolated at an earlier timepoint are

colored in red with the most worsened antibodies in dark red. Antibodies that did not change their

IC50 by more than ~3-fold are shown in yellow.

METHODS

Study participants.

Participants were healthy volunteers receiving either the Moderna (mRNA-1273) or Pfizer-BioNTech (BNT162b2) mRNA vaccines against SARS-CoV-2 who were recruited for serial blood donations at Rockefeller University Hospital in New York between January 21 and June 23, 2021. Participants indicated as "Prime/Boost" were individuals who were *de novo* recruited for this study, while a subgroup of individuals (indicated as "2m/5m") were from a long-term study cohort[12]. Eligible participants were healthy adults with no history of infection with SARS-CoV-2, as determined by clinical history and confirmed through serology testing, receiving one of the two Moderna (mRNA-1273) or Pfizer-BioNTech (BNT162b2), according to current dosing and interval guidelines. Exclusion criteria included incomplete vaccination status, presence of clinical signs and symptoms suggestive of acute infection with or a positive RT-PCR results for SARS-CoV-2 in saliva, or a positive COVID-19 serology. Seronegativity for COVID-19 was established through the absence of serological activity toward the nucleocapsid protein (N) of SARS-CoV-2. Participants presented to the Rockefeller University Hospital for blood sample collection and were asked to provide details of their vaccination regimen, possible side effects, comorbidities and possible COVID-19 history. All participants provided written informed consent before participation in the study and the study was conducted in accordance with Good Clinical Practice. The study was performed in compliance with all relevant ethical regulations and the protocol (DRO-1006) for studies with human participants was approved by the Institutional Review Board of the Rockefeller University. For detailed participant characteristics see Supplementary Tables 1 and 2.

Blood samples processing and storage.

Peripheral Blood Mononuclear Cells (PBMCs) obtained from samples collected at Rockefeller

University were purified as previously reported by gradient centrifugation and stored in liquid

nitrogen in the presence of FCS and DMSO[3,7]. Heparinized plasma and serum samples were

aliquoted and stored at -20°C or less. Prior to experiments, aliquots of plasma samples were heat-

inactivated (56°C for 1 hour) and then stored at 4°C.

ELISAs

ELISAs[48,49] to evaluate antibodies binding to SARS-CoV-2 RBD were performed by coating of

high-binding 96-half-well plates (Corning 3690) with 50 μl per well of a 1μg/ml protein solution

in PBS overnight at 4 °C. Plates were washed 6 times with washing buffer (1× PBS with 0.05%

Tween-20 (Sigma-Aldrich)) and incubated with 170 μl per well blocking buffer (1× PBS with

2% BSA and 0.05% Tween-20 (Sigma)) for 1 h at room temperature. Immediately after

blocking, monoclonal antibodies or plasma samples were added in PBS and incubated for 1 h at

room temperature. Plasma samples were assayed at a 1:66 starting dilution and 10 additional

threefold serial dilutions. Monoclonal antibodies were tested at 10 μg/ml starting concentration

and 10 additional fourfold serial dilutions. Plates were washed 6 times with washing buffer and

then incubated with anti-human IgG, IgM or IgA secondary antibody conjugated to horseradish

peroxidase (HRP) (Jackson Immuno Research 109-036-088 109-035-129 and Sigma A0295) in

blocking buffer at a 1:5,000 dilution (IgM and IgG) or 1:3,000 dilution (IgA). Plates were

developed by addition of the HRP substrate, TMB (ThermoFisher) for 10 min (plasma samples)

or 4 minutes (monoclonal antibodies). The developing reaction was stopped by adding 50 μl of 1

M H_2SO_4 and absorbance was measured at 450 nm with an ELISA microplate reader (FluoStar

Omega, BMG Labtech) with Omega and Omega MARS software for analysis. For plasma

samples, a positive control (plasma from participant COV72, diluted 66.6-fold and ten additional

threefold serial dilutions in PBS) was added to every assay plate for validation. The average of

its signal was used for normalization of all the other values on the same plate with Excel

software before calculating the area under the curve using Prism V9.1(GraphPad). For

monoclonal antibodies, the EC50 was determined using four-parameter nonlinear regression

(GraphPad Prism V9.1). EC50s above 2000 ng/mL were considered non-binders.

Proteins

Mammalian expression vector encoding the RBD of SARS-CoV-2 (GenBank MN985325.1; S

protein residues 319-539) was previously described[50].

SARS-CoV-2 pseudotyped reporter virus

A panel of plasmids expressing RBD-mutant SARS-CoV-2 spike proteins in the context of

pSARS-CoV-2-S $_{\Delta 19}$ has been described[12,22,51]. Variant pseudoviruses resembling variants of

concern B.1.1.7 (first isolated in the UK), B.1.351 (first isolated in South-Africa), B.1.526 (first

isolated in New York City), P.1 (first isolated in Brazil) and B.1.617.2 (first isolated in India)

were generated by introduction of substitutions using synthetic gene fragments (IDT) or overlap

extension PCR mediated mutagenesis and Gibson assembly. Specifically, the variant-specific

deletions and substitutions introduced were:

B.1.1.7: ΔH69/V70, ΔY144, N501Y, A470D, D614G, P681H, T761I, S982A, D118H

B.1.351: D80A, D215G, L242H, R246I, K417N, E484K, N501Y, D614G, A701V

B.1.526: L5F, T95I, D253G, E484K, D614G, A701V.

P.1: L18F, T20N, P26S, D138Y, R190S, K417T, E484K, N501Y, D614G, H655Y, T1027I,

V1167F

B.1.617.2: T19R, Δ156-158, L452R, T478K, D614G, P681R, D950N

The E484K, K417N/E484K/N501Y (KEN), L452R/E484Q and L452R/T478K substitution, as

well as the deletions/substitutions corresponding to variants of concern listed above were

incorporated into a spike protein that also includes the R683G substitution, which disrupts the

furin cleaveage site and increases particle infectivity. Neutralizing activity against mutant

pseudoviruses were compared to a wildtype SARS-CoV-2 spike sequence (NC_045512),

carrying R683G where appropriate.

SARS-CoV-2 pseudotyped particles were generated as previously described[3,8]. Briefly, 293T

cells were transfected with pNL4-3ΔEnv-nanoluc and pSARS-CoV-2-S$_{\Delta 19}$, particles were

harvested 48 hpt, filtered and stored at -80°C.

Pseudotyped virus neutralization assay

Fourfold serially diluted plasma from COVID-19-convalescent individuals or monoclonal

antibodies were incubated with SARS-CoV-2 pseudotyped virus for 1 h at 37 °C. The mixture

was subsequently incubated with 293T$_{Ace2}$ cells[3] (for all WT neutralization assays) or

HT1080Ace2 cl14 (for all mutant panels and variant neutralization assays) cells[12] for 48h after

which cells were washed with PBS and lysed with Luciferase Cell Culture Lysis 5× reagent

(Promega). Nanoluc Luciferase activity in lysates was measured using the Nano-Glo Luciferase

Assay System (Promega) with the Glomax Navigator (Promega). The relative luminescence units

were normalized to those derived from cells infected with SARS-CoV-2 pseudotyped virus in the

absence of plasma or monoclonal antibodies. The half-maximal neutralization titers for plasma

(NT_{50}) or half-maximal and 90% inhibitory concentrations for monoclonal antibodies (IC_{50} and

IC_{90}) were determined using four-parameter nonlinear regression (least squares regression

method without weighting; constraints: top=1, bottom=0) (GraphPad Prism).

Biotinylation of viral protein for use in flow cytometry

Purified and Avi-tagged SARS-CoV-2 RBD or SARS-CoV-2 RBD KEN mutant (K417N,

E484K, N501Y) was biotinylated using the Biotin-Protein Ligase-BIRA kit according to

manufacturer's instructions (Avidity) as described before[3]. Ovalbumin (Sigma, A5503-1G) was

biotinylated using the EZ-Link Sulfo-NHS-LC-Biotinylation kit according to the manufacturer's

instructions (Thermo Scientific). Biotinylated ovalbumin was conjugated to streptavidin-BV711

(BD biosciences, 563262) and RBD to streptavidin-PE (BD Biosciences, 554061) and

streptavidin-AF647 (Biolegend, 405237)[3].

Flow cytometry and single cell sorting

Single-cell sorting by flow cytometry was described previously[3]. Briefly, peripheral blood

mononuclear cells were enriched for B cells by negative selection using a pan-B-cell isolation kit

according to the manufacturer's instructions (Miltenyi Biotec, 130-101-638). The enriched B

cells were incubated in FACS buffer ($1\times$ PBS, 2% FCS, 1 mM EDTA) with the following anti-

human antibodies (all at 1:200 dilution): anti-CD20-PECy7 (BD Biosciences, 335793), anti-

CD3-APC-eFluro 780 (Invitrogen, 47-0037-41), anti-CD8-APC-eFluor 780 (Invitrogen, 47-

0086-42), anti-CD16-APC-eFluor 780 (Invitrogen, 47-0168-41), anti-CD14-APC-eFluor 780

(Invitrogen, 47-0149-42), as well as Zombie NIR (BioLegend, 423105) and fluorophore-labelled

RBD and ovalbumin (Ova) for 30 min on ice. Single CD3-CD8-CD14-CD16−CD20+Ova−RBD-

PE+RBD-AF647+ B cells were sorted into individual wells of 96-well plates containing 4 μl of

lysis buffer (0.5× PBS, 10 mM DTT, 3,000 units/ml RNasin Ribonuclease Inhibitors (Promega,

N2615) per well using a FACS Aria III and FACSDiva software (Becton Dickinson) for

acquisition and FlowJo for analysis. The sorted cells were frozen on dry ice, and then stored at

−80 °C or immediately used for subsequent RNA reverse transcription. For plasmablast single-

cell sorting, in addition to above antibodies, B cells were also stained with anti-CD19-BV605

(Biolegend, 302244), and single CD3-CD8-CD14-CD16-CD19+CD20-Ova-RBD-PE+RBD-

AF647+ plasmablasts were sorted as described above. For B cell phenotype analysis, in addition

to above antibodies, B cells were also stained with following anti-human antibodies: anti-IgD-

BV421 (Biolegend, 348226), anti-CD27-FITC (BD biosciences, 555440), anti-CD19-BV605

(Biolegend, 302244), anti-CD71- PerCP-Cy5.5 (Biolegend, 334114), anti- IgG-PECF594 (BD

biosciences, 562538), anti-IgM-AF700 (Biolegend, 314538), anti-IgA-Viogreen (Miltenyi

Biotec, 130-113-481).

Antibody sequencing, cloning and expression

Antibodies were identified and sequenced as described previously[3,52]. In brief, RNA from single

cells was reverse-transcribed (SuperScript III Reverse Transcriptase, Invitrogen, 18080-044) and

the cDNA was stored at −20 °C or used for subsequent amplification of the variable IGH, IGL

and IGK genes by nested PCR and Sanger sequencing. Sequence analysis was performed using

MacVector. Amplicons from the first PCR reaction were used as templates for sequence- and

ligation-independent cloning into antibody expression vectors. Recombinant monoclonal

antibodies were produced and purified as previously described[3].

Biolayer interferometry

Biolayer interferometry assays were performed as previously described[3]. Briefly, we used the

Octet Red instrument (ForteBio) at 30 °C with shaking at 1,000 r.p.m. Affinity measurement of

anti-SARS-CoV-2 IgGs binding were corrected by subtracting the signal obtained from traces

performed with IgGs in the absence of WT RBD. The kinetic analysis using protein A biosensor

(ForteBio 18-5010) was performed as follows: (1) baseline: 60sec immersion in buffer. (2)

loading: 200sec immersion in a solution with IgGs 10 μg/ml. (3) baseline: 200sec immersion in

buffer. (4) Association: 300sec immersion in solution with WT RBD at 20, 10 or 5 μg/ml (5)

dissociation: 600sec immersion in buffer. Curve fitting was performed using a fast 1:1 binding

model and the Data analysis software (ForteBio). Mean KD values were determined by

averaging all binding curves that matched the theoretical fit with an R^2 value ≥ 0.8.

Computational analyses of antibody sequences

Antibody sequences were trimmed based on quality and annotated using Igblastn v.1.14. with

IMGT domain delineation system. Annotation was performed systematically using Change-O

toolkit v.0.4.540[53]. Heavy and light chains derived from the same cell were paired, and

clonotypes were assigned based on their V and J genes using in-house R and Perl scripts. All

scripts and the data used to process antibody sequences are publicly available on GitHub

(https://github.com/stratust/igpipeline/tree/igpipeline2_timepoint_v2).

The frequency distributions of human V genes in anti-SARS-CoV-2 antibodies from this study

was compared to 131,284,220 IgH and IgL sequences generated by[54] and downloaded from cAb-

Rep[55], a database of human shared BCR clonotypes available at https://cab-

rep.c2b2.columbia.edu/. Based on the 112 distinct V genes that make up the 7936 analyzed

sequences from Ig repertoire of the 11 participants present in this study, we selected the IgH and

IgL sequences from the database that are partially coded by the same V genes and counted them

according to the constant region. The frequencies shown in Extended Data Fig. 4 are relative to

the source and isotype analyzed. We used the two-sided binomial test to check whether the

number of sequences belonging to a specific IgHV or IgLV gene in the repertoire is different

according to the frequency of the same IgV gene in the database. Adjusted p-values were

calculated using the false discovery rate (FDR) correction. Significant differences are denoted

with stars.

Nucleotide somatic hypermutation and CDR3 length were determined using in-house R and Perl

scripts. For somatic hypermutations, IGHV and IGLV nucleotide sequences were aligned against

their closest germlines using Igblastn and the number of differences were considered nucleotide

mutations. The average number of mutations for V genes was calculated by dividing the sum of

all nucleotide mutations across all participants by the number of sequences used for the analysis.

Data availability statement: Data are provided in Supplementary Tables 1-9. The raw

sequencing data and computer scripts associated with Figure 2 and Extended Data Fig. 3 have

been deposited at Github (https://github.com/stratust/igpipeline/tree/igpipeline2_timepoint_v2).

This study also uses data from "A Public Database of Memory and Naive B-Cell Receptor

Sequences" (https://doi.org/10.5061/dryad.35ks2), PDB (6VYB and 6NB6) and from "High

frequency of shared clonotypes in human B cell receptor repertoires"

(https://doi.org/10.1038/s41586-019-0934-8).

Code availability statement: Computer code to process the antibody sequences is available at

GitHub (https://github.com/stratust/igpipeline/tree/igpipeline2_timepoint_v2).

548
549

Data presentation

Figures arranged in Adobe Illustrator 2020.

Competing interests: The Rockefeller University has filed a provisional patent application in

connection with this work on which M.C.N.is an inventor (US patent 63/021,387). The patent

has been licensed by Rockefeller University to Bristol Meyers Squib.

Acknowledgments: We thank all study participants who devoted time to our research; The

Rockefeller University Hospital nursing staff and Clinical Research Support Office and nursing

staff. Mayu Okawa Frank, Marissa Bergh, and Robert B. Darnell for SARS-CoV-2 saliva PCR

testing. Charles M. Rice, and all members of the M.C.N. laboratory for helpful discussions, Maša

Jankovic for laboratory support, and Kristie Gordon for technical assistance with cell-sorting

experiments. This work was support by NIH grant P01-AI138398-S1 (M.C.N.) and

2U19AI111825 (M.C.N.). R37-AI64003 to P.D.B.; R01AI78788 to T.H. F.M. is supported by

the Bulgari Women & Science Fellowship in COVID-19 Research. C.G. was supported by the

Robert S. Wennett Post-Doctoral Fellowship, in part by the National Center for Advancing

Translational Sciences (National Institutes of Health Clinical and Translational Science Award

program, grant UL1 TR001866), and by the Shapiro-Silverberg Fund for the Advancement of

Translational Research. P.D.B. and   M.C.N. are Howard Hughes Medical Institute Investigators.

Author Contributions: P.D.B., T.H., and M.C.N. conceived, designed and analyzed

the experiments. M. Caskey and C.G. designed clinical protocols. A.C, F.M., D.S.B., Z.W., S.F.

M.A., E.B., J.D.S., I.S., J.D. F.S., F.Z., and T.B.T. carried out experiments. A.G. and M. Cipolla

produced antibodies. D.S.B., M.D., M.T., K.G.M., C.G. and M. Caskey recruited participants,

executed clinical protocols. T.Y.O. and V.R. performed bioinformatic analysis. A.C., F.M,

D.S.B., Z.W., S.F., and M.C.N. wrote the manuscript with input from all co-authors.

  EXTENDED FIGURES

Extended Data Fig 1: Plasma ELISA and neutralization.

a,b, Graph shows area under the curve (AUC, Y-axis) for plasma IgM (**a**) or IgA (**b**) antibody

binding to SARS-CoV-2 RBD after prime and boost for paired samples. **c,d,** Graph shows area

under the curve (AUC, Y-axis) for plasma IgM (**c**) or IgA (**d**) antibody binding to SARS-CoV-2

RBD of paired samples obtained 2 and 5 months after the boost. **e,f,** IgG antibody binding (AUC,

X-axis) after prime vs. IgG antibody binding (AUC, X-axis) after boost (**e**) and NT50 values after

prime vs. NT50 values after boost (**f**) in individuals receiving two doses of an mRNA vaccine. **g,**

NT50 values after prime and boost in females and males receiving 2 doses of an mRNA vaccine.

h, NT50 values (Y-axis) vs. IgG antibody binding (AUC, X-axis) 5 months after boost in

individuals receiving two doses of an mRNA vaccine. All experiments were performed at least in

duplicate. Red values or bar in **a-d** and **g** represent geometric mean values. Statistical significance

in **a-d** was determined by Wilcoxon test, in **e, f, and h** by spearman correlation test and in **g** by

Kruskall-Wallis test with subsequent Dunn's multiple comparisons.

Extended Data Fig. 2: Flow Cytometry. a, Gating strategy for phenotyping. Gating was on singlets that were CD19$^+$ or CD20$^+$ and CD3-CD8-CD16-Ova-. Anti-IgG, IgM, IgA, IgD, CD71 and CD27 antibodies were used for B cell phenotype analysis. Antigen-specific cells were detected based on binding to RBD WT-PE$^+$ and RBD WT/KEN-AF647$^+$. **b-d,** Flow cytometry plots showing the frequency of **b,** RBD WT-binding memory B cells, and **c,** RBD-binding memory B cells cross-reactive with WT and K417N/E484K/N501Y mutant RBD and **d,** RBD-binding plasmablasts, in 10 selected vaccinees after prime or boost. **e,** Gating strategy for single-cell sorting for CD20+ memory B cells (top panel) or CD19+CD20- plasmablasts (bottom panel) which were double positive for RBD-PE and RBD-AF647. **f-g,** Representative flow cytometry plots showing dual AlexaFluor-647-RBD and PE-RBD-binding, single-cell sorted B cells from **f,** 6 individuals after prime or boost or 5 months post-vaccination and **g,** 5 individuals from 2- or 5-months post-vaccination. Percentage of RBD-specific B cells is indicated. Invidiuals in red text indicate those that were sampled at prime, boost, and again 5 months post-vaccination.

Extended Data Fig 3: anti-SARS-CoV-2 RBD-specific plasmablast and MBC responses after vaccination. a, Pie charts show the distribution of antibody sequences from 6 individuals after prime (upper panel) or boost (lower panel). Sequences derived from IgG plasmablast, IgM MBC, and IgG MBC compartments were analyzed after prime, while only IgG MBCs were analyzed after boost, as indicated to the left of the plots. The number inside the circle indicates the number

of sequences analyzed for the individual denoted above the circle. Pie slice size is proportional to

the number of clonally related sequences. The black outline indicates the frequency of clonally

expanded sequences detected in each patient. Colored slices indicate persisting clones (same IGHV

and IGLV genes, with highly similar CDR3s) found in multiple compartments and/or timepoints

within the same patient. Grey slices indicate clones unique to the compartment. White indicates

sequences isolated once. **b-c,** Number of somatic nucleotide mutations in the IGVH and IGVL,

separately, in antibodies detected in **b,** different B cell compartments after prime or boost and **c,**

2^{12} or 5 months post-vaccination compared to convalescent infected individuals after 1.3[3] and 6.2[7]

months post-infection (also Supplementary Table 4). Red horizontal bars and numbers indicate

mean number of nucleotide mutations in each compartment at each time point. Statistical

significance was determined using a Kruskal Wallis test with subsequent Dunn's multiple

comparisons.

Extended Data Fig. 4: Frequency distribution of human V genes. a, Circos plot depicting

relationship between antibodies that share V and J gene usage in both IgH and IgL when comparing

all prime/boost IgG MBC sequences. Purple, green, and grey lines connect related clones, clones

and singlets, and singlets to each other, respectively. **b,** Graph shows relative abundance of human

heavy chain IGVH (top), light chain IGVK (middle) or IGVL (bottom) genes comparing Sequence

Read Archive accession SRP010970 (orange), and IgG MBCs after prime (blue) or boost (green)

doses of vaccination. Statistical significance was determined by two-sided binomial test. * =

$p \leq 0.05$, ** = $p \leq 0.01$, *** = $p \leq 0.001$, **** = $p \leq 0.0001$. Color of stars indicates: black - comparing

Database versus Prime; blue - comparing Database versus Boost; red - comparing Prime versus

Boost. **c,** Circos plot depicting relationship between antibodies that share V and J gene usage in

both IgH and IgL when comparing 2 mo/5 mo IgG MBC sequences. Purple, green, and grey lines

connect related clones, clones and singlets, and singlets to each other, respectively. **d,** Graph shows

relative abundance of human heavy chain IGVH (top), light chain IGVK (middle) or IGVL

(bottom) genes comparing Sequence Read Archive accession SRP010970 (orange), and IgG

MBCs after 2 months (blue) or 5 months (green) post-vaccination. Statistical significance was

determined by two-sided binomial test. * = $p \leq 0.05$, ** = $p \leq 0.01$, *** = $p \leq 0.001$, **** = $p \leq 0.0001$.

Color of stars indicates: black - comparing Database versus 2 months; blue - comparing Database

versus 5 months; red - comparing 2 months versus 5 months.

Extended Data Fig. 5: Somatic hypermutation of anti-SARS-CoV-2 RBD antibody clones after prime or boost. Clonal evolution of RBD-binding B cells from 3 individuals for which plasmablasts, IgM memory B cells, and IgG memory B cells were analyzed after prime, and IgG memory B cells were analyzed after boost (as described in Extended Data Fig. 3). The number of

653   somatic nucleotide mutations found in shared clonal families found in at least 2 different

654   compartments is graphed to the right of each donut plot. Color of dot plots match the color of pie

655   slices within the donut plot, which indicate persisting clones. nd – clone was Not Detected in the

656   indicated compartment. Black horizontal line indicates median number of SHM.

Extended Data Fig. 6: Anti-SARS-CoV-2 RBD monoclonal antibodies ELISAs. a-j, Graphs

show anti-SARS-CoV-2 binding activity of monoclonal antibodies measured by ELISA against

RBD. **a-e,** ELISA half-maximal concentration (EC_{50}) values for all antibodies (**a**), all clones (**b**),

non-persisting clones (**c**), singlets (**d**) and shared clones (**e**) isolated after prime or boost. **f-j,**

ELISA half-maximal concentration (EC_{50}) values for all antibodies (**f**), all clones (**g**), shared clones

(**h**), non-persisting clones (**i**) and singlets (**j**) isolated from COVID-19 convalescent individuals

1.3[3] and 6.2[7] months after infection or from vaccinated individuals 2m[12] or 5m after vaccination.

Each dot represents one antibody. Red horizontal bars and numbers indicate geometric mean

values. Statistical significance in **a-d, f, g, i, j** was determined by Mann-Whitney test and in **e** and

h by Wilcoxon test. All experiments were performed at least twice.

Extended Data Fig. 7. Affinity. Biolayer interferometry measurements. **a**, IC50 values for randomly selected antibodies isolated from vaccinees after prime (n=30) of boost (n=30). Red horizontal lines and numbers indicate geometric mean. **b,** Graphs depict affinity measurements of antibodies described in **a. c,** Graphs show affinities (Y axis) plotted against neutralization activity (X axis) for antibody isolated after vaccination. **d-f,** same as in **a-c,** but showing monoclonal antibodies isolated from follow-up of 2 and 5 months post-vaccination. **d,** IC50 values for randomly selected antibodies isolated from vaccinees after 2- (n=31) or 5- (n=30) months post-vaccination. **e**, Graphs depicting affinity measurements of antibodies described in **d. f,** Graph shows affinity (Y axis) plotted against neutralization activity (X axis) for antibody isolated after vaccination.

a

	wt	R683G	R346S	K417N	N440K	A475V	E484K R683G	Q493R	N501Y	KEN R683G	L452R E484Q R683G
C2159	12.2	6.2	9.1	3.3	11.2	8.8	627.1	17.1	20.6	212.0	36.9
C2029	25.1	7.6	>1000	5.9	23.0	19.5	1.8	14.6	57.7	1.6	>1000
C2139	46.0	25.4	33.0	>1000	48.7	>1000	153.6	>1000	>1000	>1000	105.1
C2033	46.2	25.8	32.0	>1000	41.4	868.4	79.5	183.4	668.2	>1000	52.1
C2209	52.6	26.7	40.7	12.5	48.2	214.0	107.3	341.9	214.9	58.8	44.0
C2020	65.8	37.8	164.0	>1000	53.5	71.5	131.3	>1000	119.0	>1000	83.7
C2221	69.3	26.4	47.6	>1000	58.8	>1000	355.2	>1000	46.3	>1000	124.0
C2019	88.5	96.3	>1000	25.5	90.8	70.1	726.4	92.3	125.5	380.3	>1000
C2110	118.4	106.6	92.6	>1000	109.9	797.4	254.9	198.0	301.5	>1000	218.3
C2018	118.9	37.2	55.4	33.4	110.8	98.0	284.5	123.8	135.8	140.0	>1000
C2022	153.4	61.6	114.5	>1000	130.5	>1000	247.7	358.7	162.8	>1000	139.2
C2113	348.4	127.8	242.4	166.8	267.1	347.9	>1000	922.3	339.4	>1000	>1000
C2149	376.8	178.6	259.3	>1000	331.2	>1000	724.8	>1000	>1000	>1000	608.0
C2026	433.1	25.6	258.1	400.1	350.5	925.7	188.0	368.0	387.1	190.1	147.3
C2150	591.0	57.5	672.9	496.9	413.7	783.8	240.8	406.4	543.2	202.1	199.1
C2013	593.3	204.6	391.0	>1000	484.3	117.9	>1000	>1000	>1000	>1000	>1000
C2185	670.6	116.1	440.0	239.1	494.6	818.2	412.8	759.7	485.4	246.7	251.1
C2004	722.5	117.4	529.5	521.1	468.7	>1000	400.1	496.2	928.0	318.2	345.1
C2140	840.9	124.4	706.9	839.5	778.2	>1000	648.7	866.7	815.5	481.7	497.8
C2109	1000.0	198.7	572.6	825.0	336.9	>1000	960.0	762.2	620.2	691.7	464.7

IC50 (ng/ml)

b

	wt	R683G	R346S	K417N	N440K	A475V	E484K R683G	Q493R	N501Y	KEN R683G	L452R E484Q R683G
C2039	1.9	0.5	1.0	0.7	1.2	1.0	>1000	1.4	2.3	>1000	>1000
C2237	6.7	0.7	3.8	2.3	4.7	3.8	342.4	9.1	4.5	815.8	>1000
C2049	10.0	5.2	7.1	319.9	9.6	65.5	10.9	17.5	12.8	>1000	7.2
C2065	11.6	9.8	>1000	4.0	11.1	6.4	387.7	9.6	11.1	123.4	>1000
C2319	13.3	6.3	8.7	>1000	10.2	131.2	7.7	28.4	297.3	>1000	8.5
C2175	17.6	4.9	12.1	5.1	14.2	8.0	506.1	>1000	17.5	347.0	23.5
C2219	20.7	9.8	13.0	5.2	19.4	35.1	>1000	369.6	13.7	>1000	>1000
C2227	48.3	28.5	94.8	20.6	45.0	36.8	>1000	8.7	>1000	>1000	14.3
C2047	49.1	40.4	145.2	>1000	48.8	53.2	168.6	>1000	96.8	>1000	123.4
C2045	52.0	41.2	46.4	>1000	61.1	375.8	62.9	>1000	60.3	>1000	76.4
C2188	90.8	45.7	64.3	>1000	56.6	743.4	134.8	>1000	>1000	>1000	119.6
C2037	148.1	74.1	53.1	32.7	88.0	103.4	378.3	147.5	157.4	246.4	>1000
C2228	178.4	145.9	124.1	70.2	132.2	>1000	770.5	>1000	197.9	886.3	785.3
C2167	200.4	140.1	156.4	13.6	144.2	143.0	287.9	233.5	183.3	28.4	243.6
C2318	351.9	126.5	262.1	113.7	241.7	335.9	286.4	311.0	477.8	244.8	231.4
C2210	366.2	145.7	236.9	188.6	270.5	297.1	382.6	333.2	276.1	381.7	363.0
C2317	429.2	549.9	>1000	105.5	296.6	282.1	>1000	305.6	387.5	>1000	>1000
C2172	451.6	246.0	324.0	214.0	257.3	>1000	363.5	>1000	486.9	>1000	199.7
C2070	584.0	532.7	856.2	260.7	529.0	709.6	884.3	629.7	802.5	838.8	578.2
C2321	843.9	254.3	>1000	648.9	>1000	627.1	400.1	316.5	693.4	445.2	>1000

IC50 (ng/ml)

Extended data Fig. 8: Breadth of anti-SARS-CoV-2 RBD antibodies elicited after prime and boost vaccination. **a-b**, IC_{50} values for n=40 neutralizing antibodies isolated after prime **(a)** or boost **(b)** against indicated mutant SARS-CoV-2 pseudoviruses. Color gradient indicates IC_{50} values ranging from 0 (white) to 1000 ng/ml (red).

REFERENCES

Wang, Z. *et al.* Naturally enhanced neutralizing breadth against SARS-CoV-2 one year after infection. *Nature*, doi:10.1038/s41586-021-03696-9 (2021).

Goel, R. R. *et al.* Distinct antibody and memory B cell responses in SARS-CoV-2 naive and recovered individuals following mRNA vaccination. *Sci Immunol* **6**, doi:10.1126/sciimmunol.abi6950 (2021).

Robbiani, D. F. *et al.* Convergent antibody responses to SARS-CoV-2 in convalescent individuals. *Nature* **584**, 437-442, doi:10.1038/s41586-020-2456-9 (2020).

Apostolidis, S. A. *et al.* Altered cellular and humoral immune responses following SARS-CoV-2 mRNA vaccination in patients with multiple sclerosis on anti-CD20 therapy. *medRxiv*, 2021.2006.2023.21259389, doi:10.1101/2021.06.23.21259389 (2021).

Sokal, A. *et al.* Memory B cells control SARS-CoV-2 variants upon mRNA vaccination of naive and COVID-19 recovered individuals. *bioRxiv*, 2021.2006.2017.448459, doi:10.1101/2021.06.17.448459 (2021).

Turner, J. S. *et al.* SARS-CoV-2 mRNA vaccines induce persistent human germinal centre responses. *Nature*, doi:10.1038/s41586-021-03738-2 (2021).

Gaebler, C. *et al.* Evolution of antibody immunity to SARS-CoV-2. *Nature* **591**, 639-644, doi:10.1038/s41586-021-03207-w (2021).

Schmidt, F. *et al.* Measuring SARS-CoV-2 neutralizing antibody activity using pseudotyped and chimeric viruses. *J Exp Med* **217**, doi:10.1084/jem.20201181 (2020).

Pilishvili, T. *et al.* Interim Estimates of Vaccine Effectiveness of Pfizer-BioNTech and Moderna COVID-19 Vaccines Among Health Care Personnel - 33 U.S. Sites, January-March 2021. *MMWR Morb Mortal Wkly Rep* **70**, 753-758, doi:10.15585/mmwr.mm7020e2 (2021).

Lopez Bernal, J. *et al.* Effectiveness of Covid-19 Vaccines against the B.1.617.2 (Delta) Variant. *N Engl J Med*, doi:10.1056/NEJMoa2108891 (2021).

Reynolds, C. J. *et al.* Prior SARS-CoV-2 infection rescues B and T cell responses to variants after first vaccine dose. *Science*, doi:10.1126/science.abh1282 (2021).

Wang, Z. *et al.* mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. *Nature* **592**, 616-622, doi:10.1038/s41586-021-03324-6 (2021).

Stamatatos, L. *et al.* mRNA vaccination boosts cross-variant neutralizing antibodies elicited by SARS-CoV-2 infection. *Science*, doi:10.1126/science.abg9175 (2021).

West, A. P. *et al.* Detection and characterization of the SARS-CoV-2 lineage B.1.526 in New York. *bioRxiv*, doi:10.1101/2021.02.14.431043 (2021).

Edara, V. V. *et al.* Infection and Vaccine-Induced Neutralizing-Antibody Responses to the SARS-CoV-2 B.1.617 Variants. *N Engl J Med*, doi:10.1056/NEJMc2107799 (2021).

Planas, D. *et al.* Reduced sensitivity of SARS-CoV-2 variant Delta to antibody neutralization. *Nature*, doi:10.1038/s41586-021-03777-9 (2021).

Victora, G. D. & Nussenzweig, M. C. Germinal centers. *Annu Rev Immunol* **30**, 429-457, doi:10.1146/annurev-immunol-020711-075032 (2012).

Brouwer, P. J. M. *et al.* Potent neutralizing antibodies from COVID-19 patients define multiple targets of vulnerability. *Science* **369**, 643-650, doi:10.1126/science.abc5902 (2020).

733   19   Kreer, C. *et al.* Longitudinal Isolation of Potent Near-Germline SARS-CoV-2-
734   Neutralizing Antibodies from COVID-19 Patients. *Cell* **182**, 843-854 e812,
735   doi:10.1016/j.cell.2020.06.044 (2020).

736   20   Seydoux, E. *et al.* Analysis of a SARS-CoV-2-Infected Individual Reveals Development
737   of Potent Neutralizing Antibodies with Limited Somatic Mutation. *Immunity* **53**, 98-105
738   e105, doi:10.1016/j.immuni.2020.06.001 (2020).

739   21   Taylor, J. J., Pape, K. A., Steach, H. R. & Jenkins, M. K. Humoral immunity. Apoptosis
740   and antigen affinity limit effector cell differentiation of a single naive B cell. *Science* **347**,
741   784-787, doi:10.1126/science.aaa1342 (2015).

742   22   Muecksch, F. *et al.* Development of potency, breadth and resilience to viral escape
743   mutations in SARS-CoV-2 neutralizing antibodies. *bioRxiv*,
744   doi:10.1101/2021.03.07.434227 (2021).

745   23   Amanna, I. J., Carlson, N. E. & Slifka, M. K. Duration of humoral immunity to common
746   viral and vaccine antigens. *N Engl J Med* **357**, 1903-1915, doi:10.1056/NEJMoa066092
747   (2007).

748   24   Davis, C. W. *et al.* Influenza vaccine-induced human bone marrow plasma cells decline
749   within a year after vaccination. *Science* **370**, 237-241, doi:10.1126/science.aaz8432
750   (2020).

751   25   Halliley, J. L. *et al.* Long-Lived Plasma Cells Are Contained within the CD19(-
752   )CD38(hi)CD138(+) Subset in Human Bone Marrow. *Immunity* **43**, 132-145,
753   doi:10.1016/j.immuni.2015.06.016 (2015).

754   26   Slifka, M. K., Antia, R., Whitmire, J. K. & Ahmed, R. Humoral immunity due to long-
755   lived plasma cells. *Immunity* **8**, 363-372, doi:10.1016/s1074-7613(00)80541-5 (1998).

756   27   Dan, J. M. *et al.* Immunological memory to SARS-CoV-2 assessed for up to 8 months
757   after infection. *Science* **371**, doi:10.1126/science.abf4063 (2021).

758   28   Sakharkar, M. *et al.* Prolonged evolution of the human B cell response to SARS-CoV-2
759   infection. *Sci Immunol* **6**, doi:10.1126/sciimmunol.abg6916 (2021).

760   29   Widge, A. T. *et al.* Durability of Responses after SARS-CoV-2 mRNA-1273
761   Vaccination. *N Engl J Med* **384**, 80-82, doi:10.1056/NEJMc2032195 (2021).

762   30   Wajnberg, A. *et al.* Robust neutralizing antibodies to SARS-CoV-2 infection persist for
763   months. *Science* **370**, 1227-1230, doi:10.1126/science.abd7728 (2020).

764   31   Pegu, A. *et al.* Durability of mRNA-1273-induced antibodies against SARS-CoV-2
765   variants. *bioRxiv*, doi:10.1101/2021.05.13.444010 (2021).

766   32   Chen, R. E. *et al.* Resistance of SARS-CoV-2 variants to neutralization by monoclonal
767   and serum-derived polyclonal antibodies. *Nat Med* **27**, 717-726, doi:10.1038/s41591-021-
768   01294-w (2021).

769   33   Abu-Raddad, L. J., Chemaitelly, H., Butt, A. A. & National Study Group for, C.-V.
770   Effectiveness of the BNT162b2 Covid-19 Vaccine against the B.1.1.7 and B.1.351
771   Variants. *N Engl J Med* **385**, 187-189, doi:10.1056/NEJMc2104974 (2021).

772   34   Hacisuleyman, E. *et al.* Vaccine Breakthrough Infections with SARS-CoV-2 Variants. *N
773   Engl J Med* **384**, 2212-2218, doi:10.1056/NEJMoa2105000 (2021).

774   35   Lumley, S. F. *et al.* Antibody Status and Incidence of SARS-CoV-2 Infection in Health
775   Care Workers. *N Engl J Med* **384**, 533-540, doi:10.1056/NEJMoa2034545 (2021).

776   36   McHeyzer-Williams, L. J., Milpied, P. J., Okitsu, S. L. & McHeyzer-Williams, M. G.
777   Class-switched memory B cells remodel BCRs within secondary germinal centers. *Nat
778   Immunol* **16**, 296-305, doi:10.1038/ni.3095 (2015).

37 Mesin, L. *et al.* Restricted Clonality and Limited Germinal Center Reentry Characterize
Memory B Cell Reactivation by Boosting. *Cell* **180**, 92-106 e111,
doi:10.1016/j.cell.2019.11.032 (2020).
38 Pape, K. A., Taylor, J. J., Maul, R. W., Gearhart, P. J. & Jenkins, M. K. Different B cell
populations mediate early and late memory during an endogenous immune response.
Science **331**, 1203-1207, doi:10.1126/science.1201730 (2011).
39 Viant, C. *et al.* Antibody Affinity Shapes the Choice between Memory and Germinal
Center B Cell Fates. *Cell* **183**, 1298-1311 e1211, doi:10.1016/j.cell.2020.09.063 (2020).
40 Sokal, A. *et al.* Maturation and persistence of the anti-SARS-CoV-2 memory B cell
response. *Cell* **184**, 1201-1213 e1214, doi:10.1016/j.cell.2021.01.050 (2021).
41 Feng, L. *et al.* An adenovirus-vectored COVID-19 vaccine confers protection from
SARS-COV-2 challenge in rhesus macaques. *Nat Commun* **11**, 4207,
doi:10.1038/s41467-020-18077-5 (2020).
42 Hassan, A. O. *et al.* A Single-Dose Intranasal ChAd Vaccine Protects Upper and Lower
Respiratory Tracts against SARS-CoV-2. *Cell* **183**, 169-184 e113,
doi:10.1016/j.cell.2020.08.026 (2020).
43 Greaney, A. J. *et al.* Antibodies elicited by mRNA-1273 vaccination bind more broadly
to the receptor binding domain than do those from SARS-CoV-2 infection. *Sci Transl
Med* **13**, doi:10.1126/scitranslmed.abi9915 (2021).
44 Pardi, N. *et al.* Expression kinetics of nucleoside-modified mRNA delivered in lipid
nanoparticles to mice by various routes. *J Control Release* **217**, 345-351,
doi:10.1016/j.jconrel.2015.08.007 (2015).
45 Wu, K. *et al.* Preliminary Analysis of Safety and Immunogenicity of a SARS-CoV-2
Variant Vaccine Booster. *medRxiv*, 2021.2005.2005.21256716,
doi:10.1101/2021.05.05.21256716 (2021).
46 Khoury, D. S. *et al.* Neutralizing antibody levels are highly predictive of immune
protection from symptomatic SARS-CoV-2 infection. *Nat Med* **27**, 1205-1211,
doi:10.1038/s41591-021-01377-8 (2021).
47 Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China.
Nature **579**, 265-269, doi:10.1038/s41586-020-2008-3 (2020).
48 Amanat, F. *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans.
Nat Med **26**, 1033-1036, doi:10.1038/s41591-020-0913-5 (2020).
49 Grifoni, A. *et al.* Targets of T Cell Responses to SARS-CoV-2 Coronavirus in Humans
with COVID-19 Disease and Unexposed Individuals. *Cell* **181**, 1489-1501 e1415,
doi:10.1016/j.cell.2020.05.015 (2020).
50 Barnes, C. O. *et al.* Structures of Human Antibodies Bound to SARS-CoV-2 Spike
Reveal Common Epitopes and Recurrent Features of Antibodies. *Cell* **182**, 828-842 e816,
doi:10.1016/j.cell.2020.06.025 (2020).
51 Weisblum, Y. *et al.* Escape from neutralizing antibodies by SARS-CoV-2 spike protein
variants. *Elife* **9**, doi:10.7554/eLife.61312 (2020).
52 Wang, Z. *et al.* Enhanced SARS-CoV-2 neutralization by dimeric IgA. *Sci Transl Med*
13, doi:10.1126/scitranslmed.abf1555 (2021).
53 Gupta, N. T. *et al.* Change-O: a toolkit for analyzing large-scale B cell immunoglobulin
repertoire sequencing data. *Bioinformatics* **31**, 3356-3358,
doi:10.1093/bioinformatics/btv359 (2015).

54   Soto, C. *et al.* High frequency of shared clonotypes in human B cell receptor repertoires.
Nature **566**, 398-402, doi:10.1038/s41586-019-0934-8 (2019).
55   Guo, Y., Chen, K., Kwong, P. D., Shapiro, L. & Sheng, Z. cAb-Rep: A Database of
Curated Antibody Repertoires for Exploring Antibody Diversity and Predicting Antibody
Prevalence. *Front Immunol* **10**, 2365, doi:10.3389/fimmu.2019.02365 (2019).
829