Cryptic genetic structure and copy-number variation in the ubiquitous forest symbiotic fungus *Cenococcum geophilum*

Authors:

Benjamin Dauphin[†], Maíra de Freitas Pereira[†,¶], Annegret Kohler[¶], Igor V. Grigoriev[§,#], Kerrie Barry[#], Hyunsoo Na[#], Mojgan Amirebrahimi[#], Anna Lipzen[#], Francis Martin[¶], Martina Peter[†,*], Daniel Croll[‡,*]

Affiliations:

[†]Swiss Federal Research Institute WSL, Birmensdorf, Switzerland

[¶]INRAE, UMR 1136 INRAE-University of Lorraine, Interactions Arbres/Microorganismes, Laboratory of Excellence ARBRE, INRAE-Grand Est, Champenoux, France

[§]Department of Plant and Microbial Biology, University of California, Berkeley, USA

[#]U.S. Department of Energy Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, USA

[‡]Laboratory of Evolutionary Genetics, Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

[*]Corresponding authors: daniel.croll@unine.ch, martina.peter@wsl.ch

[*]Co-last authors contributed equally to this study

ORCID:

24   B.D. 0000-0003-0982-4252; M.F. 0000-0002-7659-957X; A.K. 0000-0002-9575-9567; I.G.

25   0000-0002-3136-8903; K.B. 0000-0002-8999-6785; C.V. 0000-0001-9778-2659; F.M. 0000-

26   0002-4737-3715; M.P. 0000-0002-6365-6889; D.C. 0000-0002-2072-380X

28   Running title: Comparative genomics of *Cenococcum geophilum*

Originality-Significance Statement

We provide the first report on the genetic structure and copy-number variation of the globally ubiquitous and key forest symbiotic fungus *Cenococcum geophilum* using whole-genome sequencing data. We found divergent lineages within sampling sites, while closely related lineages appear over large geographic distances on a continental scale. Even though no sexual spore forming structures have been reported to date, we provide evidence of recombination in a specific lineage suggesting mating activity. Our findings help explain the high genetic diversity occurring within populations and their resilience to changing and adverse environmental conditions. Furthermore, we identify a single *MAT1* idiomorph per genome, confirming heterothallism, and discover that major genomic rearrangements are found in their flanking regions based on chromosomal synteny analysis. Intriguingly, a pseudogene of the opposite functional idiomorph has been characterised in each genome, suggesting a common homothallic ancestor to the species. As *Cenococcum geophilum* is a pivotal mycorrhizal associate of a broad range of trees and shrubs providing nutrition and water supply to their hosts, we highlight and discuss the potential role of the large genome-wide structural variations in environmental selection.

Summary

Ectomycorrhizal (ECM) fungi associated with plants constitute one of the most successful symbiotic interactions in forest ecosystems. ECM support trophic exchanges with host plants and are important factors for the survival and stress resilience of trees. However, ECM clades often harbour morpho-species and cryptic lineages, with weak morphological differentiation. How this relates to intraspecific genome variability and ecological functioning is poorly known. Here, we analysed 16 European isolates of the ascomycete *Cenococcum geophilum,* an extremely ubiquitous forest symbiotic fungus with no known sexual or asexual spore forming structures but with a massively enlarged genome. We carried out whole-genome sequencing to

identify single-nucleotide polymorphisms. We found no geographic structure at the European

scale but divergent lineages within sampling sites. Evidence for recombination was restricted

to specific cryptic lineages. Lineage differentiation was supported by extensive copy-number

variation. Finally, we confirmed heterothallism with a single *MAT1* idiomorph per genome.

Synteny analyses of the *MAT1* locus revealed substantial rearrangements and a pseudogene of

the opposite *MAT1* idiomorph. Our study provides the first evidence for substantial genome-

wide structural variation, lineage-specific recombination and low continent-wide genetic

differentiation in *C. geophilum.* Our study provides a foundation for targeted analyses of intra-

specific functional variation in this major symbiosis.

Keywords: Ascomycete, copy-number variation, Dothideomycetes, ectomycorrhizal fungi,

genomics, phylogenetics, recombination, symbiosis

Introduction

Ectomycorrhizal (ECM) fungi associated with plants constitute one of the most successful symbiotic interactions in forest ecosystems (Martin *et al.*, 2016). Abundant ECM partners support the nutrition of many host tree species by facilitating the acquisition of essential nutrients, which is a key determinant of plant survival and productivity under abiotic stress or changing environmental conditions. With exceptional species richness (~20,000; Martin *et al.*, 2016), ECM fungi carry high functional diversity among species that is a key factor of ecosystem resilience (Larsen *et al.*, 2016). The environment, through biotic and abiotic factors, plays a major role in shaping ECM communities and their host range. Environmental variation supports ECM species diversity at both small (e.g. host plants) and large spatial scales, and underlies phenotypic plasticity of functional importance for host trees (van der Linde *et al.*, 2018). Although intraspecific diversity of important symbiotic traits was highlighted in a handful of studies (Cairney, 1999; Rineau & Courty, 2011), the extent to which such variation is genetically driven and contributes to functional diversity within ECM communities remains largely unknown (Johnson *et al.*, 2012).

Cenococcum geophilum Fr. (Dothideomycetes, Ascomycota; Spatafora *et al.*, 2012) is one of the most ubiquitous and abundant ECM fungi in the world (LoBuglio, 1999). *C. geophilum* colonises a large variety of hosts and occupies a wide ecological niche, from alpine to tropical regions (Trappe, 1962). The species tolerates adverse environmental conditions, as it is commonly found in soils with limited water availability, contaminated by heavy metals, or in sites with stressful climatic conditions such as at the timberline (LoBuglio, 1999; Hasselquist *et al.*, 2005; Gonçalves *et al.*, 2009; Herzog *et al.*, 2013). Increased drought tolerance and resistance to desiccation have been shown for *C. geophilum* mycelia and ECM root-tips as compared to other ECM species (Coleman *et al.*, 1989; di Pietro *et al.*, 2007; Fernandez *et al.*, 2013). These studies also revealed high intraspecific variability and

demonstrated that melanin content is one of the key functional traits conferring an advantage under water stress. In nature, *C. geophilum* isolates have shown patterns of local adaptation to serpentine soils with a significant effect of nickel concentrations on fitness-related traits (Gonçalves *et al.*, 2009; Bazzicalupo *et al.*, 2020). Collectively, these results suggest that this cosmopolitan ECM fungus has undergone selection pressures imposed by various environmental factors.

Cenococcum geophilum is typically recognised based on morphology, i.e., a morpho-species. It forms black melanised mycorrhizae with thick, darkly pigmented hyphae spreading from the root tips into the surrounding soil. Although rare in other ECM species (Smith *et al.*, 2015), *C. geophilum* produces sclerotia that act as a "sclerotial bank" in the soil, which are resistant propagules that are melanised and can withstand desiccation for several decades and sequester carbon as dead sclerotia for centuries (Watanabe *et al.*, 2007; Obase *et al.*, 2014, 2018). Although being common, the biology and life cycle remains poorly understood. Phylogenetic analyses have revealed multiple divergent *C. geophilum* lineages on small spatial scales, even at the level of soil core samples (Douhan and Rizzo, 2005; Obase *et al.*, 2016). Combined with the high genetic diversity within populations (ECM root tips and sclerotia), this suggests that *C. geophilum* has undergone cryptic speciation (Obase *et al.*, 2017; Vélez *et al.*, 2021).

Our understanding of *C. geophilum* reproduction is limited. Whilst no spore-producing structures were observed, indicating strictly asexual reproduction through vegetative mycelial propagation, including germination of sclerotia, population genetic studies suggested recombination events that would have involved sexual mating (LoBuglio and Taylor, 2002; Douhan *et al.*, 2007). However, mating may well be restricted to certain environmental conditions or to a few specific lineages with distinct gene pathways involved in sexuality, as suggested in arbuscular mycorrhizal fungi (Riley and Corradi, 2013). This is supported by

phylogenetic studies that showed little gene flow within *C. geophilum* populations. Subtle

geographic structure with long-distance disjunction suggests (Obase *et al.*, 2016) a complex

alternation of sexual and asexual reproduction over space and time (Obase *et al.*, 2017).

Besides further sampling efforts, analyses at a whole genome level are essential to determine

the prevalence of outcrossing and to better understand patterns of cryptic speciation.

The genome of *C. geophilum,* the only known ECM member within Dothideomycetes,

is uniquely large for this class (Peter *et al.*, 2016). A high-quality reference genome (strain

1.58) revealed that about 75% of the genome is composed of transposable elements (TEs) and

this high proportion is unevenly distributed in the genome, with TE-rich and TE-poor regions

(de Freitas Pereira *et al.*, 2018). Its closest sequenced relatives within Gloniaceae,

Mytilinidiales, the saprotrophic species *Glonium stellatum* and *Lepidopterella palustris,* for

example, showed a fourfold smaller genome size (Peter *et al.*, 2016). TE proliferation seems

to be a hallmark for the transition from saprotrophy to symbiosis and in fact, *C. geophilum* is

the ECM species with the highest proportion of TEs known to date among mycorrhizal fungi

with one of the largest genomes, i.e., 178 Mbp; Miyauchi *et al.*, 2020). Nevertheless, the gene

content (14,748) is similar to the average number of genes found in fungi (Miyauchi *et al.*,

2020). Intraspecific genome size variation suggests important structural rearrangements or

gene duplications and deletions (Bourne *et al.*, 2014). Investigation of the reference genome

sequence highlighted the presence of sex-related genes and an intact *MAT1* locus containing

one of the two idiomorphs (here *MAT1-1*) typical for a heterothallic bipolar mating system of

ascomycetes (Peter *et al.*, 2016). The mating-type locus carrying the opposite idiomorph, i.e.,

MAT1-2, has not been analysed in *C. geophilum* so far.

In this study, we used whole-genome sequencing to determine the population genetic

structure of 16 *C. geophilum* isolates across Europe and test for recombination events. We

reconstructed genome-wide phylogenetic relationships within the genus and characterised gene

duplication and deletion events to investigate structural patterns of genetic diversity. Then, we

analysed patterns of selection at the gene and genome level using nucleotide diversity, Tajima's

D, and the ratio of non-synonymous to synonymous polymorphisms. Lastly, we analysed the

genetic architecture of the mating type locus and its recent sequence evolution.

Results

Read mapping, variant calling, and polymorphism filtering

Illumina sequencing of 16 *C. geophilum* strains resulted in 348.9 million read pairs (Fig. 1A;

Table S1). Of those, 293.6 million read pairs (84.16%) were successfully mapped on the

reference strain 1.58. The highest and lowest number of raw read pairs was available for the

strains 1.101 and 1.094, respectively, and the highest (95.54%) and lowest (64.37%) overall

alignment with the reference strain was found in strains 1.061 and 1.096, respectively (Table

S1). An average of 11.3 million read pairs were used for mapping (Table S1). We used the

Genome Analysis Toolkit (GATK) to detect genetic variants from the 16 sequenced strains and

found, after filtering, a total of 345,358 single nucleotide polymorphisms (SNPs), including

284,479 bi-allelic SNPs, of which 67,186 were synonymous variant sites based on the SnpEff

analysis.

[PASTE TABLE 1]

Phylogenetic tree reconstructions

We inferred phylogenetic trees based on the GAPDH and ITS loci using maximum-likelihood

(ML) analyses to identify the main lineages of *C. geophilum* strains. Our two-locus nuclear

dataset contained alignment lengths of 597 and 949 bp for GAPDH and ITS, respectively, with

a total of 1,546 aligned bp for the 40 strains (including outgroups; Table S2). The GAPDH

locus had the most variable sites. The inferred GAPDH and ITS ML tree topologies revealed

several incongruencies, but the concatenated partitions generated a fairly well-resolved

phylogenetic tree, with a majority of branches with BS values > 80; Figs. 1B, S1–2). With

minor exceptions, we found strong support for several monophyletic clades but those were not

congruent with the geographical origin of the samples (Fig. 1B). Overall, the clades that

emerged from the sequenced European strains were depicted in the worldwide clades 5 and 6

described by Obase et al. (2016).

To improve resolution, we used genome-wide data and found highly congruent tree

topologies compared to the GAPDH and ITS phylogeny. The phylogenomic tree was robust to

the use of two near neutral datasets of either synonymous SNPs or randomly-selected genome-

wide SNPs (Fig. 1C). Both genome-based ML trees were well-resolved with most of branches

being strongly supported (BS > 90; Fig. 1C). This indicates that neutral divergence in the

coding regions matches the phylogenetic relationships given by the divergence in non-coding

or coding regions.

[PASTE FIGURE 1]

Recombination events and genetic structure

We analysed patterns of recombination using a Neighbour-Net network in SplitsTree and found

evidence for significant events among the genomes of *C. geophilum* strains (Fig. 2A). We

identified two major genetic groups connected by two strains (1.005 and 1.076). The Φ-test

supported recombination across the genome when all *C. geophilum* were included ($P < 0.001$;

Table S3). However, applying the Φ test to only the cluster not containing the reference genome

(non-ref. cluster) yielded no significant evidence for recombination ($P = 0.641$), suggesting

one clonal and one recombinant genetic group.

As for the phylogenetic analyses, we found no clear geographic structure based on

either principal components (PC) or Bayesian clustering analysis. For example, isolates from

the Pfynwald (CHE) sampling site (1.057, 1.58, 1.060, 1.061, and 1.064) clustered into the two

major groups, reporting a new case of highly divergent lineages present in the same sampling

site (Fig. 2A). In addition, isolates from the Blizyn (POL) sampling site (1.094, 1.096, and

1.097) also showed divergent lineages, but these were only represented in the clonal group with

both mating types (Fig. 2A). No fungal lineages exclusive to certain host tree species were

observed in this dataset. The reference genome 1.58 appeared at the edge of the recombinant

genetic group. The first axes of the PCA captured 61.9% and 9.6% of the genetic variance (PC1

and PC2, respectively; Fig. 2B). In total, the first three axes explained 76.8% of the overall

genotypic variation (Fig. S3). Consistent with the Neighbour-Net network, PC1 and PC2

separated three to four genetic groups with substantial differentiation between the Pfynwald

strains. The intermediate genetic group consisting of the 1.005 and 1.076 strains also clustered

together independently. In addition, Bayesian clustering analyses were congruent between

replicates (Fig. S4) and we found the optimal number of clusters to be $K = 2$ (Fig. 2C). At $K =$

2, two major groups were clearly identified and the intermediate group (strains 1.005 and

1.076), displayed in the PCA and network analysis, belonged to the Pfynwald group containing

the sequenced strain 1.58. At $K = 3$, the intermediate group still exhibited patterns of admixture

between the two major groups, and two Polish genotypes (strains 1.094 and 1.096 belonging

to the phylogenetic Clade 6 in Obase *et al.*, 2016) indicate contributions from a third genetic

group. At higher K values, no significant changes were observed in genotype ancestry

memberships.

[PASTE FIGURE 2]

Detection of copy-number variations

We conducted copy-number variation (CNV) analyses comparing read depth of *C. geophilum*

strain sequencing sets against the reference genome. We improved the robustness of CNV calls

by removing any CNV calls in regions where self-alignments of strain 1.58 reads against the

1.58 reference genome produced candidate CNVs. Overall, we found that gene duplications

occurred ~3.5 fold more frequently than deletions (Fig. 3A). This predominance of gene

duplications was only detected at the highest rates of overlap between genes and CNV regions,

i.e., >95% similarity. In total, we identified 217 and 72 gene duplications and deletions,

respectively, using a threshold of >50% similarity. By inferring the overall tree topology of

based on CNV genotypes, we found patterns consistent with the outcomes of the genetic

clustering and phylogenetic analyses (Fig. 3B). First, five strains (1.022, 1.094, 1.096, 1.097,

and 1.101) formed a monophyletic clade that was paraphyletic in either the two-locus or

genome-based phylogenies (Figs. 1C, 3B). Second, we identified a monophyletic clade with

six strains (1.060, 1.075, 1.100, 1.102, 1.104, and 1.58) that constituted a paraphyletic clade in

the previous analyses. Furthermore, we tested whether the distance between genes affected by

CNVs and the closest TEs are smaller than those for all genes as indicated for other species

(e.g. Hartmann and Croll, 2017). Interestingly, CNV genes were at a slightly larger distance

compared to the background ($P = 0.028$, 719 and 448 median base pairs, respectively; Fig. 4).

[PASTE FIGURE 3]

[PASTE FIGURE 4]

Within-species genetic diversity and signatures of selection

We assessed within-species genetic diversity across the 16 *C. geophilum* genomes and

identified genomic regions with elevated diversity rates along the two longest scaffolds, i.e.,

#1 and #2; Fig. 5A,C, on which the density of coding sequences and transposable elements

(TEs) were negatively correlated (Fig. 5B,D). At the gene level, we estimated the nucleotide

diversity per site (π) to be 0.000–0.020, 0.000–0.028, and 0.000–0.013 for the "all", "ref.

cluster", and "non-ref. cluster" datasets, respectively (Fig. 6A). The genome-wide average π

value was 3.31×10^{-3}, 1.31×10^{-3}, and 1.01×10^{-3}, respectively (Fig. 6A). These results support

that the ref. cluster is genetically more diverse than the non-ref. cluster, although the sample

size is nearly double in the latter. At the gene-level, we calculated Tajima's D values ranging

from -1.90–2.86, -0.87–2.32, and -1.97–2.42 for the "all", "ref. cluster", and "non-ref. cluster"

dataset, respectively (Fig. 6B). At the genome-wide level, the average Tajima's D estimates

were 0.14 (ref. cluster) and 0.31 (non-ref. cluster), with a value of 1.64 for the "all" dataset

(Fig. 6B). This strongly skewed Tajima's D distribution when all strains are included towards

positive values indicates strong population substructure (Fig. 6B). However, the "ref. cluster"

dataset showed a non-normal distribution of Tajima's D with an excess of negative values,

which is indicative of a population expansion. The non-ref. cluster showed a distribution

largely centred around 0 under neutrality and constant population size. A slight deviation

towards positive values suggests a historic decrease in population size or some population

substructure. The distribution of the pN/pS ratio assessed from the all dataset suggests

purifying selection (Fig. 6C).

[PASTE FIGURE 5]

[PASTE TABLE 2]

Then, we characterised the top-candidate genes putatively under selection by

identifying those with the lowest Tajima's D estimates in either the "ref. cluster" or "non-ref.

cluster" (see Experimental procedures). Based on functional annotation, we found that those

top-candidate genes differed to some extent between the ref. and non-ref. clusters (Table 2). In

the recombining "ref. cluster", increased numbers of genes were detected that are putatively

involved in self/non-self recognition containing HET, NACHT, WD40- and Ankyrin-repeat

domains, as well as genes that are members of multigene-families that are expanded in *C.

geophilum* or unique in this species and its closest relatives *G. stellatum* and *L. palustris* as

compared to other Dothideomycetes (Peter *et al.*, 2016). In the "non-ref. cluster", we found

increased number of genes involved in the secondary metabolism (SM; e.g. Lofgren *et al.*,

2021) including two genes found in SM clusters for prenyltransferase synthesis (DMATs;

Khaldi *et al.*, 2010). In both ref. and non-ref. clusters, we found one gene coding for a

carbohydrate-active enzyme (CAZyme; Cantarel *et al.*, 2009). In the "ref. cluster", it codes for

a secreted β-galactosidase (Zeiner *et al.*, 2016), a member of the glycoside hydrolyse family

35 (GH35) implicated in hemicellulose degradation (van den Brink and de Vries, 2011) that is

contracted in the *C. geophilum* genome as compared to other Dothideomycetes. In the "non-

ref. cluster", it is a member of the lytic polysaccharide monooxigenases (LPMO) family A11

that are almost exclusively present in ascomycetes and the one functionally characterised so

far involved in chitin degradation (Hemsworth *et al.*, 2014). Other genes found in both groups

encode for proteins involved in information storage and processing as well as cellular processes

and signalling (Table S4). All these genes were expressed in free-living mycelium and/or

functioning ECM root-tips, but most of them not differentially regulated in these tissues and if

so, then mostly down-regulated in ECMs (Table S4).

[PASTE FIGURE 6]

Chromosomal synteny analysis of the MAT1-1 and MAT1-2 idiomorphs

Our *de novo* draft genome assemblies of other strains than 1.58 generated 151,48 million base

pairs (Mbp) on average, with a mean number of 26,963 contigs per assembly and a mean largest

contig size of 165,938 bp. The value range for N50 and L50 were 5,720–24,382 and 2,247–

4,759, respectively (Table S5). The GC content varied from 33 to 40%, with an average of

37%. Mismatches were lowest and highest with 61 and 260 per 100 kbp for the 1.101 and 1.094

strains, respectively.

We identified two idiomorphs at the *MAT1* locus in *C. geophilum* isolates and found

genetic rearrangements in its flanking coding and non-coding regions based on the synteny

analysis (Fig. 7). A single functional *MAT1* idiomorph was detected in each strain, supporting

the heterothallism of the species (Figs. 1, 7; Table 1). In both *MAT1* loci types, we found a

pseudogene of the opposite *MAT1* idiomorph close to the functional one showing conserved

strand orientation as compared to strains carrying the functional opposite idiomorphs with

unchanged relative position of their open reading frames (ORF). The same was observed for

the closely related *G. stellatum (MAT1-2*: Glost2|444166), whereas *L. palustris*, a homothallic

species and also member of the Mytilinidiales, carries both functional *MAT1* idiomorphs with

conserved   orientation   but   rearranged   position   (*MAT1-1*:   Leppa1|291743;   *MAT1-2*:

Leppa1|291655). The *MAT1-1* and *MAT1-2* idiomorphs were surrounded by a conserved gene

encoding a MFS transporter (555485) and a gene coding for a protein (72340) of unknown

function, both of which showing evidence for an insertion polymorphism generating a non-

syntenic region among the *MAT1-1* and *MAT1-2* strains. Furthermore, the flanking *APN2* gene

(686539) coding for a putative DNA lyase, and the conserved protein TEX2 (634502) coding

for a putative PH domain-containing integral membrane protein were also present in the *MAT1*

loci of the closely related *G. stellatum* and *L. palustris* and are often found adjacent to *MAT1*

idiomorphs in ascomycetes (Debuchy *et al.*, 2014; Wang *et al.*, 2016; APN2: Glost2|386243,

Leppa1|414878; TEX2: Glost2|444168, Leppa1|322867). We further found a gene (555074)

encoding a cysteine-type peptidase and a protein of unknown function (72410).

[PASTE FIGURE 7]

Discussion

Cryptic population genetic structure and evidence for recombination

Our genomic analyses of European *C. geophilum* strains, which showed that divergent lineages co-occur in single sampling sites without major geographic structuring, support previous reports from North America that were based on a restricted number of markers (Douhan and Rizzo, 2005; Vélez *et al.*, 2021). Interestingly, our network analysis indicated that the two main genetic groups had possibly different levels of recombination, with most of the sampled strains being clustered into the clonal group with little gene flow. It is unclear whether this restricted gene flow is due to incompatible mating systems or clonal proliferation. Since neither sexual nor asexual spore forming structures have been reported for *C. geophilum*, this suggests a vegetative mycelial propagation in the soil with limited migration ability of the species. In the recombinant group, cryptic sexual events must have occurred at some point in the near or distant past. In case of a asexually reproducing population, i.e., mating incompatibility, low genetic similarities are expected between strains sampled in the same site, which could explain the observed pattern within the Pfynwald site, where one strain (1.060) clearly differs and did not group in the recombinant cluster (Fig. 2A). Our synteny analyses of the *MAT1* loci showed important rearrangements in close vicinity to the *MAT1* genes, but whether this affects mating compatibility is unclear. Apart from mating incompatibility or sexual versus vegetative propagation as well as isolation by geographic distance, other evolutionary processes could explain the current genetic structure, such as demographic history, past fluctuations in effective population size (e.g. population expansion or contraction), or processes of adaptation to environmental conditions during glacial and interglacial periods, as reported in other ECM fungi (Geml *et al.*, 2006; Sánchez-Ramírez *et al.*, 2015).

Alternatively, recent adaptation of *C. geophilum* to new hosts or environmental

conditions may have led to the emergence of cryptic lineages with strong genetic structure over

short geographical distances. While *C. geophilum* is known to have a broad host tree spectrum,

some lineages may nevertheless show host preferences, as reported in other ECM species

(Rochet *et al.*, 2011; Lofgren *et al.*, 2021). In this respect, the tree-fungus interaction might

have shaped differentiation and genetic isolation of these cryptic lineages, in combination with

recent selection. The present data do not support such a mechanism because no lineage

grouping according to the dominant host tree species was observed. Also, plant-fungus

synthesis experiments showed that functional ectomycorrhizas are formed with diverse tree

species, both broadleaves and conifers, for the reference strain 1.58 but also for others (e.g. de

Freitas Pereira *et al.*, 2018). This shows that the symbiosis can be established with diverse

hosts, even if there are possibly host preferences under natural conditions. To shed light on

host specificity as a potential driver of genetic structuring within and among sites, detailed

information on *C. geophilum* populations associated with roots of different host tree species is

needed. Regarding the environment, adaptation to soil characteristics may prevent wide

distribution, foster differentiation through geographical isolation and, ultimately, lead to

independent and clonal lineages. Note that edaphic specialisation can be coupled with, or even

supplanted by, the competition avoidance strategy of ECM species within soil communities

(Mujic *et al.*, 2016). Clearly, our dataset with 16 strains is limited and further population

genomic studies, with more isolates per populations, are required to be conclusive about the

prevalence of sexual reproduction under varying ecological conditions throughout the species

range.

Massive gene duplications and functional regions under positive selection

Our CNV analysis revealed broad genomic alterations among *C. geophilum* lineages with about

3.5-fold more gene duplications than deletions. Interestingly, the genome-wide pattern of CNV

genotypes is congruent with the tree topology from phylogenetic reconstructions suggesting

that CNVs have accumulated gradually since the origin of the species. However, the genomic

distances between the CNV genes and their closest TEs were in the same range as those for all

genes. This is unexpected since previous studies showed that genes affected by CNVs are often

close to TEs due to their mutagenic potential (e.g. Hartmann and Croll, 2017). A reason for

this finding might be the high abundance of TEs throughout the *C. geophilum* genome.

Although de Freitas *et al.* (2018) found islands of TE-rich and TE-poor regions, these were

rather small and scattered throughout the scaffolds so that most genes have TEs in close

vicinity. With about 75% of its genome comprising TEs, *C. geophilum* exhibits the highest

proportion of transposons and retrotransposons found in ECM fungi (18–75%; Miyauchi *et al.*,

2020) and other Dothideomycetes (1–75%; Peter *et al.*, 2016). Our findings corroborate the

recent proliferation of species-specific TEs and its evolutionary significance in affecting

functional rearrangements of the genome (Daboussi and Capy, 2003; Muszewska *et al.*, 2019).

However, other mechanisms such as aneuploidy, chromosomal duplication, unequal crossing

over may also be responsible for these gene duplications (Bourne *et al.*, 2014). Such high

intraspecific genomic diversity has lately been observed for other long-considered asexual

fungi, such as the ancient arbuscular mycorrhizal glomeromycetes, and TE activity was one of

the explanatory mechanism (Chen *et al.*, 2018). Likewise, genomic studies indicated the

presence of cryptic sex in these species, which, as in our case, might explain the high

adaptability of these fungi to a wide range of hosts and habitats and likely has functional

significance in host-environment interactions (Chen *et al.*, 2018; Reinhardt *et al.*, 2021).

As in another ascomycete ECM fungus, *Tuber melanosporum* (Payen *et al.*, 2015), we

found a large proportion of genes that exhibited an excess of low frequency polymorphisms

relative to neutral expectation (Tajima's $D < -1$, non-ref. cluster; Fig. 6B). Candidate genes

identified as putatively under selection need to be interpreted with caution given our limited

sample size and heterogeneous genetic structure. Nevertheless, a large number of genes

identified in the recombining ref. cluster are putatively implicated in self/non-self recognition

containing HET, NACHT, WD40- and Ankyrin-repeat domains and are members of expanded

gene families. Such HET domain proteins were also identified as putative candidates under

selection for the outcrossing ECM ascomycete *T. melanosporum* in a similar analysis of six

isolates covering its geographical distribution (Payen *et al.*, 2015). Fungal genomes contain a

large number of genes encoding HET domains of unknown function, but resolved functions

often relate to vegetative incompatibility (e.g. conspecific recognition; Paoletti, 2016), hetero-

specific, or even inter-kingdom recognition as seen for characterised NOD-like receptors

(Dyrka *et al.*, 2014). These genes were not up- or down-regulated in ECM root tips as compared

with free-living mycelium in a synthesis experiment (Peter *et al.*, 2016) and, hence, are likely

not involved in specific symbiotic functions in *C. geophilum*. The repeat domains have been

shown to evolve under positive diversifying selection (Dyrka et al 2014) as were some groups

of HET domain containing proteins that displayed higher allelic diversity and a restricted

phylogenetic distribution (Paoletti et al 2016). In the more clonal non-ref. cluster, for which

Tajima's *D* values were generally more negative and centred around 0, we found increased

number of candidate genes putatively under selection, which are involved in the secondary

metabolism. This matches findings from fungal pathogens (e.g. Hartmann *et al.*, 2018).

Interestingly, clearer signatures of positive selection were found for an outcrossing fungal

pathogen species as compared to a more clonally reproducing generalist fungal pathogen

(Derbyshire, 2020).

Genomic rearrangements of the MAT1 locus

The bipolar mating-type (*MAT*) system governs sexual reproduction in ascomycetes and has a

key role in establishing cell identity and determining the sexual cycle. Two fungal partners can

mate only if they carry different *MAT* idiomorphs. Based on our chromosomal synteny

analysis, we showed that both mating-types exist in *C. geophilum* isolates with the *MAT1-1* or

the *MAT1-2* idiomorph. In fact, after screening the sixteen genome sequences, a single *MAT1*

idiomorph was consistently found in each strain, thus confirming the heterothallism of the

species by excluding autogamy. Since the first in-depth genomic analysis of the *MAT1* locus

of the *C. geophilum* reference genome (strain 1.58; Peter *et al.*, 2016), the genomic architecture

of the intact *MAT1* locus was assumed to be stable and no intraspecific reorganisation was

reported in the species. Here, we highlight that both *MAT1-1* and *MAT1-2* loci have undergone

important rearrangements with gene expansions in surrounding regions. Surprisingly, we

identified a *MAT1-1* pseudogene close to the *MAT1-2* idiomorph, i.e., about 3 kbp, and

reciprocally, a *MAT1-2* pseudogene close to the *MAT1-1* idiomorph, i.e., about 2 kbp. This

raises the question of whether a homothallic ancestor may have existed and been the precursor

of independent heterothallic lineages. Interestingly, this *MAT1-1* pseudogene, also found in *G.*

stellatum, is relatively similar to the functional *MAT1-1* idiomorph characterised in the

homothallic outgroup *L. palustris* (Fig. 7), with the same orientation and an equivalent gene

length. Moreover, the gene sequence of the *MAT1-2* idiomorph appears to be enlarged in both

C. geophilum and *G. stellatum* compared to *L. palustris*, indicating important protein

modification. Overall, the genetic makeup of the *MAT1-1* and *MAT1-2* pseudogenes most

likely originates from a common ancestor shared with *L. palustris*, from which distinct mating

types evolved. Such mating shifts, from heterothallism to homothallism, were highlighted in

the *Neurospora* genus, with up to six independent transition events assessed from 43 taxa

(Nygren *et al.*, 2011). Distinct genetic mechanisms enabled these transitions (e.g. translocation

and unequal crossover) with the mediation of specific transposons that reshaped the genetic

architecture of the *MAT* locus (Gioti *et al.*, 2012). The high mobility of genetic material in the

surrounding *MAT* locus is intriguing and points to the role of TE activities and associated

silencing mechanisms such as RIP in shaping the gene architecture and diversity that may

orchestrate sexual reproduction in *C. geophilum* (Serrato-Capuchina and Matute, 2018). In

several fungal species, TEs were described within or in flanking regions of the *MAT* locus, e.g.

in *Blastomyces dermatitidis* (Li *et al.*, 2013), *Ceratocystis* sp. (Simpson *et al.*, 2018), or *T.*

melanosporum (Martin *et al.*, 2010). Together, these studies support the idea that TEs are

frequently inserted at *MAT* loci. They are therefore key evolutionary forces for genome

dynamics likely important not only in pathogenic fungal host and environment interactions as

earlier stated (Raffaele and Kamoun, 2012), but also for symbiotic interactions, as indicated by

increased proliferation of these elements in ECM fungal species, above all in *C. geophilum*

(Miyauchi *et al.*, 2020). Further population genomic studies including genomic architecture

and prevalence of *MAT* loci within populations are needed to fully understand the success of

this widespread and abundant ECM species and its role in forest ecosystems.

Conclusions

Our study provides one of the first intraspecific genomic comparison of an ECM fungal species

sampled on a regional and continental scale. Using whole-genome sequencing from 16

European isolates of the ubiquitous forest symbiotic fungus *Cenococcum geophilum*, we found

divergent lineages in geographically restricted located sampling sites and no strong geographic

structuring. Our genome-wide polymorphism analyses support subdivisions within the species

and suggest two distinct main genetic groups, one more clonal and one recombinant. The

phylogeny of lineages and groupings are largely supported by the abundant copy-number

variations found among genomes. Our gene diversity analyses indicate higher genetic diversity

in the recombinant group even though the clonal cluster contained almost double as many

strains. Although these results need to be interpreted with caution, because of the limited

number of strains studied and an uneven sampling, the top-candidate genes putatively under

positive selection based on Tajima's *D* statistics differed between the two groups. More genes

from lineage-specific, expanded gene families involved in self/non-self recognition were found

in the recombinant cluster, whereas genes involved in the secondary metabolism detected in

the more clonal cluster. Lastly, we confirmed the heterothallism of *C. geophilum* based on the

chromosomal synteny analysis of the *MAT1-1* and *MAT1-2* idiomorphs and uncovered

important genetic rearrangements in their surrounding coding and non-coding regions for both,

strains carrying the same or opposite *MAT1* idiomorphs. Collectively, these findings

demonstrate complex genome architectures in this important ECM symbiont that maybe due

to cryptic sex and/or TE-based mechanisms, as similarly suggested for the long-believed

asexual arbuscular mycorrhizal fungi. Future studies will elucidate the role that this genetic

diversity plays in symbiotic function and the potential for adaptation to changing environments.

Experimental Procedures

Study range and selection of isolates

In this study, we selected 16 *Cenococcum geophilum* isolates to cover as broad as possible the European range of the species and the divergent phylogenetic lineages previously reported (Obase *et al.*, 2016). All strains were isolated from sclerotia collected below diverse host tree species, i.e., broadleaves and conifers, to maximise intra-specific genetic diversity in close and distant sampling sites. After screening, both mating types were included to investigate sexual reproduction from natural populations (Fig. 1; Table 1). Pure cultures of *C. geophilum* were grown in liquid culture containing a Cg-Medium, i.e., a modified Melin Norkran's (MMN) medium comprising casein (Kerner *et al.*, 2012), and incubated at room temperature in the dark. We cultivated isolates for a month and harvested mycelia that were then pulverised in liquid nitrogen for storage at –80°C until processing.

DNA extraction and barcoding

Genomic DNA was extracted using the PowerMax Soil DNA extraction kit (MoBio, now DNeasy PowerMax Soil kit of Qiagen) according to the manufacturer's instructions. DNA was quantified by spectrophotometry (Qubit, Thermo Fisher Scientific, Waltham, USA), and the quality was inspected on agarose gel (1.5%). A minimum of 1 μg of DNA was sent to the US Department of Energy Joint Genome Institute (JGI) for genome resequencing.

To identify the lineage of each isolate, we amplified and sequenced two phylogenetically informative loci at the intraspecific level, the glyceraldehyde 3-phosphate dehydrogenase (GAPDH) gene and the internal transcribed spacer (ITS) region (Obase *et al.*, 2016), using Sanger dideoxy sequencing. The optimised polymerase chain reaction (PCR) programs for ITS and the GAPDH gene were conducted following Obase *et al*. (2014). For the GAPDH gene, we used universal primers, as reported in Berbee *et al*. (1999), and for ITS, we

used ITS1F and ITS4 (White *et al.*, 1990; Gardes and Bruns, 1993). PCR products were cleaned

up using the illustra™ ExoProStar™ enzymatic PCR and sequence reaction clean kit according

to the manufacturer's instructions (Merck KGaA, Darmstadt, Germany). Cycle sequencing was

performed using the BigDye™ Terminator v3.1 Cycle Sequencing kit and products were

purified with the BigDye XTerminator™ purification kit (Thermo Fisher Scientific) before to

analyse them on an Applied Biosystems ABI3130 genetic analyser. Loci were sequenced with

both forward and reverse primers to ensure unambiguous chromatogram profiles and all

sequences were deposited to NCBI (MZ570241-MZ570257; MZ577329-MZ577345).

Genome sequencing and reference genome sequences

Plate-based DNA library preparation for Illumina sequencing was performed on the Sciclone

NGS robotic liquid handling system (PerkinElmer, Waltham, USA) using KAPA Biosystems

library preparation kit (Roche, Basel, Switzerland). For each sample, 200 ng of DNA was

sheared to 600 base pairs (bp) using a Covaris LE220 focused-ultrasonicator (Woburn,

Massachusetts, USA). The sheared DNA fragments were size selected by double-SPRI and

then the selected fragments were end-repaired, A-tailed, and ligated with Illumina compatible

sequencing adaptors from IDT containing a unique molecular index barcode for each sample

library. The prepared library was quantified using KAPA Biosystem's next-generation

sequencing library qPCR kit and run on a Roche LightCycler 480 real-time PCR instrument.

The quantified library was then multiplexed with other libraries, and the pool of libraries was

then prepared for sequencing on the Illumina HiSeq sequencing platform utilising a TruSeq

paired-end cluster kit, v4, and Illumina's cBot instrument to generate a clustered flow cell for

sequencing (San Diego, California, USA). Sequencing of the flow cell was performed on the

Illumina HiSeq 2500 sequencer using HiSeq TruSeq SBS sequencing kits, v4, following a

2×100 indexed run recipe at the Joint Genome Institute (JGI; Walnut Creek, California, USA).

We accessed the *C. geophilum* raw reads data from the NCBI Short Read Archive

(Table S1) and used the *Glonium stellatum* genome as an outgroup species (Peter *et al.*, 2016;

https://genome.jgi.doe.gov/Glost2) in downstream analyses. The *C. geophilum* reference

genome sequence and gene annotation data (filtered models) for the strain 1.58 were

downloaded from the JGI (Peter *et al.*, 2016; https://genome.jgi.doe.gov/Cenge3). The

sequenced strains included the reference genome strain 1.58 to control for the quality of SNP

calling (Peter *et al.*, 2016).

Read mapping, variant calling, and polymorphism filtering

We downloaded sequencing data using fastq-dump, as implemented in the SRA-Toolkit v.

2.9.0 (http://ncbi.github.io/sra-tools/), and inspected read quality with FastQC v. 0.11

(Andrews, 2017). We performed quality trimming of raw reads with Trimmomatic v. 0.36

(Bolger *et al.*, 2014) based on the following quality filters: ILLUMINACLIP:TruSeq3-

PE.fa:2:30:10, SLIDINGWINDOW:5:10, LEADING:10, TRAILING:10, and MINLEN:50.

Trimmed reads were mapped on the reference genome 1.58 using Bowtie v. 2.3 with the -very-

sensitive and -local settings (Langmead *et al.*, 2009). The resulting BAM files were sorted by

position and indexed with SAMtools v. 1.6 (Li *et al.*, 2009).

We applied the Genome Analysis Toolkit (GATK) v. 3.7 and used the HaplotypeCaller

to perform the SNP and indel discovery (McKenna *et al.*, 2010). Individual SNP calls were

merged and jointly genotyped using the GATK GenotypeGVCFs tool. Then, we filtered the

joint Variant Call Format (VCF) file using hard filtering parameters according to GATK best

practices recommendations (Van der Auwera *et al.*, 2013). The following thresholds were

applied for filtering: $AN \geq 12$, i.e., at least 12 isolates had a callable genotype, $QUAL \geq 5,000$,

$QD \geq 5.0$, $MQ \geq 20.0$, ReadPosRankSum, MQRankSum and BaseQRankSum each comprised

between -2.0 and 2.0. We further classified SNP loci as synonymous or non-synonymous

variant sites using SnpEff v. 4.3p (Cingolani *et al.*, 2012).

 Phylogenetic tree reconstructions

To infer the phylogenetic relationships of the 16 *C. geophilum* strains within the seven main

clades described in Obase *et al.* (2016), we assembled a broader dataset (GAPDH and ITS) and

reconstructed a two-loci phylogenetic tree based on maximum likelihood (ML) analysis. The

additional strains (22 *C. geophilum*, one *Pseudocenococcum floridanum*, and one *G. stellatum*)

covered most of the phylogenetic diversity and intercontinental distribution of the species. This

includes important disjunctions in North America and potential cryptic taxa in European Clades

5 and 6 (Obase *et al.*, 2016), which were investigated using a coalescent-based species

delimitation approach (Zhang *et al.*, 2013). For each locus, we aligned sequences using

MAFFT v. 7 with default parameters (Katoh *et al.*, 2019) and excluded ambiguous regions

using Gblocks v.0.91b with the less stringent settings (Castresana, 2000). For each locus, we

implemented the optimal substitution model, based on AICc scores calculated with the program

ModelTest-NG v. 0.1.7 (Darriba *et al.*, 2019; Table S2). We ran ML tree searches with a

randomised maximum parsimony starting tree and gamma-distributed site rates using RAxML-

NG v. 1.0.0 (Kozlov *et al.*, 2019; Edler *et al.*, 2021). We performed 20 runs and set 1,000

bootstrap replicates for each run, computed on the best-scoring ML tree with a bootstrap

random-number seed. We fixed *G. stellatum* as the outgroup to root phylograms.

At the genome-wide level, we generated two SNP datasets to infer independent

phylogenetic tree reconstructions. First, from the overall SNP data genotyped in all 17 strains

(16 *C. geophilum* and one *G. stellatum*), we randomly selected a single SNP per 1 kilo base-

pair (kbp) of the reference genome sequence using VCFtools v. 0.1.15 (Danecek *et al.*, 2011).

Second, from the overall SNP dataset, we retained only synonymous variants identified in

coding regions (see below SnpEff). We used each SNP dataset for an independent ML tree

reconstruction with RAxML-NG (Kozlov *et al.*, 2019; Edler *et al.*, 2021). We ran the best ML

tree by analysing 20 different starting trees and performed 1,000 bootstrap replicates on the

best-scoring ML tree with a bootstrap random-number seed. We implemented a General Time

Reversible (GTR) substitution model and allowed for rate heterogeneity (GTR+G).

Recombination events and genetic structure

To gain insights into recombination events and genetic structure, we generated a SNP dataset

for intraspecific genetic analyses of *C. geophilum* using the following criteria: a single SNP

randomly selected per 1 kbp of the reference genome and genotyped in a least 90% of the

strains, i.e., genotyped in at least 15 out of 16 strains. We used SplitsTree v. 4.14.6 to build up

a Neighbour-Net network from all *C. geophilum* strains (Huson and Bryant, 2006), and

uncorrected p distances calculated from the SNP matrix, which is the proportion of positions

at which two sequences differ, to generate the network. The network was drawn based on equal

angle splits. Tests of recombination within genetic groups of strains were carried out using the

Φ test implemented in SplitsTree (Bruen *et al.*, 2006). The test identifies evidence of

recombination based on phylogenetic compatibilities of neighbouring SNPs in the matrix. We

ran the test with the default settings of a 100-characters window size on the SNP matrix. Three

genotype datasets were analysed to assess lineage-specific patterns; comprising all strains

(referred as "all"), a reference cluster which contained the reference genome 1.58 (referred as

"ref. cluster" including 1.005, 1.057, 1.061, 1.064, 1.076, and 1.58), and a non-reference cluster

which did not contain the reference genome (referred as "non-ref. cluster" including 1.022,

1.060, 1.075, 1.094, 1.096, 1.097, 1.100, 1.101, 1.102, and 1.104).

We conducted Bayesian unsupervised genetic clustering using STRUCTURE v. 2.3.4

(Pritchard *et al.*, 2000). We used an admixture model with correlated frequencies and no prior

regarding the population of origin (Falush *et al.*, 2003). The *K* parameter was tested for values

ranging from 1 to 5, with 10 repetitions for each tested *K* value. We used 50,000 iterations as

a burn-in period and 100,000 iterations per run for the Monte Carlo Markov Chain (MCMC)

replicates. We visually inspected the convergence of parameters and diagnosed the optimal *K*

585 value based on the stabilisation of –Ln(K) values. We performed a principal component

586 analysis (PCA) to explore genotypic variability based on the SNP matrix using the R package

587 SNPRelate v. 1.20.1 (Zheng *et al.*, 2012).

588 *Detection of copy-number variations*

589 We identified segmental deletions and duplications in sequenced isolates compared to the

590 reference genome 1.58 using CNVcaller (Wang *et al.*, 2017). CNVcaller uses normalised read-

591 depth of short read alignments to identify high-confidence copy-number variation (CNV)

592 regions. The window size for the CNV calling was set to 500 bp according to the

593 recommendations based on sequencing coverage. The lower and upper GC content limits were

594 set to 0.2 and 0.7, respectively, and the upper limit of gap content was set to 0.5. We required

595 a minimum of Pearson's correlation coefficient between two adjacent non-overlapping

596 windows to be >0.5, as recommended for less than 30 samples. We also specified that only a

597 single individual displayed a potential gain/loss event to define a candidate CNV window (-h

598 1 and -f 0.05) given the heterogeneity in genotypes and small sample size. CNV regions were

599 further filtered based on the normalised read depth. Deletions and duplications were retained

600 if the normalised read depth was <0.4 and >1.8, respectively. CNV regions called for the

601 reference genome 1.58 were aligned against the reference genome for validation, which

602 showed possible issues in read alignment and/or the local assembly quality. Hence, gene

603 deletions and duplications among *C. geophilum* strains were only retained if the reference strain

604 showed a normalised read depth between 0.7–1.3 for the same region. A gene was considered

605 affected by a duplication or deletion if more than 50% of the coding sequence length was

606 overlapping with a CNV region using the R package ggplot2 v. 3.3.0 (Wickham, 2011). One

607 CNV region may contain several duplicated or deleted genes. Then, genomic distances between

608 all genes and their closest transposable elements (TEs), and between CNV genes and their

closest TEs were calculated with the *closest* function in BEDTools v 2.28.0 (Quinlan and Hall,

2010). Significance between median values was calculated using a two-sided Wilcoxon test.

Within-species genetic diversity and signatures of selection

To improve our understanding of genomic signatures of selection, we analysed three species-

specific estimates of gene diversity and their deviations from genome-wide average; nucleotide

diversity per site (π), Tajima's *D*, and *p*N/*p*S ratios. We assessed these population genetic

statistics for individual genes using the R package PopGenome v. 2.6.1 (Pfeifer *et al.*, 2014)

and included all SNPs falling within coding sequence boundaries that were genotyped in at

least 15 out of 16 *C. geophilum* strains. At the genome-wide level, a negative Tajima's *D* is

indicative of a population size expansion (e.g., after a bottleneck), whereas a positive Tajima's

D is supporting a decrease in population size. At the gene-specific level, negative Tajima's *D*

values are indicating of recent positive selection, while positive values are imputable to

balancing selection (Hohenlohe *et al.*, 2010). For π estimates, a decrease in gene-specific

values in comparison to the average genome-wide level suggests a signature of positive

selection (Hohenlohe *et al.*, 2010). Regarding *p*N/*p*S, a ratio greater than one implies positive

selection, whereas a value less than one assumes purifying selection (Tanaka and Nei, 1989).

The three genotype datasets (see above) were analysed to evaluate the robustness of the results.

To further examine the genes putatively under selection, we identified top-candidate genes

based on the 20 lowest Tajima's *D* values in each of the two datasets, the ref. and non-ref.

cluster. Functional annotation of the top-candidate genes was carried as outlined in Peter *et al.*,

(2016) and according to the JGI annotations (mycocosm.jgi.doe.gov), including predicted gene

ontology (GO), EuKaryotic Orthologous Groups (KOG), protein family clarification using the

InterPro database (Blum *et al.*, 2021), Carbohydrate-Active enZYme (CAZY; Cantarel *et al.*,

2009) and secondary metabolism genes and cluster prediction (Khaldi *et al.*, 2010). Gene

expression data from free-living mycelium and ECM root-tips were taken from the study of

Peter *et al*. 2016 and presented here. In addition, we examined the affiliation to multigene

families, their PFAM domains and family size changes according to expansion/depletion

analysis as detailed in Ohm *et al*. (2012) and CAFE analyses (De Bie *et al.*, 2006) using the

species set outlined in Peter *et al*. 2016 and available on the JGI *C. geophilum* 1.58 portal under

the MCL Clusters Run "Cenococcum + 50 genomes 03/21/14.399".

Chromosomal synteny analysis of the MAT1 locus

We analysed chromosomal synteny of the *MAT1* locus containing the *MAT1-1* and *MAT1-2*

idiomorphs based on our *de novo* genome assemblies of individual *C. geophilum* isolates. We

used SPAdes v. 3.11.1 to assemble genomic scaffolds based on the paired-end Illumina

sequencing data described above (Bankevich *et al.*, 2012). The SPAdes pipeline includes the

BayesHammer read error correction module and builds contigs in a stepwise procedure based

on increasing kmer lengths. We defined a kmer range of 21, 33, 45, 67 and 89 and enabled the

--careful option to reduce mismatches and indel errors in the assembly. Assembled scaffolds

were error-corrected based on read realignments. We used the tool QUAST v. 4.6.0 to retrieve

assembly statistics for each strain (Gurevich *et al.*, 2013) and performed a blastn to identify

assembled scaffolds that spanned the *MAT1* loci of the reference genome strain 1.58 (Camacho

et al., 2009; Peter *et al.*, 2016). For this, three genes in each direction of the *MAT1* locus were

inspected. Once matching scaffolds were identified in individual genome assemblies, we

performed pairwise blastn analyses between these scaffolds. Segmental blast hits between pairs

of scaffolds known to contain the *MAT1* locus were leniently filtered for a minimum identity

of 50% and a minimum alignment length of 50 bp to assess synteny patterns. For the

comparison of *G. stellatum*, *L. palustris* and *C. geophilum*, we performed a tblastx search

translated nucleotide and filtered additionally for a minimum bitscore of 100. We visualised

pairwise synteny among scaffolds using the R package GenoPlotR v. 0.8.7 (Guy *et al.*, 2011).

Acknowledgements

We thank Stephanie Pfister and Barbara Meier for help in the laboratory; the staff at the Joint

Genome Initiative for sequencing. We thank Claire Veneault-Fourrey for helpful comments

and discussions. This project was supported within the framework of the ARBRE/WSL project

"Blacksecret" by grants from the WSL and the French National Agency of Research (ANR),

and as part of the "Investissement d'Avenir" program (ANR-11_LABX-0002-01) of Labex

ARBRE (CFP15; to AK and MP) and the EC-supported Network of Excellence Evoltree

(GOCE-016322 to MP).

References

Andrews, S. (2017) FastQC: a quality control tool for high throughput sequence data.

Van der Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine,
A., et al. (2013) From FastQ data to high-confidence variant calls: The genome analysis
toolkit best practices pipeline. *Curr Protoc Bioinforma* **43**: 1–33.

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., et al. (2012)
SPAdes: A new genome assembly algorithm and its applications to single-cell
sequencing. *J Comput Biol* **19**: 455–477.

Bazzicalupo, A.L., Ruytinx, J., Ke, Y.H., Coninx, L., Colpaert, J. V., Nguyen, N.H., et al.
(2020) Fungal heavy metal adaptation through single nucleotide polymorphisms and
copy-number variation. *Mol Ecol* **29**: 4157–4169.

Berbee, M.L., Pirseyedi, M., and Hubbard, S. (1999) Cochliobolus phylogenetics and the origin
of known, highly virulent pathogens, inferred from ITS and glyceraldehyde-3-phosphate
dehydrogenase gene sequences. *Mycologia* **91**: 964–977.

De Bie, T., Cristianini, N., Demuth, J.P., and Hahn, M.W. (2006) CAFE: A computational tool
for the study of gene family evolution. *Bioinformatics* **22**: 1269–1271.

Blum, M., Chang, H.Y., Chuguransky, S., Grego, T., Kandasaamy, S., Mitchell, A., et al.
(2021) The InterPro protein families and domains database: 20 years on. *Nucleic Acids
Res* **49**: D344–D354.

Bolger, A.M., Lohse, M., and Usadel, B. (2014) Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**: 2114–2120.

Bourne, E.C., Mina, D., Gonçalves, S.C., Loureiro, J., Freitas, H., and Muller, L.A. (2014)
Large and variable genome size unrelated to serpentine adaptation but supportive of
cryptic sexuality in *Cenococcum geophilum*. *Mycorrhiza* **24**: 13–20.

van den Brink, J. and de Vries, R.P. (2011) Fungal enzyme sets for plant polysaccharide
degradation. *Appl Microbiol Biotechnol* **91**: 1477–1492.

Bruen, T.C., Philippe, H., and Bryant, D. (2006) A simple and robust statistical test for
detecting the presence of recombination. *Genetics* **172**: 2665–2681.

Cairney, J.W. (1999) Intraspecific physiological variation: Implications for understanding
functional diversity in ectomycorrhizal fungi. *Mycorrhiza* **9**: 125–135.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden,
T.L. (2009) BLAST+: Architecture and applications. *BMC Bioinformatics* **10**: 1–9.

Cantarel, B.I., Coutinho, P.M., Rancurel, C., Bernard, T., Lombard, V., and Henrissat, B.
(2009) The Carbohydrate-Active EnZymes database (CAZy): An expert resource for
glycogenomics. *Nucleic Acids Res* **37**: 233–238.

Castresana, J. (2000) Selection of conserved blocks from multiple alignments for their use in
phylogenetic analysis. *Mol Biol Evol* **17**: 540–552.

Charlesworth, D. and Willis, J.H. (2009) The genetics of inbreeding depression. *Nat Rev Genet*
10: 783–796.

Chen, E.C.H., Morin, E., Beaudet, D., Noel, J., Yildirir, G., Ndikumana, S., et al. (2018) High
intraspecific genome diversity in the model arbuscular mycorrhizal symbiont
Rhizophagus irregularis. *New Phytol* **220**: 1161–1171.

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., et al. (2012) A program
for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff:
SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly* **6**: 80–92.

Coleman, M.D., Bledsoe, C.S., and Lopushinsky, W. (1989) Pure culture response of
ectomycorrhizal fungi to imposed water stress. *Can J Bot* **67**: 29–39.

Daboussi, M.J. and Capy, P. (2003) Transposable elements in filamentous fungi. *Annu Rev*
Microbiol **57**: 275–299.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., et al. (2011)
The variant call format and VCFtools. *Bioinformatics* **27**: 2156–2158.

Darriba, D., Posada, D., Kozlov, A.M., Stamatakis, A., Morel, B., and Flouri, T. (2020)
ModelTest-NG: A new and scalable tool for the selection of DNA and protein
evolutionary models. *Mol Biol Evol* **37**: 291–294.

Debuchy, R., Berteaux-Lecellier, V., and Silar, P. (2014) Mating systems and sexual
morphogenesis in ascomycetes. In *Cellular and molecular biology of filamentous fungi*.
Borkovich, K.A. and Ebbole, D.J. (eds). Washington, U.S.A.: ASM Press, pp. 499–535.

Derbyshire, M.C. (2020) Bioinformatic detection of positive selection pressure in plant
pathogens: The neutral theory of molecular sequence evolution in action. *Front Microbiol*
11: 1–14.

Douhan, G.W., Martin, D.P., and Rizzo, D.M. (2007) Using the putative asexual fungus
Cenococcum geophilum as a model to test how species concepts influence recombination
analyses using sequence data from multiple loci. *Curr Genet* **52**: 191–201.

Douhan, G.W. and Rizzo, D.M. (2005) Phylogenetic divergence in a local population of the
ectomycorrhizal fungus *Cenococcum geophilum*. *New Phytol* **166**: 263–271.

Dyrka, W., Lamacchia, M., Durrens, P., Kobe, B., Daskalov, A., Paoletti, M., et al. (2014)
Diversity and variability of NOD-like receptors in fungi. *Genome Biol Evol* **6**: 3137–3158.

Edler, D., Klein, J., Antonelli, A., and Silvestro, D. (2021) raxmlGUI 2.0: A graphical interface
and toolkit for phylogenetic analyses using RAxML. *Methods Ecol Evol* **12**: 373–377.

Falush, D., Stephens, M., and Pritchard, J.K. (2003) Inference of population structure using
multilocus genotype data: Linked loci and correlated allele frequencies. *Genetics* **164**:
1567–1587.

Fernandez, C.W., McCormack, M.L., Hill, J.M., Pritchard, S.G., and Koide, R.T. (2013) On
the persistence of *Cenococcum geophilum* ectomycorrhizas and its implications for forest
carbon and nutrient cycles. *Soil Biol Biochem* **65**: 141–143.

de Freitas Pereira, M., Veneault-Fourrey, C., Vion, P., Guinet, F., Morin, E., Barry, K.W., et
al. (2018) Secretome analysis from the ectomycorrhizal ascomycete *Cenococcum*
geophilum. *Front Microbiol* **9**: 1–17.

Gardes, M. and Bruns, T.D. (1993) ITS primers with enhanced specificity for basidiomycetes
- application to the identification of mycorrhizae and rusts. *Mol Ecol* **2**: 113–118.

Geml, J., Laursen, G.A., O'Neill, K., Nusbaum, H.C., and Taylor, D.L. (2006) Beringian
origins and cryptic speciation events in the fly agaric (*Amanita muscaria*). *Mol Ecol* **15**:
225–239.

Gioti, A., Mushegian, A.A., Strandberg, R., Stajich, J.E., and Johannesson, H. (2012)
Unidirectional evolutionary transitions in fungal mating systems and the role of

transposable elements. *Mol Biol Evol* **29**: 3215–3226.

Gonçalves, S.C., Martins-Loução, M.A., and Freitas, H. (2009) Evidence of adaptive tolerance
to nickel in isolates of *Cenococcum geophilum* from serpentine soils. *Mycorrhiza* **19**: 221–
230.

Gurevich, A., Saveliev, V., Vyahhi, N., and Tesler, G. (2013) QUAST: Quality assessment tool
for genome assemblies. *Bioinformatics* **29**: 1072–1075.

Guy, L., Kultima, J.R., Andersson, S.G.E., and Quackenbush, J. (2011) genoPlotR:
comparative gene and genome visualization in R. *Bioinformatics* **27**: 2334–2335.

Hartmann, F.E. and Croll, D. (2017) Distinct trajectories of massive recent gene gains and
losses in populations of a microbial eukaryotic pathogen. *Mol Biol Evol* **34**: 2808–2822.

Hartmann, F.E., McDonald, B.A., and Croll, D. (2018) Genome-wide evidence for divergent
selection between populations of a major agricultural pathogen. *Mol Ecol* **27**: 2725–2741.

Hasselquist, N., Germino, M.J., McGonigle, T., and Smith, W.K. (2005) Variability of
Cenococcum colonization and its ecophysiological significance for young conifers at
alpine-treeline. *New Phytol* **165**: 867–873.

Hemsworth, G.R., Henrissat, B., Davies, G.J., and Walton, P.H. (2014) Discovery and
characterization of a new family of lytic polysaccharide monooxygenases. *Nat Chem Biol*
10: 122–126.

Herzog, C., Peter, M., Pritsch, K., Günthardt-Goerg, M.S., and Egli, S. (2013) Drought and air
warming affects abundance and exoenzyme profiles of *Cenococcum geophilum* associated
with *Quercus robur*, *Q. petraea* and *Q. pubescens*. *Plant Biol* **15**: 230–237.

Hohenlohe, P.A., Phillips, P.C., and Cresko, W.A. (2010) Using population genomics to detect
selection in natural populations: Key concepts and methodological considerations. *Int J*
Plant Sci **171**: 1059–1071.

Huson, D.H. and Bryant, D. (2006) Application of phylogenetic networks in evolutionary
studies. *Mol Biol Evol* **23**: 254–267.

Johnson, D., Martin, F., Cairney, J.W., and Anderson, I.C. (2012) The importance of
individuals: Intraspecific diversity of mycorrhizal plants and fungi in ecosystems. *New*
Phytol **194**: 614–628.

Katoh, K., Rozewicki, J., and Yamada, K.D. (2019) MAFFT online service: Multiple sequence
alignment, interactive sequence choice and visualization. *Brief Bioinform* **20**: 1160–1166.

Kerner, R., Delgado-Eckert, E., Del Castillo, E., Müller-Starck, G., Peter, M., Kuster, B., et al.
(2012) Comprehensive proteome analysis in *Cenococcum geophilum* Fr. as a tool to
discover drought-related proteins. *J Proteomics* **75**: 3707–3719.

Khaldi, N., Seifuddin, F.T., Turner, G., Haft, D., Nierman, W.C., Wolfe, K.H., and Fedorova,
N.D. (2010) SMURF: Genomic mapping of fungal secondary metabolite clusters. *Fungal*
Genet Biol **47**: 736–741.

Kozlov, A.M., Darriba, D., Flouri, T., Morel, B., and Stamatakis, A. (2019) RAxML-NG: A
fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference.
Bioinformatics **35**: 4453–4455.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009) Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol* **10**: 1–10.

Larsen, P.E., Sreedasyam, A., Trivedi, G., Desai, S., Dai, Y., Cseke, L.J., and Collart, F.R.

(2016) Multi-omics approach identifies molecular mechanisms of plant-fungus mycorrhizal interaction. *Front Plant Sci* **6**: 1–17.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009) The sequence alignment/map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Li, W., Sullivan, T.D., Walton, E., Averette, A.F., Sakthikumar, S., Cuomo, C.A., et al. (2013) Identification of the mating-type (*MAT*) locus that controls sexual reproduction of blastomyces dermatitidis. *Eukaryot Cell* **12**: 109–117.

van der Linde, S., Suz, L.M., Orme, D. l., Cox, F., Andreae, H., Asi, E., et al. (2018) Environment and host as large-scale controls of ectomycorrhizal fungi. *Nature* **558**: 243–248.

LoBuglio, K.F. (1999) *Cenococcum*. In *Ectomycorrhizal Fungi: Key Genera in Profile*. Cairney, J.W.G. and Chambers, S.M. (eds). Heidelberg, Germany: Springer, pp. 287–309.

LoBuglio, K.F. and Taylor, J.W. (2002) Recombination and genetic differentiation in the mycorrhizal fungus *Cenococcum geophilum* Fr. *Mycologia* **94**: 772–780.

Lofgren, L.A., Nguyen, N.H., Vilgalys, R., Ruytinx, J., Liao, H.L., Branco, S., et al. (2021) Comparative genomics reveals dynamic genome evolution in host specialist ectomycorrhizal fungi. *New Phytol* **230**: 774–792.

Martin, F., Kohler, A., Murat, C., Balestrini, R., Coutinho, P.M., Jaillon, O., et al. (2010) Périgord black truffle genome uncovers evolutionary origins and mechanisms of symbiosis. *Nature* **464**: 1033–1038.

Martin, F., Kohler, A., Murat, C., Veneault-Fourrey, C., and Hibbett, D.S. (2016) Unearthing the roots of ectomycorrhizal symbioses. *Nat Rev Microbiol* **14**: 760–773.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A.Y., Cibulskis, K., Kernytsky, A.M., et al. (2010) The genome analysis toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* **20**: 1297–1303.

Miyauchi, S., Kiss, E., Kuo, A., Drula, E., Kohler, A., Sánchez-García, M., et al. (2020) Large-scale genome sequencing of mycorrhizal fungi provides insights into the early evolution of symbiotic traits. *Nat Commun* **11**: 1–17.

Mujic, A.B., Durall, D.M., Spatafora, J.W., and Kennedy, P.G. (2016) Competitive avoidance not edaphic specialization drives vertical niche partitioning among sister species of ectomycorrhizal fungi. *New Phytol* **209**: 1174–1183.

Muszewska, A., Steczkiewicz, K., Stepniewska-Dziubinska, M., and Ginalski, K. (2019) Transposable elements contribute to fungal genes and impact fungal lifestyle. *Sci Rep* **9**: 1–10.

Nygren, K., Strandberg, R., Wallberg, A., Nabholz, B., Gustafsson, T., García, D., et al. (2011) A comprehensive phylogeny of *Neurospora* reveals a link between reproductive mode and molecular evolution in fungi. *Mol Phylogenet Evol* **59**: 649–663.

Obase, K., Douhan, G.W., Matsuda, Y., and Smith, M.E. (2014) Culturable fungal assemblages growing within *Cenococcum* sclerotia in forest soils. *FEMS Microbiol Ecol* **90**: 708–717.

Obase, K., Douhan, G.W., Matsuda, Y., and Smith, M.E. (2018) Isolation source matters: Sclerotia and ectomycorrhizal roots provide different views of genetic diversity in *Cenococcum geophilum*. *Mycologia* **110**: 473–481.

Obase, K., Douhan, G.W., Matsuda, Y., and Smith, M.E. (2017) Progress and challenges in

understanding the biology, diversity, and biogeography of *Cenococcum geophilum*. In *Biogeography of mycorrhizal symbiosis*. Tedersoo, L. (ed)., pp. 299–317.

Obase, K., Douhan, G.W., Matsuda, Y., and Smith, M.E. (2016) Revisiting phylogenetic diversity and cryptic species of *Cenococcum geophilum* sensu lato. *Mycorrhiza* **26**: 529–540.

Ohm, R.A., Feau, N., Henrissat, B., Schoch, C.L., Horwitz, B.A., Barry, K.W., et al. (2012) Diverse lifestyles and strategies of plant pathogenesis encoded in the genomes of eighteen Dothideomycetes fungi. *PLoS Pathog* **8**:.

Paoletti, M. (2016) Vegetative incompatibility in fungi: From recognition to cell death, whatever does the trick. *Fungal Biol Rev* **30**: 152–162.

Payen, T., Murat, C., Gigant, A., Morin, E., De Mita, S., and Martin, F. (2015) A survey of genome-wide single nucleotide polymorphisms through genome resequencing in the Périgord black truffle (*Tuber melanosporum* Vittad.). *Mol Ecol Resour* **15**: 1243–1255.

Peter, M., Kohler, A., Ohm, R.A., Kuo, A., Krützmann, J., Morin, E., et al. (2016) Ectomycorrhizal ecology is imprinted in the genome of the dominant symbiotic fungus *Cenococcum geophilum*. *Nat Commun* **7**: 12662.

Pfeifer, B., Wittelsbürger, U., Ramos-Onsins, S.E., and Lercher, M.J. (2014) PopGenome: An efficient Swiss army knife for population genomic analyses in R. *Mol Biol Evol* **31**: 1929–1936.

di Pietro, M., Churin, J.L., and Garbaye, J. (2007) Differential ability of ectomycorrhizas to survive drying. *Mycorrhiza* **17**: 547–550.

Pritchard, J.K., Stephens, M., and Donnelly, P. (2000) Inference of population structure using multilocus genotype data. *Genetics* **155**: 945–959.

Quinlan, A.R. and Hall, I.M. (2010) BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**: 841–842.

Raffaele, S. and Kamoun, S. (2012) Genome evolution in filamentous plant pathogens: Why bigger can be better. *Nat Rev Microbiol* **10**: 417–430.

Reinhardt, D., Roux, C., Corradi, N., and Di Pietro, A. (2021) Lineage-specific genes and cryptic sex: Parallels and differences between arbuscular mycorrhizal fungi and fungal pathogens. *Trends Plant Sci* **26**: 111–123.

Riley, R. and Corradi, N. (2013) Searching for clues of sexual reproduction in the genomes of arbuscular mycorrhizal fungi. *Fungal Ecol* **6**: 44–49.

Rineau, F. and Courty, P.E. (2011) Secreted enzymatic activities of ectomycorrhizal fungi as a case study of functional diversity and functional redundancy. *Ann For Sci* **68**: 69–80.

Rochet, J., Moreau, P.A., Manzi, S., and Gardes, M. (2011) Comparative phylogenies and host specialization in the alder ectomycorrhizal fungi Alnicola, Alpova and Lactarius (Basidiomycota) in Europe. *BMC Evol Biol* **11**: 1–14.

Sánchez-Ramírez, S., Tulloss, R.E., Guzmán-Dávalos, L., Cifuentes-Blanco, J., Valenzuela, R., Estrada-Torres, A., et al. (2015) In and out of refugia: Historical patterns of diversity and demography in the North American Caesar's mushroom species complex. *Mol Ecol* **24**: 5938–5956.

Serrato-Capuchina, A. and Matute, D.R. (2018) The role of transposable elements in speciation. *Genes* **9**: 1–29.

Simpson, M.C., Coetzee, M.P.A., van der Nest, M.A., Wingfield, M.J., and Wingfield, B.D.
(2018) Ceratocystidaceae exhibit high levels of recombination at the mating-type (*MAT*)
locus. *Fungal Biol* **122**: 1184–1191.

Smith, M.E., Henkel, T.W., and Rollins, J.A. (2015) How many fungi make sclerotia? *Fungal*
Ecol **13**: 211–220.

Spatafora, J.W., Owensby, C.A., Douhan, G.W., Boehm, E.W.A., and Schoch, C.L. (2012)
Phylogenetic placement of the ectomycorrhizal genus *Cenococcum* in Gloniaceae
(Dothideomycetes). *Mycologia* **104**: 758–765.

Tanaka, T. and Nei, M. (1989) Positive Darwinian selection observed at the variable-region
genes of immunoglobulins. *Mol Biol Evol* **6**: 447–459.

Trappe, J.M. (1962) *Cenococcum graniforme* - Its distribution, ecology, mycorrhiza formation,
and inherent variation.

Vélez, J., Morris, R., Vilgalys, R., Labbé, J., and Schadt, C. (2021) Phylogenetic diversity of
200+ isolates of the ectomycorrhizal fungus *Cenococcum geophilum* associated with
Populus trichocarpa soils in the Pacific Northwest, USA and comparison to globally
distributed representatives. *PLoS One* **16**: e0231367.

Wang, N.Y., Zhang, K., Huguet-Tapia, J.C., Rollins, J.A., and Dewdney, M.M. (2016) Mating
type and simple sequence repeat markers indicate a clonal population of *Phyllosticta*
citricarpa in Florida. *Phytopathology* **106**: 1300–1310.

Wang, X., Zheng, Z., Cai, Y., Chen, T., Li, C., Fu, W., and Jiang, Y. (2017) CNVcaller: Highly
efficient and widely applicable software for detecting copy number variations in large
populations. *Gigascience* **6**: 1–12.

Watanabe, M., Sato, H., Matsuzaki, H., Kobayashi, T., Sakagami, N., Maejima, Y., et al.
(2007) 14C ages and δ13C of sclerotium grains found in forest soils. *Soil Sci Plant Nutr*
53: 125–131.

White, T., Bruns, T., Lee, S., and Taylor, J. (1990) Amplification and direct sequencing of
fungal ribosomal RNA genes for phylogenetics. In *PCR protocols: a guide to methods*
and applications. pp. 315–322.

Wickham, H. (2011) ggplot2. *Wiley Interdiscip Rev Comput Stat* **3**: 180–185.

Zeiner, C.A., Purvine, S.O., Zink, E.M., Paša-Tolić, L., Chaput, D.L., Haridas, S., et al. (2016)
Comparative analysis of secretome profiles of manganese(II)-oxidizing Ascomycete
fungi. *PLoS One* **11**:.

Zhang, J., Kapli, P., Pavlidis, P., and Stamatakis, A. (2013) A general species delimitation
method with applications to phylogenetic placements. *Bioinformatics* **29**: 2869–2876.

Zheng, X., Levine, D., Shen, J., Gogarten, S.M., Laurie, C., and Weir, B.S. (2012) A high-
performance computing toolset for relatedness and principal component analysis of SNP
data. *Bioinformatics* **28**: 3326–3328.

Data Sharing and Data Availability

Raw sequence data used in this study are available in NCBI under the Sequence Read Archive identifiers listed in Table S1.

Competing Interests

The authors declare no competing interests.

Tables

924 **Table 1: Details of the *Cenococcum geophilum* individuals analysed in this study.**

Strain	*MAT1* locus	Isolate origin	Code	Site	Forest	Isolation year	Latitude [° N]	Longitude [° E]
1.104	*MAT1-2*	Finland	FIN	Punkaharju	*Pinus sylvestris*	20.08.08	61.800000	29.316667
1.076	*MAT1-1*	France	FRA	La Crête	*Fagus sylvatica*	1999	48.111111	5.356944
1.075	*MAT1-1*	France	FRA	La Crête	*Fagus sylvatica*	1999	48.111111	5.356944
1.100	*MAT1-1*	France	FRA	Ventoux	Mixed	30.07.09	44.179497	5.241850
1.094	*MAT1-1*	Poland	POL	Blizyn	Mixed	01.03.09	51.037222	20.869083
1.096	*MAT1-2*	Poland	POL	Blizyn	Mixed	13.08.09	51.037111	20.868694
1.097	*MAT1-1*	Poland	POL	Blizyn	Mixed	13.08.09	51.037222	20.870250
1.022	*MAT1-2*	Switzerland	CHE	Nessental	*Picea abies*	13.06.80	46.722500	8.296667
1.101	*MAT1-2*	Switzerland	CHE	Parabock	*Picea abies*	15.05.08	46.690960	7.236940
1.102	*MAT1-2*	Switzerland	CHE	Parabock	*Picea abies*	10.10.08	46.690960	7.236940
1.057	*MAT1-1*	Switzerland	CHE	Pfynwald	*Pinus sylvestris*	23.04.08	46.301389	7.611944
1.58	*MAT1-1*	Switzerland	CHE	Pfynwald	*Pinus sylvestris*	30.04.08	46.301389	7.611944
1.061	*MAT1-1*	Switzerland	CHE	Pfynwald	*Pinus sylvestris*	30.04.08	46.301389	7.611944
1.064	*MAT1-1*	Switzerland	CHE	Pfynwald	*Pinus sylvestris*	04.09.08	46.301389	7.611944
1.060	*MAT1-1*	Switzerland	CHE	Pfynwald	*Pinus sylvestris*	30.04.08	46.301389	7.611944
1.005	*MAT1-2*	Switzerland	CHE	Strada	*Picea abies*	03.11.77	46.767500	9.178889

927 **Table 2: Characteristics of the 20 top-candidate genes putatively under selection in the**
928 **ref. and non-ref. cluster based on Tajima's *D* statistics.** Estimation of the nucleotide
929 diversity per site is shown for each candidate gene.
930

Dataset	Protein ID	Predicted function	Multi-gene family			*D*	*π*
			Size (No. proteins in Cenge)	Size evolution (CAFE)	Conservation (in Cenge, Glost, Leppa)		
Ref. cluster	166793	Signal transduction mechanisms	4			-0.87	0.00851
	738677	Self/non-self recognition (NACHT domain)	66	expanded		-0.869	0.00765
	701966	Cytoskeleton	1			-0.869	0.00736
	486968	Self/non-self recognition (HET, Ankyrin)	97		over-represented	-0.868	0.01591
	700681	Translation, ribosomal structure and biogenesis	1		unique	-0.868	0.00427
	665335	Chromatin structure and dynamics	2		unique	-0.867	0.00925
	552403	Self/non-self recognition (NACHT domain)	56	expanded		-0.867	0.01231
	608632	Self/non-self recognition (NACHT domain)	14			-0.867	0.00503
	353610	Nucleotide transport and metabolism	2			-0.866	0.00933
	222924	-	1		unique	-0.866	0.00612
	653800	Cytoskeleton	6			-0.865	0.01017
	702114	Transcription	2			-0.865	0.00546
	605605	CAZyme (Glycoside Hydrolase 35)	2	contracted		-0.865	0.00269
	597082	Energy production and conversion	1			-0.864	0.00768
	672908	Chromatin structure and dynamics	1		unique	-0.864	0.00595
	487620	-	4			-0.864	0.00901
	674647	Transcription	1			-0.864	0.00227
	661861	RNA processing and modification	1			-0.863	0.00408
	637860	Replication, recombination and repair	1			-0.863	0.00517
	237852	-	1			-0.863	0.00686
Non-ref. cluster	675988	-	1			-1.971	0.00055
	144717	Secondary metabolism (DMAT cluster, decorating ABC transporter)	6	contracted		-1.955	0.00019
	682552	Methyltransferase	1			-1.884	0.00072
	343761	Protein binding	5			-1.853	0.00128
	650216	Secondary metabolism	5			-1.852	0.00049
	650873	Transcription	1			-1.849	0.00042
	674343	Secondary metabolism	9			-1.844	0.0003
	676801	Secondary metabolism	1			-1.832	0.00064
	698845	Cell wall/membrane/envelope biogenesis	1			-1.831	0.00064
	202303	Secondary metabolism (DMAT cluster, DMAT backbone gene)	3			-1.804	0.00098
	672500	CAZyme (LPMO AA11)	6			-1.803	0.0005
	691644	Amino acid transporter	33			-1.777	0.00094
	650097	-	1			-1.768	0.00048

671949	Posttranslational modification, protein turnover, chaperones	15	-1.767	0.00077
707138	Zinc finger, mRNA export	1	-1.766	0.00029
366609	RNA processing and modification	2	-1.763	0.00076
563055	RNA processing and modification	1	-1.734	0.00082
653324	-	1	-1.731	0.00032
706589	Secondary metabolism (Beta-lactamase, response to antibiotic)	4	-1.714	0.0006
663021	-	2	-1.709	0.00234

Note: Protein Id: JGI Cenge3 portal; Predicted function: KOG class (EuKaryotic Orthologous Groups) and other: detailed prediction see Table S4; Multi-gene family: JGI MCL Cluster Run *Cenococcum* + 50 genomes 03/21/14.399 (https://mycocosm.jgi.doe.gov/clm/run/cenococcum-and-50-genomes.399; Ohm *et al.*, 2012; Peter *et al.*, 2016) Conservation: protein orthologues only present (unique) or over-represented ($P \leq 0.05$; Wilcoxon test, two-sided, Benjamini-Hochberg multiple testing correction) in group Cenge (*Cenococcum geophilum*), Glost (*Glonium stellatum*), Leppa (*Lepidopterella palustris*) as compared to other Dothideomycetes (outgroup); Tajima's D (D); nucleotide diversity per site (π)

Figure 1: Study range, phylogenetic relationships, and genomic relatedness of the
Cenococcum geophilum **strains sampled. A**) Sampling sites of the *C. geophilum* isolates with
an illustration of the ecto-mycorrhiza morphology (Photo: M. Peter). Geographic distribution
of MAT loci is given for each sampled location. **B**) Phylogenetic relationships within the genus
Cenococcum based on two nuclear loci, i.e., ITS and GAPDH. One outgroup species, i.e.,
Glonium stellatum, was used to root the phylogenetic tree. The scale bar shows the substitution
rate. Tips in bold are the 16 samples analysed in this study using whole-genome sequencing
data. **C**) Genomic relatedness among the *C. geophilum* sampled, including the reference
genome (strain 1.58). Whole-genome SNP data capture both repetitive and coding regions
while synonymous SNPs only reflect neutral divergence in coding regions. Topologies are
rooted with *Glonium stellatum* as outgroup. Branch supports represent bootstrap values from
the maximum likelihood tree reconstruction. Strain codes are described in Table S1.

Figure 2: Recombination analysis and population genetic differentiation among the *Cenococcum geophilum* strains sampled. A) Network-based phylogenetic reconstruction supporting recombination events in one of the two main lineages. **B**) Principal component (PC) analysis showing the genomic variation without clear congruence with geography. **C**) Ancestry inferences for $K = 2$–5 indicating some admixed genotypes between the major genetic groups. Strain codes are described in Table S1. Colours refer to the geographic origin and genetic group of each strain for A–B and C, respectively.

Figure 3: Analysis of copy-number variation (CNV) regions among the sampled _Cenococcum geophilum_ strains. A) Gene content including CNV with the percentage of length covered by deletion or duplication >50%. Deletions refer to any gene locus showing at least one deletion in a strain while duplications include any gene locus showing only duplication or normal copy-numbers without deletion. **B**) Gene comparison affected by deletion or duplication among _C. geophilum_ strains. Tree topologies on the left and top show the grouping according to gene CNVs. The y-axis shows gene identifiers of CNV-affected genes. Note that the strain 1.58 displays only single-copy genes because the CNV calls were filtered to contain only those genes that were confirmed as single-copy from the reference genome. Strain codes are described in Table S1.

973 **Figure 4: Distribution of genomic distances between all genes and their closest
974 transposable elements (TEs), and between copy-number variation (CNV) genes and their
975 closest TEs.** The significance between the median values was calculated using the Wilcoxon
976 test. The red dots and blue squares represent mean and median values, respectively.

Figure 5: Genetic diversity analysis and coding sequence variation along scaffolds #1 and #2 in *Cenococcum geophilum* **strains used in this study. A)** and **C)** Nucleotide diversity per site is given for each gene identified in the 1.58 genome assembly. **B)** and **D)** Density of gene content and transposable element is presented. Results are shown for the complete dataset ("all") and subsets ("ref. cluster" and "non-ref. cluster") of *Cenococcum geophilum* strains used in this study (see section Experimental Procedures).

Figure 6: Within-species genetic diversity and signatures of selection. A) Distribution of nucleotide diversity per site for each gene. **B**) Distribution of gene-specific Tajima's *D*. **C**) Distribution of the gene-specific ratio of non-synonymous to synonymous polymorphisms (*p*N/*p*S). Results are shown for the complete dataset ("all") and subsets ("ref. cluster" and "non-ref. cluster") of *Cenococcum geophilum* strains used in this study (see section Experimental Procedures). Note that the reference cluster is genetically more diverse than the non-ref. cluster.

Figure 7: Chromosomal synteny analyses of *MAT1-1* and *MAT1-2* loci. Intraspecific analyses were performed with the reference genome strain (1.58) and interspecific comparisons were carried out between *G. stellatum*, *L. palustris* and *C. geophilum* strains. Colour gradients (white to red) correspond to sequence similarities ranging from 50–100%, with a minimum bit-score of 100.