Amazonian run-of-river dam reservoir impacts underestimated: Evidence from a Before-After Control-Impact study of freshwater turtle nesting areas

Andrea Bárcenas-García [1,2] | Fernanda Michalski [1,2,3] | James P. Gibbs[4] | Darren Norris[1,2,5] *

[1] Postgraduate Programme in Tropical Biodiversity, Federal University of Amapá, Macapá, Amapá, Brazil

[2] Ecology and Conservation of Amazonian Vertebrates Research Group, Federal University of Amapá, Macapá, Amapá, Brazil

[3] Instituto Pró-Carnívoros, Atibaia, São Paulo, Brazil

[4] Department of Environmental Biology, State University of New York, Syracuse, NY, USA

[5] Coordenação de Ciências Ambientais, Federal University of Amapá, Macapá, Amapá, Brazil

*Corresponding Author: Darren Norris, Email: dnorris75@gmail.com

Authors' Contribution statement (CRediT):

AB: Data Curation (supporting); Formal Analysis (lead); Investigation (supporting); Visualization (lead); Writing – original draft (lead); Writing – review and editing (equal)

FM: Conceptualization (equal); Data Curation (equal); Funding Acquisition (equal); Investigation (lead); Methodology (equal); Writing – review and editing (equal)

JPG: Conceptualization (equal); Funding Acquisition (equal); Writing – review and editing (equal)

DN: Conceptualization (equal); Funding Acquisition (equal); Methodology (equal); Investigation (supporting); Data Curation (equal); Formal Analysis (supporting); Visualization (supporting); Writing – original draft (supporting); Writing – review and editing (equal)

Abstract

1.   Construction of hydropower dams is associated with negative impacts on biodiversity, yet

there remains a lack of robust scientific evidence documenting the magnitude of these impacts

particularly across highly biodiverse tropical waterways. Hydropower expansion is an increasing

threat to the Endangered yellow-spotted river turtle (*Podocnemis unifilis*) across its tropical

South American range.

2.   Yellow-spotted river turtle nesting-areas were monitored as an indicator of dry-season

river level changes following run-of-river dam reservoir filling. A Before-After-Control-Impact

(BACI) study design was used with multi-year field campaigns monitoring turtle nesting-areas

along 66 km of river upstream of the dam, separated into 33 km control and impact zones.

3.   The cause and extent of changes in nesting-areas were established using Generalized

Additive Models. Nesting-area density was evaluated in relation to: time (before versus after),

treatment (control versus impact), time treatment interaction (BACI), distance to the dam and

precipitation. The extent of changes was examined by comparing the proportion of nesting-areas

remaining in subzones during the four years after reservoir filling.

4.   Dam construction generated a permanent dry season river level rise that extended more

than 20 km beyond impact assessment limits. On average the density of nesting-areas declined

69% (from 0.48 to 0.15 per km) across 33 km of river directly impacted by the dam. This loss

was reflected in a significant BACI interaction. Nesting-area density was not explained by

seasonal precipitation.

5.   Standardized monitoring of freshwater turtle nesting-areas provided an effective means to

quantify impacts of hydropower developments across biodiverse yet rapidly changing

waterways. The negative impacts documented in this study should be preventable by mitigation

actions including habitat restoration and dry season flow regulation. Such measures would also

likely benefit multiple species elsewhere in tropical rivers increasingly impacted by run-of-river

dams.

KEY WORDS: Amazon, Before-After-Control-Impact, conservation evidence, environmental

evidence, freshwater turtle, hydropower dam, nesting, *Podocnemis unifilis*, turtle conservation.

1 | INTRODUCTION

Mitigating dam impacts is a priority for the protection and restoration of freshwater biodiversity (Harper et al., 2021). Hydropower developments are among the most significant threats to Amazonian freshwater ecosystems (Castello & Macedo, 2016; Latrubesse et al., 2017) and biodiversity (Dudgeon, 2019; Latrubesse et al., 2020; Vasconcelos et al., 2020). Although hydropower supplies the demand of basic human necessities, hydropower dams result in impacts that produce negative effects and feedbacks that modify freshwater ecosystem processes, functions and services (Dudgeon, 2019; Valiente-Banuet et al., 2015). Generating changes not only to biological cycles and species ecological interactions (Latrubesse et al., 2020) but also creating drastic alterations to local communities, livelihoods and cultures (Del Bene, Scheidel, & Temper, 2018; Fearnside, 2018). Whilst negative impacts have been widely suggested, more robust evidence is required to examine the sustainability of the increasing number of dams across Brazilian Amazonia (Rodrigues dos Santos, Michalski, & Norris, 2021; Tundisi et al., 2014).

The ecological structure and complexity of Amazonian freshwater ecosystems is a result of myriad factors operating across different temporal and spatial scales, including flood pulse (Junk & Wantzen, 2006; Melack & Coe, 2021) and the spatial-temporal heterogeneity of seasonal rainfall (Melack & Coe, 2021; Reis et al., 2019; Siddiqui et al., 2021). The structure, composition, functions and services of globally important Amazonian ecosystems are threatened due to expansion of hydropower developments. To date a total of 29 dams (operational capacity > 30 MW) have been installed across Brazilian Amazonia (SIGEL, 2021) and another 750 are planned or under construction (Vasconcelos et al., 2020). This increase is incentivized by increasing energy demand, national and regional politics and international investment (Castello, 2021; Fearnside, 2018; Gerlak et al., 2020). Although hydropower dams require Environmental Impact Assessments, those conducted in the Brazilian Amazonia are typically characterized by localized, unstandardized and short-term biodiversity monitoring (Fearnside, 2014, 2018; Pelicice & Castello, 2021).

There is no standardized typology of dam types or sizes, which can often generate confusion in the scientific literature. For example the International Commission of Large Dams (https://www.icold-cigb.org/, accessed 9 July 2021) defines a large dam as: "A dam with a height of 15 meters or greater from lowest foundation to crest or a dam between 5 meters and 15 meters impounding more than 3 million cubic meters", but such definitions vary in different regional

and national contexts. "Run-of-river" dams are widely considered to generate relatively less

negative impacts compared with impoundment dams as the production of energy works with the

natural river flow (Gaudard, Avanzi, & De Michele, 2018). Yet, "run-of-river" is not

synonymous with small or low impact hydropower operations (Kuriqi et al., 2021), causing loss

of longitudinal connectivity, habitat degradation, and simplification of the biota community,

which leads to increasing need for more economically and environmentally sustainable options

such as in-stream turbines across Amazonia (Chaudhari et al., 2021).

Although adoption of newer technologies is likely to reduce environmental impacts in the

future (Chaudhari et al., 2021), currently a lack of evidence limits the development of effective

mitigating actions of Amazonian run-of-river hydropower dams (Gerlak et al., 2020; Gomes et

al., 2020). Run-of-river dams maybe constructed with or without reservoirs and range in size to

large "mega-dams" such as the Belo Monte dam complex in Brazil (total installed capacity of

11,233 MW, dam heights 36 – 90 m). The use of natural hydrological flows reduces the energy

production capacity of run-of-river dams during the dry season, which is a period when

supplying the energy demands can influence dry season river flow and freshwater habitats

(Gierszewski et al., 2020; Latrubesse et al., 2020). Indeed, although impacts of large hydropower

developments could lead to species extinction (Turvey et al., 2010) and extirpation in impacted

basins (Gomes et al., 2020; Santos et al., 2020) there remains a lack of information about the

impacts and responses of freshwater vertebrates to hydropower dams (He et al., 2018).

Aquatic species, like freshwater turtles are directly impacted by dam construction (Le Duc

et al., 2020; Stanford et al., 2020; Tucker, Guarino, & Priest, 2012). Although many studies

focus on fishes there are few studies considering the diversity of Amazonian aquatic and semi-

aquatic vertebrates (Rodrigues dos Santos, Michalski, & Norris, 2021). Turtles represent some of

the largest and most threatened freshwater vertebrates (Stanford et al., 2020) and are vulnerable

to habitat changes because their life history (Shine & Iverson, 1995) and dependence on

environmental factors for embryonic development, which limits the adaptive response for

population recruitment against anthropic impacts (Quintana et al., 2019). Little is known about

the impacts of hydropower dams on freshwater turtles at temporal and spatial scales relevant to

develop effective management actions. As freshwater turtles are typically long-lived there is a

need for long-term studies to detect life-history trends (Iverson, 1991; Shine & Iverson, 1995),

but there are relatively few studies available documenting tropical freshwater turtle survival (Rachmansah, Norris, & Gibbs, 2020).

The Endangered (Rhodin et al., 2018) yellow-spotted river turtle (*Podocnemis unifilis,* Figure 1) is the most widespread of six *Podocnemis* species found in South America, all of which nest on dry land (Turtle Taxonomy Working Group, 2017). Although widespread, *P. unifilis* is increasingly threatened (Rhodin et al., 2018) and without effective conservation actions severe ($\geq$50%) and rapid (<50 year) population losses were projected across 60% of the species 5.3 M km^2 pan-Amazonian range (Norris et al., 2019). Studies emphasize the need to focus on adult survival for the conservation of freshwater turtles (Páez et al., 2015; Rachmansah, Norris, & Gibbs, 2020), but early stage (e.g., egg and hatchling) is also important for the conservation and recovery of freshwater turtles (Rachmansah, Norris, & Gibbs, 2020) and *P. unifilis* populations (Norris et al., 2019). Nesting patterns are useful to understand responses of chelonian populations (Iverson, 1991) to environmental disturbance because nesting is key for female turtles to guarantee their survival and sustain populations (Refsnider & Janzen, 2010). Changes in dry season flood pulse and/or flow rates, could therefore result in negative impacts on *Podocnemis* populations due to changes in nest site selection and hatchling success (Eisemberg et al., 2016; Norris, Michalski, & Gibbs, 2020).

Figure 1. Study species. Adult and hatchlings of the yellow-spotted river turtle: (a) Adult females basking and (b) Hatchlings emerging from a nest laid in the sand of a riverside nesting-area.

Freshwater turtle nesting-areas are important for multiple Amazonian species not only for
the turtles themselves (Campos-Silva et al., 2018). The seasonal reduction in river flow exposes
river banks and islands, creating ephemeral habitats that represent essential seasonal resources
for both freshwater and terrestrial biodiversity across Amazonia (Castello, 2021; Junk &
Wantzen, 2006; Melack & Coe, 2021). These seasonal habitats are used not only by diverse
oviparous wildlife (Campos-Silva et al., 2018; Campos-Silva et al., 2021) and their predators (da
Costa Reis et al., 2021), but can also facilitate movements e.g. as stepping-stones to cross rivers
(Mamalis et al., 2018), and provide locations for reproductive/territorial behaviours (Michalski et
al., 2021).

In the present study *P. unifilis* nesting areas were monitored as a biologically relevant
indicator of river level changes following run-of-river dam reservoir filling. The objective was to
expand on previous results from the first year after construction of a large run-of river dam
(Norris, Michalski, & Gibbs, 2018), comparing annual numbers of turtle nesting-areas to
examine if upstream patterns in the submersion of nesting-areas remained consistent after four
years post-reservoir filling and to test the influence of potentially confounding variables
(seasonal precipitation). For the first time establishing robust causal evidence documenting the
extent of impacts of a run-of-river dam reservoir on an Endangered vertebrate species in the
Amazon. The study documents permanent dry season changes beyond established limits with
implications for global efforts to monitor free flowing rivers (Grill et al., 2019). The findings are
used to suggest important mitigation actions that are likely to benefit multiple species in highly
diverse Amazonian rivers increasingly impacted by run-of-river dams.

2 | MATERIALS & METHODS

2.1 | Study area

This study was conducted in the Araguari river basin, eastern equatorial Amazonia, in the central
region of the Brazilian State of Amapá (Figure 2). The basin includes rivers classified as "clear
water" (Junk et al., 2015). The Araguari River has an extension of 560 km (Ziesler & Ardizzone,
1979) and rises in the Guiana Shield at the base of the Tumucumaque uplands, bifurcates and
discharges into the Amazon River and the Atlantic Ocean. The region's climate is humid tropical
["Am" Tropical monsoon (Kottek et al., 2006)], with a dry season from September to November

(<150 mm of monthly rain) and the rainy season (>300 mm of monthly rain) from February to

April (Supporting Information S1).

There are currently three dams with a combined installed capacity of 549 MW (78, 252 and

219 MW) along an 18 km stretch of the Araguari River (ANEEL, 2020; EDP-Brasil, 2018;

INESC, 2017). Coaracy Nunes was the first dam installed along the Araguari River in 1975 and

was also the first dam in the Brazilian Amazon. The present study focuses on the 219 MW

Cachoeira Caldeirão dam (51°17'W, 0°51'N), which became operational in 2016 and is the most

recent and most upstream dam in the Araguari river. Cachoeira Caldeirão is a large run-of-river

dam, with a height of 20.6 m and length of 730 m (EDP-Brasil, 2018). It was installed and is

currently operated by the Empresa de Energia Cachoeira Caldeirão S.A, which is a joint venture

between China Three Gorges Brasil and Energias de Portugal- Brasil S.A (EDP-Brasil, 2018).

Upstream of the dam, the Araguari River flows between two sustainable-use conservation areas

(Figure 2): Amapá National Forest ("Floresta Nacional do Amapá" - FLONA) and the Amapá

State Forest ("Floresta Estadual do Amapá" - FLOTA).

Figure 2. Study area. **a**. State of Amapá in Brazil. **b**. Location of the study river (black line)
within Amapá. **c**. Location of the Impact (circles) and Control (triangles) subzones upstream of
the Cachoeira Caldeirão hydropower dam (square). Hashed shading is the area of direct impact
identified in the Environmental Impact Assessment. Grey shaded polygons show the protected
areas: Amapá National Forest ("Floresta Nacional do Amapá" - FLONA) and the Amapá State
Forest ("Floresta Estadual do Amapá" - FLOTA).

2.2 | Nesting-area surveys

Monthly boat surveys were conducted to locate nesting-areas from September to December, which corresponds to the nesting season of *P.unifilis* in the study area (Norris, Michalski, & Gibbs, 2018). Surveys were carried out upstream of the Cachoeira Caldeirão hydropower dam before (2011 and 2015) and after (2016, 2017, 2018, 2019) reservoir filling. Sampling and data collection in 2019 followed that from previous years (2011, 2015, 2016, 2017 and 2018) as described previously (Michalski et al., 2020; Norris, Michalski, & Gibbs, 2018; Norris, Michalski, & Gibbs, 2020; Quintana et al., 2019). Here the standardised sampling methods are briefly summarised. Areas suitable for turtle nesting (Figure 3) were located and identified based on the following features: areas >1m^2, with exposed sand/gravel substrates, raised above river level, without being waterlogged to a depth of 15 cm (Norris, Michalski, & Gibbs, 2018). For each nesting-area it was recorded whether the surface was exposed or submerged below water. Nesting-areas that were flooded subsequent to exposure/nesting were identified as flooded and not included as "submerged" i.e. submerged includes only those nesting-areas that were never exposed during the nesting season (September to December).

Figure 3 Nesting-areas. Representative yellow-spotted river turtle (*Podocnemis unifilis*) nesting-areas examples showing (a, b): potential nesting areas with suitable habitat conditions for nesting, but no nests (i.e., locations that females could use for nesting but nests were not found); (c, d): actual nesting areas (i.e., locations where females actually nested), and (e, f): unsuitable habitat areas for nesting (i.e., where substrate conditions were not appropriate for females to nest). Reproduced from Norris, Michalski, and Gibbs (2018).

To confirm the presence of turtle nests, searches were conducted by a team of three observers. One member was a local resident with over 30 years of knowledge of nesting-areas. Searches were made at a standardized walking speed (mean 0.8 km per hour, range 0.2-1.3) with time spent searching areas ranged from 10 to 97 minutes depending on the size of the area (Norris, Michalski, & Gibbs, 2018; Quintana et al., 2019). Nests were located by following turtle tracks on the substrates (sand/gravel) and systematic substrate searches with a wooden stick (Norris, Michalski, & Gibbs, 2018; Quintana et al., 2019). Nests that had been depredated were identified by the presence of broken eggshells outside the nest, disturbed/uncovered nests and the presence of excavation marks (Escalona & Fa, 1998). The presence of all nests was considered including those depredated by humans or wildlife.

2.3 | Explanatory variables

Distance from dam and precipitation were used to explain temporal and spatial patterns in nesting-area density. Distance to the hydropower dam was calculated along the river network using the riverdist package (Tyers, 2020) in R(R Core Team, 2020). Because rainfall is an important determinant of annual and seasonal variation in Amazonian river flow rates (Melack & Coe, 2021; Siddiqui et al., 2021) such that elevated precipitation during the months preceding nesting could result in nesting-areas remaining submerged. The total and standard deviation of precipitation (PPT) during each of the three months (July, August and September) prior to the peak of the nesting season in the six monitored years was calculated (Supporting Information S1). Precipitation data was obtained from a weather station (Serra do Navio, id: 8052000, 52°00'W, 0°53'N), located approximately 115 km upstream from the Cachoeira Caldeirão dam; data available in the virtual database of the Brazilian National Water Agency ("Agência Nacional de Águas": http://www.snirh.gov.br/hidroweb/serieshistoricas).

2.4 | Data Analysis

Two different datasets were analysed, one to establish the cause of changes in *P. unifilis* nesting-area density (Supporting Information S2) and another to quantify the extent of changes (Supporting Information S3).

2.4.1 | Cause of changes

To establish the cause of changes the response of actual nesting-area density was modelled using Generalized Additive Models [GAMs, (Marra & Wood, 2011; Pedersen et al., 2019)]. Actual nesting-areas were identified prior to reservoir filling as those where females nested in 2015 and/or in the previous five years [2010 – 2014, (Norris, Michalski, & Gibbs, 2018)]; post-reservoir filling the actual areas included those where females nested in the current nesting season and/or during the previous five years. To represent the Before-After-Control-Impact (BACI) sampling design (Smith, 2006; Underwood, 1993), the response of actual nesting-areas per km was modelled by including two fixed factors, (i) before-after and (ii) control-impact and their interaction, with a significant interaction indicating differences between the impacted compared with the control zone associated with reservoir filling (Underwood, 1993). In this study, the impacted zone (Figure 2) corresponds to 33 km of the Araguari River, within the upstream area of direct dam impact [as defined by the Environmental Impact Assessment (Ecotumucumaque, 2013; EDP energy, 2017)]. A 33 km stretch of the Falsino River 72 - 105 km upstream of the dam (Figure 2), which corresponds to a region with little anthropogenic disturbance was established as the control zone. For the GAM analysis each zone (control, impact) was divided into five subzones with an equal length of 6.6 km (Supporting Information S2).

Table 1. Hypotheses and variables used to explain the impacts of a run-of-river dam on yellow-spotted river turtle (*Podocnemis unifilis*) nesting-areas in the eastern Brazilian Amazon.

Category	Variable	Hypothesis	Description
Null	Intercept only		Response variable modelled without predictors.
Null	Random only		Includes years and subzone as random effects.
Environmental	Precipitation	Precipitation contributes to increase the river level, modifying the availability of nesting-areas for *P. unifilis*.	Continuous. Total and variation (SD) in monthly precipitation (July to September) before peak *P. unifilis* nesting.

Anthropic	Distance to the hydropower dam	Areas closer to the dam have a higher negative impact.	Continuous. Distance in kilometres from the hydropower dam.
Anthropic	Zone (Control-Impact)	The impacted zone will have reduced nesting-areas compared with the control zone.	Categorical with two levels. Impact Zone (Araguari River) and Control Zone (Falsino River).
Anthropic	When (Before-After)	After four years operating, the dam has affected the nesting patterns for the *P. unifilis.*	Categorical with two levels. Years before (2011 and 2015) and after (2016, 2017, 2018 and 2019) reservoir filling.
Anthropic	BACI (Before-After-Control-Impact)	Significant interaction effect means that differences in nesting patterns are a direct result of dam impact.	Categorical interaction. The river section and period variables interact to represent the Before-After-Control-Impact (BACI) sampling design.

Spatial autocorrelation was quantified using Mantel test and semi-variogram (Fortin &

Dale, 2014), which confirmed that each subzone could be considered as a spatially independent

sample unit (Supporting Information 4). An information theoretic model selection approach

(Burnham & Anderson, 2002) was applied to evaluate the alternative working hypothesis (Table

1). A set of seven models represented the working hypotheses to explain the patterns in nesting-

area density (Table 1). The five explanatory variables (total precipitation, standard deviation

precipitation, distance to the hydropower dam, before-after, control-impact and the BACI

interaction) were modelled separately to test their relationship with nesting-areas per kilometre.

All models included subzone and year as random effects to account for unexplained variation

that was not part of the sampling design and enable estimation of the mean and variance of the

global distribution of explanatory variables (Pedersen et al., 2019). An intercept only null model

(without predictors) and a random only model (subzone and year only) were also included for

comparisons. GAMs were run with Tweedie error distribution to examine which predictors

described changes in nesting-area density. Tweedie a probability distribution family that includes

the purely continuous normal and gamma distributions, the discrete Poisson distribution, and the

class of compound Poisson gamma distributions (Jorgensen, 1997), was used the default

Restricted Maximum Likelihood (REML) for all the models.

To identify support for the seven alternative candidate models two Information Criteria

were compared: Akaike Information Criterion corrected for small-sample sizes (AICc) and

Bayesian Information Criterion (BIC). Model weights were calculated representing the

likelihood of a given model, with evidence ratios constructed as the ratio of weights for the

alternative models being compared (Burnham & Anderson, 2002). Model selection was followed

by variable selection to establish the strength and direction of individual variables. Variable

selection was based on a full model (including distance, precipitation and random variables) that

was evaluated using a double penalty approach to determine which variables had the strongest

effect on the response variable (Marra & Wood, 2011). As preliminary analysis suggested

differences following reservoir filling, an interaction with the Before-After factor was included

in each of the smooth terms for distance to dam, total precipitation and SD precipitation in this

variable selection analysis.

2.4.2 | Extent of changes

The extent of changes in nesting-areas was examined by comparing the number of nesting-areas

remaining in subzones during the four years after reservoir filling (2016, 2017, 2018 and 2019).

As there was potential for new nesting-areas to emerge or to be lost due to natural changes in

river flow both actual and potential nesting-areas were included for this analysis. Potential areas

were those where environmental conditions matched those described in the literature and/or

those found at the nesting-areas from 2011, but without evidence or history of nest sites (Norris,

Michalski, & Gibbs, 2018). A nesting-area could be either potential or actual during the nesting

period. That is, if a potential area was identified and in a subsequent visit a nest was found then

the area was counted only as actual during the nesting season. Starting in 2015 a continuous

stretch of river was surveyed (completing the gap between impact and control zones initially

monitored in 2011), providing a total of 15 equally spaced subzones. From 2016 to 2020 the

proportion of remaining nesting-areas in each of the 15 subzones was calculated. The proportion

of remaining nesting-areas from 2016 to 2020 was then related to the distance of the subzone

midpoint from the dam using a GAM with quasibinomial error family. All the analysis were

performed in R 4.0.1(R Core Team, 2020), with the "mgcv" package to generate GAMs (Wood,

2020) and the "MuMIn" package (Barton, 2020) used for model selection.

3 | RESULTS

Figure 4. Nesting-areas before and after run-of-river dam construction. Observed patterns in
yellow-spotted river turtle nesting-areas in the eastern Brazilian Amazon. Density of *P. unifilis*
nesting-areas in nesting seasons before (2011, 2015) and after (2016 to 2019) dam reservoir
filling in two zones (control and impact). Points show the number of nesting-areas per kilometre
in five 6.6 km river subzones in each of the impact (triangles) and control (circles) zones. Dashed
lines join the yearly observations for each 6.6 km river subzone. Solid lines (black and grey) are
predictions from Generalized Additive Model. Vertical dashed line represents when the dam
reservoir was filled.

Nesting-area density decreased following dam construction, with the average density dropping

from 0.48 areas per km before to 0.15 areas per km after reservoir filling at the Impact zone

(Figure 4). Although Control subzones tended to have more nesting-areas (Figure 4), there was

variation in values between subzones and years, such that prior to reservoir filling there was no

significant difference in mean nesting-area density between Control and Impact zones (GAM, *P*

= 0.279). In the first year following reservoir filling nesting-area density declined to 0.06

nesting-areas per kilometre in the impacted zone, with mean values remaining low in all

following years (Figure 4). Although there was still some variation between years and subzones

(values in the Impact subzones ranged from 0 to 0.91 nesting-areas per km) there was a

significant difference (GAM, $P = 0.0061$) in the average of nesting-area density between Impact

and Control zones after reservoir filling. Indeed, following reservoir filling the observed Control

zone nesting-area density was five times greater than that found in the Impact zone (0.74 and

0.15 areas per km, Control and Impact zones respectively).

Table 2. Explaining patterns in yellow-spotted river turtle nesting in the eastern Brazilian

Amazon. Generalized Additive Models were used to explain patterns in turtle nesting-areas per

km of river from 2011 – 2019. Models ranked according to an Information criteria approach

using AICc (Akaike Information Criterion corrected for small sample sizes) and BIC (Bayesian

Informative Criterion). K= number of parameters, Δ =delta for respective criterion values, and *wi*

= the model selection weights. Models ordered by AICc values.

Hypothesis tested	Model construction	K	R^2_{adj}	Dev. Exp. (%)	AICc			BIC		
					AICc	Δ	*wi*	BIC	Δ	*wi*
BACI	Y ~ BA:CI + s(Year†)+s(Subzone†)	12	0.5	55.0	64.7	0.0	0.8	83.6	0.0	0.7
Before-After	Y~ BA + s(Year†)+s(Subzone†)	12	0.4	50.1	69.2	4.5	0.0	88.0	4.4	0.0
Random	Y~ s(Year†)+s(Subzone†)	12	0.4	48.0	71.0	6.2	0.0	89.7	6.0	0.0
Control-Impact	Y~ CI + s(Year†)+s(Subzone†)	11	0.4	44.6	71.3	6.5	0.0	89	5.3	0.0
Distance to dam	Y~ Dist. to dam + s(Year †)+s(Subzone †)	11	0.4	46.2	71.5	6.8	0.0	89.8	6.2	0.0
Precipitation	Y~ s(PPTtotal)+s(PPTsd) + s(Year †)+s(Subzone †)	12	0.4	49.0	71.6	6.8	0.0	90.7	7.0	0.0
Intercept	Y~ 1	3	0.0	0.0	95.3	30.6	0.0	101.2	17.5	0.0

s: Non-parametric smooth terms
† Random effects
R^2_{adj}: Adjusted R squared for the model
Dev. Exp. (%): Percent of total deviance explained

The model selection analysis indicated that the best model explaining patterns in nesting-

area density was the BACI (Table 2). All other models were only weakly supported (Table 2,

ΔAICc > 4 and *wi* = 0.0 for models ranked by both AICc and BIC). The model including only

random effects performed better compared with models including precipitation or distance to

dam, with the random model having greater deviance explained and lower information criteria

values (Table 2).

Table 3. Variable selection analysis. Smooth terms from a Generalized Additive Model used to

explain patterns in yellow-spotted river turtle nesting-areas per km of river from 2011 – 2019.

Non-parametric smooth terms	EDF		P-value	
	Before	After	Before	After
Distance to the dam	0.00	0.90	0.5951	0.0022
PPT total	0.00	0.48	0.7239	0.1769
PPT sd	0.84	0.00	0.0147	0.8809
Year†	0.00		0.7801	
Subzone†	6.47		0.0049	
R^2_{adj}	0.4			
Dev. Exp. (%)	55.8			

EDF: Estimated degrees of freedom for the model terms. Values close to zero indicate no

relationship with the response, close to 1 may suggest a linear relationship and values greater

than 1 suggest a non-linear relationship.

† Random effects

R^2_{adj}: Adjusted R squared from the model

Dev. Exp. (%): Percent of total deviance explained

A closer examination of the variables using a variable selection analysis indicated that

distance to the dam was only significant after reservoir filling (Table 3, *P* = 0.0022, Supporting

Information 5). There was no relationship between nesting-area density and total precipitation.

Prior to reservoir filling SD precipitation was weakly related to nesting-area density (Table 3, P

= 0.0147). Following reservoir filling there was no relationship between nesting-area density and

either of the measures of precipitation (Table 3, P = 0.1769 and 0.8809, total and SD

respectively).

Figure 5. Extent of reservoir impacts. Remaining nesting-areas following reservoir formation at the run-of-river Cachoeira Caldeirão Dam, Amapá, Brazil. (a) The proportion of nesting-areas remaining in four years following reservoir formation was calculated along 15 equally spaced river sections (graphs generated excluding one section without nesting-areas). Dashed vertical line represents the limit (upstream river confluence) used to derive free flowing river metrics (Grill et al., 2019) and the dotted vertical line represents the limit of direct impact defined by the Environmental Impact Assessment. Lines and shaded areas are mean values and 95% confidence intervals from Generalized Additive Models (formula = y ~ s(x, bs = "cs"), family = quasibinomial) that are added as a visual aid to illustrate trends in the values. Photos illustrating dry season river level and nesting-area (b) before (October 2015) and (c) after (October 2018) reservoir filling (51°36'16W, 0°53'6N, approximately 70 km upstream of the Cachoeira Caldeirão dam).

Comparison of nesting-areas post reservoir filling showed significant non-linear increase in the proportion of remaining areas further from the dam in all four years post-reservoir filling (GAM, EDF = 8.61, P < 0.0001, Figure 5). Impacts were at least 20 km beyond those established by the Environmental Impact Assessment and 30 km beyond those used to establish free flowing

river status (Figure 5). Within 50 km of the dam between 76 and 100% of nesting-areas were

submerged following reservoir filling, but more areas remained further from the dam (20 to 76 %

from 49 to 72 km). There was much less variation in the control zone, with the proportion of

remaining areas ranging from 87 to 100% in the five subzones beyond 75 km from the dam

(Figure 5). Four years after reservoir filling an average of 25 % of nesting-areas were submersed

75 km upstream of the dam some 20 km beyond recognized limits. This loss was at least double

that experienced in the five control subzones (mean 1 – 9 % of nesting-areas submersed, Figure

5).

4 | DISCUSSION

This study provided robust evidence of a reduction in *P. unifilis* nesting-areas caused by the

filling of a run-of-the-river dam reservoir in the Eastern Amazon. The availability of nesting-

areas was not related to seasonal precipitation, but was directly linked to the distance from the

hydropower dam. We suggest that mitigation actions should include habitat restoration and dry

season flow rate regulation to increase the availability of nesting-areas and avoid possible

negative impacts to *P. unifilis* populations at least within 70 km upstream of the dam.

The present study showed that the distance to the dam has implications for the nesting

patterns of *P. unifilis*. The density of nesting-areas in the Control zone remained stable, whereas

in the Impact zone the reduction in the density of nesting-areas was pronounced after reservoir

filling. Even after four years of operation, the nesting-areas of *P.unifilis* in the Impact zone were

not restored to levels registered for the same area in the period before dam reservoir filling

(Norris, Michalski, & Gibbs, 2018). The lack of nesting habitat may lead females to select poor

quality habitat (e.g. clay substrate) or areas with increased predation risk to nesting females or

nests/eggs (Quintana et al., 2019), which may contribute to increase the extinction vulnerability

of this species in the study region.

A permanent rise in dry season river levels may increase impacts of seasonal losses of

nesting-areas due to increasing occurrence of extreme flooding events across Amazonia

(Barichivich et al., 2018) that can have negative impacts on Amazonian freshwater turtle

populations that depend on terrestrial areas for nesting (Eisemberg et al., 2016; Norris,

Michalski, & Gibbs, 2020). In addition to flooding nests, total rainfall can impact the amount of

nesting habitat available or the availability of some other limiting resource, as demonstrated for

Batagur kachuga where it had a significant negative correlation with nest number and monsoon

magnitude (Sirsi et al., 2017). In contrast to Sirsi et al. (2017), the present study found that

seasonal precipitation was not related to the density of nesting areas over seven years of nesting-

area monitoring along more than 100 km of river.

The submersion of nesting-areas agrees with previous studies that suggest the impacts of

run-of-river dam reservoirs are likely underestimated (Norris, Michalski, & Gibbs, 2018). Run-

of-river dams are known to alter flow rates upstream of the reservoir, reducing flow rates which

can lead to increasing upstream river levels and water residence times (Almeida et al., 2019).

Although run-of-river dams are considered to generate hydropower with relatively reduced

impacts (Gaudard, Avanzi, & De Michele, 2018), due to highly seasonal rainfall and river flow

rates the vast majority of Amazonian run-of-river dams include reservoirs e.g. Belo Monte (Hall

& Branford, 2012) and can therefore generate drastic impacts on flowrates (Mendes et al., 2021).

The results of the present study suggest that the dry season extent of river level rise is

permanently at least 20 km beyond recognized limits. This finding that impacts extended beyond

those of the environmental impact assessment adds to the growing consensus regarding the link

between superficial impact assessments and a lack of social and environmental sustainability of

Amazonian hydropower developments (Fearnside, 2014, 2018; Gerlak et al., 2020).

In the case of the Cachoeira Caldeirão dam, the environmental impact assessment (EIA) of

reservoir impacts was based on simplistic predictions without any consideration of uncertainty or

seasonality. The spatial extent of impacts was predicted based on a single value for a "normal"

flood quota of 58.3 m, not considering the range to the estimated maximum flood quota of 59.6

m (Ecotumucumaque, 2013). As such, impacts beyond those indicated in the EIA are to be

expected and must be included in conservation management and mitigation actions. The

comparison with global scale estimates (Grill et al., 2019) suggests that the impact of dams could

be widely underestimated for biodiverse and complex tropical rivers. The analysis was based on

a single species and tropical rivers include innumerable biodiversity that remains to be evaluated.

Indeed, even for arguably the most intensely studied group (fishes) a recent review established a

lack of results documenting impacts generated by run-of-river dams in tropical regions (Turgeon

et al., 2021). Therefore, there is an urgent need for studies to generate more robust evidence

documenting the biological impacts of hydropower developments in tropical regions.

Quantifying changes in nesting-areas is important but additional studies needed to establish

direct link with species population viability. For example, boat surveys could also provide an

effective means of monitoring female populations during the nesting season as they tend to bask

on logs and rocks (Conway-Gómez, Reibel, & Mihiar, 2014; Coway-Gomez, 2007). Although

still incipient, studies show that boat surveys can be used to monitor freshwater turtle

abundances at a local scale [e.g. 6 km surrounding islands in Florida (Breininger et al., 2019)]

and more widely [e.g. *P. unifilis* populations along > 50 km of rivers in Bolivia (Coway-Gomez,

2007; Yapu-Alcázar et al., 2018) and Peru (Norris et al., 2011; Pitman et al., 2011)]. As *P.*

unifilis is widespread and ubiquitous across Amazonian river basins (Norris et al., 2019; Turtle

Taxonomy Working Group, 2017), integrating nesting-area surveys with river based population

monitoring could therefore provide robust empirical estimates to parameterize population

viability analysis in basins and rivers impacted by hydropower developments across Amazonia.

A robust understanding of hydropower development impacts is necessary to enable

development of effective minimization and mitigation actions. Among the short- and medium-

term mitigation measures that should be adopted are: i) creation of new turtle nesting habitat, ii)

improved dry season regulation of water discharge by the hydropower plant to enable upstream

river levels to drop during four months needed for turtle nesting and hatching (September,

October, November and December in the study region) and iii) integrated monitoring and

conservation of turtle populations, particularly with the local community. Although the impacts

and the recommendations listed above were shared with the operating company in 2018, years of

meetings and negotiations have resulted in no interventions or financial recompense by the

company to address the impacts that went beyond limits established in the Environmental Impact

Assessment. The lack of conservation and restoration projects materialising as mitigation actions

limits the ability of this run-of river dam to be considered as a sustainable source of energy

generation.

The results show that monitoring nesting-areas provides useful insight into the impacts of

dams. Considering the widespread distribution of *P. unifilis* such monitoring can be widely

applied to enable basin level evaluations of river level changes following dam construction.

Continuous and standardized monitoring of nesting areas can be implemented and strengthened

through the establishment of community management programs (Harju, Sirén, & Salo, 2018;

Rivera et al., 2021). Several studies have shown the benefits of this type of management focused

on the conservation of nesting areas and maintaining the recruitment of turtle populations

(Campos-Silva et al., 2018; Norris, Michalski, & Gibbs, 2020; Norris et al., 2019). In addition,

these actions bring conservation co- benefits for other species, for example migratory waterfowl,

which also use the turtles' nesting habitat (Campos-Silva et al., 2021). Similarly, local

communities can also obtain economic and food benefits (Harju, Sirén, & Salo, 2018; Rivera et

al., 2021).

CONCLUSION

Hydropower developments should adopt flow regulation schemes that are compatible with the

persistence of impacted species. Assessments of the spatial extents of hydropower development

impacts must consider relevant range of flood quota values in order to avoid underestimation of

impacts to local and regional biodiversity. Due to their social and cultural importance freshwater

turtles represent conservation "flagship" species across tropical South America. Monitoring

nesting-areas is straight forward and should be a regional/ basin wide focus to not only rapidly

identify impacts but also develop effective mitigation actions for biodiversity conservation in

tropical rivers impacted by hydropower dams. Any management program to be implemented

must be designed considering the social, economic and environmental characteristics of each

region, as well as having the participation of the local population, research groups, government

and companies in charge of dam management.

ACKNOWLEDGEMENTS

We are grateful to Alvino Pantoja and Gilberto Souza dos Santos for their invaluable assistance during field-work in 2019. We are also grateful to the field assistants, students and volunteers who have helped to collect data during the previous nesting seasons. We would like to thank the Universidade Federal do Amapá and the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) for providing logistical support; IBAMA for authorization (IBAMA/SISBIO permit 49632-1) to conduct research around FLONA.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

FUNDING

This study was funded by The United States National Academy of Sciences and the United States Agency for International Development through the Partnership for Enhanced Research (http://sites.nationalacademis.org/pga/peer/index.htm), award number AID-OAA-A11-00012 to Darren Norris, James P. Gibbs and Fernanda Michalski. Andrea Bárcenas was financed with a scholarship provided by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). FM receives a productivity scholarship from CNPq (Grant: Process 302806/2018-0) and was funded by CNPq (Grant: 403679/2016-8).

REFERENCES

Almeida, R. M., Hamilton, S. K., Rosi, E. J., Arantes, J. D., Barros, N., Boemer, G., et al. (2019). Limnological effects of a large Amazonian run-of-river dam on the main river and drowned tributary valleys. *Scientific Reports, 9*(1), 16846. doi:10.1038/s41598-019-53060-1

ANEEL. (2020). Listas de Usinas por estado/Municipio. Retrieved from https://app.powerbi.com/view?r=eyJrIjoiNjc4OGYyYjQtYWM2ZC00YjllLWJlYmEtYz dkNTQ1MTc1NjM2IiwidCI6IjQwZDZmOWI4LWVjYTctNDZhMi05MmQ0LWVhNG U5YzAxNzBlMSIsImMiOjR9

Barichivich, J., Gloor, E., Peylin, P., Brienen, R. J. W., Schöngart, J., Espinoza, J. C., et al. (2018). Recent intensification of Amazon flooding extremes driven by strengthened Walker circulation. *Science Advances, 4*(9), eaat8785. doi:https://doi.org/10.1126/sciadv.aat8785

Barton, K. (2020). MuMIn: Multi-model inference. (Version 1.43.17). Retrieved from https://cran.r-project.org/web/packages/MuMIn/MuMIn.pdf

Breininger, D. R., Stolen, E. D., Breininger, D. J., & Breininger, R. D. (2019). Sampling rare and elusive species: Florida east coast diamondback terrapin population abundance. *Ecosphere, 10*(8), e02824. doi:https://doi.org/10.1002/ecs2.2824

Burnham, K., & Anderson, D. (2002). *Model selection and multi-model inference* (2nd ed.). New York, NY.: Springer

Campos-Silva, J. V., Hawes, J. E., Andrade, P. C. M., & Peres, C. A. (2018). Unintended multispecies co-benefits of an Amazonian community-based conservation programme. *Nature Sustainability, 1*(11), 650-656. doi:https://doi.org/10.1038/s41893-018-0170-5

Campos-Silva, J. V., Peres, C. A., Hawes, J. E., Abrahams, M. I., Andrade, P. C. M., & Davenport, L. (2021). Community-based conservation with formal protection provides large collateral benefits to Amazonian migratory waterbirds. *PloS one, 16*(4), e0250022. doi:https://doi.org/10.1371/journal.pone.0250022

Castello, L. (2021). Science for conserving Amazon freshwater ecosystems. *Aquatic Conservation: Marine Freshwater Ecosystems, 31*(5), 999-1004. doi:https://doi.org/10.1002/aqc.3615

Castello, L., & Macedo, M. N. (2016). Large-scale degradation of Amazonian freshwater ecosystems. *Global change biology, 22*(3), 990-1007. doi:https://doi.org/10.1111/gcb.13173

Chaudhari, S., Brown, E., Quispe-Abad, R., Moran, E., Müller, N., & Pokhrel, Y. (2021). In-stream turbines for rethinking hydropower development in the Amazon basin. *Nature Sustainability*. doi:10.1038/s41893-021-00712-8

Conway-Gómez, K., Reibel, M., & Mihiar, C. (2014). A predictive model of yellow spotted river turtle (Podocnemis unifilis) encounter rates at basking sites in lowland eastern Bolivia. *Applied Geography, 53*, 332-340. doi:https://doi.org/10.1016/j.apgeog.2014.06.026

Coway-Gomez, K. (2007). Effects of human settlements on abundance of *Podocnemis unifilis* and *P. expansa* turtles in Nortbeastern Bolivia. *Chelonian Conservation and Biology, 6*(2), 199-205. doi:https://doi.org/10.2744/1071-8443(2007)6[199:EOHSOA]2.0.CO;2

da Costa Reis, M., Reintjes, J. M., Chen, D., & O'Donnell, S. (2021). Predation on nests of three species of Amazon River turtles (Podocnemis) by underground-foraging army ants (Labidus coecus). *Insectes Sociaux*. doi:https://doi.org/10.1007/s00040-021-00814-8

Del Bene, D., Scheidel, A., & Temper, L. (2018). More dams, more violence? A global analysis on resistances and repression around conflictive dams through co-produced knowledge. *Sustainability Science, 13*(3), 617-633. doi:https://doi.org/10.1007/s11625-018-0558-1

Dudgeon, D. (2019). Multiple threats imperil freshwater biodiversity in the Anthropocene. *Current Biology, 29*(19), R960-R967. doi:https://doi.org/10.1016/j.cub.2019.08.002

Ecotumucumaque. (2013). *Estudo de Impacto Ambiental: Analise Socioabiental Integrada, impactos e medidas, prognosticos e programas*. Retrieved from https://www.cachoeiracaldeirao.com.br/en/environment#eis-rima

EDP-Brasil. (2018). Cachoeira Caldeirão a usina. Retrieved from https://www.cachoeiracaldeirao.com.br/en/edp-cachoeira-caldeirao-power-plant

EDP energy. (2017). Cachoeira Caldeirão Technical data (Cachoeira Caldeirão Dados Técnicos). Retrieved from http://www.edp.com.br/geracao-renovaveis/geracao/amapa-para/uhe-cachoeira-caldeirao/a-usina/Paginas/dados-tecnicos.aspx

Eisemberg, C. C., Machado Balestra, R. A., Famelli, S., Pereira, F. F., Diniz Bernardes, V. C., & Vogt, R. c. (2016). Vulnerability of Giant South American Turtle (Podocnemis expansa) nesting habitat to climate-change-induced alterations to fluvial cycles. *Tropical Conservation Science, 9*(4), 1-12. doi:https://doi.org/10.1177/1940082916667139

Escalona, T., & Fa, J. E. (1998). Survival of nests of the terecay turtle (Podocnemis unifilis) in the Nichare-Tawadu rivers, Venezuela. *Journal of Zoology, 244*(2), 303-312. doi:https://doi.org/10.1111/j.1469-7998.1998.tb00034.x

Fearnside, P. M. (2014). Impacts of Brazil's Madeira River Dams: Unlearned lessons for hydroelectric development in Amazonia. *Environmental Science & Policy, 38*, 164-172. doi:https://doi.org/10.1016/j.envsci.2013.11.004

Fearnside, P. M. (2018). Challenges for sustainable development in Brazilian Amazonia. *Sustainable Development, 26*(2), 141-149. doi:https://doi.org/10.1002/sd.1725

Fortin, M.-J., & Dale, M. R. (2014). *Spatial analysis: a guide for ecologists*: Cambridge University Press.

Gaudard, L., Avanzi, F., & De Michele, C. (2018). Seasonal aspects of the energy-water nexus: The case of a run-of-the-river hydropower plant. *Applied Energy, 210*, 604-612. doi:http://dx.doi.org/10.1016/j.apenergy.2017.02.003

Gerlak, A. K., Saguier, M., Mills-Novoa, M., Fearnside, P. M., & Albrecht, T. R. (2020). Dams, Chinese investments, and EIAs: A race to the bottom in South America? *Ambio, 49*(1), 156-164. doi:https://doi.org/10.1007/s13280-018-01145-y

Gierszewski, P. J., Habel, M., Szmańda, J., & Luc, M. (2020). Evaluating effects of dam operation on flow regimes and riverbed adaptation to those changes. *Science of The Total Environment, 710*, 136202. doi:https://doi.org/10.1016/j.scitotenv.2019.136202

Gomes, C. J. P., Nepomuceno Salvador, G., Casarim, R., Santos Pompeu, P., Guedes Brito, M. F., Andrade, F. R., et al. (2020). Hydropower construction plans threaten the largest Brazilian national river. *Aquatic Conservation: Marine Freshwater Ecosystems, 30*(12), 2464-2465. doi:https://doi.org/10.1002/aqc.3485

Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., et al. (2019). Mapping the world's free-flowing rivers. *Nature, 569*(7755), 215-221. doi:https://doi.org/10.1038/s41586-019-1111-9

Hall, A., & Branford, S. (2012). Development, Dams and Dilma: the Saga of Belo Monte. *Critical Sociology, 38*(6), 851-862. doi:https://doi.org/10.1177/0896920512440712

Harju, E., Sirén, A. H., & Salo, M. (2018). Experiences from harvest-driven conservation:
Management of Amazonian river turtles as a common-pool resource. *Ambio, 47*(3), 327-
339. doi:https://doi.org/10.1007/s13280-017-0943-5
Harper, M., Mejbel, H. S., Longert, D., Abell, R., Beard, T. D., Bennett, J. R., et al. (2021).
Twenty-five essential research questions to inform the protection and restoration of
freshwater biodiversity. *Aquatic Conservation: Marine and Freshwater Ecosystems,*
n/a(n/a). doi:https://doi.org/10.1002/aqc.3634
He, F., Bremerich, V., Zarfl, C., Geldmann, J., Langhans, S. D., David, J. N. W., et al. (2018).
Freshwater megafauna diversity: patterns, status and threats. *Diversity Distributions,*
24(10), 1395-1404. doi:https://doi.org/10.1111/ddi.12780
INESC. (2017). Banco de dados de hidrelétricas na Amazônia, Cachoeira Caldeirão Retrieved
from http://amazonia.inesc.org.br/banco-de-dados-hidreletricas-na-amazonia/cachoeira-
calderao/
Iverson, J. B. (1991). Patterns of survivorship in turtles (order Testudines). *Journal of Zoology,*
69(2), 385-391. doi:https://doi.org/10.1139/z91-060
Jorgensen, B. (1997). *The theory of dispersion models*. . London, UK: CRC Press.
Junk, W. J., & Wantzen, K. M. (2006). Flood pulsing and the development and maintenance of
biodiversity in floodplains. In B. University of California Press (Ed.), *Ecology of*
freshwater estuarine wetlands. (pp. 407-435). Berkeley SN.
Junk, W. J., Wittmann, F., Schöngart, J., & Piedade, M. T. F. (2015). A classification of the
major habitats of Amazonian black-water river floodplains and a comparison with their
white-water counterparts. *Wetlands Ecology and Management, 23*, 677-693.
doi:https://doi.org/10.1007/s11273-015-9412-8
Kottek, M., Grieser, J., Beck, C., Rudolf, B., & Rubel, F. (2006). World Map of the Köppen-
Geiger climate classification updated. *Meteorologische Zeitschrift, 15*(3), 259-263.
doi:https://doi.org/10.1127/0941-2948/2006/0130
Kuriqi, A., Pinheiro, A. N., Sordo-Ward, A., Bejarano, M. D., & Garrote, L. (2021). Ecological
impacts of run-of-river hydropower plants—Current status and future prospects on the
brink of energy transition. *Renewable and Sustainable Energy Reviews, 142*, 110833.
doi:https://doi.org/10.1016/j.rser.2021.110833
Latrubesse, E. M., Arima, E. Y., Dunne, T., Park, E., Baker, V. R., d'Horta, F. M., et al. (2017).
Damming the rivers of the Amazon basin. *Nature, 546*, 363-369.
doi:https://doi.org/10.1038/nature22333
Latrubesse, E. M., d'Horta, F. M., Ribas, C. C., Wittmann, F., Zuanon, J., Park, E., et al. (2020).
Vulnerability of the biota in riverine and seasonally flooded habitats to damming of
Amazonian rivers. *Aquatic Conservation: Marine Freshwater Ecosystems*, 1-14.
doi:https://doi.org/10.1002/aqc.3424
Le Duc, O., Van, T. P., Leprince, B., Bordes, C., Tuan, A. N., Benansio, J. S., et al. (2020).
Fishers, dams, and the potential survival of the world's rarest turtle, Rafetus swinhoei, in
two river basins in northern Vietnam. *Aquatic Conservation: Marine and Freshwater*
Ecosystems, 30(6), 1074-1087. doi:https://doi.org/10.1002/aqc.3317
Mamalis, L., Sousa da Silva, S., Bezerra Valerio, A., Norris, D., & Michalski, F. (2018).
Stepping stones facilitate river crossings by Myrmecophaga tridactyla in the north-eastern
Brazilian Amazon. *Edentata, 19*, 30-34.
doi:https://doi.org/10.2305/IUCN.CH.2018.EDENTATA-19-1.3.en

Marra, G., & Wood, S. N. (2011). Practical variable selection for generalized additive models. *Computational Statistics Data Analysis, 55*(7), 2372-2387.

Melack, J. M., & Coe, M. T. (2021). Amazon floodplain hydrology and implications for aquatic conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems, 31*(5), 1029-1040. doi:https://doi.org/10.1002/aqc.3558

Mendes, Y. A., Oliveira, R. S., Montag, L. F. A., Andrade, M. C., Giarrizzo, T., Rocha, R. M., et al. (2021). Sedentary fish as indicators of changes in the river flow rate after impoundment. *Ecological Indicators, 125*, 107466. doi:https://doi.org/10.1016/j.ecolind.2021.107466

Michalski, F., Martins, C. B., Rheingantz, M. L., & Norris, D. (2021). New Scent Marking Behavior of Neotropical Otter (Lontra longicaudis) in the Eastern Brazilian Amazon. *IUCN Otter Specialist Group Bulletin, 38*(1), 28 -37.

Michalski, F., Norris, D., Quintana, I., Valerio, A., & Gibbs, J. P. (2020). Substrate influences human removal of freshwater turtle nests in the eastern Brazilian Amazon. *Scientific Reports, 10*. doi:https://doi.org/10.1038/s41598-020-65074-1

Norris, D., Michalski, F., & Gibbs, J. P. (2018). Beyond harm's reach? Submersion of river turtle nesting areas and implications for restoration actions after Amazon hydropower development. *Peerj, 6*, e4228. doi:https://doi.org/10.7717/peerj.4228

Norris, D., Michalski, F., & Gibbs, J. P. (2020). Community based actions save Yellow-spotted river turtle (Podocnemis unifilis) eggs and hatchlings flooded by rapid river level rises. *Peerj, 8*, e9921. doi:https://doi.org/10.7717/peerj.9921

Norris, D., Peres, C. A., Michalski, F., & Gibbs, J. P. (2019). Prospects for freshwater turtle population recovery are catalyzed by pan-Amazonian community-based management. *Biological Conservation, 233*, 51-60. doi:https://doi.org/10.1016/j.biocon.2019.02.022

Norris, D., Pitman, N. C. A., Martínez Gonzalez, J., Torres, E., Pinto, F., Collado, H., et al. (2011). Abiotic modulators of Podocnemis unifilis (Testudines: Podocnemididae) abundances in the Peruvian Amazon. *Zoologia (Curitiba), 28*, 343-350. doi:http://dx.doi.org/10.1590/S1984-46702011000300008

Páez, V. P., Lipman, A., Bock, B. C., & Heppell, S. S. (2015). A plea to redirect and evaluate conservation programs for South America's Podocnemidid River Turtles. *Chelonian Conservation and Biology, 14*(2), 205-216. doi:https://doi.org/10.2744/CCB-1122.1

Pedersen, E. J., Miller, D. L., Simpson, G. L., & Ross, N. (2019). Hierarchical generalized additive models in ecology: an introduction with mgcv. *Peerj, 7*, e6876. doi:https://doi.org/10.7717/peerj.6876

Pelicice, F. M., & Castello, L. (2021). A political tsunami hits Amazon conservation. *Aquatic Conservation: Marine Freshwater Ecosystems, 31*(5), 1221-1229. doi:https://doi.org/10.1002/aqc.3565

Pitman, N. C. A., Norris, D., Gonzalez, J. M., Torres, E., Pinto, F., Collado, H., et al. (2011). Four years of vertebrate monitoring on an upper Amazonian river. *Biodiversity and Conservation, 20*(4), 827-849. doi:https://doi.org/10.1007/s10531-010-9982-y

Quintana, I., Norris, D., Valerio, A., Becker, F., P Gibbs, J., & Michalski, F. (2019). Nest removal by humans creates an evolutionary trap for Amazonian freshwater turtles. *Journal of Zoology, 309*(2), 94-105. doi:https://doi.org/10.1111/jzo.12689

R Core Team. (2020). R: A Language and Environment for Statistical Computing (Version 4.0.1). Vienna, Austria: R Foundation for Statistical Computing. Retrieved from https://www.R-project.org/

Rachmansah, A., Norris, D., & Gibbs, J. P. (2020). Population dynamics and biological feasibility of sustainable harvesting as a conservation strategy for tropical and temperate freshwater turtles. *PloS one, 15*(2), e0229689. doi:https://doi.org/10.1371/journal.pone.0229689

Refsnider , J. M., & Janzen, F. J. (2010). Putting eggs in one basket: Ecological and evolutionary hypotheses for variation in oviposition-site choice. *Annual Review of Ecology, Evolution, and Systematics, 41*, 39-57. doi:https://doi.org/10.1146/annurev-ecolsys-102209-144712

Reis, V., Hermoso, V., Hamilton, S. K., Bunn, S. E., Fluet-Chouinard, E., Venables, B., et al. (2019). Characterizing seasonal dynamics of Amazonian wetlands for conservation and decision making. *Aquatic Conservation: Marine Freshwater Ecosystems, 29*(7), 1073-1082. doi:https://doi.org/10.1002/aqc.3051

Rhodin, A. G. I., Stanford, C. B., van Dijk, P. P., Eisemberg, C., Luiselli, L., Mittermeier, R. A., et al. (2018). Global conservation status of turtles and tortoises (order Testudines). Chelonian Conservation and Biology. *Chelonian Conservation and Biology, 17*, 135-161. doi:https://doi.org/10.2744/CCB-1348.1

Rivera, C. J., Macey, S. K., Blair, M. E., & Sterling, E. J. (2021). Assessing Ecological and Social Dimensions of Success in a Community-based Sustainable Harvest Program. *Environmental Management, 67*(4), 731-746. doi:https://doi.org/10.1007/s00267-021-01425-6

Rodrigues dos Santos, E., Michalski, F., & Norris, D. (2021). Understanding hydropower impacts on Amazonian wildlife is limited by a lack of robust evidence: results from a systematic review. *bioRxiv*, 2021.2007.2001.450737. doi:https://doi.org/10.1101/2021.07.01.450737

Santos, R. E., Pinto-Coelho, R. M., Drumond, M. A., Fonseca, R., & Zanchi, F. B. (2020). Damming Amazon Rivers: Environmental impacts of hydroelectric dams on Brazil's Madeira River according to local fishers' perception. *Ambio, 49*, 1612–1628. doi:https://doi.org/10.1007/s13280-020-01316-w

Shine, R., & Iverson, J. B. (1995). Patterns of survival, growth and maturation in turtles. *Oikos, 72*(3), 343-348. doi:https://doi.org/10.2307/3546119

Siddiqui, S. F., Zapata-Rios, X., Torres-Paguay, S., Encalada, A. C., Anderson, E. P., Allaire, M., et al. (2021). Classifying flow regimes of the Amazon basin. *Aquatic Conservation: Marine Freshwater Ecosystems, 31*(5), 1005-1028. doi:https://doi.org/10.1002/aqc.3582

SIGEL. (2021). Sistema de Informações Georreferenciadas do Setor Elétrico. Available from Agência Nacional de Energia Elétrica Sistema de Informações Georreferenciadas do Setor Elétrico Retrieved 30 March 2021, from ANEEL https://sigel.aneel.gov.br/Down/

Sirsi, S., Singh, S., Tripathi, A., McCracken, S. F., Forstner, M. R. J., & Horne, B. (2017). Variation in reproductive output of the red-crowned roofed turtle (Batagur kachuga) and the three-striped roofed turtle (Batagur dhongoka) in the Chambal River of North India. *Chelonian Conservation Biology, 16*(2), 203-214. doi:https://doi.org/10.2744/CCB-1236.1

Smith, E. P. (2006). BACI Design. In *Encyclopedia of Environmetrics*. Chichester, UK: John Wiley & Sons, Ltd.

Stanford, C. B., Iverson, J. B., Rhodin, A. G. J., van Dijk, P. P., Mittermeier, R. l. A., Kuchling, G., et al. (2020). Turtles and tortoises are in trouble. *Current Biology, 30*(12), R721-R735. doi:https://doi.org/10.1016/j.cub.2020.04.088

Tucker, A. D., Guarino, F., & Priest, T. E. (2012). Where lakes were once rivers: contrasts of freshwater turtle diets in dams and rivers of southeastern Queensland. *Chelonian Conservation Biology, 11*(1), 12-23. doi:https://doi.org/10.2744/CCB-0906.1

Tundisi, J. G., Goldemberg, J., Matsumura-Tundisi, T., & Saraiva, A. C. F. (2014). How many more dams in the Amazon? *Energy Policy, 74*, 703-708. doi:https://doi.org/10.1016/j.enpol.2014.07.013

Turgeon, K., Trottier, G., Turpin, C., Bulle, C., & Margni, M. (2021). Empirical characterization factors to be used in LCA and assessing the effects of hydropower on fish richness. *Ecological Indicators, 121*, 107047. doi:https://doi.org/10.1016/j.ecolind.2020.107047

Turtle Taxonomy Working Group. (2017). Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (8th Ed.). In A. G. J. Rhodin, J. B. Iverson, P. P. van Dijk, R. A. Saumure, K. A. Buhlmann, P. C. H. Pritchard, & R. A. Mittermeier (Eds.), Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs 7:1–292. doi:https://doi.org/10.3854/crm.7.checklist.atlas.v8.2017

Turvey, S. T., Barrett, L. A., Hart, T., Collen, B., Yujiang, H., Lei, Z., et al. (2010). Spatial and temporal extinction dynamics in a freshwater cetacean. *Proceedings of the Royal Society B: Biological Sciences, 277*(1697), 3139-3147. doi:10.1098/rspb.2010.0584

Tyers, M. (2020). Package 'riverdist' (Version 0.15.3).

Underwood, A. J. (1993). The mechanics of spatially replicated sampling programmes to detect environmental impacts in a variable world. *Australian Journal of Ecology, 18*(1), 99-116. doi:https://doi.org/10.1111/j.1442-9993.1993.tb00437.x

Valiente-Banuet, A., Aizen, M. A., Alcántara, J. M., Arroyo, J., Cocucci, A., Galetti, M., et al. (2015). Beyond species loss: the extinction of ecological interactions in a changing world. *Functional Ecology, 29*(3), 299-307. doi:https://doi.org/10.1111/1365-2435.12356

Vasconcelos, L. P., Alves, D. C., da Câmara, L. F., & Hahn, L. (2020). Dams in the Amazon: The importance of maintaining free-flowing tributaries for fish reproduction. *Aquatic Conservation: Marine Freshwater Ecosystems*, 1-11. doi:https://doi.org/10.1002/aqc.3465

Wood, S. N. (2020). mgcv: Mixed GAM Computation Vehicle with Automatic Smoothness Estimation (Version 1.8-34). Retrieved from https://cran.r-project.org/web/packages/mgcv/mgcv.pdf

Yapu-Alcázar, M., Tellería, L., Vaca, C., García, J., Arias, R., & Pacheco, L. F. (2018). Bases para el manejo de la tortuga de río Podocnemis unifilis en la Reserva de la Biosfera Estación Biológica Beni, Bolivia. *Ecología en Bolivia, 53*(1), 52-72.

Ziesler, R., & Ardizzone, G. D. (1979). Amazon River System. In The Inland waters of Latin America: Food and Agriculture Organization of the United Nations, COPESCAL, Doc. Téc./COPESCAL Tech.Pap. Retrieved from http://www.fao.org/docrep/008/ad770b/AD770B05.htm.

    Supporting Information

Supporting information 1. Precipitation. Historical series of precipitation obtained from the

Serra de Navio weather station (id: 8052000), available in the virtual database of the National

Water Agency/Agência Nacional de Águas (ANA-

http://www.snirh.gov.br/hidroweb/serieshistoricas).

Figure S1. Monthly trends in precipitation during the sampling years. Boxplots show means and

95% confidence limits estimated via nonparametric bootstrap.

811
812
813
814

815 Figure S1.2. Annual precipitation patterns during the sampling years. The lines and shaded areas
816 are the mean value and 95% confidence intervals from Generalized Additive Models [formula
817 $=y \sim s(x, k=10)$] that are added to illustrate trends in the values.
818

819
820
821

References

823 ANA. (2020). Hidroweb v3.1.1, Séries Históricas de Estações. Agência Nacional de Águas.
824 Available at:http://www.snirh.gov.br/hidroweb/serieshistoricas. [Accessed 16 September 2020].
825
826

Supporting information 2. Data used to establish cause of changes in nesting-area density
(nesting-areas per km of river).

Lat	Long	Zone	Subzone	When	Year	Areas_per_km	Precip_total	Precip_sd	Dist_dam_km
0.9434	-51.5948	control	1	before	2011	0.57	293.2	24.9	77.7
0.9722	-51.5964	control	2	before	2011	0.46	293.2	24.9	84.41
1.0123	-51.5942	control	3	before	2011	0.91	293.2	24.9	91.58
1.0438	-51.5636	control	4	before	2011	0.23	293.2	24.9	98.24
1.0577	-51.5241	control	5	before	2011	0.91	293.2	24.9	105.8
0.7529	-51.3660	impact	6	before	2011	0.46	293.2	24.9	16.96
0.7206	-51.4113	impact	7	before	2011	0.46	293.2	24.9	23.68
0.7257	-51.4567	impact	8	before	2011	0.68	293.2	24.9	30.24
0.7415	-51.4987	impact	9	before	2011	0.68	293.2	24.9	36.92
0.7257	-51.5515	impact	10	before	2011	0.00	293.2	24.9	43.61
0.9434	-51.5948	control	1	before	2015	0.68	53.1	4.8	77.7
0.9722	-51.5964	control	2	before	2015	0.68	53.1	4.8	84.41
1.0123	-51.5942	control	3	before	2015	1.60	53.1	4.8	91.58
1.0438	-51.5636	control	4	before	2015	0.46	53.1	4.8	98.24
1.0577	-51.5241	control	5	before	2015	0.57	53.1	4.8	105.8
0.7529	-51.3660	impact	6	before	2015	1.37	53.1	4.8	16.96
0.7206	-51.4113	impact	7	before	2015	0.23	53.1	4.8	23.68
0.7257	-51.4567	impact	8	before	2015	0.46	53.1	4.8	30.24
0.7415	-51.4987	impact	9	before	2015	0.46	53.1	4.8	36.92
0.7257	-51.5515	impact	10	before	2015	0.00	53.1	4.8	43.61
0.9434	-51.5948	control	1	after	2016	1.03	113.4	35.01	77.7
0.9722	-51.5964	control	2	after	2016	0.23	113.4	35.01	84.41
1.0123	-51.5942	control	3	after	2016	0.91	113.4	35.01	91.58
1.0438	-51.5636	control	4	after	2016	0.23	113.4	35.01	98.24
1.0577	-51.5241	control	5	after	2016	0.46	113.4	35.01	105.8
0.7529	-51.3660	impact	6	after	2016	0.11	113.4	35.01	16.96
0.7206	-51.4113	impact	7	after	2016	0.00	113.4	35.01	23.68
0.7257	-51.4567	impact	8	after	2016	0.23	113.4	35.01	30.24
0.7415	-51.4987	impact	9	after	2016	0.00	113.4	35.01	36.92
0.7257	-51.5515	impact	10	after	2016	0.00	113.4	35.01	43.61
0.9434	-51.5948	control	1	after	2017	1.36	289.1	32.53	77.7
0.9722	-51.5964	control	2	after	2017	0.46	289.1	32.53	84.41
1.0123	-51.5942	control	3	after	2017	0.46	289.1	32.53	91.58
1.0438	-51.5636	control	4	after	2017	0.76	289.1	32.53	98.24
1.0577	-51.5241	control	5	after	2017	1.08	289.1	32.53	105.8
0.7529	-51.3660	impact	6	after	2017	0.00	289.1	32.53	16.96
0.7206	-51.4113	impact	7	after	2017	0.91	289.1	32.53	23.68
0.7257	-51.4567	impact	8	after	2017	0.15	289.1	32.53	30.24

Lat	Long	Zone	Subzone	When	Year	Areas_per_km	Precip_total	Precip_sd	Dist_dam_km
0.7415	-51.4987	impact	9	after	2017	0.00	289.1	32.53	36.92
0.7257	-51.5515	impact	10	after	2017	0.00	289.1	32.53	43.61
0.9434	-51.5948	control	1	after	2018	1.21	234	60.52	77.7
0.9722	-51.5964	control	2	after	2018	0.46	234	60.52	84.41
1.0123	-51.5942	control	3	after	2018	0.30	234	60.52	91.58
1.0438	-51.5636	control	4	after	2018	0.76	234	60.52	98.24
1.0577	-51.5241	control	5	after	2018	0.92	234	60.52	105.8
0.7529	-51.3660	impact	6	after	2018	0.00	234	60.52	16.96
0.7206	-51.4113	impact	7	after	2018	0.61	234	60.52	23.68
0.7257	-51.4567	impact	8	after	2018	0.15	234	60.52	30.24
0.7415	-51.4987	impact	9	after	2018	0.15	234	60.52	36.92
0.7257	-51.5515	impact	10	after	2018	0.00	234	60.52	43.61
0.9434	-51.5948	control	1	after	2019	1.21	275	56.97	77.7
0.9722	-51.5964	control	2	after	2019	0.46	275	56.97	84.41
1.0123	-51.5942	control	3	after	2019	0.76	275	56.97	91.58
1.0438	-51.5636	control	4	after	2019	0.76	275	56.97	98.24
1.0577	-51.5241	control	5	after	2019	0.92	275	56.97	105.8
0.7529	-51.3660	impact	6	after	2019	0.15	275	56.97	16.96
0.7206	-51.4113	impact	7	after	2019	0.46	275	56.97	23.68
0.7257	-51.4567	impact	8	after	2019	0.00	275	56.97	30.24
0.7415	-51.4987	impact	9	after	2019	0.15	275	56.97	36.92
0.7257	-51.5515	impact	10	after	2019	0.00	275	56.97	43.61

830
831

832   **Supporting information 3.** Data used to establish extent of changes in nesting-areas, with
833   values of the proportion of nesting-areas remaining in subzones during the four years after
834   reservoir filling.
835

Zone	Year	dist_dam_km	area_remain_prop
impact	2016	17.0	NA
impact	2017	17.0	NA
impact	2018	17.0	NA
impact	2019	17.0	NA
impact	2016	23.7	0.10
impact	2017	23.7	0.10
impact	2018	23.7	0.10
impact	2019	23.7	0.10
impact	2016	30.2	0.24
impact	2017	30.2	0.24
impact	2018	30.2	0.24
impact	2019	30.2	0.24
impact	2016	36.9	0.13
impact	2017	36.9	0.13
impact	2018	36.9	0.13
impact	2019	36.9	0.13
impact	2016	43.6	0.00
impact	2017	43.6	0.00
impact	2018	43.6	0.00
impact	2019	43.6	0.00
intermediate	2016	49.3	0.20
intermediate	2017	49.3	0.20
intermediate	2018	49.3	0.20
intermediate	2019	49.3	0.20
intermediate	2016	55.0	0.50
intermediate	2017	55.0	0.50
intermediate	2018	55.0	0.50
intermediate	2019	55.0	0.50
intermediate	2016	60.7	0.43
intermediate	2017	60.7	0.43
intermediate	2018	60.7	0.43
intermediate	2019	60.7	0.43
intermediate	2016	66.4	0.75
intermediate	2017	66.4	0.50
intermediate	2018	66.4	0.50
intermediate	2019	66.4	0.25
intermediate	2016	72.1	0.77

Zone	Year	dist_dam_km	area_remain_prop
intermediate	2017	72.1	0.65
intermediate	2018	72.1	0.65
intermediate	2019	72.1	0.65
control	2016	77.7	1.00
control	2017	77.7	0.96
control	2018	77.7	0.96
control	2019	77.7	0.96
control	2016	84.4	1.00
control	2017	84.4	0.94
control	2018	84.4	0.94
control	2019	84.4	0.88
control	2016	91.6	1.00
control	2017	91.6	0.97
control	2018	91.6	0.97
control	2019	91.6	0.97
control	2016	98.2	1.00
control	2017	98.2	1.00
control	2018	98.2	1.00
control	2019	98.2	1.00
control	2016	105.8	1.00
control	2017	105.8	0.88
control	2018	105.8	0.88
control	2019	105.8	0.88

836
837

Supporting information 4. Spatial Autocorrelation analysis.
Semi-variograms were used to test for spatial autocorrelation (Fortin & Dale, 2014) in nesting-areas per
kilometre. The tests were performed in R.4.0.1 (R Development Core Team, 2021), using the "gstat"
package (Pebesma & Graeler, 2020) to calculate semi-variograms of the generalized additive model
residuals (Figure S4). The semi-variograms for all models suggest no autocorrelation as uncorrelated
residuals give a more or less flat semi-variogram (Fortin & Dale, 2014).
844
Figure S4. Semi-variograms from residuals of generalized additive models (see main text Table 2 and Table 3 for
model definitions and results respectively). Distances calculated from geographic coordinates (decimal degrees, 0.01
≈ 900m).

References

Fortin, M.-J., & Dale, M. R. (2014). Spatial analysis: a guide for ecologists: Cambridge
University Press.

Pebesma, E., & Graeler, B. (2020). gstat: Spatial and Spatio-Temporal Geostatistical Modelling,
Prediction and Simulation.

R Development Core Team. (2021). The R Project for Statistical Computing. 4.0.1. Available at:
https://www.r-project.org/. Vienna, Austria: R Foundation for Statistical Computer.

Supporting information 5. Effects on distance to the dam and precipitation on the density of
nesting-areas.
859

Effect of distance to the dam before and after reservoir filling.

Effect of total precipitation before and after reservoir filling.

Effect of standard deviation of precipitation before and after reservoir filling.

860
861
862
863