# Identification of an isoflavonoid transporter required for the nodule establishment of the *Rhizobium-Fabaceae* symbiotic interaction

Wanda Biała-Leonhard[1]†, Laura Zanin[2]†, Stefano Gottardi[2]†, Rita de Brito Francisco[3], Silvia Venuti[2], Fabio Valentinuzzi[2,4], Tanja Mimmo[4], Stefano Cesco[4], Barbara Bassin[3], Enrico Martinoia[3], Roberto Pinton[1], Michał Jasiński[1,5*], Nicola Tomasi[2*]

[1]Department of Plant Molecular Physiology, Institute of Bioorganic Chemistry, Polish Academy of Sciences, Noskowskiego 12/14, 61-704 Poznań, Poland.

[2]Dipartimento di Scienze Agro-Alimentari, Ambientali e Animali, University of Udine, Via delle Scienze 206, 33100 Udine, Italy.

[3]Institute of Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107, 8008 Zurich, Switzerland.

[4]Faculty of Science and Technology, Free University of Bozen Bolzano, Piazza Università 5, 39100 Bolzano, Italy.

[5]Department of Biochemistry and Biotechnology, Poznań University of Life Sciences, Dojazd 11, 60-637 Poznań, Poland.

†These authors contributed equally

*To whom correspondence should be addressed: Nicola Tomasi, Email: nicola.tomasi@uniud.it, Phone: +39 0432 558637; Michał Jasiński, Email: jasinski@ibch.poznan.pl , Phone: +48-61-852-85-03

## SUMMARY

Nitrogen (N) as well as Phosphorus (P) are key nutrients determining crop productivity. Legumes have developed strategies to overcome nutrient limitation by e.g., forming a symbiotic relationship with N-fixing *rhizobia* and the release of P-mobilizing exudates and are thus able to grow without supply of N or P fertilizers. The legume-rhizobial symbiosis starts with root release of isoflavonoids, that act as signaling molecules perceived by compatible bacteria. Subsequently, bacteria release nod factors, which induce signaling cascades allowing the formation of functional N-fixing nodules.

We report here the identification and functional characterization of a plasma membrane-localized MATE-type transporter (LaMATE2) involved in the release of genistein from white lupin roots.

The *LaMATE2* expression in the root is upregulated under N deficiency as well as low phosphate availability, two nutritional deficiencies that induce the release of this isoflavonoid. *LaMATE2* silencing reduced genistein efflux and even more the formation of symbiotic nodules, supporting the crucial role of LaMATE2 in isoflavonoid release and nodulation. Furthermore, silencing of LaMATE2 limited the P-solubilization activity of lupin root exudates. Transport assays in yeast vesicles demonstrated that LaMATE2 acts as a proton-driven isoflavonoid transporter.

## KEY WORDS

## INTRODUCTION

One of the major challenges of sustainable agriculture comprises the production of high-quality plant material with preservation of soil components and reduced application of chemical fertilizers without penalizing yield. Nitrogen (N) and phosphorus (P) are limiting nutrients in most natural soils (Tilman, 1999). High input of N-fertilizers is required to sustain crop growth in conventional agriculture. However, this feature may contaminate soils and groundwater and markedly contribute to the release of greenhouse gases (Tilman et al., 2001). Much of the P in soils is not available to plants due to its tendency to interact with calcium and magnesium salts or iron and aluminum oxides. It is mainly present as sparingly soluble rock phosphate, or it is immobilized in slowly mineralizable P-containing organic compounds, such as phytates. Phosphorous is a non-renewable resource which is mined at an increasing rate to meet the demand for fertilizers (Vance et al., 2003).

*Leguminous* plants (*Fabaceae*) such as soybean and lupin, have evolved several strategies to survive in low nutrient soils. In the case of P, in many ecosystems and, in particular, in acidic soils, the plant`s response consists mainly in the association with mycorrhizal fungi or the formation of particular root structures, such as cluster roots (Purnell, 1960; Neumann and Martinoia, 2002; Lambers et al., 2015). Cluster roots release huge amounts of exudates into the rhizosphere which are mainly composed of carboxylates and flavonoids. Flavonoids are involved both in the mobilization of nutrients and in the modulation of soil microbial activities (Cesco et al., 2010). The production and release of flavonoids, are also essential for the establishment of the symbiotic interaction between legumes and N-fixing bacteria such as *Ensifer*, *Bradyrhizobium* or *Mesorhizobium*, leading to the fixation of atmospheric N. Moreover it has been hypothesized that flavonoids are involved in the initiation

of the nodule through their action on the plant hormone auxin and

could thus play a developmental role in addition to their action as

nod gene regulators (Subramanian et al., 2006; Wasson et al., 2006;

Li et al., 2016).

The mutualistic fungal and bacterial symbionts are striking

examples of soil microorganisms that have successfully coevolved

with their hosts since plants adapted to terrestrial ecosystems. They

promote plant growth by facilitating the acquisition of scarce

nutrients. The most commonly established symbiosis in plants is the

mycorrhizal association, with 80–90% of all land-plant species able

to enter this interaction. Around 100 Mio years ago, certain

angiosperms evolved a bias toward the evolution of nodulation with

the so-called N-fixing soil bacteria. Among those angiosperms are

legumes (*Fabales*) and one non-legume genus, *Parasponia*

(*Cannabaceae*, *Rosales*) which can establish mutualistic symbioses

with *Rhizobia*, a polyphyletic group of proteobacteria and diverse

group of plants belonging to the orders *Fagales*, *Rosales*, and

*Cucurbitales* which can associate symbiotically with filamentous

actinobacteria of the genus *Frankia* (Martin et al., 2017). By

forming symbiotic associations, plants obtain mineral nutrients. In

turn, they supply the symbiont with organic compounds, sugars and

lipids in the case of mycorrhiza, mostly carboxylates to N-fixing

bacteria (Udvardi and Poole, 2013; Jiang et al., 2017; Luginbuehl et

al., 2017). The establishment of the symbiosis is a complex event

and requires coordinated regulation of the corresponding genes and

release of signaling molecules into the rhizosphere. For the legume-

*rhizobia* symbiosis, it is expected that a flavonoid transporter must

be present in the plasma membrane of root cells to release

isoflavonoids into the rhizosphere (Sugiyama et al., 2007). Up to

now, transporters for flavonoids have been mainly described at the

vacuolar membrane (Zhao, 2015). Furthermore, an ABC (ATP-

Binding Cassette) transporter from *Medicago* was shown to transport flavonoids. However, this transporter is localized in the vasculature (Biala et al., 2017). Using a biochemical approach Sugiyama and coworkers (2007) presented evidence that, in soybean, genistein transmembrane transport is mediated by an ABC-type transport system. But its contribution to genistein root release and the legume-*rhizobia* symbiosis establishment is still unclear.

Despite all the research performed on this symbiotic interaction, a transporter releasing flavonoids into the rhizosphere and initiating the first step of this symbiosis awaits its identification.

**MATERIALS AND METHODS**

**Plant growth and transformation**

White lupin seeds (*Lupinus albus* L. cv. Amiga; Südwestdeutsche Saatzucht, Rastatt, Germany) were soaked for 24 hours in aerated water and germinated on a plastic net placed at the surface of an aerated 0.5 mM $CaSO_4$ solution in a growth chamber at 25 °C in the dark. Thereafter, 7-day-old seedlings were transferred to a hydroponic system, containing a P-free nutrient solution (μM): 5000 $Ca(NO_3)_2$, 1250 $MgSO_4$, 1750 $K_2SO_4$, 250 KCl, 20 Fe(III)EDTA, 25 $H_3BO_4$, 1.25 $MnSO_4$, 1.5 $ZnSO_4$, 0.5 $CuSO_4$, 0.025 $(NH_4)_6Mo_7O_4$. Phosphorus-deficient plants were grown on P-free nutrient solution, while 0.25 mM $KH_2PO_4$ were added to nutrient solution for P-sufficient condition. Plants were grown under controlled conditions for 4 weeks (day/night photoperiod, 16/8 h; radiation, 220 $\mu E\ m^{-2}s^{-1}$; day/night temperature, 25/20 °C; relative humidity, 70-80%).

In P-deficiency, lupin plants modify the root architecture developing particularly root structures, called cluster roots or proteoid roots. In order to differentiate the developmental stages of root clusters (Figure S1), the root system was immerged in a pH-indicator solution (0.04% w/v bromocresol purple). Depending on their morphology and capability to acidify the solution, at the end of the growing period different regions of cluster root were sampled from P-deficient plants (juvenile, immature, mature, senescent), as described by Massonneau and coworkers (2001). Root apices and cluster root parts were sampled from a pool of 12-16 P-sufficient and P-deficient plants. The samples were immediately frozen in liquid nitrogen for the RNA extraction or were rinsed twice in 0.5 mM $CaSO_4$ solution and the root exudates were collected for one hour in 0.5 mM $CaSO_4$ 10 mM 2-[N-Morpholino]ethanesulfonic acid (MES)-KOH pH 6.0 at a ratio 1:10 W/V. The samples were conserved at –80 °C until processing. Three independent experiments were performed.

To investigate the plant response to N-deficiency, white lupin seeds were grown for two weeks on watered Whatman paper in Petri dishes and then transferred to the magenta boxes containing PFR N-free solution described by Stróżycki and coworkers (2003). For N-sufficient conditions $KNO_3$ (0.1 mM), $NH_4H_2PO_4$ (5 mM), $Ca(NO_3)_2$ (2.46 mM) were added to the PFR medium. Plants were grown for further two weeks under controlled conditions (day/night photoperiod, 16/8 h; radiation, 220 $\mu E\ m^{-2}s^{-1}$; day/night temperature, 23/20 °C; relative humidity, 70-80 %). The samples were collected at 7 and 14 days after the transfer in N-deficient condition. Two biological repetition and 4/8 technical repetition were performed.

White lupin seedlings were transformed using *Agrobacterium*

*rhizogenes* ARqua1 strain (Quandt et al., 1993) carrying binary

vector pRedRoot::*LaMATE2 RNAi* or empty vector (EV) pRedRoot.

After germination 5 mm root tips were removed from radicles. The

sectioned surface was coated with *A. rhizogenes* and seedlings were

placed on solid Fahraeus medium supplemented with kanamycin (15

mg $L^{-1}$).

Nitrogen-deficient plants were grown and analyzed as described

above. For P-deficiency, after 3 weeks post germination, plants were

transferred to P-free nutrient solution (as described above). After 2

weeks, white lupin plants were moved in hydroponic solutions in P-

free nutrient solution and grown for 3 additional weeks. *LaMATE2*

expression analyses were performed, and root release was collected

as described previously on the fully developed cluster root

(immature, mature). Six biological replicates were performed for

each sample.

**Nodulation and effects of genistein exogenous addition**

Three-week-old lupin composite plants were transferred to the pots

filled with sterile perlite (0.75 L). Plants were nourished with PFR

N-free solution. After 2 week of N-deficiency plants were

inoculated with *B. japonicum* (strain UPP 133, (Stępkowski et al.,

2011)) and grown for further two weeks under controlled conditions

(day/night photoperiod, 16/8 h; radiation, 220 $\mu E\ m^{-2}s^{-1}$; day/night

temperature, 23/20 °C; relative humidity, 70-80 %). For

complementation experiments 1 h before inoculation with *B.*

*japonicum* plants were additionally supplemented with 1 μM

genistein solution. Fourteen-day post-inoculation plants were

removed from pots, nodules were counted for each single root, and

roots were collected for each plant. Afterwards all roots of single

plant were grounded and divided into aliquots dedicated for gene
expression and metabolomic analysis. Only plants revealing
expression of marker gene encoding fluorescent protein DsRed were
considered during analysis. These experiments were performed  6,
8 or 21.

**Mobilization of P from vivianite**

The vivianite suspension ($Fe_3(PO_4)_2 \cdot 8H_2O$) was prepared by mixing
$H_3PO_4$ with $FeSO_4 \cdot$ as described by Eynard et al. (1992). A solution
of 0.035 M $H_3{}^{32}PO_4$ and 0.05 M $FeSO_4 \cdot 7H_2O$ were brought to pH
6.0 with 0.05 M KOH under stirring. A bluish suspension of
vivianite was obtained, after precipitation via centrifugation (8000
g, 10 min), the precipitate was washed (by 3 successive
centrifugations and decantations) with deionized water.
Mobilization of $PO_4{}^{2-}$ from Vivianite ($Fe_3(PO_4)_2 \cdot 8H_2O$) by exudates
after purification with the SPE methods, in order to remove
carboxylates, from *LaMATE2 RNAi* or empty vector (EV) pRedRoot
transformed P-deficient roots was performed as followed: one mL
of a suspension containing ${}^{32}P$-vivianite (70 µmol $PO_4$, specific
activity 60 KBq µmol${}^{-1}$ P) was transferred into a dialysis tube
(ZelluTrans/Roth 6.0, $\varnothing$ 16mm, exclusion limit of 4÷6 kDa, ROTH)
and mixed with 5 mL of mobilization solution [0.5mM $CaSO_4$,
20mM Mes-KOH (pH 6.0), 50 µl methanol]. Thereafter, the dialysis
tube was transferred into continuously mixed 34 ml of mobilization
solution. At the beginning of the experiment, exudates
corresponding to the amount release for 1 hour by 1 gram of juvenile
cluster root tissues (Figure 1) were added to the external solution.
After 15, 30, 45 and 60 min, samples from the solution outside the
dialysis tube were collected and the amount of ${}^{32}P$ was measured by
liquid scintillation counting. The ${}^{32}P$ mobilization was estimated
from the difference between the ${}^{32}P$ concentration measured in the

presence of the mock and in the absence of exudates and was expressed as µmol P h$^{-1}$.

**Extraction and LC/MS analysis**

Frozen root tissue (500 mg) was grounded and extracted with 80% methanol. Root exudates were extracted from the medium by SPE (Solid Phase Extraction) method using octadecylosilane matrix and methanol according to Staszków and coworkers (2011). Luteolin was used as an internal standard. Samples were analyzed by liquid chromatography-electrospray ionization tandem mass spectrometry (LC/ESI/MS) using a Waters UPLC coupled with Bruker micrOTOF-Q mass spectrometer. The analysis was performed in a gradient mobile phase consisting of 0.5% formic acid (v/v) in water (A) and 0.5% formic acid (v/v) in acetonitrile (B). The m/z range of the recorded spectra was 50-1000. Analyses were performed in the ion-positive mode.

**RNA extraction and cDNA synthesis**

RNA extractions were performed using the InviTrap Spin Plant RNA Mini Kit (Stratec Molecular, Berlin, Germany) following manufacturer's instructions and contaminant genomic DNA was removed using 10 U of DNase I (GE Healthcare, Munich, Germany). The quantity and the quality of RNA was checked using a spectrophotometer, followed by a migration in a 1 % agarose gel. One microgram of total RNA for each sample was retro-transcribed using 1 pmol Oligo d(T)23 (Sigma Aldrich, Saint Louis, USA) and 10 U M-MulV RNase H (Finnzymes, Helsinki, Finland) following manufacturers' instruction.

**Isolation of the *LaMATE2* sequence**

269 The partial sequence was isolated via a cDNA-AFLP approach
270 starting from RNA extracted from juvenile cluster roots which were
271 compared to mature and senescent cluster roots, for details see
272 Massonneau and coworkers (2001). The full Open Reading Frame
273 (ORF) of *LaMATE2* (*LaMATE2*$_{ORF}$) was isolated from the cDNA of
274 juvenile cluster root tissues from P-deficient plants using the 5'/3'
275 RACE Kit (2nd generation, Roche Diagnostics S.p.a., Monza, Italy)
276 following the manufacturer's instructions. *LaMATE2*$_{ORF}$ sequence
277 was cloned in pGEM-T easy vector (for primers see Table S1,
278 Promega Italia Srl, Milan, Italy) and deposited in the NCBI database
279 (KY464927).

281 **Gene expression analysis**

282 The reaction was performed by adding 0.1 µl of cDNA to
283 RT complete reaction mix, Fluocycle$^{TM}$ sybr green (20-µl final
284 volume; Euroclone, Pero, Italy). Specific primers were designed for
285 the target, the two closest homologues ([www.whitelupin.fr](www.whitelupin.fr);
286 Hufnagel *et al.*, 2020) and the housekeeping gene using Primer3
287 software (Koressaar and Remm, 2007; Untergasser et al., 2012) and
288 were synthesized by Sigma Aldrich (Table S1). Gene expression
289 analyses were performed using CFX96$^{TM}$ Real-Time System
290 (C1000TM Thermal Cycler, BioRad) and CFX Manager™
291 Software (v 2.0, BioRad). The two closest homologues
292 (*Lalb_Chr06g0165721, Lalb_Chr02g0145611*) have a low
293 expression in the root tissues and their expression levels is shown in
294 Figure S5. Data were normalized in respect to the transcript level of
295 the housekeeping gene (*Ubiquitin* gene, *LaUBI*) using the $2^{-\Delta\Delta CT}$
296 method (Livak and Schmittgen, 2001). The efficiency of
297 amplification was calculated using R program (version 2.9.0;
298 http://www.r-project.org/) with the qPCR package (version 1.1-8),

299  following the authors' indications (Ritz and Spiess, 2008). Six
300  biological replicates were performed for each sample.
301
**Genistein transport by LaMATE2 in *Saccharomyces cerevisiae***
**vesicles**
The construction of the yeast expression vector pNEV (Sauer and
Stolz, 1994) containing $LaMATE2_{ORF}$ was performed by amplifying
$LaMATE2_{ORF}$ sequence with primers having both a *NotI*-restriction
sites at the 5'end. The cloning of the PCR product ($LaMATE2_{ORF}$)
was performed into the *NotI* site of pNEV (pNEV::$LaMATE2_{ORF}$),
the orientation was verified by sequencing. The transformation of
competent yeast cells (*S. cerevisiae* YPH499 strain) was performed
following a standard procedure (Gietz and Woods, 2002) and
transformants were selected on synthetic dextrose minimal medium
lacking uracil (SD-Ura medium; Burke *et al.*, 2000).
The yeast cells, strain YPH499, were transformed with the pNEV
empty vector or pNEV::LaMATE2$_{ORF}$ and selected on liquid-SD
Ura-medium. Thereafter, cells were incubated in YPD medium for
30 min, collected by centrifugation, and digested with lyticase
(1,000 U $g^{-1}$ fresh weight cells; Sigma Aldrich), and subsequently
microsomal vesicles were isolated as described by Klein and
coworkers (2002). Transport assays were performed to study the
genistein transport using the rapid filtration technique with
nitrocellulose filters (0.45 µm pore size; Millipore, Millipore Co.,
Bedford, USA). The transport experiment was carried out in
presence of isolated vesicles, transport buffer (0.4 M glycerol, 0.1
M KCl, 1 mM DTT, 1 mM EDTA, 5 mM ATP, 10 mM MgCl$_2$, 10
mM creatine phosphate, 0.1 mg ml$^{-1}$ creatine kinase, 20 mM Tris-
MES pH 7.4) and 5 µM of labelled $^3$H-genistein (American
Radiolabeled Chemicals, Saint Louis, USA; 1850 Bq filter$^{-1}$). Only
for the kinetic experiments, the $^3$H-genistein concentration ranged

from 5 µM up to 100 µM (5, 7, 10, 15, 33, 50 and 100 µM $^3$H-genistein) and the incubation time was 30 s. The kinetic parameters of $^3$H-genistein uptake were calculated by subtracting uptake rates recorded in the empty-vector vesicles. The kinetic parameters of $^3$H-genistein uptake were calculated between 7-100 µM by subtracting uptake rates recorded in the empty-vector vesicles using the Hanes–Woolf plot.

To test the dependence of genistein transport on a proton gradient, 25 mM NH$_4$Cl was included in the transport buffer assay and the incubation time ranging between 15 and 120 s (15, 30, 60 and 120 s). Moreover, the capability of LaMATE2 to mediate the uptake of different flavonoids was tested by UPLC under the same experimental conditions reported above (5 µM of flavonoid: genistein, genistin, hydroxygenistein, biochanin A, daidzein, or kaempferol; the incubation time was 30 s). The mixture was loaded on a pre-wetted filter and removed by suction at the end of incubation time. The membranes were rapidly washed twice with 2 ml of ice-cold transport buffer.

The radioactive measurements were determined with a beta-counter (Tri-Carb 1900CA, Packard, Downers Grove, USA). As standards, solutions with known amounts of $^3$H-genistein were used. Vesicle protein content was quantified with BioRad Protein Assay Dye Reagent (BioRad, Hercules, CA, USA), and the data are shown as pmol $^3$H-genistein µg$^{-1}$ protein. Data are shown as net pmol $^3$H-genistein µg$^{-1}$ protein after removing background, i.e. unspecific adsorption of genistein onto empty-vector yeast membrane. Three independent transformations were performed for each sample.

**LaMATE2 subcellular localization in *Arabidopsis thaliana* protoplasts**

For transient expression of *LaMATE2*~ORF~ in Arabidopsis protoplasts,

the plasmid harboring the sequence for the *Green Fluorescent*

*Protein* (*GFP*) was fused at the C-terminus of *LaMATE2*~ORF~ inside

the pUC18-Sp-GFP6 vector (Komarova et al., 2008) using *NheI* and

*SphI* restriction sites via PCR amplification. A plasmid harboring

the sequence for *mCherry-fluorescent protein* was fused with

*AtPIP2a*, a gene coding for an aquaporin used as a plasma

membrane marker (Nelson et al., 2007). Arabidopsis protoplasts

were co-transformed with both constructs, *LaMATE2*~ORF~-*GFP* and

*AtPIP2a-mCherry*, using the polyethylene glycol method (Jin et al.,

2001). Protoplasts were examined with a TCS SP5 confocal

microscope (Leica Microsystems, Wetzlar, Germany), excited with

an argon laser at 458 nm for GFP and 540–552 nm for mCherry (for

GFP: excitation BP458, beamsplitter FT500, emission BP 492-511

nm; for mCherry: excitation BP 540–552, beamsplitter FT560,

emission BP 575–640).

**RNAi-based silencing of *LaMATE2* gene in white lupin roots.**

*LaMATE2* silencing was adapted from Uhde-Stone and coworkers

(2005) using binary transformation vectors, pRNAi and pRedRoot

(Limpens et al., 2004). The target region (350 bp long, covering the

5' end of *LaMATE2*~ORF~) was first amplified with PCR, and

subsequently cloned into pRNAi vector between the restriction sites

*NcoI–SwaI* and *BamHI–SpeI*. The cloned sequence (*LaMATE2*

*RNAi*) was regulated by a double CaMV35s promoter and OCS-3'

terminator. Using the *KpnI–PacI* restriction sites, the *LaMATE2*

*RNAi* cassette from the previously produced in pRNAi vector was

transfered into the pRedRoot binary vector (pRedRoot::*LaMATE2*

*RNAi*).

**Phylogenetic and Statistical Analyses**

Phylogenetic analyses were conducted using MEGA software, version 6 (Tamura et al., 2013). The tree was constructed by aligning the protein sequences by Clustal-W and the evolutionary history was inferred using the Neighbor-Joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown in Fig. **S1** next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Poisson correction method and are in the units of the number of amino acid substitutions per site.

The topology prediction of LaMATE2 was performed by using the PROTTER program (Omasits et al., 2014), while the alignment of LaMATE2 protein sequence with orthologous MATE-protein sequences was generated by Clustal-WS using Jalview software version 2 (Waterhouse et al., 2009).

Statistical significance was determined by one-way analysis of variances (ANOVA) using Holm–Sidak test, P < 0.05 or 0.01. Statistical analyses were calculated using SigmaPlot (Systat Software Inc., San Jose, CA, USA).

## RESULTS AND DISCUSSION

### Release of flavonoids and expression of *LaMATE2*

Former work from our laboratories showed that flavonoids are released mainly from the so-called juvenile, and immature (young states) cluster roots and that genistein and its derivative exudation was induced by phosphate (P) deficiency (Weisskopf et al., 2006). Furthermore, in 2001, we published research on genes differentially expressed amongst different stages of cluster roots in P-deficient white lupin (Massonneau et al., 2001). Within these genes, we

identified two LaMATE-type transporters, *LaMATE1* was

predominantly expressed in mature (Uhde-Stone et al., 2005), citrate

excreting cluster roots, while the second (*LaMATE2*) was

predominantly present in juvenile cluster roots.

To identify a flavonoid exporter possibly playing an important role

in the establishment of nodules in *Fabaceae* plants, we looked first

whether a link between the expression of *LaMATE2* and release of

flavonoids exists. Under P-deficiency, the *LaMATE2* was

predominantly expressed in the root apex and in the juvenile stage

of cluster roots (Figure 1a). Nitrogen (N) deficiency, on the other

hand, induced *LaMATE2* expression mainly in apex and nodule

(Figure 2a). Analyses of the root exudates of white lupin revealed

that genistein release was strongly induced by P deficiency,

particularly in young cluster root tissues (Figure 1b), confirming

previous results (Weisskopf et al., 2006). In order to see whether a

similar behavior could be observed under N deficiency, plants were

grown under N-deficient condition. In this case, cluster root

formation was not induced. Hence, we compared the whole roots of

plants grown in the presence or absence of N. Nevertheless, we

could observe that under N deficiency, genistein release was also

increased significantly after 1 and even more pronounced after 2

weeks (Figure 2b).

**Characterization of LaMATE2**

As already shown for other MATE transporters, LaMATE2 contains

12-transmembrane helical domains (Figure S2) (He et al., 2010).

Indeed, LaMATE2 exhibits a good homology to other plant MATE

transporters (Figure 3, S2), including the functionally characterized

vacuolar flavonoid transporters MtMATE1, MtMATE2 and

AtTT12 (Marinova et al., 2007; Zhao and Dixon, 2010; Zhao et al.,

2011). Up to date, only few MATE transporters have been

characterized in roots and most of them mediate either the efflux of

citrate, such as AtFRD3, HvAACT1, SbMATE1 (Durrett et al.,

2007; Furukawa et al., 2007; Magalhaes et al., 2007) or act as

vacuolar flavonoid transporters in *Arabidopsis*, *Medicago* or

grapevine (for a review see Zhao and Dixon (2010) and Zhao

(2015)). Interestingly all citrate transporters contain a large

cytosolic loop. In white lupin roots and in agreement with our

previous results (Massonneau et al., 2001), the expression of another

MATE transporter, LaMATE1, a homolog of the citrate transporter

AtFRD3, is highest at the mature stage where a burst of citrate

exudation occurs (Uhde-Stone et al., 2005; Wang et al., 2014). Also,

this transporter contains a large cytosolic loop.

At the amino acid level, LaMATE2 exhibits a high similarity to a

cluster of MATE proteins, which includes a heterogeneous group of

transporters (different substrates, subcellular localization and

physiological role). Among the characterized members of this

cluster, NtJAT1 and AtDTX1 are the closest homologs of

LaMATE2, 55 and 50% of identities, respectively (Figure 3). Both

are localized at the plasma membrane (PM) of root cells and

function as efflux carriers for plant-derived alkaloids and other toxic

compounds (Li et al., 2002; Morita et al., 2009).

Several ABC transporters and MATE proteins mediating the export

of compounds into the soil have been described (Weston et al., 2012;

Baetz and Martinoia, 2014). However, none of them was shown so

far to be responsible for the root release of flavonoids and

isoflavonoids. Our data showing a connection between flavonoid

exudation and *LaMATE2* expression together with the fact that

LaMATE2 fused with GFP co-localizes with the PM marker

AtPIP2a-mCherry (Figure 4) prompted us to investigate whether

LaMATE2 acts as a flavonoid transporter.

**LaMATE2 silencing reduces genistein release, P mobilization and nodule number**

Genistein is one of the major released isoflavonoids of *Fabaceae* inducing *NOD* genes in *Rhizobium* bacteria, attracting them and inducing nodule formation (Zhang and Smith, 1995; Lang et al., 2008; Liu and Murray, 2016). Therefore, to assess whether the LaMATE2 might affect genistein release and nodule formation, we have used RNA-dependent gene silencing (*LaMATE2*-RNAi) in roots of white lupin plants. The effectiveness of the silencing was confirmed by a significant reduction (approximatively -80 and 65 % of *LaMATE2* expression in *LaMATE2*-RNAi transformants grown in N and P deficiency, respectively (Figure 5a, g). Analysis of these roots revealed that the release of genistein was strongly reduced in these transformants (Figure 5b, h) and concomitantly isoflavonoids, the uppermost as glycosides, were accumulating in the cell content of silenced roots, which can be regarded as a direct consequence of the reduced genistein exudation (Figure 5d-f). To investigate whether the impaired genistein exudation influences nodulation, we used the *LaMATE2*-RNAi plants to compare the nodule number of silenced and control plants. Our results highlight the importance of LaMATE2-dependent isoflavonoid release in the early step of nodulation, since *LaMATE2* silencing leads to a highly significant reduction (a. -80 %) of nodules in *LaMATE2*-RNAi compared to empty-vector transformed roots (Figure 5c, 6b, S4). Interestingly exogenous application of genistein onto the LaMATE2-RNAi roots partially restored the nodulation efficiency (Figure 6 a, b). Similarly, *LaMATE2* expression, genistein release patterns and capability of root exudates to mobilize P were measured under P-deficient conditions in silenced plants (Figure 5g-i). Cluster roots of *LaMATE2*-RNAi transformants grown under P deficiency released 85% less genistein along with the lower level of *LaMATE2*

expression (a. -60%). The SPE-purified exudates from those roots also exhibited a limited (a. -66%) mobilization capability from a poorly soluble P source (vivianite). These results suggest that LaMATE2 is the major if not the sole genistein exporter from roots of white lupin plants and in P-deficient condition it contributes to the P solubilization process.

The results showing that genistein export is dependent on the presence of LaMATE2 indicates that within the root this transporter is at least partially localized in the cortex. Attempts in our laboratory to get a more detailed picture on its localization, either using in-situ hybridization or a GUS-promoter construct unfortunately failed. However, it should be mentioned that interestingly, excretion of strigolactones into the soil was independent, whether the transporter was localized specifically in hypodermal passage cells as in Petunia or in the whole cortex as in Medicago (Kretzschmar et al., 2012; Banasiak et al., 2020).

**Transport of phenolics by LaMATE2.**

To demonstrate that the reduced genistein release of LaMATE2 silenced plants depends directly on the activity of LaMATE2, we expressed it heterologously in *Saccharomyces cerevisiae* and performed transport assays using isolated membrane vesicles (Figure 7). Indeed, using this system, we could observe that LaMATE2 mediates an efficient transport of genistein which was characterized by a saturable kinetic exhibiting an apparent Km of 16.2 μM (Figure 7b). This Km is in the same order of magnitude as determined for the vacuolar flavonoid transporters characterized so far (e.g. AtTT12 and MtMATE1: Km of 50 and 36 μM for epicatechin 3′-O-glucoside, respectively; MtMATE2: Km of 88 μM for cyanidin 3-O-glucoside) (Zhao and Dixon, 2010; Zhao et al., 2011). In plasmalemma vesicles isolated from soybean roots,

genistein transport exhibited a Km of 160 µM and was strongly

dependent on the presence of ATP (Sugiyama et al., 2007).

Up to date, certain MATEs have been shown to exhibit a broad range

of substrate specificity (Takanashi et al., 2014), while LaMATE2

displays a strong specificity for the substrate genistein (Figure 6c).

Besides genistein, LaMATE2 was able to transport only daidzein at

a low rate, but not other tested flavonoids such as genistin (a

glycosylated form of genistein), hydroxygenistein, biochanin A (a

methylated form of genistein) or kaempferol. Previous observations

in soybean showed that the presence of daidzein in the external

media strongly limited genistein transport in plasma membrane

vesicles of root cells, indicating a possible competition between

those molecules (Sugiyama et al., 2007).

A feature of MATE proteins is to couple the substrate transport to

an electrochemical gradient, working as $H^+$ or $Na^+$/substrate

antiporters (He et al., 2010; Lu et al., 2013; Tanaka et al., 2013). Our

results showed that ATP-dependent $^3$H-genistein accumulation in

vesicles deriving from *LaMATE2*$_{ORF}$-expressing yeast was strongly

reduced in presence of ammonium chloride, which dissipates the

transmembrane proton gradient (Marinova et al., 2007; Zhao and

Dixon, 2010) (Figure 7a), indicating that LaMATE2 acts as a

substrate-proton co-transporter, similarly to AtTT12 (Marinova et

al., 2007), AtDTX1 (Li et al., 2002) and NtJAT1 (Morita et al.,

2009).

**Conclusions**

In this work the long-sought-after isoflavonoid plasma membrane

exporter required to attract symbiotic bacteria for N fixation has

been identified. Different *leguminous* plants release different sets of

isoflavonoids to induce nodulation (Liu and Murray, 2016).

Released genistein has been indicated in faba bean and soybean as

*rhizobia* attractant (Zhang and Smith, 1995; Li et al., 2016). Since

*Medicago truncatula* and soybean both encode one MATE protein

that exhibits high homology to LaMATE2, it is tempting to

speculate that these homologues could also act as isoflavonoid

exporters, possibly showing slightly different substrate preferences

according to the isoflavonoid produced by the plant to initiate the

symbiosis. However, it cannot be excluded that in addition to

MATE-type transporters and some ABC transporters are also

involved in isoflavonoid release. Since isoflavonoids can also

release Pi from minerals (Cesco et al., 2010) and organic complexes

these transporters may play a dual role in N and P supply.

During the last years, we learned a lot about the signaling pathways

and the effectors involved in establishing the legume-*rhizobia*

symbiosis. Surprisingly molecular identity of membrane

transporters responsible for the release of phenolic compounds

initiating interactions was not known. With the identification of

LaMATE2 we succeeded to identify the very initial step leading to

this symbiosis. Since there is a huge interest to transfer this complex

mechanism to other crop plants to grow them in the absence of

artificial N-sources we do believe that it is a valuable piece of the

puzzle to consider in such an ambitious project.

**ACKNOWLEDGEMENTS**

Research was supported by grants from Italian Ministry of

University and Research-MIUR (FIRB-Programme Futuro in

Ricerca, RBFR08L2ZT and RBFR127WJ9), from funds of the

Zurich University; and statutory funds from the Polish Ministry of

Science and Higher Education. **Competing interests:** Authors

declare no competing interests; and **Data and materials**

**availability:** Sequence data from this article can be found in the

EMBL/GenBank data libraries under accession number KY464927,

LaMATE2 of *Lupinus albus*.

**AUTHOR CONTRIBUTION**

NT, EM, RP, SC, MJ: designed and oversaw the research; WB,

LZ, SG, RBF, SV, FV, TM, BB, NT: performed experiments and

analyzed data; NT, EM, RP, MJ: wrote the paper

**REFERENCES**

Baetz, U., and Martinoia, E. (2014). Root exudates: the hidden part
of plant defense. *Trends in Plant Science* 19**,** 90-98.
Banasiak, J., Borghi, L., Stec, N., Martinoia, E., and Jasiński, M.
(2020). The Full-Size ABCG Transporter of *Medicago*
*truncatula* Is Involved in Strigolactone Secretion, Affecting
Arbuscular Mycorrhiza. 11.
Biala, W., Banasiak, J., Jarzyniak, K., Pawela, A., and Jasinski, M.
(2017). Medicago truncatula ABCG10 is a transporter of 4-
coumarate and liquiritigenin in the medicarpin biosynthetic
pathway. *Journal of experimental botany* 68**,** 3231-3241.
Burke, D., Dawson, D., and Stearns, T. (2000). Methods in yeast
genetics: a Cold Spring Harbor Laboratory course manual,
2000 ed. *Cold Spring Harbor Press: Plainview, NY* 17**,**
205.
Cesco, S., Neumann, G., Tomasi, N., Pinton, R., and Weisskopf, L.
(2010). Release of plant-borne flavonoids into the
rhizosphere and their role in plant nutrition. *Plant and Soil*
329**,** 1-25.
Durrett, T.P., Gassmann, W., and Rogers, E.E. (2007). The FRD3-
Mediated Efflux of Citrate into the Root Vasculature Is
Necessary for Efficient Iron Translocation. *Plant*
*Physiology* 144**,** 197-205.
Furukawa, J., Yamaji, N., Wang, H., Mitani, N., Murata, Y., Sato,
K., Katsuhara, M., Takeda, K., and Ma, J.F. (2007). An
Aluminum-Activated Citrate Transporter in Barley. *Plant*
*and Cell Physiology* 48**,** 1081-1091.
Gietz, R.D., and Woods, R.A. (2002). Transformation of yeast by
lithium acetate/single-stranded carrier DNA/polyethylene
glycol method. *Methods in enzymology* 350**,** 87-96.

He, X., Szewczyk, P., Karyakin, A., Evin, M., Hong, W.-X.,
Zhang, Q., and Chang, G. (2010). Structure of a cation-
bound multidrug and toxic compound extrusion transporter.
*Nature* 467**,** 991.
Hufnagel, B., Marques, A., Soriano, A., Marquès, L., Divol, F.,
Doumas, P., Sallet, E., Mancinotti, D., Carrere, S.,
Marande, W., Arribat, S., Keller, J., Huneau, C., Blein, T.,
Aimé, D., Laguerre, M., Taylor, J., Schubert, V., Nelson,
M., Geu-Flores, F., Crespi, M., Gallardo, K., Delaux, P.-
M., Salse, J., Bergès, H., Guyot, R., Gouzy, J., and Péret,
B. (2020). High-quality genome sequence of white lupin
provides insight into soil exploration and seed quality.
*Nature Communications* 11**,** 492.
Jiang, Y., Wang, W., Xie, Q., Liu, N., Liu, L., Wang, D., Zhang,
X., Yang, C., Chen, X., Tang, D., and Wang, E. (2017).
Plants transfer lipids to sustain colonization by mutualistic
mycorrhizal and parasitic fungi. *Science* 356**,** 1172-1175.
Jin, J.B., Kim, Y.A., Kim, S.J., Lee, S.H., Kim, D.H., Cheong,
G.W., and Hwang, I. (2001). A new dynamin-like protein,
ADL6, is involved in trafficking from the *trans*-Golgi
network to the central vacuole in Arabidopsis. *The Plant*
*cell* 13**,** 1511-1526.
Klein, M., Mamnun, Y.M., Eggmann, T., Schüller, C., Wolfger,
H., Martinoia, E., and Kuchler, K. (2002). The ATP-
binding cassette (ABC) transporter Bpt1p mediates
vacuolar sequestration of glutathione conjugates in yeast.
*FEBS letters* 520**,** 63-67.
Komarova, N.Y., Thor, K., Gubler, A., Meier, S., Dietrich, D.,
Weichert, A., Suter Grotemeyer, M., Tegeder, M., and
Rentsch, D. (2008). AtPTR1 and AtPTR5 Transport
Dipeptides in Planta. *Plant Physiology* 148**,** 856-869.
Koressaar, T., and Remm, M. (2007). Enhancements and
modifications of primer design program Primer3.
*Bioinformatics* 23**,** 1289-1291.
Kretzschmar, T., Kohlen, W., Sasse, J., Borghi, L., Schlegel, M.,
Bachelier, J.B., Reinhardt, D., Bours, R., Bouwmeester,
H.J., and Martinoia, E. (2012). A petunia ABC protein
controls strigolactone-dependent symbiotic signalling and
branching. *Nature* 483**,** 341-344.
Lambers, H., Martinoia, E., and Renton, M. (2015). Plant
adaptations to severely phosphorus-impoverished soils.
*Current Opinion in Plant Biology* 25**,** 23-31.
Lang, K., Lindemann, A., Hauser, F., and Göttfert, M. (2008). The
genistein stimulon of *Bradyrhizobium japonicum*.
*Molecular Genetics and Genomics* 279**,** 203-211.

Li, B., Li, Y.-Y., Wu, H.-M., Zhang, F.-F., Li, C.-J., Li, X.-X., Lambers, H., and Li, L. (2016). Root exudates drive interspecific facilitation by enhancing nodulation and $N_2$ fixation. *Proceedings of the National Academy of Sciences* 113**,** 6496-6501.

Li, L., He, Z., Pandey, G.K., Tsuchiya, T., and Luan, S. (2002). Functional Cloning and Characterization of a Plant Efflux Carrier for Multidrug and Heavy Metal Detoxification. *Journal of Biological Chemistry* 277**,** 5360-5368.

Limpens, E., Ramos, J., Franken, C., Raz, V., Compaan, B., Franssen, H., Bisseling, T., and Geurts, R. (2004). RNA interference in *Agrobacterium rhizogenes*-transformed roots of *Arabidopsis* and *Medicago truncatula*. *Journal of Experimental Botany* 55**,** 983-992.

Liu, C.-W., and Murray, J. (2016). The Role of Flavonoids in Nodulation Host-Range Specificity: An Update. *Plants* 5**,** 33.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the $2-\Delta\Delta CT$ Method. *Methods* 25**,** 402-408.

Lu, M., Radchenko, M., Symersky, J., Nie, R., and Guo, Y. (2013). Structural insights into $H^+$-coupled multidrug extrusion by a MATE transporter. *Nature Structural & Molecular Biology* 20**,** 1310-1317.

Luginbuehl, L.H., Menard, G.N., Kurup, S., Van Erp, H., Radhakrishnan, G.V., Breakspear, A., Oldroyd, G.E.D., and Eastmond, P.J. (2017). Fatty acids in arbuscular mycorrhizal fungi are synthesized by the host plant. *Science* 356**,** 1175-1178.

Magalhaes, J.V., Liu, J., Guimaraes, C.T., Lana, U.G.P., Alves, V.M.C., Wang, Y.H., Schaffert, R.E., Hoekenga, O.A., Pineros, M.A., Shaff, J.E., Klein, P.E., Carneiro, N.P., Coelho, C.M., Trick, H.N., and Kochian, L.V. (2007). A gene in the multidrug and toxic compound extrusion (MATE) family confers aluminum tolerance in sorghum. *Nature Genetics* 39**,** 1156-1161.

Marinova, K., Kleinschmidt, K., Weissenbock, G., and Klein, M. (2007). Flavonoid Biosynthesis in Barley Primary Leaves Requires the Presence of the Vacuole and Controls the Activity of Vacuolar Flavonoid Transport. *Plant Physiology* 144**,** 432-444.

Martin, F.M., Uroz, S., and Barker, D.G. (2017). Ancestral alliances: Plant mutualistic symbioses with fungi and bacteria. *Science* 356**,** eaad4501.

Massonneau, A., Langlade, N., Leon, S., Smutny, J., Vogt, E., Neumann, G., and Martinoia, E. (2001). Metabolic changes

738           associated with cluster root development in white lupin

739           (*Lupinus albus* L.): relationship between organic acid

740           excretion, sucrose metabolism and energy status. *Planta*

741           213**,** 534-542.

742  Morita, M., Shitan, N., Sawada, K., Van Montagu, M.C.E., Inzé,

743           D., Rischer, H., Goossens, A., Oksman-Caldentey, K.-M.,

744           Moriyama, Y., and Yazaki, K. (2009). Vacuolar transport

745           of nicotine is mediated by a multidrug and toxic compound

746           extrusion (MATE) transporter in *Nicotiana tabacum*.

747           *Proceedings of the National Academy of Sciences of the*

748           *United States of America* 106**,** 2447-2452.

749  Nelson, B.K., Cai, X., and Nebenführ, A. (2007). A multicolored

750           set of in vivo organelle markers for co-localization studies

751           in *Arabidopsis* and other plants. *The Plant Journal* 51**,**

752           1126-1136.

753  Neumann, G., and Martinoia, E. (2002). Cluster roots - an

754           underground adaptation for survival in extreme

755           environments. *Trends in Plant Science* 7**,** 162-167.

756  Omasits, U., Ahrens, C.H., Müller, S., and Wollscheid, B. (2014).

757           Protter: interactive protein feature visualization and

758           integration with experimental proteomic data.

759           *Bioinformatics* 30**,** 884-886.

760  Purnell, H.M. (1960). Studies of the family Proteaceae. Anatomy

761           and morphology of the roots of some Victorian species.

762           *Australian Journal of Botany* 8**,** 38-50.

763  Quandt, H.J., Pühler, A., and Broer, I. (1993). Transgenic root

764           nodules of *Vicia hirsuta:* a fast and efficient system for the

765           study of gene expression in indeterminate-type nodules.

766           *MPMI-Molecular Plant Microbe Interactions* 6**,** 699-706.

767  Ritz, C., and Spiess, A.N. (2008). qpcR: an R package for

768           sigmoidal model selection in quantitative real-time

769           polymerase chain reaction analysis. *Bioinformatics* 24**,**

770           1549-1551.

771  Sauer, N., and Stolz, J. (1994). SUC1 and SUC2: two sucrose

772           transporters from *Arabidopsis thaliana*; expression and

773           characterization in baker's yeast and identification of the

774           histidine-tagged protein. *The Plant Journal* 6**,** 67-77.

775  Staszków, A., Swarcewicz, B., Banasiak, J., Muth, D., Jasiński,

776           M., and Stobiecki, M. (2011). LC/MS profiling of

777           flavonoid glycoconjugates isolated from hairy roots,

778           suspension root cell cultures and seedling roots of

779           Medicago truncatula. *Metabolomics* 7**,** 604-613.

780  Stępkowski, T., Żak, M., Moulin, L., Króliczak, J., Golińska, B.,

781           Narożna, D., Safronova, V.I., and Mądrzak, C.J. (2011).

782           *Bradyrhizobium canariense* and *Bradyrhizobium*

783           *japonicum* are the two dominant *rhizobium* species in root

784    nodules of lupin and serradella plants growing in Europe.
785    *Systematic and Applied Microbiology* 34**,** 368-375.

786 Strozycki, P.M., Skąpska, A., Szcześniak, K., Sobieszczuk, E.,
787    Briat, J.-F., and Legocki, A.B. (2003). Differential
788    expression and evolutionary analysis of the three ferritin
789    genes in the legume plant *Lupinus luteus*. *Physiologia*
790    *Plantarum* 118**,** 380-389.

791 Subramanian, S., Stacey, G., and Yu, O. (2006). Endogenous
792    isoflavones are essential for the establishment of symbiosis
793    between soybean and *Bradyrhizobium japonicum*. *The*
794    *Plant Journal* 48**,** 261-273.

795 Sugiyama, A., Shitan, N., and Yazaki, K. (2007). Involvement of a
796    soybean ATP-binding cassette-Type transporter in the
797    secretion of genistein, a signal flavonoid in legume-
798    *Rhizobium* Symbiosis. *Plant Physiology* 144**,** 2000-2008.

799 Takanashi, K., Shitan, N., and Yazaki, K. (2014). The multidrug
800    and toxic compound extrusion (MATE) family in plants.
801    *Plant Biotechnology* 31**,** 417-430.

802 Tamura, K., Stecher, G., Peterson, D., Filipski, A., and Kumar, S.
803    (2013). MEGA6: Molecular Evolutionary Genetics
804    Analysis version 6.0. *Molecular biology and evolution* 30**,**
805    2725-2729.

806 Tanaka, Y., Hipolito, C.J., Maturana, A.D., Ito, K., Kuroda, T.,
807    Higuchi, T., Katoh, T., Kato, H.E., Hattori, M., Kumazaki,
808    K., Tsukazaki, T., Ishitani, R., Suga, H., and Nureki, O.
809    (2013). Structural basis for the drug extrusion mechanism
810    by a MATE multidrug transporter. *Nature* 496**,** 247-251.

811 Tilman, D. (1999). Global environmental impacts of agricultural
812    expansion: The need for sustainable and efficient practices.
813    *Proceedings of the National Academy of Sciences* 96**,** 5995-
814    6000.

815 Tilman, D., Fargione, J., Wolff, B., D'antonio, C., Dobson, A.,
816    Howarth, R., Schindler, D., Schlesinger, W.H., Simberloff,
817    D., and Swackhamer, D. (2001). Forecasting Agriculturally
818    Driven Global Environmental Change. *Science* 292**,** 281-
819    284.

820 Udvardi, M., and Poole, P.S. (2013). Transport and Metabolism in
821    Legume-*Rhizobia* Symbioses. *Annual Review of Plant*
822    *Biology* 64**,** 781-805.

823 Uhde-Stone, C., Liu, J., Zinn, K.E., Allan, D.L., and Vance, C.P.
824    (2005). Transgenic proteoid roots of white lupin: a vehicle
825    for characterizing and silencing root genes involved in
826    adaptation to P stress. *Plant Journal* 44**,** 840-853.

827 Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth,
828    B.C., Remm, M., and Rozen, S.G. (2012). Primer3—new

capabilities and interfaces. *Nucleic Acids Research* 40**,**
e115.
Vance, C.P., Uhde-Stone, C., and Allan, D.L. (2003). Phosphorus
acquisition and use: critical adaptations by plants for
securing a nonrenewable resource. *New Phytologist* 157**,**
423-447.
Wang, Z., Straub, D., Yang, H., Kania, A., Shen, J., Ludewig, U.,
and Neumann, G. (2014). The regulatory network of
cluster-root function and development in phosphate-
deficient white lupin (*Lupinus albus*) identified by
transcriptome sequencing. *Physiologia Plantarum* 151**,**
323-338.
Wasson, A.P., Pellerone, F.I., and Mathesius, U. (2006). Silencing
the flavonoid pathway in *Medicago truncatula* inhibits root
nodule formation and prevents auxin transport regulation
by rhizobia. *Plant Cell* 18**,** 1617-1629.
Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M., and
Barton, G.J. (2009). Jalview Version 2—a multiple
sequence alignment editor and analysis workbench.
*Bioinformatics* 25**,** 1189-1191.
Weisskopf, L., Tomasi, N., Santelia, D., Martinoia, E., Langlade,
N.B., Tabacchi, R., and Abou-Mansour, E. (2006).
Isoflavonoid exudation from white lupin roots is influenced
by phosphate supply, root type and cluster-root stage. *New*
*Phytologist* 171**,** 657-668.
Weston, L.A., Ryan, P.R., and Watt, M. (2012). Mechanisms for
cellular transport and release of allelochemicals from plant
roots into the rhizosphere. *Journal of Experimental Botany*.
Zhang, F., and Smith, D.L. (1995). Preincubation of
Bradyrhizobium japonicum with Genistein Accelerates
Nodule Development of Soybean at Suboptimal Root Zone
Temperatures. *Plant Physiology* 108**,** 961-968.
Zhao, J. (2015). Flavonoid transport mechanisms: how to go, and
with whom. *Trends in Plant Science* 20**,** 576-585.
Zhao, J., and Dixon, R.A. (2010). The 'ins' and 'outs' of flavonoid
transport. *Trends in Plant Science* 15**,** 72-80.
Zhao, J., Huhman, D., Shadle, G., He, X.-Z., Sumner, L.W., Tang,
Y., and Dixon, R.A. (2011). MATE2 Mediates Vacuolar
Sequestration of Flavonoid Glycosides and Glycoside
Malonates in *Medicago truncatula*. *The Plant Cell* 23**,**
1536-1555.
870
871
872
873

**Figure 1. *LaMATE2* expression level and genistein release in roots of P-deficient white lupin**

*LaMATE2* expression analyses in white lupin grown under control (+N+P, black bars) condition or P deficiency (+N-P, white bars) (**a**). Gene expression was evaluated in control apex (+N+P), P-deficient apex and cluster roots (separated depending on developing stages: Juvenile, Immature, Mature, Senescent cluster-root stages; example of cluster root shown in Figure S1). Expression levels are shown relative to the *LaMATE2* expression level of the root apex under control conditions (+N+P). Release of genistein from different root tissues of 4-week-old P-deficient plants, release from the control apex was below detectable value <LOD (**b**). Data are means +SD (* refer to statistically significant differences among the mean value of the sample *vs* control or -P apex for genistein release, ANOVA Holm–Sidak, N=6, *P <0.05, ** P <0.01).

**Figure 2. *LaMATE2* expression analyses and genistein release in roots of N-deficient white lupin**

*LaMATE2* expression analyses in white lupin grown under control (+N+P, black bars) condition or N-deficiency (-N+P, dark grey bars) (**a**). Gene expression was evaluated in control and N-deficient (other) root, root apex and nodules. Expression level is shown relative to *LaMATE2* expression in control (+N+P) root. Root release of genistein from white lupin plants grown under 1- or 2-week-old N-sufficient and -deficient condition (**b**). Genistein release in 7 and 14 days of N deficiency (-N+P) or sufficiency (+N+P) . Data are means +SD (* refers to statistically significant differences among the mean value of the sample *vs* control, ANOVA Holm–Sidak, N=6, * P <0.05, ** P <0.01).

0.2

**Figure 3. Phylogenetic tree of MATE transporters**

A phylogenetic analysis was performed using the LaMATE2 amino
acid sequences of *Lupinus albus* (LaMATE1, LaMATE2,
Lalb_Chr17g0335991,　　　　　　　　Lalb_Chr06g0165721,
Lalb_Chr02g0145611, LaALMT1); AtFRD3 AtEDS5, AtTT12,
AtDTX1,10,11,12,13,14, AtALF5 and AtFFT of *Arabidopsis*
*thaliana*; Glyma.10G267700 of *Glycine max*; HvAACT1 of
*Hordeum vulgare*; MtMATE1, MtMATE2, Medtr1g108840,
Medtr1g108990, Medtr1g109060.2, Medtr1g109060.1 of *Medicago*
*truncatula*; NtJAT1, NtMATE1 and NtMATE2 of *Nicotiana*

*tabacum*; SlMTP77 of *Solanum lycopersicum*; SbMATE1 of
*Sorghum bicolor*; VvAM1 and VvAM3 of *Vitis vinifera*; VcMATE1
and VcMATE4 of *Vaccinium corymbosum*. The tree was
constructed by aligning the protein sequences by Clustal-W and the
evolutionary history was inferred using the Neighbour-Joining
method. The percentage of replicate trees in which the associated
taxa clustered together in the bootstrap test (1000 replicates) are
shown next to the branches. The tree is drawn to scale, with branch
lengths in the same units as those of the evolutionary distances used
to infer the phylogenetic tree. The evolutionary distances were
computed using the Poisson correction method and are in the units
of the number of amino acid substitutions per site.

**Figure 4. Subcellular localization of LaMATE2**

Co-localization of LaMATE2 fused with green fluorescent protein (GFP) and mCherry-labeled plasma membrane marker AtPIP2A in an Arabidopsis mesophyll protoplast. White scale bars = 20 μm. Fluorescence intensity over distance plot of LaMATE2-GFP (green) and AtPIP2a-mCherry (red).

**Figure 5. Effect of *LaMATE2* silencing in N and P deficient roots**

Alteration of *LaMATE2* expression, flavonoid content and release,
nodule number or P mobilization due to *LaMATE2* silencing in N
(**a-f**) or P (**g-i**) deficiency. *LaMATE2* relative expression (**a, g**),

genistein release (**b, h**), number of nodules per plant (pictures shown in Figure S4) (**c**), cell content: genistein (**d**), genistein 7-O-glucoside (**e**), genistein malonylglucoside (**f**), P mobilization from a poorly soluble P source (**i**). The analyses were performed on roots of N- or P-deficient lupin plants independently transformed with either pRedRoot::*LaMATE2* RNAi (*LaMATE2* RNAi) or empty-vector pRedRoot (Empty vector).

All data are expressed relative to Empty-vector-transformed roots, with the exceptions of the number of nodule per plants and P mobilization ($\mu$mol P h$^{-1}$). Data are means+SD (* refers to statistically significant differences among the mean value of RNAi and Empty vector, ANOVA Holm–Sidak, N=6-21, *P <0.05; ** for P<0.01).

953
954
**Figure 6. Effect of exogenous genistein on *LaMATE2* expression and nodule numbers**

955
956

*LaMATE2* relative expression (**a**) and number of nodules per plant
(**b**) in roots of N-deficient lupin plants independently transformed
with either pRedRoot::*LaMATE2* RNAi (*LaMATE2* RNAi) or
empty-vector pRedRoot (Empty vector) or RNAi roots treated with
1 μM genistein (*LaMATE2* RNAi + genistein).
Expression data are shown relative to *LaMATE2* expression level in
empty vector-transformed roots. Data are means+SD (* refers to
statistically significant differences among the mean value of RNAi
and EV or RNAi+G values, ANOVA Holm–Sidak, N=6, *P <0.05;
** for P<0.01).
967

**Figure 7. Transport of phenylpropanoid compounds mediated by LaMATE2**

LaMATE2-mediated transport rate of phenylpropanoids in yeast microsomal membrane vesicles. (**A**) Time-dependent accumulation of $^3$H-genistein in yeast microsomal membrane vesicles. Membrane vesicles were isolated from yeast transformed with the empty vector (pNEV empty vector) or transformed with pNEV-*LaMATE2*$_{ORF}$. The yeast vesicles were incubated up to 120 s in presence of 5 µM $^3$H-genistein: To determine the effect of the pH gradient, 25 mM NH$_4$Cl was added in an assay solution (pNEV-*LaMATE2*$_{ORF}$ + NH$_4$Cl), as uncoupler of the proton gradient. (**B**) Concentration-dependent transport of $^3$H-genistein by LaMATE2 in yeast vesicles. Vesicles were incubated for 30 s in the assay solution containing $^3$H-genistein at different concentrations (from 5 to 100 µM). The kinetic parameters of $^3$H-genistein uptake were calculated by subtracting uptake rates recorded in the empty-vector vesicles using the Hanes–Woolf plot. (**C**) Substrate specificity of LaMATE2 transporter was evaluated in presence of genistein, hydroxygenistein, genistin (glycosylated genistein), biochanin A (methylgenistein), daidzein or kaempferol (5 µM, 30 s). All data are expressed relative to pNEV-*LaMATE2*$_{ORF}$-transformed yeast. Data are means±SD of three independent experiments (*refers to statistically significant differences among the mean value in comparison to empty vector sample within each time point; capital letters refer to statistically significant differences among the mean value, ANOVA Holm–Sidak, N=6, *P <0.05, **P <0.01, <LOD: below detectable value).

**Supplementary materials**

1000

**Figure S1. Cluster roots of P-deficient white lupin**

Picture of a lateral root of a 4-week-old white lupin grown under P-deficient condition (+N-P). The different developing stages: Juvenile, Immature, Mature, Senescent cluster-root stage and Apex tissue [10 mm from the root tips] are highlighted and named. White scale bar = 10 mm.

1008

1009

1010

1011

1012 **Figure S2. Predicted membrane topology for LaMATE2 from**
1013 **white lupin**

1014

1015 The model was obtained by using the PROTTER program (Omasits
1016 *et al.,* 2014).

1017

1018

1019

1020
1021

1022

**Figure S3. Multiple sequence alignment of amino acid sequence**
**of LaMATE2 with selected MATE transporter orthologs in**
**higher plants**
Protein sequence alignment was performed using Clustal-WS using
Jalview software version 2 (Waterhouse *et al.,* 2009). Amino acids
with only highly conservative substitutions are highlighted in colour
blue. Thin grey lines below the consensus sequence (D 1-5) indicate
five short stretches of conservative amino acids reported for all 56
Arabidopsis MATE proteins. The red box indicates the residue E290
(TT12), constituting the cation-binding site in the pore. Twelve
putative-transmembrane helical domains (TM 1-12) of LaMATE2
are indicated above the alignment
1035

1036
1037

1038
1039

**Figure S4. Pictures of *LaMATE2*-silenced roots grown under N-deficient conditions**

Roots are shown in the following order: wild type, empty-vector-transformed roots and *LaMATE2*-RNAi-transformed roots. Nodule localizations are highlighted with arrows and close-up inserts are shown in the bottom right corners.

1046

1047

**Figure S5. Gene expression in the silencing experiments**

Relative expression of *LaMATE2* and the two closest homologues:
*Lalb_Chr06g0165721* and *Lalb_Chr02g0145611* in P-deficient (**a**)
and N-deficient (**b**) condition in pRedRoot::*LaMATE2* RNAi
(RNAi) or empty-vector pRedRoot (EV) transformed roots.
Expression data are expressed relative to *LaMATE2* expression in
EV-transformed roots. Data are means+SD (* refers to statistically
significant differences among the mean value of RNAi and EV,
ANOVA Holm–Sidak, N=3, **P <0.01).

1057

1058

1059 **Table S1.**
1060 List of primers

| | FORWARD | REVERSE |
|---|---|---|
| *LaMATE2* | cttgttttgctgtctgggct | tgcagcaccaattccaaagg |
| *Lalb_Chr06g0165721* | gtgaaacgatgaacatagaac | ctttgttgctgttgatgatg |
| *Lalb_Chr02g0145611* | agtgtcaacaccattgca | gatctaaccttgctgcag |
| *LaMATE2-GFP* | atagctagcatggaagagagtctaatacc | atacatgcctcatgctagactattttctgc |
| pNEV-*LaMATE2*ORF | atgcggccgcatggaagagagtctaatacc | agcggccgctcatgctagactattttctg |
| *LaMATE2* RNAi a | atccatggggcacgagatttctatct | gtgcatttaaatccacaaatagtttccagtc |
| *LaMATE2* RNAi b | atggatccccacaaatagtttccagtc | tgcactagtggcacgagatttctatct |
| *LaUbiquitin* | gcaccctagccgactacaac | ccggtaagggtcttgacaaa |

1061

1062

1063