The Pace of Modern Life, Revisited

Sarah Sanderson[1], Marc-Olivier Beausoleil[1], Rose E. O'Dea[1,2], Zachary T. Wood[3], Cristian Correa[4,5], Victor Frankel[1], Lucas D. Gorné[1,6,7,8], Grant E. Haines[1], Michael T. Kinnison[3], Krista B. Oke[9], Fanie Pelletier[10], Felipe Pérez-Jvostov[1], Winer Daniel Reyes-Corral[1], Yanny Ritchot[10], Freedom Sorbara[1], Kiyoko M. Gotanda[8,10*†], and Andrew P. Hendry[1*]

1. Department of Biology and Redpath Museum, McGill University, 859 Sherbrooke Street West, Montréal, Québec H3A 2K6, Canada

2. Evolution & Ecology Research Centre, UNSW, Sydney, Australia

3. School of Biology and Ecology and Maine Center for Genetics in the Environment, University of Maine, Orono, ME, USA.

4. Facultad de Ciencias Forestales y Recursos Naturales, Instituto de Conservación Biodiversidad y Territorio, Universidad Austral de Chile, Valdivia, Chile

5. Centro de Humedales Río Cruces, Universidad Austral de Chile, Valdivia, Chile

6. Universidad Nacional de Córdoba, Facultad de Ciencias Exactas Físicas y Naturales. Córdoba, Argentina.

7. Consejo Nacional de Investigaciones Científicas y Técnicas, CONICET, IMBiV. Córdoba, Argentina.

8. Department of Biological Sciences, Brock University, St. Catharine's, ON J2S 3A1

9. College of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Juneau, AK, USA

10. Département de biologie, Université de Sherbrooke, Sherbrooke, QC J1K 2R1, Canada

*Authors contributed equally

†Corresponding author: kgotanda@brocku.ca

ABSTRACT

Wild populations must continuously adapt to environmental changes or they risk extinction. Such adaptations can be measured as phenotypic rates of change and can allow us to predict patterns of contemporary evolutionary change. About two decades ago, a dataset of phenotypic rates of change in wild populations was compiled. Since then, researchers have used (and expanded) this dataset to look at microevolutionary processes in relation to specific types of human disturbances. Here, we have updated the dataset adding 5257 estimates of phenotypic changes and used it to revisit established patterns of contemporary evolutionary change. Using this newer version, containing 6920 estimates of phenotypic changes, we revisit the conclusions of four published articles. We then synthesize the expanded dataset to compare rates of change across different types of human disturbance. Analyses of this expanded dataset suggests that: 1) a small absolute difference in rates of change exists between human disturbed and natural populations, 2) harvesting by humans results in larger rates of change than other types of disturbances, 3) introduced populations have increased rates of change, and 4) body size does not increase through time. Overall, findings from earlier analyses have largely held-up in analyses of our new dataset that encompass a much larger breadth of species, traits, and human disturbances. Lastly, we found that types of human disturbances affect rates of phenotypic change and we call for this database to serve as a stepping stone for further analyses to understand patterns of contemporary evolution.

KEYWORDS

Phenotypic traits, contemporary evolution, rapid evolution, anthropogenic disturbance, human disturbance, Darwins, Haldanes

INTRODUCTION

The rate at which populations can adapt to environmental change will determine their ability to persist, thrive, and expand in a changing world. This adaptation is defined by changes in organismal phenotypes (as opposed to genotypes) – because phenotypes interface with the environment and thus are the direct determinants of fitness (Alberti et al., 2017; Endler, 1986; Hendry, 2017; Schluter, 2000). Historically, adaptive phenotypic changes were thought to be very slow, as exemplified by Charles Darwin's statement that "*we see nothing of these slow changes in progress until the hand of time has marked the long lapse of ages*" (Darwin, 1859). This assumption began to crumble with the accumulation of studies documenting so-called "rapid evolution." Some famous early examples of this phenomenon included industrial melanism in peppered moths (Kettlewell, 1973), body size in mice colonizing islands (Berry, 1964), body size and colour in house sparrows invading North America (Johnston & Selander, 1964), and resistance of plants to pollutants found in mine tailings (Antonovics & Bradshaw, 1970). In most of these early examples from nature, it was unclear whether the observed phenotypic changes were genetic as opposed to plastic. It was therefore game-changing when a series of common-garden experiments confirmed that at least some rapid phenotypic changes seen in nature are, in fact, genetically based (Al-Hiyaly, Mcneilly, & Bradshaw, 1990; Reznick, 1982; Stearns, 1983; Wu & Kruckeberg, 1985).

At the end of the 20th century, it remained unclear if the documented cases of rapid phenotypic change were rare exceptions or the tip of the iceberg. Resolving this uncertainty required broader literature surveys and a quantitative standard for calculating and comparing rates of phenotypic change. A precedent already existed in the literature because paleontologists had long been calculating rates of phenotypic change in fossil time series (Gingerich, 1983, 1993; Haldane, 1949). Following that precedent, Hendry and Kinnison (1999) combed the literature for examples of phenotypic change on contemporary time scales and calculated rates of change using two classic metrics ("darwins" and

"haldanes"). The authors concluded that "[…] *evolution as hitherto considered "rapid" may often be*

the norm and not the exception" (Hendry & Kinnison, 1999). They further advocated use of the general

term "contemporary evolution" because "rapid evolution" requires formal confirmation of

exceptionally rapid rates. This original paper, and the follow-up analysis of Kinnison and Hendry

(2001), opened the flood-gates to a series of influential papers analyzing the dataset of rates of

phenotypic change to answer a series of evolutionary questions (Alberti et al., 2017; Crispo et al., 2010;

Gorné & Díaz, 2019; Gotanda, Correa, Turcotte, Rolshausen, & Hendry, 2015; Hendry, Farrugia, &

Kinnison, 2008; Hendry & Kinnison, 2001; Palkovacs, Wasserman, & Kinnison, 2011; Westley, 2011).

The studies also made the case that evolution operates on scales large enough to have ecological

outcomes, leading the way into the field of eco-evolutionary dynamics (Des Roches et al., 2018;

Fitzpatrick et al., 2015).

The number of studies available for calculating rates of phenotypic change has increased

dramatically over the last decade or so. Compared to the last published version of the dataset that

contained only populations in the wild (Alberti et al., 2017), the new dataset has increased from 1663 to

6920 rates, 89 to 184 studies, 175 to 1540 systems (Box 1), and 155 to 297 species. Hence, our first

goal in the current paper is to present a new quality-controlled and much-expanded database of

contemporary phenotypic changes in nature – The Phenotypic Rates of Change Evolutionary and

Ecological Database (PROCEED) Version 5.0. This new database is publicly available at Dryad and

https://proceeddatabase.weebly.com/. Our second goal was to use the new dataset to revisit previous

analyses and conclusions based on earlier versions of the dataset. Specifically, we wanted to ask four

questions: 1. Does the evidence still support the conclusion from Hendry et al., (2008) that phenotypic

change is greater in human-disturbed systems than in more "natural" systems (see also Alberti et al.,

2017)? 2. Does the evidence still support the conclusion from Darimont et al., (2009) that phenotypic

changes are most rapid when humans act as predators, such as during harvesting (see also Sharpe &

Hendry 2009)? 3. Does the evidence still support the conclusion from Westley (2011) that introduced

populations do not drive particularly rapid phenotypic changes, relative to non-introduced populations?

4. Does the evidence still support the conclusion from Gotanda et al., (2015) that no evidence exists for

microevolutionary trends toward increasing body size – as had been suggested to follow from "Cope's

Rule" (J. Kingsolver & Pfennig, 2004)?

The inclusion of 5257 additional data entries might change earlier conclusions for either of two

reasons: (1) biases in earlier data compilations such as underrepresented or missing taxa and

disturbances, or (2) statistical limitations including small sample sizes associated to taxonomic levels or

types of studies (i.e., genetic; Box 1). To assess the first possibility, we re-analyze the new dataset

using the same methods as the original authors: that is, Hendry et al., (2008), Darimont et al., (2009),

Westley (2011), and Gotanda et al., (2015). Through this approach, we can assess if previous

approaches yield similar conclusions following the accumulation of more data. To assess the second

possibility, we use updated statistical models in a comprehensive approach to ask: 5. Do any types of

human disturbances stand out in terms of their effects on contemporary rates of phenotypic change

(Pelletier & Coltman, 2018). We envision this last analysis as a precursor to what will surely be

additional analyses of the new (and future) dataset with current and future statistical approaches. To

synthesize, we propose a new perspective on how to study contemporary rates of change in future

studies.

METHODS & RESULTS

Database development

The current dataset has a series of notable changes relative to earlier versions of the dataset. First, we

added new data that met the necessary criteria (phenotypic traits were quantified from natural

populations of the same species either at two time points in the same populations or in two populations

with known divergence time), and recorded meta-data as described in Box 1. Second, we modified and

expanded the categorization of types of human disturbances as defined in Box 2. Third, we checked

both old and new data entries to fix any errors. These efforts were facilitated by – whenever possible –

obtaining summary data (means, sample sizes, and standard deviations) directly from tables, figures, or

by contacting authors, from which we calculated rates of change (details below).

The data

Here we outline processes common to all analyses. First, all statistical analyses were performed

in R environment 4.0.5 (R Core Team, 2021). Second, we only include studies where the number of

generations elapsed was 300 or fewer, which is suitable for analyses of "contemporary" change

(Hendry et al., 2008). Third, most analyses were conducted using both darwins and haldanes – because

the two metrics have different biological and statistical properties, as well as different data

requirements (Gingerich, 1993; Hendry & Kinnison, 1999; Hunt, 2012). Darwins are defined as the

proportional change in the mean trait value in units of *e* per million years and are appropriate for data

on a ratio scale, but not an interval scale (Box 1). Darwins were calculated as

$$D = \frac{ln\left(\underline{X_2}\right) - ln\left(\underline{X_1}\right)}{10^6\, years},$$

133   where $\underline{X_1}$ and $\underline{X_2}$ are either mean trait values for one population at times 1 and 2 or mean trait values

134   for two populations that had a common ancestor at a known time in the past, which then scales "time"

135   as million years in the denominator. Haldanes are the change in the mean trait value in standard

136   deviations per generation. Haldanes were calculated as

$$H = \frac{\left(\frac{X_2}{SD_p}\right) - \left(\frac{X_1}{SD_p}\right)}{g},$$

where $\underline{X}_1$ and $\underline{X}_2$ are the mean trait values as explained above, SD_p is the pooled standard deviation of

the two samples, and g is the elapsed time in generations (i.e., number of years divided by generation

length (Box 1)).

The response variable in all subsequent analyses was the numerator of the rate metric (i.e., darwin

or haldane numerator). For all analyses (except questions 3 and 4) we used the mean amount of

phenotypic change for a given system (Box 1) to avoid nonindependence of data points within a system

(Hendry & Kinnison, 2001). Finally, we calculated effect sizes using partial η^2 to compare the original

studies to the updated ones. We now present analyses specific to each question, their respective results,

and a brief discussion of results.

THE QUESTIONS

Question 1: Are rates of phenotypic change greater in human-disturbed systems?

Humans cause particularly dramatic environmental changes, and so we might expect human

disturbance to accelerate rates of phenotypic change. Consistent with this idea, Hendry et al., (2008)

reported that phenotypic rates of change were higher for populations in human-disturbed systems than

for populations in more "natural" systems that were not subject to direct human disturbance. To

replicate that original analysis with our new dataset, we used analysis of covariance (ANCOVA) to

assess whether the absolute amount of phenotypic change (darwin or haldane numerator) differed

between the two general contexts (human-disturbed or natural) while controlling for the length of the

time interval (years for darwins, generations for haldanes).

As in Hendry et al., (2008), our new dataset suggests that rates of change were generally higher

in human-disturbed systems compared to natural systems (Fig. 1; Fig. S1). The difference between

contexts (human-disturbed versus natural) was, however, reduced in our new dataset, $\eta^2 = 0.018$,

compared to the original analysis, $\eta^2 = 0.115$, (Fig. 1; Fig. S1; Table 1). This smaller difference could

result from confounding effects of multiple types of disturbances (Galton, 1886; Kelly & Price, 2017;

Pelletier & Coltman, 2018). That is, it can be difficult to assign a single type of human disturbance to a

particular system. For example, climate change is likely to affect all systems indirectly, including

systems we have classified as "natural." Here, we classified disturbance as climate change only if the

original study specifically tested for an effect of climate change. We will later return to the influence of

these multiple disturbances on inferences about contemporary evolution. Our expanded dataset could

also mean that effects like winnowing, which would reduce the number of populations with low rates

of phenotypic change because they are more likely to go extinct (Hendry et al., 2008), do not appear to

have a strong overall effect.

By comparing estimates from wild populations versus those from common-garden or animal

model analyses (Box 1), Hendry et al., (2008) concluded that plasticity likely contributed substantially

to the rate differences between human-disturbed and more natural contexts (Fig. S2; Fig. S3). This

suggestion arose because the difference between contexts was lower when common-garden or animal

model studies were used – and because large changes could sometimes be seen immediately after a

disturbance (Hendry et al., 2008). Such patterns also occurred in the present study supporting those

original inferences (Fig. S2; Fig. S3). At the same time, it is important to note that genetic changes

definitely occurred in a number of studies, but that many of the most disturbed contexts (e.g.,

harvesting) are not particularly amenable to the assessment of genetically-based phenotypic change.

Only 24% of the phenotypic change entries in our dataset could be labelled are genetically based,

provided an adequate experimental design.

Question 2: Are particularly rapid and consistent changes associated with harvesting?

A particularly strong and consistent disturbance that directly impacts some populations occurs

when humans act as a predator, such as in cases of harvesting. To explore this idea, Darimont et al.,

(2009) took the human-disturbed versus natural distinction of Hendry et al., (2008) and divided the

human-disturbed systems into those experiencing direct harvesting versus those experiencing other

forms of human disturbances. After adding more data, especially to the harvesting category, Darimont

et al., (2009) reported that populations subject to harvesting had increased phenotypic rates of change

compared to other types of human disturbance and also compared to natural systems. To replicate that

original analysis with our updated dataset, we used ANCOVA to compare darwin numerators

associated with harvesting to those associated with other types of human disturbances (combined) and

those associated with more natural systems – while including time (years) as a covariate. We ran

ANCOVAs using both mean and maximum rates of change per system to replicate the original study.

As in Darimont et al., (2009), our new data suggested that phenotypic changes associated with

harvesting are greater than those associated with other types of human disturbances or natural systems

(Fig. 2; Fig S4; Table 1). However, the effect size (partial η^2) of human disturbance (three contexts:

harvesting, other human disturbance, or natural) decreased from 0.180 in the original study to 0.010 in

the updated dataset (Table 1). We suggest that this decrease in effect size is mainly driven by large

datasets containing many study systems with high variation. As an example, when we remove the Oke

et al., (2020) and Clark et al., (2018) datasets, the effect size of human disturbance increases to 0.139

(Fig. S5). These datasets correspond to salmonid data for 1072 populations from four species over a

time frame of up to 77 years. Of these populations, 60% are (or were) likely subject to harvest – and so

needed to be added to that category in the present analysis. The remaining 40% of populations were

added to the natural category. However, the strength and type of harvest is expected to be highly

variable among those species and populations. We note that this variability in harvest strength is likely

true for most harvested populations. This issue again highlights the difficulty of unambiguously

assigning human disturbances to systems that are surely experiencing multiple types of disturbance that

vary in type and intensity across systems.

Despite the lower effect size for harvesting in our new dataset, we note that harvested systems

represented 35 of the 50 largest mean phenotypic changes in the dataset. Hence, it seems likely that

humans as predators generate a diversity of rates of change – from many slow rates to some truly

exceptionally high rates. Our findings continue to support the claims of fisheries and hunting wildlife

scientists who have long argued the lasting effects of harvesting on some (but not all) natural

populations (Kuparinen & Festa-Bianchet, 2017; Morrissey, Hubbs, & Festa-Bianchet, 2021; Pigeon,

Festa-Bianchet, Coltman, & Pelletier, 2016; Van de Walle, Pigeon, Zedrosser, Swenson, & Pelletier,

2018).

Question 3: Do introduced populations show particularly rapid rates of change?

When a species is introduced into a new location, it often experiences massive shifts in biotic

and abiotic conditions, which are expected to cause particularly rapid phenotypic changes (Carroll,

2007; Cox, 2004; Huey, Gilchrist, & Hendry, 2005; Kinnison, Unwin, & Quinn, 2008). However,

Westley (2011) used an earlier version of this dataset to report that introduced populations do not – in

fact – evince particularly strong or consistent phenotypic changes when compared to non-introduced

populations. To replicate their analysis, we first averaged rates by species and then used ANCOVA to

compare the magnitude of phenotypic change (darwin and haldane numerators) between introduced

(disturbance classified as introduced) and non-introduced (all other disturbance categories) populations

with time (years or generations) as a covariate (Westley, 2011).

We found that, on average, introduced populations have higher rates of change than non-

introduced populations (Fig. 3; Fig. S6). Although the difference between introduced and non-

introduced populations is marginal, the addition of our new data increased the effect size (Table 1).

This finding is consistent with the evidence that introduced populations do sometimes show very rapid

rates of change (Fig. 3). For example, zebra mussels (*Dreissena polymorpha*) introduced in European

lakes rapidly shifted their growth rates (Czarnołeski et al., 2005) and Eastern grey kangaroos,

(*Macropus giganteus*) introduced from Tasmania to Maria Island rapidly shifted their behaviour when

no longer subject to predation (Blumstein & Daniel, 2003). This is consistent with Westley's

conclusion that a small number of introduced species having very high rates of change drove the

perception that introduced species show rapid change. Indeed, the updated dataset has both very high

rates and very low rates of change in introduced populations (Fig. 3).

Populations introduced to a new habitat likely experience abrupt directional selection that drives

rapid evolutionary rates compared to non-introduced populations (Carroll, 2007; Cox, 2004). Once a

population achieves adaptive optima and is more locally adapted, phenotypic rates of change are

expected to decline with time (Fig. 3). This expectation is consistent with the declining rates of change

with time since introduction (Figs. 1 & 2 in Westley 2011), although that pattern is also consistent with

other processes (Hendry & Kinnison, 2001). These processes include averaged rates over longer time

spans or depletion of genetic variation (Kinnison & Hairston, 2007). Finally, we note that both

introduced and non-introduced populations are also experiencing other types of human disturbances

that can influence their rates of evolution, thus potentially obscuring effects of introduction per se.

Question 4: Are body sizes increasing through time as suggested by Cope's Rule?

Cope's Rule states that lineages generally evolve larger body sizes over evolutionary time

(Cope, 1885). Based on an analysis of selection estimates, Kingsolver and Pfennig (2004) argued that

evidence of directional selection for larger body size in contemporary populations is consistent with a

microevolutionary explanation for Cope's Rule. However, when looking for evidence of general trends

toward larger mean body size in contemporary populations, Gotanda et al., (2015) found no such trend

and instead found a general trend towards decreasing body size.

To replicate the analysis of Gotanda et al., (2015), we used the raw (i.e., signed, rather than

absolute value) estimates of haldane and darwin numerators for allochronic studies (Box 1). The

original analysis did not average per system or species, and so all new analyses are similarly based on

individual rates. We first square-root transformed 2-D traits (e.g., surface area) and cube-root

transformed 3-D traits (e.g., volume or mass) and then compared rates for body size traits to rates for

other types of traits in a one-tailed Wilcoxon rank-sum test (see Gotanda et al., 2015 for details). We

next conducted a sign test to determine whether the change in body size within populations was more

commonly positive or negative. Finally, we re-ran the sign tests excluding rates calculated from

populations known to be subject to harvesting – as harvesting is expected to cause particularly rapid

decreases in body size (Darimont et al., 2009; Sharpe & Hendry, 2009; see question 2).

When compared to other traits, rates of change for body size are not larger (Table 2; Fig S7). As

in the original analysis of Gotanda et al., (2015), rates of change in body size are more often negative

(organisms are getting smaller overall), even when excluding populations subject to harvesting (Table

3; Fig. 4; Fig S8). Further, rates of change for body size are not more positive (or less negative) when

compared to other traits (Table 3; Fig. S8), and the type of human disturbance does not appear to affect

rates of change on body size (Fig. 4). These results are matched by other recent analyses of body size

trends in a variety of taxa (Gardner, Peters, Kearney, Joseph, & Heinsohn, 2011; Sheridan & Bickford,

2011). For instance, recent research suggests that body sizes are broadly decreasing as a response to

climate change perhaps due to effects of increasing temperature and variable precipitation on

organismal development and growth (Fryxell et al., 2020; Sheridan & Bickford, 2011).

Question 5: Does any type of disturbance stand out with respect to rates of change?

Based on the previous studies we have now revisited, as well as more recent reviews (Pelletier

& Coltman, 2018), we expect associations between high rates of phenotypic change and particular

types of human disturbances such as pollution (Hamilton, Rolshausen, Uren Webster, & Tyler, 2017),

harvesting (Sullivan, Bird, & Perry, 2017), or landscape change (Legrand et al., 2017). Using our

extensive dataset, we are now able to ask if these or other types of human disturbances stand out with

respect to rates of phenotypic change.

283   To answer this question, we used a more advanced analysis – in contrast to the above questions.

We used general linear models in which the response variables were log10-transformed absolute

darwin (or haldane) numerators, and independent variables included type of human disturbance, time

(years and generations, log10-transformed), and type of study (genetic or phenotypic). As in question 4,

we first square-root transformed 2-D traits (e.g., surface area) and cube-root transformed 3-D traits

(e.g., volume or mass) before log transforming the rates. Finally, we used Tukey post-hoc tests

(Hothorn, Bretz, & Westfall, 2008) to explore the differential effects of disturbances on differences in

rates of evolution.

291   This new analysis suggests that time (years, but not generations) had a significant positive

impact on darwin rates of change, but type of study (genetic vs. phenotypic) did not (Table 4; Table

S2). We also found that systems associated with pollution have the highest rates of change and that

systems associated with climate change have the slowest rates of change (Fig. 5; Fig S9). We should

note that, within our dataset, most systems looking at the impact of pollution were plants, which might

or might not influence the high rates for this type of disturbance. In short, more work needs to be done

to explore potential interactive effects of disturbance type and taxonomic group on rates of change.

298   Surprisingly, on average, phenotypic changes associated with climate change were amongst the

slowest. We suspect that climate change has broad reaching effects and is especially difficult to assign

as a particular sole causal force. Indeed, climate change must be – at some level – influencing all or

most systems included in the dataset, regardless of the disturbance category to which we assigned

them. Furthermore, studies focusing on trait change in response to climate change are likely to focus on

the more gradual aspects of climate change, such as shifting seasonality and temperature increases

(Parmesan & Yohe, 2003), rather than abrupt aspects such as heat waves, storms, and droughts. This

focus could therefore underestimate the evolutionary impacts of climate change. More importantly,

climate change can be a particularly noisy environmental driver and so, is especially prone to temporal

averaging (Hendry & Kinnison, 1999). In fact, few studies in nature follow populations long enough

and at a fine enough temporal resolution to detect fast and large phenotypic changes (e.g., Grant &

Grant, 2002). For these reasons, we replicate the analyses by including all populations affected by

climate change as "natural" (Fig S10). The results from this analysis confirm that whether we have a

disturbance category dedicated to climate change, or if we include climate change with "natural"

systems, our conclusions do not change: climate change and natural systems are still amongst the

slowest rates of evolution. Regardless, our new analysis provides the most comprehensive hypothesis

(Fig. 5) for how various types of disturbance differ in their effects on rates of phenotypic change.

DISCUSSION

Conclusions from analyses of earlier datasets of contemporary phenotypic change have largely

been upheld in analyses of our new dataset that encompasses a much larger breadth of species, traits,

and human disturbances. 1) Human disturbed systems have slightly larger rates of phenotypic change

then do natural systems (Fig. 1). 2) Harvesting by humans results in larger rates of change compared to

other types of disturbances (Fig. 2). 3) Introduced populations have higher rates of change than non-

introduced populations (Fig. 3). 4) There is no trend for increasing body size through time (Fig. 4). 5)

Systems affected by pollution have larger rates of change compared to other types of disturbances (Fig.

5).

Overall, contemporary rates of phenotype change range from very slow to very fast (relative to other rates), with the latter typically gaining the most attention. This pattern is found in other datasets focusing on estimates of selection in natural populations (Kingsolver et al., 2001; Siepielski et al., 2013). Our database includes some striking examples of rapid phenotypic rates of change. For example, horn size in bighorn sheep (*Ovis canadensis*) decreased by 10% over 19 years when targeted by trophy hunters (Pigeon et al., 2016); body length in harvested Chinook salmon (*Oncorhynchus tshawytscha*) in the Yukon River of Alaska has decreased by 5-7% since the late 1970s (Ohlberger et al., 2020); zinc tolerance in tufted hairgrass (*Deschampsia cespitosa*) in zinc-contaminated soils increased by 80% over 26 years (Al-Hiyaly et al., 1990); and total egg count in soapberry bugs (*Jadera haematoloma*) adapting to an introduced host decreased by 8% over 38 years (Carroll, Klassen, & Dingle, 1998). Human influences clearly shape these and many other phenotypic responses; yet the high levels of variation in rates of response make the most consistent and dramatic influences hard to confirm.

Our primary goal in the present paper is to make the new database available to any researchers seeking to leverage phenotypic rates of change to answer questions in ecology, evolution, or conservation biology. For instance, we anticipate that researchers will use the new database to answer a series of questions relevant to contemporary evolution and for better understanding and predicting how wild populations will adapt (or not) to human disturbances. Some examples of questions left unanswered, or only lightly broached in prior reviews include: Do different organisms or trait types evolve at different rates? Are different disturbances generating confounding or synergistic effects? What are the upper and lower limits to sustainable versus unsustainable evolutionary rates in nature?

The use of this new database should be accompanied with an understanding of its limits and the resulting caveats of inference.

1. We suspect a strong ascertainment bias where researchers tend to study systems they suspect are strongly affected by human disturbances (e.g., harvested fish populations) or where phenotypic changes are already documented (e.g., phenology). Thus, average effect sizes for a given disturbance might well decrease with the accumulation of more diverse and objective sets of data – as we have shown above with the decrease in average rates of change for harvested systems after we added the 1320 rates from Oke et al., (2020) and Clark et al., (2018) (Fig. S5). These rates associated with body size declines in salmonids were calculated from data collected by the Alaska Department of Fish and Game and collaborators. As such, these rates represent a very broad sampling of populations across Alaska, not a narrow dataset on just the most impacted populations. Another form of ascertainment bias occurs when disturbances cause some populations to go extinct, in which case their rate of phenotypic change cannot be measured (Hendry et al., 2008). It is currently unknown whether this winnowing effect of extinction (Hendry et al., 2008) biases rates upward (i.e., slower-changing populations are more likely to perish) or downward (i.e., faster-changing populations are more likely to perish – likely because they are experiencing more disturbance).

2. Seeking to attribute a particular phenotypic change to a single disturbance (e.g., climate change) is problematic because most populations will be subject to multiple disturbances – and the degree of a given disturbance will vary dramatically among systems. Thus, we encourage future work to consider variation in disturbance intensity (rather than just presence). As examples, one could relate the strength of harvesting on populations (e.g., local catch rates data) to the rate of change in size or age (e.g., Sharpe & Hendry, 2009) or the rate of climate change experienced by populations (e.g., local

temperature change) to their specific rate of trait change (e.g., Franks, Sim, & Weis,

2007; Jenni & Kéry, 2003).

3. The genetic and plastic contributions to trait change remain uncertain for most traits in

most systems (Merilä & Hendry, 2014). As more studies accumulate, we might be able

to profitably analyze only genetically based phenotypic change, such as from common-

garden, reciprocal transplants, or animal-model studies. Regardless, we highlight the

importance of controlled experiments in combination with phenotypic change in the

wild for several reasons: some populations cannot be analyzed with common-garden or

animal-model approaches, phenotypes measured in the lab might not be natural, and the

"wild" is where organisms interact with their environment (Hendry, 2017).

The study of eco-evolutionary dynamics was born from the recognition of widespread

contemporary evolution and the cyclical feedback with ecological processes (Hendry, 2017). With this

dataset, we identified some patterns of contemporary evolution (e.g., pollution is the strongest human

influence on phenotypic rates of change) and we can use this information to broaden our understanding

of eco-evolutionary feedbacks in the wild. We suggest that the next step is to understand how the

feedback dynamics implicit to eco-evolutionary dynamics mechanistically shape emergent patterns of

contemporary evolution: are some systems more feedback prone and thus have faster or slower rates of

evolution? To address such feedback dynamics, we support the development of a comprehensive

database which includes not only phenotypic rates of change but also ecological and environmental

rates of change and estimates of selection.

Whilst this dataset can be a steppingstone to further our understanding of eco-evolutionary

dynamics, we also advocate that this dataset can allow us to move beyond 'traditional' ecological and

evolutionary patterns and start to consider societal consequences. In the past, the starting point for the

study of contemporary evolution was often population dynamics: "*perhaps the greatest contribution*

that evolutionary rate estimates will ultimately make is an awareness of our own role in the present

microevolution of life and cautious consideration of whether populations and species can adapt rapidly

enough to forestall the macroevolutionary endpoint of extinction" (Hendry & Kinnison, 1999). What is

now needed is a widespread and formal assessment of how trait changes, alone and in aggregate (e.g.,

Kinnison et al., 2008; Westley, Ward, & Fleming, 2013, Calow, 1987), affect fitness, function, and

species interactions in ways that shape ecological processes such as population dynamics, communities,

and ecosystems, as well as the societal consequences such as the health and well-being of people (Des

Roches et al., 2018; Hendry, 2017; Hendry, Gotanda, & Svensson, 2017; Palkovacs, Kinnison, Correa,

Dalton, & Hendry, 2012; Stange, Barrett, & Hendry, 2021; Fig. 6). For example, we propose that more

studies need to empirically study such feedbacks while also assessing the impacts on nature's

contributions to people (NCPs). Oke et al., (2020) provide a compelling example, by comparing the

average body size of Alaskan chinook salmon pre-1990 to the average post-2010. The authors estimate

that the overall average 8% decrease in body length could – all else being equal – translate into a 16%

decrease in number of eggs per female, a 28% decrease in transport of marine-derived phosphorous

into freshwater, a 26% reduction in average number of meals provided per fish for people in

subsistence communities, and a 21% decrease in price per pound for commercial fishers.

We would like to close by re-emphasizing that the most dramatic progress will be made through

studies that explicitly examine the *consequences* of phenotypic change. That is, more studies should

formally calculate the importance of observed (and predicted) phenotypic change for all levels of

ecology (populations, communities, ecosystems) and for people (nature's contributions to people).

Determining the genetic and plastic contributions to that change (and its consequences) can then help to

determine the limits and opportunities for enhancing or arresting trait changes via conservation and

management actions. We here present a hypothetical scenario of such a study using observed rates of

change (Fig. 6) where the amount of phenotypic change is correlated with an ecological process that is

linked to nature's contributions to people. The present paper is not the end of an inspiring era of

contemporary evolution – it is instead part of the start of an upcoming critical era of contemporary eco-

evolutionary dynamics.

AKNOWLEDGEMENTS

This work has been supported by NSF OIA 1826777 and NSF OAI 1849227 received by Micheal

Kinnison. We thank the numerous researchers that shared their data and contributed to the development

of The Phenotypic Rates of Change Evolutionary and Ecological Database (PROCEED) Version 5.0.

We are especially grateful to both the Carnegie Museum of Natural History and the Powdermill Avian

Research Center for sharing their data.

[dataset] Sarah Sanderson, Marc-Olivier Beausoleil, Rose E. O'Dea, Zachary T. Wood, Cristian

Correa, Victor Frankel, Lucas D. Gorné,Grant E. Haines, Michael T. Kinnison, Krista B. Oke, Fanie

Pelletier, Felipe Pérez-Jvostov, Winer Daniel Reyes-Corral, Yanny Ritchot, Freedom Sorbara, Kiyoko

M. Gotanda, and Andrew P. Hendry; 2021; Database reference: Phenotypic Rates of Change

Evolutionary and Ecological Database (PROCEED). https://proceeddatabase.weebly.com; Version 5.0

Box 1. Definitions of the different types of data included in the data set.

1. System: Each system has its own unique species, disturbance, and location (population). Within a given system, you can have multiple traits – for example, tarsus length and fledging date.
2. Disturbance: See Box 2.
3. Trait Classification: Traits were classified as determined by Kingsolver and Diamond 2011: size, other morphology, phenology, other life history traits, behaviour, or physiology.
4. Study type: Studies were determined to be phenotypic in nature if traits were studied in natural populations and studies were determined to be genetic in nature if they used common-garden or quantitative genetic methods (i.e., animal model analyses). We note that studies classified as phenotypic could have a genetic basis (see introduction), but that we could not determine that from the methods of the study.
5. Design: Data were determined to be either allochronic (same population/different time points) or synchronic (populations with known divergence time).
6. Data scale: Data were determined to be ratio (constant interval with a precise zero; e.g., mass or length) or interval (constant interval with an arbitrary zero; e.g., temperature or time of day).
7. Generation time: The amount of time to reproductive age, given in years.

Box 2. Definitions of the different types of human disturbance categories used in the updated database.

🟣 Introductions: "*when humans transferred a species to a new geographical location, and comparisons were then made between introduced and ancestral populations* (Carroll et al., 2005)".

🔵 Response to introductions: when a local population of a species is responding to the introduction of a species.

🟡 Landscape change: when any type of modification to the habitat of a population occurs.

🟣 Hunting/Harvesting: when there is hunting or harvesting of a species by humans.

⚫ Pollution: when any type of pollutant enters a system.

🟠 Climate change: when the objectives of the study are directly linked to climate change.

🟢 Natural: established populations that are not subject to obvious human impacts (as listed above). Generally, these studies were the long-term monitoring of natural populations.

REFERENCES

Alberti, M., Correa, C., Marzluff, J. M., Hendry, A. P., Palkovacs, E. P., Gotanda, K. M., … Zhou, Y.
(2017). Global urban signatures of phenotypic change in animal and plant populations.
Proceedings of the National Academy of Sciences, *114*(34), 8951–8956. doi:
10.1073/pnas.1606034114
Al-Hiyaly, S. A. K., Mcneilly, T., & Bradshaw, A. D. (1990). The effect of zinc contamination from
electricity pylons. Contrasting patterns of evolution in five grass species. *New Phytologist*,
114(2), 183–190. doi: 10.1111/j.1469-8137.1990.tb00389.x
Antonovics, J., & Bradshaw, A. D. (1970). Evolution in closely adjacent plant populations. VIII. Clinal
patterns at a mine boundary. *Heredity*, *25*, 349–362. doi: 10.1038/hdy.1970.36
Berry, R. J. (1964). The evolution of an island population of the house mouse. *Evolution*, *18*(3), 468–
483. doi: 10.2307/2406357
Blumstein, D. T., & Daniel, J. C. (2003). Foraging behavior of three Tasmanian macropodid marsupials
in response to present and historical predation threat. *Ecography*, *26*(5), 585–594.
Calow, P. (1987). Towards a definition of functional ecology. Functional Ecology, 1(1), 57–61. doi:
10.2307/2389358
Carroll, S. P. (2007). Natives adapting to invasive species: Ecology, genes, and the sustainability of
conservation. Ecological Research, 22(6), 892–901. doi: 10.1007/s11284-007-0352-5
Carroll, S. P., Klassen, S. P., & Dingle, H. (1998). Rapidly evolving adaptations to host ecology and
nutrition in the soapberry bug. *Evolutionary Ecology*, *12*(8), 955–968. doi:
10.1023/A:1006568206413
Carroll, S. P., Loye, J. E., Dingle, H., Mathieson, M., Famula, T. R., & Zalucki, M. P. (2005). And the
beak shall inherit—Evolution in response to invasion. *Ecology Letters*, *8*(9), 944–951. doi:
10.1111/j.1461-0248.2005.00800.x
Clark, J., Brenner, R., & Lewis, B. (2018). Compiled age, sex, and length data for Alaskan salmon,
1922-2017. doi: 10.5063/F1707ZTM
Cope, E. D. (1885). On the evolution of the Vertebrata, progressive and retrogressive (continued). The
American Naturalist, 19, 341–353.
Cox, G. W. (2004). *Alien species and evolution: The evolutionary ecology of exotic plants, animals,*
microbes, and interacting native species. Washington: Island Press.
Crispo, E., Dibattista, J. D., Correa, C., Thibert-plante, X., Mckellar, A. E., Schwartz, A. K., …
Hendry, A. P. (2010). The evolution of phenotypic plasticity in response to anthropogenic
disturbance. *Evolutionary Ecology Research*, *12*, 47–66. doi: 10.1088/0004-637X/742/2/126
Czarnołeski, M., Kozlowski, J., Lewandowski, K., Mikolajczyk, M., Müller, T., & Stanczykowska, A.
(2005). Optimal resource allocation explains changes in the zebra mussel growth pattern
through time. *Evolutionary Ecology Research*, *7*, 821–835.
Darimont, C. T., Carlson, S. M., Kinnison, M. T., Paquet, P. C., Reimchen, T. E., & Wilmers, C. C.
(2009). Human predators outpace other agents of trait change in the wild. *Proceedings of the*
National Academy of Sciences, *106*(3), 952–954. doi: 10.1073/pnas.0809235106
Darwin, C. (1859). *On the Origin of Species by means of natural selection, or preservation of flavoured*
races in the struggle for life. London: John Murray.
Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., … Palkovacs,
E. P. (2018). The ecological importance of intraspecific variation. *Nature Ecology & Evolution*,
2(1), 57–64. doi: 10.1038/s41559-017-0402-5
Endler, J. A. (1986). *Natural selection in the wild*. Princeton University Press.

Fitzpatrick, C. R., Agrawal, A. A., Basiliko, N., Hastings, A. P., Isaac, M. E., Preston, M., & Johnson, M. T. J. (2015). The importance of plant genotype and contemporary evolution for terrestrial ecosystem processes. *Ecology*, *96*(10), 2632–2642. doi: 10.1890/14-2333.1

Franks, S. J., Sim, S., & Weis, A. E. (2007). Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. *Proceedings of the National Academy of Sciences*, *104*(4), 1278–1282. doi: 10.1073/pnas.0608379104

Fryxell, D. C., Hoover, A. N., Alvarez, D. A., Arnesen, F. J., Benavente, J. N., Moffett, E. R., … Palkovacs, E. P. (2020). Recent warming reduces the reproductive advantage of large size and contributes to evolutionary downsizing in nature. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1928), 20200608. doi: 10.1098/rspb.2020.0608

Galton, F. (1886). Regression towards mediocrity in hereditary stature. *The Journal of the Anthropological Institute of Great Britain and Ireland*, *15*, 246–263.

Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., & Heinsohn, R. (2011). Declining body size: A third universal response to warming? *Trends in Ecology & Evolution*, *26*(6), 285–291. doi: 10.1016/j.tree.2011.03.005

Gingerich, P. (1983). Rates of evolution: Effects of time and temporal scaling. *Science*, *222*, 159–162. doi: 10.1126/science.222.4620.159

Gingerich, P. (1993). Quantification and comparison of evolutionary rates. *American Journal of Science*, *293-A*, 453–478.

Gorné, L. D., & Díaz, S. (2019). Meta-analysis shows that rapid phenotypic change in angiosperms in response to environmental change is followed by stasis. *The American Naturalist*, *194*(6), 840–853. doi: 10.1086/705680

Gotanda, K. M., Correa, C., Turcotte, M. M., Rolshausen, G., & Hendry, A. P. (2015). Linking macrotrends and microrates: Re-evaluating microevolutionary support for Cope's rule. *Evolution*, *69*(5), 1345–1354. doi: 10.1111/evo.12653

Grant, P. R., & Grant, B. R. (2002). Unpredictable evolution in a 30-year study of Darwin's finches. *Science*, *296*(5568), 707–711. doi: 10.1126/science.1070315

Haldane, J. B. S. (1949). Suggestions as to quantitative measurement of rates of evolution. *Evolution*, *3*(1), 51–56. doi: 10.2307/2405451

Hamilton, P. B., Rolshausen, G., Uren Webster, T. M., & Tyler, C. R. (2017). Adaptive capabilities and fitness consequences associated with pollution exposure in fish. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *372*(1712), 20160042. doi: 10.1098/rstb.2016.0042

Hendry, A. P. (2017). *Eco-evolutionary dynamics*. Princeton University Press.

Hendry, A. P., Farrugia, T. J., & Kinnison, M. T. (2008). Human influences on rates of phenotypic change in wild animal populations. *Molecular Ecology*, *17*(1), 20–29. doi: 10.1111/j.1365-294X.2007.03428.x

Hendry, A. P., Gotanda, K. M., & Svensson, E. I. (2017). Human influences on evolution, and the ecological and societal consequences. Philosophical Transactions of the Royal Society B: Biological Sciences, 372(1712). doi: 10.1098/rstb.2016.0028

Hendry, A. P., & Kinnison, M. T. (1999). Perspective: The pace of modern life: Measuring rates of contemporary microevolution. *Evolution*, *53*(6), 1637–1653. doi: 10.2307/2640428

Hendry, A. P., & Kinnison, M. T. (2001). An introduction to microevolution: Rate, pattern, process. *Genetica*, *112*(1), 1–8. doi: 10.1023/A:1013368628607

Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous inference in general parametric models. *Biometrical Journal*, *50*(3), 346–363. doi: 10.1002/bimj.200810425

Huey, R., Gilchrist, G., & Hendry, A. P. (2005). Using Invasive Species to Study Evolution: Case
studies with *Drosophila* and Salmon. In *Species invasions: Insights into ecology, evolution and*
biogeography (p. 139-164). Sinauer Associates.
Hunt, G. (2012). Measuring rates of phenotypic evolution and the inseparability of tempo and mode.
Paleobiology, *38*(03), 351–373. doi: 10.1666/11047.1
Jenni, L., & Kéry, M. (2003). Timing of autumn bird migration under climate change: Advances in
long-distance migrants, delays in short-distance migrants. *Proceedings. Biological Sciences*,
270(1523), 1467–1471. doi: 10.1098/rspb.2003.2394
Johnston, R. F., & Selander, R. K. (1964). House Sparrows: Rapid evolution of races in North
America. *Science*, *144*(3618), 548–550. doi: 10.1126/science.144.3618.548
Kelly, C., & Price, P. D. (2017). Correcting for Regression to the Mean in Behavior and Ecology. *The*
American Naturalist, *166*(6), 700–700. doi: 10.2307/3491232
Kettlewell, H. (1973). *Evolution of melanism: The study of a recurring necessity*. Oxford: Clarendon
Press.
Kingsolver, J. G., Hoekstra, H. E., Hoekstra, J. M., Berrigan, D., Vignieri, S. N., Hill, C. E., … Beerli,
P. (2001). The strength of phenotypic selection in natural populations. *The American Naturalist*,
157(3), 245–261. doi: 10.1086/319193
Kingsolver, J., & Pfennig, D. (2004). Individual-level selection as a cause of Cope's Rule of phyletic
size increase. *Evolution; International Journal of Organic Evolution*, *58*, 1608–1612. doi:
10.1111/j.0014-3820.2004.tb01740.x
Kinnison, M. T., & Hairston, N. G. (2007). Eco-evolutionary conservation biology: Contemporary
evolution and the dynamics of persistence. *Functional Ecology*, *21*(3), 444–454. doi:
10.1111/j.1365-2435.2007.01278.x
Kinnison, M. T., Unwin, M. J., & Quinn, T. P. (2008). Eco-evolutionary vs. Habitat contributions to
invasion in salmon: Experimental evaluation in the wild. *Molecular Ecology*, *17*(1), 405–414.
doi: 10.1111/j.1365-294X.2007.03495.x
Kuparinen, A., & Festa-Bianchet, M. (2017). Harvest-induced evolution: Insights from aquatic and
terrestrial systems. *Philosophical Transactions of the Royal Society B: Biological Sciences*,
372(1712), 20160036. doi: 10.1098/rstb.2016.0036
Law, W., & Salick, J. (2005). Human-induced dwarfing of Himalayan snow lotus, S*aussurea laniceps*
(Asteraceae). *Proceedings of the National Academy of Sciences*, *102*(29), 10218–10220. doi:
10.1073/pnas.0502931102
Legrand, D., Cote, J., Fronhofer, E. A., Holt, R. D., Ronce, O., Schtickzelle, N., … Clobert, J. (2017).
Eco-evolutionary dynamics in fragmented landscapes. Ecography, 40(1), 9–25. doi:
10.1111/ecog.02537
Merilä, J., & Hendry, A. P. (2014). Climate change, adaptation, and phenotypic plasticity: The problem
and the evidence. *Evolutionary Applications*, *7*(1), 1–14. doi: 10.1111/eva.12137
Morrissey, M. B., Hubbs, A., & Festa-Bianchet, M. (2021). Horn growth appears to decline under
intense trophy hunting, but biases in hunt data challenge the interpretation of the evolutionary
basis of trends. *Evolutionary Applications*, *14*(6), 1519–1527. doi: 10.1111/eva.13207
Ohlberger, J., Schindler, D. E., Brown, R. J., Harding, J. M. S., Adkison, M. D., Munro, A. R., …
Spaeder, J. (2020). The reproductive value of large females: Consequences of shifts in
demographic structure for population reproductive potential in Chinook salmon. *Canadian*
Journal of Fisheries and Aquatic Sciences, *77*(8), 1292–1301. doi: 10.1139/cjfas-2020-0012
Oke, K. B., Cunningham, C. J., Westley, P. A. H., Baskett, M. L., Carlson, S. M., Clark, J., …
Palkovacs, E. P. (2020). Recent declines in salmon body size impact ecosystems and fisheries.
Nature Communications, *11*(1), 4155. doi: 10.1038/s41467-020-17726-z

Palkovacs, E. P., Kinnison, M. T., Correa, C., Dalton, C. M., & Hendry, A. P. (2012). Fates beyond traits: Ecological consequences of human-induced trait change. *Evolutionary Applications*, *5*(2), 183–191. doi: 10.1111/j.1752-4571.2011.00212.x

Palkovacs, E. P., Wasserman, B. A., & Kinnison, M. T. (2011). Eco-evolutionary trophic dynamics: Loss of top predators drives trophic evolution and ecology of prey. *PLoS ONE*, *6*(4), e18879–e18879. doi: 10.1371/journal.pone.0018879

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, *421*(6918), 37–42. doi: 10.1038/nature01286

Pelletier, F., & Coltman, D. W. (2018). Will human influences on evolutionary dynamics in the wild pervade the Anthropocene? *BMC Biology*, *16*(1), 1–10. doi: 10.1186/s12915-017-0476-1

Pigeon, G., Festa-Bianchet, M., Coltman, D. W., & Pelletier, F. (2016). Intense selective hunting leads to artificial evolution in horn size. *Evolutionary Applications*, *9*(4), 521–530. doi: 10.1111/eva.12358

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Reznick, D. (1982). The impact of predation on life history evolution in Trinidadian Guppies: Genetic basis of observed life history patterns. *Evolution*, *36*(6), 1236–1250. doi: 10.2307/2408156

Schluter, D. (2000). Ecological character displacement in adaptive radiation. *The American Naturalist*, *156*(4), S4–S16. doi: 10.2307/3079223

Sharpe, D. M. T., & Hendry, A. P. (2009). Life history change in commercially exploited fish stocks: An analysis of trends across studies. *Evolutionary Applications*, *2*(3), 260–275. doi: 10.1111/j.1752-4571.2009.00080.x

Sheridan, J. A., & Bickford, D. (2011). Shrinking body size as an ecological response to climate change. *Nature Climate Change*, *1*(8), 401–406. doi: 10.1038/nclimate1259

Siepielski, A. M., Gotanda, K. M., Morrissey, M. B., Diamond, S. E., DiBattista, J., & Carlson, S. M. (2013). The spatial patterns of directional phenotypic selection. *Ecology Letters*, *16*(11), 1382–1392. doi: 10.1111/ele.12174

Stange, M., Barrett, R. D. H., & Hendry, A. P. (2021). The importance of genomic variation for biodiversity, ecosystems and people. *Nature Reviews Genetics*, *22*(2), 89–105. doi: 10.1038/s41576-020-00288-7

Stearns, S. C. (1983). The genetic basis of differences in life-history traits among six populations of Mosquitofish (*Gambusia affinis*) that shared ancestors in 1905. *Evolution*, *37*(3), 618–627. doi: 10.2307/2408274

Sullivan, A. P., Bird, D. W., & Perry, G. H. (2017). Human behaviour as a long-term ecological driver of non-human evolution. *Nature Ecology & Evolution*, *1*(3), 1–11. doi: 10.1038/s41559-016-0065

Van de Walle, J., Pigeon, G., Zedrosser, A., Swenson, J. E., & Pelletier, F. (2018). Hunting regulation favors slow life histories in a large carnivore. *Nature Communications*, *9*(1), 1100. doi: 10.1038/s41467-018-03506-3

Westley, P. A. H. (2011). What invasive species reveal about the rate and form of contemporary phenotypic change in nature. *The American Naturalist*, *177*(4), 496–509. doi: 10.1086/658902

Westley, P. A. H., Ward, E. J., & Fleming, I. A. (2013). Fine-scale local adaptation in an invasive freshwater fish has evolved in contemporary time. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1751), 20122327. doi: 10.1098/rspb.2012.2327

Wu, L., & Kruckeberg, A. L. (1985). Copper tolerance in two legume species from a copper mine habitat. *New Phytologist*, *99*(4), 565–570. doi: 10.1111/j.1469-8137.1985.tb03684.x

AUTHOR CONTRIBUTIONS:

Ideas: SS, MOB, ROE, VF, MTK, APH, KMG, CC

Data contributions: SS, MOB, GH, REO, WDR, KBO, KMG, YR, LG, FP, ZTW, APH

Analyses: SS, MOB, CC, REO, ZTW, KMG

Writing: SS, KMG, APH, with contributions from all authors

CONFLIT OF INTEREST

The authors declare no conflict of interest

DATA AVAILABILITY STATEMENT

The dataset used in this manuscript will be archived on DRYAD and is available at

https://proceeddatabase.weebly.com/.

677
Figure 1. Violin plot of rates of phenotypic change (log-transformed darwin numerator) comparing
natural (green) and human disturbed (black) systems. The two left violins are data from Hendry et al.,
(2008) and the two right violins are from our updated dataset. Axes were cut for better visualization (2
outliers removed from figure).

690
691

Figure 2. Rates of phenotypic change (mean darwin numerator) for hunted/harvested systems (purple),
natural systems (green), and other types of human disturbed systems (black). Rates of change are
averaged values per study system. Left panel is data from Darimont et al., (2009) and the right panel is
our current dataset.

698
699   **Figure 3.** Absolute phenotypic change in introduced species (pink) and non-introduced species (green)
700   measured in absolute darwin numerators. Each point is an expression at the species taxonomic level.
701   Darwin numerators are plotted as a function of years. The left panel is data from Westley (2011) and
702   the right panel is our updated dataset.
703

705
706
707 **Figure 4**. Boxplots depicting rates of change associated with body size measured in darwin numerators
708 for each disturbance category (climate change, hunting/harvesting, introduction, landscape change,
709 response to introductions, natural, and pollution). Light boxes are results from Gotanda et al., (2015)
710 and dark boxes are results from our updated dataset. Y axis was truncated to aid in visual assessment.

725 **Figure 5.** Rates of evolution—in log-transformed absolute darwin numerators—for six types of
726 disturbances (pollution, response to introductions, hunting/harvest, introductions, landscape change,
727 climate change) and natural populations. Points show individual data, lines show smoothed data
728 distributions, and crosses show GLM estimates ± standard errors. Letters indicate characterisations
729 based on Tukey HSD tests. Each point represents a system and reference-specific average.

Trait change (%)

Figure 6. Hypothetical effects of observed phenotypic change for all levels of ecology (populations, communities, ecosystems, and nature's contribution to people (NCPs)). A) 5-7% decrease in body size in harvested Chinook salmon (*Oncorhynchus tshawytscha*) in the Yukon River (Ohlberger et al., 2020) could decrease fish fecundity, affecting fishery yields. B) 8% increase in time spent foraging in Forester kangaroos (*Macropus giganteus)* introduced to Maria Island (Blumstein & Daniel, 2003) could affect the composition makeup of native vegetation and have effects on ecosystem processes like pollination. C) 5-7% decrease in body size in harvested Chinook salmon in the Yukon River (Ohlberger et al., 2020) could decrease nutrient transport affecting ecosystem properties. D) 43 % decrease in plant length in harvested Himalayan snow lotus (*Saussurea laniceps*) (Law & Salick, 2005) will decrease plant availability for medicinal use by humans.

Table 1. Partial η^2 values for darwin numerators and their appropriate models for each study revisited. Type refers to if the data were from a phenotypic study or a genotypic study (e.g., common garden).

Question	Data processing	Model	N original	N updated	Explanatory variable	Effect size original	Effect size updated
Hendry et al., (2008)	Raw data	Absolute Darwin numerators ~ years * human influence	2844	6957	Years	0.004	0.006
					Human Influence	0.115	0.018
Darimont et al., (2009)	Averaged by system, phenotypic data	Absolute Darwin numerators (means) ~ human influence + years	87	1529	Years	0.010	0.002
					Human Influence	0.180	0.010
Westley (2011)	Averaged by species	Absolute Darwin numerators ~ introductions * years	90	276	Years	0.613	0.135
					Introductions	0.781	0.906
New	Averaged by system and polynomial transformations	Log absolute Darwin numerators ~ disturbance + type + years +generations		2109	Years		0.959
					Disturbance		0.998
					Type		0.041

752
753
754

Table 2. Pairwise Wilcoxon signed-rank test results for size *versus* a different phenotypic trait (one-sided) to see if rates of evolution for body size were higher than other traits. Bold indicates significant P values where body size rates are higher than the other phenotypic trait, though not necessarily *positive*. Trait classification followed the definitions found in Kingsolver and Diamond (2011).

		2015				2021			
		Other morphology	Phenology	Other life history traits	Physiology	Other morphology	Phenology	Other life history traits	Physiology
Darwins	W	69532	6326	1313	3311	999128	89790	192019	50292
	P value	1	1	**0.025**	0.997	1	0.938	0.180	0.959
Darwin numerators	W	69428	6339	1712	3525	1008067	89545	195065	55501
	P value	1	1	0.429	1	1	0.932	0.290	0.999
Haldanes	W	26355	2560	835	1316	779302	73446	50376	38219
	P value	0.821	0.084	0.607	0.922	1	0.982	0.661	0.924
Haldane numerators	W	26151	3525	773	1289	765025	70326	49636	40013
	P value	0.790	0.983	0.390	0.889	1	0.912	0.585	0.979

767 **Table 3.** Sign-test results for rates of evolution testing whether body size rates were significantly different from zero. Results are shown for
768 all data and for data with harvesting data removed. Median rates are given, and bold values mean the median is significantly different from
769 zero. Due to the nature of the sign test, numerators yield the exact same results, and so are not reported. Body size classification followed the
770 trait classification definitions found in Kingsolver and Diamond (2011).
771
772

		2015	2021	2015 (no harvesting)	2021 (no harvesting)
Darwins	N (body size)	146	1902	87	705
	Median	-1763	-1306	-711	-372.842
	p-value	**<0.001**	**<0.001**	**0.018**	**<0.001**
Haldanes	N (body size)	70	1576	67	682
	Median	-0.00028	-0.0396	-0.00197	-0.0167
	p-value	0.403	**<0.001**	0.625	**<0.001**

779 **Table 4**. Estimates, standard errors, Tukey test categorizations, and type II likelihood ratio test results
780 for models predicting $\log_{10}$ absolute value darwins.
781
782
783 **$\log_{10}$ darwins**

	Variable	Est.	SE	Tukey	χ^2	df	p
Disturbance	Climate change	-2.20	0.08	a			
	Hunting / harvesting	-1.54	0.07	c			
	Introduction	-1.32	0.06	d			
	Landscape change	-1.52	0.13	bcd	361.03	6	**< 0.001**
	Local response to introduced	-1.13	0.11	d			
	Other	-1.69	0.07	b			
	Pollution	-0.64	0.14	e			
Phenotypic (vs. genetic)		0.01	0.06		0.06	1	0.80
$\log_{10}$ years		0.15	0.02		34.76	1	**< 0.001**
$\log_{10}$ generations		0.05	0.04		1.48	1	0.22

784
785
786