1 **Ecological specialisation, rather than the island effect, explains morphological**
2 **diversification in an ancient radiation of geckos**
3

4 Héctor Tejero-Cicuéndez[1*#], Marc Simó-Riudalbas[1*], Iris Menéndez[2,3], Salvador Carranza[1]
5
6 [1] Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la
7 Barceloneta 37-49, 08003 Barcelona, Spain.
8
9 [2] Departamento de Geodinámica, Estratigrafía y Paleontología, Facultad de Ciencias
10 Geológicas, Universidad Complutense de Madrid, C/ José Antonio Novais 12, Madrid,
11 28040, Spain.
12
13 [3] Departamento de Cambio Medioambiental, Instituto de Geociencias (UCM, CSIC),
14 C/Severo Ochoa 7, Madrid, 28040, Spain.
15
16 [*] These authors contributed equally.
17
18 [#] Correspondence to be sent to: Institute of Evolutionary Biology (CSIC-Universitat Pompeu
19 Fabra), Passeig Marítim de la Barceloneta 37-49, 08003 Barcelona, Spain.
20 E-mail: hector.tejero@ibe.upf-csic.es

21
22
ABSTRACT

Island colonists are often assumed to experience higher levels of phenotypic diversification
than continental taxa. However, empirical evidence has uncovered exceptions to this "island
effect". Here, we tested this pattern using the geckos of the genus *Pristurus* from continental
Arabia and Africa and the Socotra Archipelago. Using a recently published phylogeny and an
extensive morphological dataset, we explore the differences in phenotypic evolution between
Socotran and continental taxa. Moreover, we reconstructed ancestral habitat occupancy to
examine if ecological specialisation is correlated with morphological change, comparing
phenotypic disparity and trait evolution between habitats. We found a heterogeneous outcome
of island colonisation. Namely, only one of the three colonisation events resulted in a body
size increase. However, in general, Socotran species do not present higher levels or rates of
morphological diversification than continental groups. Instead, habitat specialisation explains

better the body size and shape evolution in *Pristurus*. Particularly, the colonisation of ground
habitats appears as the main driver of morphological change, producing the highest disparity
and evolutionary rates. Additionally, arboreal species show very similar body size and head
proportions. These results reveal a determinant role of ecological mechanisms in
morphological evolution and corroborate the complexity of ecomorphological dynamics in
continent-island systems.
41
42 **Keywords**
43 Body size; disparity; evolutionary rate; island colonisation; morphospace; *Pristurus* geckos.
44
45
INTRODUCTION
The life history and population biology of continental and insular taxa of a specific
evolutionary radiation are fundamentally distinct [1,2]. In the continent, communities are
often assumed to be complex and composed of many species that share a long history of
coevolution [3]. In such a scenario, most of the ecological niches will be filled, and high
levels of interspecific competition are expected [4]. These factors, together with higher
predation pressures, will tend to limit niche expansion and, consequently, morphological
diversification [5]. In contrast, insular groups are usually exposed to higher levels of
ecological opportunity and thus, they can occupy the new or relatively unexploited adaptive
landscapes that islands provide [6,7]. As a result, island species may experience increased
rates of phenotypic diversification and higher levels of morphological disparity compared to
continental taxa [8]. Moreover, the divergence in body size found in insular taxa relative to
their continental counterparts is a pervasive and widely studied pattern across vertebrates,
known as the "island rule" [9]. However, empirical evidence outlines a more complex
scenario in which island colonists might not necessarily experience great levels of
evolutionary divergence [10]. Instead, these potential outcomes of island colonisation — i.e.,
body size and shape divergence, increased phenotypic disparity, and increased rates of
morphological evolution; from now on referred to as "island effect" — might depend on
multiple *extrinsic* factors (mostly modulated by the geography or geology of the island), as
well as *intrinsic* factors (i.e., the biological characteristics of the group concerned) [3,11].
Thus, ecological specialisation is expected to be more pronounced when island colonisation
results in an expansion into novel ecological contexts, and such specialisation might carry
morphological changes depending on species and system-specific factors [4,6].

Whether ecological specialisation follows island colonisation or not, the study of
habitat occupancy is essential to understand the evolution of associated traits and the
structuring of ecological communities [12]. In particular, specialisation in substrate use can
promote morphological diversity through deterministic body size evolution and
diversification [13]. Moreover, microhabitat use can be strongly correlated with convergent
phenotypic evolution resulting in recurrent ecomorphs beyond the effect of history and clade
membership [14].

Arid regions, generally considered relatively depauperate in terms of animal diversity,
have been successfully inhabited by some vertebrate groups and harbour especially high
levels of reptile diversity. Among reptiles, geckos are particularly prominent due to their
outstanding diversity in ecological features, exhibiting a wide variety of morphological and
behavioural adaptations [15]. Afro-Arabian geckos have been recently prominent in studies
of the role of arid biomes in generating biodiversity [16–21]. However, the outcomes of
morphological diversification in these animals have only been properly investigated within
the genus *Hemidactylus,* which is the best-studied Arabian reptile group with well-resolved
taxonomy and reliably reconstructed biogeographic history [18,19,22–27]. In particular, two
recent studies including continental and insular taxa from the Socotra Archipelago indicated
that the genus *Hemidactylus* shows an island effect at least regarding body size evolution
[28,29]. In contrast, the geckos of the genus *Pristurus*, which have also colonised and
diversified within the same archipelago, seem to show lower rates of body size diversification
in Socotra than other genera in the archipelago, but also compared to their continental
relatives [28]. Despite these preliminary results, a more nuanced analysis of the
morphological evolution in *Pristurus*, including undescribed diversity, morphological and
ecological data, is still lacking. Interestingly, the genus *Pristurus* not only has representatives
in both the Socotra Archipelago and continental Africa and Arabia, but it also has diversified
ecologically to exploit a variety of habitats, including rocky and sandy surfaces, gravel plains,
and trees [30,31].

Here we use a recently inferred phylogenetic assessment of Afro-Arabian reptiles
including undescribed diversity, together with an extensive morphological sampling and
detailed ecological information, to explore the morphological evolution in *Pristurus* geckos.
Specifically, we test alternative scenarios of body size and shape evolution in this genus, to
determine the role of the colonisation of the Socotra Archipelago and the ecological
specialisation in generating the morphological diversity observed. The independent
diversification of both Socotran and continental taxa, the ecological and behavioural

diversity, and the unique phenotypic dataset compiled in this study, make this group of
geckos an exceptional system to investigate keystone dynamics in evolutionary biology such
as the island effect and ecological adaptation, and their impact on morphological evolution.
106
107

MATERIALS AND METHODS

Phylogenetic and ecological data
We used a recently published phylogenetic tree of Afro-Arabian squamates [32]. This tree
contains all the species of *Pristurus* for which there exists genetic data, including some
species currently in the process of being described, resulting in a total of 30 species. We
extracted the *Pristurus* clade from the squamate tree, both for the consensus and for 1,000
trees randomly selected from the posterior distribution generated in the cited study. Using a
sample of posterior trees allowed us to take into account the phylogenetic and temporal
uncertainty in the subsequent analyses.
Each species was defined as present in the Socotra Archipelago or in the continent
(Africa and Arabia). For habitat specialisation, each species was categorized based on
substrate preferences into one of three groups: ground-dwelling, rock climber, or arboreal
[30,33]. Additionally, the ground-dwelling species were divided into "soft-ground" (sandy
surfaces) and "hard-ground" (gravel plains) categories to further characterize the
morphological adaptations to each type of ground habitat. Nevertheless, the disparity
dynamics and rates of trait evolution were estimated with the original categorizations
(continent - Socotra; and the three habitat states) which, due to the limited number of species,
were more appropriate for the analyses.

Ancestral reconstructions
We studied the colonisation of the Socotra Archipelago and the habitat specialisation through
time by reconstructing ancestral states across the phylogeny. First, we fitted several models
of character evolution across the phylogeny in order to select the best-fit model for both
traits. With such a purpose, we used the function *fitDiscrete* from the R package geiger v2.0
[34,35]. We fitted three models: an equal-rates model (ER), a symmetrical model (SYM), and
an all-rates-different model (ARD). We selected the best-fit model in each case based on the
Akaike information criterion, correcting for small sample size (AICc; Akaike 1973). We then
used the function *make.simmap* from the R package phytools [37], which simulates plausible

stochastic character histories after fitting a continuous-time reversible Markov model for the
evolution of the character states assigned to the tips of the tree. We ran 1,000 simulations
with the previously selected model of character evolution (ER model for both traits).
Additionally, we randomly selected 100 trees from the posterior distribution, and we ran 100
stochastic character histories on each of them for both traits (presence in Socotra and habitat
occupancy).

Phenotypic data
For the 30 species included in the phylogenetic tree, a total of 697 specimens were examined
and measured, with a minimum of one, a maximum of 56, and a mean of 23 specimens per
species (Table S1). All vouchers were obtained from the following collections: Institute of
Evolutionary Biology (CSIC-UPF), Barcelona, Spain (IBE), Natural History Museum,
London, UK (BM), Museo Civico di Storia Naturale, Carmagnola, Turin, Italy (MCCI),
Università di Firenze, Museo Zoologico "La Specola", Firenze, Italy (MZUF), Oman Natural
History Museum (ONHM), Laboratoire de Biogéographie et Écologie des Vertébrés de
l'École Pratique des Hautes Etudes, Montpellier, France (BEV), and National Museum
Prague, Czech Republic (NMP). The following measurements were taken by the same person
(MSR) using a digital caliper with accuracy to the nearest 0.1 mm: snout-vent length (SVL;
distance from the tip of the snout to the cloaca), trunk length (TrL; distance between forelimb
and hindlimb measured posterior to the forelimb and anterior to the hindlimb insertion
points), head length (HL; taken axially from tip of the snout to the anterior ear border), head
width (HW; taken at anterior ear border), head height (HH; taken laterally at anterior ear
border), humerus length (Lhu; from elbow to axilla), ulna length (Lun; from wrist to elbow),
femur length (Lfe; from knee to groin) and tibia length (Ltb; from ankle to knee). Tail length
was not measured because most of the specimens had regenerated tails or had lost it.

Morphological differentiation
As body size and shape evolution might be affected by the colonisation of Socotra and/or the
ecological specialisation, we investigated whether Socotran colonists are morphologically
distinct from their relatives in the continent (one potential outcome under the "island effect"
framework) and likewise whether the differential habitat use is reflected in species
morphology. For each species, the mean of each morphological variable was calculated and
$\log_{10}$-transformed in order to improve normality and homoscedasticity prior to subsequent
analyses.

To characterize shape differentiation, we performed a phylogenetic regression of each

trait on snout-vent length (SVL) to remove the effect of the body size on the other variables.

The residuals of these regressions were used to implement a phylogenetically controlled

principal component analysis (pPCA) using the functions *phyl.resid* and *phyl.pca* from the R

package phytools with the method set to 'lambda' [37]. We used the principal components

representing 75% of the cumulative proportion of variance as shape variables and we

visualised the 2D shape morphospace. Additionally, we performed a principal component

analysis (PCA) with the shape data from all the specimens measured, after correcting for

body size through regressions on SVL similarly to our processing of the species data. We

generated per-species boxplots of size and shape variation with the specimen data.

For body size, we used the function *phenogram* from phytools [37] to map and visualize

variation in SVL across the species tree. For all the visualisations, we categorised the species

according to their presence in the Socotra Archipelago (Socotra or continent) and to their

habitat use (ground, rock or tree) separately, to have a detailed perspective of the extent of the

morphological differentiation undergone in each category. Furthermore, to test the island rule

prediction that body size in insular species is different from the continent, we performed a

phylogenetic ANOVA with the *phylANOVA* function from phytools [37]. Additionally,

according to the island rule, we would expect to see body size differences before and after a

colonisation event. To test this, we performed paired t-tests on the estimated body size for

each pair of parental and descendent nodes in which a change from continent to Socotra

occurred. We used the function *OUwie.anc* from the R package OUwie [38] to estimate the

ancestral body size under a multi-rate Brownian Motion model, which was the best-fit model

for body size evolution (see below "Differences in tempos of phenotypic diversification").

Exploring differences in phenotypic disparities

Since one of the main possible outcomes of island colonisation and/or ecological

specialisation is the increase in phenotypic disparity, we tested this assumption following

previous research on other geckos [29] and defining disparity as the average squared

Euclidean distance between all pairs of species in a group for a given continuous variable [7],

in our case body size (SVL) and two variables of shape (pPC1 and pPC2). First, we tested

whether disparity is higher in Socotran species than in the continent, which would indicate

the existence of an island effect. Then, considering our results of morphological

differentiation according to habitat use, in which ground-dwelling species are widespread

throughout most of the morphospace of the genus, we tested whether the phenotype of

ground-dwelling species is significantly more disparate than that of species in all the other
habitats together (i.e., rock and arboreal species). We calculated the observed disparity ratios
(Socotra/continent and ground/no-ground) for each morphological variable, using the
function *disparity* from the R package geiger [34]. In the case of higher disparity in Socotra
or in ground species (disparity ratio Socotra/continent or ground/no-ground higher than 1),
we then compared the observed ratios with a null distribution of disparity ratios obtained
from 10,000 simulations of the evolution of a continuous character according to a Brownian
motion model across the phylogeny. These simulations were performed by applying the
function *sim.char* after estimating the empirical rate parameter for body size and shape from
the best-fit model (Brownian motion) with the function *fitContinuous*, both from the package
geiger [34]. This approach allowed us to test, in the case of an observed higher disparity in
Socotran or ground species, whether this is a significant increase considering the rate of
evolution of the character, or rather this is not evidence of effectively increased disparity. In
order to test the sensitivity of the results to low intraspecific sampling, we performed the
analyses described above also with a dataset in which the species with less than five
specimens had been removed: *Pristurus adrarensis*, *P. flavipunctatus*, *P.* sp. 4, *P.* sp. 9, and
P. sp. 12 (see Table S1).
222

Differences in tempos of phenotypic diversification
In order to test the effect of Socotran colonisation and ecological specialisation in the tempo
of phenotypic evolution, we fitted different models of character evolution across the
phylogeny in which the evolutionary rates of body size and shape might or might not differ
between categories (i.e., between Socotra and continent, and between ground, rock and tree
habitats). For body size, we used the R package OUwie [38] to fit three alternative models:
BM1 (Brownian motion single rate, i.e., assuming one single rate regime for all lineages in
the phylogeny), OU1 (Ornstein-Uhlenbeck single-rate value with a phenotypic optimum and
a selective pressure towards it), and BMS (Brownian motion multi-rate model, with different
rate values for each of the regimes specified, i.e., Socotran different from continental
lineages, and differences between habitats). Similarly, we studied the rates of phenotypic
evolution for body shape, but in this case we fitted multivariate models including the first two
principal components resulting from the phylogenetic PCA (pPC1 and pPC2; 77% of the
variance, see Results section). We used the R package mvMorph [39] to fit four alternative
models: BM1, OU1, and BMM (analogous to BMS in the OUwie package), and OUM
(multi-rate Ornstein-Uhlenbeck model). We fitted these models in the 1,000 stochastic

character maps generated for the consensus tree, and also in the 100 character maps on each

of the 100 posterior trees used to reconstruct ancestral states (see above Ancestral

reconstructions). We then selected the best-fit models based on the AICc distributions and

means, and we extracted the distributions of rate values estimated for each regime (Socotra,

continent, ground, rock and tree) in the multi-rate models, to unravel the effect of each trait in

morphological rates. All the visualisations of the disparity and phenotypic rate analyses were

built with the R packages ggplot2 [40], patchwork [41], and cowplot [42].

RESULTS

Ancestral reconstructions and morphological variation

The ancestral reconstructions following an equal-rates model (ER), with the probabilities of

each state in ancestral nodes (Socotra and continent; ground, rock, and tree) can be found in

the Supplementary Material (Fig. S1). For the Socotra-continent reconstruction, the most

probable ancestral state for the root was "continent" (continent: 0.649; Socotra: 0.351). The

ecological reconstruction shows rocky habitats as the ancestral state in *Pristurus* (rock: 0.978;

ground: 0.012; tree: 0.010), with several transitions to arboreal habitats and one colonisation

of the ground in the ancestor of the clade known as "Spatalura group" (Arnold 1993, 2009;

Fig. S1). One of the subclades of this group later colonised more compact, harder substrates,

shown in our more detailed analysis separating soft- and hard-ground species (Fig. S1).

The pPCA analysis of body shape resulted in two first components explaining 77% of

the total variance: pPC1 (61% of the variance) mainly representing limb dimensions

(variables Lhu, Lun, Lfe, Ltb), and pPC2 (16% of the variance) mostly representing head

proportions (variables HL, HW, HH) (Fig. 1a; Table S2; see Fig. S2 and Table S3 for results

on body size and shape differentiation using the specimen data and PCA). The morphospace

occupied by continental species is notably larger than that occupied by Socotran species, and

they overlap almost completely (Fig. 1a left). When visualizing the shape phylomorphospace

along with habitat categories, we observe a wide portion occupied by the ground-dwelling

species, especially in pPC2 (head dimensions) (Fig. 1a right). These eight species of the

"Spatalura group" essentially occupy almost as much of the morphospace as all the rest of the

species together, with morphologies specialised to arboreal habits localized in a narrow area,

especially for head proportions (Fig. 1a right).

We found a similar pattern for body size. Body size variability of Socotran species is
completely contained in the range occupied by continental species (Fig. 1b left). On the
contrary, ground-dwelling species show a size variability higher than all the species from
other habitats together, being the largest and the smallest species of *Pristurus* specialised to
ground habitats (Fig. 1b right). As with head proportions, arboreal species have apparently
constrained body sizes, being restricted to specific intermediate values within the genus' size
range. The phylogenetic ANOVA showed no statistical differences in body size between
Socotran and continental species (F = 7.535, p = 0.215). Although one of the colonisation
events of the Socotra Archipelago was followed by an apparent increase in body size (the
clade composed by *Pristurus insignis*, *P. insignoides*, and *P.* sp. 12; see Fig. S3), our paired t-
test analysis did not find significant differences between parental and descendant nodes in
which a change from continent to Socotra occurred (t = -1.201, p = 0.353; Fig. S3).

When separating ground species into hard- and soft-ground habitats, we observed a
clear morphological differentiation, especially in body size. Hard-ground species seem to be
highly specialised, with some of the largest body sizes of the genus, long limbs and large
heads (Fig. S4).

Phenotypic disparity
We found that the morphological disparity in the Socotra Archipelago is lower than in the
continent for the three variables, with disparity ratios Socotra/continent below 1 (SVL: 0.88;
pPC1: 0.52; pPC2: 0.53). When comparing disparity between ground and no-ground habitats,
we found a higher observed disparity in ground for body size (SVL) and head proportions
(pPC2), with disparity ratios ground/no-ground of 2.25 for SVL, 0.86 for pPC1, and 2.47 for
pPC2. Furthermore, both for size and head proportions, the increased disparity in ground
habitats was significant compared with the null distribution of simulated disparity ratios (p_{size}
= 0.03; p_{head} = 0.01; Fig. 2). These results were virtually identical to those from the analyses
performed after removing the species with less than five specimens (Table S4).

Rates of morphological evolution
For body size, a multi-rate Brownian motion model (BMS) was the best fit both for presence
in Socotra and for ecological specialisation (lowest AICc; Fig. 3a), suggesting differences in
the rate of morphological evolution across discrete categories (i.e., Socotra vs. continent, and
different habitats). For body shape, however, we did not find evidence for differences in
evolutionary rates, being the single-rate Brownian motion model (BM1) the best fit both for

limb (pPC1) and head (pPC2) dimensions, although the overlap across all models was
considerable (Fig. 3b).

We extracted the rates of body size evolution from the Brownian motion multi-rate
models, and we found that Socotran species present lower rates than species in the continent
(Fig. 4a top), although the bimodal distribution of the rates in Socotran taxa suggests rate
heterogeneity among those clades. For ecological specialisation, we found increased rates of
body size evolution in the ground-dwelling species relative to the other habitats, with arboreal
habitats showing the lowest rates (Fig. 4a bottom). We also extracted per-category rates of
body shape evolution according to the BMM model in mvMorph, even though the multi-rate
models were not the best supported, and we found a similar scenario, in which shape
evolution (both for limbs and for head proportions) would be notably faster in ground-
dwelling species (Fig. 4b bottom). Results from the analyses performed with the 1,000
stochastic character maps on the consensus tree and with the 100 maps on each of 100
posterior trees lead to the same conclusions, so we show the ones from the posterior trees on
the main text. Results from analyses with the consensus tree can be found in the
Supplementary Material (Fig. S5).

DISCUSSION
The present study represents the first comprehensive comparative work on the genus
Pristurus, including undescribed diversity and extensive morphological (size and shape) and
ecological data. We tested the relative roles of the colonisation of the Socotra Archipelago
and the ecological specialisation in the evolution of the phenotypic diversity observed within
the genus. We did not find evidence for the existence of an island effect in this radiation of
geckos, since Socotran species do not present a notably different morphology, higher
disparity, nor increased rates of morphological evolution, relative to species in the continent.
On the contrary, ecological specialisation emerges as a determinant factor in generating
morphological diversity, with the colonisation of ground habitats as the main driver of
phenotypic divergence. Our results reveal a complex scenario in which different
morphological traits interact with ecological characteristics of the species in different ways,
suggesting a differential relevance of body size and shape proportions for the adaptation to
specific habitats.

The tendency of island taxa to diverge in morphology compared to their continental
relatives is a general pattern in terrestrial vertebrates, especially concerning body size

[43,44]. In fact, recent studies on Afro-Arabian geckos colonising the Socotra Archipelago

found support for this island effect, particularly in the genus *Hemidactylus* [28,29].

Nevertheless, preliminary results on *Pristurus* geckos failed to find this phenomenon in this

genus [28]. Here we corroborated and extended those preliminary results, incorporating the

most complete phylogenetic and morphological sampling within *Pristurus* up to date. We did

not find the predicted effects of thecolonisation of the Socotra Archipelago in phenotypic

diversification. Even though one of the Socotran clades (the one including *P. insignis*, *P.*

insignoides and *P.* sp. 12) has effectively undergone an increase in body size, we did not find

significant differences in body size between the first nodes of each colonisation event and

their direct ancestral nodes (Fig. S3). There is also no apparent divergence in body shape,

with limb and head proportions of Socotran species being completely encompassed into the

morphospace of continental species (Fig. 1a). Moreover, neither size nor shape disparities

observed in the Socotra Archipelago are higher than those of the continental species. Finally,

our analyses failed to find another expected outcome of island colonisation, as is the increase

in rates of phenotypic evolution. Consistent with our results of body size divergence, in

which one of the Socotran clades is shown to have undergone a body size increase (Fig. S3),

we found heterogeneity in the rates of body size evolution, with some species evolving faster

than others producing a bimodal distribution (Fig. 4a). Nevertheless, species in the continent

show higher rates of body size evolution than all Socotran species (Fig. 4a). For body shape,

our evolutionary model fitting showed no support for differences in shape rates between

continental and Socotran species (BM1 was the best-fit model; Fig. 3a). When extracting the

rate values for shape from the multi-rate Brownian motion model, we found little or no

difference between continental and Socotran taxa (Fig. 4b). The lack of a general island effect

in *Pristurus* after the colonisation of Socotra, opposed to other similar diversifications of

geckos, may indicate the existence of different ecological contexts even in the same physical

settings, which would imply different ecological opportunities in the same island [11].

Namely, some life-history traits may have limited niche expansion in *Pristurus* species in the

Socotra Archipelago. For instance, the fact that the ground-dwelling clade of *Pristurus* is also

the only one that has adopted partially nocturnal habits (Arnold 2009) may indicate a

potentially related development of these two traits. This would explain that diurnal Socotran

Pristurus have not colonised ground habitats and therefore their morphology has not changed

substantially, while some nocturnal geckos that have colonised the Socotra Archipelago, like

Hemidactylus or *Haemodracon*, show a marked phenotypic divergence [29,45]. , This is

consistent with results on global insular vertebrate communities suggesting that the

prevalence of the island rule is subjected to system-specific ecological and environmental

dynamics [44]. Furthermore, a recent study on the anole radiation in the Greater Antilles did

not find evidence for an island effect, pointing instead to ecological opportunity and key

innovations as the drivers of the adaptive radiation [46]. In *Pristurus*, one of the clades

colonising the Socotra Archipelago (*P. insignis*, *P. insignoides* and *P.* sp. 12) seems to have

undergone to a certain extent an increase in body size coupled with increased evolutionary

rates compared with the other Socotran *Pristurus*. This could be indicative of some kind of

ecological release derived from the island colonisation but, in any case, the degree of

evolutionary change of this Socotran clade is not comparable to that observed with the

colonisation of ground habitats.

Following that reasoning, ecological specialisation gives us a much more nuanced

insight on *Pristurus* phenotypic evolution. The relationship between habitat use and

morphological traits is well recognized in lizards [3,47,48]. In fact, preceding observations on

Pristurus geckos suggested that many morphological changes might be functionally

associated with shifts in ecology and behavior [30]. Our results are consistent with this notion

and provide strong evidence that novel ecological opportunities produced high levels of

phenotypic disparity associated with increased rates of trait evolution in some forms of

Pristurus, particularly the species exploiting ground habitats. Even though ground-dwelling

species do not show an extremely divergent body shape relative to species inhabiting other

habitats (i.e., rocky and arboreal habitats; Fig. 1a), they do comprise the largest and the

smallest sizes of the genus (Fig. 1b) and show some extreme values of limb and head

proportions (pPC1 and pPC2 respectively; Fig. 1a), as well as they occupy a very large

portion of the genus' entire morphospace (Fig. 1a). This extreme variability within ground-

dwelling species is reflected in our disparity results. While ground species do not present

higher disparity in limb dimensions (pPC1; ratio ground/no-ground < 1), they have more than

twice as much disparity as all the rest of the species in body size and head proportions, with

these ratios being highly significant compared to the null model generated from simulations

of character evolution (Fig. 2). Furthermore, we found increased rates of body size evolution

in ground-dwelling species, followed first by rocky and last by arboreal habitats (Fig. 4a).

Although rate heterogeneity across habitat categories was not the best-fit model for body

shape (Fig. 3b), the rate values extracted from the multi-rate Brownian motion model show a

similar pattern, especially for head proportions, with highest rates in ground-dwelling species

(Fig. 4b). Taken together, these results point to the existence of a morphological response to

the ecological context, especially in body size. This is consistent with the idea that the main

driver of morphological divergence, even in an island colonisation event, is habitat diversity
[49,50]. If habitat heterogeneity in the Socotra Archipelago is lower than in continental
Africa and Arabia (e.g., no particularly large gravel plains in Socotra), or if the access to
those habitats is limited for *Pristurus* geckos in Socotra and not in the continent (e.g., due to
the interplay of other ecological and historical factors), phenotypic evolution after island
colonisation would not be as extreme as expected under the island effect framework. This
could imply a tight relationship between morphology and structural habitat, a pattern
observed in other Arabian geckos such as *Ptyodactylus* or *Asaccus*, where niche conservatism
is associated with a very conserved morphology [17,51–54]. This would be further supported
by the fact that within ground habitats, species show a clear morphological segregation
between 'hard' and 'soft' substrates, suggesting a particularly conspicuous specialisation to
the former (large bodies, long limbs and large heads in the hard-ground species: *P. carteri*, *P.*
collaris, *P. ornithocephalus*; Fig. S4). In fact, these hard-ground species that have developed
partially nocturnal habits have evolved into rough ecological analogues of some small diurnal
ground-dwelling desert iguanians (e.g., the rock agamas of the genus *Pseudotrapelus*, the
toad-headed agamas of the genus *Phrynocephalus*, and the yellow-spotted agama of the
genus *Trapelus*), with the long limbs favouring the movement in the newly colonised grounds
and with large heads able to accommodate large eyes advantageous in those open habitats.
Alternatively, the lack of an island effect in Socotran *Pristurus* might be explained by
climatic divergences replacing ecomorphological differentiation, or by a low morphological
evolvability [28].

Another interesting result is the morphological affinities of arboreal species,
especially in body size and head proportions, where they present intermediate values within a
very restricted range (Fig. 1). Consistently, we found notably reduced evolutionary rates in
body size and head proportions in these species relative to other habitats with the multi-rate
models (Fig. 4). This might corroborate the idea of adaptive processes leading to a tight
relationship between ecological traits and phenotype, since this scenario is expected if a
specific habitat constrains the morphology towards optimum values of body size and shape
[14].

Ultimately, our results provide evidence of the determinant role of habitat
specialisation in phenotypic evolution. This has important implications for understanding the
prevalence of the island effect in the context of differential ecological opportunity and,
combined with previous results on other similar systems, shows the complex nature of the
relationships between ecological mechanisms and morphology and their reliance on system-

specific dynamics. More detailed ecological and morphological data (e.g., dietary habits, geometric morphometrics of head shape) might help for a deeper understanding of the evolutionary dynamics of this and other groups of arid-adapted lizards.

ACKNOWLEDGEMENTS

We are very grateful to J. Roca, M. Metallinou, K. Tamar, J. Šmid, R. Vasconcelos, R. Sindaco, F. Amat, Ph. de Pous, L. Machado, J. Garcia-Porta, J. Els, T. Mazuch, T. Papenfuss, B. Burriel and all the people from the Environment Authority, Oman, for their help in different aspects of the work. This work was supported by grants CGL2015-70390-P (MINECO/FEDER, UE) and PGC2018-098290-B-I00 (MCIU/AEI/FEDER, UE), Spain and grant 2017-SGR-00991 from the Secretaria d'Universitats i Recerca del Departament d'Economia i Coneixement de la Generalitat de Catalunya to SC. H.T.-C. was funded by an FPI grant (BES-2016-078341) (MINECO/AEI/FSE), Spain. I.M. was funded by a predoctoral grant from the Complutense University of Madrid (CT27/16-CT28/16), Spain, and the project PGC2018-094955-A-I00 from the Spanish Ministry of Science and Innovation. M.S.-R was funded by an FPI grant (BES-2013-064248) (MINECO/AEI/FSE), Spain.

DATA AVAILABILITY STATEMENT

Supplementary material, data and R scripts used for this study are available at the online public repository https://doi.org/10.5061/dryad.xwdbrv1f6.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Baeckens S, Van Damme R. The island syndrome. Current Biology 2020;30:R338–9.

2. Foster JB. Evolution of mammals on islands. Nature 1964;202:234–5.

3. Losos J. Lizards in an evolutionary tree: Ecology and adaptive radiation of anoles. University of California Press; 2009.

4.   Losos JB, Ricklefs RE. Adaptation and diversification on islands. Nature 2009;457:830–6.

5.   Yoder JB, Clancey E, Des Roches S, Eastman JM, Gentry L, Godsoe W, et al. Ecological
opportunity and the origin of adaptive radiations. Journal of evolutionary biology
2010;23:1581–96.

6.   Schluter D. The ecology of adaptive radiation. Oxford University Press; 2000.

7.   Harmon LJ, Melville J, Larson A, Losos JB, Oakley T. The Role of Geography and
Ecological Opportunity in the Diversification of Day Geckos (*Phelsuma*). Systematic Biology
2008;57:562–73.

8.   Whittaker RJ, Triantis KA, Ladle RJ. A general dynamic theory of oceanic island
biogeography. Journal of Biogeography 2008;35:977–94.

9.   Lomolino MV. Body Size of Mammals on Islands: The Island Rule Reexamined. The
American Naturalist 1985;125:310–6.

10.   Rundell RJ, Price TD. Adaptive radiation, nonadaptive radiation, ecological speciation and
nonecological speciation. Trends in Ecology and Evolution 2009;24:394–9.

11.   Losos JB. Adaptive radiation, ecological opportunity, and evolutionary determinism.
American Naturalist 2010;175:623–39.

12.   Mahler DL, Revell LJ, Glor RE, Losos JB. Ecological opportunity and the rate of
morphological evolution in the diversification of Greater Antillean anoles. Evolution
2010;64:2731–45.

13.   Reynolds RG, Collar DC, Pasachnik SA, Niemiller ML, Puente-Rolón AR, Revell LJ.
Ecological specialization and morphological diversification in Greater Antillean boas.
Evolution 2016;70:1882–95.

14.   Moen DS, Morlon H, Wiens JJ. Testing Convergence Versus History: Convergence
Dominates Phenotypic Evolution for over 150 Million Years in Frogs. Systematic Biology
2016;65:146–60.

15.   Gamble T, Greenbaum E, Jackman TR, Bauer AM. Into the light: diurnality has evolved
multiple times in geckos. Biological Journal of the Linnean Society 2015;115:896–910.

16.   Metallinou M, Arnold EN, Crochet P-A, Geniez P, Brito JC, Lymberakis P, et al. Conquering
the Sahara and Arabian deserts: systematics and biogeography of *Stenodactylus* geckos
(Reptilia: Gekkonidae). BMC evolutionary biology 2012;12:258–258.

17.   Metallinou M, Červenka J, Crochet PA, Kratochvíl L, Wilms T, Geniez P, et al. Species on
the rocks: Systematics and biogeography of the rock-dwelling *Ptyodactylus* geckos
(Squamata: Phyllodactylidae) in North Africa and Arabia. Molecular Phylogenetics and
Evolution 2015;85:208–20.

18.   Šmíd J, Moravec J, Kratochvíl L, Nasher AK, Mazuch T, Gvoždík V, et al. Multilocus
phylogeny and taxonomic revision of the *Hemidactylus robustus* species group (Reptilia,
Gekkonidae) with descriptions of three new species from Yemen and Ethiopia. Systematics
and Biodiversity 2015;13:346–68.

19.   Šmíd J, Shobrak M, Wilms T, Joger U, Carranza S. Endemic diversification in the mountains:
514   genetic, morphological, and geographical differentiation of the *Hemidactylus* geckos in
515   southwestern Arabia. Organisms Diversity & Evolution 2017;17:267–85.

516   20.   Garcia-Porta J, Simó-Riudalbas M, Robinson M, Carranza S. Diversification in arid
517   mountains: biogeography and cryptic diversity of *Pristurus rupestris rupestris* in Arabia.
518   Journal of Biogeography 2017;44:1694–704.

519   21.   Machado L, Salvi D, James Harris D, Brito JC, Crochet PA, Geniez P, et al. Systematics,
520   biogeography and evolution of the Saharo-Arabian naked-toed geckos genus *Tropiocolotes*.
521   Molecular Phylogenetics and Evolution 2021;155:106969.

522   22.   Carranza S, Arnold EN. A review of the geckos of the genus *Hemidactylus* (Squamata:
523   Gekkonidae) from Oman based on morphology, mitochondrial and nuclear data, with
524   descriptions of eight new species. Zootaxa 2012;95:1–95.

525   23.   Gómez-Díaz E, Sindaco R, Pupin F, Fasola M, Carranza S. Origin and in situ diversification
526   in *Hemidactylus* geckos of the Socotra Archipelago. Molecular Ecology 2012;21:4074–92.

527   24.   Šmíd J, Carranza S, Kratochvíl L, Gvoždík V, Nasher AK, Moravec J. Out of Arabia: A
528   Complex Biogeographic History of Multiple Vicariance and Dispersal Events in the Gecko
529   Genus *Hemidactylus* (Reptilia: Gekkonidae). PLoS ONE 2013;8:e64018.

530   25.   Šmíd J, Moravec J, Kratochvíl L, Gvoždík V, Nasher AK, Busais SM, et al. Two newly
531   recognized species of *Hemidactylus* (Squamata, Gekkonidae) from the Arabian Peninsula and
532   Sinai, Egypt. ZooKeys 2013;79–107.

533   26.   Šmíd J, Uvizl M, Shobrak M, Salim AFA, AlGethami RHM, Algethami AR, et al. Swimming
534   through the sands of the Sahara and Arabian deserts: Phylogeny of sandfish skinks (Scincidae,
535   *Scincus*) reveals a recent and rapid diversification. Molecular Phylogenetics and Evolution
536   2020;107012.

537   27.   Vasconcelos R, Carranza S. Systematics and biogeography of *Hemidactylus homoeolepis*
538   Blanford, 1881 (Squamata: Gekkonidae), with the description of a new species from Arabia.
539   Zootaxa 2014;3835:501–27.

540   28.   Garcia-Porta J, Morales HE, Gómez-Díaz E, Sindaco R, Carranza S. Patterns of
541   diversification in islands: A comparative study across three gecko genera in the Socotra
542   Archipelago. Molecular Phylogenetics and Evolution 2016;98:288–99.

543   29.   Garcia-Porta J, Šmíd J, Sol D, Fasola M, Carranza S. Testing the island effect on phenotypic
544   diversification: insights from the *Hemidactylus* geckos of the Socotra Archipelago. Scientific
545   Reports 2016;6:23729.

546   30.   Arnold EN. Relationships, evolution and biogeography of Semaphore geckos, *Pristurus*
547   (Squamata, Sphaerodactylidae) based on morphology. Zootaxa 2009;21:1–21.

548   31.   Badiane A, Garcia-Porta J, Červenka J, Kratochvíl L, Sindaco R, Robinson MD, et al.
549   Phylogenetic relationships of Semaphore geckos (Squamata: Sphaerodactylidae: *Pristurus*)
550   with an assessment of the taxonomy of *Pristurus rupestris*. Zootaxa 2014;3835:33–58.

551   32.   Tejero-Cicuéndez H, Patton AH, Caetano DS, Šmíd J, Harmon LJ, Carranza S.
552   Reconstructing Squamate Biogeography in Afro-Arabia Reveals the Influence of a Complex

553   and Dynamic Geologic Past. Systematic Biology 2021;Advance Publication.

554   33.   Arnold EN. Historical changes in the ecology and behaviour of semaphore geckos (*Pristurus*,
555   Gekkonidae) and their relatives. Journal of Zoology 1993;229:353–84.

556   34.   Pennell MW, Eastman JM, Slater GJ, Brown JW, Uyeda JC, Fitzjohn RG, et al. Geiger v2.0:
557   An expanded suite of methods for fitting macroevolutionary models to phylogenetic trees.
558   Bioinformatics 2014;30:2216–8.

559   35.   R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
560   Austria: R Foundation for Statistical Computing; 2019. Available from: https://www.r-
561   project.org/

562   36.   Akaike H. Information theory and an extension of the maximum likelihood principle. In:
563   Petrov B, Csaki F, editors. Second International Symposium on Information Theory. Budapest
564   (Hungary): Akademia Kiado; 1973. page 267-281.

565   37.   Revell LJ. phytools: An R package for phylogenetic comparative biology (and other things).
566   Methods in Ecology and Evolution 2012;3:217–23.

567   38.   Beaulieu JM, O'Meara B. OUwie: analysis of evolutionary rates in an OU framework. R
568   package, version 2.6. 2021;

569   39.   Clavel J, Escarguel G, Merceron G. mvMORPH: An R package for fitting multivariate
570   evolutionary models to morphometric data. Methods in Ecology and Evolution 2015;6:1311–
571   9.

572   40.   Wickham H. ggplot2. Wiley Interdisciplinary Reviews: Computational Statistics 2011;3:180–
573   5.

574   41.   Pedersen TL. Patchwork: The composer of plots. R package version 1.1.1. 2020;

575   42.   Wilke CO. cowplot: streamlined plot theme and plot annotations for 'ggplot2'. R package
576   version 1.1. 2020;

577   43.   Lomolino MV. Body size evolution in insular vertebrates: Generality of the island rule.
578   Journal of Biogeography 2005;32:1683–99.

579   44.   Benítez-López A, Santini L, Gallego-Zamorano J, Milá B, Walkden P, Huijbregts MAJ, et al.
580   The island rule explains consistent patterns of body size evolution in terrestrial vertebrates.
581   Nature Ecology and Evolution 2021;5:768–86.

582   45.   Tamar K, Simó-Riudalbas M, Garcia-Porta J, Santos X, Llorente G, Vasconcelos R, et al. An
583   integrative study of island diversification: Insights from the endemic *Haemodracon* geckos of
584   the Socotra Archipelago. Molecular Phylogenetics and Evolution 2019;133:166–75.

585   46.   Burress ED, Muñoz MM. Ecological Opportunity from Innovation, not Islands, Drove the
586   Anole Lizard Adaptive Radiation. Systematic Biology 2021;Advance online publication.

587   47.   Goodman BA, Miles DB, Schwarzkopf L. Life on the rocks: Habitat use drives morphological
588   and performance evolution in lizards. Ecology 2008;89:3462–71.

48.   Higham TE, Russell AP. Divergence in locomotor performance, ecology, and morphology

between two sympatric sister species of desert-dwelling gecko. Biological Journal of the
Linnean Society 2010;101:860–9.

49. Lack D. Island biology illustrated by the land birds of Jamaica. Berkeley: University of
California Press; 1976.

50. Losos JB, Parent CE. The speciation-area relationship. In: Losos JB, Ricklefs RE, editors. The
Theory of Island Biogeography Revisited. Princeton University Press; 2009. page 415–38.

51. Carranza S, Simó-Riudalbas M, Jayasinghe S, Wilms T, Els J. Microendemicity in the
northern Hajar Mountains of Oman and the United Arab Emirates with the description of two
new species of geckos of the genus *Asaccus* (Squamata: Phyllodactylidae). PeerJ
2016;4:e2371.

52. Simó-Riudalbas M, Metallinou M, de Pous P, Els J, Jayasinghe S, Péntek-Zakar E, et al.
Cryptic diversity in *Ptyodactylus* (Reptilia: Gekkonidae) from the northern Hajar Mountains
of Oman and the United Arab Emirates uncovered by an integrative taxonomic approach.
PloS ONE 2017;12:e0180397.

53. Simó-Riudalbas M, Tarroso P, Papenfuss T, Al-Sariri T, Carranza S. Systematics,
biogeography and evolution of *Asaccus gallagheri* (Squamata, Phyllodactylidae) with the
description of a new endemic species from Oman. Systematics and Biodiversity 2018;16:323–
39.

54. Tamar K, Mitsi P, Carranza S. Cryptic diversity revealed in the leaf-toed gecko *Asaccus*
montanus (Squamata, Phyllodactylidae) from the Hajar Mountains of Arabia. Journal of
Zoological Systematics and Evolutionary Research 2019;57:369–82.

Figure 1. Morphological variability in *Pristurus*, with insight from insularity (left) and habitat use (right). a) Morphospace with phylogenetic relationship between the species,

642   showing body shape differentiation. b) Traitgram showing body size (SVL) through time on
643   the summary phylogenetic tree of *Pristurus*, mapped by the discrete categories of presence in
644   Socotra or the continent (left) and by ecological specialisation (right). Photos (proportional to
645   species' SVL): *Pristurus carteri* (top) and *P. masirahensis* (bottom), the largest and smallest
646   species of the genus, respectively.

653   **Figure 2**. Observed (red arrows) and simulated (grey bars) ratios of phenotypic disparity
654   between ground versus no-ground habitats. a) Body size disparity ratios. b) Head proportions
655   (pPC2) disparity ratios.

Figure 3. AICc distributions from the model fitting for a) body size and b) shape evolution of the genus *Pristurus*. These results correspond to model fitting on 100 stochastic character maps (insularity in the top panels and habitat use in the bottom) on 100 trees from the phylogenetic posterior distribution. BM1: Brownian motion single rate. OU1: Ornstein-Uhlenbeck single rate. BMS (OUwie) / BMM (mvMorph): Brownian motion multi-rate. OUM: Ornstein-Uhlenbeck multi-rate. For body size, the best supported model is a Brownian motion with rate heterogeneity across categories. For body shape, a single-rate Brownian motion model was the best-fit, although there is an extensive overlap across all models.

Rates of morphological evolution

Figure 4. Rates of a) body size and b) body shape evolution in the genus *Pristurus*, extracted from multi-rate Brownian motion models fitted on a total of 10,000 character maps (100 stochastic character maps on 100 posterior trees) of Socotra vs. continent and habitat use (ground, rock, and tree).

688