Changes in movement characteristics in response to private and social

information acquisition of socially foraging fish

Geoffrey P. F. Mazué[1*], Maxim W. D. Adams[1], Frank Seebacher[1], Ashley J. W. Ward[1]

[1] School of Life and Environmental Sciences, University of Sydney, Sydney, NSW, 2006,

Australia

* Correspondence should be addressed to G. P. F. Mazué (geoffrey.mazue@sydney.edu.au)

ORCID

Geoffrey Mazue: https://orcid.org/0000-0003-3254-1605

Frank Seebacher: https://orcid.org/0000-0002-2281-9311

Ashley Ward: https://orcid.org/0000-0003-0842-533X

ACKNOWLEDGEMENTS

This work was supported by a Barry Jonassen Award granted by the Australian Society for

Fish Biology to G.P.F.M. and an ARC Discovery Project Grant DP190100660 to A.J.W.W.

AUTHORS' CONTRIBUTIONS

G.P.F.M. and A.J.W.W. designed the experiment; G.P.F.M. performed the experiment;

G.P.F.M. and M.W.D.A. scored the video recordings and performed the tracking; G.P.F.M.

analysed the data with inputs from F.S. and A.J.W.W.; G.P.F.M. wrote the manuscript. All

authors edited the manuscript.

CONFLICT OF INTEREST

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

ABSTRACT

1. To overcome the cost of competition resulting from close social proximity while foraging in a group, individuals may balance their use of private (i.e. acquired from personal sampling) and social (i.e. acquired by watching other individuals) information in order to adjust their foraging strategy accordingly. Reliability of private information about environmental characteristics, such as the spatial distribution of prey, is thus likely to affect individual movement and social interactions during foraging. Our aim was to investigate how movement characteristics of foraging individuals changed as they acquired reliable private information about the spatial occurrence of prey in a foraging context.

2. We allowed guppies (*Poecilia reticulata*) to develop the reliability of their private knowledge about prey spatial occurrence by repeatedly testing shoals in a foraging task under three experimental distributions of prey: 1) aggregated prey forming three patches located in fixed locations, 2) scattered distribution of prey with random locations, or 3) no prey (used as control). We then applied tracking methods to obtain individual time series of spatial coordinates from which we computed a suite of movement variables reflecting search effort, social proximity and locomotion characteristics during foraging, in order to examine changes occurring over repeated trials and to investigate which best explained foraging success.

3. We show that foraging shoals became more efficient at finding and consuming food over the first three days by increasing their time spent active. Over time, individuals foraging on either scattered or aggregated prey travelled greater distances, showed an increasing distance

to their closest neighbour and became slightly more stochastic in their acceleration profile,

compared to control individuals.

4. We found that behaviour changed as private information increased over time. Social

proximity was the major predictor of foraging success in the absence of prior foraging

information, while stochasticity in acceleration and search effort became the most important

predictors of foraging success as information increased.

5. In conclusion, we show that individual movement patterns changed as they acquire private

information. Contrary to our predictions, the spatial distribution of prey did not affect any of

the movement variables of interest. Our results emphasise the importance of information, both

private and social, in shaping movement behaviour in animals.

Keywords: social foraging, movement, private information, prey spatial distribution, fish.

INTRODUCTION

Individuals benefit from foraging socially through diverse mechanisms such as local

enhancement (e.g., cues from successful foragers provide information about food location) or

social facilitation (e.g., initiating foraging quicker in the presence of foraging conspecifics),

which allow an increase in food detection and foraging success (Galef & Giraldeau, 2001;

Giraldeau & Beauchamp, 1999; Giraldeau & Caraco, 2018; Grand & Dill, 1999; Pitcher et

al., 1982; Snijders et al., 2021; Ward & Zahavi, 1973). Such benefits are thought to be due to

an increased sensorial scope, which allows for more information to be collected and

disseminated within the group (Clark & Mangel, 1986). In a social foraging scenario, two

types of information are available: 'private' information, which individuals acquire from their

own experience within the environment such as the recent observation of prey in the vicinity

or the memory of a location associated with recent successful foraging events (Valone, 1991);

and 'social' information which is acquired from watching other foragers and gauging their success and the level of competition at a patch (Coolen et al., 2003; Templeton, & Giraldeau, 1995; Valone, 1989). At any moment, an individual faces the choice of continuing to search for food on its own or joining other foraging conspecifics. Such a decision therefore results from balancing private and social information to reduce uncertainty about the environment and so maximise foraging success (Templeton & Giraldeau, 1996). Social information is typically ubiquitous in social foraging and it is assumed to be less costly to acquire and use in terms of time and energy than private information. Yet, relying excessively on social information can lead individuals to make maladaptive decisions (Giraldeau et al., 2002; Laland & Williams, 1998), and foraging within close social proximity can generate non-negligible costs of competition (Cresswell, 1998). Additionally, individual nine-spined sticklebacks (*Pungitius pungitius*) tended to disregard social information when making foraging decisions if their private information was reliable and up-to-date (Van Bergen et al., 2004). Similarly, guppies (*Poecilia reticulata*) relied on private knowledge and preferentially foraged on patches they had visited before, rather than joining an alternative patch occupied by conspecifics (Kendal et al., 2004). Therefore, individuals might benefit from relying preferentially on private information when its acquisition is recent and when its cost is low (i.e. time and energy allocated to private sampling, risk of predation).

In most complex natural environments, food availability may vary both over time and in its spatial distribution (Riotte-Lambert & Matthiopoulos, 2020), which necessitates continuous gathering and processing of information in order to adjust foraging behaviour (Higginson et al., 2012). Without any prior knowledge about the spatial or temporal occurrence of resources in the environment, foraging individuals must rely primarily on social information (Harel et al., 2017; Lewis et al., 2013; Reebs, 2000). As they sample the environment, individuals increase their private information about food distribution.

Experience improves a forager's ability to predict the outcome of future sampling at a given location and/or time. Private information that promotes successful foraging may then be considered as reliable information. Such reliability is thus likely to affect the decisions that individuals make, such as how they move in the environment and how they interact with other foragers (Riotte-Lambert & Matthiopoulos, 2020), and it is likely to be affected by the distribution patterns of the resources. For example, foraging on unpredictable patchy resources may promote grouping behaviour and the use of social information to maximize chances of discovery and minimize the latency to arrive at a newly-discovered food patch (Beauchamp & Ruxton, 2014; Ioannou et al., 2011). However, in the presence of private information about patch location, foraging asocially may reduce costs of competition within a foraging patch. Alternatively, when food discoveries lead to relatively small food intakes, as is the case when resources are distributed more evenly across the environment, individuals should ignore social information to rely mostly on private information. Searching asocially in a Levy flight-type pattern is known to be efficient when resources are sparse and unpredictable (Humphries et al., 2012; Viswanathan et al., 2000).

In this study, we investigated the effect of reinforced private information on movement characteristics of individuals foraging on different spatial distributions of prey. We analysed high resolution trajectory data collected from shoals of guppies (*Poecilia reticulata*) that were allowed to forage repeatedly on prey that were either aggregated in patches at constant locations or scattered at random locations in the testing arena. First, we examined the changes occurring in a suite of variables describing different aspects of individual foraging behaviour including search effort, social proximity and stochasticity of locomotion, and compared it to those of control shoals tested in absence of food. Then we estimated relative importance of these variables at explaining individual foraging success. We hypothesised that (*i*) in the absence of prior information about the prey distribution and

competition, individuals initially forage as a group to enhance their exposure to social

information, regardless of the prey distribution; and (*ii*) changes of behaviour occur as private

information about spatial occurrence of prey and perceived level of competition increases,

before reaching a phase of fixation when individuals' knowledge of the foraging conditions

has become highly reliable (i.e., reinforced). A corollary from (*ii*) is that (*iii*) the spatial

distribution of prey should affect behavioural adjustments.

MATERIALS AND METHODS

Experimental subjects

We used 140 female domestic guppies (*Poecilia reticulata*) as experimental subjects

(obtained from Pisces Aquatics, Kenmore, QLD, Australia). Individuals had a body length of

33.7 ± 2.8 mm (standard length, mean $\pm$ sd). Only females were used in this experiment as

previous research showed that sex affects movement, food discovery and information

spreading in foraging shoals of guppies; with females having greater success at discovering

foraging patches than males (both under laboratory and natural conditions) and enhanced

spreading of foraging information (i.e. route to access the foraging patch) in groups of

females than groups of males (Croft et al., 2003; Reader & Laland, 2000; Snijders et al.,

2019). Individuals were split between two large 180L white plastic tanks placed in a

temperature and photoperiod (12h Light: 12h Dark cycle) controlled room for 22 days before

commencement of the experiment. Holding tanks were filled with dechlorinated aged water

maintained at a constant temperature of 23°C and furnished with air powered sponge filters

and plastic plants. Fish were fed tropical fish flakes (SERA Vipan, Heinsberg, Germany)

twice daily during the first 12 days, and then received flakes in the morning and defrosted

bloodworms (*Chironomidae*, Ocean Nutrition) in the afternoon for the remaining 10 days in

order to ensure the fish were familiar with the prey later used in the experiment (Godin,

1978). The day before the commencement of the experiment, we gently netted fish from their

holding tank and distributed them across 12 experimental groups of 4 fish. Each fish within a

group had a unique coloration patterns on their abdomen and their caudal fin, enabling

individual identification in video recordings. Each experimental group was photographed

from above in a small container containing a ruler placed in shallow water to enable the

measurement of each individual's standard length from the photograph using ImageJ. Each

experimental group was placed in an opaque green plastic tank (33 cm L x 24 cm W x 18 cm

H, 14L) containing an air powered sponge filter, and two plastic plants, and filled with

dechlorinated aged water maintained at a constant temperature of 23°C and kept under the

same light cycle as before. Two blocks of 12 groups and a third block of 10 groups were

tested within eight consecutive weeks (October-December 2019), and treatments were

balanced within each block.

Experimental design

We used two identical experimental arenas made of opaque white Perspex (inner

dimensions 118 cm L x 30 cm W x 20 cm H). The bottom of the arenas was covered with a 8

mm deep white plastic grid with each square of the grid measuring 15 mm x 15 mm (referred

to as 'foraging pits'). Each of the 1200 foraging pits contained either a single prey item or

none, and ensured that prey were held in assigned locations and did not drift in currents

generated by nearby swimming fish. We used a standardized quantity (n=12) of defrosted

bloodworms (*Chironomidae*, Ocean Nutrition, Essen, Belgium) as prey items, that were

distributed across the grid according to one of the three experimental treatments: 1) the

'aggregated' treatment (n=13 groups) consisting of 12 bloodworms divided into three

aggregations of 4 adjacent foraging pits, at fixed locations across the grid; 2) the 'scattered'

treatment (n=12 groups) consisting of 12 bloodworms scattered across the arena with a

pattern of distribution changing daily, obtained by haphazardly scattering bloodworms while

avoiding any aggregation between two worms; and 3) the 'control' treatment (n=10 groups)

consisting of an empty foraging grid containing no prey items (Figure 1). The arena was

filled with 7 cm of dechlorinated aged water, leaving 6.2 cm of water column for the fish to

navigate above the grid. Such a low water depth reduced the ability of the fish to see multiple

foraging pits while swimming, forcing them to discover which foraging pits contained a prey

item by exploring the entire arena. A filtered solution from 10 crushed bloodworms was

added to the arena each day before the trials commenced to standardize the amount of

foraging cues in the arena for the first groups to be tested on the day. Three sandstone

cylinders (diameter 5.8 cm and height 8 cm) were placed at fixed locations (across replicates

and treatments) of the grid to create environmental enrichment and break-up the line of sight

of foraging individuals. A square container made of transparent plastic (15 cm side lengths)

was used as a starting box (Figure 1). The arenas were illuminated overhead with fluorescent

tubes and were surrounded by white plastic sheets to minimize external stimuli.

Experimental groups were randomly assigned to one of the three treatments on the

day prior to the commencement of the experiment and were tested daily for 10 consecutive

days. Testing order and arenas were randomized daily to prevent any effect on foraging

behaviour. Tested groups were gently netted out of their holding tank to be placed inside the

starting box. After 2 minutes of acclimation, one edge of the starting box was raised remotely

using a pulley system, freeing the fish to explore the arena. All trials were filmed for 25 min

at 30 frames per second and at a resolution of 1280x720 pixels, using a Canon G1X camera

suspended above the arena. After completion of a trial, each group was returned to its holding

tank. To standardize the food intake, all groups assigned to the control treatment received 12

bloodworms in their holding tank. All groups were offered tropical fish flakes (SERA Vipan)

at the end of the day.

195 **Figure 1.** Top view of the experimental arena with the spatial distributions of prey used as

196 experimental treatments: a) control, b) scattered, and c) aggregated. The dashed-line square

197 symbolises the starting box used to release the fish, set on a platform symbolized by the grey

198 area. Sandstone cylinders are symbolized by black disks. Prey items (i.e. bloodworms) are

199 symbolized by small black crosses.

Data extraction and processing

From the video recordings, we recorded the latency (in frames, measured from the

frame when the starting box was raised) for each bloodworm to be eaten and the identity of

the fish that consumed it. We then quantified the time (in seconds) for the group to consume

the 12 bloodworms (later referred as 'depletion time'). Groups that failed to find and

consume all bloodworm within that a maximal time of 25 min were given a score of 1500sec

(3 groups on day 1). Early inspection of the data revealed that the depletion time decreased

significantly over the first 4 days to reach a plateau at day 5 (Figure 2A). Subsequently,

further data extraction and analyses were performed solely on the days 1, 3, 5 and 7 of the

experiment.

211   To track individual movement, video recordings from the aggregated and scattered

treatments were cropped from the 'starting frame', defined as the frame at which the first

bloodworm was picked by a fish (as a standardization method), onwards to reach a total of

1800 frames (i.e., 60 sec). This period was chosen because most of bloodworms were

consumed during the first minute. Videos from the control treatment, were instead cropped to

1800 frames from a starting frame equal to the averaged value of the starting frames for the

aggregated and scattered videos of the same day. Cropped video files were then converted to

.avi using VirtualDub (version 1.10.4) and tracked using the CTRAX automated tracking

software (Caltech Ethonomics Project, The Caltech Multiple Fly Tracker, Version 0.5.18)

(Branson et al., 2009). The time series of (x, y) coordinates produced for each individual

were then screened for any abnormalities and manually corrected using the FixErrors GUI in

MATLAB. Pixel coordinates were converted to mm using a conversion ratio calculated by

measuring the distance between two points of known distance on a video frame. Within each

experimental group for each daily video recording, the four time-series of coordinates (of

length 1800 frames each), were matched to the identity of the fish based on the coloration

patterns of their abdomen and caudal fins.

227   From the tracking data we performed different calculations to characterize three

228   distinct aspects of individual movement in a foraging context and to describe grouping

patterns observed at the level of the group (see SM). First, we calculated each individuals'

speed (standardized in body length per second) at each time step. Within this time series for

each individual, we identified all bouts (defined as sequences of consecutive frames) for

which it was moving by using a criterion of speed greater than 0.5 $BL.s^{-1}$ (Jolles et al., 2020),

and further quantified the proportion of time spent moving. Additionally, we calculated the

distance between each pair of fish (in averaged body length unit) at each time step, from

which we extracted the distance of each fish to its nearest neighbour (further referred to as

distance to NN, or NND) at each frame. Subsequently, we quantified the cumulative distance

(in cm) travelled by each individual over all frames for which it was identified to be moving,

as a measure of search effort. The average distance to NN of each individual, calculated as

the mean distance to NN over all frames for which it was identified to be moving, was used

as a measure of social proximity while foraging. Additionally, we computed the entropy

values for the time series of acceleration for each bout of at least 15 consecutive frames (i.e.,

half a second) during which a fish was identified as moving (a threshold determined to be

highly conservative, as it contained over 80% of the frames spent moving within each day),

using the 'pracma' package in R (Borchers, 2019). Approximate entropy estimates the

amount of regularity and unpredictability of fluctuations in a time series, enabling us to

determine whether individuals' locomotion was more deterministic (low values of

approximate entropy) or more stochastic (i.e., less predictable, high values of approximate

entropy) while searching for food in the arena (Pincus, 1991). The median value of all bouts'

entropy values was used as an overall measure of unpredictability in acceleration of a

foraging fish.

Each fish was given a +2 score for each bloodworm that was consumed in its entirety

during the 60s of the tracked video segments. Because of a non-negligible number of

kleptoparasitism events, a score of +1 was given to both individuals that were involved in the

consumption of a single bloodworm (e.g. stealing or dropping/picking up a half-consumed

prey).

Statistical analysis

Sample sizes presented until day 5 included 13 shoals (52 fish) in the aggregated

treatment, 12 shoals (48 fish) in the scattered treatment and 10 shoals (40 fish) in the control

treatment; but only 12 and 11 shoals (48 and 44 fish) in aggregated and scattered treatments

(respectively) on day 7 due to the exclusion of two groups from the experiment caused by the

loss of two individuals. Our general approach was to analyse the dependent variable using

models fitted according to their initial distribution, and no transformation was applied on any

of the dependent variables.

Analysis of depletion time and proportion of time moving

We first analysed the depletion time using a Generalized Linear Mixed Effects model

(GLMM) fitted with a negative binomial error distribution. The model included day and

treatment (with the exclusion of the control level as there were no prey in the arena) as fixed

effects, and shoal identity as random factor. We then explored the influence of treatment and

day on the individual proportion of time spent in motion using a GLMM fitted with a

binomial error distribution and the logit link function. Fish identity was nested within the

shoal identity as random effect. Attempts to simplify the models were made by comparing the

models with and without interaction terms based on their AIC values. The significance of the

terms kept in the reduced models were assessed using type III ANOVAs. Model assumptions

were verified through visual inspection of the Q-Q plot of the residuals and by plotting the

predicted residual against the observed residuals.

Analysis of movement characteristics

We analysed the effect of repeated testing (i.e. day) and prey distribution (i.e.,

treatment), and their interaction, on each of the three dependent variables, using GLMMs

fitted with a Gaussian or Gamma error distribution depending on the estimated distribution of

the dependent variables analysed. The model included treatment and day and their interaction

as fixed effects, and fish identity nested within the shoal identity as random effects. Attempts

to simply the models, assessment of terms' significance, and assumption verifications were

performed as described previously.

Analysis of the foraging success

We assessed the relative effect of four variables of interest, *viz*. treatment, total

distance travelled, average distance to NN, acceleration entropy, on individual foraging

success by calculating estimates of model-averaged coefficients (with their 95%CIs) from a

set of candidate models generated from reducing an initial model containing all four variables

of interests as main effects (Burnham & Anderson, 2002; Grueber et al., 2011). Foraging

success was thus analysed as a response variable in a generalized mixed effect model

(GLMM) fitted with Poisson error distribution. This initial complete model included all four

non-colinear (see Table S1) variables of interest as fixed terms, and shoal identity as random

term. A set of 16 candidate models was generated from using these four independent

variables, including the complete model and a null model. Predictors were standardized

(mean =0, SD = 0.5) prior to the analysis in order to simplify interpretation of model

averaged parameter estimates (Gelman et al., 2009), as recommended by Grueber et al.

(2011). Akaike information criterion corrected for low values of the ratio between sample

size and parameters (i.e., AICc) were used to evaluate each model within the set of candidate

models (Burnham & Anderson, 2002). Candidate models were then ranked from best to worst

based on their deltaAICc values. Akaike weights and evidence ratios were used to assess the

relative strengths of each candidate model.

Because we were interested in determining which factors have the strongest effect on

foraging success during a trial, we performed a model averaging procedure on the 95%

candidate set of models (i.e., top models with a cumulative weight of 95%) using a full

averaging method (i.e., 'zero method') implemented in the MuMIn package (Burnham &

Anderson, 2002; Nakagawa & Freckleton, 2010). The relative importance (RI) of the

parameters was calculated by summing the Akaike weights from the models in which they

appear, using the 95% candidate set of models. RI values close to 1 indicate that the

parameter appears in the best supported models, while RI values close to 0 indicate that the

parameters appear in the least supported models.

We performed the same AICc model averaging procedure for each day separately in

order to investigate whether there were changes in the relative importance of the predictors

on foraging success as individuals acquire more reliable private information (i.e., over

repeated days of testing) and because 'day' had a strong effect on most of the variables of

interest. Doing so also maintained parsimony in the model construction.

All data were analysed using the 'lme4' (Bates et al., 2015), 'MASS' (Venables &

Ripley, 2002), 'car' (Fox & Weisberg, 2019) and 'MuMIn' (Barton, 2020) packages for

modelling and visualized using the 'ggplot2' (Wickham, 2016) package in R 3.6.2. (R Core

Team, 2019).

RESULTS

Depletion time and proportion of time spent moving

There was a significant effect of the interaction between treatment and day on the

depletion time measured in shoals foraging on aggregated and scattered treatments (negative

binomial GLMM: $\chi2 = 6.263$, df $= 1$, $p = 0.012$). Depletion time decreased over the first four

days of the experiment, before reaching a plateau at around 60 seconds on the fifth day of

testing (Figure 2A). There was a significant effect of the interaction between treatment and

day on the time spent moving (GLMM: $\chi2 = 13.199$, df $= 6$, $p = 0.04$). The time spent

moving decreased over days in individuals from the control treatment, whereas individuals

from aggregated and scattered treatments increased their time spent moving over the first

three repeated tests (Figure 2B).

Figure 2. Visualisation of collective foraging efficiency and individual time spent moving.

(A) Time elapsed until the depletion of the arena (i.e., when the last prey was consumed) as a

function of treatment and for the 10 consecutive days (seconds, mean ± 95%CI). Only

'scattered' (blue) and 'aggregated' (red) treatments are displayed because there was no prey

in the 'control' treatment. (B) Proportion of time spent moving (i.e. speed greater than 0.5 body lengths per second) during the first 60 seconds of the foraging trials (mean ± 95%CI).

Movement characteristics

There was a significant effect of the interaction between treatment and day on the total distance travelled (GLMM, Day: Treatment: $\chi 2 = 104.888$, df = 6, $p < 0.001$). Distances travelled increased over the first three days for individuals foraging on either aggregated or scattered prey distribution, whereas the distances travelled by individuals in the control treatment decreased over the first five tests (Figure 3A).

There was a significant effect of the interaction between treatment and day on the average distance to the nearest neighbour (GLMM, Day: Treatment: $\chi 2 = 21.882$, df = 6, $p = 0.001$). Distances to nearest neighbour tended to remain consistent over the repeated tests in fish from the control treatment, whereas individuals from both aggregated and scattered treatments increased their distance to their nearest neighbour over at least the first five repetitions of the foraging task (Figure 3B).

There was a significant effect of the interaction between treatment and day on the acceleration entropy (GLMM, Day: Treatment: $\chi 2 = 56.681$, df = 6, $p < 0.001$). Acceleration entropy increased over the first 5 days of testing before stabilising in fish from both aggregated and scattered treatments, whereas it slightly decreased over the repeated tests in fish from the control treatment (Figure 3C).

Figure 3. Temporal changes of individuals' movement characteristics as a function of
treatments (mean ± 95%CI). (A) Total distance travelled (in cm), calculated over all frames
spent moving (i.e., speed greater than 0.5 BL.s^{-1}). (B) Average distance to nearest neighbour
(in body length), calculated over all frames spent moving. (C) Acceleration entropy

calculated as the median of entropy values of all bouts of at least 15 consecutive frames spent

moving.

Foraging success

The best supported models and the relative importance of terms varied across days.

On day 1, the three best supported models included distance to NN (model 9), distance to NN

and total distance travelled (model 13), as well as distance to NN and acceleration entropy

(model 11). All three models had a delta AICc < 2 and the second and third best supported

models both had an evidence ratio smaller than 2 (Table S3). Distance to NN was identified

from the model averaging procedure to be the most important predictor of foraging success

on day 1 (Table 1). Individuals with greater average distance to their nearest neighbour had a

greater foraging success; a result which was strongly driven by the aggregated treatment

(Figure 4A).

On day 3, the two best supported models included total distance travelled, distance to

NN and acceleration entropy (model 15), and all four predictors (model 16). Both models had

a delta AICc < 2 and the second-best supported model had an evidence ratio smaller than 2

(Table S3). Acceleration entropy was identified from the model averaging procedure to be the

most important predictor of foraging success on day 3 (Table 1). Individuals with greater

values of median acceleration entropy had a greater foraging success, independently of the

treatment (Figure 4B). Note that distance travelled also had a strong relative importance

(RI=0.89), despite being marginally non-significant (Table 1). Individuals that travelled

greater distance had a greater foraging success, independently of the treatment (Figure 4C).

On day 5, a single best supported model, including total distance travelled (model 5),

was identified. All three following best ranked models had a delta AICc > 2 and evidence

ratios ranging from 2.8 to 3, making the best ranked model (model 5) 2.8 to 3 times more

likely to be the best model than any of the three following best ranked models (Table S3).

Distance travelled was identified from the model averaging procedure to be the most

important predictor of foraging success on day 5 (Table 1). Individuals that travelled greater

distance had a greater foraging success, independently of the treatment (Figure 4D).

On day 7, a single best supported model, including acceleration entropy (model 3),

was identified. All three following best ranked models had a delta AICc > 2 and evidence

ratios ranging equals to 2.9, making the best ranked model (model 3) 2.9 times more likely to

be the best model than any of the three follow-up models (Table S3). Acceleration entropy

was identified from the model averaging procedure to be the most important predictor of

foraging success on day 7 (Table 1). Individuals with greater values of median acceleration

entropy had a greater foraging success, independently of the treatment (Figure 4E).

Table 1. Standardized model averaged coefficient estimates from the 95% candidate set of

models to determine the best predictors of foraging success within each day of testing.

Occurrence denotes the number of models that contain the term. Standardized model

averaged coefficient estimates (based on the 95% candidate set of models) are reported with

the standard error (SE) and the 95% confidence interval (95%CI). The relative importance

(RI) of each term is the sum of the Akaike weights over the models in which that tern appears

(based on the 95% candidate set of models). Important terms of significant effects are

indicated in bold within each distinct day, and marginally non-significant effects are

indicated by *.

Day	Terms	Occurrence	Estimates	SE	95%CI	RI
1	Intercept	10	0.954	0.111	[0.736, 1.172]	-
1	Treatment	4	-0.018	0.114	[-0.241, 0.204]	0.25
1	Distance travelled	5	-0.079	0.147	[-0.367, 0.208]	0.401
1	**Distance to NN**	**8**	**0.33**	**0.145**	**[0.046, 0.614]**	**0.96**
1	Acceleration entropy	4	-0.036	0.104	[-0.24, 0.168]	0.303
3	Intercept	8	1.618	0.049	[1.521, 1.714]	-
3	Treatment	3	-0.04	0.079	[-0.195, 0.116]	0.362
3	Distance travelled*	5	0.259	0.146	[-0.027, 0.545]	0.888
3	Distance to NN	5	0.13	0.113	[-0.092, 0.351]	0.713
3	**Acceleration entropy**	**7**	**0.296**	**0.136**	**[0.029, 0.562]**	**0.965**
5	Intercept	6	1.708	0.044	[1.622, 1.793]	-
5	Treatment	3	-0.009	0.047	[-0.101, 0.084]	0.264
5	**Distance travelled**	**6**	**0.395**	**0.089**	**[0.221, 0.569]**	**1**
5	Distance to NN	2	0.001	0.04	[-0.078, 0.079]	0.199
5	Acceleration entropy	2	0.007	0.047	[-0.086, 0.099]	0.21
7	Intercept	16	1.732	0.045	[1.644, 1.819]	-
7	Treatment	8	-0.005	0.046	[-0.095, 0.084]	0.274
7	Distance travelled	8	0.011	0.061	[-0.108, 0.13]	0.254
7	Distance to NN	8	0.004	0.046	[-0.085, 0.093]	0.252
7	**Acceleration entropy**	**8**	**0.239**	**0.104**	**[0.036, 0.443]**	**0.944**

Figure 4. Visualisation of the effects of the movement characteristics identified to be

significant predictors of foraging success from the model averaging procedure performed

within days 1, 3 and 5. (A) The effect of the average distance to nearest neighbour (in

average body length unit) on individual foraging success on day 1. (B) The effect of the

average acceleration entropy of movement bouts of at least 15 frames on individual foraging

success on day 3. (C) The effect of total distance travelled (in cm) on individual foraging

success on day 3. (D) The effect of total distance travelled (in cm) on individual foraging

success on day 5. (E) The effect of the average acceleration entropy of movement bouts of at

least 15 frames on individual foraging success on day 7. All visualisations are presented for

individuals foraging on aggregated and scattered distribution of prey distinctively.

DISCUSSION

We showed that the movement characteristics of individuals changed over repeated foraging trials. These changes mostly occurred within the first 3 to 5 days of our repeated tests and differed between control and foraging treatments (i.e., scattered and aggregated treatments). Unlike control individuals, fish foraging on scattered and aggregated prey significantly increased their time spent moving over the first 3 days of testing, resulting in an increase of collective foraging efficiency. Notable changes in the behaviour of foraging individuals over this time were observable in all three movement variables considered in this study. Overall, we found increases in both distance travelled and distance to nearest neighbour in fish from scattered and aggregated treatments, with pronounced changes occurring between days 1 and 3, which contrasted with control fish that maintained consistent values of distance travelled and distance to NN over this time. Acceleration entropy also increased in foraging fish over the first 5 days, with the most pronounced increase occurring between days 3 and 5, which differs from control fish whose entropy tended to remain unchanged over time. As hypothesised, all three variables seemed to reach a plateau initiated somewhere between days 3 and 5. We found no effect of the spatial distribution of prey on the movement of foraging individuals. Furthermore, we found that the relative importance of the variables considered here changed over time; with an important effect of social proximity on the first test, while search effort and unpredictability in acceleration had more important effects on variation in foraging success in the later tests.

The changes occurring over time suggest an important effect of the knowledge individuals possess about their foraging context on their movement characteristics. Without any prior information on the surrounding environment (i.e., day 1), individuals from all treatments travelled similar distances, were equally unpredictable in their variation in speed, and positioned themselves within approximately 3 body lengths from their nearest neighbour.

Such relatively close proximity to the nearest neighbour is typically observed in species

known for their strong schooling behaviour when exploring unfamiliar experimental or

natural environments (e.g., Calovi et al., 2018; Schaerf et al., 2017; Viscido et al., 2004;

Ward et al., 2017). As hypothesised, our results suggest that individuals displayed grouping

behaviour in absence of prior information, by maintaining close social proximity and with

individuals leaving or joining others on relatively few occasions (see supplementary

material). Social information is assumed to enhance integration of information, particularly

when exploring novel environments (Berdahl et al., 2013; Couzin, 2007; Sumpter et al.,

2008). In the absence of prior information, grouping behaviour in fish has been found to

enhance both individual foraging and the ability to detect and avoid potential predators

(Pitcher et al., 1982; Ward et al., 2011). As hypothesised, social proximity was the most

important predictor of foraging success on the first foraging test. Although we found that

grouping behaviour was most pronounced on that day, individuals that maintained greater

distances to their nearest neighbour tended to have a higher foraging success throughout; an

effect driven by the aggregated treatment. This surprising result suggests that competition

may be an important driver of movement characteristics even in the absence of private

information about the quantity of food available.

Over the repeated tests, individuals were able to reinforce their private information,

which included the presence (or not, in the case of the control shoals) of food in the arena, its

spatial occurrence, and the presence of a consistent number of competitors. This is likely to

have driven the changes observed in search effort, unpredictability in acceleration and social

proximity. Within each repeated test, all fish within an experimental group may have

responded to expected presence of food (for scattered and aggregated treatments only) by

increasing the search effort (i.e., swimming greater distances) within the first minutes of the

foraging test, and may have attempted to mitigate competition by maintaining greater

distances to conspecifics. Yet, the low relative importance of social proximity in explaining

variation in foraging success suggest that other mechanisms are of primary importance.

Rather than monitoring positions of local neighbours at any time and maintaining appropriate

distances, which may alter individuals' efficiency at searching for prey, staying away from

other foragers may be achieved by performing pronounced and sudden accelerations upon

arrival of a conspecific within close proximity. An individual which just discovered a prey

and is in the process of consuming a prey provides attracts conspecifics (Harpaz &

Schneidman, 2020). Joining a successful forager may increase the probability of making a

local discovery or offers an opportunity for kleptoparasitism (Hansen et al., 2016). Fleeing

from an approaching competitor would then allow an individual to completely ingest a prey,

minimizing the risk of kleptoparasitism. Such interactions based on social information and

driven by competition would explain the changes in acceleration entropy observed in

scattered and aggregated treatments from the third day onwards. In addition, grouping events,

where fish were in close proximity, were frequent but tended to be short-lived (see supplementary

material). This interpretation is supported by our model averaging analysis, which showed

that the distance travelled and acceleration entropy were the most important predictors of

foraging success during the repeated tests.

Our results show evidence that while the presence of food affects individual

movement, there was no effect of the spatial distribution of prey. It is known that fish can

acquire a variety of information about their current ecological context by sensing olfactory

cues diluted in the water, which can lead to behavioural changes (e.g., Hoare et al., 2004;

Ferrari et al., 2010; Sosna et al. 2019). Previous work on schooling fish suggested that

individuals are less frequently observed at close proximities form one another and have

higher magnitude of acceleration in the presence of food odour (Schaerf et al., 2017), which

is in line with our findings. Furthermore, fish can retain information about temporal and

spatial food occurrence and use this to improve foraging (Reebs, 2000; Van Bergen et al.,

2004). Our results show no difference in changes of movement characteristics between fish

from the aggregated and scattered treatments, which suggest that knowledge of prey

availability triggers behavioural responses that are independent of their spatial distribution.

This surprising result goes against our initial working hypothesis and differs from previous

work on juvenile walleye pollock (*Theragra chalcogramma)* conducted by Ryer & Olla

(1995), as well as a study on a broad range of open-ocean predatory fish (Humphries et al.,

2010), but is in line with a study on foraging groups of three-spined sticklebacks

(*Gasterosteus aculeatus*) conducted by Hansen et al. (2016). Although our arena is consistent

in size with those used in prior studies, it is possible that the distance between prey items in

the scattered treatment did not differ sufficiently from the distance separating each pair of

patches used in the aggregated treatment. When foraging in the wild, individuals may have to

explore larger portions of the environment between successful samplings, inducing higher

energetic and time costs from search and transport. Such costs may be compensated by

moving in a manner that maximizes food discovery and reduces costs of competition (e.g.,

grouping when foraging on clumped patches of food, foraging asocially when the food is

distributed across the environment). Additionally, the temporal uncertainty of food

occurrence in natural environments may reinforce the need for pronounced behavioural

adjustment in response to different spatial distribution of resources. Therefore, the scale used

to test the two distributions of prey in our study, combined with the experimental control for

temporal uncertainty by making food available within each daily test, may not have imposed

high enough costs of transport and search required to observe distinct movement patterns.

Overall, we found that behavioural changes appeared to be driven by individual

information about the foraging context, including learned presence of food and competitors,

but were unaffected by the spatial distribution of that food. As individuals reinforced their

information about foraging conditions, they actively avoided close contact with other foragers

through sudden changes of speed in order to minimize competition, which results in dynamic

grouping association patterns. Further research considering temporal unpredictability and a

larger experimental scale, where scattered (or patches of) prey would be separated by greater

distances, would likely promote the use of fundamentally different movement strategies at the

individual level and would ultimately provide a better understanding of how resource-driven

individual movement strategies affect group self-organisation.

REFERENCES

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects
models Usinglme4. *Journal of Statistical Software*, *67*(1), 1-48.
https://doi.org/10.18637/jss.v067.i01

Barton, K. (2020). MuMIn: multi-model inference. R package version 1.43.17

Beauchamp, G., & Ruxton, G. D. (2014). Frequency-dependent conspecific attraction to food
patches. *Biology letters*, *10*(8), 20140522. https://doi.org/10.1098/rsbl.2014.0522

Berdahl, A., Torney, C. J., Ioannou, C. C., Faria, J. J., & Couzin, I. D. (2013). Emergent
sensing of complex environments by mobile animal groups. *Science*, *339*(6119), 574-576.
DOI: 10.1126/science.1225883

Borchers, H.W. (2019). Pracma: Practical Numerical Math Functions. R package version
2.2.9.

Branson, K., Robie, A.A., Bender, J., Perona, P. & Dickinson, M.H. (2009) High-throughput
ethomics in large groups of Drosophila. *Nature Methods*, 6, 451–457.
DOI:10.1038/NMETH.1328

Burnham, K.P. & Anderson, D.R. (2002) Model Selection and Multimodel Inference: A
Practical Information-Theoretic Approach, 2nd edn. Springer, New York.
Calovi, D. S., Litchinko, A., Lecheval, V., Lopez, U., Pérez Escudero, A., Chaté, H., ... &
Theraulaz, G. (2018). Disentangling and modeling interactions in fish with burst-and-coast

swimming reveal distinct alignment and attraction behaviors. *PLoS computational*

biology, *14*(1), e1005933. https://doi.org/10.1371/journal.pcbi.1005933

Clark, C. W., & Mangel, M. (1986). The evolutionary advantages of group

foraging. *Theoretical population biology*, *30*(1), 45-75. https://doi.org/10.1016/0040-

5809(86)90024-9

Coolen, I., Bergen, Y. V., Day, R. L., & Laland, K. N. (2003). Species difference in adaptive

use of public information in sticklebacks. *Proceedings of the Royal Society of London. Series*

B: Biological Sciences, *270*(1531), 2413-2419. https://doi.org/10.1098/rspb.2003.2525

Couzin, I. (2007). Collective minds. *Nature*, *445*(7129), 715-715.

Cresswell, W. (1998). Relative competitive ability changes with competitor density: evidence

from feeding blackbirds. *Animal Behaviour*, *56*(6), 1367-1373.

https://doi.org/10.1006/anbe.1998.0902

Croft, D. P., Albanese, B., Arrowsmith, B. J., Botham, M., Webster, M., & Krause, J. (2003).

Sex-biased movement in the guppy (*Poecilia reticulata*). *Oecologia*, *137*(1), 62-68. DOI

10.1007/s00442-003-1268-6

Ferrari, M. C., Wisenden, B. D., & Chivers, D. P. (2010). Chemical ecology of predator–prey

interactions in aquatic ecosystems: a review and prospectus. *Canadian Journal of*

Zoology, *88*(7), 698-724. https://doi.org/10.1139/Z10-029

Fox, J. & Weisberg, S. (2019). *An R Companion to Applied Regression*, Third edition. Sage,

Thousand Oaks CA.

Galef, Jr, B. G., & Giraldeau, L. A. (2001). Social influences on foraging in vertebrates:

causal mechanisms and adaptive functions. *Animal behaviour*, *61*(1), 3-15.

https://doi.org/10.1006/anbe.2000.1557

Gelman, A., Su, Y. S., Yajima, M., Hill, J., Pittau, M. G., Kerman, J., Zheng, T., Dorie, V. &

Su, M.Y.S. (2015). Package 'arm'. *Data Analysis Using Regression and*

Multilevel/Hierarchical Models. R package, version 1.11-2.

Giraldeau, L. A., & Beauchamp, G. (1999). Food exploitation: searching for the optimal

joining policy. *Trends in Ecology & Evolution*, *14*(3), 102-106.

https://doi.org/10.1016/S0169-5347(98)01542-0

Giraldeau, L. A., & Caraco, T. (2018). *Social foraging theory* (Vol. 73). Princeton University

Press.

Giraldeau, L. A., Valone, T. J., & Templeton, J. J. (2002). Potential disadvantages of using

socially acquired information. *Philosophical Transactions of the Royal Society of London.*

Series B: Biological Sciences, *357*(1427), 1559-1566. https://doi.org/10.1098/rstb.2002.1065

Godin, J. G. J. (1978). Behavior of juvenile pink salmon (*Oncorhynchus gorbuscha*

Walbaum) toward novel prey: influence of ontogeny and experience. *Environmental Biology*

of Fishes, *3*(3), 261-266. https://doi.org/10.1007/BF00001451

Grand, T. C., & Dill, L. M. (1999). The effect of group size on the foraging behaviour of

juvenile coho salmon: reduction of predation risk or increased competition?. *Animal*

Behaviour, *58*(2), 443-451. https://doi.org/10.1006/anbe.1999.1174

Grueber, C. E., Nakagawa, S., Laws, R. J., & Jamieson, I. G. (2011). Multimodel inference in

ecology and evolution: challenges and solutions. *Journal of evolutionary biology*, *24*(4), 699-

711. https://doi.org/10.1111/j.1420-9101.2010.02210.x

Hansen, M. J., Ward, A. J., Fürtbauer, I., & King, A. J. (2016). Environmental quality

determines finder-joiner dynamics in socially foraging three-spined sticklebacks

(*Gasterosteus aculeatus*). *Behavioral Ecology and Sociobiology*, *70*(6), 889-899.

https://doi.org/10.1007/s00265-016-2111-5

Harel, R., Spiegel, O., Getz, W. M., & Nathan, R. (2017). Social foraging and individual

consistency in following behaviour: testing the information centre hypothesis in free-ranging

vultures. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1852), 20162654.

https://doi.org/10.1098/rspb.2016.2654

Harpaz, R., & Schneidman, E. (2020). Social interactions drive efficient foraging and income

equality in groups of fish. *Elife*, *9*, e56196. https://doi.org/10.7554/eLife.56196

Higginson, A. D., Fawcett, T. W., Trimmer, P. C., McNamara, J. M., & Houston, A. I.

(2012). Generalized optimal risk allocation: foraging and antipredator behavior in a

fluctuating environment. *The American Naturalist*, *180*(5), 589-603.

https://doi.org/10.1086/667885

Hoare, D. J., Couzin, I. D., Godin, J. G., & Krause, J. (2004). Context-dependent group size choice in fish. *Animal Behaviour*, *67*(1), 155-164. https://doi.org/10.1016/j.anbehav.2003.04.004

Humphries, N. E., Queiroz, N., Dyer, J. R., Pade, N. G., Musyl, M. K., Schaefer, K. M., ... & Sims, D. W. (2010). Environmental context explains Lévy and Brownian movement patterns of marine predators. *Nature*, *465*(7301), 1066-1069. https://doi.org/10.1038/nature09116

Humphries, N. E., Weimerskirch, H., Queiroz, N., Southall, E. J., & Sims, D. W. (2012). Foraging success of biological Lévy flights recorded in situ. *Proceedings of the National Academy of Sciences*, *109*(19), 7169-7174. https://doi.org/10.1073/pnas.1121201109

Ioannou, C. C., Bartumeus, F., Krause, J., & Ruxton, G. D. (2011). Unified effects of aggregation reveal larger prey groups take longer to find. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1720), 2985-2990. https://doi.org/10.1098/rspb.2011.0003

Jolles, J. W., Mazué, G. P., Davidson, J., Behrmann-Godel, J., & Couzin, I. D. (2020). Schistocephalus parasite infection alters sticklebacks' movement ability and thereby shapes social interactions. *Scientific reports*, *10*(1), 1-11. https://doi.org/10.1038/s41598-020-69057-0

Kendal, R. L., Coolen, I., & Laland, K. N. (2004). The role of conformity in foraging when personal and social information conflict. *Behavioral Ecology*, *15*(2), 269-277. https://doi.org/10.1093/beheco/arh008

Laland, K. N., & Williams, K. (1998). Social transmission of maladaptive information in the guppy. *Behavioral Ecology*, *9*(5), 493-499. https://doi.org/10.1093/beheco/9.5.493

Lewis, J. S., Wartzok, D., & Heithaus, M. R. (2013). Individuals as information sources: Could followers benefit from leaders' knowledge?. *Behaviour*, *150*(6), 635-657. https://doi.org/10.1163/1568539X-00003075

Nakagawa, S., & Freckleton, R. P. (2011). Model averaging, missing data and multiple imputation: a case study for behavioural ecology. *Behavioral Ecology and Sociobiology*, *65*(1), 103-116. https://doi.org/10.1007/s00265-010-1044-7

Pincus, S. M. (1991). Approximate entropy as a measure of system complexity. *Proceedings of the National Academy of Sciences*, *88*(6), 2297-2301. https://doi.org/10.1073/pnas.88.6.2297

Pitcher, T. J., Magurran, A. E., & Winfield, I. J. (1982). Fish in larger shoals find food

faster. *Behavioral Ecology and Sociobiology*, *10*(2), 149-151.

https://doi.org/10.1007/BF00300175

R Core Team (2019). R A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienne, Austria.

Reader, S. M., & Laland, K. N. (2000). Diffusion of foraging innovations in the

guppy. *Animal Behaviour*, *60*(2), 175-180. https://doi.org/10.1006/anbe.2000.1450

Reebs, S. G. (2000). Can a minority of informed leaders determine the foraging movements

of a fish shoal?. *Animal behaviour*, *59*(2), 403-409. https://doi.org/10.1006/anbe.1999.1314

Riotte-Lambert, L. & Matthiopoulos, J. (2020) Environmental Predictability as a Cause and

Consequence of Animal Movement. *Trends in ecology and evolution*, 35, 163–174.

https://doi.org/10.1016/j.tree.2019.09.009

Ryer, C. H., & Olla, B. L. (1995). Influences of food distribution on fish foraging

behaviour. *Animal Behaviour*, *49*(2), 411-418. https://doi.org/10.1006/anbe.1995.0054

Schaerf, T. M., Dillingham, P. W., & Ward, A. J. (2017). The effects of external cues on

individual and collective behavior of shoaling fish. *Science advances*, *3*(6), e1603201. https:/

doi.org/10.1126/sciadv.1603201

Snijders, L., Kurvers, R. H., Krause, S., Tump, A. N., Ramnarine, I. W., & Krause, J. (2019).

Females facilitate male food patch discovery in a wild fish population. *Journal of Animal*

Ecology, *88*(12), 1950-1960. https://doi.org/10.1111/1365-2656.13086

Snijders, L., Krause, S., Tump, A. N., Breuker, M., Ortiz, C., Rizzi, S., ... & Kurvers, R. H.

(2021). Causal evidence for the adaptive benefits of social foraging in the

wild. *Communications Biology*, *4*(1), 1-8. https://doi.org/10.1111/1365-2656.13086

Sosna, M. M., Twomey, C. R., Bak-Coleman, J., Poel, W., Daniels, B. C., Romanczuk, P., &

Couzin, I. D. (2019). Individual and collective encoding of risk in animal

groups. *Proceedings of the National Academy of Sciences*, *116*(41), 20556-20561.

https://doi.org/10.1073/pnas.1905585116

Sumpter, D., Buhl, J., Biro, D., & Couzin, I. (2008). Information transfer in moving animal

groups. *Theory in biosciences*, *127*(2), 177-186. https://doi.org/10.1007/s12064-008-0040-1

Templeton, J. J., & Giraldeau, L. A. (1995). Patch assessment in foraging flocks of European

starlings: evidence for the use of public information. *Behavioral Ecology*, *6*(1), 65-72.

https://doi.org/10.1093/beheco/6.1.65

Templeton, J. J., & Giraldeau, L. A. (1996). Vicarious sampling: the use of personal and

public information by starlings foraging in a simple patchy environment. *Behavioral Ecology*

and Sociobiology, *38*(2), 105-114. https://doi.org/10.1007/s002650050223

Valone, T. J. (1989). Group foraging, public information, and patch estimation. *Oikos*, 357-

363. https://doi.org/10.2307/3565621

Valone, T. J. (1991). Bayesian and prescient assessment: foraging with pre-harvest

information. *Animal Behaviour*, *41*(4), 569-577. https://doi.org/10.1016/S0003-

3472(05)80897-8

Van Bergen, Y., Coolen, I., & Laland, K. N. (2004). Nine-spined sticklebacks exploit the

most reliable source when public and private information conflict. *Proceedings of the Royal*

Society of London. Series B: Biological Sciences, *271*(1542), 957-962.

https://doi.org/10.1098/rspb.2004.2684

Vernables, W. N. & Ripley, B. D. (2002). Modern applied statistics with S, 4th ed. Springer,

New York. ISBN 0-387-95457-0.

Viscido, S. V., Parrish, J. K., & Grünbaum, D. (2004). Individual behavior and emergent

properties of fish schools: a comparison of observation and theory. *Marine Ecology Progress*

Series, *273*, 239-249. https://doi.org/10.3354/meps273239

Viswanathan, G. M., Afanasyev, V., Buldyrev, S. V., Havlin, S., Da Luz, M. G. E., Raposo,

E. P., & Stanley, H. E. (2000). Lévy flights in random searches. *Physica A: Statistical*

Mechanics and its Applications, *282*(1-2), 1-12. https://doi.org/10.1016/S0378-

4371(00)00071-6

Ward, A. J., Herbert-Read, J. E., Sumpter, D. J., & Krause, J. (2011). Fast and accurate

decisions through collective vigilance in fish shoals. *Proceedings of the National Academy of*

Sciences, *108*(6), 2312-2315. https://doi.org/10.1073/pnas.1007102108

Ward, A. J., Schaerf, T. M., Herbert-Read, J. E., Morrell, L., Sumpter, D. J., & Webster, M.

M. (2017). Local interactions and global properties of wild, free-ranging stickleback

shoals. *Royal Society open science*, *4*(7), 170043. https://doi.org/10.1098/rsos.170043

Ward, P. & Zahavi, A.1973. The importance of certain assemblages of birds as 'information

centres' for food-finding. Ibis, 115, 517–534. https://doi.org/10.1111/j.1474-

919X.1973.tb01990.x

Wickham, H. (2016). ggplot2: elegant graphics for data analysis, Springer, New York.