A new paradigm for leprosy diagnosis based on host gene expression

Insights from leprosy lesions transcriptomics

Thyago Leal-Calvo[1], Charlotte Avanzi[2,#a], Mayara Abud Mendes[1], Andrej Benjak[2,#b],

Philippe Busso[2], Roberta Olmo Pinheiro[1], Euzenir Nunes Sarno[1], Stewart T. Cole[2,3],

Milton O. Moraes[1*]

Affiliations

[1] Laboratório de Hanseníase, Instituto Oswaldo Cruz, FIOCRUZ, Rio de Janeiro, Rio

de Janeiro, Brazil

[2] Global Health Institute, École Polytechnique Fédérale de Lausanne, Lausanne,

Switzerland

[3] Institut Pasteur, Paris, France

[#a]Current address: Department of Microbiology, Immunology and Pathology,

Mycobacteria Research Laboratories, Colorado State University, Fort Collins,

Colorado, United States of America

[#b]Current address: Department for BioMedical Research, Oncogenomics Laboratory,

University of Bern, Bern, Switzerland

* Corresponding author

E-mail: milton.moraes@fiocruz.br (MOM)

Abstract

Transcriptional profiling is a powerful tool to investigate and detect human diseases. In this study, we used bulk RNA-sequencing (RNA-Seq) to compare the transcriptomes in skin lesions of leprosy patients or matched controls affected by other dermal conditions such as granuloma annulare, a confounder for paucibacillary leprosy. We identified five genes capable of accurately distinguishing multibacillary and paucibacillary leprosy from other skin conditions. Indoleamine 2,3-dioxygenase 1 (*IDO1*) expression alone was highly discriminatory, followed by *TLR10*, *BLK*, *CD38*, and *SLAMF7*, whereas the *HS3ST2* and *CD40LG* mRNA separated multi- and paucibacillary leprosy. Finally, from the main differentially expressed genes (DEG) and enriched pathways, we conclude that paucibacillary disease is characterized by epithelioid transformation and granuloma formation, with an exacerbated cellular immune response, while multibacillary leprosy features epithelial-mesenchymal transition with phagocytic and lipid biogenesis patterns in the skin. These findings will help catalyze the development of better diagnostic tools and potential host-based therapeutic interventions. Finally, our data may help elucidate host-pathogen interplay driving disease clinical manifestations.

Author Summary

Despite effective treatment, leprosy is still a significant public health issue in more than 120 countries, with more than 200 000 new cases yearly. The disease is caused mainly by *Mycobacterium leprae*, a slow-growing bacillus still uncultivable in axenic media. This limitation has hampered basic research into host-pathogen

interaction and the development of new diagnostic assays. Currently, leprosy is

diagnosed clinically, with no standalone diagnostic assay accurate enough for all

clinical forms. Here, we use RNA-seq transcriptome profiling in leprosy lesions and

granuloma annulare to identify mRNA biomarkers with potential diagnostic

applications. Also, we explored new pathways that can be useful in further

understanding the host-pathogen interaction and how the bacteria bypass host

immune defenses. We found that *IDO1*, a gene involved with tryptophan catabolism,

is an excellent candidate for distinguishing leprosy lesions from other dermatoses.

Additionally, we observed that a previous signature of keratinocyte development and

cornification negatively correlates with epithelial-mesenchymal transition genes in the

skin, suggesting new ways in which the pathogen may subvert its host to survive and

spread throughout the body. Our study identifies new mRNA biomarkers that can

improve leprosy diagnostics and describe new insights about host-pathogen

interactions in human skin.

Introduction

Leprosy is a chronic infectious disease caused mainly by the slow-growing intracellular pathogen *Mycobacterium leprae* that does not grow in axenic media. This bacterium resides preferentially in skin macrophages and Schwann cells in peripheral nerves, inducing dermatosis and/or neuritis. Patients can present several distinct clinical forms according to their immune response, histopathological characterization, and bacterial load. A localized tuberculoid form (TT) is characterized by low bacterial counts and a strong cellular immune response. Conversely, in the opposite lepromatous (LL) pole, a disseminated form, patients exhibit several lesions, a predominantly humoral response, and a high bacterial load in the tissues [1–3]. Borderline forms are classified according to their proximity to the poles. For operational and treatment purposes, leprosy is classified by the World Health Organization as paucibacillary (PB) or multibacillary (MB), based on the number of skin lesions, associated with nerve involvement or the bacilli detection in slit-skin smears [4].

Early and precise diagnosis is instrumental to leprosy control since delay in diagnosis leads to late multidrug therapy, higher disability risk, and continuing transmission, as highlighted by the 200,000 new cases consistently reported annually in the last 10 years [4,5]. However, bacteriological, immunological, genetics or molecular methods are not sufficient for specific diagnosis when used alone. Diagnosis most commonly relies on clinical evaluation, occasionally complemented with histopathological examination and bacterial counts, but these procedures are mostly performed in national reference centers [4,6].

Efforts have been deployed to improve leprosy diagnostics using cutting-edge

technologies, such as molecular identification of *M. leprae*, serological tests for

specific bacterial antigens, and quantification of host biomarkers in plasma or *in vitro*

whole blood assays (WBA) [7–9]. Overall, all methods outperform standard clinical

diagnosis and can compensate for the low accuracy in detecting PB patients

[4,7,8,10–14]. Yet, until now such investigations involved comparing confirmed leprosy

cases against healthy endemic controls, who are not representative of individuals with

suspected leprosy. Here, other skin conditions represent a better comparator.

Identification of markers for early infection is hindered by our poor

understanding of pathogenicity and the mechanism by which patients develop one or

the other form of leprosy, and nerve injuries [15]. Gene expression signatures have

been used as diagnostic tools for several illnesses, from infectious [10–12,14] and

autoimmune diseases [16,17] to cancer [18–20]. Some signatures have already been

approved for clinical use [12,21–23]. In leprosy, findings from past studies indicate the

great potential of expression profiling for disease diagnosis [24–27]. Nonetheless, they

were limited by the number of patients [28], or lacked proper epidemiological controls,

such as differential diagnosis groups.

Here, we applied a combination of bulk RNA sequencing and quantitative

validation by RT-qPCR on RNA extracted from skin biopsies of various leprosy forms

and from non-leprosy patients to define a specific leprosy host signature applicable to

diagnosis. Then, we explored gene expression patterns to improve our understanding

of the immunopathogenic mechanisms towards leprosy polarization.

Results

Discrimination of leprosy *vs.* non-leprosy lesions based on mRNA expression

RNA sequencing was used for pinpointing host candidate genes capable of differentiating leprosy lesions from one of the commonest differential diagnoses of leprosy, granuloma annulare (GA), and from healthy skin. RNA from skin lesions of all leprosy clinical forms (n=33), plus GA (n=4) and healthy skin (n=5) were sequenced (S1 Table). Differentially expressed genes (DEG) in leprosy *vs.* non-leprosy (GA + healthy skin) samples resulted in 1160 DEG with a $|\log_2 FC| \geq 1$ and FDR ≤ 0.01, with 961 upregulated in leprosy forms compared to non-leprosy (Fig 1A-B and S2 Table). Exploratory hierarchical clustering of the DEG with $|\log_2 FC| \geq 1$ and FDR < 0.01 grouped all patients' samples into roughly two clusters, except for two: one BL leprosy and one GA that clustered apart from samples with the same diagnosis (Fig 1C). Gene Ontology enrichment analysis of up-regulated genes in leprosy compared to non-leprosy showed enrichment for biological processes associated with leukocyte activation, T-cell activation, immune response, response to the bacterium, neutrophil degranulation, cell killing, cytokine secretion, purinergic receptor signaling pathway, and regulation of defense response to viruses by the host (Fig 1D and S3 Table).

Fig 1. Differentially expressed genes from RNA-seq in leprosy *vs.* GA and

leprosy *vs.* non-leprosy. (A) MA plots showing the number of differentially expressed

genes (red points) with FDR < 0.1 for the comparison of leprosy *vs.* non-leprosy. (B)

Volcano plot depicting DEG from leprosy vs. non-leprosy, where violet dashed line

marks $|\log_2 FC| = 2.5$ and green horizontal dashed line the FDR = 0.01. (C) Heatmap

with hierarchical clustering of samples based on expression of the DEG from leprosy

vs. non-leprosy comparison. Color scale ranges from lower expression (blue) to higher

expression (red). Library size is given in millions. LIB, logarithmic index of bacilli. (D)

Biological processes from GO enriched for up-regulated DEG from leprosy vs. non-

leprosy comparison. FDR, false discovery rate; NL, non-leprosy; GA, granuloma

annulare; non-leprosy: GA + healthy individuals.

A total of 15 genes with the largest effect size ($|\log_2 FC| \geq 1.5$, FDR < 0.001),

highest area under the curve (AUC), and plausible involvement with leprosy

pathogenesis (S4 Table) were then validated using a two-step RT-qPCR with a new,

larger, and more heterogeneous dataset including skin lesion samples from leprosy

patients (n=24), and other common dermatoses (n=30) (S1 Table). Other

dermatological diseases (ODD) included dermatitis (n=11), eczema (n=1), erythema

(n=5), GA (n=6), lichen planus (n=2), psoriasis (n=2) and pityriasis alba (n=2) (S1

Table). A total of 12 samples per group was estimated to be sufficient to attain a power

of 85% based on the Welch t-test (PB *vs.* ODD, MB *vs.* ODD) with alpha set at 0.03

to replicate the standardized effect size ($\log_2 FC/SD$) estimated from RNA sequencing.

Relative expression using the new sample set by RT-qPCR is shown in Fig 2A. Indeed,

the validation data are in agreement with RNA sequencing, because most tested

genes were replicated by RT-qPCR in differentiating leprosy from other dermatoses

(Fig 2A) and by their mean differences (Fig 2B-C, S5 Table).

Fig 2. Technical and biological validation for selected DEG discovered from RNA

sequencing. (A) Tukey boxplots with RT-qPCR normalized (2-3 reference genes) $\log_2$

expression values (A.U) according to clinical and histopathological diagnosis. (B)

$\log_2$FC from MB-ODD and PB-ODD comparisons estimated from Bayesian linear

mixed models and their 95% credible intervals. (C) Tukey boxplot highlighting *IDO1*

RT-qPCR normalized $\log_2$ expression values by final diagnosis grouped into ODD

category. Missing values are omitted.

Next, hierarchical clustering with RT-qPCR data including missing values for

some genes (no target gene amplification by RT-qPCR) was performed to examine all

samples simultaneously. The analysis roughly revealed three major clusters (Fig 3A).

At the highest tree subdivision, one small cluster (n=6) with the dendrogram grouped

in light brown was composed of ODD samples with lower expression levels (Fig 3A).

Due to several ODD having missing values, we confirmed that these samples had

similar gene expression for the reference genes, thereby eliminating the possibility of

insufficient cDNA input. Another cluster, grouped in the light purple dendrogram,

included all MB and most PB samples (except three in light yellow dendrogram). GA

samples displayed two patterns, the first with two samples showing undetectable *IDO1*

expression (Fig 3A, bottom star symbols). The second set (n=4) is scattered among

other ODD samples (Fig 3A). It can be seen that GA and PB samples show highly

similar expression profiles for some genes (Fig 3A bottom diamond symbols),

reinforcing the difficulty in clinically discriminating between these two conditions, and

underlining the relevance of their inclusion in our comparisons [29–31]*.*

Then, by applying principal component analysis (PCA) to the 15 gene signature

obtained with the expanded sample panel tested by RT-qPCR, we uncovered two

major patterns separating leprosy lesions from ODD (Fig 3B). As expected, MB

samples appeared more homogeneous than PB and ODD samples, while the latter

were more dispersed revealing heterogeneous expression patterns (Fig 3B).

Next, we quantified the individual classification potential of these genes in

distinguishing leprosy from ODD using ROC analysis on RT-qPCR data. *IDO1*

expression alone was found to be 100% accurate using an arbitrary threshold,

followed by *BLK* (exon 11), *TLR10, CD38,* and *SLAMF7* (Fig 3C and S6 Table)*.* Finally,

to confirm the causal link between mycobacteria and our gene-set, we evaluated the

mRNA profiles induced by other live-mycobacteria using a public RNA-seq dataset

[32]. We observed that most gene expression signatures, including *IDO1,* could be

successfully replicated as induced by either *M. leprae* and/or other mycobacteria (Fig

1 in Appendix S1 and S7 Table). By contrast, some of the tested genes such as *BLK*,

CXCL9, *MS4A1,* and *TLR10* were not differentially expressed in any of the *in vitro*

assays with mycobacteria (Fig 1 in Appendix S1 and S7 Table).

Fig 3. Hierarchical clustering of RT-qPCR replicated DEG and ROC analysis. (A) Hierarchical clustering with scaled and centered normalized log$_2$ RT-qPCR expression values (arbitrary units) and annotated according to group and specific diagnosis. Dendrogram tree was cut arbitrarily and cluster analysis is for hypothesis generating purposes only. (B) Principal component analysis (PCA) with 15 genes measured by RT-qPCR and using log$_2$ normalized scaled data. For PCA only, missing values were imputed by the gene arithmetic mean. NA, not amplified, i.e., Cp > 40. In this regard,

197   there were two outliers (psoriasis and erythema), which are samples with high

198   numbers of NA values and that were imputed using the gene arithmetic mean. (C)

199   Receiver operating characteristic analysis for genes with largest AUC (95%

200   confidence intervals) from RT-qPCR replication samples (complete data are shown in

201   S6 Table). See also S1 Appendix and S1 Fig.

MB and PB gene expression profiling and mRNA-based classifier

To define a small subset of genes with high classificatory potential (i.e. with

non-overlapping expression values) to distinguish MB from PB lesions, we performed

a penalized logistic regression (LASSO) model with k-fold cross-validation trained on

the public microarray dataset [24]. This dataset was chosen because of the higher

number of PB/MB samples compared to our RNA-seq dataset. As a result, three genes

with non-zero coefficients were selected by the cross-validated LASSO model:

HS3ST2, CD40LG, and *CCR6*, but only the first two genes were most frequently

(~80%) selected across 10,000 bootstrapped samples within the training dataset (Fig

4A-B). The median misclassification error estimated by the resampling was about 4%

(±5.4% median absolute deviation), ranging from 0% to 32% (Fig 4C). Instability

assessment in the number of selected genes by LASSO (Fig 4D) showed that most

iterations resulted in four non-zero genes (range, 1-20). The final model containing the

three genes (*HS3ST2*, *CD40LG,* and *CCR6*) was evaluated on two test RNA-seq

datasets: our dataset, and the one from Montoya *et al.* including MB (n=9) and PB

(n=6) groups [28]. Penalized logistic regression demonstrated an accuracy of 100%

(lower 95% CIs: 86.8% and 78.2%, respectively) in classifying MB from PB samples

in both test RNA-seq datasets (Fig 4E-F). The *HS3ST2* gene was consistently more

expressed in MB leprosy lesions compared to PB, whereas the opposite was observed

for *CD40LG* (Fig 4E-H) and *CCR6* (S2 Fig). In both datasets, the combined expression

levels of *HS3ST2* and *CD40LG* showed good discrimination between the two groups

(Fig 4E-H). However, given the sample size and the bootstrapped estimates, it is not

currently possible to exclude *CCR6* from the model without additional replication.

Fig 4. Gene candidates identified with the penalized logistic regression (LASSO)

model as the most important to distinguish PB and MB leprosy lesions. (A)

Coefficients (log odds) from the top 10 most selected genes (i.e., non-zero) across

10,000 bootstrap samples using the microarray from Belone *et al.* as training dataset.

(B) Frequency of non-zero coefficients across all bootstrap samples. (C)

Misclassification error distribution estimated from 4-fold cross-validation (k-) across

10,000 bootstrap samples, with median error of 3.70% (±5.4% median absolute

deviation). (D) Number of genes kept across all resamples. Predicted probability from

the final model performance on this study test RNA-seq (E) and Montoya *et al.* RNA-

seq. (F) Normalized $\log_2$ gene expression (z-score) of the two most frequently selected

variables for distinguishing MB from PB samples in the (G) microarray training dataset

and (H) this study test RNA-seq. PB, paucibacillary leprosy; MB, multibacillary leprosy.

Tukey box plots with 1st, 2nd and 3rd quartiles ± 1.5 × inter quartile range (IQR)

whiskers. See also S2 Fig.

Next, to assess the dichotomy beyond cellular *vs.* humoral response in leprosy

lesions [33,34], a comparison of gene expression in MB leprosy (LL+BL+BB) *vs.* PB

(TT+BT) skin lesions was performed. Differential expression analysis with $|\log_2 FC| \geq$

1 and FDR ≤ 0.01 resulted in 112 DEGs; 69 up-regulated and 43 down-regulated (Fig

5A and S8 Table). In addition, we compared DEG to the public microarray data

available in Gene Expression Omnibus (GEO) from Belone *et al.* [24,35] using only

the FDR cutoff. With an FDR < 0.01, 161 DEGs were common to both studies, all

except one showed concordant modulation characterized by an overall high

correlation coefficient and concordance index, irrespective of the technology used, the

sample processing, and the data analysis methods (Fig 5B). Functional enrichment

analysis of the RNA-seq up-regulated genes (i.e., more expressed in MB than PB)

revealed processes involved with regulation of immune response, humoral immunity,

phagocytosis, cholesterol metabolism, complement activation among others (Fig 5C

and S9 Table). On the contrary, enrichment analysis of genes more expressed in PB

revealed biological processes such as leukocyte differentiation, lymphocyte

differentiation, lymphocyte-mediated immunity, B cell activation, STAT cascade

activation/regulation, and JAK-STAT cascade activation (Fig 5D and S10 Table),

which are consistent with exacerbated responses in granulomatous diseases.

Localized clinical forms, i.e., BT and TT, show a gene expression pattern indicative of

differentiation towards epithelioid transformation and granuloma assembly, which is

also observed in cutaneous or pulmonary sarcoidosis [36,37].

Fig 5. Differentially expressed genes from multibacillary (MB) vs. paucibacillary (PB) leprosy lesions. (A) Volcano plot showing DEG from the MB vs. PB comparison, where blue points are DE with |log₂FC| ≥1 and FDR < 0.1. (B) Scatter plots with the 161 DEG common between this study and Belone *et al.* (24) microarray for the same comparison. Red and green dashed lines indicate log₂FC of -1 and 1, respectively. Blue points are genes with the same modulation signal and red indicates discordancy. Rho, Spearman's rank correlation coefficient. CCC, Lin's concordance correlation

271   coefficient. Venn diagram on the right displays the number of DEG in each study

272   according to FDR < 0.01. (C) Biological processes from GO enriched from up-

273   regulated and (D) down-regulated DEG. FDR, false discovery rate.

Epithelial-mesenchymal transition (EMT) in the skin of multibacillary leprosy patients

In a previous microarray meta-analysis, our group has identified a consistent

down-regulation of cornification, keratinocyte differentiation, and epidermal

development-related genes in leprosy lesions, predominantly in MB patients [35]. We

hypothesized that such regulation could result from *M. leprae* inducing

dedifferentiation of keratinocytes, similar to the phenomenon described previously in

infected Schwann cells [38], and also described in skin cancer by a process known as

epithelial-mesenchymal transition (EMT) [39,40]. To test the hypothesis that such

modulation was involved with EMT, we correlated the previously identified down-

regulated genes in leprosy [35] with the list of reported genes [38], together with

canonical genes involved with "stem-cell biology" (Reactome, R-HAS-452723) and

"TGF-beta in epithelial-mesenchymal transition" (R-HAS-2173791). We found a

consistent,   moderate   negative   correlation   mostly   between

keratinization/cornification/epidermal development genes (*KRT5, KRT1, SCEL,*

DMKN, GJA1, AQP3, TWIST1, VDR, IVL, PKPL) with those involved with canonical

and alternative EMT and mesenchymal phenotypes (such as *TGFB1, TGIF2, RHOA,*

ZEB2, PSAP, CTSZ, MMP9) in our new RNA-seq dataset (Fig 6A-B, stars), and also

replicated these observations with an external microarray [24] and RNA-seq datasets

[28], respectively (Fig 6C, S3 and S4 Fig). Our results also showed a pattern in the

expression of the EMT-related genes as opposed to cytokeratins and epidermal

differentiation genes with low expression in healthy skin samples, and a linear

expression increase in PB and MB patients, especially with the microarray dataset,

except for *MMP9* (Fig 6C). From this result we hypothesized that in addition to TGFβ-

dependent immunosuppression in MB, activation of this pathway could be slowing or

arresting keratinocyte cornification processes in leprosy lesions thereby both

facilitating survival and/or spread of *M. leprae*.

Fig 6. Strongest correlations between keratinocyte and dedifferentiation genes

in leprosy lesions. (A) Heat plot with Spearman's rho correlation coefficient of the

strongest correlations after multiple testing adjustment (FDR ≤ 0.0001 and rho ≤ -0.8).

Correlations with FDR > 0.1 are filled with white. Row and column colored squares

identify gene categories (overlaps occur). Scatter plot of average $\log_2$ expression

calculated with keratinocyte-related genes previously documented as down-regulated

in leprosy skin against dedifferentiation-related genes using either (B) this study RNA-

seq dataset or (C) Belone *et al.* microarray (GSE74481). Lines were drawn based on

intercept and beta estimated from robust linear regression for all samples (black) or

separately for PB (blue), and MB (red). Spearman's rho coefficient along with 95%

nominal confidence intervals are shown inside scatter plots. See also S3 Fig and S4

Fig.

Discussion

One of the priorities in leprosy research is the development of reliable and

accurate laboratory diagnosis tools for all leprosy forms to provide efficient treatment

and prevent disability [41]. This goal includes diagnosing patients with early forms of

the disease, those with low or mild apparent symptoms, thus assisting with ambiguous

differential diagnoses, and even classifying the disease for treatment (MB *vs.* PB) [4].

Host response to infection as measured by gene expression in skin biopsies

offers diagnostic, prognostic and predictive potential. By applying host transcriptomics

to skin lesions from leprosy patients and other common confounding dermatoses that

challenge clinicians and pathologists [9,30], we identified a small set of genes that

provide a promising expression signature capable of distinguishing PB leprosy cases

from other confounding dermatological diseases. The top candidate, *IDO1,* is a gene

involved in nutritional immunity and metabolism [42–45]. Alone, the expression of this

gene was able to differentiate leprosy from non-leprosy lesions with high accuracy in

our dataset and in others. According to the latest data from single-cell analysis [46],

IDO1 has been shown to be differentially expressed in Langerhans cells from leprosy

lesions compared to healthy skin, corroborating our findings. However, *IDO1*

expression is also increased in other mycobacterial diseases such as tuberculosis

[47,48], which might decrease its specificity. The accuracy of classification could be

improved by combining measurement of *IDO1* expression with that of four other

biomarker genes *BLK, TLR10, CD38, and SLAMF7,* which also showed high

classification accuracy in the replication dataset. In parallel, the penalized logistic

regression model, evaluated on two independent datasets, demonstrated that

HS3ST2 and *CD40LG* hold great potential to differentiate between MB and PB lesions.

Considering the functional evidence for *HS3ST2* [49], it is possible that this gene may

be involved with granuloma disassembly, tissue permeability, and cellular migration in

leprosy, which would explain its overexpression in MB lesions. On the contrary,

CD40LG (also known as CD154) is more expressed in PB patients when compared to

MB with a predominant role in the activation of the microbicidal *Th1* response

associated with PB lesions [50]. After mechanistic validation of our findings,

quantifying expression levels of *HS3ST2* and *CD40LG* from leprosy lesions could be

useful to assess immune responsiveness against *M. leprae*, help patient stratification

and/or provide a basis for host-based adjuvant treatment for leprosy lesions.

One of the challenges in translating gene expression signatures into medical

diagnosis is the cost of measuring a large number of genes and transforming these

values into a unique continuous or binary classifier. So far, we were able to reproduce

the finding using both bulk RNA-sequencing and relative RT-qPCR, with the latter

being more accessible to clinicians in reference centers or central hospitals. Although

there are successful RT-qPCR gene expression-based diagnostic tests currently

approved for diagnosing sepsis [12], clinical support for prostate [22], and breast

cancer [18], there is a need for alternatives to reduce the cost and complexity of such

assays. Quantification of mRNA based on isothermal amplification either with NASBA

[51,52], RT-LAMP [53,54] or CRISPR-Cas12 [55] is conceivable for less specialized

settings without high-end equipment. Besides, combining a multi-target expression-

based diagnostic test with qPCR detection of *M. leprae* DNA could increase the

specificity and sensitivity of leprosy diagnosis [56]. Alternatively, an ELISA assay

measuring the levels of IDO1 protein from skin interstitial fluid, for example, could be

proven useful [57].

In parallel with poor diagnosis, lack of fundamental understanding of leprosy

pathogenesis has misled scientists for centuries [5,6]. Herein, we also compared the

two leprosy poles, MB and PB, and identified several pathways already known to be

associated with leprosy, such as the humoral immune response, phagocytosis, and

complement activation. Genes involved with cholesterol and fatty acids were more

expressed in MB lesions, as already reported [58–60]. Interestingly, B-cell-related

genes were more expressed in PB than MB. In fact, it seems that both poles modulate

this pathway by a distinct set of genes. Involvement of B lymphocytes in PB leprosy

pathogenesis has been described by a few groups, which may indicate differential

involvement of such cells depending on the disease pole [61,62].

M. leprae subverts host cell metabolism [63] by inducing lipid biosynthesis,

while avoiding type II (IFN-gamma) responses through a type I IFNs mechanism,

following the phagolysosomal breach that releases DNA into the cytosol [64].

However, exactly how the bacilli spread throughout the body and bypass the

microbicidal immune response remains unknown. Here, we provide robust evidence

indicating that *M. leprae* may induce EMT in the skin within keratinocytes and

macrophages, as described in Schwann cells [38]. Indeed, *M. leprae* induced

dedifferentiation of infected Schwann cells into an immature stage resembling

progenitor/stem-like phenotype [38]. These reprogramming events induced by long-

term infection with *M. leprae* resulted in mesenchymal cells capable of migratory and

immune-permissive behavior, which in turn facilitated *M. leprae* spread to skeletal and

smooth muscles and furthered macrophage recruitment [38,65]. In our previous work,

we identified a down-regulated signature of keratinocyte differentiation and

cornification gene markers in MB skin lesions [35]. Here, we showed that such genes

are inversely correlated with genes involved with EMT, especially the members of the

TGFβ-EMT pathway, such as *TGFB1, TGFBR1, TGIF2, PSAP, ZEB2* [66,67]. Some

of these genes are directly or indirectly associated with EMT, such as a *PSAP* [68],

WAS [69], *RHOA* [70–73], *CTSZ* [74], *MMP9* [75], *LOXL3* [76], *HIF1A* [77,78] among

others.

Our hypothesis that *M. leprae* is inducing dedifferentiation or slowing the

cornification process in keratinocytes is plausible, given the evidence in Schwann cells

and a few reports of infection in this cell type (Fig 7) [79,80]. Nevertheless, other

phenomena could explain EMT's role in leprosy pathogenesis, such as wound healing

or loss of the epithelial barrier. Although, given its obligatory intracellular lifestyle, *M.*

leprae induces dedifferentiation in other cell types, either directly as in Schwann cells

or indirectly via chemokine and cytokine production in lesions. Besides inducing

keratinocyte dedifferentiation to mesenchymal cells, *M. leprae* might benefit from a

decreased or alternative immune activation of these cells [81,82]. Further functional

confirmatory experiments should elucidate the causality of this correlation and provide

definitive evidence of the relationship between the bacilli and other cell types, such as

keratinocytes, fibroblasts, and epithelial cells.

Our preliminary data also showed that the enriched pathways among PB skin

lesions were consistent with profiles observed in other granulomatous diseases, such

as noninfectious sarcoidosis and granuloma annulare, or chronic infectious diseases

like tuberculosis [37,83–85]. Our findings revealed that PB (TT/BT) lesions have,

among others, JAK-STAT cascade activation, which has been implicated in

sarcoidosis and GA. Remarkably, the JAK-STAT specific biological inhibitor,

tofacitinib, has a potent effect promoting rebalance of exacerbated immunity among

sarcoidosis and granuloma annulare patients reestablishing homeostasis [83].

Another compound, everolimus, has been shown in experimental models to achieve

the same response [37] suggesting that these drugs could be useful to treat PB, but

not MB, leprosy.

To conclude, our combined findings provide highly discriminatory mRNA

signatures from skin lesions that could distinguish leprosy from other dermatological

diseases and allow disease classification by monitoring only a handful of genes. In

addition, we report new genes and pathways that are likely informative regarding how

M. leprae interacts with and subverts host cells to promote its spread within the body

and subsequent transmission.

Fig 7. Hypothetical hourglass model contextualizing the observed findings for leprosy clinical outcomes. The host-pathogen interaction in the skin leads to opposing leprosy clinical forms. Upon infection, *M. leprae* induces baseline metabolic alterations such as an increase in glucose uptake, modulation of lipid biosynthesis, reduction of mitochondrial metabolism, and upregulation of IDO-1 and type I IFN. Eventually, progression towards an unspecified inflammatory state can be observed where three ways could be anticipated: I) self-healing; II) progression towards the tuberculoid pole; or III) progression to lepromatous pole. These outcomes are driven

by specific environmental and host genetic factors. It is expected that lower (or shorter)

M. leprae exposure, food shortage, BCG vaccination, and polymorphisms in genes

controlling autophagy/granuloma formation (*NOD2*, *LRRK2*, *PRKN*) all contribute to

developing leprosy per se. Excessive inflammation is one phenotype observed, that is

also seen in other granulomatous diseases (e.g., cutaneous sarcoidosis, granuloma

annulare), especially in paucibacillary lesions. On the other pole, epithelial-

mesenchymal transition and local immunosuppression are present due to a probably

higher (and/or longer) *M. leprae* exposure, combined with host single-nucleotide

polymorphisms (SNPs) at key genes, like lipid biogenesis (*APOE*) and central

metabolism (*HIF1A, LACC1/FAMIN*), culminating in disease progression.

Materials and Methods

Patient cohort

All patients were enrolled after informed written consent was obtained with

approval from the Ethics Committee of the Oswaldo Cruz Foundation, number 151/01.

Leprosy clinical forms were classified according to the criteria of Ridley and Jopling

[2]. Leprosy patients were treated according to the operational criteria established by

the World Health Organization [4]. Leprosy and patients with other dermatological

diseases were eligible if their diagnosis was confirmed by clinical and histopathological

findings. Additionally, detection of *M. leprae* DNA by qPCR routinely performed in our

laboratory could be employed to support diagnosis [56,86]. HIV and hepatitis B

positive patients were not included in this study, in addition, we excluded individuals

with a current or previous history of tuberculosis. No other comorbidities were used to

exclude patients and further individual information is available in S1 Table. Skin biopsy

specimens containing both epidermis and dermis were obtained with 3 mm (diameter)

sterile punches following local anesthesia from the lesion site. Skin biopsies were

immediately stored in one milliliter of RNALater (Ambion, Thermo Fisher Scientific Inc.,

MA, USA) according to the manufacturer's instructions and stored in liquid nitrogen

until RNA isolation. Normal skin biopsies were from lesion-free sites of patients

diagnosed with indeterminate or pure neural leprosy.

Study Design

The main objective of this research was to identify host gene expression

patterns capable of distinguishing leprosy (including the PB forms) from other

differential diagnosis of skin lesions. Our working hypothesis was that leprosy lesions,

despite their morphological and histopathological similarity to other skin diseases, may

induce distinct patterns of gene expression in at a small subset. We predefined the

comparison of leprosy (PB+MB) from non-leprosy including GA in addition to healthy

patients for RNA sequencing experiment. In addition, we predetermined comparisons

between leprosy poles: MB vs. PB. Our samples are representative of a population of

individuals attending the Sousa Araujo Outpatient Clinic based in Rio de Janeiro,

Brazil, which also receives patients from surrounding municipalities.

RNA isolation

Snap frozen skin biopsies were thawed in wet ice and processed using TRIzol

Reagent (Ambion, Thermo Fisher Scientific Inc., MA, USA) according to the

manufacturer's instructions with the help of Polytron Homogenizer PT3100

(Kinematica AG, Switzerland). RNA was treated with DNAse using the DNAfree kit

(Thermo Fisher Scientific Inc., MA, USA) according to the standard manufacturer's

protocol, prior to use for library preparation and RT-qPCR. RNA integrity was

assessed in 1% agarose gel electrophoresis or TapeStation RNA ScreenTape (Agilent

Technology, CA, USA). During RNA isolation, samples were randomly assigned to

extraction batches and freeze-thaw cycles to minimize batch effects and the

introduction of technical artifacts. All procedures applied to samples were carried out

using reagents from the same lot. The first author conducted the experiments aware

of each sample group during the entire process, therefore, no blinding scheme was

used, although we do not rely on perceptual/abstract measurements or analyses nor

did we purposefully exclude samples.

497   Library preparation and Illumina RNA sequencing

RNA-seq libraries were prepared with 1 µg of total RNA for each sample using

the Illumina TruSeq mRNA kit (Illumina, USA) as recommended by the manufacturer

using the Illumina CD RNA indexes (Illumina, USA). Libraries were quantified and

qualified using a qPCR quantification protocol guide (KAPA Library Quantification Kits

for Illumina Sequencing platforms) and TapeStation D1000 ScreenTape (Agilent

Technologies, USA), respectively. The resulting libraries (fragment size 200-350bp)

were multiplexed (17, 17, and 19 libraries, respectively) and sequenced using the

NextSeq 500 platform (Illumina, USA), generating approximately 520 million single-

end reads of 75 nucleotides in length.

RNA-sequencing analysis

RAW bcl files were converted into .fastq using Illumina's bcl2fastq script. Then,

read quality was assessed using FastQC version 0.11.8 [87]. Next, transcript counts

were estimated using Salmon (v.1.13.0) quasi-mapping (human transcriptome

GRCh38_cdna sourced from Ensembl/RefGenie plus pre-computed salmon index,

http://refgenomes.databio.org/#hg38_cdna) with default settings and --seqBias flag

set [88]. Transcript counts were summarized into ENSEMBL gene counts using the R

v.3.6.1 package tximport v.1.12.0 [89,90] and biomaRt v.2.40.5 [91]. The expression

of sex-chromosome-specific genes, such as *UTY* and *XIST,* was used to rule out

sample mislabeling. Differential expression was estimated using DESEq2 v.1.24.0,

after filtering out weakly expressed genes with less than 10 counts per million and less

than 15 total counts in 70% of samples [92–94]. In addition to the patient's biological

sex, extraction batch and sequencing run, three surrogate variables estimated with

RUVseq v.1.18.0 were included in DESeq2's generalized linear model [95,96].

Nominal P-values were inspected with histograms and adjusted for multiple testing

according to the method [97] proposed for controlling the false discovery rate (FDR).

All $\log_2$ fold-changes were shrunken prior to DE filtering with the apeglm [94] or normal

algorithms. For visualization, counts per million (CPM) were computed with edgeR's

cpm function v.3.26.1 and variance stabilized with the parametric method [92]. Then,

surrogate variables and covariates were regressed out from the expression matrix

using limma's removeBatchEffect [98–100] before being visualized with ggplot2

v.3.3.0 [101]. Hierarchical clustering, heatmaps, and ROC analysis were all performed

with the previously processed expression matrix. Heatmap with hierarchical clustering

was drawn with ComplexHeatmap v.2.0.0 [102] or pheatmap v.1.0.12 [103] using

gene-wise scaled and centered matrix with Euclidean distance and average

agglomeration method. Overrepresentation analysis (ORA) was used to test for Gene

Ontology Biological Process (GO BP) enrichment with clusterProfiler v.3.12.0 [104]

and org.Hs.eg.db v.3.8.2 annotations [105]. Up and down-regulated lists were used as

inputs and the background list was composed of all genes subjected to differential

expression. P-values were adjusted for multiple testing using the Benjamini-Hochberg

method [97]. Raw and normalized RNA sequencing data are available in EMBL-EBI's

ENA and ArrayExpress under accessions ERP128243 and E-MTAB-10318,

respectively.

RT-qPCR

A total of 2.5 µg of RNA was reversed transcribed into cDNA using 4 µL of Vilo

Master Mix (Thermo Fisher Scientific Inc., USA) according to the manufacturer's

instructions. Then, cDNA was diluted to a final concentration of 5 ng/µL using TE buffer

(10 mM Tris-HCL and 0.1 mM EDTA in RNAse-free water). RT-qPCR was performed

using Fast Sybr Master Mix (Thermo Fisher Scientific Inc., USA) in a final reaction

volume of 10 µL. For each reaction, performed in duplicate, 5 µL of Fast Sybr Green

were combined with 200 nM of each primer, 10 ng of cDNA, and q.s.p of injection-

grade water. Thermal cycling and data acquisition were performed on Viia7 with 384

well block (Applied Biosystems, Thermo Fisher Scientific Inc., USA) following the

master mix manufacturer cycling preset with a final melting curve analysis (65 °C to

95 °C, captured at every 0.5 °C). All primers were designed with NCBI Primer-Blast

[106–109] to either flank intron(s) or span exon-exon junction(s) to avoid gDNA

amplification (S11 Table). Further, primers were quality checked for specificity, dimers

and hairpin with MFEPrimer v.3.0 [110,111] and IDT's oligoAnalyzer (https://www.idtdna.com/calc/analyzer). Data were exported from QuantStudio software v.1.3 in RDML format, which was imported to LinRegPCR v.2020.0 for RT-qPCR efficiency determination and calculation of the N_0 value [112,113]. Finally, N_0 values were imported to R and normalized using as the denominator the normalization factor (NF) calculated from the geometric mean of at least three reference genes (*RPS16*, *RPL35* and *QRICH1*), which were previously tested for stability [114]. These N_0 normalized values were used for visualization in Fig 2A. For mean difference estimation between groups, RT-qPCR data were analyzed in a Bayesian framework (Markov Chain Monte Carlo sampling, MCMC) using generalized linear mixed effect models under lognormal-Poisson error with MCMC.qpcr v.1.2.4 [115,116]. Per-gene efficiency estimates from LinRegPCR were used in conjunction with Cp (crossing point) calculated in QuantStudio software v.1.3 to generate the counts table. Then, the generalized linear mixed-effect model was fitted using three reference genes (allowing up to 20% between-group variation) with 550,000 iterations, thin = 100, and burn-in of 50,000. The model specification included the sample (factor with 51 levels) as a random effect and the diagnosis group (factor with 3 levels) as a fixed effect. MCMC diagnostics were done by inspecting chain mixing plots and linear mixed model diagnostic plots. Ninety-five percent credible intervals were drawn around the posterior means and MCMC equivalent P-values were also computed.

Reanalysis of public gene expression datasets

Belone and collaborators GSE74481 [24] and de Toledo-Pinto and cols. GSE35423 [64] microarray datasets were reanalyzed as described elsewhere [35]. Blischak and cols. [32] RNA-seq dataset (GSE67427) was reanalyzed from counts per

sample file from the author's Bitbucket repository (https://bitbucket.org/jdblischak/tb-

data/src/master/). Briefly, a normalized $\log_2$ expression matrix was regressed out for

RNA integrity number and extraction batch variables. Then, differences in gene

expression (48h post-infection) for specific genes and treatments were tested using a

gene-wise linear mixed model with a random intercept per sample (replicate) followed

by Dunnet comparison against a "mock" group using emmeans v.1.5.3. Montoya and

collaborators' dataset was retrieved from GEO (GSE125943) already normalized

(DESeq2 median ratio method) and transformed with base 2 logarithm with no further

processing [28].

Correlation analyses

For RNA-seq datasets, normalized $\log_2$ counts-per-million values were used

and $\log_2$ normalized intensities for microarray. Spearman's rank correlation method

was chosen because it is robust against outliers, does not rely on normality

assumption, and also identifies monotonic but non-linear relationships. Initially, a list

of keratinocyte/cornification/epidermal development genes that were DE in the meta-

analysis was assembled [35]. Then, lists of target genes were compiled from results

of Masaki *et al.* [38]: EMT and non-EMT; and from Reactome: R-HSA-452723

(Transcriptional regulation of pluripotent stem cells), R-HAS-5619507.3 (Activation of

HOX genes during differentiation), R-HAS-2173791 (TGFβ receptor signaling in EMT);

as well as Gene Ontology GO:0001837 (EMT), and literature for EMT canonical

markers. Pairwise Spearman correlation was calculated using the Hmisc's rcorr

function v.4.2-0 for each of the gene sets assembled. P-values were adjusted for

multiple testing using the BH method for FDR control [97]. Additionally, 95% nominal

confidence intervals were calculated using the Fieller method implemented by

correlation R package v.0.5.0 [117,118]. Next, a full correlation was computed using

the union of all gene sets against the keratinocyte/cornification signature. Scatter plots

were drawn with ggplot2 v.3.3.3 showing lines from coefficients estimated using

default robust regression (MASS::rlm v.7.3-51.4). No outliers were omitted.

Regularized (LASSO) logistic regression classification

Normalized $\log_2$ expression matrices regressed out for covariates and batches

were used as input predictors. The model was trained using the microarray dataset

from Belone et al. [24] with penalized regression (L1-norm, LASSO) and 4-fold cross-

validation (k-fold CV) with the negative binomial log-likelihood link function, glmnet

v.4.1 [119–121]. Predictors were standardized to have mean zero and unit variance

inside the cv.glmnet function. We opted for L1-norm because it results in a smaller

number of genes (#features ≤ n) with non-zero coefficients, as compared to elastic-

net or ridge regression counterparts. In addition, this model is suitable for high-

dimensional data as it combines feature selection during model tuning and training,

mitigating the effects of predictors' collinearity and reducing overfitting. To assess the

coefficients' error, misclassification error rate, feature stability and model size we used

non-parametric bootstrap (boot v.1.3.25) with 10,000 samples, with 4-fold cross-

validation inside each loop [122,123]. The final LASSO model selected by 4-fold cross-

validation contained three non-zero genes. Finally, independent RNA-seq test

datasets were used to compute the accuracy of the final model. Alternatively, the

whole process was repeated with leave-one-out cross-validation instead of k-fold. The

results were practically indistinguishable, especially regarding the feature stability

(data not shown).

Sample sizes

The sample size for RNA sequencing was selected based on previous leprosy

work with microarrays, aiming at detecting genes with at least a differential fold-change

of two. For RT-qPCR validation, sample size calculation was performed using the per-

gene standardized effect size estimated from the RNA-seq data, aiming at a power of

85% and alpha = 0.03. No samples were discarded after successful data collection

(i.e. outliers). In the end, the sample sizes per group for RT-qPCR were: MB = 13,

PB=11, ODD = 30. All RT-qPCR reactions were conducted in duplicate for each

biological unit (here, a fragment of a skin biopsy derived from an individual).

RT-qPCR and ROC statistical analyses

Normalized RT-qPCR gene expression data were $\log_2$ transformed before use

in data visualization. Additionally, we checked if the Bayesian results remained

consistent using a more common procedure (data not shown). For this, the mean

normalized expression (from N_0) was compared pairwise for the prior stipulated groups

using Welch's t-test implemented in R language, using the predetermined alpha of

0.03. Normality assumption was verified with normal quantile-quantile plots (qqplots,

car v. 3.0-2). In cases where quantile-quantile plots showed huge deviation from

theoretical normal distribution, the Wilcoxon Rank Sum was used to verify results.

Receiver Operating Curve (ROC) analysis was used to determine the accuracy

(measured by the area under the curve, AUC) and its respective best classification

threshold, aiming at maximizing AUC with equal importance for sensitivity and

specificity. Confidence intervals (95%) for AUC were calculated using the Delong non-

parametric method as implemented in pROC v.1.15.3 [124–126].

Data and code reporting

Raw .fastq data are available in EMBL-EBI European Nucleotide Archive (ENA)

database (ERP128243). Raw Salmon counts and normalized batch cleaned

expression matrices are available in EMBL-EBI ArrayExpress, under E-MTAB-10318,

along with experimental and phenotypic metadata. R source code and accompanying

intermediate data used in all analyses in this manuscript are also readily available

through Zenodo, doi.org/10.5281/zenodo.4682010.

Acknowledgements

The authors wish to acknowledge Suelen Justo Moreira (MSc) and Rhana Prata

(PhD) for assistance with skin biopsy RNA isolation. Helen Ferreira (MSc) and José

Augusto for their technical and logistic support. The Gene Expression Core Facility

(GECF) at EPFL, Lausanne, Switzerland, especially Drs. Elisa Cora and Bastien

Mangeat for sequencing assistance. All patients and staff (physicians, nurses and

technicians) from Sousa Araujo Outpatient clinic at FIOCRUZ, Rio de Janeiro, Brazil.

References

1. Britton WJ, Lockwood DN. Leprosy. The Lancet. 2004;363: 1209–1219.

doi:10.1016/S0140-6736(04)15952-7

2.   Ridley DS, Jopling WH. Classification of leprosy according to immunity. A five-

group system. Int J Lepr Mycobact Dis Off Organ Int Lepr Assoc. 1966;34: 255–73.

3.   Scollard DM, Adams LB, Gillis TP, Krahenbuhl JL, Truman W, Williams DL.

The Continuing Challenges of Leprosy The Continuing Challenges of Leprosy. Clin

Microbiol Rev. 2006;19: 338–381. doi:10.1128/CMR.19.2.338

4.   WHO. Guidelines for the Diagnosis, Treatment and Prevention of Leprosy.

Geneva: World Health Organization; 2018 p. 106.

5.   WHO. Global leprosy (Hansen disease) update, 2019: time to step-up

prevention initiatives. Wkly Epidemiol Rec. 2020;95: 417–440.

6.   Nath I, Saini C, Valluri VL. Immunology of leprosy and diagnostic challenges.

Clin Dermatol. 2015;33: 90–98. doi:10.1016/j.clindermatol.2014.07.005

7.   van Hooij A, Tjon Kon Fat EM, Batista da Silva M, Carvalho Bouth R, Cunha

Messias AC, Gobbo AR, et al. Evaluation of Immunodiagnostic Tests for Leprosy in

Brazil, China and Ethiopia. Sci Rep. 2018;8: 1–9. doi:10.1038/s41598-018-36323-1

8.   van Hooij A, van den Eeden S, Richardus R, Tjon Kon Fat E, Wilson L,

Franken KLMC, et al. Application of new host biomarker profiles in quantitative point-

of-care tests facilitates leprosy diagnosis in the field. EBioMedicine. 2019;47: 301–

308. doi:10.1016/j.ebiom.2019.08.009

9.   Manta FS de N, Leal-Calvo T, Moreira SJM, Marques BLC, Ribeiro-Alves M,

Rosa PS, et al. Ultra-sensitive detection of Mycobacterium leprae: DNA extraction

and PCR assays. Poonawala H, editor. PLoS Negl Trop Dis. 2020;14: e0008325.

doi:10.1371/journal.pntd.0008325

10.   Gliddon HD, Herberg JA, Levin M, Kaforou M. Genome-wide host RNA

signatures of infectious diseases: discovery and clinical translation. Immunology.

2018;153: 171–178. doi:10.1111/imm.12841

11.   Ko ER, Yang WE, McClain MT, Woods CW, Ginsburg GS, Tsalik EL. What

was old is new again: Using the host response to diagnose infectious disease.

Expert Rev Mol Diagn. 2015;15: 1143–1158. doi:10.1586/14737159.2015.1059278

12.   Miller RR, Lopansri BK, Burke JP, Levy M, Opal S, Rothman RE, et al.

Validation of a host response assay, SeptiCyte LAB, for discriminating sepsis from

systemic inflammatory response syndrome in the ICU. Am J Respir Crit Care Med.

2018;198: 903–913. doi:10.1164/rccm.201712-2472OC

13.   Van Hooij A, Fat EMTK, Van Den Eeden SJF, Wilson L, Da Silva MB,

Salgado CG, et al. Field-friendly serological tests for determination of M. Leprae-

specific antibodies. Sci Rep. 2017;7: 1–8. doi:10.1038/s41598-017-07803-7

14.   Warsinske H, Vashisht R, Khatri P. Host-response-based gene signatures for

tuberculosis diagnosis: A systematic comparison of 16 signatures. PLoS Med.

2019;16. doi:10.1371/journal.pmed.1002786

15.   Röltgen K, Pluschke G, Spencer JS, Brennan PJ, Avanzi C. The immunology

of other mycobacteria: M. ulcerans, M. leprae. Semin Immunopathol. 2020;42: 333–

353. doi:10.1007/s00281-020-00790-4

16.   Mesko B, Poliska S, Nagy L. Gene expression profiles in peripheral blood for

the diagnosis of autoimmune diseases. Trends Mol Med. 2011;17: 223–233.

doi:10.1016/j.molmed.2010.12.004

17.   Wang B, Chen S, Zheng Q, Gao Z, Chen R, Xuan J, et al. Development and

initial validation of diagnostic gene signatures for systemic lupus erythematosus. Ann

Rheum Dis. 2019. doi:10.1136/annrheumdis-2019-216695

18.   Carlson JJ, Roth JA. The impact of the Oncotype Dx breast cancer assay in

clinical practice: A systematic review and meta-analysis. Breast Cancer Res Treat.

2013;141: 13–22. doi:10.1007/s10549-013-2666-z

19.   Gordon GJ, Jensen RV, Hsiao LL, Gullans SR, Blumenstock JE, Ramaswamy

S, et al. Translation of microarray data into clinically relevant cancer diagnostic tests

using gene expression ratios in lung cancer and mesothelioma. Cancer Res.

2002;62: 4963–4967.

20.   Narrandes S, Xu W. Gene expression detection assay for cancer clinical use.

J Cancer. 2018;9: 2249–2265. doi:10.7150/jca.24744

21.   Clark-Langone KM, Sangli C, Krishnakumar J, Watson D. Translating tumor

biology into personalized treatment planning: analytical performance characteristics

of the Oncotype DX®Colon Cancer Assay. BMC Cancer. 2010;10: 691.

doi:10.1186/1471-2407-10-691

22.   Knezevic D, Goddard AD, Natraj N, Cherbavaz DB, Clark-Langone KM,

Snable J, et al. Analytical validation of the Oncotype DX prostate cancer assay - a

clinical RT-PCR assay optimized for prostate needle biopsies. BMC Genomics.

2013;14: 1–12. doi:10.1186/1471-2164-14-690

23.   Laible M, Schlombs K, Kaiser K, Veltrup E, Herlein S, Lakis S, et al. Technical

validation of an RT-qPCR in vitro diagnostic test system for the determination of

breast cancer molecular subtypes by quantification of ERBB2 , ESR1 , PGR and

MKI67 mRNA levels from formalin- fixed paraffin-embedded breast tumor

specimens. BMC Cancer. 2016; 1–14. doi:10.1186/s12885-016-2476-x

24.   Belone A de FF, Rosa PS, Trombone APF, Fachin LRV, Guidella CC, Ura S,

et al. Genome-wide screening of mRNA expression in leprosy patients. Front Genet.

2015;6: 1–12. doi:10.3389/fgene.2015.00334

25.   Jorge KTOS, Souza RP, Assis MTA, Araújo MG, Locati M, Jesus AMR, et al.

Characterization of MicroRNA Expression Profiles and Identification of Potential

Biomarkers in Leprosy. J Clin Microbiol. 2017;55: 1516–1525.

doi:10.1128/JCM.02408-16

26.   Tió-Coma M, van Hooij A, Bobosha K, van der Ploeg-van Schip JJ, Banu S,

Khadge S, et al. Whole blood RNA signatures in leprosy patients identify reversal

reactions before clinical onset: a prospective, multicenter study. Sci Rep. 2019;9:

17931. doi:10.1038/s41598-019-54213-y

27.   Tió-Coma M, Kiełbasa SM, van den Eeden SJF, Mei H, Roy JC, Wallinga J, et

al. Blood RNA signature RISK4LEP predicts leprosy years before clinical onset.

EBioMedicine. 2021;68: 103379. doi:10.1016/j.ebiom.2021.103379

28.   Montoya DJ, Andrade P, Silva BJA, Teles RMB, Ma F, Bryson B, et al. Dual

RNA-Seq of Human Leprosy Lesions Identifies Bacterial Determinants Linked to

Host Immune Response. Cell Rep. 2019;26: 3574-3585.e3.

doi:10.1016/j.celrep.2019.02.109

29.   Bhatia S, Shenoi SD, Pai K, Srilatha PS. Granuloma multiforme: an

uncommon differential for leprosy. Trop Doct. 2019;49: 55–58.

doi:10.1177/0049475518803191

30.   Kundakci N, Erdem C. Leprosy: A great imitator. Clin Dermatol. 2019;37:

200–212. doi:10.1016/j.clindermatol.2019.01.002

31.   Zhu TH, Kamangar F, Silverstein M, Fung MA. Borderline Tuberculoid

Leprosy Masquerading as Granuloma Annulare: A Clinical and Histological Pitfall.

Am J Dermatopathol. 2017;39: 296–299. doi:10.1097/DAD.0000000000000698

32.   Blischak JD, Tailleux L, Mitrano A, Barreiro LB, Gilad Y. Mycobacterial

infection induces a specific human innate immune response. Sci Rep. 2015;5: 1–16.

doi:10.1038/srep16882

33.   Modlin RL. Th1-Th2 paradigm: insights from leprosy. J Invest Dermatol.

1994;102: 828–832. doi:10.1111/1523-1747.ep12381958

34.   Yamamura M, Uyemura K, Deans RJ, Weinberg K, Rea TH, Bloom BR, et al.

Defining protective responses to pathogens: Cytokine profiles in leprosy lesions.

Science. 1991;254: 277–279. doi:10.1126/science.1925582

35.   Leal-Calvo T, Moraes MO. Reanalysis and integration of public microarray

datasets reveals novel host genes modulated in leprosy. Mol Genet Genomics.

2020;295: 1355–1368. doi:10.1007/s00438-020-01705-6

36.   Judson MA, Marchell RM, Mascelli M, Piantone A, Barnathan ES, Petty KJ, et

al. Molecular profiling and gene expression analysis in cutaneous sarcoidosis: the

role of interleukin-12, interleukin-23, and the T-helper 17 pathway. J Am Acad

Dermatol. 2012;66: 901–910, 910.e1–2. doi:10.1016/j.jaad.2011.06.017

37.   Linke M, Pham HTT, Katholnig K, Schnöller T, Miller A, Demel F, et al.

Chronic signaling via the metabolic checkpoint kinase mTORC1 induces

macrophage granuloma formation and marks sarcoidosis progression. Nat Immunol.

2017;18: 293–302. doi:10.1038/ni.3655

38. Masaki T, Qu J, Cholewa-Waclaw J, Burr K, Raaum R, Rambukkana A.

Reprogramming adult Schwann cells to stem cell-like cells by leprosy bacilli

promotes dissemination of infection. Cell. 2013;152: 51–67.

doi:10.1016/j.cell.2012.12.014

39. Brabletz T, Kalluri R, Nieto MA, Weinberg RA. EMT in cancer. Nat Rev

Cancer. 2018;18: 128–134. doi:10.1038/nrc.2017.118

40. Pastushenko I, Blanpain C. EMT Transition States during Tumor Progression

and Metastasis. Trends Cell Biol. 2019;29: 212–226. doi:10.1016/j.tcb.2018.12.001

41. Khazai Z, Van Brakel W, Essink D, Gillis T, Kasang C, Kuipers P, et al.

Reviewing Research Priorities of the Leprosy Research Initiative (LRI): a

stakeholder's consultation. Lepr Rev. 2019;90: 3–30. doi:10.47276/lr.90.1.3

42. Chen W. IDO: more than an enzyme. Nat Immunol. 2011;12: 809–811.

doi:10.1038/ni.2088

43. Greco FA, Coletti A, Camaioni E, Carotti A, Marinozzi M, Gioiello A, et al. The

Janus-faced nature of IDO1 in infectious diseases: challenges and therapeutic

opportunities. Future Med Chem. 2016;8: 39–54. doi:10.4155/fmc.15.165

44. Melé M, Ferreira PG, Reverter F, DeLuca DS, Monlong J, Sammeth M, et al.

The human transcriptome across tissues and individuals. Science. 2015;348: 660–

665. doi:10.1126/science.aaa0355

45.   Yamazaki F, Kuroiwa T, Takikawa O, Kido R. Human indolylamine 2,3-

dioxygenase. Its tissue distribution, and characterization of the placental enzyme.

Biochem J. 1985;230: 635–638. doi:10.1042/bj2300635

46.   Hughes TK, Wadsworth MH, Gierahn TM, Do T, Weiss D, Andrade PR, et al.

Second-Strand Synthesis-Based Massively Parallel scRNA-Seq Reveals Cellular

States and Molecular Features of Human Inflammatory Skin Pathologies. Immunity.

2020;53: 878-894.e7. doi:10.1016/j.immuni.2020.09.015

47.   Gautam US, Foreman TW, Bucsan AN, Veatch AV, Alvarez X, Adekambi T, et

al. In vivo inhibition of tryptophan catabolism reorganizes the tuberculoma and

augments immune-mediated control of Mycobacterium tuberculosis. Proc Natl Acad

Sci U S A. 2018;115: E62–E71. doi:10.1073/pnas.1711373114

48.   Yeung AWS, Terentis AC, King NJC, Thomas SR. Role of indoleamine 2,3-

dioxygenase in health and disease. Clin Sci. 2015;129: 601–672.

doi:10.1042/CS20140392

49.   Denys A, Allain F. The emerging roles of heparan sulfate 3-O-

sulfotransferases in cancer. Front Oncol. 2019;9. doi:10.3389/fonc.2019.00507

50.   Yamauchi PS, Bleharski JR, Uyemura K, Kim J, Sieling PA, Miller A, et al. A

Role for CD40-CD40 Ligand Interactions in the Generation of Type 1 Cytokine

Responses in Human Leprosy. J Immunol. 2000;165: 1506–1512.

doi:10.4049/jimmunol.165.3.1506

51.   Heim A. Highly sensitive detection of gene expression of an intronless gene:

amplification of mRNA, but not genomic DNA by nucleic acid sequence based

amplification (NASBA). Nucleic Acids Res. 1998;26: 2250–2251.

doi:10.1093/nar/26.9.2250

52.   Patterson SS, Casper ET, Garcia-Rubio L, Smith MC, Paul JH. Increased

precision of microbial RNA quantification using NASBA with an internal control. J

Microbiol Methods. 2005;60: 343–352. doi:10.1016/j.mimet.2004.10.011

53.   Ganguli A, Ornob A, Spegazzini N, Liu Y, Damhorst G, Ghonge T, et al.

Pixelated spatial gene expression analysis from tissue. Nat Commun. 2018;9.

doi:10.1038/s41467-017-02623-9

54.   Pandey M, Singh D, Onteru SK. Reverse transcription loop-mediated

isothermal amplification (RT-LAMP), a light for mammalian transcript analysis in low-

input laboratories. J Cell Biochem. 2018;119: 4334–4338. doi:10.1002/jcb.26624

55.   Broughton JP, Deng X, Yu G, Fasching CL, Servellita V, Singh J, et al.

CRISPR–Cas12-based detection of SARS-CoV-2. Nat Biotechnol. 2020;38: 870–

874. doi:10.1038/s41587-020-0513-4

56.   Barbieri RR, Manta FSN, Moreira SJM, Sales AM, Nery JAC, Nascimento

LPR, et al. Quantitative polymerase chain reaction in paucibacillary leprosy

diagnosis: A follow-up study. PLoS Negl Trop Dis. 2019;13: e0007147.

doi:10.1371/journal.pntd.0007147

57.   Strassner JP, Rashighi M, Ahmed Refat M, Richmond JM, Harris JE. Suction

blistering the lesional skin of vitiligo patients reveals useful biomarkers of disease

activity. J Am Acad Dermatol. 2017;76: 847-855.e5. doi:10.1016/j.jaad.2016.12.021

58.   Elamin AA, Stehr M, Singh M. Lipid Droplets and Mycobacterium leprae

Infection. J Pathog. 2012;10. doi:10.1155/2012/361374

59.   Lobato LS, Rosa PS, Ferreira J da S, Neumann A da S, da Silva MG, do

Nascimento DC, et al. Statins increase rifampin mycobactericidal effect. Antimicrob

Agents Chemother. 2014;58: 5766–74. doi:10.1128/AAC.01826-13

60.   Wang D, Zhang D-F, Li G-D, Bi R, Fan Y, Wu Y, et al. A pleiotropic effect of

the APOE gene: association of APOE polymorphisms with multibacillary leprosy in

Han Chinese from Southwest China. Br J Dermatol. 2018;178: 931–939.

doi:10.1111/bjd.16020

61.   Fabel A, Giovanna Brunasso AM, Schettini AP, Cota C, Puntoni M, Nunzi E,

et al. Pathogenesis of Leprosy. Am J Dermatopathol. 2019;41: 422–427.

doi:10.1097/DAD.0000000000001310

62.   Iyer AM, Mohanty KK, van Egmond D, Katoch K, Faber WR, Das PK, et al.

Leprosy-specific B-cells within cellular infiltrates in active leprosy lesions. Hum

Pathol. 2007;38: 1065–1073. doi:10.1016/j.humpath.2006.12.017

63.   Medeiros RCA, Girardi K do C de V, Cardoso FKL, Mietto B de S, Pinto TG

de T, Gomez LS, et al. Subversion of Schwann Cell Glucose Metabolism by

Mycobacterium leprae. J Biol Chem. 2016;291: 21375–21387.

doi:10.1074/jbc.M116.725283

64.   de Toledo-Pinto TG, Ferreira ABR, Ribeiro-Alves M, Rodrigues LS, Batista-

Silva LR, Silva BJ de A, et al. STING-Dependent 2′-5′ Oligoadenylate Synthetase–

Like Production Is Required for Intracellular Mycobacterium leprae Survival. J Infect

Dis. 2016;214: 311–320. doi:10.1093/infdis/jiw144

65.   Hess S, Rambukkana A. Bacterial-induced cell reprogramming to stem cell-

like cells: new premise in host–pathogen interactions. Curr Opin Microbiol. 2015;23:

179–188. doi:10.1016/j.mib.2014.11.021

66.   Vandewalle C, Comijn J, De Craene B, Vermassen P, Bruyneel E, Andersen

H, et al. SIP1/ZEB2 induces EMT by repressing genes of different epithelial cell-cell

junctions. Nucleic Acids Res. 2005;33: 6566–6578. doi:10.1093/nar/gki965

67.   DaSilva-Arnold SC, Kuo CY, Davra V, Remache Y, Kim PCW, Fisher JP, et

al. ZEB2, a master regulator of the epithelial-mesenchymal transition, mediates

trophoblast differentiation. Mol Hum Reprod. 2018;25: 61–75.

doi:10.1093/molehr/gay053

68.   Jiang Y, Zhou J, Hou D, Luo P, Gao H, Ma Y, et al. Prosaposin is a biomarker

of mesenchymal glioblastoma and regulates mesenchymal transition through the

TGF-β1/Smad signaling pathway. J Pathol. 2019;249: 26–38. doi:10.1002/path.5278

69.   Frugtniet BA, Martin TA, Zhang L, Jiang WG. Neural Wiskott-Aldrich

syndrome protein (nWASP) is implicated in human lung cancer invasion. BMC

Cancer. 2017;17. doi:10.1186/s12885-017-3219-3

70.   Bendris N, Arsic N, Lemmers B, Blanchard JM. Cyclin A2, Rho GTPases and

EMT. Small GTPases. 2012;3: 225–228. doi:10.4161/sgtp.20791

71.   Bhowmick NA, Ghiassi M, Bakin A, Aakre M, Lundquist CA, Engel ME, et al.

Transforming growth factor-β1 mediates epithelial to mesenchymal

transdifferentiation through a RhoA-dependent mechanism. Mol Biol Cell. 2001;12:

27–36. doi:10.1091/mbc.12.1.27

72.   Salvi A, Thanabalu T. WIP promotes in-vitro invasion ability, anchorage

independent growth and EMT progression of A549 lung adenocarcinoma cells by

regulating RhoA levels. Biochem Biophys Res Commun. 2017;482: 1353–1359.

doi:10.1016/j.bbrc.2016.12.040

73.   Wang Q, Yang X, Xu Y, Shen Z, Cheng H, Cheng F, et al. RhoA/Rho-kinase

triggers epithelial-mesenchymal transition in mesothelial cells and contributes to the

pathogenesis of dialysis-related peritoneal fibrosis. Oncotarget. 2018;9: 14397–

14412. doi:10.18632/oncotarget.24208

74.   Wang J, Chen L, Li Y, Guan XY. Overexpression of cathepsin Z contributes to

tumor metastasis by inducing epithelial-mesenchymal transition in hepatocellular

carcinoma. PLoS ONE. 2011;6. doi:10.1371/journal.pone.0024967

75.   Lin CY, Tsai PH, Kandaswami CC, Lee PP, Huang CJ, Hwang JJ, et al.

Matrix metalloproteinase-9 cooperates with transcription factor Snail to induce

epithelial-mesenchymal transition. Cancer Sci. 2011;102: 815–827.

doi:10.1111/j.1349-7006.2011.01861.x

76.   Peinado H, del Carmen Iglesias-de la Cruz M, Olmeda D, Csiszar K, Fong

KSK, Vega S, et al. A molecular role for lysyl oxidase-like 2 enzyme in Snail

regulation and tumor progression. EMBO J. 2005;24: 3446–3458.

doi:10.1038/sj.emboj.7600781

77.   Tam SY, Wu VWC, Law HKW. Hypoxia-Induced Epithelial-Mesenchymal

Transition in Cancers: HIF-1α and Beyond. Front Oncol. 2020;10.

doi:10.3389/fonc.2020.00486

78. Zhu Y, Tan J, Xie H, Wang J, Meng X, Wang R. HIF-1α regulates EMT via the

Snail and β-catenin pathways in paraquat poisoning-induced early pulmonary

fibrosis. J Cell Mol Med. 2016;20: 688–697. doi:10.1111/jcmm.12769

79. Lyrio ECD, Campos-Souza IC, Corrêa LCD, Lechuga GC, Verícimo M, Castro

HC, et al. Interaction of Mycobacterium leprae with the HaCaT human keratinocyte

cell line: new frontiers in the cellular immunology of leprosy. Exp Dermatol. 2015;24:

536–542. doi:10.1111/exd.12714

80. Okada S, Komura J, Nishiura M. Mycobacterium leprae found in epidermal

cells by electron microscopy. IntJLeprOther MycobactDis. 1978;46: 30–34.

81. Pivarcsi A, Kemény L, Dobozy A. Innate Immune Functions of the

Keratinocytes. Acta Microbiol Immunol Hung. 2004;51: 303–310.

doi:10.1556/AMicr.51.2004.3.8

82. Pivarcsi A, Nagy I, Lajos K. Innate Immunity in the Skin: How Keratinocytes

Fight Against Pathogens. Curr Immunol Rev. 2005;1: 29–43.

doi:10.2174/1573395052952941

83. Damsky W, Thakral D, McGeary MK, Leventhal J, Galan A, King B. Janus

kinase inhibition induces disease remission in cutaneous sarcoidosis and granuloma

annulare. J Am Acad Dermatol. 2020;82: 612–621. doi:10.1016/j.jaad.2019.05.098

84. Flynn JL, Chan J, Lin PL. Macrophages and control of granulomatous

inflammation in tuberculosis. Mucosal Immunol. 2011;4: 271–278.

doi:10.1038/mi.2011.14

85. Locke LW, Crouser ED, White P, Julian MW, Caceres EG, Papp AC, et al. IL-

13–regulated Macrophage Polarization during Granuloma Formation in an In Vitro

Human Sarcoidosis Model. Am J Respir Cell Mol Biol. 2019;60: 84–95.

doi:10.1165/rcmb.2018-0053OC

86.   Manta FSN, Barbieri RR, Moreira SJM, Santos PTS, Nery JAC, Duppre NC,

et al. Quantitative PCR for leprosy diagnosis and monitoring in household contacts:

A follow-up study, 2011–2018. Sci Rep. 2019;9. doi:10.1038/s41598-019-52640-5

87.   Brabaham Bioinformatics. FastQC: A Quality Control Tool for High

Throughput Sequence Data [Online]. 2015. Available:

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

88.   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast

and bias-aware quantification of transcript expression. Nat Methods. 2017;14: 417–

419. doi:10.1038/nmeth.4197

89.   R Core Team. R: A language and environment for statistical computing.

Vienna, Austria; 2017. Available: https://www.r-project.org/

90.   Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq:

transcript-level estimates improve gene-level inferences. F1000Research. 2016;4:

1521. doi:10.12688/f1000research.7563.2

91.   Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, et al.

BioMart and Bioconductor: a powerful link between biological databases and

microarray data analysis. Bioinformatics. 2005;21: 3439–3440.

doi:10.1093/bioinformatics/bti525

92.   Anders S, Huber W. Differential expression analysis for sequence count data.

Genome Biol. 2010;11: R106. doi:10.1186/gb-2010-11-10-r106

93.   Love MI, Huber W, Anders S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550.

doi:10.1186/s13059-014-0550-8

94.   Zhu A, Ibrahim JG, Love MI. Heavy-Tailed prior distributions for sequence

count data: Removing the noise and preserving large differences. Bioinformatics.

2019;35: 2084–2092. doi:10.1093/bioinformatics/bty895

95.   Gagnon-Bartsch JA, Speed TP. Using control genes to correct for unwanted

variation in microarray data. Biostatistics. 2012;13: 539–552.

doi:10.1093/biostatistics/kxr034

96.   Risso D, Ngai J, Speed TP, Dudoit S. Normalization of RNA-seq data using

factor analysis of control genes or samples. Nat Biotechnol. 2014;32: 896–902.

doi:10.1038/nbt.2931

97.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical

and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society.

Series B (Methodological). WileyRoyal Statistical Society; 1995.

doi:10.2307/2346101

98.   Phipson B, Lee S, Majewski IJ, Alexander WS, Smyth GK. Robust

hyperparameter estimation protects against hypervariable genes and improves

power to detect differential expression. Ann Appl Stat. 2016;10: 946–963.

doi:10.1214/16-AOAS920

99.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers

differential expression analyses for RNA-sequencing and microarray studies. Nucleic

Acids Res. 2015;43: e47. doi:10.1093/nar/gkv007

100.   Smyth GK. Linear Models and Empirical Bayes Methods for Assessing

Differential Expression in Microarray Experiments Linear Models and Empirical

Bayes Methods for Assessing Differential Expression in Microarray Experiments.

Stat Appl Genet Mol Biol. 2004;3: 1–26. doi:10.2202/1544-6115.1027

101.   Wickham H. ggplot2-Elegant Graphics for Data Analysis. 1st ed. New York,

NY: Springer New York; 2009. doi:10.1007/978-0-387-98141-3

102.   Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and

correlations in multidimensional genomic data. Bioinformatics. 2016;32: 2847–2849.

doi:10.1093/bioinformatics/btw313

103.   Kolde R. pheatmap: Pretty Heatmaps. 2015. Available: https://cran.r-

project.org/package=pheatmap

104.   Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R Package for

Comparing Biological Themes Among Gene Clusters. OMICS J Integr Biol. 2012;16:

284–287. doi:10.1089/omi.2011.0118

105.   Carlson M. org.Hs.eg.db: Genome wide annotation for Human. 2019.

Available: 10.18129/B9.bioc.org.Hs.eg.db

106.   Koressaar T, Remm M. Enhancements and modifications of primer design

program Primer3. Bioinformatics. 2007;23: 1289–1291.

doi:10.1093/bioinformatics/btm091

107.   Kõressaar T, Lepamets M, Kaplinski L, Raime K, Andreson R, Remm M.

Primer3_masker: integrating masking of template sequence with primer design

software. Bioinformatics. 2018;34: 1937–1938. doi:10.1093/bioinformatics/bty036

108.   Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al.

Primer3—new capabilities and interfaces. Nucleic Acids Res. 2012;40: e115–e115.

doi:10.1093/nar/gks596

109.   Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. Primer-

BLAST: a tool to design target-specific primers for polymerase chain reaction. BMC

Bioinformatics. 2012;13: 134. doi:10.1186/1471-2105-13-134

110.   Qu W, Shen Z, Zhao D, Yang Y, Zhang C. MFEprimer: Multiple factor

evaluation of the specificity of PCR primers. Bioinformatics. 2009;25: 276–278.

doi:10.1093/bioinformatics/btn614

111.   Wang K, Li H, Xu Y, Shao Q, Yi J, Wang R, et al. MFEprimer-3.0: Quality

control for PCR primers. Nucleic Acids Res. 2019;47: W610–W613.

doi:10.1093/nar/gkz351

112.   Ramakers C, Ruijter JM, Lekanne Deprez RH, Moorman AFM. Assumption-

free analysis of quantitative real-time polymerase chain reaction (PCR) data.

Neurosci Lett. 2003;339: 62–66. doi:10.1016/S0304-3940(02)01423-4

113.   Ruijter JM, Ramakers C, Hoogaars WMH, Karlen Y, Bakker O, Van den hoff

MJB, et al. Amplification efficiency: Linking baseline and bias in the analysis of

quantitative PCR data. Nucleic Acids Res. 2009;37. doi:10.1093/nar/gkp045

114.   Vandesompele J, De Preter K, Pattyn ilip, Poppe B, Van Roy N, De Paepe A,

et al. Accurate normalization of real-time quantitative RT-PCR data by geometric

averaging of multiple internal control genes. Genome Biol. 2002;3: 34–1.

doi:10.1186/gb-2002-3-7-research0034

115.   Matz MV, Wright RM, Scott JG. No control genes required: Bayesian analysis

of qRT-PCR data. PloS One. 2013;8: 1–12. doi:10.1371/journal.pone.0071448

116.   Steibel JP, Poletto R, Coussens PM, Rosa GJM. A powerful and flexible linear

mixed model framework for the analysis of relative quantification RT-PCR data.

Genomics. 2009;94: 146–152. doi:10.1016/j.ygeno.2009.04.008

117.   Fieller EC, Hartley HO, Pearson ES. TESTS FOR RANK CORRELATION

COEFFICIENTS I. Biometrika. 1957;44: 470–481. doi:10.1093/biomet/44.3-4.470

118.   Makowski D, Ben-Shachar MS, Patil I, Lüdecke D. Methods and Algorithms

for Correlation Analysis in R. J Open Source Softw. 2020;5: 2306.

doi:10.21105/joss.02306

119.   Friedman J, Hastie T, Tibshirani R. Regularization paths for generalized linear

models via coordinate descent. J Stat Softw. 2010;33: 1–22.

doi:10.18637/jss.v033.i01

120.   Simon N, Friedman JH, Hastie T, Tibshirani R. Regularization Paths for Cox's

Proportional Hazards Model via Coordinate Descent. J Stat Softw. 2011;39: 1–13.

doi:10.18637/jss.v039.i05

121.   Tibshirani R. Regression Shrinkage and Selection via the Lasso. J R Stat Soc

Ser B Methodol. 1996;58: 267–288.

122.   Hastie T, Tibshirani R, Wainwright M. Statistical Learning with Sparsity. 1st

ed. Chapman and Hall/CRC; 2015.

123.   Davison AC, Hinley DV. Bootstrap Methods and Their Application. Cambrige

University Press; 1997. Available: http://statwww.epfl.ch/davison/BMA/

124. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the Areas under

Two or More Correlated Receiver Operating Characteristic Curves: A Nonparametric

Approach. Biometrics. 1988;44: 837. doi:10.2307/2531595

125. Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, et al. pROC:

An open-source package for R and S+ to analyze and compare ROC curves. BMC

Bioinformatics. 2011;12: 77. doi:10.1186/1471-2105-12-77

126. Sun X, Xu W. Fast implementation of DeLong's algorithm for comparing the

areas under correlated receiver operating characteristic curves. IEEE Signal Process

Lett. 2014;21: 1389–1393. doi:10.1109/LSP.2014.2337313

Supporting Information

1051

S1 Appendix. Linking expression profiles to mycobacteria species.

S1 Fig. Gene expression in MB and PB groups from test and training datasets.

Normalized $\log_2$ expression values per group from (A) this study RNA-seq dataset or

(B) Belone *et al.* (GSE74481) [24]. The genes shown were selected in 25%–50% of

the LASSO models (Fig 4B) according to the bootstrap. MB, multibacillary leprosy; PB,

paucibacillary leprosy; TT, tuberculoid leprosy; BT, borderline-tuberculoid; BB,

borderline-borderline; BL, borderline-lepromatous; LL, lepromatous. Each point

represents an independent skin biopsy from a patient. Y-axis values are not

comparable between panels A and B.

S2 Fig. Strongest correlations between the average expression of genes

associated with keratinocyte/cornification against dedifferentiation-related

genes using Montoya *et al.* RNA-seq dataset [28]**.** Scatter plots of scores (average

normalized $\log_2$ expression) calculated from genes with previously documented down-

regulation in leprosy skin lesions against dedifferentiation-related genes with Montoya

et al. RNA-seq dataset (GSE125943) [28]. Lines were drawn based on intercept and

beta estimates from robust linear regression for all samples (black) or separately for

TL (tuberculoid leprosy, blue), and LL (lepromatous leprosy, red). X-axis shows $\log_2$

normalized expression values. Spearman's rho are shown along with nominal 95%

confidence intervals inside the plots. Most genes shown have FDR < 0.1 and rho ≤ -

0.6. Related to figure 6.

S3 Fig. Strongest correlations between modulated genes from

keratinocyte/cornification and dedifferentiation-related genes using Belone et

al. microarray dataset (GSE74481) [24]**.** Heat plot with Spearman's rho correlation

coefficient of the strongest correlations from all ontologies screened after multiple

testing adjustment (BH-FDR). Most genes shown have FDR ≤ 0.0001 and rho ≤ -0.7.

Bottom colored rectangles indicate which category the gene was present (some genes

co-occur). Related to figure 6.

S1 Table. Demographic and clinical metadata from human participants.

S2 Table. Genes differentially expressed from leprosy vs. non-leprosy with

|$\log_2$FC| ≥ 1 and FDR ≤ 0.01.

S3 Table. Over-representation analysis (ORA) for leprosy vs. non-leprosy (up-

regulated) differentially expressed genes.

S4 Table. ROC analysis from RNA-seq dataset using leprosy vs. non-leprosy

samples.

S5 Table. Posterior log$_2$FC estimates, 95% credible intervals and MCMC P-
values from PB-OD and MB-OD comparisons.

S6 Table. ROC analysis results using RT-qPCR with the validation dataset
(Related to Fig 3). 95% confidence intervals are shown, except for AUCs of 1.0. The
table is sorted from highest to lowest AUC.

S7 Table. Log$_2$FC estimates, confidence intervals, and Dunnet *P*-values from
distinct mycobacterial stimuli in human macrophages *in vitro*.

S8 Table. Genes differentially expressed from multibacillary paucibacillary
leprosy with |log$_2$FC| ≥ 1 and FDR ≤ 0.01.

S9 Table. Over-representation analysis (ORA) for MB vs. PB (up-regulated)
differentially expressed genes.

S10 Table. Over-representation analysis (ORA) for MB vs. PB (down-regulated)
differentially expressed genes.

S11 Table. Oligonucleotide sequences.