A model of photosynthetic CO_2 assimilation in C_3 leaves accounting for respiration and

energy recycling by the plastidial oxidative pentose phosphate pathway

Thomas Wieloch[a,*], Angela Augusti[b], Jürgen Schleucher[c]

Author affiliations

[a] Department of Forest Genetics and Plant Physiology, Swedish University of Agricultural

Sciences, Umeå Plant Science Centre, 90183 Umeå, Sweden

[b] Research Institute on Terrestrial Ecosystems, National Research Council, 05010 Porano (TR),

Italy.

[c] Department of Medical Biochemistry and Biophysics, Umeå University, 90187 Umeå,

Sweden

* To whom correspondence may be addressed.

email: thomas.wieloch@slu.se

phone: +46706460369

Twitter: https://twitter.com/WielochThomas

ResearchGate: https://www.researchgate.net/profile/Thomas-Wieloch

GoogleScholar: https://scholar.google.de/citations?user=g0ZB7JMAAAAJ&hl=de&oi=sra

Main body of the text: 6548 words

Summary: 199 words

Introduction: 677 words

Theory: 1089 words

Materials and Methods: 684 words

Results: 1529 words

Discussion: 2569 words

7 Figures (Figs. 1, 4, 5 and 7 in colour), 4 Tables, Supporting Information Notes S1

Author	Email address	ORCID
Thomas Wieloch	thomas.wieloch@umu.se	0000-0001-9162-2291
Angela Augusti	angela.augusti@cnr.it	0000-0002-9591-693X
Jürgen Schleucher	jurgen.schleucher@umu.se	0000-0002-4815-3466

Summary

- Recently, we reported estimates of anaplerotic carbon flux through the oxidative pentose

phosphate pathway (OPPP) in chloroplasts into the Calvin-Benson cycle. These estimates were

based on intramolecular hydrogen isotope analysis of sunflower leaf starch. However, the

isotope method is believed to underestimate actual flux at low atmospheric CO_2 concentration

(C_a).

- Since the OPPP releases CO_2 and reduces $NADP^+$, it can be expected to affect leaf gas

exchange under both rubisco- and RuBP-regeneration-limited conditions. Therefore, we

expanded Farquhar-von Caemmerer-Berry models to account for OPPP metabolism. Based on

model parameterisation with values from the literature, we estimated OPPP-related effects on

leaf carbon and energy metabolism in the sunflowers analysed previously.

- We found that flux through the plastidial OPPP increases both above and below $C_a \approx 450$ ppm

(the condition the plants were acclimated to). This is qualitatively consistent with our previous

isotope-based estimates, yet gas-exchange-based estimates are larger at low C_a.

- We discuss our results in relation to regulatory properties of the plastidial and cytosolic OPPP,

the proposed variability of CO_2 mesophyll conductance, and the contribution of day respiration

to the A/C_i curve drop at high C_a. Furthermore, we critically examine the models and

parameterisation and derive recommendations for follow-up studies.

Keywords: Calvin-Benson cycle; CO_2 mesophyll conductance; oxidative pentose phosphate

pathway; glucose-6-phosphate shunt; Farquhar-von Caemmerer-Berry photosynthesis model;

net CO_2 assimilation; day respiration; ATP-to-NADPH ratio

Introduction

Anaplerotic carbon flux into the Calvin-Benson cycle (CBC; for a list of abbreviations and symbols see Table 1) can occur by two metabolic pathways (Fig. 1). The first pathway (here denoted as cytosolic pathway) was proposed by Eicks *et al.* (2002) on discovering the xylulose 5-phosphate/phosphate translocator in *Arabidopsis thaliana*. This translocator enables exchange of pentose phosphates, and inorganic phosphate between chloroplasts and the cytosol. Screening *Arabidopsis* genome databases, Eicks *et al.* (2002) found genes encoding for the cytosolic oxidative branch of the pentose phosphate pathway (OPPP) as well as cytosolic ribulose-5-phosphate epimerase and ribose-5-phosphate isomerase. Genes encoding for cytosolic transketolase and transaldolase required to process pentose phosphates were missing. Because of the apparent lack of fate for pentose phosphates in the cytosol, Eicks *et al.* (2002) proposed their translocation into chloroplasts and injection into the CBC. Recently, studies on *Camelina sativa* leaf metabolism reported that this pathway carries high flux ($\approx 5\%$ relative to net CO_2 assimilation, A) and supplies significant amounts of cytosolic NADPH ($\approx 10\%$ relative to A) at an atmospheric CO_2 concentration (C_a) of ≈ 400 ppm (Xu *et al.*, 2022; Wieloch & Sharkey, 2022).

The second pathway (here denoted as plastidial pathway) was proposed by Sharkey and Weise (2016). It involves glucose 6-phosphate (G6P) to ribulose 5-phosphate (Ru5P) conversion by the plastidial OPPP and injection of the latter into the CBC. For *Pinus nigra*, we reported evidence consistent with flux through this pathway under drought (Wieloch *et al.*, 2018, 2022b). In sunflower, significant flux occurs when plants raised at $C_a \approx 450$ ppm are moved into low or high C_a ($\approx 180, 280, 700, 1500$ ppm) whereas flux at $C_a \approx 450$ ppm was not significantly greater than zero (Wieloch *et al.*, 2022a; Wieloch, 2022). By contrast, Xu *et al.* (2021) reported, in *Camelina sativa*, significant flux occurs at $C_a \approx 400$ ppm corresponding to the conditions the plants were raised in. However, several assumptions of this study were criticised (Wieloch, 2021) and, after implementing amendments, the result was not confirmed (Xu *et al.*, 2022).

In sunflower, we estimated flux through the plastidial anaplerotic pathway based on a hydrogen isotope signal in leaf starch (the term isotope signal denotes systematic variability in relative isotope abundance) (Wieloch *et al.*, 2022a; Wieloch, 2022). Assuming 50% of the net assimilated carbon becomes leaf starch (Sharkey *et al.*, 1985), anaplerotic flux proceeds at $>$ 7%, $>$ 5%, 0%, $\approx$ 2%, and $\approx$ 5% relative to A at $C_a \approx 180, 280, 450, 700,$ and 1500 ppm, respectively (Wieloch, 2022). However, this method is thought to strongly underestimate flux

at 180 and 280 ppm because much of the hydrogen isotope signal can be expected to not arrive in starch under low-C_a conditions (Wieloch *et al.*, 2022a). Therefore, the present study aims first and foremost at developing an alternative method for estimating flux through the plastidial anaplerotic pathway and secondly at exploring its capabilities.

For each molecule Ru5P synthesised anaplerotically, one molecule CO_2 is released from metabolism while two molecules $NADP^+$ are reduced (Fig. 1). Thus, we hypothesise anaplerotic flux affects leaf gas exchange and can be traced by gas exchange modelling. To test this, we expand Farquhar-von Caemmerer-Berry (FvCB) photosynthesis models by terms accounting for anaplerotic respiration and energy recycling. Using gas exchange data of sunflower, we estimate anaplerotic flux rates and associated effects on leaf carbon and energy metabolism. These data are unique in that they were collected during the same experiment and from the same plants which previously provided isotope evidence for flux through the plastidial anaplerotic pathway (Wieloch *et al.*, 2022a; Wieloch, 2022). The results are discussed (*inter alia*) in relation to (i) previously reported flux estimates, (ii) regulatory properties of the plastidial and cytosolic anaplerotic pathway, (iii) the proposed variability of CO_2 mesophyll conductance (g_m) with C_a and during photosynthetic induction, and (iv) the contribution of day respiration to the often-seen A/C_i curve drop at high C_a (C_i, CO_2 concentration in intercellular air spaces). We close with a paragraph about weaknesses of the developed models and the chosen parameterisation and future research directions.

Theory

Photosynthesis models accounting for anaplerotic flux into the Calvin-Benson cycle

In FvCB photosynthesis models (Farquhar *et al.*, 1980), net CO_2 assimilation is given as

$$A = V_c - 0.5V_o - R_d = V_c(1 - 0.5\,V_o/V_c) - R_d = V_c(1 - 0.5\Phi) - R_d \qquad \text{Eqn 1}$$

where R_d denotes day respiration, V_o and V_c denote the oxygenation and carboxylation rates of ribulose 1,5-bisphosphate (RuBP), respectively, and Φ denotes the V_o/V_c ratio. Making day respiration by the plastidial anaplerotic pathway ($R_{a,p}$) explicit yields

$$A = V_c(1 - 0.5\Phi) - R_{a,p} - R_x \qquad \text{Eqn 2}$$

where R_x denotes day respiration by pathways other than the plastidial anaplerotic pathway. Please note that we also use $R_{a,p}$ to denote flux through the plastidial anaplerotic pathway.

Different biochemical processes may exert control over A. First, rubisco-limited CO_2 assimilation (A_c) denotes assimilation limited by CO_2 supply. Under these conditions, rubisco carboxylation rate is given as

$$V_c = \frac{V_{cmax} C_c}{C_c + K_c(1 + O_c/K_o)} \qquad \text{Eqn 3}$$

where V_{cmax} denotes the CO_2-saturated maximum rate of rubisco carboxylation, C_c and O_c denote CO_2 and O_2 partial pressures at the active site of rubisco, respectively, and K_c and K_o denote the Michaelis-Menten constants of rubisco carboxylation and oxygenation, respectively (Farquhar *et al.*, 1980). According to Farquhar and von Caemmerer (1982), Φ is related to the CO_2-compensation point in the absence of day respiration (Γ_*) as

$$\Phi = \frac{2\Gamma_*}{C_c} \qquad \text{Eqn 4}$$

Using equations 3 and 4 to respectively remove V_c and Φ from equation 2 yields

$$A_c = \frac{V_{cmax}(C_c - \Gamma_*)}{C_c + K_c(1 + O_c/K_o)} - R_{a,p} - R_x \qquad \text{Eqn 5}$$

Second, RuBP-regeneration-limited CO_2 assimilation (A_j) presumably results from a shortage of incoming electrons supplied to the CBC and photorespiration as NADPH and ATP (Farquhar *et al.*, 1980). Here, we assume NADPH rather than ATP supply is limiting. Electrons for $NADP^+$ reduction come from the linear electron transport pathway (1 mol NADPH requires 2 mol electrons). Additionally, per 1 mol CO_2 respired by anaplerotic pathways, 2 mol NADPH (4 mol electrons) are recycled (Fig. 1). However, only NADPH from the plastidial pathway can be recycled into the CBC and photorespiration. NADPH from the cytosolic pathway cannot enter chloroplasts (Wieloch & Sharkey, 2022). Thus, overall electron supply is given as

$$e^- \text{supply rate} = J + J_{a,p} = J + 4R_{a,p} \qquad \text{Eqn 6}$$

149 where J denotes the rate of linear electron transport, and $J_{a,p}$ denotes the rate of electron

150 recycling by the plastidial anaplerotic pathway. According to Farquhar *et al.* (1980), the rate of

151 electron consumption by the CBC and photorespiration is given as

$$e^- consumption\ rate = V_c(4 + 4\Phi) \qquad \text{Eqn 7}$$

154 Balancing electron supply (Eqn. 6) with electron consumption (Eqn. 7) yields

$$J + 4R_{a,p} = V_c(4 + 4\Phi) \qquad \text{Eqn 8}$$

157 Solving equation 8 for V_c yields

$$V_c = \frac{J + 4R_{a,p}}{4 + 4\Phi} \qquad \text{Eqn 9}$$

160 Using equation 9 to remove V_c from equation 2 yields

$$A_j = \frac{J + 4R_{a,p}}{4 + 4\Phi}(1 - 0.5\Phi) - R_{a,p} - R_x \qquad \text{Eqn 10}$$

163 CO_2 assimilation with electrons recovered by the plastidial anaplerotic pathway in the form of

164 NADPH (Fig. 1) is given as

$$A_{a,p} = \frac{4R_{a,p}}{4 + 4\Phi}(1 - 0.5\Phi) = \frac{R_{a,p}}{1 + \Phi}(1 - 0.5\Phi) \qquad \text{Eqn 11}$$

167 Using equation 4 to remove Φ from equation 10 yields

$$A_j = \frac{J + 4R_{a,p}}{4 + 8\,\Gamma_*/C_c}(1 - \Gamma_*/C_c) - R_{a,p} - R_x \qquad \text{Eqn 12}$$

170 To quantify flux through the plastidial anaplerotic pathway and associated respiration, equation

171 12 is solved for $R_{a,p}$ as

$$R_{a,p} = -\frac{(A_j + R_x)(1 + 2\,\Gamma_*/C_c) - 0.25J(1 - \Gamma_*/C_c)}{3\,\Gamma_*/C_c}$$

<div style="text-align:right">Eqn 13</div>

174   To calculate V_c, Φ is removed from equation 9 using equation 4 as

$$V_c = \frac{J + 4R_{a,p}}{4 + 8\,\Gamma_*/C_c}$$

<div style="text-align:right">Eqn 14</div>

177   Using Φ (Eqn. 4), V_o is calculated as

$$V_o = \Phi * V_c = \frac{2\Gamma_* V_c}{C_c}$$

<div style="text-align:right">Eqn 15</div>

Effects of respiration by the plastidial anaplerotic pathway on A/C_i curves

Respiration by the plastidial anaplerotic pathway can be expected to affect RuBP-regeneration-

limited and rubisco-limited parts of A/C_i curves differently. In the following, effects associated

with RuBP-regeneration-limited CO_2 assimilation are discussed first. Anaplerotic flux of 1 mol

is associated with the release of 1 mol CO_2 and the supply of 2 mol NADPH (Fig. 1). In the

complete absence of photorespiration, all this NADPH is used by the CBC (requires 2 NADPH

per rubisco carboxylation) resulting in complete CO_2 reassimilation. Thus, in this theoretical

scenario, A_j does not change in response to $R_{a,p}$, and gas exchange data do not convey any

information on $R_{a,p}$ no matter how large $R_{a,p}$ is. By contrast, complete photorespiratory use of

NADPH from the anaplerotic pathway (requires 2 NADPH per rubisco oxygenation) involves

the release of 0.5 mol CO_2 by photorespiration per 1 mol CO_2 from the anaplerotic pathway

(i.e., a total of 1.5 mol CO_2 is released). Thus, in the RuBP-regeneration-limited case, $R_{a,p}$

affects A/C_i curves because of photorespiration. The absolute effect of flux through the

plastidial anaplerotic pathway on A/C_i curves (ΔA_j) is given as

$$\Delta A_j = 1.5R_{a,p}\frac{V_o}{V_c + V_o} = \frac{1.5R_{a,p}\Phi}{1 + \Phi} = \frac{3R_{a,p}}{2 + C_c/\Gamma_*}$$

<div style="text-align:right">Eqn 16</div>

196   To calculate $R_{a,p}$ from observed A_j offsets, equation 16 is solved for $R_{a,p}$ as

<div style="text-align:center">7</div>

$$R_{a,p} = \frac{\Delta A_j (1 + \Phi)}{1.5\Phi} = \frac{\Delta A_j (2 + C_c/\Gamma_*)}{3} \qquad \text{Eqn 17}$$

The relative effect of respiration by the plastidial anaplerotic pathway on A/C_i curves is given as

$$\frac{\Delta A_j}{R_{a,p}} = \frac{1.5\Phi}{1 + \Phi} = \frac{3}{2 + C_c/\Gamma_*} \qquad \text{Eqn 18}$$

Hence, the larger Φ, the larger the $\Delta A_j / R_{a,p}$ ratio (Fig. 2, black line). That is, the higher photorespiration, the larger the effect of respiration by the plastidial anaplerotic pathway on A/C_i curves. At $\Phi = 2$ (i.e., $C_c = \Gamma_*$, see Eqn. 4), $\Delta A_j = R_{a,p}$ (i.e., $\Delta A_j / R_{a,p} = 1$). If $\Phi > 2$, then $\Delta A_j > R_{a,p}$. If $\Phi < 2$, then $\Delta A_j < R_{a,p}$.

Under rubisco-limited CO_2 assimilation, RuBP concentrations are above saturation level. Hence, NADPH from the anaplerotic pathway can be expected to exert no control on A_c. Shifts in $R_{a,p}$ will cause A_c shifts of the same size.

NADPH and ATP demand of plastidial carbon metabolism

NADPH demand by the CBC and photorespiration is given as (Farquhar *et al.*, 1980)

$$NADPH\ demand = 2V_c + 2V_o = V_c(2 + 2\Phi) \qquad \text{Eqn 19}$$

This demand is met by NADPH supply from the light reactions. In the presence of flux through the plastidial anaplerotic pathway, NADPH demand from the light reactions is reduced as

$$NADPH\ demand = 2V_c + 2V_o - 2R_{a,p} = V_c(2 + 2\Phi) - 2R_{a,p} \qquad \text{Eqn 20}$$

because anaplerotic flux of 1 mol is associated with the supply of 2 mol NADPH (Fig. 1).

Upon carboxylation of 1 mol RuBP, the CBC requires 3 mol ATP (Farquhar & von Caemmerer, 1982). Upon oxygenation of 1 mol RuBP, photorespiration requires 3 mol ATP in chloroplasts (1.5 mol by phosphoglycerate kinase, 1 mol by phosphoribulokinase, and 0.5 mol by glycerate kinase). This accounting assumes that potentially harmful ammonium released by the

226 mitochondrial glycine decarboxylase complex is immediately recapture by mitochondrial (not

227 plastidial) glutamine synthetase and then transported as glutamine to chloroplasts (Taira *et al.*,

228 2004; Buchanan *et al.*, 2015). ATP required by mitochondrial glutamine synthetase can be

229 expected to come from mitochondrial oxidative phosphorylation. Hence, ATP demand by the

230 CBC and photorespiration from the light reactions is given as

$$ATP\ demand = 3V_c + 3V_o = V_c(3 + 3\Phi) \qquad \text{Eqn 21}$$

233 Combining equations 20 and 21, the ratio at which the CBC and photorespiration require ATP

234 and NADPH is given as

$$\frac{ATP\ demand}{NADPH\ demand} = \frac{3V_c + 3V_o}{2V_c + 2V_o - 2R_{a,p}} = \frac{V_c(3 + 3\Phi)}{V_c(2 + 2\Phi) - 2R_{a,p}} \qquad \text{Eqn 22}$$

Evidently, the CBC and photorespiration have the same plastidial cofactor demands with a

plastidial ATP-to-NADPH demand ratio of 1.5. The ratio is independent of Φ. However,

NADPH demands from the light reactions vary with $R_{a,p}$ resulting in varying ATP-to-NADPH

demands from the light reactions.

Materials and Methods

Samples and leaf gas exchange measurements

The sunflower (*Helianthus annuus*) samples were described previously (Ehlers *et al.*, 2015). In

brief, the plants were raised in a greenhouse at $C_a \approx 450$ ppm and a light intensity of 300-400

µmol photons m^{-2} s^{-1} (16 h photoperiod, 21% O_2). After 7-8 weeks, they were transferred to a

growth chamber ($T = 22$ °C : 18 °C, $rH = 60\%$: 70%, day : night), kept in darkness for 1 day

to drain the starch reserves (see Notes S3 in Wieloch *et al.*, 2022a), and subsequently grown

for 2 days in groups of 8 at C_a of either $\approx 180, 280, 450, 700,$ or 1500 ppm (300-400 µmol

photons m^{-2} s^{-1}, 16 h photoperiod).

Leaf gas exchange measurements were described previously (Ehlers *et al.*, 2015; see Notes S1

in Wieloch *et al.*, 2022a). In brief, the measurements were performed with a previously

described system (Laisk & Edwards, 1997) under conditions similar to those in the growth

chamber ($C_a = 173, 269, 433, 675, 1431$ ppm; $T = 22 \pm 0.5$ °C; 300 µmol photons m^{-2} s^{-1}, 21%

O_2). C_i values obtained from gas exchange measurements (136, 202, 304, 516, 1282 ppm) are

similar to C_i values estimated for the 2-day growth chamber experiments (140, 206, 328, 531,

and 1365 ppm) (see Notes S2 in Wieloch *et al.*, 2022a). To constrain V_{cmax}, we additionally

measured gas exchange at 900 μmol photons m^{-2} s^{-1}, C_a = 173, 269, 432, 674, 1430 ppm

corresponding to C_i = 126, 188, 264, 450, 1204 ppm.

Parameterisation of photosynthesis models

C_c was calculated according to Fick's first law as

$$C_c = C_i - \frac{A}{g_m} \qquad \text{Eqn 23}$$

265 using g_m = 0.45 mol CO_2 m^{-2} s^{-1} reported for sunflower at 20 °C, 600 μmol photons m^{-2} s^{-1}

266 (Schäufele *et al.*, 2011). For rubisco extracted from sunflower, Genkov *et al.* (2010) reported

267 K_c = 19 μM CO_2, K_o = 640 μM O_2, and a relative CO_2/O_2 specificity ($S_{c/o}$) of 77 [M/M] at 25

268 °C. Γ_* at 25 °C was calculated based on published procedures (von Caemmerer, 2000) as

$$\Gamma_* = \frac{0.5 O_c}{S_{c/o}} \qquad \text{Eqn 24}$$

with O_c = 256 μM. To convert from concentration to partial pressure, solubilities for CO_2 of

0.0334 mol l^{-1} bar^{-1} and O_2 of 0.00126 mol l^{-1} bar^{-1} were used. Since correction factors for

sunflower are unavailable, we used Q_{10} values reported in Table 2.3 of von Caemmerer (2000)

and equation 2.32 in von Caemmerer (2000) to correct for leaf temperature (22 °C). This gave

K_c = 447 μbar CO_2, K_o = 439 mbar O_2, and Γ_* = 45.3 μbar CO_2. To estimate V_{cmax}, equation 5

was fitted to gas exchange data obtained at 900 μmol photons m^{-2} s^{-1}, C_a = 173, 269, 432 ppm

(Fig. 3). Under standard conditions (400 ppm CO_2, 21% O_2, 20-25 °C), R_d usually proceeds at

≈ 5% relative to the rate of A (Tcherkez *et al.*, 2017; Xu *et al.*, 2022). Fixing $R_{a,p} + R_x$ at this

rate (= 0.83 μmol CO_2 m^{-2} s^{-1}), we found V_{cmax} = 83.2 μmol m^{-2} s^{-1}. This is similar to a previously

reported estimate for sunflower of 80 μmol m^{-2} s^{-1} at 20 °C (Jacob & Lawlor, 1991;

Wullschleger, 1993).

Based on Farquhar and Wong (1984), J is given as

$$J = \frac{I_2 + J_{max} - \sqrt{(I_2 + J_{max})^2 - 4\theta I_2 J_{max}}}{2\theta} \qquad \text{Eqn 25}$$

286   where θ denotes the empirical curvature factor of the light response of electron transport

287   commonly set to 0.7 (Evans, 1989), J_{max} denotes the light-saturated rate of linear electron

288   transport, and I_2 denotes incident quanta utilised in electron transport through photosystem II

289   given as

$$I_2 = I * \text{abs}(1 - f)/2 \qquad \text{Eqn 26}$$

292   where I denotes incident photon flux, abs denotes absorptance of leaves commonly set to 0.85,

293   and f denotes a correction factor for the spectral quality of light commonly set to 0.15 (Evans,

294   1987). Since gas exchange data collected at very high irradiance are unavailable, J_{max} was

295   estimated based on its close relationship with V_{cmax} according to published procedures (Walker

296   *et al.*, 2014) as

$$\ln(J_{max}) = \beta_1 + \beta_2 \ln(V_{cmax}) \qquad \text{Eqn 27}$$

Using parameter estimates pertaining to data reported by Wullschleger (1993) ($\beta_1 = 1.425$, $\beta_2 =$

0.837, $R^2 = 87.2$, $p < 0.001$, $n = 110$), we estimate $J_{max} = 168.3$ μmol m^{-2} s^{-1}.

Results

Metabolic phases of the A/C_i curve

A/C_i curves enable identification of metabolic processes limiting net CO_2 assimilation

(Farquhar *et al.*, 1980). Initial linear increases at low C_i are commonly attributed to rubisco-

limited assimilation. A change point in the response of A to C_i marks the onset of RuBP-

regeneration-limited assimilation. Furthermore, under high light, A/C_i curves may level out at

high C_i which is commonly attributed to limited triose phosphate utilisation (Sharkey, 1985).

Despite the relatively low resolution of our A/C_i curve, two phases are evident (Fig. 4a). An

initial linear increase (dashed line) is followed by RuBP-regeneration-limited assimilation

whereas levelling out at high C_i is not evident. Hence, triose phosphate utilisation did not affect

A_m (denoting measured net CO_2 assimilation). This is as expected considering the low

irradiance applied during data collection (300 μmol photons m^{-2} s^{-1}).

Previously, we have shown that flux through the plastidial anaplerotic pathway is not significantly different from zero at $C_a \approx 450$ ppm (Wieloch *et al.*, 2022a; Wieloch, 2022). Hence, RuBP-regeneration-limited assimilation for this datapoint as derived from equation 12 is given as

$$A_j = \frac{J}{4 + 8\,\Gamma_*/C_c}(1 - \Gamma_*/C_c) - R_x \qquad \text{Eqn 28}$$

which is equivalent to the original FvCB model. Equation 28 was fitted to A_m by varying R_x until the offset between modelled and measured ≈ 450 ppm values was zero (Fig. 4b, red line). The condition was met at $R_x = 0.96$ μmol CO_2 m^{-2} s^{-1}. Hence, at ≈ 450 ppm, R_d proceeds at 7.8% relative to A_m which agrees well with previous reports. Under standard conditions (400 ppm CO_2, 21% O_2, 20-25 °C), leaf R_d commonly proceeds at $\geq 5\%$ relative to A_m (Tcherkez *et al.*, 2017). This agreement lends some credibility to the validity of our chosen model parameterisation.

Using $R_x = 0.96$ μmol CO_2 m^{-2} s^{-1} and assuming $R_{a,p} = 0$, we next modelled rubisco-limited assimilation based on equation 5 (Fig. 4b, black line). Above $C_a \approx 133$ ppm ($C_i \approx 110$ ppm), we found that $A_c > A_j \geq A_m$ (Fig. 4b, black line versus red line versus black dots). This indicates that A_m at $C_a > 133$ ppm does not result from rubisco-limited but RuBP-regeneration-limited assimilation.

Effects of varying rates of day respiration on the A/C_i curve

Based on changes in hydrogen isotope fractionation in starch extracted from the leaves studied here, we previously estimated $R_{a,p}$ proceeds at > 7%, > 5%, 0%, $\approx 2\%$, and $\approx 5\%$ relative to A_m at $C_a \approx 180, 280, 450, 700,$ and 1500 ppm, respectively (Table 2) (Wieloch, 2022). That is, $R_{a,p}$ exhibits a change point in response to C_a at $C_a \approx 450$ ppm. Seeing that there are positive offsets between A_m and A_j both above and below $C_a \approx 450$ ppm (Table 2; Fig. 4b, red line versus black dots), this same change point appears to be present in A_m, and $R_{a,p}$ can be expected to contribute to the offsets.

Using equation 13, we estimate $R_{a,p}$ proceeds at 54%, 29%, 0%, 5%, and 29% relative to A_m at $C_a \approx 180, 280, 450, 700,$ and 1500 ppm, respectively (Table 2). Estimates at low C_a (≈ 180, and

280 ppm) are consistent with isotope-derived estimates because, based on properties of the biochemical system, the isotope approach can be expected to underestimates $R_{a,p}$ at low C_a (see Wieloch *et al.*, 2022a). By contrast, the isotope approach is believed to not strongly underestimate $R_{a,p}$ at high C_a (Wieloch, 2022). However, gas-exchange estimates of $R_{a,p}$ at high C_a (≈ 700, and 1500 ppm) are considerably larger than isotope-derived estimates. This may suggest that gas exchange modelling overestimates $R_{a,p}$ at high C_a.

Overestimation of $R_{a,p}$ at high C_a may derive from variability of R_x. To follow up on this, we solve the A_j model (Eqn. 12) for R_x and substitute A_j by A_m as

$$R_x = \frac{J + 4R_{a,p}}{4 + 8\,\Gamma_*/C_c}(1 - \Gamma_*/C_c) - R_{a,p} - A_m \qquad \text{Eqn 29}$$

Using isotope-derived $R_{a,p}$ values as inputs, we estimate that absolute rates of R_x increase from 450 to 1500 ppm while relative rates remain approximately constant (Table 2). Based on regulatory properties of leaf respiratory processes, this finding seems reasonable (see 'Metabolic origin of day respiration at high C_a'). Taken together, our results suggest A_m is a function of RuBP-regeneration-limited assimilation and variability of $R_{a,p}$ and R_x (Eqn. 12). Figure 5 summarises estimated carbon and associated NADPH fluxes. Uncertainties associated with these estimates are discussed below ('Model criticism and future research directions').

Integration of day respiration in carbon metabolism
Based on estimations and assumptions given above, $R_{a,p}$ proceeds at $\approx 54\%$, 29%, 0%, 2%, and 5% relative to A_m at $C_a \approx 180$, 280, 450, 700, and 1500 ppm, respectively (Fig. 6a). Thus, it governs R_d at low C_a ($\approx 73\%$ contribution at ≈ 180, and 280 ppm) and contributes substantially to R_d at high C_a ($\approx 22\%$ at ≈ 700 ppm, and $\approx 39\%$ at ≈ 1500 ppm; Fig. 6b). Taken together, R_d proceeds at $\approx 72\%$, 40%, 8%, 9%, and 13% relative to A_m at $C_a \approx 180$, 280, 450, 700, and 1500 ppm, respectively (Table 2). While $R_{a,p}$ exceeds photorespiratory CO_2 release at high C_a (≈ 1500 ppm), it proceeds at $> 55\%$ relative to the rate of photorespiratory CO_2 release at low C_a (≈ 180, and 280 ppm; Fig. 6c). Furthermore, $R_{a,p}$ accounts for $> 40\%$ of the total CO_2 release at both low and high C_a (≈ 180, 280, and 1500 ppm; Fig. 6d). At low C_a (≈ 180, and 280 ppm), $R_{a,p}$ accounts for $> 15\%$ of V_c (Fig. 6e) and thus causes significant futile carbon cycling involving CO_2 uptake by the CBC and release by the plastidial anaplerotic pathway. At the same C_a, $> 10\%$ of all RuBP is provided by the plastidial anaplerotic pathway (Fig. 6f). When plotted

378 against A_m, $R_{a,p}$ increases nonlinearly both below and above $C_a \approx 450$ ppm (Fig. 6g). Towards

379 very low A_m (< 5 µmol m^{-2} s^{-1}, $C_a < 180$ ppm), $R_{a,p}$ may approach a maximum. Above $A_m = 12.3$

380 µmol m^{-2} s^{-1} ($C_a \approx 450$ ppm), $R_{a,p}$ increases exceed R_x increases by approximately a factor of

381 two (Fig. 6g-h).

Energy recovery by the plastidial anaplerotic pathway and associated CO₂ assimilation

NADPH inputs required to assimilate CO_2 respired by the plastidial anaplerotic pathway exceed

NADPH outputs of this pathway (Table 3). However, relative NADPH loss decreases as Φ

decreases with relative NADPH recovery converging to one. Accordingly, CO_2 release by the

anaplerotic pathway ($R_{a,p}$) exceeds CO_2 recovery with NADPH from the anaplerotic pathway

($A_{a,p}$, Table 3). Relative CO_2 recovery increases with decreasing Φ converging to one ($A_{a,p}$ /

$R_{a,p}$). Hence, $R_{a,p}$ is larger than the effect that flux through the plastidial anaplerotic pathway

has on A/C_i curves (ΔA_j, Table 3). With decreasing Φ, the relative effect of flux through the

plastidial anaplerotic pathway on the A/C_i curve decreases converging to zero (ΔA_j / $R_{a,p}$, Table

3, black circles in Fig. 2). Taken together, anaplerotic flux is energetically wasteful and thus

decreases net CO_2 assimilation but this is partly offset by NADPH recovery. Recovered

NADPH is most efficiently used for CO_2 recovery when photorespiration is low since

photorespiration is energetically wasteful and releases additional CO_2 (see 'Theory').

Effects of the plastidial anaplerotic pathway on the light reactions of photosynthesis

In the present study, light was maintained at 300 µmol photons m^{-2} s^{-1} across all C_a treatments.

Accordingly, linear electron transport supplied a steady amount of NADPH (Fig. 5b, yellow

bars). Above and below $C_a \approx 450$ ppm, the plastidial anaplerotic pathway supplied additional

NADPH yet no ATP (Fig. 5b, light green bars). Balancing the ensuing NADPH surplus with

the demands of the CBC and photorespiration requires additional inputs of ATP from

photophosphorylation (Fig. 7, black triangles). Hence, the ratio of ATP-to-NADPH demand

from the light reactions increases both above and below $C_a \approx 450$ ppm with increasing flux

through the plastidial anaplerotic pathway (Fig. 7, red circles).

Assuming the synthesis of 3 mol ATP requires 12 mol H$^+$ as derived from the thermodynamics

of the ATPase reaction (Steigmiller *et al.*, 2008), we find that ATP-to-NADPH supply by linear

electron transport including Q cycling is perfectly aligned with the demands of the CBC and

photorespiration at $C_a \approx 450$ ppm (Table 4, *cf*. Eqn. 22). By contrast, assuming the synthesis of

3 mol ATP requires 14 mol H$^+$ as derived from ATPase subunit composition (Seelert *et al.*,

2000), linear electron transport including Q cycling results in an ATP deficit at $C_a \approx 450$ ppm.

In both cases, ATP deficits increase significantly both above and below $C_a \approx 450$ ppm. This

suggests that increasing flux through the plastidial anaplerotic pathway increases the

requirement for ATP synthesis via cyclic electron transport around photosystem I. Assuming

an H^+-ATP synthesis ratio of 12/3, cyclic electron transport including Q cycling is not required

when the plastidial anaplerotic pathway is inactive, but it may account for up to 17% of the rate

of whole electron transport when this pathway is active. Assuming an H^+-ATP synthesis ratio

of 14/3, cyclic electron transport including Q cycling accounts for $\approx 20\%$ of the rate of whole

electron transport when the plastidial anaplerotic pathway is inactive and may increase to above

30% when this pathway is active. Thus, flux through the plastidial anaplerotic pathway may

affect the light reactions of photosynthesis.

Discussion

Here, we expanded FvCB models by terms accounting for respiration and energy recycling by

the plastidial anaplerotic pathway (Eqs. 5, and 12). We fitted the model of RuBP-regeneration-

limited assimilation to gas exchange data of sunflower leaves by adjusting day respiration. This

approach and previous reports (Wieloch *et al.*, 2022a; Wieloch, 2022) challenge a longstanding

FvCB modelling assumption, namely, the treatment of day respiration as independent, constant

term. If this assumption is false, then several FvCB-model-based methods require revision,

including methods for estimating day respiration (e.g., Laisk, 1977) and mesophyll conductance

(see below). Moreover, initial linear increases in A/C_i curves are commonly interpreted in terms

of rubisco-limited assimilation (Farquhar *et al.*, 1980). Here, we propose RuBP-regeneration-

limited assimilation and varying rates of respiration from the plastidial anaplerotic pathway as

alternative explanation. Our modelling results suggest that the plastidial anaplerotic pathway

can be an important player in plant carbon and energy metabolism affecting properties such as

the rate of day respiration, net CO_2 assimilation, net carboxylation, net oxygenation

(photorespiration), RuBP regeneration, NADPH availability, the ATP-to-NADPH demand ratio

of plastidial carbon metabolism, and probably the contribution of the cyclic electron pathway

to supplying ATP (Figs. 5-7, Tables 2-4).

Consistent evidence in support of flux through the plastidial anaplerotic pathway

According to our gas exchange analysis, flux through the plastidial anaplerotic pathway

proceeds at 0, 2.44, and 2.75 μmol m^{-2} s^{-1} at $C_a \approx 450$, 280, and 180 ppm, respectively (Table

2, Fig. 5a). That is, flux increases with decreasing C_a from $\approx 0\%$ over $\approx 29\%$ to $\approx 54\%$ relative

to the rate of net CO_2 assimilation (Fig. 6a). This is consistent with regulatory properties of the

pathway and previously reported flux estimates as discussed next.

Flux through the plastidial anaplerotic pathway is controlled at its first enzyme, glucose-6-

phosphate dehydrogenase (G6PD, Fig. 1). Under illumination, G6PD is downregulated by a

thioredoxin dependent mechanism (Née et al., 2009). However, downregulation can be reversed

allosterically by G6P (Cossar *et al.*, 1984; Preiser *et al.*, 2019). Under medium to high C_a, the

reaction catalysed by plastidial phosphoglucose isomerase (PGI) was found to be removed from

equilibrium on the side of F6P (Schleucher *et al.*, 1999; Wieloch *et al.*, 2022a; Wieloch, 2022)

resulting in low G6P concentration (Dietz, 1985; Gerhardt *et al.*, 1987; Kruckeberg *et al.*, 1989).

With shifts to low C_a, the PGI reaction moves towards equilibrium (Wieloch *et al.*, 2022a;

Wieloch, 2022) and plastidial G6P concentrations increase (Dietz, 1985). Furthermore,

phosphorolytic starch breakdown increases with photorespiration providing additional G6P

(dotted arrow in Fig. 1; Weise *et al.*, 2006). These mechanisms are consistent with reports of

negligible flux through the anaplerotic pathway at medium C_a (low G6P concentration) and flux

increases with decreasing C_a, increasing photorespiration, and increasing drought (high G6P

concentration; Wieloch *et al.*, 2018, 2022a,b; Wieloch, 2022).

Previously, we estimated flux through the plastidial anaplerotic pathway in the leaves studied

here proceeds at 0%, $> 5\%$ and $> 7\%$ relative to the rate of net CO_2 assimilation at $C_a \approx 450$,

280, and 180 ppm, respectively (Wieloch *et al.*, 2022a; Wieloch, 2022). These estimates are

based on a hydrogen isotope signal in starch introduced in the starch biosynthesis pathway at

the level of G6P H[1]. G6P may be converted back to fructose 6-phosphate (F6P) by PGI, and

F6P may leave the starch biosynthesis pathway via transketolase. Combinedly, G6P to F6P

conversion and F6P use by transketolase can be expected to cause washout of the isotope signal

from the starch biosynthesis pathway. As evident from an isotope signal at starch glucose H[2],

the plastidial PGI reaction is strongly removed from equilibrium on the side of F6P at $C_a \geq 450$

ppm (Wieloch *et al.*, 2022a; Wieloch, 2022). This is thought to impede back conversion of G6P

to F6P and concomitant signal washout. However, the PGI reaction moves progressively

towards equilibrium as C_a decreases below ≈ 450 ppm (Wieloch *et al.*, 2022a; Wieloch, 2022).

This is thought to result in increasing back conversion of G6P to F6P and signal washout. Thus,

increasing offsets between $R_{a.p}$ estimates from isotope and gas exchange analysis with

decreasing C_a probably derive from increasing washout of the isotope signal.

Physiological function of flux through the plastidial anaplerotic pathway

Flux through the anaplerotic pathway lowers net CO_2 assimilation below theoretically possible values (Fig. 4b, black dots versus red line). As a result of this flux, sunflower leaves merely achieve $\approx 88\%$ and $\approx 75\%$ of their theoretical assimilatory potential at $C_a \approx 280$ ppm and ≈ 180 ppm, respectively. Similarly, the anaplerotic pathway recovers only part of the energy it consumes (Fig. 5b, Table 3), and energy dissipation is surely not physiologically beneficial under RuBP-regeneration-limited conditions. Thus, for the conditions and period studied here, the anaplerotic pathway appears to be detrimental for both leaf carbon and energy balances.

However, the sunflowers studied here were raised at $C_a \approx 450$ ppm. Hence, heterotrophic carbon demands were probably adapted to carbon inputs at $C_a \approx 450$ ppm. Moving the plants into low-C_a environments may have caused divergences from the previously established whole-plant steady state with leaf carbon exports exceeding net CO_2 assimilation (Fig. 1). This source limitation may have resulted in excess export of triose phosphates from chloroplasts. Plastidial triose phosphate shortage would impede RuBP regeneration. To maintain the supply of RuBP, plants may inject G6P from phosphorolytic starch breakdown into the CBC (Weise *et al.*, 2006). G6P-derived carbon can enter the CBC via transketolase without respiratory carbon loss and via the anaplerotic pathway with respiratory carbon loss. However, transketolase and other enzymes in the regeneration part of the CBC require triose phosphate. When RuBP regeneration is impeded by triose phosphate shortage, the anaplerotic pathway may help to ensure RuBP supply.

RuBP may undergo either oxygenation or carboxylation at rubisco. With decreasing C_a, oxygenation becomes substantial (Table 3, ≈ 1 and ≈ 1.5 mol RuBP oxygenation per 2 mol RuBP carboxylation at $C_a \approx 280$ ppm and ≈ 180 ppm, respectively). Thus, much of the anaplerotically supplied RuBP feeds into photorespiration. As proposed previously, maintaining photorespiration through inputs of RuBP might make sense physiologically in the context of the concept of photorespiration-linked *de novo* nitrogen assimilation (Wieloch *et al.*, 2022a). Reportedly, photorespiration rate correlates positively with the rate of *de novo* nitrate assimilation (Bloom, 2015), and it seems plausible that plants would shift metabolism towards nitrogen assimilation when carbon assimilation is impeded (at low C_a). Assimilated nitrogen may support *de novo* protein synthesis which may result in increased photosynthetic capacity at the enzyme level and help to overcome source limitations.

In this context, it is interesting to note that mutants incapable of phosphorolytic starch breakdown show severe phenotypes especially under drought (dotted arrow in Fig. 1; Zeeman *et al.*, 2004). This was explained by the loss of G6P injection from starch into the CBC at night (Zeeman *et al.*, 2004). Based on findings presented here and previously, impaired G6P injection during the day may contribute to this phenotype (Wieloch *et al.*, 2018, 2022a,b). Lastly, G6P injection from starch into the CBC by the anaplerotic pathway may support photosynthetic induction (see below).

Metabolic origin of day respiration at high C_a

In our gas exchange analysis, respiration by the plastidial anaplerotic pathway was fixed at 0%, 2%, and 5% relative to the rate of net CO_2 assimilation at $C_a \approx 450$, 700, and 1500 ppm, respectively (Fig. 5a, Table 2). These $R_{a,p}$ estimates come from a previously published isotope analysis of the samples studied here (Wieloch, 2022). Increases in $R_{a,p}$ above $C_a \approx 450$ ppm are consistent with regulatory properties of the plastidial anaplerotic pathway (Wieloch, 2022). Towards high C_a, plastidial G6P concentrations increase with net CO_2 assimilation (Dietz, 1985). This can be expected to cause increased G6PD activity and flux through the plastidial anaplerotic pathway (see above). Note, in contrast to the low-C_a case (see above), $R_{a,p}$ increases at high C_a are not caused by PGI regulation (Wieloch, 2022).

Here, we estimated that respiration by processes other than the plastidial anaplerotic pathway proceeds at 0.96, 1.08, and 1.38 μmol m^{-2} s^{-1} at $C_a \approx 450$, 700, and 1500 ppm, respectively (Fig. 5a, Table 2). These increases are consistent with regulatory properties of the cytosolic OPPP (Wieloch, 2022) which has recently been identified as a major source of day respiration (Xu *et al.*, 2022). Flux through the cytosolic OPPP and associated respiration is controlled at the level of the first OPPP enzyme, G6PD (Fig. 1). Under illumination, cytosolic G6PD activity increases with glucose concentration through *de novo* enzyme synthesis (Hauschild & von Schaewen, 2003). In line with this, respiration at 5 °C was shown to increase with glucose concentration and overall leaf soluble sugar concentration (Tjoelker *et al.*, 2009; Wieloch, 2022). Generally, increasing C_a results in increasing leaf soluble sugar concentration (Ainsworth & Long, 2005) and may thus cause increasing respiration by the cytosolic OPPP.

Does g_m vary with C_a?

Using the online carbon isotope discrimination method and/or the variable J method (Evans *et*
al., 1986; Harley *et al.*, 1992), several authors reported nonlinear relationships between g_m and
C_a with similar patterns across different species including sunflower (Flexas *et al.*, 2007;
Hassiotou *et al.*, 2009; Vrábl *et al.*, 2009; Xiong *et al.*, 2015). As C_a increases, g_m initially
increases up to a change point beyond which it decreases again. This response is strikingly
similar to the C_a-response of day respiration reported here. As C_a increases, $R_{a,p}$ initially
decreases up to a change point beyond which it increases again along with R_x (Fig. 5a, Table
2). Day respiration and g_m have opposite effects on net CO_2 assimilation. That is, both processes
could explain offsets between measured and theoretically possible values of net CO_2
assimilation (Fig. 4b, black dots versus red line). However, changes in day respiration estimated
here based on gas exchange analysis are corroborated by results from an independent isotope
analysis (Wieloch *et al.*, 2022a; Wieloch, 2022). Furthermore, underlying metabolic
mechanisms seem straight forward (see above). By contrast, mechanisms underlying the C_a-
response of g_m have remained unclear despite considerable research interest over the past three
decades (Nadal *et al.*, 2021). Several authors suggested that much of the observed variability
of g_m may be due to methodological shortcomings (Tholen *et al.*, 2012; Gu & Sun, 2014; Yin
et al., 2020; Nadal *et al.*, 2021). Our results corroborate this viewpoint. Therefore, we
recommend modifying the variable J method to allow for variability in day respiration.
Furthermore, we recommend modifying the online carbon isotope discrimination method to
allow for variability in isotope fractionation of photorespiration and day respiration. Adequate
(yet probably challenging) parameterisation of these models may show whether g_m varies with
C_a. Similarly, considering day respiratory variability may advance the discussion about
relationships between g_m and other environmental variables (drought, irradiance, ozone,
temperature; see Nadal *et al.*, 2021).

Respiration by the plastidial anaplerotic pathway during photosynthetic induction
Flux through the plastidial anaplerotic pathway and associated respiration depend on G6PD
activity. Plastidial G6PD is inhibited by reduced thioredoxin (Née et al., 2009). However,
reduction of the plastidial thioredoxin pool after illumination is not instantaneous but builds up
over the course of several minutes (Scheibe, 1981). This was shown to result in a gradual
activation of plastidial malate dehydrogenase (Scheibe, 1981). Similarly, gradual inhibition of
plastidial G6PD can be expected. Thus, flux through the plastidial anaplerotic pathway may
persist over the course of several minutes after illumination decreasing from initially higher to
lower levels. This would affect day respiration, net CO_2 assimilation, NADPH supply, the

isotopic composition of respired CO_2, etc. Future studies on the variability of photosynthetic

parameters such as g_m during photosynthetic induction (*cf.* Kaiser *et al.*, 2017; Sakoda *et al.*,

2021; Liu *et al.*, 2022) are encouraged to consider this as a possibility.

Does day respiration explain part of the A/C_i curve drop at high C_a?

If R_d is held constant, rubisco-limited and RuBP-regeneration-limited net CO_2 assimilation

modelled by the canonical FvCB equations increase continuously with increasing C_a (Farquhar

et al., 1980). However, especially under high light, measured A/C_i curves are frequently seen

to level out or even decline at high C_a (e.g., Sharkey, 1985; Harley & Sharkey, 1991). According

to our analysis, day respiration increases from $\approx 8\%$ relative to the rate of net CO_2 assimilation

at $C_a \approx 450$ ppm to $\approx 13\%$ at $C_a \approx 1500$ ppm (Fig. 5a, Table 2). This is consistent with positive

responses of R_d to C_a reported by some authors (e.g., Wang *et al.*, 2001). By contrast, other

authors reported R_d is independent of C_a (e.g., Ayub *et al.*, 2011).

Increases in R_d are consistent with regulatory properties of both the plastidial and cytosolic

OPPP (see above). As C_a increases from medium to high values, plastidial G6P concentrations

and leaf soluble sugar concentrations increase (Dietz, 1985; Ainsworth & Long, 2005). This

can be expected to activate both plastidial and cytosolic G6PD (Cossar *et al.*, 1984; Hauschild

& von Schaewen, 2003; Preiser *et al.*, 2019), increase respiration by the plastidial and cytosolic

OPPP and explain part of the A/C_i curve drop at high C_a. Furthermore, it may explain part of

the lower than expected stimulation of net CO_2 assimilation by increasing C_a (Wieloch, 2022).

However, high sink strengths (e.g., in vigorously growing plants) may counteract the build-up

of leaf soluble sugar concentrations and respiration by the plastidial and cytosolic OPPP. This

may explain why some studies found R_d increases at high C_a whereas others did not.

Model criticism and future research directions

Generally, modelling net CO_2 assimilation by FvCB-type models relies on numerous

assumptions (Tcherkez & Limami, 2019), and the present study is no exception. For instance,

our models assume that photorespiration is a closed cycle. However, some of the carbon

entering photorespiration may actually be diverted into other pathways such as *de novo* nitrogen

assimilation (Busch *et al.*, 2018). Similarly, in our A_j model (Eqn. 12), the CBC and

photorespiration are assumed to consume all energy. However, part of the available energy can

be expected to be utilised by other plastidial processes such as the reduction of nitrite to

ammonia (Buchanan *et al.*, 2015) or NAD(P)H shuttling to other cell compartments (Scheibe,

2004). Furthermore, our modelling assumes constant absolute R_x at $C_a \leq 450$ ppm (Table 2).

However, leaf soluble sugar concentration may decrease with decreasing C_a (Ainsworth &

Long, 2005). This may cause a reduction in G6PD activity and respiration by the cytosolic

OPPP (Hauschild & von Schaewen, 2003). Lastly, our modelling relies on sunflower parameter

estimates from the literature. Such estimates can vary strongly among species (Orr *et al.*, 2016),

and there is a lack of data on parameter variability in sunflower. Considering these uncertainties,

results reported here require validation (Supporting Information, Notes S1). Hence, we

recommend follow-up studies that combine gas exchange measurements, isotope

measurements, and model parameter measurements from the same plants.

That said, our estimate of R_x at $C_a \approx 450$ ppm is consistent with previously reported estimates

(Tcherkez *et al.*, 2017). Furthermore, estimates of flux through the plastidial anaplerotic

pathway from gas exchange analyses presented here are qualitatively consistent with previously

reported isotope-derived estimates (Wieloch *et al.*, 2022a; Wieloch, 2022). Lastly, all estimates

of day respiration are consistent with regulatory properties of the plastidial and cytosolic

anaplerotic pathway. Hence, we hope that the model extensions and ideas presented here will

help to advance photosynthesis modelling. Future research directions of interest include the

study of $R_{a,p}$ at and below the CO_2 compensation point, and attempts to model $R_{a,p}$ as function

of underlying biochemical mechanisms.

Table 1 List of abbreviations and symbols

Abbreviation	Definition
CBC	Calvin-Benson cycle
F6P	fructose 6-phosphate
FvCB	Farquhar-von Caemmerer-Berry
G6P	glucose 6-phosphate
G6PD	glucose-6-phosphate dehydrogenase
OPPP	oxidative branch of the pentose phosphate pathway
PGI	phosphoglucose isomerase
Ru5P	ribulose 5-phosphate
RuBP	ribulose 1,5-bisphosphate

Symbol	Definition
A	rate of net CO_2 assimilation
A_c	rate of rubisco-limited CO_2 assimilation
A_j	rate of RuBP-regeneration-limited CO_2 assimilation
A_m	measured net CO_2 assimilation
abs	absorptance of leaves
C_a	atmospheric CO_2 concentration
C_c	CO_2 partial pressure or concentration at the active site of rubisco
C_i	CO_2 partial pressure or concentration in intercellular air spaces
g_m	CO_2 mesophyll conductance
I	incident photon flux
I_2	incident quanta utilised in electron transport through photosystem II
J	rate of linear electron transport
$J_{a,p}$	rate of electron recycling by the plastidial anaplerotic pathway
J_{max}	light-saturated rate of linear electron transport
K_c	Michaelis-Menten constants of rubisco carboxylation
K_o	Michaelis-Menten constants of rubisco oxygenation
O_c	oxygen partial pressure at the active site of rubisco
R_d	rate of day respiration
$R_{a,p}$	rate of respiration by and flux through the plastidial anaplerotic pathway
R_x	rate of respiration by pathways other than the plastidial anaplerotic pathway
$S_{c/o}$	relative CO_2/O_2 specificity of rubisco
T	temperature
V_c	rubisco carboxylation rate
V_{cmax}	CO_2-saturated maximum rate of rubisco carboxylation
V_o	rubisco oxygenation rate
Γ_*	CO_2-compensation point in the absence of day respiration
ΔA_j	absolute effect of flux through the plastidial anaplerotic pathway on A/C_i curves
θ	empirical curvature factor of the light response of electron transport
Φ	V_o/V_c ratio

636 **Table 2** Rates of day respiration in sunflower leaves.

C_a [ppm]	A_m [μmol m^{-2} s^{-1}]	A_j - A_m [a] [μmol m^{-2} s^{-1}]	$R_{a,p}$ [b] [μmol m^{-2} s^{-1}]	$R_{a,p}$ / A_m %	$R_{a,p}$ / A_m [d] %	R_x [e] [μmol m^{-2} s^{-1}]	R_x / A_m %
≈ 180	5.13	1.73	2.75	53.7	> 7	0.96	18.8
≈ 280	8.46	1.20	2.44	28.9	> 5	0.96	11.4
≈ 450	12.29	0	0	0	0	0.96	7.8
≈ 700	14.88	0.19	0.80 [c]	5.3 [c]	≈ 2	1.08	7.2
≈ 1500	17.33	0.51	5.03 [c]	29.0 [c]	≈ 5	1.38	8.0

Plants were grown in a greenhouse at an atmospheric CO_2 concentration (C_a) of ≈ 450 ppm and

then moved to growth chambers. After a day in darkness to drain the starch reserves, the plants

were grown at different levels of C_a (180, 280, 450, 700, 1500 ppm) corresponding to different

levels of intercellular CO_2 concentration (C_i = 140, 206, 328, 531, and 1365 ppm) for two days.

During gas exchange measurements slightly lower C_i levels prevailed (136, 202, 304, 516, 1282

ppm). Symbols: A_j, modelled RuBP-regeneration-limited CO_2 assimilation; A_m, measured CO_2

assimilation; $R_{a,p}$, respiration by the plastidial anaplerotic pathway; R_x, day respiration by

processes other than the plastidial anaplerotic pathway.

[a] A_j estimates based on equation 28 with R_x = 0.96 µmol CO_2 m^{-2} s^{-1}.

[b] Estimates based on gas exchange data (Eqn. 13).

[c] Probably overestimated (see text).

[d] Estimates based on previously published isotope data (Wieloch, 2022).

[e] Estimates based on equation 28 for C_a ≈ 180, 280, and 450 ppm, and equation 29 for C_a ≈

700, and 1500 ppm.

Table 3 Energy recovery by the plastidial anaplerotic pathway and associated CO_2 assimilation in sunflower leaves.

C_a [ppm]	Φ	$R_{a,p}$ [a] [µmol m^{-2} s^{-1}]	abs. NADPH input [b] [µmol m^{-2} s^{-1}]	abs. NADPH output [c] [µmol m^{-2} s^{-1}]	rel. NADPH recovery [d]	abs. CO_2 recovery ($A_{a,p}$) [e] [µmol m^{-2} s^{-1}]	rel. CO_2 recovery [f]	ΔA_j [µmol m^{-2} s^{-1}]	$\Delta A_j / R_{a,p}$
≈ 180	0.72	2.75	9.46	5.51	0.58	1.03	0.37	1.73	0.63
≈ 280	0.49	2.44	7.27	4.89	0.67	1.24	0.51	1.20	0.49
≈ 450	0.32	0	0	0	---	0	---	0.00	---
≈ 700	0.18	0.30	0.70	0.60	0.84	0.23	0.77	0.07	0.23
≈ 1500	0.07	0.87	1.86	1.73	0.93	0.78	0.90	0.09	0.10

Plants were grown in a greenhouse at an atmospheric CO_2 concentration (C_a) of ≈ 450 ppm and then moved to growth chambers. After a day in

darkness to drain the starch reserves, the plants were grown at different levels of C_a (180, 280, 450, 700, 1500 ppm) corresponding to different

levels of intercellular CO_2 concentration (C_i = 140, 206, 328, 531, and 1365 ppm) for two days. During gas exchange measurements slightly lower

C_i levels prevailed (136, 202, 304, 516, 1282 ppm). Symbols: $A_{a,p}$, CO_2 assimilation with electrons recovered by the plastidial anaplerotic pathway

in the form of NADPH; $R_{a,p}$, respiration by the plastidial anaplerotic pathway; ΔA_j, absolute effect of flux through the plastidial anaplerotic pathway

on the A/C_i curve (RuBP-regeneration-limited case); Φ, oxygenation-to-carboxylation ratio of rubisco.

[a] Estimates based on equation 13 with R_x values from Table 2.

[b] = $(2+2\Phi)R_{a,p}$ (derivation of Eqn. 7, 2 electrons per NADPH)

[c] = $2R_{a,p}$ (see Fig. 1)

[d] = NADPH output / NADPH input

[e] Estimates based on equation 11.

[f] = $A_{a,p} / R_{a,p}$

Table 4 Effects of the plastidial anaplerotic pathway on electron transport in sunflower leaves.

H$^+$-ATP ratio of ATPase	12/3					14/3				
C_a [ppm]	180	280	450	700	1500	180	280	450	700	1500
LET [µmol m^{-2} s^{-1}]	83.6	83.6	83.6	83.6	83.6	83.6	83.6	83.6	83.6	83.6
ATP demand [µmol m^{-2} s^{-1}]	71.0	70.0	62.7	63.6	65.3	71.0	70.0	62.7	63.6	65.3
ATP from LET [µmol m^{-2} s^{-1}]	62.7	62.7	62.7	62.7	62.7	53.8	53.8	53.8	53.8	53.8
ATP deficit [µmol m^{-2} s^{-1}]	8.3	7.3	0.0	0.9	2.6	17.2	16.3	9.0	9.9	11.6
CET [µmol m^{-2} s^{-1}]	16.5	14.7	0.0	1.8	5.2	40.2	38.0	20.9	23.0	27.0
CET/(CET+LET)	0.17	0.15	0.0	0.02	0.06	0.32	0.31	0.20	0.22	0.24

Plants were grown in a greenhouse at an atmospheric CO$_2$ concentration (C_a) of $\approx$ 450 ppm and then moved to growth chambers. After a day in darkness to drain the starch reserves, the plants were grown at different levels of C_a (180, 280, 450, 700, 1500 ppm) for two days. Gas exchange measurements were performed at a light intensity of 300 µmol photons m^{-2} s^{-1}. Estimations pertaining to linear and cyclic electron transport (LET and CET, both including Q cycling) assumed transmembrane transport of 3 and 2 H$^+$ per electron, respectively. Synthesis of 3 mol ATP by ATPase was assumed to require either 14 or 12 mol H$^+$ (Seelert *et al.*, 2000; Steigmiller *et al.*, 2008). Processes not considered include the water-water cycle (affects the ATP-to-NADPH supply ratio of electron transport), GAP/3PGA cycles and the chloroplast malate valve (affect ATP-to-NADPH demand ratios of carbon metabolism).

Figure 1 Anaplerotic carbon flux into the Calvin-Benson cycle. Dotted arrow, phosphorolytic starch breakdown. Abbreviations: 3PGA, 3-phosphoglycerate; 6PG, 6-phosphogluconate; 6PGL, 6-phosphogluconolactone; ATP, adenosine triphosphate; F6P, fructose 6-phosphate; FBP, fructose 1,6-bisphosphate; G1P, glucose 1-phosphate; G6P, glucose 6-phosphate; G6PD, glucose-6-phosphate dehydrogenase; NADPH, nicotinamide adenine dinucleotide phosphate; PGI, phosphoglucose isomerase; Ru5P, ribulose 5-phosphate; RuBP, ribulose 1,5-bisphosphate; TP, triose phosphate (glyceraldehyde 3-phosphate, dihydroxyacetone phosphate); Xu5P, Xylulose 5-phosphate. Chloroplast and cytosolic metabolism inside and outside the green box, respectively. Solid and dashed arrows represent transformations catalysed by a single enzyme or multiple enzymes, respectively.

Figure 2 Relative effect of respiration by the plastidial anaplerotic pathway on A/C_i curves

under RuBP-regeneration-limited conditions ($\Delta A_j / R_{a,p}$) as function of the oxygenation-to-

carboxylation ratio of rubisco (Φ). Black line: Relationship based on theory (equation 18).

Black circles: $\Delta A_j / R_{a,p}$ ratios pertaining to gas exchange data presented here where ΔA_j denotes

the observed difference between modelled and measured RuBP-regeneration-limited CO_2

assimilation (Fig. 4b, red line versus black dots, respiration by the plastidial anaplerotic

pathway not considered by the model) and $R_{a,p}$ denotes respiration by the plastidial anaplerotic

pathway estimated based on equation 13.

Figure 3 Net CO_2 assimilation (A) of sunflower leaves as function of intercellular CO_2

concentration (C_i). Black dots: measured values (A_m, $n = 8$). Solid line: Best fit between the

data at $C_a = 173, 269, 432$ ppm and a model of rubisco-limited A (Eqn. 5). Plants were raised

in a greenhouse at an atmospheric CO_2 concentration (C_a) of ≈ 450 ppm and a light intensity of

300-400 μmol photons m^{-2} s^{-1}. Gas exchange measurements were performed at 22 °C, 900 μmol

photons m^{-2} s^{-1}, and atmospheric CO_2 concentrations of $C_a = 173, 269, 432, 674, 1430$ ppm

corresponding to intercellular CO_2 concentrations of $C_i = 126, 188, 264, 450, 1204$ ppm. Model

inputs: $C_c = 111, 168, 226$ μbar, $K_c = 447$ μbar CO_2, $K_o = 439$ mbar O_2, and $\Gamma_* = 45.3$ μbar

CO_2, $O_c = 206373$ μbar, $R_{a,p} + R_x = 0.83$ μmol CO_2 m^{-2} s^{-1}. Model output: $V_{cmax} = 83.2$ μmol m$^-$

^{2} s^{-1}.

Figure 4 Measured **(a)** and modelled **(b)** net CO_2 assimilation (A) of sunflower leaves as function of intercellular CO_2 concentration (C_i). Black dots: measured values (A_m, $n = 8$). Dashed black line: Trendline passing through the two lowest A_m values. Solid black and red lines: modelled rubisco-limited assimilation (A_c) and RuBP-regeneration-limited assimilation (A_j), respectively (respiration by the plastidial anaplerotic pathway not considered). Red circle: intersection between A_c and A_j. Plants were grown in a greenhouse at an atmospheric CO_2 concentration (C_a) of ≈ 450 ppm and then moved to growth chambers. After a day in darkness to drain the starch reserves, plants were grown at different levels of C_a (180, 280, 450, 700, 1500 ppm) corresponding to different levels of C_i (140, 206, 328, 531, and 1365 ppm) for two days. During gas exchange measurements slightly lower C_i levels prevailed (136, 202, 304, 516, 1282 ppm). A_j-model inputs: $\Gamma_* = 45.3$ μbar CO_2, $J_{max} = 168.3$ μmol m^{-2} s^{-1}, $\theta = 0.7$, $I = 300$ μmol photons m^{-2} s^{-1}, abs $= 0.85$, and f $= 0.15$. A_j-model output: $R_{a,p} + R_x = 0.96$ μmol CO_2 m^{-2} s^{-1}. A_c-model inputs: $K_c = 447$ μbar CO_2, $K_o = 439$ mbar O_2, and $\Gamma_* = 45.3$ μbar CO_2, $O_c = 207548$ μbar, $V_{cmax} = 83.2$ μmol m^{-2} s^{-1}, $R_{a,p} + R_x = 0.96$ μmol CO_2 m^{-2} s^{-1}.

Figure 5 Carbon **(a)** and NADPH fluxes **(b)** at varying atmospheric CO_2 concentration (C_a) in sunflower leaves. Estimates derived from modelling RuBP-regeneration-limited CO_2 assimilation. The model considers the Calvin-Benson cycle, the photorespiration cycle, the plastidial anaplerotic pathway, and respiration by processes other than the plastidial anaplerotic pathway. Plants were grown in a greenhouse at $C_a \approx 450$ ppm and then moved to growth chambers. After a day in darkness to drain the starch reserves, plants were grown at different levels of C_a (180, 280, 450, 700, 1500 ppm) corresponding to different levels of C_i (140, 206, 328, 531, and 1365 ppm) for two days. During gas exchange measurements slightly lower C_i levels prevailed (136, 202, 304, 516, 1282 ppm). NADPH demands of individual carbon fluxes (see Fig. 5a) were calculated as $(2 + 2\Phi)$ * flux with flux = $\{-A_m; 0.5V_o; R_x; R_{a,p}\}$. Symbols: A_m, measured net CO_2 assimilation; A_m', NADPH demand of net CO_2 assimilation; J', NADPH supply by whole chain electron transport ($0.5J$); $R_{a,p}$, respiration by the plastidial anaplerotic pathway; $R_{a,p}$', NADPH supply (light green) and demand (dark green) by the plastidial anaplerotic pathway; R_x, day respiration by processes other than the plastidial anaplerotic pathway; R_x', NADPH demand for assimilation of CO_2 respired by processes other than the plastidial anaplerotic pathway; V_c, rubisco carboxylation; $0.5V_o$, photorespiratory CO_2 release; $0.5V_o$', NADPH demand for assimilation of CO_2 respired by photorespiration; Φ, oxygenation-to-carboxylation ratio of rubisco.

Figure 6 Integration of day respiration in carbon metabolism of sunflower leaves. Plants were
grown in a greenhouse at $C_a \approx 450$ ppm and then moved to growth chambers. After a day in
darkness to drain the starch reserves, plants were grown at different levels of C_a (180, 280, 450,
700, 1500 ppm) corresponding to different levels of C_i (140, 206, 328, 531, and 1365 ppm) for
two days. During gas exchange measurements slightly lower C_i levels prevailed (136, 202, 304,
516, 1282 ppm). Symbols: A_m, measured CO_2 assimilation; C_a, atmospheric CO_2 concentration;
C_i, intercellular CO_2 concentration; $R_{a,p}$, respiration by and flux through the plastidial
anaplerotic pathway; R_d, total day respiration; R_x, day respiration by processes other than the
plastidial anaplerotic pathway; V_c, rubisco carboxylation; V_o, rubisco oxygenation. All panels
show discrete data. Dashed lines in added to guide the eye.
761

Figure 7 Energy demand from the light reactions of photosynthesis at varying atmospheric CO_2 concentration (C_a) in sunflower leaves. Black triangles: ATP demand. Red circles: ATP-to-NADPH demand ratio. Energy fluxes were estimated based on carbon flux estimates from modelling RuBP-regeneration-limited CO_2 assimilation. The model considers the Calvin-Benson cycle, the photorespiration cycle, the plastidial anaplerotic pathway, and respiration by processes other than the plastidial anaplerotic pathway. Plants were grown in a greenhouse at $C_a \approx 450$ ppm and then moved to growth chambers. After a day in darkness to drain the starch reserves, plants were grown at different levels of C_a (180, 280, 450, 700, 1500 ppm) corresponding to different levels of C_i (140, 206, 328, 531, and 1365 ppm) for two days. During gas exchange measurements slightly lower C_i levels prevailed (136, 202, 304, 516, 1282 ppm).

Acknowledgements

We thank Prof. Jun Yu (Umeå University) for helpful discussions and bioRxiv (Cold Spring

Harbor Laboratory, NY, USA) for publishing preprints of the present paper

(https://doi.org/10.1101/2021.07.30.454461).

Author contributions

T.W. conceived the study, led the research, expanded FvCB models, and analysed the data.

A.A., and J.S. prepared samples, and acquired gas exchange data. T.W. wrote the paper with

input from all authors. T.W. revised the paper.

Competing interests: The authors declare no competing interests.

Data availability statement: Data available on request from the authors.

References

Ainsworth EA, Long SP. **2005**. What have we learned from 15 years of free-air CO_2

enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy

properties and plant production to rising CO_2. *New Phytologist* **165**: 351–372.

Ayub G, Smith RA, Tissue DT, Atkin OK. **2011**. Impacts of drought on leaf respiration in

darkness and light in *Eucalyptus saligna* exposed to industrial-age atmospheric CO_2 and growth

temperature. *New Phytologist* **190**: 1003–1018.

Bloom AJ. **2015**. Photorespiration and nitrate assimilation: A major intersection between plant

carbon and nitrogen. *Photosynthesis Research* **123**: 117–128.

Buchanan BB, Gruissem W, Jones RL (Eds.). **2015**. *Biochemistry and molecular biology of*

plants. Chichester: John Wiley & Sons, Ltd.

Busch FA, Sage RF, Farquhar GD. **2018**. Plants increase CO_2 uptake by assimilating nitrogen

via the photorespiratory pathway. *Nature Plants* **4**: 46–54.

von Caemmerer S. **2000**. *Biochemical models of leaf photosynthesis*. Collingwood: CSIRO.

Cossar JD, Rowell P, Stewart WDP. **1984**. Thioredoxin as a modulator of glucose-6-

phosphate dehydrogenase in a N_2-fixing cyanobacterium. *Microbiology* **130**: 991–998.

Dietz K-J. **1985**. A possible rate-limiting function of chloroplast hexosemonophosphate
isomerase in starch synthesis of leaves. *Biochimica et Biophysica Acta* **839**: 240–248.

Ehlers I, Augusti A, Betson TR, Nilsson MB, Marshall JD, Schleucher J. **2015**. Detecting
long-term metabolic shifts using isotopomers: CO_2-driven suppression of photorespiration in
C_3 plants over the 20th century. *Proceedings of the National Academy of Sciences of the United*
States of America **112**: 15585–15590.

Eicks M, Maurino V, Knappe S, Flügge U-I, Fischer K. **2002**. The plastidic pentose
phosphate translocator represents a link between the cytosolic and the plastidic pentose
phosphate pathways in plants. *Plant Physiology* **128**: 512–522.

Evans J. **1987**. The dependence of quantum yield on wavelength and growth irradiance.
Functional Plant Biology **14**: 69–79.

Evans JR. **1989**. Photosynthesis and nitrogen relationships in leaves of C_3 plants. *Oecologia*
78: 9–19.

Evans JR, Farquhar GD, Sharkey TD, Berry JA. **1986**. Carbon isotope discrimination
measured concurrently with gas exchange to investigate CO_2 diffusion in leaves of higher
plants. *Australian Journal of Plant Physiology* **13**: 281–292.

Farquhar GD, Caemmerer S, Berry JA. **1980**. A biochemical model of photosynthetic CO_2
assimilation in leaves of C_3 species. *Planta* **149**: 78–90.

Farquhar GD, von Caemmerer S. **1982**. Modelling of photosynthetic response to
environmental conditions. In: Lange OL, Nobel PS, Osmond CB, Ziegler H, eds. Physiological
plant ecology II: Water relations and carbon assimilation. Berlin, Heidelberg: Springer, 549–
587.

Farquhar G, Wong S. **1984**. An empirical model of stomatal conductance. *Functional Plant*
Biology **11**: 191–210.

Flexas J, Diaz-Espejo A, GalméS J, Kaldenhoff R, Medrano H, Ribas-Carbo M. **2007**.
Rapid variations of mesophyll conductance in response to changes in CO_2 concentration around
leaves. *Plant, Cell & Environment* **30**: 1284–1298.

Genkov T, Meyer M, Griffiths H, Spreitzer RJ. **2010**. Functional hybrid Rubisco enzymes
with plant small subunits and algal large subunits. *Journal of Biological Chemistry* **285**: 19833–
19841.

Gerhardt R, Stitt M, Heldt HW. **1987**. Subcellular metabolite levels in spinach leaves:
Regulation of sucrose synthesis during diurnal alterations in photosynthetic partitioning. *Plant*
Physiology **83**: 399–407.

Gu L, Sun Y. **2014**. Artefactual responses of mesophyll conductance to CO_2 and irradiance
estimated with the variable J and online isotope discrimination methods. *Plant, Cell &*
Environment **37**: 1231–1249.

Harley PC, Loreto F, Di Marco G, Sharkey TD. **1992**. Theoretical considerations when
estimating the mesophyll conductance to CO_2 flux by analysis of the response of photosynthesis
to CO_2. *Plant Physiology* **98**: 1429–1436.

Harley PC, Sharkey TD. **1991**. An improved model of C_3 photosynthesis at high CO_2:
Reversed O_2 sensitivity explained by lack of glycerate reentry into the chloroplast.
Photosynthesis Research **27**: 169–178.

Hassiotou F, Ludwig M, Renton M, Veneklaas EJ, Evans JR. **2009**. Influence of leaf dry
mass per area, CO_2, and irradiance on mesophyll conductance in sclerophylls. *Journal of*
Experimental Botany **60**: 2303–2314.

Hauschild R, von Schaewen A. **2003**. Differential regulation of glucose-6-phosphate
dehydrogenase isoenzyme activities in potato. *Plant Physiology* **133**: 47–62.

Jacob J, Lawlor DW. **1991**. Stomatal and mesophyll limitations of photosynthesis in
phosphate deficient sunflower, maize and wheat plants. *Journal of Experimental Botany* **42**:
1003–1011.

Kaiser E, Kromdijk J, Harbinson J, Heuvelink E, Marcelis LFM. **2017**. Photosynthetic
induction and its diffusional, carboxylation and electron transport processes as affected by CO_2
partial pressure, temperature, air humidity and blue irradiance. *Annals of Botany* **119**: 191–205.

Kruckeberg AL, Neuhaus HE, Feil R, Gottlieb LD, Stitt M. **1989**. Decreased-activity

mutants of phosphoglucose isomerase in the cytosol and chloroplast of *Clarkia xantiana*. *The*

Biochemical Journal **261**: 457–467.

Laisk A. **1977**. Kinetics of photosynthesis and photorespiration in C$_3$ plants. Nauka, Moscow,

Russia.

Laisk A, Edwards GE. **1997**. CO$_2$ and temperature-dependent induction in C$_4$ photosynthesis:

an approach to the hierarchy of rate-limiting processes. *Functional Plant Biology* **24**: 505–516.

Liu T, Barbour MM, Yu D, Rao S, Song X. **2022**. Mesophyll conductance exerts a significant

limitation on photosynthesis during light induction. *New Phytologist* **233**: 360–372.

Nadal M, Carriquí M, Flexas J. **2021**. Chapter 3 Mesophyll conductance to CO$_2$ diffusion in

a climate change scenario: Effects of elevated CO$_2$, temperature and water stress. In: Becklin

KM, Ward JK, Way DA, eds. Photosynthesis, Respiration, and Climate Change. Cham:

Springer International Publishing, 49–78.

Orr DJ, Alcântara A, Kapralov MV, Andralojc PJ, Carmo-Silva E, Parry MAJ. **2016**.

Surveying Rubisco diversity and temperature response to improve crop photosynthetic

efficiency. *Plant Physiology* **172**: 707–717.

Preiser AL, Fisher N, Banerjee A, Sharkey TD. **2019**. Plastidic glucose-6-phosphate

dehydrogenases are regulated to maintain activity in the light. *Biochemical Journal* **476**: 1539–

1551.

Sakoda K, Yamori W, Groszmann M, Evans JR. **2021**. Stomatal, mesophyll conductance,

and biochemical limitations to photosynthesis during induction. *Plant Physiology* **185**: 146–

160.

Schäufele R, Santrucek J, Schnyder H. **2011**. Dynamic changes of canopy-scale mesophyll

conductance to CO$_2$ diffusion of sunflower as affected by CO$_2$ concentration and abscisic acid.

Plant, Cell & Environment **34**: 127–136.

Scheibe R. **1981**. Thioredoxinm in pea chloroplasts: Concentration and redox state under light

and dark conditions. *FEBS Letters* **133**: 301–304.

Scheibe R. **2004**. Malate valves to balance cellular energy supply. *Physiologia Plantarum* **120**:
21–26.

Schleucher J, Vanderveer P, Markley JL, Sharkey TD. **1999**. Intramolecular deuterium
distributions reveal disequilibrium of chloroplast phosphoglucose isomerase. *Plant, Cell and*
Environment **22**: 525–533.

Seelert H, Poetsch A, Dencher NA, Engel A, Stahlberg H, Müller DJ. **2000**. Proton-powered
turbine of a plant motor. *Nature* **405**: 418–419.

Sharkey TD. **1985**. O_2-insensitive photosynthesis in C_3 plants: Its occurrence and a possible
explanation. *Plant Physiology* **78**: 71–75.

Sharkey TD, Berry JA, Raschke K. **1985**. Starch and sucrose synthesis in *Phaseolus vulgaris*
as affected by light, CO_2, and abscisic acid. *Plant Physiology* **77**: 617–620.

Sharkey TD, Weise SE. **2016**. The glucose 6-phosphate shunt around the Calvin–Benson
cycle. *Journal of Experimental Botany* **67**: 4067–4077.

Steigmiller S, Turina P, Gräber P. **2008**. The thermodynamic H^+/ATP ratios of the H^+-
ATPsynthases from chloroplasts and *Escherichia coli*. *Proceedings of the National Academy of*
Sciences of the United States of America **105**: 3745–3750.

Taira M, Valtersson U, Burkhardt B, Ludwig RA. **2004**. *Arabidopsis thaliana* GLN2-
encoded glutamine synthetase is dual targeted to leaf mitochondria and chloroplasts. *The Plant*
Cell **16**: 2048–2058.

Tcherkez G, Gauthier P, Buckley TN, Busch FA, Barbour MM, Bruhn D, Heskel MA,
Gong XY, Crous KY, Griffin K, *et al.* **2017**. Leaf day respiration: low CO_2 flux but high
significance for metabolism and carbon balance. *New Phytologist* **216**: 986–1001.

Tcherkez G, Limami AM. **2019**. Net photosynthetic CO_2 assimilation: more than just CO_2 and
O_2 reduction cycles. *New Phytologist* **223**: 520–529.

Tholen D, Ethier G, Genty B, Pepin S, Zhu X-G. **2012**. Variable mesophyll conductance
revisited: theoretical background and experimental implications. *Plant, Cell & Environment* **35**:
2087–2103.

Tjoelker MG, Oleksyn J, Lorenc-Plucinska G, Reich PB. **2009**. Acclimation of respiratory
temperature responses in northern and southern populations of *Pinus banksiana*. *New*
Phytologist **181**: 218–229.

Vrábl D, Vašková M, Hronková M, Flexas J, Šantrůček J. **2009**. Mesophyll conductance to
CO_2 transport estimated by two independent methods: effect of variable CO_2 concentration and
abscisic acid. *Journal of Experimental Botany* **60**: 2315–2323.

Walker AP, Beckerman AP, Gu L, Kattge J, Cernusak LA, Domingues TF, Scales JC,
Wohlfahrt G, Wullschleger SD, Woodward FI. **2014**. The relationship of leaf photosynthetic
traits - V_{cmax} and J_{max} - to leaf nitrogen, leaf phosphorus, and specific leaf area: a meta-analysis
and modeling study. *Ecology and evolution* **4**: 3218–3235.

Wang X, Lewis JD, Tissue DT, Seemann JR, Griffin KL. **2001**. Effects of elevated
atmospheric CO_2 concentration on leaf dark respiration of *Xanthium strumarium* in light and in
darkness. *Proceedings of the National Academy of Sciences of the United States of America* **98**:
2479–2484.

Weise SE, Schrader SM, Kleinbeck KR, Sharkey TD. **2006**. Carbon balance and circadian
regulation of hydrolytic and phosphorolytic breakdown of transitory starch. *Plant Physiology*
141: 879–886.

Wieloch T. **2021**. The next phase in the development of ^{13}C isotopically nonstationary
metabolic flux analysis. *Journal of Experimental Botany* **72**: 6087–6090.

Wieloch T. **2022**. High atmospheric CO_2 concentration causes increased respiration by the
oxidative pentose phosphate pathway in chloroplasts. *New Phytologist* **235**: 1310–1314.

Wieloch T, Augusti A, Schleucher J. **2022a**. Anaplerotic flux into the Calvin-Benson cycle.
Hydrogen isotope evidence for *in vivo* occurrence in C_3 metabolism. *New Phytologist* **234**: 405–
411.

Wieloch T, Ehlers I, Yu J, Frank D, Grabner M, Gessler A, Schleucher J. **2018**.
Intramolecular ^{13}C analysis of tree rings provides multiple plant ecophysiology signals covering
decades. *Scientific Reports* **8**: 5048.

Wieloch T, Grabner M, Augusti A, Serk H, Ehlers I, Yu J, Schleucher J. **2022b**. Metabolism is a major driver of hydrogen isotope fractionation recorded in tree-ring glucose of *Pinus nigra*. *New Phytologist* **234**: 449–461.

Wieloch T, Sharkey TD. **2022**. Compartment-specific energy requirements of photosynthetic carbon metabolism in *Camelina sativa* leaves. *Planta* **255**: 103.

Wullschleger SD. **1993**. Biochemical limitations to carbon assimilation in C_3 plants. A retrospective analysis of the A/C_i curves from 109 Species. *Journal of Experimental Botany* **44**: 907–920.

Xiong D, Liu X, Liu L, Douthe C, Li Y, Peng S, Huang J. **2015**. Rapid responses of mesophyll conductance to changes of CO_2 concentration, temperature and irradiance are affected by N supplements in rice. *Plant, Cell & Environment* **38**: 2541–2550.

Xu Y, Fu X, Sharkey TD, Shachar-Hill Y, Walker BJ. **2021**. The metabolic origins of non-photorespiratory CO_2 release during photosynthesis: A metabolic flux analysis. *Plant Physiology* **186**: 297–314.

Xu Y, Wieloch T, Kaste JAM, Shachar-Hill Y, Sharkey TD. **2022**. Reimport of carbon from cytosolic and vacuolar sugar pools into the Calvin–Benson cycle explains photosynthesis labeling anomalies. *Proceedings of the National Academy of Sciences of the United States of America* **119**: e2121531119.

Yin X, van der Putten PEL, Belay D, Struik PC. **2020**. Using photorespiratory oxygen response to analyse leaf mesophyll resistance. *Photosynthesis Research* **144**: 85–99.

Zeeman SC, Thorneycroft D, Schupp N, Chapple A, Weck M, Dunstan H, Haldimann P, Bechtold N, Smith AM, Smith SM. **2004**. Plastidial α-glucan phosphorylase is not required for starch degradation in Arabidopsis leaves but has a role in the tolerance of abiotic stress. *Plant Physiology* **135**: 849–858.