The effects of a synthetic and biological surfactant on the community composition and

metabolic activity of a freshwater biofilm

Running Title: Surfactant effects: Biofilm composition and metabolism

Authors: Stephanie P. Gill[1], William R. Hunter[2], Laura Coulson[3,4], and Jakob Schelker[3,5]

[1]Department of Geography and Environmental Studies, Ulster University Coleraine, Northern

Ireland

[2]Fisheries and Aquatic Ecosystems Branch, Agri-Food and Biosciences Institute, Belfast,

Northern Ireland

[3]WasserCluster Lunz, Lunz Am See, Austria

[4]Institute of Hydrobiology and Aquatic Ecosystem Management, University of Natural

Resources and Life Sciences, Vienna, Austria

[5] Department of Functional and Evolutionary Ecology, University of Vienna, Vienna, Austria

*Correspondence: Stephanie P. Gill

18   School of Geography and Environmental Science
19   Ulster University, Coleraine Campus
20   BT52 1SA. United Kingdom
21   Email: gill-S5@ulster.ac.uk
22   Telephone: 07541-036849
23

24 Keywords (5): Biofilm, surfactants, freshwater, metabolic, sequencing

Originality-Significance Statement

To our knowledge this is the first study that examines the effects of the chemical surfactant

sodium dodecyl sulphate and the biological surfactant rhamnolipid on a mixed species freshwater

biofilm. Research typically focuses on single species biofilms in medical or industrial settings.

Therefore, we are also the first study that examines the community composition and metabolic

activity of a mixed freshwater biofilm exposed to these surfactants, and are the only study that

has examined the carbon and nitrogen content of an exposed aquatic biofilm. Our results will

help to shape policy surrounding the input of contaminants of emerging concern such as

surfactants into waterways, and potentially to help manage the use of these chemicals within

industrial settings as a means to control biofilms.

Summary

Surfactants are used to control microbial biofilms in industrial and medical settings. Their

known toxicity on aquatic biota and their longevity in the environment has encouraged research

on biodegradable alternatives such as rhamnolipids. While previous research has investigated the

effects of biological surfactants on single species biofilms, there remains a lack of information

regarding the effects of synthetic and biological surfactants in freshwater ecosystems. We

conducted a mesocosm experiment to test how the surfactant sodium dodecyl sulphate (SDS) and

the biological surfactant rhamnolipid altered community composition and metabolic activity of

freshwater biofilms. Biofilms were cultured in the flumes using lake water from Lake Lunz in

Austria, under high (300 ppm) and low (150 ppm) concentrations of either surfactant over a four

week period. Our results show that both surfactants significantly affected microbial diversity. Up

to 36% of microbial OTUs were lost after surfactant exposure. Rhamnolipid exposure also

increased the production of the extracellular enzymes leucine aminopeptidase and glucosidase,

while SDS exposure reduced leucine aminopeptidase and glucosidase. This study demonstrates

that exposure of freshwater biofilms to chemical and biological surfactants caused a reduction of

microbial diversity and changes in biofilm metabolism, exemplified by shifts in extracellular

enzyme activities.

Introduction

In aquatic ecosystems, microorganisms live preferentially in microbial biofilms and act as major drivers of the biogeochemical cycling of carbon, nitrogen, and other nutrients (Paul *et al.,* 1991; Decho, 2000; Dang and Lovell, 2016; Flemming *et al.,* 2016; Battin *et al.,* 2016). In streams and rivers, microbial biofilms play an important role in the carbon cycle (Lyon and Ziegler, 2009; Qu *et al.,* 2017). Microorganisms within biofilms are able to break down dissolved organic matter and utilize the dissolved organic carbon (DOC) within the compounds (Romani *et al.,* 2003). Through mineralization, aquatic biofilms are then able to produce large quantities of carbon dioxide for release back into the atmosphere (Kroukamp and Wolfaardt, 2008). Similarly, depending on the nitrogen levels in the system, microbial biofilms resident in fluvial sediments can substantially reduce inorganic nitrogen (N) loads in N-rich aquatic ecosystems through denitrification (Zhang *et al.,* 2016; Cui *et al.,* 2017). In N-limited aquatic systems, they may help to facilitate the transfer of N to higher trophic levels (Dodds *et al.,* 2000).

While microbial biofilms play an important ecological role, they are also of concern in industrial and medical applications. For example, biofilms can facilitate the growth of antibiotic resistant bacteria (Lindsay and von Holy, 2006; Srey *et al.,* 2013; Bowler *et al.,* 2020). Synthetic surfactants, such as sodium dodecyl sulphate (SDS), are widely used in personal care and cleaning products. They reduce biofilm growth in a wide range of industrial, medical, and environmental contexts (Waaler *et al.,* 1993; Landa *et al.,* 1999; Rigotti *et al.,* 2017; Sloup *et al.,* 2016; Ueda *et al.,* 2019). Surfactants are amphiphilic chemicals that reduce surface tension within a liquid through the formation of micelles. Micelles occur when molecules form an aggregate with hydrophobic tails on the inside (Schmitz, 2018) and hydrophilic heads on the outside of an amphiphilic compound (Milovanovic *et al.,* 2017). An early study estimated the

global use of surfactants at 36.4 million tons per year (Hauthal, 2004; Mungray and Kumar,

2009), suggesting that synthetic surfactant may be commonly transferred into freshwater

ecosystems. A previous report by the European Commission (Thornton *et al.,* 2001) has

highlighted surfactants as one of the top contaminants found in wastewater treatment effluent.

Given the mass of surfactants produced every year, it can well be anticipated that surfactants still

represent some of the most abundant chemicals in municipal and industrial wastewater outflow.

Wastewater treatment plants (WWTPs) have been observed to have very high levels of

surfactants in their inflows (for example, up to 1900 ug L^{-1} for LAS and 40,000 ug L^{-1} for alkyl

ethoxysulfates (AES) (Gomez *et al.,* 2011)) with removal rates of only 40-90% during waste-

water treatment (Gomez *et al.,* 2011; Kruszelnicka *et al.,* 2019). Similarly, surfactant levels of

raw water intakes of drinking water treatment plants (DWTPs) can be high (up to 49 ug L^{-1} for

LAS and 114 ug L^{-1} for AES) (Gomez *et al.,* 2011) with only partial removal of surfactants by

DWTPs. Finally, high levels of surfactants have also been identified in surface sediments

obtained near wastewater treatment plant discharge areas into coastal waters, such as the Jamaica

Bay of New York (Lara-Martín *et al.,* 2012). This underlines the presence and persistence of

surfactants in aquatic ecosystems downstream of urban and industrial environments.

Research on the effects of surfactants on microbial biofilms has primarily focused on

single-species environmental biofilms or those commonly found in medical and industrial

environments. Direct applications of the anionic synthetic surfactant SDS has shown significant

disruption of growth and development of *Pseudomonas aeruginosa* biofilms (Díaz De Rienzo *et*

al., 2016; Nguyen *et al.,* 2020). Exposure to SDS has also decreased biofilm density and

flattened its shape (Robinson *et al.,* 2006). Effects on other single species or medical biofilms

have also been investigated including methicillin-resistant *Staphylococcus aureus* (Ueda *et al.,*

2019)*, Pseudomonas fluorescens* (Simoes *et al.,* 2008)*, and Salmonella enterica* serovar

typhimurium (Mireles II *et al.,* 2001). Inhibited biofilm formation was seen to occur within all

investigations.

The use of synthetic surfactants has been questioned due to the potential environmental

impact, specifically the longevity and toxicity to aquatic biota. For example, previous research

has shown that synthetic surfactants, such as Triton X-100 (Octylphenol Ethoxylate), are not

easily biodegradable under aerobic nor anaerobic conditions (Mohan *et al.,* 2006). Other

synthetic surfactants such as SDS have also been noted as toxic to a variety of microorganisms

such as *Vibrio fischeri,* fish and shellfish such as *Clarias batrachus* and *Mytilus*

galloprovincialis, yeast, and also small invertebrates, such as *Daphnia magna,* and *Tigriopus*

fulvus (Mariani *et al.,* 2006; Lima *et al.,* 2011; Franzetti *et al.,* 2012; Messina *et al.,* 2014; Cao *et*

al., 2020; Kumar *et al.,* 2020). Research has therefore shifted towards identifying a natural

alternative that may be less harmful to the aquatic environment.

Biofilm-dwelling microorganisms naturally produce a number of surfactants to control

the surrounding extracellular polymeric substance (EPS) matrix. The most common substances

of this group are rhamnolipids and sophorolipids which are glycolipids formed by the

combination of either a long-chain aliphatic acid or hydroxyaliphatic acid with a carbohydrate

(Desai and Banat, 1997). When exposing single-cell biofilms of *P. aeruginosa* to rhamnolipids,

growth and development were disrupted, and cell viability was reduced (De Rienzo *et al.,* 2015).

Disruptions by rhamnolipids have also been shown for single species biofilms of *Staphylococcus*

aureus (Quinn *et al.,* 2013; Aleksic *et al.,* 2017; Tambone *et al.,* 2021)*, Bacillus pumilus*

(Dusane *et al.,* 2010)*, and Rhodococcus erythropolis* (Schreiberova *et al.,* 2012). Yet,

rhamnolipids are biodegradable in a variety of environments, including under aerobic and

anaerobic conditions (Mohan *et al.,* 2006; Ramos da Silva *et al.,* 2019). Biological surfactants

can also function in more varied environmental conditions such as large pH and temperature

ranges (Muthusamy *et al.,* 2008). Given that previous research primarily addressed the effects of

biological surfactants on single species biofilms of bacteria relevant to medical settings, there

remains a knowledge gap on how the exposure of multi-species freshwater biofilms to

surfactants affects biofilm microbial community composition and function.

There is a growing body of evidence that anthropogenic trace contaminants such as

pharmaceuticals, herbicides, and antimicrobial agents can have major impacts upon the

composition and functioning of aquatic microbial communities (Rosi-Marshall *et al.,* 2013;

Harjung *et al.,* 2020; McClean and Hunter, 2020; Ke *et al.,* 2020; Kusi *et al.,* 2020). Exposure to

contaminants such as the antibacterial agent Tricolsan (Hay *et al.,* 2001; Drury *et al.,* 2013) and

the herbicide Diuron (Sumpono *et al.,* 2003; Seghers *et al.,* 2003) have been observed to place

microbial communities under a selective pressure that favors microorganisms that can use these

contaminants as an additional carbon source. Given that levels of surfactants and their

metabolites are consistently among the top contaminants identified within wastewater influent

and groundwater (Lara-Martín *et al.,* 2014; Schaider *et al.,* 2014), it is reasonable to infer that

they could represent a major control upon the microbial communities within human affected

inland waters.

This study investigated the effects of the synthetic surfactant SDS and the biological

surfactant rhamnolipid on microbial biofilm community composition, microbial diversity, and

metabolic activity. Using recirculating flume mesocosms, we aimed to answer the following

overarching scientific question: What are the effects of a synthetic and biological surfactant on a

natural aquatic microbial biofilm, and how do these effects compare? We hypothesized that

biofilm exposure to a surfactant will decrease bacterial biodiversity, as the surfactants impact

microbial community composition and causes the loss of sensitive bacterial taxa. Secondly, we

hypothesize that the changes in microbial community composition will alter biofilm metabolic

activity. The loss of sensitive taxa will allow generalists to dominate who do not have the same

primary functions and efficiency as more specialized taxa, reducing biofilm performance,

expressed by metabolic activity. From this, we predict that the concentrations of extracellular

enzymes produced by the biofilms, used as a proxy for community-scale metabolic activity

within the biofilms, will be lower in the surfactant treatments. We tested these hypotheses by

quantifying changes in community composition through 16s rRNA analysis and changes in

biofilm structure and metabolic activity through different analytical methods, such as changes in

gross primary productivity (GPP), community respiration (CR), ash free dry mass (AFDM),

Carbon: Nitrogen ratios (C:N), and extracellular enzyme activity (EEA).

Results

Primary productivity and community respiration

We found no significant differences in primary productivity (ANOVA, $F=0.022$,

$p=0.978$) and community respiration (ANOVA, $F=0.148$, $p=0.865$) of the treatments versus the

controls (Figures 1A, 1B). All treatments showed substantial variation among replicates (Figure

1), with the most pronounced variability being present in respiration. Coefficients of variation

were highest for respiration in the controls (1.56, see Table 1) and lowest (0.021) in the

rhamnolipid 150 treatment. Rhamnolipid treated systems overall had the lowest variation for CR,

while the controls had the highest.

Ash free dry mass and C:N ratios

We found no significant difference among treatments and controls (ANOVA F=0.045,

p=0.957) for AFDM. The coefficients of variation, however, showed differences in variation

among the treatments (Table 1), with higher surfactant concentrations decreasing variation (e.g.,

rhamno 150 has greater variation than rhamno 300) (Figure 1C). Control variation was

comparable to both low dose surfactant variations. However, C (ANOVA F=18.36, p=0.0028)

and N (ANOVA F=5.471, p=0.044) content was significantly different by mass among

treatments (Figure 2A). Differences lie between the rhamno 300 and the control for both C

(Tukey HSD p=0.0063) and N (Tukey HSD p=0.041) content, and between SDS 300 and the

control for C (Tukey HSD p=0.0036) content.

We found no significant differences among treatments and controls for C:N ratios by

mass (Kruskal-Wallis p=0.193) present in the biofilms (Figure 2B). There were differences in

variation among treatments and controls (Table 1). Rhamnolipid treated flumes had lower

variation than either SDS treated flumes or the controls, however within each treatment a higher

surfactant concentration increased variation.

Bacterial community composition

Shannon-Wiener Diversity was significantly different among treatments (ANOVA

F=71.702, p<0.0001) with a higher diversity in the control flumes, second highest in the

rhamnolipid flumes, and lowest in the SDS flumes, while for Chao1 the differences were not

significant (ANOVA F=3.7748, p=0.087). Shannon diversity showed significant differences

among all three treatments tested (Tukey HSD rhamnolipid and control; p=0.043, SDS and

control; p<0.0001, SDS and rhamnolipid; p<0.001). Significance was driven by surfactant

presence and type, but not concentration. The control had the greatest bacterial diversity, while

both rhamnolipid treatments had the next highest, and the lowest belonged to both SDS

treatments (Figure 3A).

By testing beta diversity and community differences on the rarefied data, we identified

distinct groupings between all treatments and controls, as evident by the NMDS (Figure 3B). The

ANOSIM results confirmed the differences among both surfactants and the controls (R=0.9112,

p=0.001). Community differences were further identified by looking at only the taxa in the top

80% of all taxa present across all treatments, which was 18 taxa. We identified several genera

that were heavily dominant in rhamnolipid-treated flumes. These were: *Brevundimonas spp*.,

OTU005, *Pseudomonas,* and *Arthrobacter*. In SDS flumes we observed *Brevundimonas terrae,*

Flavobacterium spp. A, and in the control flumes *Exiguobacterium, Arenimonas,* and

Stenotrophomona. We also identified 32 genera only present in the controls, 5 there were

exclusive to rhamno 150, 6 in rhamno 300, 19 in SDS 150, and 7 in SDS 300.

Total OTUs within each treatment and OTUs within the top 80% of all taxa per treatment

were also calculated (Table 2). The control flumes had the greatest number of OTUs present, and

the greatest number of OTUs within their top 80% of taxa. The SDS and rhamnolipid flumes had

similar numbers of total OTUs present, but the SDS flumes had 80% of their total OTUs within

half the amount of taxa as the rhamnolipids. This indicated a large difference in evenness

between treatments, while abundance was fairly similar.

Extracellular Enzyme Activity

According to the DRA, EEA differed among all treatments and controls, with the

exception of the rhamnolipid treatments. The differences in DRA-distances were mainly caused

by the differences in EEA of leucine aminopeptidase and phosphatase, although the results were

not significant (ANOSIM F=2.9907, p=0.058). The effects of rhamno 300 and SDS 300 on

leucine aminopeptidase were significantly different (ANOVA F=5.914 p=0.038, Tukey HSD

p=0.032). The addition of rhamnolipid yielded a higher amount of the extracellular enzymes

when compared to the control and the addition of SDS reduced production. Rhamnolipid

additions also yielded a higher production of glucosidase, while SDS additions reduced

production (ANOVA F=5.741 p=0.0404, Tukey HSD p=0.034). While phosphatase did not have

any significant differences among treatments (ANOVA F=2.934 p=0.129), with the exception of

SDS 150 addition of any surfactant decreases the production of phosphatase when compared to

the control (Figure 4).

Discussion

Our experiment showed that exposure to biological and synthetic surfactants alters the

metabolic activity and the community composition of a mixed-species freshwater biofilm. The

effects of SDS were more pronounced than the effects of rhamnolipid. Our experiment is the first

study of our knowledge that examines the effects of these two surfactants on a multi-species

near-natural freshwater biofilm, which contrasts with the commonly performed single-species

studies. This study was also novel as it directly compared the effects of a synthetic and a

biological surfactant and explicitly studied for the first time the C:N ratios and C and N contents

of biofilms that are exposed to surfactants.

Primary Productivity, Respiration, Ash Free Dry Mass, and C:N ratios

Chemical exposure decreased biofilm respiratory activity in previous experiments,

including surfactant exposure (Simoes *et al.,* 2008) and pharmaceutical exposure (Rosi-Marshall

et al., 2013). Biofilm volume has also been reduced after a single surfactant exposure event, with

decreases of 26.5-98.1% occurring between 22-48 hours after exposure (Mireles II *et al.,* 2001;

Quinn *et al.,* 2013). However, previous work on surfactant effects have also shown that biofilms

demonstrate resistance and can recover full respiratory activity 12 hours after the initial exposure

(Simoes *et al.,* 2008) and biomass 5 hours after initial exposure (Nguyen *et al.,* 2020). Therefore,

the length of our experiment of 32 days may have been long enough where GPP, CR, and AFDM

could be stable over time, despite the permanent exposure.

Altered C:N ratios are indicative of changes in microbial productivity (where reduced

C:N ratios may yield greater productivity) (Li *et al.,* 2016) and biofilm formation (where lower

C:N ratios increase biofilm volume) (Thompson *et al.,* 2006). As we did not have any alterations

in C:N ratios, and no changes in productivity (GPP and CR) or growth (AFDM) among any

treatments or controls, our results support these studies. The addition of surfactants provided an

added source of C, however as there were consistent C:N ratios across all flumes it can be

assumed that our system was C limited. It is also likely that a thickened EPS matrix, which may

be used as a defense mechanism in biofilms in response to surfactant exposure, could be another

source of extra carbon (Jefferson, 2004; Zakaria and Dhar, 2020). In contrast, in systems where

C is not limiting, an increase in C inputs through surfactants may increase C:N ratios, and

therefore negatively affect biofilm productivity and growth.

Diversity and bacterial community composition

Our controls had taxa present that were minimally identified within the surfactant flumes.

These taxa included species (spp.) of *Exiguobacterium*, *Acinetobacter, Arenimonas,* and

Stenotrophomonas. The addition of either surfactant created an environment unsuitable for these

bacteria, demonstrating how surfactant exposure can alter biofilm community composition. The

control flumes had the highest amount of OTUs present (*n*=203) and the highest diversity, which

indicates that both the diversity and abundance of bacteria present in the biofilms decreases after

surfactant exposure.

Rhamnolipids are studied for their ability to reduce pre-existing biofilms and prevent

future biofilm growth with some success (Quinn *et al.,* 2013; Dusane *et al.,* 2010; De Rienzo *et*

al., 2015). Current research has also shown that rhamnolipid exposure on soil bacteria can shift

community composition, although many studies do not examine differences to the genus level

(Lu *et al.,* 2019; Wei *et al.,* 2020; Akbari *et al.,* 2021). Our study showed that, much like soil

communities, aquatic biofilms experience significant community changes after exposure to

rhamnolipids. Further examination into the bacterial genera indicate large increases in a

Brevundimonas spp., *Pseudomonas* spp., *Arthrobacter* spp., and OTU005 when compared to all

other treatments and to the controls. One OTU of *Arthrobacter* spp. was only present within the

rhamnolipid flumes, with the exception of one control flume that had low levels of that particular

OTU.

SDS caused the largest reduction in microbial diversity when compared with the other

treatments and controls. SDS has been researched and used for its ability to degrade biofilms in

applications where biofilms are undesirable (Izano *et al.,* 2007; Ueda *et al.,* 2019; Nguyen *et al.,*

2020). This is particularly due to their ability to bind to hydrophobic protein components,

introducing a negative charge to the structure which unfolds the bound protein (Gudiksen *et al.,*

2006; Fong and Yildiz, 2015). While rhamnolipids are also able to denature proteins, they are

considered weak binders, and typically have a reduced denaturing ability when compared to

many other surfactants (Otzen, 2017). This may explain why biofilms exposed to SDS had lower

diversity than those in the rhamnolipid treatments or the controls. SDS also had the lowest

amount of OTUs within the top 80% of taxa present in the treatments. We believe that the low

diversity, and community domination by few taxa, demonstrates that SDS exposure creates an

environment inhospitable to many potential specialist taxa. The few remaining and insensitive

taxa can then thrive as more resources are available. Following our findings, these taxa would

include *Brevundimonas terrae,* and an *Acinetobacter* and *Flavobacterium* spp. While the

rhamnolipid treated flumes also had less diversity than the controls, the rhamnolipid treatments

still appeared to favor more taxa and a higher diversity than the SDS treated flumes.

Extracellular Enzyme Activity

Our experiment showed that substantial changes in biofilm EEA can occur after

surfactant exposure. Although phosphatase activity, or the separation of phosphate groups from a

larger compound such as an acid or protein (Margalef *et al.,* 2017), generally decreased after

surfactant exposure, we did have similar levels of activity between the control and SDS 150

flumes. Examination of the bacterial communities within SDS 150ppm flumes and control

flumes show several parallel genera present that are not as common in the other systems,

including *Nocardioides*. Treatments were laid out randomly across our experimental flumes

within the experimental setup, and water was allocated to all systems from the same source,

which minimized potential experimental alterations by relative location. We thus assume that all

changes in community composition and EEA refer to the actual biofilm response to treatments.

The presence of a low concentration SDS allowed for the development of bacterial genera like

Nocardioides which typically have both acid and alkaline phosphatase (Xie *et al.,* 2017).

However, the addition of high concentration SDS, or rhamnolipid at either a high or low

concentration, reduced phosphatase activity.

Beta-glucosidase (rate-limiting enzyme that converts cellulase to glucose (Singh *et al.,*

2016)) and leucine aminopeptidase (enzyme that aids in the hydrolysis of leucine and other N-

terminal residues at the end of peptides and proteins (Matsui *et al.,* 2006)) activities were also

affected by surfactant exposure. Previous research has shown that rhamnolipids can act as an

enhancer for the enzyme beta-glucosidase (Zhang *et al.,* 2009; Yao *et al.,* 2020). Rhamnolipids

are able to protect proteins and limit the degradation of cellulase (Zhang *et al.,* 2009; Wang *et*

al., 2011). Yin *et al.* (2019) showed that the presence of rhamnolipid can increase genes related

to leucine and amino acid synthesis. Addition of rhamnolipid to our flumes likely protected

cellulase and leucine in the system which led to enhanced beta-glucosidase and leucine

aminopeptidase activity. Beta-glucosidase is also known for its ability to degrade *Pseudomonas*

aeruginosa, a rhamnolipid producing bacteria (Banar *et al.,* 2019; Pirlar *et al.,* 2020). The flumes

treated with rhamnolipid had a higher abundance of *P. aeruginosa* when compared with the other

flumes. Therefore, we assume that the higher beta-glucosidase levels within the rhamnolipid

flumes were likely caused by a combination of the presence of rhamnolipid to act as an enhancer,

and more abundant *P. aeruginosa.* However, we observed somewhat contrasting results in the

SDS treated flumes, where a decrease in enzyme activity occurred. SDS is an anionic surfactant

that can be used in denaturation of polypeptides (Bhuyan, 2009; Schlager *et al.,* 2012). SDS is

commonly used in an analysis called SDS-polyacrylamide gel electrophoresis (SDS-PAGE)

where an SDS infused gel is used to separate proteins and enzymes based on their molecular

weight. This analytical method has been used with experiments examining beta-glucosidase

previously (Laemmli, 1970; Bai *et al.,* 2013), which demonstrates that SDS is able to denature

beta-glucosidase. It is possible that the denaturing ability of SDS has degraded beta-glucosidase

and leucine aminopeptidase, leading to reduced enzyme activity, and likely also reduced biofilm

metabolic activity.

Conclusion

Our experiment successfully showed that both the biologic surfactant rhamnolipid and

the synthetic surfactant, SDS, can significantly affect a mixed freshwater microbial biofilm in

terms of community composition and metabolic activity. Rhamnolipids acted as an enhancer for

beta-glucosidase and leucine aminopeptidase, increasing metabolic activity within the biofilm,

while SDS likely denatured the enzymes and reduced metabolic activity within the biofilm. Both

surfactants decreased the diversity of the biofilm bacterial community. There were unique genera

in the biofilms of each treatment, including the controls which demonstrates that the presence of

either surfactant creates an environment that is inhospitable to sensitive taxa. These results

highlight two important areas of concern. First, rhamnolipids may not be a suitable alternative to

SDS for use in industrial and medical biofilm control because of its lower effectiveness. Second,

in natural ecosystems where biofilms are necessary for key ecosystem processes, such as

degradation of organic matter, the presence of these surfactants, (for example through the release

from WWTPs) may lead to negative effects on microbial biofilms. Altered biofilms may have

decreased nutrient uptake efficiency due to shifts in their metabolic activity. This could have

negative implications for the aquatic ecosystem as a whole, such as reduced denitrification.

Experimental Procedures

Experimental Set up

We set up 14 recirculating experimental flumes near Lunz Am See, Austria (following

Roche *et al.,* 2017). Each flume was 2.5m long and had an approximate slope of 0.8%. In each

flume, ~5L of water recirculated, which was renewed with unfiltered lake water twice per week.

We added rhamnolipid (R90-100G, Sigma-Aldrich, Darmstadt, Germany) and SDS (sodium

dodecyl sulphate, Sigma-Aldrich, Darmstadt, Germany) as treatments of 150 and 300 ppm to

three flumes each (hereafter referred to as either rhamno 150, rhamno 300, SDS 150, or SDS

300), while three flumes acted as controls. After the first week, 5 ug L^{-1} of Phosphorus in the

form of Na_2HPO_4 was also added to help growth. Surfactant concentrations were chosen to

represent the effective dose of rhamnolipid, as well as the double of that. Due to malfunction of

one flume, *n* for the rhamnolipid treatment was reduced to two. Treatments were randomly

placed among flumes, and light averaged 9.5 umol $m^{-2} s^{-1}$, with a light cycle of 12:12 and 18:6

hours of light versus dark early and later in the experiment., Each flume was equipped with

microscope slides (6) and clay tiles (8) to quantify biofilm growth.

We measured pH, DO (mg/L), water temperature (° C), and flow (by bucket

measurement) every other day to ensure stable conditions. Twice per week we analysed

dissolved organic carbon (DOC), ammonium (NH_4), NO_x, and PO_4 of flume water to ensure they

remained stable. In addition, water temperature was monitored by automatic loggers in drainage

tanks. After 32 days the experiment was stopped and biofilm samples retrieved from the flumes

for further analysis.

Sample Collection

Two microscope slides were removed from each flume and placed in 120mL Schott

bottles with oxygen sensors. Bottles were filled with filtered lake water and sealed to remove all

air bubbles. Three control bottles with no slides were also set up to test for potential activity

within the filtered lake water. All bottles were placed in an Imago 500 Controller Environmental

Chamber (Snijders Scientific B.V., Tilburg, Netherlands) using a 12:12 light cycle for 72 hours.

Oxygen measurements were taken at the start and end of each light and dark cycle to obtain data

on GPP and CR.

We determined AFDM by weighing 2 clay tiles per flume after drying (9h at 70 $^{\circ}$C), and

combustion (4h at 450 $^{\circ}$C). AFDM was obtained by subtracting the remaining ash weight from

the dry weight. Two tiles from SDS 150 flumes were damaged during drying, and AFDM could

not be determined. Biofilms covering two tiles per flume were then scraped into sterilized and

pre-weighed 2mL Eppendorf tubes using a sterile razor blade, lyophilised (-0.001mbar; -76 □)

and then analysed on a Thermo Scientific FLASH 2000 Elemental Analyser (Thermo Fisher

Scientific GmbH, Bremen, Germany) to quantify carbon (% C) and nitrogen (% N) content.

Quantities of C and N by mass were then examined in ratios to identify changes in ecological

stoichiometry (the relationship between elements, energy, and organisms and their interactions in

an ecological system (Sterner and Elser, 2002; Van de Waal *et al.,* 2018)).

EEA of β-D-1,4-glucosidase (glucosidase), phosphatase, and leucine-aminopeptidase

were measured in the biofilms following the method described in Romaní and Sabater (2001).

Four mL of sterile water was added to approximately 0.5 g of biofilm per flume that was scraped

from the tiles. The mixture was homogenized by shaking. EEA was measured via artificial

substrates: Glucosidase and phosphatase using 4-methylumbelliferyl (MUF)- substrates (MUF-β-

D-glucopyranoside and MUF-phosphate, respectively (Sigma-Aldrich, Darmstadt, Germany))

and leucine-aminopeptidase using a 7-amino-4-methylcoumarin (AMC)- substrate (L-leucine-

AMC, Sigma-Aldrich, Darmstadt, Germany). Artificial substrates were added to the homogenate

at a final concentration of MUF-glucosidase; 135 mg L^{-1}, MUF-phosphatase; 102.46 mg L^{-1}, and

AMC-leucine-aminopeptidase; 129.94 mg L^{-1} and incubated for 1 hour in the dark. Fluorescence

was measured at the beginning and end of the incubation with a platereader (Varioskan Flash,

Thermo Fischer Scientific, Vaanta, Finland) at 365/455 nm excitation/emission wavelengths for

the MUF-substrates and 380/440 excitation/emission wavelengths for the AMC-substrate.

Two microscope slides per flume were scraped using sterile procedures into a sterile

1.5mL microcentrifuge tube for 16s rRNA analysis. DNA was immediately extracted from

collected samples using the Dneasy Power Soil Kit (Qiagen GmbH, Hilden, Germany) following

the manufacturer's methodology. We included internal replicates per flume (*n*=2), and

experimental controls were used to control for any potential unaccounted sample contamination.

Extracted products were sent to LGC Biosearch Technologies (LGC Genomics GmbH, Berlin,

Germany). DNA was quantified using gel electrophoresis before amplifying the V3-V4 region of

the 16s rRNA gene with polymerase chain reactions using 341F and 785R primers. Resultant

DNA were sequenced using the Illumina Mi-Seq platform (300 bp paired-end reads) for

amplicon creation (sequencing results will be made available online after manuscript acceptance

and publication).

We used Mothur version 1.44.3 (Schloss *et al.,* 2009), and the Mothur MiSeq SOP

(Kozich *et al.,* 2013) with some modification, to analyze returned reads. A maximum

homopolymer of 8 was chosen due to the limitation of the reference database (SILVA version

132) with a 97% similarity and above required for identification of operational taxonomic units

(OTUs). UCHIME (Edgar *et al.,* 2011) was also used to identify and remove chimeric fragments.

Sequences that could not be directly identified through SILVA, were manually run through

NCBI's BLAST (Madden, 2002). We were able to identify all but one OTU with 97% similarity;

OTU005 is thus reported as 'unnamed' in this study.

 Data analysis

All data, except the 16s rRNA and C:N results, met assumptions of normality and

homogeneity of variance. We considered $p < 0.05$ as significant and analyzed data using R

version 4.0.3 (R Core Team, 2020) and the R packages car (Fox and Weisberg, 2019), vegan

(Oksanen *et al.,* 2020), ggplot2 (Wickham, 2016), and entropart (Marcon and Herault, 2015).

Non metric multidimensional scaling analyses (NMDS) were run using the PAST 3 software

(Hammer *et al.,* 2001). Due to replication error early in the experiment only the controls and 300

ppm concentration variables were used in statistical difference testing. All GPP and CR data

were normalized to mg O_2 in 120mL per mg of biofilm before being analyzed. C:N data was

normalized to mass (mg C/N per mg biofilm). A one way analysis of variance (ANOVA)

analysis was used to test for significant differences among treatments and the controls for the

GPP, CR, and AFDM. A Kruskal-Wallis test was used to test for significant differences among

the C:N ratios. Coefficients of variation were also calculated for individual flumes for each

analysis.

Changes in EEA were analysed through both a distance based redundancy analysis

(DRA) and three one-way ANOVAs, one per extracellular enzyme. All the data was used within

the DRA and only the maximum concentrations of each chemical and control were used for

analysis in the ANOVA. Where appropriate, we applied a Tukey HSD post-hoc test to the

ANOVA output.

Bacterial OTUs were analyzed for alpha and beta diversity as well as differences in the

overall bacterial community composition. We used the Chao1 (Chao, 1984) and the Shannon-

Wiener Diversity Indices to quantify bacterial Alpha diversity (Jost, 2007). Chao 1 and Shannon-

Wiener results were then analyzed using an ANOVA followed by a Tukey HSD post-hoc test to

identify differences in diversity among treatments. Following this, all singletons and doubletons

were removed and the data was rarefied based on the smallest number of sequences identified

within a sample. Beta diversity was then analysed using a non-metric multidimensional scaling

analysis (NMDS) on all data present to identify differences in the biofilm bacterial community

among treatments and controls. An analysis of similarities (ANOSIM) was performed with 999

permutations to identify if these differences were significant. Families and genera were

compared graphically, focusing on the top 80% of OTUs identified within the biofilms.

significant differences).

Acknowledgements

We thank Gertraud Steniczka and Hermann Hofreiter for help in the laboratory and with setting

up the experiment. SG is funded by an Ulster University Vice Chancellors Doctoral Research

Fellowship, and received additional support through an Ulster University Broadening Horizons

Travel Bursary. Analytical costs were partly supported by the HYDRO-DIVERSITY project

funded by the Environmental Systems Sciences Program of the Austrian Academy of Sciences

(ÖAW) to JS, and core funding of the AFBI Aquatic Chemistry Laboratory (BH).

Conflict of Interest

The authors declare no conflict of interest.

References

Akbari, A., Kasprzyk, A., Galvez, R., and Ghoshal, S. (2021) A rhamnolipid biosurfactant

increased bacterial population size but hindered hydrocarbon biodegradation in

weathered contaminated soils. *Sci Total Environ* **778:** 145441.

Aleksic, I., Petkovic, M., Jovanovic, M., Milivojevic, D., Vasiljevic, B., Nikodinovic-Runic, J.,

and Senerovic, L. (2017) Anti-biofilm Properties of Bacterial Di-Rhamnolipids and Their

Semi-Synthetic Amide Derivatives. *Frontiers in Microbiology* **8:** 2454.

Bai, H., Wang, H., Sun, J., Irfan, M., Han, M., Huang, Y. *et al.* (2013) Production, purification

and characterization of novel beta glucosidase from newly isolated Penicillium

simplicissimum H-11 in submerged fermentation. *EXCLI journal* **12:** 528-540.

Banar, M., Emaneini, M., Beigverdi, R., Fanaei Pirlar, R., Node Farahani, N., van Leeuwen, W.

B., and Jabalameli, F. (2019) The efficacy of lyticase and β-glucosidase enzymes on

biofilm degradation of Pseudomonas aeruginosa strains with different gene profiles. *BMC*

Microbiol **19:** 291-9.

Battin, T. J., Besemer, K., Bengtsson, M. M., Romani, A. M., and Packmann, A.I. (2016) The

ecology and biogeochemistry of stream biofilms. *Nature reviews. Microbiology* **14:** 251-

263.

Bhuyan, A. K. (2010) On the mechanism of SDS-induced protein denaturation. *Biopolymers* **93:**

186-199.

Bowler, P., Murphy, C., and Wolcott, R. (2020) Biofilm exacerbates antibiotic resistance: Is this
a current oversight in antimicrobial stewardship? *Antimicrobial Resistance* and *Infection*
Control **9:** 162.

Cao, C., Cao, Z., Yu, P., and Zhao, Y. (2020) Genome-wide identification for genes involved in
sodium dodecyl sulfate toxicity in Saccharomyces cerevisiae. *BMC Microbiology* **20:** 34.

Chao, A. (1984) Nonparametric Estimation of the Number of Classes in a Population.
Scandinavian Journal of Statistics **11:** 265-270.

Cui, B., Liu, X., Yang, Q., Li, J., Zhou, X., and Peng, Y. (2017) Achieving partial denitrification
through control of biofilm structure during biofilm growth in denitrifying biofilter.
Bioresour Technol **238:** 223-231.

Dang, H., and Lovell, C. R. (2016) Microbial surface colonization and biofilm development in
marine environments. *Microbiology and molecular biology reviews* **80:** 91-138.

Decho, A. W. (2000) Microbial biofilms in intertidal systems: an overview. *Continental shelf*
research **20:** 1257-1273.

Desai, J. D., and Banat, I. M. (1997) Microbial production of surfactants and their commercial
potential. *Microbiology and Molecular Biology Reviews* **61:** 47-64.

Díaz De Rienzo, M. A., Stevenson, P. S., Marchant, R., and Banat, I. M. (2016) Pseudomonas
aeruginosa biofilm disruption using microbial surfactants. *J Appl Microbiol* **120:** 868-
876.

Díaz De Rienzo, Mayri A, Banat, I. M., Dolman, B., Winterburn, J., and Martin, P. J. (2015)

Sophorolipid biosurfactants: Possible uses as antibacterial and antibiofilm agent. *New*

Biotechnol **32:** 720-726.

Dodds, W. K., Evans-White, M., Gerlanc, N. M., Gray, L., Gudder, D. A., Kemp, M. J. *et al.*

(2000) Quantification of the nitrogen cycle in a prairie stream. *Ecosystems* **3:** 574-589.

Drury, B., Scott, J., Rosi-Marshall, E. J., and Kelly, J. J. (2013) Triclosan exposure increases

triclosan resistance and influences taxonomic composition of benthic bacterial

communities. *Environ Sci Technol* **47:** 8923-8930.

Dusane, D. H., Zinjarde, S. S., Venugopalan, V. P., Mclean, R. J., Weber, M. M., and Rahman,

P. K. S. M. (2010) Quorum sensing: implications on Rhamnolipid biosurfactant

production. *Biotechnol Genet Eng Rev* **27:** 159-184.

Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., and Knight, R. (2011) UCHIME improves

sensitivity and speed of chimera detection. *Bioinformatics* **27:** 2194-2200.

Fanaei Pirlar, R., Emaneini, M., Beigverdi, R., Banar, M., van Leeuwen, W.,B., and Jabalameli,

F. (2020) Combinatorial effects of antibiotics and enzymes against dual-species

Staphylococcus aureus and Pseudomonas aeruginosa biofilms in the wound-like medium.

PLOS ONE **15:** e0235093.

Flemming, H., Wingender, J., Szewzyk, U., Steinberg, P., Rice, S. A., and Kjelleberg, S. (2016)

Biofilms: an emergent form of bacterial life. *Nat Rev Microbiol* **14:** 563-575.

Fong, J. N. C., and Yildiz, F. H. (2015) Biofilm Matrix Proteins. *Microbiol Spectr* **3:**

10.1128/microbiolspec.MB-2014.

Fox, J., and Weisberg S. (2019) An R Companion to Applied Regression, Third Edition.

Thousand Oaks California, USA: Sage.

Franzetti, A., Gandolfi, I., Raimondi, C., Bestetti, G., Banat, I. M., Smyth, T. J. *et al.* (2012)

Environmental fate, toxicity, characteristics and potential applications of novel

bioemulsifiers produced by Variovorax paradoxus 7bCT5. *Bioresour Technol* **108:** 245-

251.

Gomez, V., Ferreres, L., Pocurull, E., and Borrull, F. (2011) Determination of non-ionic and

anionic surfactants in environmental water matrices. *Talanta* **84:** 859-866.

Gudiksen, K. L., Gitlin, I., and Whitesides, G. M. (2006) Differentiation of proteins based on

characteristic patterns of association and denaturation in solutions of SDS. *Proc Natl*

Acad Sci USA **103:** 7968.

Hammer, Ø, Harper, D. A. T., and Ryan, P. D. (2001) PAST: Paleontological statistics software

package for education and data analysis. *Palaeontol Electron* **4:**

Harjung, A., Attermeyer, K., Aigner, V., Krlovic, N., Steniczka, G., Švecová, H. *et al.* (2020)

High Anthropogenic Organic Matter Inputs during a Festival Increase River Heterotrophy

and Refractory Carbon Load. *Environ Sci Technol* **54:** 10039-10048.

Hauthal, H. G. (2004) CESIO 2004: Dynamic surfactants and nanostructured surfaces for an

innovative industry. *SÖFW Journal* **130:** 3-17.

Hay, A. G., Dees, P. M., and Sayler, G. S. (2001) Growth of a bacterial consortium on triclosan.

FEMS Microbiol Ecol **36:** 105-112.

Izano, E. A., Wang, H., Ragunath, C., Ramasubbu, N., and Kaplan, J. B. (2007) Detachment and

Killing of Aggregatibacter actinomycetemcomitans Biofilms by Dispersin B and SDS. *J*

Dent Res **86:** 618-622.

Jefferson, K. K. (2004) What drives bacteria to produce a biofilm? *FEMS Microbiol Lett* **236:**

163-173.

Jost, L. (2007) Partitioning diversity into independent alpha and beta components. *Ecology* **88:**

2427-2439.

Ke, Y., Chen, J., Hu, X., Tong, T., Huang, J., and Xie, S. (2020) Emerging perfluoroalkyl

substance impacts soil microbial community and ammonia oxidation. *Environ Pollut*

257: 113615.

Kozich James, J., Westcott Sarah, L., Baxter Nielson, T., Highlander Sarah, K., and Schloss

Patrick, D. (2013) Development of a Dual-Index Sequencing Strategy and Curation

Pipeline for Analyzing Amplicon Sequence Data on the MiSeq Illumina Sequencing

Platform. *Appl Environ Microbiol* **79:** 5112-5120.

Krishnaswamy Muthusamy, Subbuchettiar Gopalakrishnan, Thiengungal Kochupappy Ravi, and

Panchaksharam Sivachidambaram. (2008) Biosurfactants: Properties, commercial

production and application. *Curr Sci (Bangalore)* **94:** 736-747.

Kroukamp, O., and Wolfaardt, G. M. CO□ Production as an Indicator of Biofilm Metabolism.

Appl Environ Microbiol **75:** 4391-4397.

Kruszelnicka, I., Ginter-Kramarczyk, D., Wyrwas, B., and Idkowiak, J. (2019) Evaluation of

surfactant removal efficiency in selected domestic wastewater treatment plants in Poland.

J Environ Health Sci Eng **17:** 1257-1264.

Kumar, A., Yadav, C. B., and Bajpeyee, A. K. (2020) Glucose, protein, cholesterol and

immunohaematological alterations with acute (SDS) Sodium Dodecyl Sulfate toxicity in

Clarias batrachus. *J Exp Zoo (India)* **23:** 401-404.

Kusi, J., Scheuerman, P. R., and Maier, K. J. (2020) Emerging environmental contaminants

(silver nanoparticles) altered the catabolic capability and metabolic fingerprinting of

microbial communities. *Aquat Toxicol* **228:** 105633.

Laemmli, U. K. (1970) Cleavage of structural proteins during the assembly of the head of

bacteriophage T4. *Nature* **227:** 680-685.

Landa, A. S., De Belt-Gritter, B. v., Der Mei, Henny C. van, and Busscher, H. J. (1999)

Recalcitrance of Streptococcus mutans biofilms towards detergent-stimulated

detachment. *Eur J Oral Sci* **107:** 236-243.

Lara-Martín, P. A., González-Mazo, E., and Brownawell, B. J. (2012) Environmental analysis of

alcohol ethoxylates and nonylphenol ethoxylate metabolites by ultra-performance liquid

chromatography–tandem mass spectrometry. *Anal Bioanal Chem* **402:** 2359-2368.

Lara-Martín, P. A., González-Mazo, E., Petrovic, M., Barceló, D., and Brownawell, B. J. (2014)

Occurrence, distribution and partitioning of nonionic surfactants and pharmaceuticals in

the urbanized Long Island Sound Estuary (NY). *Mar Pollut Bull* **85:** 710-719.

Li, Y., Wu, J., Shen, J., Liu, S., Wang, C., Chen, D. *et al.* (2016) Soil microbial C:N ratio is a

robust indicator of soil productivity for paddy fields. *Sci Rep* **6:** 35266.

Lima, T. M. S., Procópio, L. C., Brandão, F. D., Leão, B. A., Tótola, M. R., and Borges, A. C.

(2011) Evaluation of bacterial surfactant toxicity towards petroleum degrading

microorganisms. *Bioresour Technol* **102:** 2957-2964.

Lindsay, D., and von Holy, A. (2006) Bacterial biofilms within the clinical setting: what

healthcare professionals should know. *J Hosp Infect* **64:** 313-325.

Lu, L., Zhang, J., and Peng, C. (2019) Shift of Soil Polycyclic Aromatic Hydrocarbons (PAHs)

Dissipation Pattern and Microbial Community Composition due to Rhamnolipid

Supplementation. *Water Air Soil Pollut* **230:** 107.

Lyon, D. R., and Ziegler, S. E. (2009a) Carbon cycling within epilithic biofilm communities

across a nutrient gradient of headwater streams. *Limnol Oceanogr* **54:** 439-449.

Lyon, D. R., and Ziegler, S. E. (2009b) Carbon cycling within epilithic biofilm communities

across a nutrient gradient of headwater streams. *Limnol Oceanogr* **54:** 439-449.

Madden, T. (2002) Chapter 16: The BLAST sequence analysis tool. In *The NCBI handbook.* J.

McEntyre, and J. Ostell (eds). Bethesda, National Center for Biotechnology Information.

Marcon, E., and Herault, B. (2015) Entropart: An R Package to Measure and Partition Diversity.

J Stat Softw **67**: 1-26.

Margalef, O., Sardans, J., Fernández-Martínez, M., Molowny-Horas, R., Janssens, I. A., Ciais, P.

et al. (2017) Global patterns of phosphatase activity in natural soils. *Sci Rep* **7:** 1337.

Mariani, L., De Pascale, D., Faraponova, O., Tornambè, A., Sarni, A., Giuliani, S. *et al.* (2006)

The use of a test battery in marine ecotoxicology: The acute toxicity of sodium dodecyl

sulfate. *Environ Toxicol* **13:** 373-379.

Matsui, M., Fowler, J. H., and Walling, L. L. (2006) Leucine aminopeptidases: diversity in

structure and function. *Biol Chem* **387:** 1535-1544.

McClean, P., and Hunter, W. R. (2020) 17α-ethynylestradiol (EE2) limits the impact of

ibuprofen upon respiration by streambed biofilms in a sub-urban stream. *Environ Sci*

Pollut Res **27:** 37149-37154.

Messina, C. M., Faggio, C., Laudicella, V. A., Sanfilippo, M., Trischitta, F., and Santulli, A.

(2014) Effect of sodium dodecyl sulfate (SDS) on stress response in the Mediterranean

mussel (Mytilus Galloprovincialis): Regulatory volume decrease (Rvd) and modulation

of biochemical markers related to oxidative stress. *Aquat Toxicol* **157:** 94-100.

Milovanovic, M., Arsenijevic, A., Milovanovic, J., Kanjevac, T., and Arsenijevic, N. (2017)

Chapter 14 : Nanoparticles in antiviral therapy. In *Antimicrobial nanoarchitectonics.*

Elsevier Inc, pp. 383-410.

Mireles II, J. R., Toguchi, A., and Harshey, R. M. (2001) Salmonella enterica Serovar

Typhimurium Swarming Mutants with Altered Biofilm-Forming Abilities: Surfactin

Inhibits Biofilm Formation. *J Bacteriol* **183:** 5848-5854.

Mohan, P. K., Nakhla, G., and Yanful, E. K. (2006) Biokinetics of biodegradation of surfactants

under aerobic, anoxic and anaerobic conditions. *Water Res (Oxford)* **40:** 533-540.

Mungray, A. K., and Kumar, P. (2009) Fate of linear alkylbenzene sulfonates in the

environment: A review. *Int Biodeterior Biodegrad* **63:** 981-987.

Nguyen, B. V. G., Nagakubo, T., Toyofuku, M., Nomura, N., and Utada, A. S. (2020) Synergy

between Sophorolipid Biosurfactant and SDS Increases the Efficiency of P. aeruginosa

Biofilm Disruption. *Langmuir* **36:** 6411-6420.

Oksanen, J.F., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., *et al.* (2020)

Vegan: Community Ecology Package. R package version 2.5-7.

Otzen, D. E. (2017) Biosurfactants and surfactants interacting with membranes and proteins:

Same but different? *Biochim Biophys Acta Biomembr* **1859:** 639-649.

Paul, B. J., Duthie, H. C., and Taylor, W. D. (1991) Nutrient Cycling by Biofilms in Running

Waters of Differing Nutrient Status. *J N Amer Benthol Soc* **10:** 31-41.

Qu, X., Ren, Z., Zhang, H., Zhang, M., Zhang, Y., Liu, X., and Peng, W. (2017) Influences of

anthropogenic land use on microbial community structure and functional potentials of

stream benthic biofilms. *Sci Rep* **7:** 15117.

Quinn, G., Maloy, A., Banat, M., and Banat, I. (2013) A Comparison of Effects of Broad-

Spectrum Antibiotics and Biosurfactants on Established Bacterial Biofilms. *Curr*

Microbiol **67:** 614-623.

R Core Team (2020) R: A language and environment for statistical computing. R Foundation for

Statistical Computing. Vienna, Austria. URL https://www.R-project.org/.

Ramos da Silva, A., Manresa, M. Á, Pinazo, A., García, M. T., and Pérez, L. (2019)

Rhamnolipids functionalized with basic amino acids: Synthesis, aggregation behavior,

antibacterial activity and biodegradation studies. *Colloids Surf B* **181:** 234-243.

Rigotti, R. T., Corrêa, J. A. F., Maia, N. J. L., Cesaro, G., Rosa, E. A. R., Macedo, Renata

Ernlund Freitas de, and Luciano, F. B. (2017) Combination of natural antimicrobials and

sodium dodecyl sulfate for disruption of biofilms formed by contaminant bacteria

isolated from sugarcane mills. *Innov Food Sci Emerg Technol* **41:** 26-33.

Robinson, C., Strafford, S., Rees, G., Brookes, S. J., Kirkham, J., Shore, R. C. *et al.* (2006)

Plaque biofilms: The effect of chemical environment on natural human plaque biofilm

architecture. *Arch Oral Biol* **51:** 1006-1014.

Roche, K. R., Drummond, J. D., Boano, F., Packman, A. I., Battin, T. J., and Hunter, W. R.

(2017) Benthic biofilm controls on fine particle dynamics in streams. *Water Resour Res*

53: 222-236.

Romaní, A. M., Guasch, H., Muñoz, I., Ruana, J., Vilalta, E., Schwartz, T. *et al.* (2004) Biofilm

Structure and Function and Possible Implications for Riverine DOC Dynamics. *Microb*

Ecol **47:** 316-328.

Romaní, A. M., and Sabater, S. (2001) Structure and activity of rock and sand biofilms in a

mediterranean stream. *Ecology (Durham)* **82:** 3232-3245.

Rosi-Marshall, E., Kincaid, D. W., Bechtold, H. A., Royer, T. V., Rojas, M., and Kelly, J. J.

(2013) Pharmaceuticals suppress algal growth and microbial respiration and alter

bacterial communities in stream biofilms. *Ecol Appl* **23:** 583-593.

Schaider, L. A., Rudel, R. A., Ackerman, J. M., Dunagan, S. C., and Brody, J. G. (2014)

Pharmaceuticals, perfluorosurfactants, and other organic wastewater compounds in public

drinking water wells in a shallow sand and gravel aquifer. *Sci Total Environ* **468-469:**

384-393.

Schlager, B., Straessle, A., and Hafen, E. (2012) Use of anionic denaturing detergents to purify

insoluble proteins after overexpression. *BMC Biotechnol* **12:** 95.

Schloss Patrick, D., Westcott Sarah, L., Thomas, R., Hall Justine, R., Martin, H., Hollister Emily,

B. *et al.* (2009) Introducing mothur: Open-Source, Platform-Independent, Community-

Supported Software for Describing and Comparing Microbial Communities. *Appl*

Environ Microbiol **75:** 7537-7541.

Schmitz, K. S. (2018) Chapter 4: Life science. In *Physical chemistry.* Boston, Elsevier, pp. 755-

832.

Schreiberova, O., Hedbavna, P., Aejkova, A., JirkA[macron], V., and Masak, J. (2012) Effect of

surfactants on the biofilm of Rhodococcus erythropolis, a potent degrader of aromatic

pollutants. *New Biotechnol* **30:** 62-68.

Seghers, D., Verthé, K., Reheul, D., Bulcke, R., Siciliano, S. D., Verstraete, W., and Top, E. M.

(2003) Effect of long-term herbicide applications on the bacterial community structure

and function in an agricultural soil. *FEMS Microbiol Ecol* **46:** 139-146.

Simões, M., Simões, L. C., Pereira, M. O., and Vieira, M. J. (2008) Sodium dodecyl sulfate

allows the persistence and recovery of biofilms of Pseudomonas fluorescens formed

under different hydrodynamic conditions. *Biofouling* **24:** 35-44.

Singh, G., Verma, A. K., and Kumar, V. (2015) Catalytic properties, functional attributes and

industrial applications of β-glucosidases. *3 Biotech* **6:** 3.

Sloup, R. E., Cieza, R. J., Needle, D. B., Abramovitch, R. B., Torres, A. G., and Waters, C. M.

(2016) Polysorbates prevent biofilm formation and pathogenesis of Escherichia coli

O104:H4. *Biofouling* **32:** 1131-1140.

Srey, S., Jahid, I. K., and Ha, S. (2013) Biofilm formation in food industries: A food safety

concern. *Food Control* **31:** 572-585.

Sterner, R. W., and Elser, J. J. (2017) *Ecological stoichiometry.* Princeton university press,

Sumpono, Perotti, P., Belan, A., Forestier, C., Lavedrine, B., and Bohatier, J. (2003) Effect of

Diuron on aquatic bacteria in laboratory-scale wastewater treatment ponds with special

reference to Aeromonas species studied by colony hybridization. *Chemosphere* **50:** 445-

455.

Tambone, E., Bonomi, E., Ghensi, P., Maniglio, D., Ceresa, C., Agostinacchio, F. *et al.* (2021)

Rhamnolipid coating reduces microbial biofilm formation on titanium implants: an in

vitro study. *BMC Oral Health* **21:** 49.

Thompson, L. J., Gray, V., Lindsay, D., and Von Holy, A. (2006) Carbon   :   nitrogen   :

phosphorus ratios influence biofilm formation by Enterobacter cloacae and Citrobacter

freundii. *J Appl Microbiol* **101:** 1105-1113.

Thornton, I., Butler, D., Docx, P., Hession, M., Makropoulos, C., McMullen, M. *et al.* (2001)

Pollutants in Urban Waste Water and Sewage Sludge. *European Commission*

Ueda, Y., Mashima, K., Miyazaki, M., Hara, S., Takata, T., Kamimura, H. *et al.* (2019)

Inhibitory effects of polysorbate 80 on MRSA biofilm formed on different substrates

including dermal tissue. *Sci Rep* **9:** 3128.

Van de Waal, D. B, Elser, J. J., Martiny, A. C., Sterner, R. W., and Cotner, J. B. (2018) Editorial:

Progress in Ecological Stoichiometry. *Front Microbiol* **9:** 1957.

Waaler, S. M., Rölla, G., Skjörland, K. K., and Ogaard, B. (1993) Effects of oral rinsing with

triclosan and sodium lauryl sulfate on dental plaque formation: a pilot study. *Scand J*

Dent Res **101:** 192-195.

Wang, H., Fan, B., Li, C., Liu, S., and Li, M. (2011) Effects of rhamnolipid on the cellulase and

xylanase in hydrolysis of wheat straw. *Bioresour Technol* **102:** 6515-6521.

Wei, Z., Wang, J. J., Gaston, L. A., Li, J., Fultz, L. M., DeLaune, R. D., and Dodla, S. K. (2020)

Remediation of crude oil-contaminated coastal marsh soil: Integrated effect of biochar,

rhamnolipid biosurfactant and nitrogen application. *J Hazard Mater* **396:** 122595.

Wickham, H. (2016) Ggplot2: Elegant Graphics for Data Analysis. New York, USA: Springer-

Verlag.

Xie, F., Yang, Y., Ma, H., Quan, S., Yue, D., Sun, Y. *et al.* Nocardioides phosphati sp. nov., an

actinomycete isolated from a phosphate mine. *Int J Syst Evol* **67:** 1522-1528.

Yao, X., Xiao, L., Liu, H., Zhou, M., Zhou, Z., Ju, X., and Li, L. (2020) Rhamnolipid enhanced

beta-glucosidase from Paenibacillus sp. LLZ1 for in situ lignocellulose saccharification

in ionic liquids. *Biomass Convers and Biorefin*

Yin, Y., Gu, J., Wang, X., Zhang, Y., Zheng, W., Chen, R., and Wang, X. (2019) Effects of

rhamnolipid and Tween-80 on cellulase activities and metabolic functions of the bacterial

community during chicken manure composting. *Bioresour Technol* **288:** 121507.

Zakaria, B. S., and Dhar, B. R. (2020) Changes in syntrophic microbial communities, EPS

matrix, and gene-expression patterns in biofilm anode in response to silver nanoparticles

exposure. *Sci Total Environ* **734:** 139395.

Zhang, Q., He, G., Wang, J., Cai, W., and Xu, Y. (2009) Mechanisms of the stimulatory effects

of rhamnolipid biosurfactant on rice straw hydrolysis. *Appl Energy* **86:** S233-S237.

Zhang, S., Pang, S., Wang, P., Wang, C., Guo, C., Addo, F. G., and Li, Y. (2016) Responses of

bacterial community structure and denitrifying bacteria in biofilm to submerged

macrophytes and nitrate. *Sci Rep* **6:** 36178.

Table and Figure Legends

Table 1 Coefficients of variation for AFDM, GPP, and CR data. Data collected from samples of

biofilm grown in either the surfactant SDS (sodium dodecyl sulphate) or rhamnolipid for 32

days.

Table 2 Biofilm OTUs per surfactant treatment and their dominant genera. Biofilms were grown

in either SDS (sodium dodecyl sulphate) or rhamnolipid for 32 days.

Figure 1 GPP (A), CR (B) and AFDM (C) of microbial biofilms exposed to SDS (sodium

dodecyl sulphate) and rhamnolipid for 32 days at 150 and 300ppm. Controls are shown in brown,

rhamno 150 in dark green, rhamno 300 in light green, SDS 150 in dark blue, and SDS 300 in

light blue.

Figure 2 Carbon (A), Nitrogen (B) and C:N ratios (C) of microbial biofilms exposed to SDS

(sodium dodecyl sulphate) and rhamnolipid for 32 days at 150 and 300ppm. Controls are shown

in brown, rhamno 150 in dark green, rhamno 300 in light green, SDS 150 in dark blue, and SDS

300 in light blue.

Figure 3 Biofilm bacterial diversity from 16s rRNA sequencing of biofilm exposed to SDS

(sodium dodecyl sulphate) or rhamnolipid after 32 days at 150 and 300ppm. A) shows Shannon

Diversity, B) shows an NMDS depicting the community differences in treatments when

examining all OTUs present within each sample after rarefication. Controls are shown in brown,

rhamno 150 in dark green, rhamno 300 in light green, SDS 150 in dark blue, and SDS 300 in

light blue.

Figure 4 EEA of phosphatase (A), glucosidase (B), and leucine aminopeptidase (C) of biofilms

after exposure to either SDS (sodium dodecyl sulphate) or rhamnolipid over 32 days at 150 and

300ppm. Controls are shown in brown, rhamno 150 in dark green, rhamno 300 in light green,

SDS 150 in dark blue, and SDS 300 in light blue (*indicates significant differences).

Table 1

Treatment	Ash free dry mass	Gross primary productivity	CR	C:N
SDS 150	0.70	0.67	.92	0.373
SDS 300	0.11	0.626	1.09	0.643
Rhamno 150	0.60	0.34	0.021	0.0875
Rhamno 300	0.32	0.52	0.59	0.0698
Control	0.59	0.61	1.56	0.569

Table 2

Surfactant	Conc. (ppm)	Total OTUs present	OTUs in top 80% per treatment	OTUs lost compared to control	OTUs gained compared to control	Dominant genera in top 80% of all taxa present						
Rhamnolipid	150	145	10	58	25	*Brevundimonas*	*Otu005*	*Pseudomonas*	*Arthrobacter*	*Flavobacterium B*	*Prosthecobacter*	*Nocardioides*
Rhamnolipid	300	159	14	44	36	*Brevundimonas*	*Otu005*	*Pseudomonas*	*Arthrobacter*	*Flavobacterium B*	*Prosthecobacter*	
SDS	150	176	7	27	37	*Brevundimonas terrae*	*Acinetobacter*	*Flavobacterium A*	*Paucibacter*	*Flavobacterium B*	*Prosthecobacter*	
SDS	300	130	6	73	23	*Brevundimonas terrae*	*Acinetobacter*	*Flavobacterium A*	*Prosthecobacter*	*Flavobacterium B*	*Prosthecobacter*	*Sphingomonas*
Control	0	203	15	--	--	*Exiguobacterium*	*Acinetobacter*	*Arenimonas*	*Stenotrophomonas*	*Nocardioides*		

827 **Figure 1**

845 **Figure 2**

Figure 3

Figure 4