Synthetic and biological surfactant effects on freshwater biofilm community composition

and metabolic activity

Authors: Stephanie P. Gill[1], William R. Hunter[2], Laura E. Coulson[3,4], Ibrahim M. Banat[5], and

Jakob Schelker[3,6]

[1]Department of Geography and Environmental Studies, Ulster University, Coleraine, N. Ireland,

UK

[2]Fisheries and Aquatic Ecosystems Branch, Agri-Food and Biosciences Institute, Belfast, N.

Ireland, UK

[3]WasserCluster Lunz, Lunz Am See, Austria

[4]Institute of Hydrobiology and Aquatic Ecosystem Management, University of Natural

Resources and Life Sciences, Vienna, Austria

[5]School of Biomedical Sciences, Ulster University, Coleraine, N. Ireland, UK

[6]Department of Functional and Evolutionary Ecology, University of Vienna, Vienna, Austria

*Correspondence: Stephanie P. Gill

16   School of Geography and Environmental Science
17   Ulster University, Coleraine Campus
18   BT52 1SA. United Kingdom
19   Email: gill-S5@ulster.ac.uk
20   Telephone: 07541-036849
21

22 Keywords (6): Biofilm, surfactants, freshwater, metabolic, sequencing, metagenomics

23 Acknowledgements: We thank Gertraud Steniczka and Hermann Hofreiter for help in the

24 laboratory and with setting up the experiment.

Abstract

Surfactants are used to control microbial biofilms in industrial and medical settings. Their known toxicity on aquatic biota and their longevity in the environment has encouraged research on biodegradable alternatives such as rhamnolipids. While previous research has investigated the effects of biological surfactants on single species biofilms, there remains a lack of information regarding the effects of synthetic and biological surfactants in freshwater ecosystems. We conducted a mesocosm experiment to test how the surfactant sodium dodecyl sulphate (SDS) and the biological surfactant rhamnolipid altered community composition and metabolic activity of freshwater biofilms. Biofilms were cultured in the flumes using lake water from Lake Lunz in Austria, under high (300 ppm) and low (150 ppm) concentrations of either surfactant over a four-week period. Our results show that both surfactants significantly affected microbial diversity. Up to 36% of microbial operational taxonomic units were lost after surfactant exposure. Rhamnolipid exposure also increased the production of the extracellular enzymes, leucine aminopeptidase and glucosidase, while SDS exposure reduced leucine aminopeptidase and glucosidase. This study demonstrates that exposure of freshwater biofilms to chemical and biological surfactants caused a reduction of microbial diversity and changes in biofilm metabolism, exemplified by shifts in extracellular enzyme activities.

Key Points

1. Microbial biofilm diversity decreased significantly after surfactant exposure.
2. Exposure to either surfactant altered extracellular enzyme activity.
3. Overall metabolic activity was not altered, suggesting functional redundancy.

Introduction

In aquatic ecosystems, microorganisms live preferentially in microbial biofilms and act as major drivers of the biogeochemical cycling of carbon, nitrogen, and other nutrients (Paul *et al.* 1991; Decho 2000; Dang and Lovell 2016; Flemming *et al.* 2016; Battin *et al.* 2016). In streams and rivers, microbial biofilms play an important role in the carbon cycle (Lyon and Ziegler 2009; Qu *et al.* 2017). Microbial biofilms resident in fluvial sediments can also substantially reduce inorganic nitrogen (N) loads in N-rich aquatic ecosystems through denitrification (Zhang *et al.* 2016; Cui *et al.* 2017). In N-limited aquatic systems, they may help to facilitate the transfer of N to higher trophic levels (Dodds *et al.* 2000).

While microbial biofilms play an important ecological role, they are of concern in industrial and medical applications. For example, biofilms can facilitate the growth of antibiotic resistant bacteria (Lindsay and von Holy 2006; Srey *et al.* 2013; Bowler *et al.* 2020). Synthetic surfactants, such as sodium dodecyl sulphate (SDS), are widely used in personal care and cleaning products. They reduce biofilm growth in a wide range of industrial, medical, and environmental contexts (Waaler *et al.* 1993; Landa *et al.* 1999; Rigotti *et al.* 2017; Sloup *et al.* 2016; Ueda *et al.* 2019). Surfactants are amphiphilic chemicals that reduce surface tension within a liquid, sometimes through the formation of micelles. Micelles occur when molecules form an aggregate with hydrophobic tails on the inside (Schmitz 2018) and hydrophilic heads on the outside of an amphiphilic compound (Milovanovic *et al.* 2017).

Research on the effects of surfactants on microbial biofilms has primarily focused on single-species biofilms or those commonly found in medical and industrial environments. Direct applications of the anionic synthetic surfactant SDS has shown significant disruption in the

growth and development of *Pseudomonas aeruginosa* biofilms (Díaz De Rienzo *et al.* 2016;

Nguyen *et al.* 2020). Effects on other single species or medical biofilms have also been

investigated including methicillin-resistant *Staphylococcus aureus* (Ueda *et al.* 2019)*,*

Pseudomonas fluorescens (Simoes *et al.* 2008)*,* and *Salmonella enterica* serovar *typhimurium*

(Mireles II *et al.* 2001). Inhibited biofilm formation was seen to occur within all investigations*.*

The use of synthetic surfactants has been questioned due to the potential environmental

impact, specifically the longevity and toxicity to aquatic biota. For example, previous research

has shown that synthetic surfactants, such as Triton X-100 (Octylphenol Ethoxylate), are not

easily biodegradable under aerobic nor anaerobic conditions (Mohan *et al.* 2006). Other synthetic

surfactants such as SDS have also been noted as toxic to a variety of microorganisms such as

Vibrio fischeri, fish and shellfish such as *Clarias batrachus* and *Mytilus galloprovincialis,* yeast,

and also small invertebrates, such as *Daphnia magna,* and *Tigriopus fulvus* (Mariani *et al.* 2006;

Lima *et al.* 2011; Franzetti *et al.* 2012; Messina *et al.* 2014; Cao *et al.* 2020; Kumar *et al.* 2020).

Research has therefore shifted towards identifying a natural alternative that may be less harmful

to the aquatic environment.

Biofilm-dwelling microorganisms naturally produce a number of surfactants to control

the surrounding extracellular polymeric substance (EPS) matrix. The most common substances

of this group are rhamnolipids and sophorolipids which are glycolipids formed by the

combination of either a long-chain aliphatic acid or hydroxyaliphatic acid with a carbohydrate

(Desai and Banat 1997, Marchant and Banat 2012). When exposing single-cell biofilms of *P.*

aeruginosa to rhamnolipids, growth and development were disrupted, and cell viability was

reduced (De Rienzo *et al.* 2015). Disruptions by rhamnolipids have also been shown for single

species biofilms of *Staphylococcus aureus* (Quinn *et al.* 2013; Aleksic *et al.* 2017; Ceresa *et al.*

2021a; Tambone *et al.* 2021)*, Bacillus pumilus* (Dusane *et al.* 2010)*, and Rhodococcus*

erythropolis (Schreiberova *et al.* 2012). Yet, rhamnolipids are biodegradable in a variety of

environments, including under aerobic and anaerobic conditions (Mohan *et al.* 2006; Naughton

et al. 2019; Ramos da Silva *et al.* 2019). Biosurfactants in general have been reported to act as an

antibiofilm agents (Banat *et al.* 2014). Given that previous research primarily addressed the

effects of biological surfactants on single species biofilms of bacteria relevant to medical

settings, there remains a knowledge gap on how the exposure of multi-species freshwater

biofilms to surfactants affects biofilm microbial community composition and function.

There is a growing body of evidence that anthropogenic trace contaminants, and

contaminants of emerging concern (CECs), such as pharmaceuticals, herbicides, and

antimicrobial agents can have major impacts upon the composition and functioning of aquatic

microbial communities (Rosi-Marshall *et al.* 2013; Argudo *et al.* 2020; Harjung *et al.* 2020;

McClean and Hunter 2020; Ke *et al.* 2020; Kusi *et al.* 2020). Exposure to contaminants such as

the antibacterial agent Triclosan (Hay *et al.* 2001; Drury *et al.* 2013) and the herbicide Diuron

(Sumpono *et al.* 2003; Seghers *et al.* 2003) have been observed to place microbial communities

under a selective pressure that favors microorganisms that can use these contaminants as an

additional carbon source. Given that levels of surfactants and their metabolites are consistently

among the top contaminants identified within wastewater influent and groundwater (Lara-Martín

et al. 2014; Schaider *et al.* 2014), it is reasonable to infer that they could represent a major

control upon the microbial communities within human affected inland waters.

This study investigated the effects of the synthetic surfactant SDS and the biological

surfactant rhamnolipid on microbial biofilm community composition, microbial diversity, and

metabolic activity. Using recirculating flume mesocosms, we aimed to answer the following

overarching scientific question: What are the effects of a synthetic and biological surfactant on a

natural aquatic microbial biofilm, and how do these effects compare? We hypothesized that

biofilm exposure to a surfactant will decrease bacterial biodiversity, as the surfactants impact

microbial community composition and causes the loss of sensitive bacterial taxa as previous

research has shown that chemical exposure can place microbial communities under selective

pressure. Secondly, we hypothesize that the resultant changes in microbial community

composition will alter biofilm metabolic activity. We planned to test these hypotheses by

quantifying changes in community composition through 16s rRNA analysis and changes in

biofilm structure and metabolic activity through different analytical methods, such as changes in

gross primary productivity (GPP), community respiration (CR), ash free dry mass (AFDM),

Carbon: Nitrogen ratios (C:N), and extracellular enzyme activity (EEA).

Materials and Methods

Experimental Set up

We set up 15 recirculating experimental flumes near Lunz Am See, Austria following the

method described by Roche *et al.* (2017). Each flume was 2.5m long and had an approximate

slope of 0.8%. In each flume, ~5L of water recirculated, which was renewed with unfiltered lake

water twice per week. We added rhamnolipid (RL) (R90-100G, Sigma-Aldrich, Darmstadt,

Germany) and SDS (sodium dodecyl sulphate, Sigma-Aldrich, Darmstadt, Germany) as

treatments of 150 and 300 ppm to three flumes each (hereafter referred to as either RL 150, RL

300, SDS 150, or SDS 300), while three flumes acted as controls. After the first week, 5 μg L^{-1}

of phosphorus in the form of Na_2HPO_4 was also added to help growth due to low phosphorus in

the utilized lake water. Surfactant concentrations were chosen to represent the effective dose of

rhamnolipid, as well as double that concentration. Due to malfunction of one flume, *n* for the

rhamnolipid treatments was reduced to 2. Treatments were randomly placed among flumes.

Artificial light was spread equally over all flumes and light density averaged 9.5 µmol m^{-2} s^{-1} and

was the maximum even light density achievable with the provided light sources. We used a light

cycle of 12:12 and 18:6 (Cheah and Chan 2021) hours of light versus dark early and later in the

experiment to mimic natural light cycles at first and then to encourage biofilm growth later in the

experiment. Each flume was equipped with microscope slides (6) and clay tiles (8) to quantify

biofilm growth.

We measured pH, dissolved oxygen (DO) (mg/L), water temperature (°C), and flow (by

bucket measurement) every other day to ensure stable conditions. Twice per week we analysed

DOC, ammonium (NH_4), NO_x, and PO_4 of flume water to ensure they remained stable. In

addition, water temperature was monitored by automatic loggers in drainage tanks. After 32 days

the experiment was stopped due to adequate biofilm growth observed, and biofilm samples were

retrieved from the flumes for further analysis.

Sample Collection

Two microscope slides were removed from each flume and placed in 120mL Schott

bottles with oxygen sensors. Bottles were filled with lake water run through glass fiber filters to

remove any microorganisms that may alter the results of the analyses and sealed to remove all air

bubbles. Three control bottles with no slides were also set up to test for potential activity within

the filtered lake water with control activity subtracted from treatments during analyses. All

bottles were placed in an Imago 500 Controller Environmental Chamber (Snijders Scientific

B.V., Tilburg, Netherlands) using a 12:12 light cycle for 72 hours. Oxygen measurements were

taken at the start and end of each light and dark cycle to obtain data on GPP and CR.

We determined AFDM by weighing 2 clay tiles per flume after drying (9h at 70°C), and

combustion (4h at 450°C). AFDM was obtained by subtracting the remaining ash weight from

the dry weight. Two tiles from SDS 150 flumes were damaged during drying, and AFDM could

not be determined. Biofilms covering two tiles per flume were then scraped into sterilized and

pre-weighed 2mL Eppendorf tubes using a sterile razor blade, lyophilised (-0.001mbar; -76☐)

(Martin Christ, Gefriertrocknungsanlagen GmbH, Osterode am Harz, Germany) and then

analysed on a Thermo Scientific FLASH 2000 Elemental Analyser (Thermo Fisher Scientific

GmbH, Bremen, Germany) to quantify carbon (% C) and nitrogen (% N) content. Quantities of C

and N by mass were then examined in ratios to identify changes in ecological stoichiometry (the

relationship between elements, energy, and organisms and their interactions in an ecological

system (Sterner and Elser 2002; Van de Waal *et al.* 2018)).

EEA of β-D-1,4-glucosidase (glucosidase), phosphatase, and leucine-aminopeptidase

were measured in the biofilms following the method described in Romaní and Sabater (2001).

Four mL of sterile water was added to approximately 0.5 g of biofilm per flume that was scraped

from the tiles. The mixture was homogenized by shaking. EEA was measured via artificial

substrates: Glucosidase and phosphatase using 4-methylumbelliferyl (MUF)- substrates (MUF-β-

D-glucopyranoside and MUF-phosphate, respectively (Sigma-Aldrich, Darmstadt, Germany))

and leucine-aminopeptidase using a 7-amino-4-methylcoumarin (AMC)- substrate (L-leucine-

AMC, Sigma-Aldrich, Darmstadt, Germany). Artificial substrates were added to the homogenate

at a final concentration of MUF-glucosidase; 135 mg L^{-1}, MUF-phosphatase; 102.46 mg L^{-1}, and

AMC-leucine-aminopeptidase; 129.94 mg L^{-1} and incubated for 1 hour in the dark. Fluorescence

was measured at the beginning and end of the incubation with a plate reader (Varioskan Flash,

Thermo Fischer Scientific, Vaanta, Finland) at 365/455 nm excitation/emission wavelengths for

the MUF-substrates and 380/440 excitation/emission wavelengths for the AMC-substrate.

Two microscope slides per flume were scraped using sterile procedures into a sterile

1.5mL microcentrifuge tube for 16s rRNA analysis. DNA was immediately extracted from

collected samples using the Dneasy Power Soil Kit (Qiagen GmbH, Hilden, Germany) following

the manufacturer's methodology. We included internal replicates per flume (*n*=2), and

experimental controls were used to control for any potential unaccounted sample contamination.

Extracted products were sent to LGC Biosearch Technologies (LGC Genomics GmbH, Berlin,

Germany). DNA was quantified using gel electrophoresis before amplifying the V3-V4 region of

the 16s rRNA gene with polymerase chain reactions using 341F and 785R primers. Resultant

DNA were sequenced using the Illumina Mi-Seq platform (300 bp paired-end reads) for

amplicon creation.

We used Mothur version 1.44.3 (Schloss *et al.* 2009), and the Mothur MiSeq SOP

(Kozich *et al.* 2013) with some modification, to analyze returned reads. A maximum

homopolymer of 8 was chosen due to the limitation of the reference database (SILVA version

132) with a 97% similarity and above required for identification of operational taxonomic units

(OTUs). UCHIME (Edgar *et al.* 2011) was also used to identify and remove chimeric fragments.

Sequences that could not be directly identified through SILVA, were manually run through

NCBI's BLAST (Madden 2002). We were able to identify all but one OTU with 97% similarity;

OTU005 is thus reported as 'unnamed' in this study.

Data analysis

All data, except the 16s rRNA and C:N results, met assumptions of normality and

homogeneity of variance. We considered $p< 0.05$ as significant and analyzed data using R

version 4.0.3 (R Core Team 2020) and the R packages car (Fox and Weisberg 2019), vegan

(Oksanen *et al.* 2020), ggplot2 (Wickham 2016), and entropart (Marcon and Herault 2015). Non

metric multidimensional scaling analyses (NMDS) were run using the PAST 3 software

(Hammer *et al.* 2001). Due to replication error early in the experiment only the controls and the

300ppm concentration variables were used in statistical difference testing. All GPP and CR data

were normalized to mg O_2 in 120mL per mg of biofilm in 12 hours before being analyzed. C:N

data was normalized to mass (mg C/N per mg biofilm). A one way analysis of variance

(ANOVA) analysis was used to test for significant differences among treatments and the controls

for the GPP, CR, and AFDM. A Kruskal-Wallis test was used to test for significant differences

among the C:N ratios. Coefficients of variation were also calculated for individual flumes for

each analysis.

Changes in EEA were analysed through both a distance-based redundancy analysis (db-

RDA) and three one-way ANOVAs, one per extracellular enzyme. All the data was used within

the db-RDA and only the maximum concentrations of each chemical and control were used for

analysis in the ANOVA. Where appropriate, we applied a Tukey HSD post-hoc test to the

ANOVA output.

Bacterial OTUs were analyzed for alpha and beta diversity as well as differences in the

overall bacterial community composition. We used the Chao1 (Chao 1984) and the Shannon-

Wiener Diversity Indices to quantify bacterial Alpha diversity (Jost 2007). Chao 1 and Shannon-

Wiener results were then analyzed using an ANOVA followed by a Tukey HSD post-hoc test to

identify differences in diversity among treatments. Following this, all singletons and doubletons

were removed, and the data was rarefied based on the smallest number of sequences identified

within a sample. Beta diversity was then analysed using a non-metric multidimensional scaling

analysis (NMDS) on all data present to identify differences in the biofilm bacterial community

among treatments and controls. An analysis of similarities (ANOSIM) was performed with 999

permutations to identify if these differences were significant. Families and genera were

compared graphically, focusing on the top 80% of OTUs identified within the biofilms.

Results

Primary productivity and community respiration

Lake control bottles had an average primary productivity of 0.0304 ± 0.0256 mg O_2 in

120mL and average community respiration of -0.016 ± 0.0177 mg O_2 in 120mL. We found no

significant differences in primary productivity (ANOVA, $F=0.022$, $p=0.978$) and community

respiration (ANOVA, $F=0.148$, $p=0.865$) of the treatments versus the controls (Figures 1A, 1B).

All treatments showed substantial variation among replicates (Figure 1), with the most

pronounced variability being present in respiration. Coefficients of variation were highest for

respiration in the controls (1.56, see Table 1) and lowest (0.021) in the rhamnolipid 150

treatment. Rhamnolipid treated systems overall had the lowest variation for CR, while the

controls had the highest.

Ash free dry mass and C:N ratios

We found no significant difference among treatments and controls (ANOVA $F=0.045$,

$p=0.957$) for AFDM. The coefficients of variation, however, showed differences in variation

among the treatments (Table 1), with higher surfactant concentrations decreasing variation (e.g.,

RL 150 has greater variation than RL 300) (Figure 1C). Control variation was comparable to

both low dose surfactant variations. However, C (ANOVA F=18.36, p=0.0028) and N (ANOVA

F=5.471, p=0.044) content was significantly different by mass among treatments (Figure 2A and

Figure 2B). Differences lie between the RL 300 and the control for both C (Tukey HSD

p=0.0063) and N (Tukey HSD p=0.041) content, and between SDS 300 and the control for C

(Tukey HSD p=0.0036) content.

There were also no significant differences among treatments and controls for C:N ratios

by mass (Kruskal-Wallis p=0.193) present in the biofilms (Figure 2C). There were differences in

variation among treatments and controls (Table 1). Rhamnolipid treated flumes had lower

variation than either SDS treated flumes or the controls, however within each treatment a higher

surfactant concentration increased variation.

Bacterial community composition

Shannon-Wiener Diversity was significantly different among treatments (ANOVA

F=71.702, p<0.0001, Tukey HSD rhamnolipid and control; p=0.043, SDS and control;

p<0.0001, SDS and rhamnolipid; p<0.001) with a higher diversity in the control flumes, second

highest in the rhamnolipid flumes, and lowest in the SDS flumes. Significance was driven by

surfactant presence and type, but not concentration. The control had the greatest bacterial

diversity, while both rhamnolipid treatments had the next highest, and the lowest belonged to

both SDS treatments (Figure 3A). Chao1 had no significant differences (ANOVA F=3.7748,

p=0.087).

By testing beta diversity and community differences on the rarefied data, we identified

distinct groupings between all treatments and controls, as evident by the NMDS (Figure 3B). The

ANOSIM results confirmed the differences among both surfactants and the controls (R=0.9112,

p=0.001). Community differences were further identified by looking at only the taxa in the top

80% of all taxa present across all treatments, which was 18 taxa. We identified several genera

that were heavily dominant in rhamnolipid-treated flumes. These were: *Brevundimonas spp.*,

OTU005, *Pseudomonas,* and *Arthrobacter*. In SDS flumes we observed *Brevundimonas terrae,*

Flavobacterium spp. A, and in the control flumes *Exiguobacterium, Arenimonas,* and

Stenotrophomona. We also identified 32 genera only present in the controls, 5 there were

exclusive to RL 150, 6 in RL 300, 19 in SDS 150, and 7 in SDS 300 (Figure 3C).

Total OTUs within each treatment and OTUs within the top 80% of all taxa per treatment

were also calculated (Table 2). The control flumes had the greatest number of OTUs present, and

the greatest number of OTUs within their top 80% of taxa. The SDS and rhamnolipid flumes had

similar numbers of total OTUs present, but the SDS flumes had 80% of their total OTUs within

half the number of taxa as the rhamnolipids. This indicated a large difference in evenness

between treatments, while the total number of OTUs was fairly similar.

Extracellular Enzyme Activity

According to the db-RDA, EEA differed among all treatments and controls, with the

exception of the rhamnolipid treatments. The differences in db-RDA-distances were mainly

caused by the differences in EEA of leucine aminopeptidase and phosphatase, although the

results were not significant (ANOSIM *F*=2.9907, *p*=0.058). While phosphatase did not have any

significant differences among treatments (ANOVA *F*=2.934 *p*=0.129), with the exception of

SDS 150 addition of any surfactant decreased the production of phosphatase when compared to

the control (Figure 4A). Rhamnolipid additions yielded a significantly higher production of

glucosidase, while SDS additions reduced production (ANOVA F=5.741 *p*=0.0404, Tukey HSD

p=0.034) (Figure 4B). The effects of RL 300 and SDS 300 on leucine aminopeptidase were also

significantly different (ANOVA *F*=5.914 *p*=0.038, Tukey HSD *p*=0.032). The addition of

rhamnolipid yielded a higher amount of the extracellular enzymes when compared to the control

and the addition of SDS reduced production (Figure 4C).

Discussion

Our experiment showed that exposure to biological and synthetic surfactants alters the

metabolic activity and the community composition of a mixed-species freshwater biofilm. The

effects of SDS were more pronounced than the effects of rhamnolipid. Our experiment is the first

study to our knowledge that examines the effects of these two surfactants on a multi-species

near-natural freshwater biofilm, which contrasts with the commonly performed single-species

studies. This study was also novel as it directly compared the effects of a synthetic and a

biological surfactant and explicitly studied for the first time the C:N ratios and C and N contents

of biofilms that are exposed to surfactants. These results differ from previous research examining

the effectiveness of biosurfactants on biofilm removal in medical (Ceresa et al. 2019; Ceresa et

al. 2021b) and industrial (Campos et al. 2013) environments.

Primary Productivity, Respiration, Ash Free Dry Mass, and C:N ratios

Although our results showed biofilms had no differences in GPP or CR in any treatments

or controls, chemical exposure decreased biofilm respiratory activity in previous experiments,

including surfactant exposure (Simoes *et al.* 2008) and pharmaceutical exposure (Rosi-Marshall

et al. 2013). Biofilm volume has also been reduced after a single surfactant exposure event, with

decreases of 26.5-98.1% occurring between 22-48 hours after exposure (Mireles II *et al.* 2001;

Quinn *et al.* 2013). Typically, biosurfactants, which are amphiphilic polymers, can affect biofilm

surface attachment through altering surface wettability (Rodrigues *et al*. 2006). Biosurfactants

such as rhamnolipids are able to make cell surfaces more hydrophobic through loss of fatty acids

such as lipopolysaccharides (Al-Tahhan *et al.* 2000). Amphiphilic anionic surfactants work

similarly through altering surface characteristics and changing solubility of certain compounds

(Cserháti *et al*. 2002). Surface changes then can alter a biofilm's ability to adhere to a surface,

reducing growth and development. However, previous work on surfactant effects have also

shown that biofilms demonstrate resistance and can recover full respiratory activity 12 hours

after the initial exposure (Simoes *et al.* 2008) and biomass 5 hours after initial exposure (Nguyen

et al. 2020). As our experiment lasted 32 days, we believe that GPP, CR, and AFDM could have

stabilized and recovered full respiratory activity as seen in previous studies (Simoes *et al.* 2008;

Nguyen *et al.* 2020).

Altered C:N ratios are indicative of changes in microbial productivity (where reduced

C:N ratios may yield greater productivity) (Li *et al.* 2016) and biofilm formation (where lower

C:N ratios increase biofilm volume) (Thompson *et al.* 2006). As we did not have any alterations

in C:N ratios, and no changes in productivity (GPP and CR) or growth (AFDM) among any

treatments or controls, our results support these studies. The addition of surfactants provided an

added source of C, however as there were consistent C:N ratios across all flumes it can be

assumed that our system was C limited. It is also likely that a thickened EPS matrix, which may

be used as a defense mechanism in biofilms in response to surfactant exposure, could be another

source of extra carbon (Jefferson 2004; Zakaria and Dhar 2020). In contrast, in systems where C

is not limiting, an increase in C inputs through surfactants may increase C:N ratios, and therefore

negatively affect biofilm productivity and growth.

Diversity and bacterial community composition

Our controls had taxa present that were minimally identified within the surfactant flumes.

These taxa included species (spp.) of *Exiguobacterium*, *Acinetobacter, Arenimonas,* and

Stenotrophomonas. Exiguobacterium is currently researched as an emerging human pathogen

and while susceptible to antibiotics appears to easily develop antibiotic resistance (Chen *et al*.

2017). *Acinetobacter* contains species such as *A. baumanni* that are very well known as human

pathogens and that demonstrate high antibiotic resistance (Rice 2008; Tewari *et al*. 2018).

However, it appears that some species may not have resistance to biological surfactants as its

presence in rhamnolipid treated flumes was extremely low, although certain species of

Acinetobacter had an increase in abundance within the SDS treated flumes. Similar results

appear with *Stenotrophomonas* as it has human pathogenic potential, and species that have

developed multi drug resistance, but appears to be sensitive to both biological and synthetic

surfactants (Chang *et al*. 2015). In our study the addition of either surfactant created an

environment unsuitable for these bacteria, demonstrating how surfactant exposure can alter

biofilm community composition and potentially help to control certain pathogenic species that

are demonstrating resistance to antibiotics. The control flumes had the highest amount of OTUs

present (n=203) and the highest diversity, which indicates that both the diversity and abundance

of bacteria present in the biofilms decreases after surfactant exposure.

Rhamnolipids are studied for their ability to reduce pre-existing biofilms and prevent

future biofilm growth with some success (Quinn *et al.* 2013; Dusane *et al.* 2010; De Rienzo *et al.*

2015; Satpute *et al.* 2018). Current research has shown that rhamnolipid exposure on soil

bacteria can shift community composition, although many studies do not examine differences to

the genus level (Lu *et al.* 2019; Wei *et al.* 2020; Akbari *et al.* 2021). Research has also shown

that aquatic microbial communities will shift in response to an outside contaminant with an

increased abundance of bacteria that are not sensitive to that contaminant or can use it as a C

source (Hay *et al.* 2001; Sumpono *et al.* 2003; Seghers *et al.* 2003; Drury *et al.* 2013). Our study

showed that, much like soil communities and aquatic communities exposed to other

contaminants, aquatic biofilms experience significant community changes after exposure to

rhamnolipids.

Further examination into the bacterial genera indicate large increases in a *Brevundimonas*

spp., *Pseudomonas* spp., *Arthrobacter* spp., and OTU005 when compared to all other treatments

and to the controls. One OTU of *Arthrobacter* spp. was only present within the rhamnolipid

flumes, with the exception of one control flume that had low levels of that particular OTU.

Brevundimonas spp., who are opportunistic pathogens (Ryan and Pembroke 2018) have

experienced accelerated growth in the presence of rhamnolipids before when testing the effects

of rhamnolipids on increasing bioremediation in soils (Lu *et al.* 2019). It is likely that these

genera were able to use rhamnolipid as a C source due to the large C quantities within

rhamnolipids (Mnif *et al.* 2018). Their adaptability may explain why they were able to increase

in abundance over time when compared to the other genera present.

SDS caused the largest reduction in microbial diversity when compared with the other

treatments and controls. SDS has been researched and used for its ability to degrade biofilms in

applications where biofilms are undesirable (Izano *et al.* 2007; Ueda *et al.* 2019; Nguyen *et al.*

2020). This is particularly due to their ability to bind to hydrophobic protein components,

introducing a negative charge to the structure which unfolds the bound protein (Gudiksen *et al.*

2006; Fong and Yildiz 2015). While rhamnolipids are also able to denature proteins, they are

considered weak binders, and typically have a reduced denaturing ability when compared to

many other surfactants (Otzen 2017). This may explain why biofilms exposed to SDS had lower

diversity than those in the rhamnolipid treatments or the controls. SDS also had the lowest

amount of OTUs within the top 80% of taxa present in the treatments. We believe that the low

diversity, and community domination by few taxa, demonstrates that SDS exposure creates an

inhospitable environment to many potential specialist taxa. The few remaining and insensitive

taxa can then thrive as more resources are available. Following our findings, these taxa would

include *Brevundimonas terrae,* and an *Acinetobacter* and *Flavobacterium spp*. While the

rhamnolipid treated flumes also had less diversity than the controls, the rhamnolipid treatments

still appeared to favor more taxa and a higher diversity than the SDS treated flumes.

Extracellular Enzyme Activity

Our experiment showed that substantial changes in biofilm EEA can occur after

surfactant exposure. Although phosphatase activity, or the separation of phosphate groups from a

larger compound such as an acid or protein (Margalef *et al.* 2017), generally decreased after

surfactant exposure, we did have similar levels of activity between the control and SDS 150

flumes. Examination of the bacterial communities within SDS 150 flumes and control flumes

show several parallel genera present that are not as common in the other systems, including

Nocardioides. Treatments were laid out randomly across our experimental flumes within the

experimental setup, and water was allocated to all systems from the same source, which

minimized potential experimental alterations by relative location. We thus assume that all

changes in community composition and EEA refer to the actual biofilm response to treatments.

The presence of a low concentration SDS allowed for the development of bacterial genera like

Nocardioides which typically have both acid and alkaline phosphatase (Xie *et al.* 2017).

However, the addition of high concentration SDS, or rhamnolipid at either a high or low

concentration, reduced phosphatase activity.

Beta-glucosidase (rate-limiting enzyme that converts cellulase to glucose (Singh *et al.*

2016)) and leucine aminopeptidase (enzyme that aids in the hydrolysis of leucine and other N-

terminal residues at the end of peptides and proteins (Matsui *et al.* 2006)) activities were also

affected by surfactant exposure. Previous research has shown that rhamnolipids can act as an

enhancer for the enzyme beta-glucosidase (Zhang *et al.* 2009; Yao *et al.* 2020). Rhamnolipids

are able to protect proteins and limit the degradation of cellulase (Zhang *et al.* 2009; Wang *et al.*

2011). Yin *et al.* (2019) showed that the presence of rhamnolipid can increase genes related to

leucine and amino acid synthesis. The addition of rhamnolipid to our flumes likely protected

cellulase and leucine in the system which led to enhanced beta-glucosidase and leucine

aminopeptidase activity. Beta-glucosidase is also known for its ability to degrade *Pseudomonas*

aeruginosa, a rhamnolipid producing bacteria (Banar *et al.* 2019; Pirlar *et al.* 2020). The flumes

treated with rhamnolipid had a higher abundance of *P. aeruginosa* when compared with the other

flumes.

We assume therefore, that the higher beta-glucosidase levels within the rhamnolipid

flumes were likely caused by a combination of the presence of rhamnolipid to act as an enhancer,

and more abundant *P. aeruginosa.* However, we observed somewhat contrasting results in the

SDS treated flumes, where a decrease in enzyme activity occurred. SDS is an anionic surfactant

that can be used in denaturation of polypeptides (Bhuyan 2009; Schlager *et al.* 2012). SDS is

commonly used in an analysis called SDS-polyacrylamide gel electrophoresis (SDS-PAGE)

where an SDS infused gel is used to separate proteins and enzymes based on their molecular

weight. This analytical method has been used with experiments examining beta-glucosidase

previously (Laemmli 1970; Bai *et al.* 2013), which demonstrates that SDS is able to denature

beta-glucosidase. It is possible that the denaturing ability of SDS has degraded beta-glucosidase

and leucine aminopeptidase, leading to reduced enzyme activity, and likely also reduced biofilm

metabolic activity.

Our experiment showed that both the biologic surfactant rhamnolipid and the synthetic

surfactant, SDS, can significantly affect a mixed freshwater microbial biofilm in terms of

community composition and EEA. Our original hypotheses predicted changes to bacterial

biodiversity, and biofilm metabolic activity after exposure to either surfactant. Rhamnolipids

acted as an enhancer for beta-glucosidase and leucine aminopeptidase, increasing EEA within

the biofilm, while SDS likely denatured the enzymes and reduced EEA within the biofilm. While

there were no differences in GPP and CR, EEA changes can be used as a proxy for community-

scale metabolic activity within the biofilms. There were alterations in biofilm metabolic activity

post surfactant exposure, supporting one of our original hypotheses. Both surfactants also

decreased the diversity of the biofilm bacterial community, which supports our other original

hypothesis. There were unique genera in the biofilms of each treatment, including the controls

which demonstrates that the presence of either surfactant creates an environment that is

inhospitable to sensitive taxa. Although we saw differences in EEA and diversity, there were no

differences in GPP and CR, potentially demonstrating functional redundancy within mixed

microbial biofilms. These results highlight several important areas of concern. Firstly,

rhamnolipids may not be a suitable alternative to SDS for use environmental biofilm control

because of its lower effectiveness in open environmental settings. Secondly, in natural

ecosystems where biofilms are necessary for key ecosystem processes, such as degradation of

organic matter, we have shown that the presence of these surfactants, (for example through the

release from WWTPs) may lead to negative effects on microbial diversity and metabolic activity

within biofilms. Finally, our results demonstrate that functional redundancy may exist within

diverse biofilms. Surfactant exposure altered community composition and EEA, but the overall

biofilm did not experience any negative respiratory effects. Therefore, while the function of the

biofilm may be altered, enough bacteria are still taking part in respiratory cycles that the overall

metabolic activity of the biofilm is not disrupted.

Statements and Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Data can be accessed at
https://zenodo.org/record/5153485#.YQgBzEAo9PY. Access is restricted and must be requested

from the page to prevent work from being copied before publishing. After publishing work will

be publicly accessible.

Competing interests

The authors declare no competing interests.

Funding

SG is funded by an Ulster University Vice Chancellors Doctoral Research Fellowship, and

received additional support through an Ulster University Broadening Horizons Travel Bursary.

Analytical costs were partly supported by the HYDRO-DIVERSITY project funded by the

472　

and core funding of the AFBI Aquatic Chemistry Laboratory (WH).

Author's contributions

SG, WH, LC, IB, and JS were all equally involved in the design, execution, and data analysis

within the study. SG wrote the first draft of the manuscript with the exception of the extracellular

enzyme analysis methodology which was written by LC. All authors contributed to commenting

on and editing the manuscript and have read and approved of the final manuscript.

References

Akbari A, Kasprzyk A, Galvez R, Ghoshal S (2021) A rhamnolipid biosurfactant increased

bacterial population size but hindered hydrocarbon biodegradation in weathered

contaminated soils. Sci Total Environ. https://doi.org/10.1016/j.scitotenv.2021.145441

Aleksic I, Petkovic M, Jovanovic M, Milivojevic D, Vasiljevic B, Nikodinovic-Runic J,

Senerovic L (2017) Anti-biofilm properties of bacterial di-Rhamnolipids and their semi-

synthetic amide derivatives. Front Microbiol. https://doi.org/10.3389/fmicb.2017.02454

Al-Tahhan R, Sandrin TR, Bodour AA, Maier RM (2000) Rhamnolipid-induced removal of

lipopolysaccharide from *Pseudomonas aeruginosa*: effect on cell surface properties and

interaction with hydrophobic substrates. Appl Environ Microbiol.

https://doi.org/10.1128/AEM.66.8.3262-3268.2000

Argudo M, Gich F, Bonet B, Espinosa C, Gutiérrez M, Guasch H (2020) Responses of resident

(DNA) and active (RNA) microbial communities in fluvial biofilms under different

polluted scenarios. Chemosphere. https://doi.org/10.1016/j.chemosphere.2019.125108

Bai H, Wang H, Sun J, Irfan M, Han M, Huang Y, Han X, Yang Q (2013) Production,

purification and characterization of novel beta glucosidase from newly isolated

Penicillium simplicissimum H-11 in submerged fermentation. EXCLI journal

Banar M, Emaneini M, Beigverdi R, Fanaei Pirlar R, Node Farahani N, van Leeuwen WB,

Jabalameli F (2019) The efficacy of lyticase and β-glucosidase enzymes on biofilm

degradation of *Pseudomonas aeruginosa* strains with different gene profiles. BMC

Microbiol. https://doi.org/10.1186/s12866-019-1662-9

Banat IM, Díaz De Rienzo MA, Quinn GA (2014). Microbial biofilms: biosurfactants as

antibiofilm agents. Appl Micobiol Biotechnol. https://doi.org/10.1007/s00253-014-6169-

6.

Battin TJ, Besemer K, Bengtsson MM, Romani AM, Packmann AI (2016) The ecology and

biogeochemistry of stream biofilms. Nat Rev Microbiol.

https://doi.org/10.1038/nrmicro.2016.15

Bhuyan AK (2010) On the mechanism of SDS-induced protein denaturation. Biopolymers.

https://doi.org/10.1002/bip.21318

Bowler P, Murphy C, Wolcott R (2020) Biofilm exacerbates antibiotic resistance: Is this a

current oversight in antimicrobial stewardship? Antimicrob Resist Infect Control.

https://doi.org/10.1186/s13756-020-00830-6

Cao C, Cao Z, Yu P, Zhao Y (2020) Genome-wide identification for genes involved in sodium

dodecyl sulfate toxicity in *Saccharomyces cerevisiae*. BMC Microbiol.

https://doi.org/10.1186/s12866-020-1721-2

Campos JM, Montenegro Stamford TL, Sarubbo LA, de Luna JM, Rufino RD, Banat IM (2013)

Microbial biosurfactants as additives for food industries. Biotechnol Progress. 29:1097-

1108. https://doi.org/10.1002/btpr.1796

Ceresa C, Tessarolo F, Maniglio D, Tambone E, Carmagnola I, Fedeli E, Caola I, Nollo G,

Chiono V, Allegrone G, Rinaldi M, Fracchia L (2019) Medical-grade silicone coated

with Rhamnolipid R89 is effective against *Staphylococcus spp*. biofilms. Molecules.

24:3843. https://doi.org/10.3390/molecules24213843

Ceresa C, Rinaldi M, Tessarolo F, Maniglio D, Fedeli E, Tambone E, Caciagli P, Banat IM, Diaz

De Rienzo MA, Fracchia L (2021a) Inhibitory effects of lipopeptides and glycolipids on

C. albicans - Staphylococcus spp. dual-species biofilms. Front Microbiol. 11:545654.

https://doi.org/10.3389/fmicb.2020.545654

Ceresa C, Fracchia L, Fedeli E, Porta C, Banat IM (2021b) Recent advances in biomedical,

therapeutic and pharmaceutical applications of microbial surfactants. Pharmaceutics,

13(4), 466. https://doi.org/10.3390/pharmaceutics13040466

Chao A (1984) Nonparametric Estimation of the Number of Classes in a Population. Scand J

Stat. 11:265-270.

Chang YT, Lin CY, Chen YH, Hsueh P (2015) Update on infections caused by

Stenotrophomonas maltophilia with particular attention to resistance mechanisms and

therapeutic options. Front Microbiol

Cheah YT, Chan DJC (2021) Physiology of microalgal biofilm: a review on prediction of

adhesion on substrates. https://doi.org/10.1080/21655979.2021.1980671

Chen X, Wang L, Zhou J, Wu H, Li D, Cui Y, Lu B (2017) *Exiguobacterium sp. A1b/GX59*

isolated from a patient with community-acquired pneumonia and bacteremia: genomic

characterization and literature review. BMC Infect Dis. https://doi.org/10.1186/s12879-

017-2616-1

Cserháti T, Forgács E, Oros G (2002) Biological activity and environmental impact of anionic

surfactants. Environ Int. https://doi.org/10.1016/S0160-4120(02)00032-6

Cui B, Liu X, Yang Q, Li J, Zhou X, Peng Y (2017) Achieving partial denitrification through

control of biofilm structure during biofilm growth in denitrifying biofilter. Bioresour

Technol. https://doi.org/10.1016/j.biortech.2017.04.034

Dang H, Lovell CR (2016) Microbial surface colonization and biofilm development in marine

environments. Microbiol Mol Biol Rev. https://doi.org/10.1128/MMBR.00037-15

Decho AW (2000) Microbial biofilms in intertidal systems: an overview. Cont Shelf Res.

https://doi.org/10.1016/S0278-4343(00)00022-4

Desai JD, Banat IM (1997) Microbial production of surfactants and their commercial potential.

Microbiol Mol Biol Rev. https://doi.org/10.1128/.61.1.47-64.1997

Díaz De Rienzo MA, Banat IM, Dolman B, Winterburn J, Martin PJ (2015) Sophorolipid

biosurfactants: Possible uses as antibacterial and antibiofilm agent. New Biotechnol.

https://doi.org/10.1016/j.nbt.2015.02.009

Díaz De Rienzo MA, Stevenson PS, Marchant R, Banat IM (2016) *Pseudomonas aeruginosa*

biofilm disruption using microbial surfactants. J Appl Microbiol.

https://doi.org/10.1111/jam.13049

Dodds WK, Evans-White M, Gerlanc NM, Gray L, Gudder DA, Kemp MJ, Lopez AL, Stagliano

D, Strauss EA, Tank JL, Whiles MR, Wollheim WM (2000) Quantification of the

nitrogen cycle in a prairie stream. Ecosystems 3:574-589.

Drury B, Scott J, Rosi-Marshall EJ, Kelly JJ (2013) Triclosan exposure increases triclosan

resistance and influences taxonomic composition of benthic bacterial communities.

Environ Sci Technol. https://doi.org/10.1021/es401919k

Dusane DH, Zinjarde SS, Venugopalan VP, Mclean RJ, Weber MM, Rahman PKSM (2010)
Quorum sensing: implications on Rhamnolipid biosurfactant production. Biotechnol
Genet Eng Rev. https://doi.org/10.1080/02648725.2010.10648149

Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity
and speed of chimera detection. Bioinformatics.
https://doi.org/10.1093/bioinformatics/btr381

Fanaei PR, Emaneini M, Beigverdi R, Banar M, van Leeuwen WB, Jabalameli F (2020)
Combinatorial effects of antibiotics and enzymes against dual-species *Staphylococcus
aureus* and *Pseudomonas aeruginosa* biofilms in the wound-like medium. PLOS ONE
15:e0235093.

Flemming H, Wingender J, Szewzyk U, Steinberg P, Rice SA, Kjelleberg S (2016) Biofilms: an
emergent form of bacterial life. Nat Rev Microbiol.
https://doi.org/10.1038/nrmicro.2016.94

Fong JNC, Yildiz FH (2015) Biofilm matrix proteins. Microbiol Spectr 3:10.
https://doi.org/10.1128/microbiolspec.MB-0004-2014

Fox J, Weisberg S (2019) An R Companion to applied regression, Third Edition. Thousand Oaks
California, USA: Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Franzetti A, Gandolfi I, Raimondi C, Bestetti G, Banat IM, Smyth TJ, Papacchini M, Cavallo M,
Fracchia L (2012) Environmental fate, toxicity, characteristics and potential applications
of novel bioemulsifiers produced by *Variovorax paradoxus 7bCT5*. Bioresour Technol.
https://doi.org/10.1016/j.biortech.2012.01.005

Gudiksen KL, Gitlin I, Whitesides GM (2006) Differentiation of proteins based on characteristic

patterns of association and denaturation in solutions of SDS. Proc Natl Acad Sci USA.

https://doi.org/10.1073/pnas.0602816103

Hammer Ø, Harper DAT, Ryan PD (2001) PAST: Paleontological statistics software package for

education and data analysis. Palaeontol Electron.

https://doi.org/10.1016/j.bcp.2008.05.025

Harjung A, Attermeyer K, Aigner V, Krlovic N, Steniczka G, Švecová H, Schagerl M, Schelker

J (2020) High anthropogenic organic matter inputs during a festival increase river

heterotrophy and refractory carbon load. Environ Sci Technol.

https://doi.org/10.1021/acs.est.0c02259

Hay AG, Dees PM, Sayler GS (2001) Growth of a bacterial consortium on triclosan. FEMS

Microbiol Ecol. https://doi.org/10.1016/S0168-6496(01)00127-1

Izano EA, Wang H, Ragunath C, Ramasubbu N, Kaplan JB (2007) Detachment and killing of

Aggregatibacter actinomycetemcomitans biofilms by Dispersin B and SDS. J Dent Res.

https://doi.org/10.1177/154405910708600707

Jefferson KK (2004) What drives bacteria to produce a biofilm? FEMS Microbiol Lett.

https://doi.org/10.1111/j.1574-6968.2004.tb09643.x

Jost L (2007) Partitioning diversity into independent alpha and beta components. Ecol (Durham).

https://doi.org/10.1890/06-1736.1

Ke Y, Chen J, Hu X, Tong T, Huang J, Xie S (2020) Emerging perfluoroalkyl substance impacts

soil microbial community and ammonia oxidation. Environ Pollut.

https://doi.org/10.1016/j.envpol.2019.113615

Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD (2013) Development of a dual-

index sequencing strategy and curation pipeline for analyzing amplicon sequence data on

the MiSeq Illumina sequencing platform. Appl Environ Microbiol.

https://doi.org/10.1128/AEM.01043-13

Kumar A, Yadav CB, Bajpeyee AK (2020) Glucose, protein, cholesterol and

immunohaematological alterations with acute (SDS) Sodium Dodecyl Sulfate toxicity in

Clarias batrachus. J Exp Zoo (India). 23:401-404.

Kusi J, Scheuerman PR, Maier KJ (2020) Emerging environmental contaminants (silver

nanoparticles) altered the catabolic capability and metabolic fingerprinting of microbial

communities. Aquat Toxicol. https://doi.org/10.1016/j.aquatox.2020.105633

Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of

bacteriophage T4. Nature. 227:680-685.

Landa AS, Van de Belt-Gritter B, Van der Mei HC, Busscher HJ (1999) Recalcitrance of

Streptococcus mutans biofilms towards detergent-stimulated detachment. Eur J Oral Sci.

https://doi.org/10.1046/j.0909-8836.1999.eos107402.x

Lara-Martín PA, González-Mazo E, Petrovic M, Barceló D, Brownawell BJ (2014) Occurrence,

distribution and partitioning of nonionic surfactants and pharmaceuticals in the urbanized

Long Island Sound Estuary (NY). Mar Pollut Bull.

https://doi.org/10.1016/j.marpolbul.2014.01.022

Li Y, Wu J, Shen J, Liu S, Wang C, Chen D, Huang T, Zhang J (2016) Soil microbial C:N ratio

is a robust indicator of soil productivity for paddy fields. Sci Rep.

https://doi.org/10.1038/srep35266

Lima TMS, Procópio LC, Brandão FD, Leão BA, Tótola MR, Borges AC (2011) Evaluation of

bacterial surfactant toxicity towards petroleum degrading microorganisms. Bioresour

Technol. https://doi.org/10.1016/j.biortech.2010.09.109

Lindsay D, von Holy A (2006) Bacterial biofilms within the clinical setting: what healthcare

professionals should know. J Hosp Infect.   https://doi.org/10.1016/j.jhin.2006.06.028

Lu L, Zhang J, Peng C (2019) Shift of soil polycyclic aromatic hydrocarbons (PAHs) dissipation

pattern and microbial community composition due to Rhamnolipid supplementation.

Water Air Soil Pollut. https://doi.org/10.1007/s11270-019-4118-9

Lyon DR, Ziegler SE (2009) Carbon cycling within epilithic biofilm communities across a

nutrient gradient of headwater streams. Limnol Oceanogr.

https://doi.org/10.4319/lo.2009.54.2.0439

Madden T (2002) Chapter 16: The BLAST sequence analysis tool. In: McEntyre J, Ostell J (eds)

The NCBI Handbook. National Center for Biotechnology Information, Bethesda.

Marchant R, Banat IM (2012) Biosurfactants: a sustainable replacement for chemical

surfactants? Biotechnol. Lett. 34: 1597-1605.

Margalef O, Sardans J, Fernández-Martínez M, Molowny-Horas R, Janssens IA, Ciais P, Goll D,

Richter A, Obersteiner M, Asensio D, Peñuelas J (2017) Global patterns of phosphatase

activity in natural soils. Sci Rep. https://doi.org/10.1038/s41598-017-01418-8

Mariani L, De Pascale D, Faraponova O, Tornambè A, Sarni A, Giuliani S, Ruggiero G, Onorati

F, Magaletti E (2006) The use of a test battery in marine ecotoxicology: The acute

toxicity of Sodium Dodecyl Sulfate. Enviro Toxicol. https://doi.org/10.1002/tox.20204

Matsui M, Fowler JH, Walling LL (2006) Leucine aminopeptidases: diversity in structure and

function. Biol Chem. https://doi.org/10.1515/BC.2006.191

McClean P, Hunter WR (2020) 17α-ethynylestradiol (EE2) limits the impact of ibuprofen upon

respiration by streambed biofilms in a sub-urban stream. Environ Sci Pollut Res

27:37149-37154.

Messina CM, Faggio C, Laudicella VA, Sanfilippo M, Trischitta F, Santulli A (2014) Effect of

Sodium Dodecyl Sulfate (SDS) on stress response in the Mediterranean mussel (Mytilus

Galloprovincialis): Regulatory volume decrease (Rvd) and modulation of biochemical

markers related to oxidative stress. Aquat Toxicol.

https://doi.org/10.1016/j.aquatox.2014.10.001

Milovanovic M, Arsenijevic A, Milovanovic J, Kanjevac T, Arsenijevic N (2017) Chapter 14:

Nanoparticles in antiviral therapy. In: Antimicrobial Nanoarchitectonics. Elsevier Inc, pp

383-410.

Mireles II JR, Toguchi A, Harshey RM (2001) *Salmonella enterica Serovar Typhimurium*

swarming mutants with altered biofilm-forming abilities: Surfactin inhibits biofilm

formation. J Bacteriol. https://doi.org/10.1128/JB.183.20.5848-5854.2001

Mnif I, Ellouz-Chaabouni S, Ghribi D (2018) Glycolipid biosurfactants, main classes, functional

properties and related potential applications in environmental biotechnology. J Polym

Environ. https://doi.org/10.1007/s10924-017-1076-4

Mohan PK, Nakhla G, Yanful EK (2006) Biokinetics of biodegradation of surfactants under

aerobic, anoxic and anaerobic conditions. Water Res (Oxford).

https://doi.org/10.1016/j.watres.2005.11.030

Naughton PJ, Marchant R, Naughton V, Banat IM (2019). Microbial biosurfactants: Current

trends and applications in Agricultural and Biomedical industries. J Appl Microbiol.

https:// doi.org/10.1111/jam.14243

Nguyen BVG, Nagakubo T, Toyofuku M, Nomura N, Utada AS (2020) Synergy between

Sophorolipid biosurfactant and SDS increases the efficiency of *P. aeruginosa* biofilm

disruption. Langmuir. https://doi.org/10.1021/acs.langmuir.0c00643

Otzen DE (2017) Biosurfactants and surfactants interacting with membranes and proteins: Same

but different? Biochim Biophys Acta Biomembr.

https://doi.org/10.1016/j.bbamem.2016.09.024

Paul BJ, Duthie HC, Taylor WD (1991) Nutrient cycling by biofilms in running waters of

differing nutrient status. J North Am Benthol Soc. 10.2307/1467762

Qu X, Ren Z, Zhang H, Zhang M, Zhang Y, Liu X, Peng W (2017) Influences of anthropogenic

land use on microbial community structure and functional potentials of stream benthic

biofilms. Sci Rep. https://doi.org/10.1038/s41598-017-15624-x

Quinn GA, Maloy AP, Banat MM, Banat IM (2013) A comparison of effects of broad-spectrum

antibiotics and biosurfactants on established bacterial biofilms. Curr Microbiol.

https://doi.org/10.1007/s00284-013-0412-8

Ramos da Silva A, Manresa MÁ, Pinazo A, García MT, Pérez L (2019) Rhamnolipids

functionalized with basic amino acids: Synthesis, aggregation behavior, antibacterial

activity and biodegradation studies. Colloids Surf B.

https://doi.org/10.1016/j.colsurfb.2019.05.037

Rigotti RT, Corrêa JAF, Maia NJL, Cesaro G, Rosa EAR, Macedo, Renata Ernlund Freitas de,

Luciano FB (2017) Combination of natural antimicrobials and sodium dodecyl sulfate for

disruption of biofilms formed by contaminant bacteria isolated from sugarcane mills.

Innov Food Sci Emerg Technol. https://doi.org/10.1016/j.ifset.2017.01.007

Rice LB (2008) Federal funding for the study of antimicrobial resistance in nosocomial

pathogens: No ESKAPE. J Infect Dis. https://doi.org/10.1086/533452

Roche KR, Drummond JD, Boano F, Packman AI, Battin TJ, Hunter WR (2017) Benthic biofilm

controls on fine particle dynamics in streams. Water Resour Res.

https://doi.org/10.1002/2016WR019041

Rodrigues L, Banat IM, Teixeira J, Oliveira R (2006) Biosurfactants: potential applications in

medicine. J Antimicrob Chemother. https://doi.org/10.1093/jac/dkl024

Romaní AM, Sabater S (2001) Structure and activity of rock and sand biofilms in a

Mediterranean stream. Ecol (Durham). https://doi.org/10.1890/0012-

9658(2001)082[3232:SAAORA]2.0.CO;2

Rosi-Marshall E, Kincaid DW, Bechtold HA, Royer TV, Rojas M, Kelly JJ (2013)

Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial

communities in stream biofilms. Ecol Appl. https://doi.org/10.1890/12-0491.1

Ryan MP, Pembroke JT (2018) *Brevundimonas spp*: Emerging global opportunistic pathogens.

Virulence. https://doi.org/10.1080/21505594.2017.1419116

Satpute SK, Mone NS, Das P, Banpurkar AG, Banat IM (2018). *Lactobacillus acidophilus*

derived biosurfactant as a biofilm inhibitor: A promising investigation using microfluidic

approach. Appl Sci. https://doi.org/10.3390/app8091555

Schaider LA, Rudel RA, Ackerman JM, Dunagan SC, Brody JG (2014) Pharmaceuticals,

perfluorosurfactants, and other organic wastewater compounds in public drinking water

wells in a shallow sand and gravel aquifer. Sci Total Environ.

https://doi.org/10.1016/j.scitotenv.2013.08.067

Schlager B, Straessle A, Hafen E (2012) Use of anionic denaturing detergents to purify insoluble

proteins after overexpression. BMC Biotechnol. https://doi.org/10.1186/1472-6750-12-95

Schloss PD, Westcott SL, Thomas R, Hall JR, Martin H, Hollister EB, Lesniewski RA, Oakley

BB, Parks DH, Robinson CJ, Sahl JW, Blaz S, Thallinger GG, Van Horn DJ, Weber CF

(2009) Introducing mothur: open-source, platform-independent, community-supported

software for describing and comparing microbial communities. Appl Environ Microbiol.

https://doi.org/10.1128/AEM.01541-09

Schmitz KS (2018) Chapter 4: Life Science. In: Physical Chemistry. Elsevier, Boston, pp 755-

832

Schreiberova O, Hedbavna P, Aejkova A, JirkA[macron] V, Masak J (2012) Effect of surfactants

on the biofilm of *Rhodococcus erythropolis*, a potent degrader of aromatic pollutants.

New Biotechnol. https://doi.org/10.1016/j.nbt.2012.04.005

Seghers D, Verthé K, Reheul D, Bulcke R, Siciliano SD, Verstraete W, Top EM (2003) Effect of

long-term herbicide applications on the bacterial community structure and function in an

agricultural soil. FEMS Microbiol Ecol. https://doi.org/10.1016/S0168-6496(03)00205-8

Simões M, Simões LC, Pereira MO, Vieira MJ (2008) Sodium Dodecyl Sulfate allows the

persistence and recovery of biofilms of *Pseudomonas fluorescens* formed under different

hydrodynamic conditions. Biofouling. https://doi.org/10.1080/08927010701730311

Singh G, Verma AK, Kumar V (2015) Catalytic properties, functional attributes and industrial

applications of β-glucosidases. 3 Biotech. https://doi.org/10.1007/s13205-015-0328-z

Sloup RE, Cieza RJ, Needle DB, Abramovitch RB, Torres AG, Waters CM (2016) Polysorbates

prevent biofilm formation and pathogenesis of *Escherichia coli O104:H4*. Biofouling.

https://doi.org/10.1080/08927014.2016.1230849

Srey S, Jahid IK, Ha S (2013) Biofilm formation in food industries: A food safety concern. Food

Control. https://doi.org/10.1016/j.foodcont.2012.12.001

Sterner RW, Elser JJ (2002) Ecological stoichiometry. Princeton university press

Sumpono, Perotti P, Belan A, Forestier C, Lavedrine B, Bohatier J (2003) Effect of Diuron on

aquatic bacteria in laboratory-scale wastewater treatment ponds with special reference to

Aeromonas species studied by colony hybridization. Chemosphere.

https://doi.org/10.1016/S0045-6535(02)00621-5

Tambone E, Bonomi E, Ghensi P, Maniglio D, Ceresa C, Agostinacchio F, Caciagli P, Nollo G,

Piccoli F, Caola I, Fracchia L, Tessarolo F (2021) Rhamnolipid coating reduces microbial

biofilm formation on titanium implants: an in vitro study. BMC Oral Health.

https://doi.org/10.1186/s12903-021-01412-7

Tewari R, Chopra D, Wazahat R, Dhingra S, Dudeja M (2018) Antimicrobial susceptibility

patterns of an emerging multidrug resistant nosocomial pathogen: *Acinetobacter*

baumannii. Malays J Med Sci. https://doi.org/10.21315/mjms2018.25.3.13

Thompson LJ, Gray V, Lindsay D, Von Holy A (2006) Carbon: nitrogen: phosphorus ratios

influence biofilm formation by *Enterobacter cloacae* and *Citrobacter freundii*. J Appl

Microbiol. https://doi.org/10.1111/j.1365-2672.2006.03003.x

Ueda Y, Mashima K, Miyazaki M, Hara S, Takata T, Kamimura H, Takagi S, Jimi S (2019)

Inhibitory effects of polysorbate 80 on MRSA biofilm formed on different substrates

including dermal tissue. Sci Rep. https://doi.org/10.1038/s41598-019-39997-3

Van de Waal, D. B, Elser JJ, Martiny AC, Sterner RW, Cotner JB (2018) Editorial: Progress in

ecological stoichiometry. Front Microbiol. https://doi.org/10.3389/fmicb.2018.01957

Waaler SM, Rölla G, Skjörland KK, Ogaard B (1993) Effects of oral rinsing with triclosan and

sodium lauryl sulfate on dental plaque formation: a pilot study. Scand J Dent Res

101:192-195.

Wang H, Fan B, Li C, Liu S, Li M (2011) Effects of Rhamnolipid on the cellulase and xylanase

in hydrolysis of wheat straw. Bioresour Technol 102:6515-6521.

Wei Z, Wang JJ, Gaston LA, Li J, Fultz LM, DeLaune RD, Dodla SK (2020) Remediation of

crude oil-contaminated coastal marsh soil: Integrated effect of biochar, Rhamnolipid

biosurfactant and nitrogen application. J Hazard Mater.

https://doi.org/10.1016/j.jhazmat.2020.122595

Wickham H (2016) Ggplot2: Elegant graphics for data analysis. New York, USA: Springer-

Verlag.

Xie F, Yang Y, Ma H, Quan S, Yue D, Sun Y, Feng F, Zhen J, Liu D (2017) *Nocardioides*

phosphati sp. nov., an *actinomycete* isolated from a phosphate mine. Int J Syst Evol.

https://doi.org/10.1099/ijsem.0.001754

Yao X, Xiao L, Liu H, Zhou M, Zhou Z, Ju X, Li L (2020) Rhamnolipid enhanced beta-

glucosidase from *Paenibacillus sp. LLZ1* for in situ lignocellulose saccharification in

ionic liquids. Biomass Convers Biorefin. https://doi.org/10.1007/s13399-020-01002-7

Yin Y, Gu J, Wang X, Zhang Y, Zheng W, Chen R, Wang X (2019) Effects of rhamnolipid and

Tween-80 on cellulase activities and metabolic functions of the bacterial community

during chicken manure composting. Bioresour Technol.

https://doi.org/10.1016/j.biortech.2019.121507

Zakaria BS, Dhar BR (2020) Changes in syntrophic microbial communities, EPS matrix, and

gene-expression patterns in biofilm anode in response to silver nanoparticles exposure.

Sci Total Environ. https://doi.org/10.1016/j.scitotenv.2020.139395

Zhang Q, He G, Wang J, Cai W, Xu Y (2009) Mechanisms of the stimulatory effects of

Rhamnolipid biosurfactant on rice straw hydrolysis. Appl Energy.

https://doi.org/10.1016/j.apenergy.2009.04.030

Zhang S, Pang S, Wang P, Wang C, Guo C, Addo FG, Li Y (2016) Responses of bacterial

community structure and denitrifying bacteria in biofilm to submerged macrophytes and

nitrate. Sci Rep. https://doi.org/10.1038/srep36178

Figure Legends

Fig. 1 GPP (A), CR (B) and AFDM (C) of microbial biofilms exposed to SDS (sodium dodecyl

sulphate) and RL (rhamnolipid) for 32 days at 150 and 300ppm.

Fig. 2 Carbon (A), Nitrogen (B) and C:N ratios (C) of microbial biofilms exposed to SDS

(sodium dodecyl sulphate) and RL (rhamnolipid) for 32 days at 150 and 300ppm.

Fig. 3 Biofilm bacterial diversity from 16s rRNA sequencing of biofilm exposed to SDS (sodium

dodecyl sulphate) or RL (rhamnolipid) after 32 days at 150 and 300ppm. A) shows Shannon

Diversity, B) shows an NMDS depicting the community differences in treatments when

examining all OTUs present within each sample after rarefication and C) depicts relative

abundance of OTUs.

Fig. 4 EEA of phosphatase (A), glucosidase (B), and leucine aminopeptidase (C) of biofilms

after exposure to either SDS (sodium dodecyl sulphate) or RL (rhamnolipid) for 32 days at 150

and 300ppm. (*indicates significant differences)

Table 1 Coefficients of variation for AFDM, GPP, and CR data. Data collected from samples of biofilm grown in either the surfactant SDS (sodium dodecyl sulphate) or RL (rhamnolipid) for 32 days.

Treatment	Ash free dry mass	Gross primary productivity	CR	C:N
SDS 150	0.70	0.67	0.92	0.373
SDS 300	0.11	0.626	1.09	0.643
RL 150	0.60	0.34	0.021	0.0875
RL 300	0.32	0.52	0.59	0.0698
Control	0.59	0.61	1.56	0.569

Table 2 Biofilm OTUs per surfactant treatment and their dominant genera. Biofilms were grown in either SDS (sodium dodecyl sulphate) or rhamnolipid for 32 days.

Surfactant	Conc. (ppm)	Total OTUs present	OTUs in top 80% per treatment	OTUs lost compared to control	OTUs gained compared to control	Dominant genera in top 80% of all taxa present						
Rhamnolipid	150	145	10	58	25	*Brevundimonas*	*Otu005*	*Pseudomonas*	*Arthrobacter*	*Flavobacterium B*	*Prosthecobacter*	*Nocardioides*
Rhamnolipid	300	159	14	44	36	*Brevundimonas*	*Otu005*	*Pseudomonas*	*Arthrobacter*	*Flavobacterium B*	*Prosthecobacter*	
SDS	150	176	7	27	37	*Brevundimonas terrae*	*Acinetobacter*	*Flavobacterium A*	*Paucibacter*	*Flavobacterium B*	*Prosthecobacter*	
SDS	300	130	6	73	23	*Brevundimonas terrae*	*Acinetobacter*	*Flavobacterium A*	*Prosthecobacter*	*Flavobacterium B*	*Prosthecobacter*	*Sphingomonas*
Control	0	203	15	--	--	*Exiguobacterium*	*Acinetobacter*	*Arenimonas*	*Stenotrophomonas*	*Nocardioides*		