The immediate early protein 1 of human herpesvirus 6B counteracts ATM activation in an NBS1-dependent manner

Vanessa Collin[1,2,7], Élise Biquand[3,4,5,6,7], Vincent Tremblay[3,4,5,7], Élise G. Lavoie[3,5], Julien Dessapt[3,4,5], Andréanne Blondeau[3,5], Annie Gravel[1,2], Louis Flamand[1,2,*], Amélie Fradet-Turcotte[3,4,5,*]

[1] Division of Infectious Disease and Immunity, Centre Hospitalier Universitaire (CHU) de Québec-Université Laval Research Center, Quebec City, Quebec, Canada, G1V 4G2;

[2] Department of microbiology, infectious disease and immunology, Faculty of Medicine, Université Laval, Quebec City, Quebec, Canada, G1V 0A6;

[3] Oncology Division, Centre Hospitalier Universitaire (CHU) de Québec-Université Laval Research Center, Quebec City, Quebec, Canada, G1R 2J6;

[4] Department of Molecular biology, Medical Biochemistry and Pathology, Faculty of Medicine, Université Laval, Québec City, Quebec, Canada, G1V 0A6;

[5] Université Laval Cancer Research Center, Université Laval, Quebec City, Quebec, Canada, G1R 3S3;

[6] Current location: INSERM, Centre d'Étude des Pathologies Respiratoires (CEPR), UMR 1100, Université de Tours, Tours, France;

[7] These authors contributed equally to this work.

*Co-corresponding authors

Author Contributions: V.C., E.B., V.T., L.F., and A.F.-T. designed research; V.C., E.B., V.T., E.G.L., J.D., A.B., and A.G. performed research and analyzed data; V.C. and A.F.-T. wrote the original draft and E.B., V.T., and L.F. edited the manuscript.

Competing Interest Statement: The authors declare that they have no conflict of interest.

Classification: Biological sciences, Microbiology

Keywords: DNA double-strand break signaling, telomere; integration; human herpesvirus 6A/B; immediate-early protein IE1.

Co-corresponding authors' information:

Amélie Fradet-Turcotte

CRCHU de Québec Research center – Hotel-Dieu de Québec

9 McMahon, room 3744, Québec, Québec, Canada, G1R 2J6

Tel : 1-418-525-4444 x16473

amelie.fradet-turcotte@crchudequebec.ulaval.ca

Louis Flamand

CRCHU de Québec Research center – CHUL

2705 boulevard Laurier, T1-64, Québec, Québec, Canada G1V 4G2

Tel : 1-418-525-4444 x46164

43   louis.flamand@crchudequebec.ulaval.ca

Abstract

Viral infection often trigger an ATM-dependent DNA damage response (DDR) in host cells that suppresses viral replication. To counteract this antiviral surveillance system, viruses evolved different strategies to induce the degradation of the MRE11/RAD50/NBS1 (MRN) complex and prevent subsequent DDR signaling. Here, we report that human herpesvirus 6B (HHV-6B) infection causes genomic instability by suppressing the host cell's ability to induce ATM-dependent signaling pathways. Expression of immediate early protein 1 (IE1) phenocopies this phenotype and blocks further homology-directed double-strand break (DSB) repair. In contrast to other viruses, IE1 does not affect the stability of the MRN complex. Instead, it uses two distinct domains to inhibit ATM serine/threonine kinase (ATM) activation at DSBs. Structure-based analyses revealed that the N-terminal domain of IE1 interacts with the BRCA1 C-terminal domain 2 of nibrin (NBN, also known as NBS1), while ATM inhibition is attributable to on its C-terminal domain. Consistent with the role of the MRN complex in antiviral responses, NBS1 depletion resulted in increased HHV-6B replication in infected cells. However, in semi-permissive cells, viral integration of HHV-6B into the telomeres was not strictly dependent on NBS1, supporting models where this process occurs via telomere elongation rather than through DNA repair. Interestingly, as IE1 expression has been detected in cells of subjects with inherited chromosomally-integrated form of HHV-6B (iciHHV-6B), a condition associated with several health conditions, our results raise the possibility of a link between genomic instability and the development of iciHHV-6-associated diseases.

Significance Statement

Many viruses have evolved ways to inhibit DNA damage signaling, presumably to prevent infected cells from activating an antiviral response. Here, we show that this is also true for human herpesvirus 6B (HHV-6B), through its immediate early protein 1 (IE1). However, in contrast to adenovirus' immediate early proteins, HHV-6B IE1 is recruited to double-strand breaks in an NBS1-dependent manner and inhibits ATM serine/threonine kinase activation. Characterizing this phenotype revealed a unique mechanism by which HHV-6B manipulates DNA damage signaling in infected cells. Consistently, viral replication is restricted by the MRN complex in HHV-6B infected cells. Viral integration of HHV-6B into the host's telomeres is not strictly dependent on NBS1, challenging current models where integration occurs through homology-directed repair.

Introduction

To infect a cell, a virus needs to successfully replicate its genetic material and produce new virions. Accordingly, cells have a sophisticated surveillance system that detects viral DNA and activates an innate antiviral response. Mounting evidence supports a role for the DNA damage response (DDR) in this process, revealing an intricate interplay between its activation and the activation of intrinsic antiviral responses (1). Some viruses, such as adenovirus, target the MRE11-RAD50-NBS1 (MRN) complex for degradation to prevent the activation of ATM serine/threonine kinase (ATM) (2, 3), while others, such as herpes simplex virus 1 (HSV-1) and human papillomavirus, rely on these proteins for efficient viral replication (4–6). How and why viruses inhibit or hijack the ATM pathway remains a mystery (7).

In mammalian cells, the MRN complex and ATM are essential to maintain genomic stability in the presence of DNA double-strand breaks (DSBs). Broken DNA ends are first detected by the MRN complex (8), where its accumulation induces a signaling cascade that activates ATM and subsequent phosphorylation of the histone variant H2AX on Ser139 (producing γ-H2AX). Mediator of DNA damage checkpoint 1 (MDC1) interacts with γ-H2AX, triggering the ubiquitylation of chromatin surrounding the break by promoting the accumulation of the E3-ubiquitin ligases ring finger protein (RNF) 8 and RNF168 (9, 10). In the G1 phase of the cell cycle, the recruitment of the DNA repair factor tumor protein p53 binding protein 1 (53BP1) at ubiquitylated chromatin promotes DNA repair *via* non-homologous end-joining (NHEJ)(11). In the S and G2 phases, BRCA1 DNA repair associated (BRCA1) and RB binding protein 8 endonuclease (CtIP) accumulate at the break and cooperate with exonuclease 1 (EXO1), BLM RecQ like helicase, and DNA replication helicase/nuclease 2 (BLM-DNA2) to facilitate end resection. This process results in extensive single-stranded (ss) DNA accumulation, which ultimately triggers the recruitment of recombinases that drive the homology searching required for homology-driven recombination (HDR) (12). HDR uses homologous sequences as templates to repair breaks in a faithful manner and includes processes such as homologous recombination (HR), single-stranded annealing (SSA), and break-induced replication (BIR) (11, 13–15).

Human herpesvirus 6B (HHV-6B) is a betaherpesvirus that infects nearly 90% of the world's population in the first 2 years of life and is responsible for roseola infantum, a pathology defined by skin rashes, high fever, and respiratory distress (16–18). In this double-stranded DNA (dsDNA) virus subfamily, HHV-6B shares 90% homology with HHV-6A, another lymphotropic virus. Although both infect CD4$^+$ T lymphocytes, they have epidemiological, biological, and immunological differences (19). Like other herpesviruses, HHV-6A and HHV-6B (HHV-6A/B) establish lifelong latency in infected hosts and can occasionally reactivate (20). However, whereas most herpesviruses achieve latency by circularizing and silencing their genome, HHV-6A/B integrate their genomes into the host's chromosomal terminal repeats (telomeres) (21, 22). The linear dsDNA genomes of HHV-6A/B are both flanked by an array of direct repeats containing 15–180 reiterations of perfect telomeric repeats (pTMRs) identical to the human telomeric sequence (5′-TTAGGG-3′), enabling viral integration (23). This process depends on the integrity of these pTMRs (23), resulting in a model in which viral integration is mediated through HDR, including the SSA or BIR DNA repair mechanisms (24). When HHV-6A/B integration occurs in a gamete before fertilization, the newborn carries a copy of HHV-6A/B in every cell of its body and can be transmitted to its offspring. This condition, called inherited chromosomally-integrated (ici)HHV-6A/B, affects ~1% of the world's population, representing almost 80 million people (25, 26). It is more prevalent in

those suffering from health issues such as high spontaneous abortion rates (27, 28), pre-eclampsia (29), and
angina pectoris (30) compared to healthy subjects (reviewed in (31, 32)). However, the reason why iciHHV-
6A/B contributes to these clinical syndromes has not been elucidated in any detail.

HHV-6B, which is better characterized than HHV-6A, sequentially expresses more than 97
genes/proteins during its lytic cycle(33). Immediate early (IE) proteins are expressed early in the viral cycle, to
regulate viral gene expression and establish a favorable environment for viral replication. In the context of HHV-
6B, IE protein 1 (IE1) is the first protein expressed during cell infection (34). It inhibits the innate antiviral
response in part by sequestering signal transducer and activator of transcription 2 (STAT2) in the nucleus,
thereby compromising type I interferon production and signaling(35, 36). In infected cells, IE1 is exclusively
localized within PML bodies (37), which were recently implicated in HDR-mediated DNA repair through an
undefined mechanism (38–41). Interestingly, PML depletion reduces HHV-6B integration (42), suggesting that
IE1-containing PML bodies also participate in viral integration.

In this study, we report that HHV-6B infection—and more specifically, IE1 expression—leads to genomic
instability in cells. Further investigations revealed that IE1 specifically prevents phosphorylation of the histone
variant H2AX and subsequent HDR repair. Structure-function analyses reveal that IE1 interacts with NBS1 and
inhibits ATM. Consistent with a role for the MRN complex in interfering with viral replication, we show that NBS1
depletion results in increased HHV-6B replication in infected cells. Although current models propose that viral
integration occurs through HDR DNA repair, we show that viral integration is not affected in NBS1-depleted
cells that elongate their telomeres in a human telomerase reverse transcriptase (hTERT)-dependent manner
(42). Thus, our findings reveal that viral integration relies on biological pathways that safeguard telomere
extension in infected cells and not on specific DNA repair pathways.

Results

HHV-6B infection and IE1 expression induce genomic instability

Our first indication that HHV-6B infection induces genomic instability in host cells was the observation
that HHV-6B infection rapidly induced micronuclei (MNi) formation in MOLT-3 cells (a lymphoblast T cell line;
Fig.1*A* and *SI Appendix*, Fig. S1*A*). In these experiments, cells infected with the HHV-6B strain Z29 accumulated
6.6-fold more MNi than non-infected cells (Mock) 24 h post-infection (Fig.1*A*). Consistent with the rapid
accumulation of genomic instability in infected cells, we observed a similar phenotype in duplicate clones of
stable U2OS cell lines containing a doxycycline (Dox)-inducible expression cassette for IE1 (C10 and C102;
Fig. 1*B* and *SI Appendix*, Fig. S1*B-C*). These clones had 4.8-fold more MNi than parental U2OS cells 48 h post-
IE1 induction (Fig.1*B*), suggesting that the instability observed in HHV-6B infected cells is at least partially
caused by IE1. Note that both U2OS clone (C10 and C102) exhibit similar levels of MNi than the parental cell
line without IE1 induction (*SI Appendix*, Fig. S1*C*). MNi arise from unresolved genomic instabilities such as
DSBs (i), lagging chromosomes (ii), and ruptured anaphase bridges (ABs) (iii) (Fig. 1*C*) (43). They are
compartmentally separated from the primary nucleus and surrounded by a nuclear envelope, as shown by the
presence of lamin B in these perinuclear structures (*SI Appendix*, Fig. S1*D-E*) (44). To determine how IE1
triggers MNi formation, we analyzed the accumulation of different markers in IE1-induced MNi, such as

centromeres and telomeric DNA (to detect lagging chromosomes and ABs caused by telomere fusion,

respectively). Interestingly, a much lower proportion of the IE1-induced MNi contained centromeres compared

with those in parental U2OS cells (~10-fold, Fig. 1*D* and *SI Appendix*, Fig. S1*F*), suggesting that they are not

induced by chromosome segregation defects. Although IE1 partially colocalizes with telomeres in host cells

(42), fluorescence *in situ* hybridization (FISH) revealed that IE1 does not specifically promote instability at

telomeres, as IE1-induced MNi accumulated similar levels of telomeric DNA as those in parental U2OS cells

(Fig. 1*E* and *SI Appendix*, Fig. S1*G*). Metaphase spread assays revealed that IE1-expressing U2OS cell lines

exhibited higher frequencies of DNA breaks than parental cells (Fig. 1*F-G*), consistent with MNi being induced

by DSB accumulation. Interestingly, IE1 was detected in only 5–10% of the MNi (*SI Appendix*, Fig. S1*H*),

suggesting that they are not arising from DSBs induced by the physical binding of IE1 at any defined DNA locus.

HHV-6B IE1 impairs DSB signaling and homology-directed DNA repair

DSB accumulation results from either an increase in DNA breaks or defective DNA DSB signaling and

repair. To determine how HHV-6B infection promotes genomic instability, we first investigated whether infected

cells accumulated Ser139-phosphorylated H2AX (*i.e.*, the DSB marker γ-H2AX). Surprisingly, while γ-H2AX foci

accumulated in > 75% of non-infected MOLT-3 cells exposed to irradiation (IR), this number was dramatically

reduced in infected cells (Fig. 2*A-B* and *SI Appendix*, Fig. S2*A*). U2OS clones stably expressing IE1 reproduced

this phenotype (Fig. 2*C-D* and *SI Appendix*, Fig. S2*B*), indicating that IE1 impairs DSB signaling. This inhibition

is independent of IE1 accumulation within PML bodies, as no γ-H2AX foci were detected in PML-deficient U2OS

cells that transiently express IE1 (*PML$^{-/-}$*, Fig. 2*E* and *SI Appendix*, Fig. S2*C-E*).

DSB signaling is essential to activate DNA repair pathways. Therefore, we tested whether DNA repair

is inhibited in HHV-6B IE1-expressing cells. As mammalian cells use several pathways to repair DSBs (45), we

performed a panel of DNA repair reporter assays in cells with stable or transient IE1 expression. These reporter

assays all rely on the detection of a fluorescent protein that is expressed only if a site-specific DSB (induced by

either I-SceI or Cas-9) is adequately repaired (Fig. 3*A-D* and *SI Appendix*, Fig. S3*A*, top panels) (46, 47). In the

DR-GFP (direct repeats) and CRISPR-LMNA HDR assays, DSBs repaired by HR either reconstitute a defective

green fluorescent protein (GFP)-reporter transgene integrated into the genome (DR-GFP) or introduce a

cassette expressing mRuby in frame with endogenous lamin A (CRISPR-LMNA HDR), respectively (48, 49). In

SA-GFP (single-strand annealing), proper annealing of a small homologous stretch reconstitutes a truncated

GFP (48). In BIR-GFP (break-induced replication), replication-mediated repair following homology searching

places a GFP coding sequence in the correct orientation (15, 50). Finally, in NHEJ-GFP, the ends of two DSBs

need to be correctly ligated to recreate a full-length (FL) GFP-expressing cassette (51, 52). In assays using

reporter transgenes integrated into the genome (DR-GFP, SA-GFP, BIR-GFP, and NHEJ-GFP) (Fig. 3*A-D* and

SI Appendix, Fig. S3*B-C*), a condition without endonuclease I-SceI or Cas-9 was used as a negative control

and the percentage of fluorescent cells obtained with I-SceI was set to 1. In each condition, a small amount of

a near-infrared fluorescent protein (iRFP)-expressing vector was transfected with I-SceI, IE1, or an empty vector

(EV) to ensure that DNA repair was measured in transfected cells only, and DNA repair was assessed 48 or 72

h post-transfection. Finally, as the clonal BIR-GFP U2OS cell line was generated in this study using a previously

described BIR-GFP reporter plasmid(50), we used short interfering (si)RNAs against RAD51 and RAD52 as

additional controls (*SI Appendix*, Fig. S3*D-F*) (13). As expected, the BIR-GFP signal was specifically inhibited
in cells depleted of RAD51 (50).

Interestingly, these analyses revealed that both transient and stable IE1 expression drastically reduced
all types of homology-directed DNA repair (Fig. 3*A-C*, lower panels, and *SI Appendix*, Fig. S3*A-C*). In contrast,
IE1 only slightly modulated DNA repair in reporter assays assessing NHEJ (Fig. 3*D*). As the choice between
HDR and NHEJ is driven by the cell cycle (45), we confirmed that cell cycle progression was not affected in
cells expressing IE1 (*SI Appendix*, Fig. S3*G*). Altogether, these results show that homology-based DNA repair
is specifically inhibited in cells expressing HHV-6B IE1.

HHV-6B IE1 interacts with NBS1 and inhibits its ability to promote ATM activation

At DSBs, homology-based DNA repair is initiated when lesions are detected by the MRN complex,
which leads to ATM auto-activation (Fig. 4*A*) through a still poorly understood mechanism (8, 53, 54). In
adenovirus-infected cells, ATM activation is inhibited through the degradation of the MRN complex, which is
mediated by the protein E4 (2). In contrast, MRN complex components were stable at steady state upon IE1
induction in U2OS clones (Fig. 4*B*). Interestingly, immunofluorescence analyses revealed that IE1 colocalizes
with NBS1 in ~75% of IE1-expressing cells (Fig. 4*C-E* and *SI Appendix*, Fig. S4*A-B*). This colocalization was
also detected in irradiated cells and *PML* knockout cells (*SI Appendix*, Fig. S4*C-E*), suggesting that the
interaction between IE1 and NBS1 is constitutive and independent of PML bodies. We also observed
constitutive colocalization between MRE11 and IE1 (Fig. 4*D-E* and *SI Appendix*, Fig. S4*A-B, F*). However, the
colocalization of transiently expressed IE1 with MRE11 was greatly reduced upon NBS1 depletion (Fig. 4*F and*
SI Appendix, Fig. S4*G-H*), supporting a model where IE1 colocalizes with the MRN complex by interacting with
NBS1. Importantly, we confirmed that colocalization between IE1 and NBS1 is also observed in HHV-6B-
infected MOLT-3 cells (Fig. 4*G-H*).

To further characterize the interplay between IE1 and the MRN complex, we took advantage of a cell-
based assay that quantifies the ability of a mCherry-LacRnls fusion protein to specifically induce the recruitment
of a "prey" to a *LacO* array integrated at a single locus in U2OS 2-6-5 cells (*(i)* No DSBs, Fig. 5A) (55, 56). This
system can also be used to study signaling at DSBs by recruiting the ER-mCherry-LacR-FOKI-DD
endonuclease to the *LacO* array (*(ii)* Localized DSBs, Fig. 5A). Although ER-mCherry-LacR-FOKI-DD is
constitutively expressed in U2OS 2-6-5 cells, the protein is cytoplasmic, and a C-terminal destabilization domain
(DD) ensures its continual degradation (55). DSBs can be rapidly induced by adding 4-hydroxytamoxifen (4-
OHT) and Shield-1 to the culture medium. 4-OHT induces nuclear relocalization of ER-mCherry-LacR-FOKI-DD
via its modified estrogen receptor (ER) domain and Shield-1 stabilizes it by inactivating the DD. When the
mCherry-LacRnls-IE1 fusion protein was transiently expressed in U2OS 2-6-5 cells, only NBS1 was efficiently
recruited to the *LacO* (Fig. 5*B-C* and *SI Appendix*, Fig. S5*A-E*). No DDR signaling proteins were recruited to the
LacO by the negative control, mCherry-LacRnls. As a positive control, we added 4-OHT and Shield-1 to the
medium for 6 h, and readily detected ATM, phospho(p)-ATM (Ser1981), γ-H2AX, RAD50, NBS1, and MRE11.
No pATM or γ-H2AX signals were detected at the array upon recruitment of mCherry-LacRnls-IE1, consistent
with a constitutive interaction between IE1 and NBS1 that is independent of DSB signaling. Intriguingly, while
an mCherry-LacRnls-NBS1 fusion protein is sufficient to induce ATM recruitment and activation, as well as

subsequent H2AX phosphorylation at the *LacO* array in NIH-3T3 cells (57), NBS1 recruitment by mCherry-
LacRnls-IE1 did not trigger ATM activation (Fig. 5B and *SI Appendix*, Fig. S5*B*). To further validate that IE1
inhibits the ability of NBS1 to activate ATM at the *LacO*, we transiently transfected a mCherry-LacRnls-NBS1
fusion protein and an untagged or FLAG-tagged IE1 vector in U2OS 2-6-5 cells. As expected, a full length (FL)
NBS1 construct (aa 1–754) specifically promoted ATM and H2AX phosphorylation at the *LacO* in approximately
75% and 50% of cells, respectively (Fig. 5*D-F* and *SI Appendix*, Fig. S5*F-H*). This function depends on its ability
to bind ATM, as an NBS1 ATM binding deficient construct (∆A) (aa 1–733) produced similar γ-H2AX levels as
the negative control (Fig. 5*D* and *SI Appendix*, Fig. S5*F*). Interestingly, NBS1-dependent accumulation of γ-
H2AX and pATM was strongly inhibited in cells expressing untagged or FLAG-tagged IE1 (Fig. 5*D-F* and *SI*
Appendix, Fig. S5*F-H*). These findings suggest that the interaction between IE1 and NBS1 at the array directly
prevents ATM activation and subsequent H2AX phosphorylation. In support of this model, we found that IE1
accumulates at the *LacO* array upon DSB induction in an NBS1-dependent manner (Fig. 5*G-H*), an observation
that can only be made in this system, as no marker of DSB signaling can be used to detect IE1 accumulation at
endogenous DSBs. In these conditions, IE1 colocalizes with 60% of mCherry-LacR-FOKI foci and this amount
is reduced to 30% in cells treated with a siRNA against NBS1.

245 Two distinct domains of IE1 interact with NBS1 and prevent ATM activation

The functional domains of IE1 are not well characterized aside from its STAT2 binding domain (aa 270–
540; Fig. 6*A*) (36). Guided by its secondary structure, we designed a series of IE1 fragments that we fused to
mCherry-LacRnls to assess their ability to recruit endogenous NBS1 to the *LacO* array (*SI Appendix*, Fig. S6*A*).
The fusion encoding aa 1–540 was the smallest fragment capable of recruiting NBS1 at the *LacO* array as
efficiently as FL IE1 (~81% of mCherry-LacRnls-IE1 1–540 colocalized with NBS1; Fig. 6*B-C* and *SI Appendix*,
Fig. S6*B*). All attempts to generate smaller fragments of this N-terminal domain of IE1 resulted in unstable
proteins in our hands. Interestingly, the 1–540 fragment was also the smallest to efficiently accumulate in PML
bodies (*SI Appendix*, Fig. S6*C-D*) suggesting that both functions are related. Consistently, we observed a
reduced accumulation of IE1 at PML bodies in cells treated with siNBS1 (*SI Appendix*, Fig. S4*E-F*).
To determine if the IE1-NBS1 interaction is sufficient to prevent ATM activation, we transiently
transfected expression vectors containing the different fragments of IE1 into U2OS cells (without the *LacO* array)
and quantified their ability to inhibit H2AX phosphorylation in irradiated cells. Surprisingly, the IE1 N-terminus
alone was unable to prevent the accumulation of γ-H2AX foci (Fig. 6*D-E* and *SI Appendix*, Fig. S6*G*). Instead,
we found that this function depends on a fragment of 268 amino acids in the IE1 C-terminus. The 810–1078
fragment inhibited H2AX phosphorylation as efficiently as the FL protein (~75% of cells transfected with
mCherry-LacRnls-IE1 810–1078 had <10 γ-H2AX foci 1 h post-irradiation with 1 Gy; Fig. 6*D-E* and *SI Appendix*,
Fig. S6*G*). Together, these results show that IE1 interacts with NBS1 and blocks ATM activation using two
distinct motifs: an N-terminal NBS1-binding domain (NBS1-BD) and a C-terminal domain that independently
inhibits the ability of NBS1 to trigger DSB signaling (Fig. 6*A*), which we have named ATM-inhibitory domain
(ATMiD).
NBS1 encodes a 95-kDa protein with multiple domains, which are required for its recruitment to DSBs
and its interactions with the ATM and ATR (58). Briefly, NBS1 contains a forkhead-associated (FHA) domain

and two breast cancer C-terminal domains (BRCTs) that are required for optimal phospho-dependent NBS1
accumulation at DNA breaks. The C-terminus contains a domain that promotes its interactions with MRE11
(MRE11-binding motif, MBM) and ATM (ATM-binding motif, ATM-BM; Fig. 6*F*). Interestingly, NBS1 also
contains an intrinsically disordered domain (IDD) that drives a species-specific interaction with the HSV-1 IE
protein ICP0 (59). To determine if this domain also promotes the interaction between NBS1 and IE1, we used
the same approach used to map the IE1-NBS1 interaction (Fig. 6*A*-*C*). Different fragments of NBS1 were fused
with the mCherry-LacRnls protein and co-expressed with an untagged version of IE1 (Fig. 6*F*-*H* and *SI*
Appendix, Fig. S6*H-M*). The mCherry-LacRnls-NBS1 construct lacking the BRCT2 domain (ΔB2, Δaa 201–327)
was unable to recruit IE1 to the array, while the construct containing only this domain was sufficient for the
interaction (Fig. 6*F*-*H* and *SI Appendix*, Fig. S6*J, M*). Consistently, immunoprecipitation of FLAG-tagged IE1
from U2OS cell lysates revealed an interaction with FL mCherry-LacRnls-NBS1 but not the ΔB2 fusion (Fig. 6*I*).
Using the LacR system, we noted that the mCherry-LacRnls-NBS1 fusion lacking the linker region of NBS1 (ΔL,
Δ328–638) significantly reduced the interaction between NBS1 and IE1 (*SI Appendix*, Fig. S6*H, J-M*). In contrast
with the BRCT2 domain, the linker alone was unable to recruit IE1 to the *LacO* array (*SI Appendix*, Fig. S6*H, J-*
M).

Altogether, our results support a model where the N-terminus of IE1 interacts with the BRCT2 domain
of NBS1 and the C-terminus of IE1 blocks ATM activation. In the LacR system, IE1 did not interact with the
domain of NBS1 that interacts with ATM (ATM-BM, aa 733–754) (Fig. 6*F*-*H* and *SI Appendix*, Fig. S6*I, L*) or
with ATM itself (Fig. 5*B*). The latter observation suggests that IE1 does not interfere with the NBS1-dependent
activation of ATM by directly competing for interactions between them or that the interaction is too weak to be
detected in our experimental setting.

290 HHV-6B integration relies on a pathway that safeguards telomere elongation

Depending on the virus, the MRN complex is either required for viral replication or it inhibits it (7). As
HHV-6B IE1 interacts with NBS1 and blocks ATM activation, the complex is likely detrimental for its replication.
HHV-6B infection has different outcomes depending on the nature of the infected cells (Fig. 7*A*). In permissive
cells (*e.g.*, MOLT-3), viral protein expression promotes replication (the lytic state). In contrast, in semi-
permissive cells, integration of the viral genome into the host's telomeres is favored, and this process has been
proposed to rely on HDR processes in the infected cells (24). To understand the interplay between HHV-6B,
DSB signaling, and HDR repair, we investigated the impacts of depleting NBS1 on viral replication and
integration. In these experiments, we depleted NBS1 from permissive cells (MOLT-3) and semi-permissive cells
(U2OS, HeLa, and GM847) by shRNA (*SI Appendix*, Fig. S6*A-D*). MOLT-3 cells treated with control and NBS1
shRNA were infected with HHV-6B and viral DNA was quantified over time by qualitative polymerase chain
reaction (qPCR; Fig. 7*B*). Viral DNA replication was 1.67-fold higher in MOLT-3 cells depleted of NBS1 72 h
post-infection (note that this is an underestimate, as CellTiter-Glo® assays revealed that the shRNA against
NBS1 was toxic in MOLT-3 cells, Fig. 7*C*). Viral integration was assessed in two types of semi-permissive cells:
HeLa cells, which lengthen their telomeres *via* hTERT-dependent mechanisms, and U2OS and GM847 cells
which rely on ALT, a telomerase-independent mechanism that uses HDR pathways for telomere elongation (60,
61). All cell lines were infected with HHV-6B at a multiplicity of infection (MOI) of 1 and passaged for 4 weeks

prior to DNA extraction and viral genome quantification by droplet digital (dd)PCR (62). Interestingly, levels of
viral integration were approximately 6-fold higher in HeLa cells depleted of NBS1 than in control HeLa cells
(Table 1). In contrast, the integration frequency was decreased by at least 2-fold in U2OS and GM847 cell lines
depleted of NBS1 vs the control lines. This difference resembles the lower integration level measured in U2OS
PML$^{-/-}$ cells, a condition previously reported to reduce viral integration (Table 1) (42). NBS1 depletion did not
further impact viral integration in these conditions. Lastly, the differences in viral integration levels between the
semi-permissive cell lines were not artefactually driven by cell death, as shNBS1 slightly decreased the viability
of all cell types used in this study (Fig. 7*C* and *SI Appendix*, Fig. S7E). Altogether, these results are consistent
with the need for functional NBS1-dependent HDR repair pathways to promote integration in ALT$^+$ cells and
support a model where viral integration in semi-permissive cells relies on the molecular mechanisms that drive
telomere elongation rather than specific DNA repair mechanisms.

Discussion

In this study, we sought to understand how HHV-6B manipulates factors that safeguard genomic
instability in infected cells, as well as its impacts on two key events of the viral life cycle: genome replication and
chromosomal integration. Using a series of microscopy- and cytometry-based approaches to track the source
of DNA breaks in infected cells and in cells expressing the HHV-6B immediate-early protein IE1, we found that
IE1 promotes the accumulation of DNA DSBs and inhibits their repair. Further structure-function analyses
revealed a molecular mechanism by which HHV-6B IE1 localizes to DSBs in an NBS1-dependent manner and
prevents HDR-mediated DNA repair by blocking ATM activation and subsequent DDR signaling. We report that
IE1 specifically interacts with NBS1 through an N-terminal NBS1-BD and that ATM activation by NBS1 is
strongly inhibited by a newly characterized domain of IE1, the C-terminal ATMiD.

ATM activation by the MRN complex requires conformational changes in ATM that expose its substrate-
binding site (53). Our findings show that, in contrast with NBS1, IE1 does not interact strongly with ATM in the
LacR-based system. Furthermore, the ATMiD domain of IE1 does not interact with NBS1. The exact mechanism
by which IE1 inhibits ATM activation thus remains unclear. Based on current literature, IE1 could be interfering
with ATM activation by preventing the interaction between the FxF/Y motif of NBS1 and ATM or by directly
blocking the substrate-binding site of ATM (53, 63). Alternatively, IE1 may block H2AX phosphorylation through
steric hindrance (*e.g.,* though chromatin binding). Viral proteins such as Kaposi's sarcoma-associated
herpesvirus LANA and adenovirus protein VII interfere with the activation of chromatin-dependent mechanisms
by directly interacting with the nucleosomes (64, 65). Further investigation will be required to elucidate how IE1
prevents ATM activation by the MRN complex.

The mechanism by which IE1 interacts with NBS1 and inhibits ATM signaling in cells differs from the
mechanisms by which other viruses manipulate this pathway (2, 59). The BRCT2 domain of NBS1 contributes
to its retention on DSBs (66), which may be reduced when IE1 binds this domain. However, the fact that the IE1
N-terminus is insufficient to inhibit ATM signaling suggests otherwise. In this study, we show that IE1 is recruited
to DSBs in an NBS1-dependent manner and that ectopic expression of the IE1 ATMiD is sufficient to inhibit
DSB signaling. Thus, it is still unclear whether IE1 needs to accumulate at DSBs in an NBS1-dependent manner

to inhibit ATM when expressed in infected cells at lower levels, or if NBS1 interaction and ATM inhibition are independent functions of IE1. A model in which IE1 inhibits ATM activation through a bi-partite mechanism is appealing, as it would provide a way for HHV-6B to inhibit ATM signaling at specific loci. This would support a recently proposed concept in which viruses prevent local ATM signaling on the viral genome and restrict viral replication, while avoiding a global inhibition of the DSB signaling cascade in infected cells (67). During the lytic cycle, the accumulation of genomic instability in the host cell genome is not a problem as these cells will die upon the lysis provoked by the virus to release new virus particles. However, more selective inhibition of ATM by IE1 during the latent cycle of HHV-6B or from iciHHV-6B would avoid a detrimental accumulation of genomic alterations in host cells. This model would be consistent with the fact that HHV-6B is not associated with a higher frequency of cancer development, as would be expected if global DSB signaling was inhibited in these cells. Alternatively, expression of IE1 upon the exit of latency may inhibit global DSB signaling, but this phenomenon is restricted to the early stages of the process, thereby minimizing the impact on the host cell's genomic stability.

In addition to its role during viral infection, Peddu et al. used RNA-seq approach to show that *IE1 (U90)* is among the most abundantly expressed genes in a variety of tissues from iciHHV-6A/B+ individuals (68). Spontaneous and inducible IE1 protein expression from integrated HHV-6A/B genomes has also been documented (62), raising the possibility that IE1 expression from integrated genomes might contribute to the development of iciHHV-6A/B associated diseases by inducing genomic instability in these cells. At present, conditions associated with iciHHV-6A/B status include increased spontaneous abortion rates (27, 28), pre-eclampsia (29) and angina pectoris (30). Further characterization of the proteins expressed from integrated genomes as well as the diseases associated with these conditions will be required to strengthen our understanding of the consequences associated with viral latency in iciHHV-6A/B subjects. Importantly, the intricate interplay between IE1, the MRN complex, and ATM pathway activation will need to be studied in a spatiotemporal manner to elucidate when and how IE1 manipulates this important pathway during viral infection and integration. Further efforts will also be required to determine if ATM inhibition by IE1 contributes to its ability to block type I interferon signaling in infected cells (36). From a mechanistic point of view, it will be interesting to investigate if the interaction between IE1 and NBS1's BRCT2 domain—a phospho-recognition domain—is regulated by phosphorylation (66, 69–72). Finally, the model presented here assumes that NBS1 and ATM activity must be inhibited to prevent their detrimental effect on viral replication. However, it is impossible to rule out that enhanced viral replication and integration result from the increased level of genomic instability induced in host cells upon viral infection. Further studies will be required to address this question.

In germline, hematopoietic, stem, and rapidly renewing cells, telomere elongation relies on hTERT, a polymerase that catalyzes the extension of telomeric DNA repeats using RNA as a template (73). While hTERT is negatively regulated in healthy somatic cells, cancer cells can overcome senescence either through its re-activation or by an alternative homology-directed mechanism called ALT(60). The HHV-6B genome contains conserved telomeric sequences that are required for viral integration (23). In this study, we show that HHV-6B integration is independent of NBS1 in ALT⁻ cells but dependent on NBS1 in ALT⁺ cells. These findings are consistent with previous reports showing that the telomerase complex is required for optimal HHV-6B integration (74) and with the reported role of NBS1 in ALT (75, 76). While PML is not required for the IE1-NBS1 interaction or the ability of IE1 to inhibit H2AX phosphorylation (this study), NBS1 is required for the assembly of functional

ALT-associated PML bodies (77). These concomitant roles are in line with the absence of phenotypes
associated with NBS1 depletion in integration assays performed on *PML*$^{-/-}$ ALT^{+} U2OS cells. Intriguingly, we
previously reported that *PML* knockout also reduces integration in ALT^{-} HeLa cells, reinforcing the hypothesis
that PML plays an ALT-independent role in this process (42). Further studies will be required to elucidate this
function.

Consistent with previous findings showing that HHV-6B integration is not altered upon inhibition of
RAD51 (78, 79), we found that IE1 inhibits HDR processes, and that integration is independent of NBS1 in ALT-
cell lines. Together, these observations argue against models where integration mechanisms rely on RAD51-
dependent BIR or SSA (24). However, it is important to note that all homology-directed reporter assays used in
this study rely on extensive DNA end resection following nuclease-induced breakage, a process that is
dependent on NBS1 (80). Thereby, integration models where SSA or RAD51-independent BIR trigger
integration following extensive accumulation of ssDNA generated at stalled replication forks remain plausible.
One attractive model is that HHV-6B integration occurs during mitotic DNA synthesis (MiDAS), a RAD52-
dependent BIR mechanism that is initiated by replication fork stalls that remain unresolved at the start of
mitosis—a problem often observed at hard-to-replicate loci, including the telomeres (13, 27, 81). Alternatively,
upon cell entry but before viral genome circularization (and before IE1 is expressed), the viral genome may be
perceived as broken DNA. Under such circumstances, the MRN complex would be recruited to the ends of the
viral genome and initiate 3′→ 5′ resections. The ssDNA ends of eroded telomeres (no longer efficiently protected
by the shelterin complex) could anneal to the near-terminal telomeric sequence at the right end of the genome
in a process analogous to an ALT mechanism described in yeast (reviewed in (13)). Once the entire viral
genome is copied, the telomeric repeats at the left end of the genome would serve as a template for the hTERT
and ALT mechanisms to regenerate a telomere of appropriate length (82).

In conclusion, we provide a detailed characterization of the HHV-6B IE1 protein as an efficient inhibitor
of DSB signaling and DDR that contributes to the favorable establishment of a productive infection. Despite
being a relatively abundant protein expressed very early upon entry, the functions of IE1 remain poorly defined.
IE1 shares very little sequence homology with proteins from other herpesviruses (except HHV-6A and HHV-7)
meaning that deductions based on primary sequence analysis are very limited. Our work adds to the growing
knowledge surrounding HHV-6B integration processes and the potential importance of IE1 during infection.

Materials and Methods

RNA interference
A SMARTPool siRNA targeting RAD51, single siRNA duplexes targeting NBS1, and a non-targeting single
siRNA duplex were purchased from Dharmacon (Horizon). Single siRNA duplexes targeting RAD52 were a kind
gift from Jean-Yves Masson (Université Laval, Québec, Canada). siRNAs were forward-transfected 24 h prior
to cell processing using RNAimax (Invitrogen) according to the manufacturer's protocol. Plasmids carrying an
NBS1 shRNA (Open Biosystems) or a control shRNA (Sigma) in the pLKO background backbone were a kind
gift from Cary A. Moody (6). Lentiviruses were produced as previously described (6). Briefly, plasmids
expressing shRNAs with vesicular stomatitis virus G (pMD2.g) and lentiviral packaging (pPAX) plasmids were

co-transfected into HEK293T cells using polyethyleneimine. Lentivirus-containing supernatants were harvested

48–72 h post-transfection, and U2OS, MOLT-3, HeLa, and GM847 cells were transduced in the presence of 8

µg/mL Polybrene (Sigma). For all relevant experiments, RAD51, RAD52, and NBS1 depletion was confirmed

by immunoblotting or qPCR analyses.

Cell cultures and transfections

Cell lines were maintained at 37°C and 5% CO_2. All culture media were supplemented with 10% fetal bovine

serum. MOLT-3 cells (American Type Culture Collection (ATCC)) were cultured in Roswell Park Memorial

Institute-1640 (RPMI) medium (Corning Cellgro), 8.85 mM HEPES, and 5 µg/mL plasmocin (Invivogen). GM847

and HeLa cell lines were obtained from ATCC and cultured in Dulbecco's modified Eagle's medium (DMEM;

Corning Cellgro), NEM (Corning Cellgro), 8.85 mM HEPES, and 5 µg/mL plasmocin. U2OS (ATCC), U2OS

PML[-/-] (42), U2OS 2-6-5 (a kind gift from Roger Greenberg, University of Pennsylvania, Philadelphia) (55), U2OS

DR-GFP, SA-GFP (a kind gift from Jeremy Stark, City of Hope National Medical Center, California) (48, 52),

and cell lines were cultured in McCoy's medium (Life Technologies).

Viral infection and integration assays

Viral infection was done as previously described (42) using HHV-6B (strain Z29) at an MOI of 1 (or not (Mock

samples)). At the indicated time points, cells were harvested and processed for DNA extraction using a QIAamp

DNA Blood Mini Kit (Qiagen) and analyzed by qPCR. Integration assays were performed as described

previously (62). Briefly, cells were infected with HHV-6B (MOI of 1) for 24 h and passaged for 4 weeks prior to

DNA extraction with the QIAamp DNA Blood Mini Kit for ddPCR.

PCR analyses

qPCR was performed as previously described (83). DNA was quantified using primers and probes against *U67-*

68 (HHV-6B) and ribonuclease P/MRP subunit p30 (*RPP30*; as a host reference gene). Data were normalized

against the corresponding genome copies of *RPP30*. ddPCR was used to quantify integration frequency as

previously described (62). Briefly, HHV-6B chromosomal integration frequencies were estimated assuming a

single integrated HHV-6/cell and calculated with the following formula: (number of HHV-6 copies)/(number of

RPP30 copies/2 copies per cell) × 100, as previously described (62). This assay has been extensively validated

and provides comparable data to single cell cloning and quantification.

Immunofluorescence microscopy

Immunofluorescence were done essentially as previously described for MOLT-3 (42) and U2OS (56) cells.

Briefly, cells were either fixed with 2% paraformaldehyde (PFA) or 100% MeOH prior to permeabilization and

incubation with primary antibody diluted in blocking buffer. DNA was counterstained with DAPI and the

coverslips were mounted onto glass slides with Prolong Diamond Mounting Agent (Invitrogen). Further

experimental details are provided as the Supplementary information.

FISH

Fixed cells were processed as described for immunofluorescence staining and then fixed for 2 min at room
temperature with 1% PFA/PBS. Coverslips were washed twice with PBS for 5 min and dehydrated for 5 min in
successive ethanol baths (70%, 95%, 100%). Once dried, coverslips were placed upside down on a drop of
hybridizing solution (70% formamide, 0.5% blocking reagent (Sigma, Cat:11096176001), 10 mM Tris-HCl pH
7.2, 1/1000 Cy5-TelC PNA probe (F1003, PNABio)). Samples were denatured for 10 min at 80°C on a heated
block, then incubated overnight at 4°C in the dark. After hybridization, coverslips were washed twice for 15 min
in washing solution (70% formamide; 10 mM Tris-HCl pH 7.2) and then washed three times for 5 min with PBS.
Coverslips were air-dried, counterstained with DAPI, washed with PBS, and mounted onto glass slides with
Prolong Gold Mounting Agent.
470
Metaphase spread analysis
U2OS SA-GFP HHV-6B IE1 cells were arrested in mitosis using 1 µM nocodazole for 3 h at 37°C and 5% CO_2.
Cells were then resuspended and incubated in pre-warmed hypotonic solution (KCl 75 mM, 15% fetal bovine
serum) at 37°C for 15 min to induce swelling and fixed in a 75% ethanol 25% acetic acid solution overnight at
4°C. Droplets of cells were spread onto glass slides pre-cooled to -20°C and dried overnight in the dark at room
temperature. Slides were then mounted with Vectashield Antifade Mounting Medium containing DAPI
(VECTH20002, MJS BioLynx Inc.). Images were acquired using a Zeiss LSM700 laser-scanning microscope
equipped with a 40× water lens. Quantification was performed on three biological replicates and 10 spreads
were quantified per experiment.
480
Immunoprecipitation
U2OS cells (1×10^7) were transfected with NBS1- or non-targeting single siRNA duplexes for 24 h, then co-
transfected with the indicated mCherry-LacR and 3×FLAG expression vectors. After 24 h, cells were lysed in
NETN lysis buffer (50 mM Tris pH 8.0, 150 mM NaCl, 1mM EDTA, 0.5% NP-40) complemented with 1×
complete, EDTA-free Protease Inhibitor Cocktail (Roche), 20 mM N-ethylmethylamine, 1 mM NaF, and 0.2 mM
Na_3VO_4. Cleared cell lysates were immunoprecipitated using 1 µg FLAG-M2 antibody coupled to 40 µL of
packed protein G Sepharose beads (Cat GE17-0618-01, Sigma) for 3 h at 4°C. Beads were washed four times
with NETN buffer and eluted in 2× Laemmli buffer for immunoblotting.
489
Reporter-based DNA repair assays
DR-GFP, NHEJ-GFP, SA-GFP, and BIR-GFP cell lines were plated at 125,000 cells/well in 6-well plates. After
24 h, cells were co-transfected with 900 ng of the I-SceI plasmid (pCBASceI, Addgene #26477) and 900 ng of
pcDNA4/TO-HHV-6B IE1 (+I-SceI, +IE1) or 900 ng of the pcDNA4/TO/Myc-His vector as a negative control (+I-
SceI, -IE1). The pcDNA4/TO/myc-His vector alone was transfected for conditions without IE1 and I-SceI (-I-
SceI/-IE1). A plasmid expressing iRFP (200 ng) was also transfected into all conditions to control for transfection
efficiency. After 48 h, cells were harvested and washed with PBS, and an Accuri C6 flow cytometer (BD
Biosciences) was used to quantify the GFP$^+$ cells in the iRFP$^+$ population. Data were analyzed using FlowJo.
The NHEJ-GFP (EJ7) assay was performed essentially as described above, but cells were co-transfected with
600 ng of each Cas9/sgRNA-expressing vector p330X-sgRNA7a, and p330X-sgRNA7b along with 600 ng of

pcDNA4/TO-HHV-6B IE1 or pcDNA4/TO/myc-His (52) and processed for flow cytometry analysis 72 h post-transfection.

Statistical analysis

Quantifications were performed on three biological replicates. Unless otherwise stated, one-way analysis of variance and Dunnett's multiple comparisons test were used to assess statistical significance.

Acknowledgements

We are grateful to Matthew D. Weitzman, Alexandre Orthwein, Cary A. Moody, and members of the Fradet-Turcotte and Flamand laboratories for critical reading of the manuscript; High-Fidelity Science Communications for manuscript editing, and Daniel Durocher, Jean-Yves Masson, Graham Dellaire, Roger Greenberg, and Jeremy Stark for essential reagents. E.B, and V.C. received postdoctoral and doctoral fellowships, respectively, from the Fonds de Recherche du Québec - Santé. V.T. received a master's fellowship from the Fonds de recherche Nature et technologies. This work was supported by three Canadian Institutes of Health Research Grants (PJT_152948 to A.F.-T.; MOP_123214 and PJT_156118 to L.F.). A.F.-T. is a Tier 2 Canada Research Chair in Molecular Virology and Genomic Instability and is supported by the Foundation J.-Louis Lévesque. We thank the Bioimaging platform of the Infectious Disease Research Centre, which is funded by an equipment and infrastructure grant from the Canadian Foundation for Innovation.

References

1. J. L. Justice, I. M. Cristea, Nuclear antiviral innate responses at the intersection of DNA sensing and DNA repair. *Trends Microbiol* **TBD** (2022).

2. T. H. Stracker, C. T. Carson, M. D. Weilzman, Adenovirus oncoproteins inactivate the Mre11-Rad50-NBs1 DNA repair complex. *Nature* **418** (2002).

3. S. S. Lakdawala, *et al.*, Differential Requirements of the C Terminus of Nbs1 in Suppressing Adenovirus DNA Replication and Promoting Concatemer Formation. *J Virol* **82** (2008).

4. C. E. Lilley, C. T. Carson, A. R. Muotri, F. H. Gage, M. D. Weitzman, DNA repair proteins affect the lifecycle of herpes simplex virus 1. *Proc Natl Acad Sci U S A* **102** (2005).

5. C. A. Moody, L. A. Laimins, Human papillomaviruses activate the ATM DNA damage pathway for viral genome amplification upon differentiation. *PLoS Pathog* **5** (2009).

6. D. C. Anacker, D. Gautam, K. A. Gillespie, W. H. Chappell, C. A. Moody, Productive Replication of Human Papillomavirus 31 Requires DNA Repair Factor Nbs1. *J Virol* **88**, 8528–8544 (2014).

7. M. D. Weitzman, A. Fradet-Turcotte, Virus DNA replication and the host DNA damage response. *Annu Rev Virol* **5** (2018).

8. A. Syed, J. A. Tainer, The MRE11-RAD50-NBS1 Complex Conducts the Orchestration of Damage Signaling and Outcomes to Stress in DNA Replication and Repair. *Annu Rev Biochem* **87** (2018).

9.   A. Fradet-Turcotte, *et al.*, 53BP1 is a reader of the DNA-damage-induced H2A Lys 15 ubiquitin mark.
Nature **499** (2013).

10.   F. Mattiroli, L. Penengo, Histone Ubiquitination: An Integrative Signaling Platform in Genome
Stability. *Trends in Genetics* (2021) https:/doi.org/10.1016/j.tig.2020.12.005.

11.   L. Krenning, J. van den Berg, R. H. Medema, Life or Death after a Break: What Determines the Choice?
Mol Cell **76** (2019).

12.   A. Maréchal, L. Zou, DNA damage sensing by the ATM and ATR kinases. *Cold Spring Harb Perspect Biol*
5 (2013).

13.   Z. W. Kockler, B. Osia, R. Lee, K. Musmaker, A. Malkova, Repair of DNA Breaks by Break-Induced
Replication. *Annu Rev Biochem* **90** (2021).

14.   R. Bhargava, D. O. Onyango, J. M. Stark, Regulation of Single Strand Annealing and its role in genome
maintenance Chromosomal break repair by the Single Strand Annealing (SSA) pathway. *Trends Genet*
32, 566–575 (2016).

15.   J. Kramara, B. Osia, A. Malkova, Break-Induced Replication: The Where, The Why, and The How.
Trends in Genetics **34**, 518–531 (2018).

16.   K. Yamanishi, *et al.*, Identification of Human Herpesvirus-6 As a Causal Agent for Exanthem Subitum.
The Lancet **331**, 1065–1067 (1988).

17.   D. M. Zerr, *et al.*, A Population-Based Study of Primary Human Herpesvirus 6 Infection. *New England*
Journal of Medicine **352**, 768–776 (2005).

18.   C. B. Hall, *et al.*, Congenital infections with human herpesvirus 6 (HHV6) and human herpesvirus 7
(HHV7). *Journal of Pediatrics* **145**, 472–477 (2004).

19.   V. Collin, L. Flamand, HHV-6A/B integration and the pathogenesis associated with the reactivation of
chromosomally integrated HHV-6A/B. *Viruses* **9** (2017).

20.   S. N. Pantry, P. G. Medveczky, Latency, integration, and reactivation of human herpesvirus-6. *Viruses*
9 (2017).

21.   J. H. Arbuckle, *et al.*, Mapping the telomere integrated genome of human herpesvirus 6A and 6B.
Virology **442**, 3–11 (2013).

22.   J. H. Arbuckle, *et al.*, The latent human herpesvirus-6A genome specifically integrates in telomeres of
human chromosomes in vivo and in vitro. *Proc Natl Acad Sci U S A* **107**, 5563–5568 (2010).

23.   N. Wallaschek, *et al.*, The Telomeric Repeats of Human Herpesvirus 6A (HHV-6A) Are Required for
Efficient Virus Integration. *PLoS Pathog* **12**, 1–15 (2016).

24.   G. Aimola, G. Beythien, A. Aswad, B. B. Kaufer, Current understanding of human herpesvirus 6 (HHV-
6) chromosomal integration. *Antiviral Res* **176** (2020).

25.   K. Tanaka-Taya, *et al.*, Human herpesvirus 6 (HHV-6) is transmitted from parent to child in an
integrated form and characterization of cases with chromosomally integrated HHV-6 DNA. *Journal of*
Medical VirologyThe prevalence of chromosomally integrated human herpesvirus 6 genomes in the
blood of UK blood donors. **73**, 465–473 (2004).

26.   M. Daibata, T. Taguchi, Y. Nemoto, H. Taguchi, I. Miyoshi, Inheritance of chromosomally integrated
human herpesvirus 6 DNA. *Blood* **94**, 1545–1549 (1999).

27.   S. Minocherhomji, *et al.*, Replication stress activates DNA repair synthesis in mitosis. *Nature* **528**
(2015).

28.   H. Miura, *et al.*, Inherited Chromosomally Integrated Human Herpesvirus 6 Is a Risk Factor for
Spontaneous Abortion. *J Infect Dis* **223** (2021).

29.   F. Gaccioli, *et al.*, Fetal inheritance of chromosomally integrated human herpesvirus 6 predisposes the
mother to pre-eclampsia. *Nat Microbiol* **5** (2020).

30.   A. Gravel, *et al.*, Inherited chromosomally integrated human herpesvirus 6 as a predisposing risk
factor for the development of angina pectoris. *Proc Natl Acad Sci U S A* **112** (2015).

31.   P. E. Pellett, Chromosomally integrated human herpesvirus 6: questions and answers. *Rev Med Virol*
22, 144–55 (2012).

32.   L. Flamand, "Chromosomal integration by human herpesviruses 6A and 6B" in *Advances in
Experimental Medicine and Biology*, (2018).

33.   A. Gravel, *et al.*, Mapping the Human Herpesvirus 6B transcriptome. *J Virol* (2021)
https://doi.org/10.1128/JVI.01335-20.

34.   U. Schiewe, F. Neipel, D. Schreiner, B. Fleckenstein, Structure and transcription of an immediate-early
region in the human herpesvirus 6 genome. *J Virol* **68**, 2978–2985 (1994).

35.   J. Jaworska, A. Gravel, K. Fink, N. Grandvaux, L. Flamand, Inhibition of Transcription of the Beta
Interferon Gene by the Human Herpesvirus 6 Immediate-Early 1 Protein. *J Virol* **81**, 5737–5748
(2007).

36.   J. Jaworska, A. Gravel, L. Flamand, Divergent susceptibilities of human herpesvirus 6 variants to type I
interferons. *Proc Natl Acad Sci U S A* **107**, 8369–8374 (2010).

37.   R. Bernardi, P. P. Pandolfi, Structure, dynamics and functions of promyelocytic leukaemia nuclear
bodies. *Nat Rev Mol Cell Biol* **8**, 1006–1016 (2007).

38.   P. L. Yeung, *et al.*, Promyelocytic leukemia nuclear bodies support a late step in DNA double-strand
break repair by homologous recombination. *J Cell Biochem* **113**, 1787–1799 (2012).

39.   K. M. Attwood, *et al.*, PML isoform expression and DNA break location relative to PML nuclear bodies
impacts the efficiency of homologous recombination. *Biochemistry and Cell Biology*, 1–42 (2019).

40.   G. Dellaire, *et al.*, Promyelocytic leukemia nuclear bodies behave as DNA damage sensors whose
response to DNA double-strand breaks is regulated by NBS1 and the kinases ATM, Chk2, and ATR.
Journal of Cell Biology **175**, 55–66 (2006).

41.   M. Vancurova, *et al.*, PML nuclear bodies are recruited to persistent DNA damage lesions in an
RNF168-53BP1 dependent manner and contribute to DNA repair. *DNA Repair (Amst)* **78** (2019).

42.   V. Collin, A. Gravel, B. B. Kaufer, L. Flamand, The promyelocytic leukemia protein facilitates human
herpesvirus 6B chromosomal integration, immediate-early 1 protein multiSUMOylation and its
localization at telomeres. *PLoS Pathog* **16** (2020).

43.   A. P. R. Cassel, R. B. Barcellos, C. M. D. da Silva, S. E. de Matos Almeida, M. L. R. Rossetti, Association
between human papillomavirus (HPV) DNA and micronuclei in normal cervical cytology. *Genet Mol*
Biol **37**, 360–363 (2014).

44.   E. M. Hatch, A. H. Fischer, T. J. Deerinck, M. W. Hetzer, Catastrophic Nuclear Envelope Collapse in
Cancer Cell Micronuclei. *Cell* **154** (2013).

45.   N. Hustedt, D. Durocher, The control of DNA repair by the cell cycle. *Nat Cell Biol* **19**, 1–9 (2017).

46.   B. van de Kooij, H. van Attikum, Genomic Reporter Constructs to Monitor Pathway-Specific Repair of
DNA Double-Strand Breaks. *Front Genet* **12** (2022).

47.   J. Pinder, J. Salsman, G. Dellaire, Nuclear domain "knock-in" screen for the evaluation and
identification of small molecule enhancers of CRISPR-based genome editing. *Nucleic Acids Res* **43**
(2015).

48.   J. M. Stark, A. J. Pierce, J. Oh, A. Pastink, M. Jasin, Genetic Steps of Mammalian Homologous Repair
with Distinct Mutagenic Consequences. *Mol Cell Biol* **24**, 9305–9316 (2004).

49.   R. A. C. M. Boonen, *et al.*, Functional analysis of genetic variants in the high-risk breast cancer
susceptibility gene PALB2. *Nat Commun* **10** (2019).

50.   L. Costantino, *et al.*, Break-induced replication repair of damaged forks induces genomic duplications
in human cells. *Science (1979)* **343**, 88–91 (2014).

51.   K. Jacquet, *et al.*, The TIP60 Complex Regulates Bivalent Chromatin Recognition by 53BP1 through
Direct H4K20me Binding and H2AK15 Acetylation. *Mol Cell* **62**, 409–421 (2016).

52.   R. Bhargava, *et al.*, C-NHEJ without indels is robust and requires synergistic function of distinct XLF
domains. *Nat Commun* **9** (2018).

53.   C. Warren, N. P. Pavletich, Structure of the human ATM kinase and mechanism of Nbs1 binding. *Elife*
11 (2022).

54.   C. B. Schiller, *et al.*, Structure of Mre11-Nbs1 complex yields insights into ataxia-telangiectasia- like
disease mutations and DNA damage signaling. *Nat Struct Mol Biol* **19** (2012).

55.   J. Tang, *et al.*, Acetylation limits 53BP1 association with damaged chromatin to promote homologous
recombination. *Nat Struct Mol Biol* **20**, 317–325 (2013).

56.   J. Sitz, *et al.*, Human papillomavirus E7 oncoprotein targets RNF168 to hijack the host DNA damage
response. *Proc Natl Acad Sci U S A* **116**, 19552–19562 (2019).

57.   E. Soutoglou, T. Misteli, Activation of the cellular DNA damage response in the absence of DNA
lesions. *Science (1979)* **320** (2008).

58.   L. Bian, Y. Meng, M. Zhang, D. Li, MRE11-RAD50-NBS1 complex alterations and DNA damage
response: Implications for cancer treatment. *Mol Cancer* **18** (2019).

59.   D. I. Lou, *et al.*, An Intrinsically Disordered Region of the DNA Repair Protein Nbs1 Is a Species-Specific
Barrier to Herpes Simplex Virus 1 in Primates. *Cell Host Microbe* **20** (2016).

60.   A. J. Cesare, R. R. Reddel, Alternative lengthening of telomeres: Models, mechanisms and
implications. *Nat Rev Genet* **11** (2010).

61.   R. L. Dilley, *et al.*, Break-induced telomere synthesis underlies alternative telomere maintenance.
648   *Nature* **539** (2016).

649   62.   A. Gravel, *et al.*, Cell Culture Systems To Study Human Herpesvirus 6A/B Chromosomal Integration. *J*
650   *Virol* **91**, e00437-17 (2017).

651   63.   Z. You, C. Chahwan, J. Bailis, T. Hunter, P. Russell, ATM Activation and Its Recruitment to Damaged
652   DNA Require Binding to the C Terminus of Nbs1. *Mol Cell Biol* **25** (2005).

653   64.   A. J. Barbera, *et al.*, The nucleosomal surface as a docking station for Kaposi's sarcoma herpesvirus
654   LANA. *Science (1979)* **311** (2006).

655   65.   D. C. Avgousti, *et al.*, A core viral protein binds host nucleosomes to sequester immune danger
656   signals. *Nature* **535** (2016).

657   66.   C. Lukas, *et al.*, Mdc1 couples DNA double-strand break recognition by Nbs1 with its H2AX-dependent
658   chromatin retention. *EMBO Journal* **23** (2004).

659   67.   G. A. Shah, C. C. O'Shea, Viral and Cellular Genomes Activate Distinct DNA Damage Responses. *Cell*
660   **162** (2015).

661   68.   V. Peddu, *et al.*, Inherited Chromosomally Integrated Human Herpesvirus 6 Demonstrates Tissue-
662   Specific RNA Expression In Vivo That Correlates with an Increased Antibody Immune Response . *J Virol*
663   **94** (2019).

664   69.   F. J. Hari, C. Spycher, S. Jungmichel, L. Pavic, M. Stucki, A divalent FHA/BRCT-binding mechanism
665   couples the MRE11-RAD50-NBS1 complex to damaged chromatin. *EMBO Rep* **11** (2010).

666   70.   J. R. Chapman, S. P. Jackson, Phospho-dependent interactions between NBS1 and MDC1 mediate
667   chromatin retention of the MRN complex at sites of DNA damage. *EMBO Rep* **9** (2008).

668   71.   K. Kim, T. W. Kirby, L. Perera, R. E. London, Phosphopeptide interactions of the Nbs1 N-terminal FHA-
669   BRCT1/2 domains. *Sci Rep* **11** (2021).

670   72.   C. Xu, *et al.*, Structure of a Second BRCT Domain Identified in the Nijmegen Breakage Syndrome
671   Protein Nbs1 and its Function in an MDC1-Dependent Localization of Nbs1 to DNA Damage Sites. *J*
672   *Mol Biol* **381** (2008).

673   73.   R. A. Wu, H. E. Upton, J. M. Vogan, K. Collins, Telomerase mechanism of telomere synthesis. *Annu Rev*
674   *Biochem* **86** (2017).

675   74.   S. Gilbert-Girard, *et al.*, Stabilization of telomere G-quadruplexes interferes with human herpesvirus
676   6A chromosomal integration. *J Virol*, JVI.00402-17 (2017).

677   75.   Z. H. Zhong, *et al.*, Disruption of telomere maintenance by depletion of the MRE11/RAD50/NBS1
678   complex in cells that use alternative lengthening of telomeres. *Journal of Biological Chemistry* **282**
679   (2007).

680   76.   W.-Q. Jiang, *et al.*, Suppression of Alternative Lengthening of Telomeres by Sp100-Mediated
681   Sequestration of the MRE11/RAD50/NBS1 Complex. *Mol Cell Biol* **25** (2005).

682   77.   G. Wu, X. Jiang, W. H. Lee, P. L. Chen, Assembly of functional ALT-associated promyelocytic leukemia
683   bodies requires nijmegen breakage syndrome 1. *Cancer Res* **63** (2003).

78.   D. J. Wight, *et al.*, Viral proteins U41 and U70 of human herpesvirus 6A are dispensable for telomere
integration. *Viruses* **10** (2018).

79.   N. Wallaschek, A. Gravel, L. Flamand, B. B. Kaufer, The putative U94 integrase is dispensable for
human herpesvirus 6 (HHV-6) chromosomal integration. *Journal of General Virology* **97** (2016).

80.   S. Sakamoto, *et al.*, Homologous recombination repair is regulated by domains at the N- and C-
terminus of NBS1 and is dissociated with ATM functions. *Oncogene* **26** (2007).

81.   R. Bhowmick, S. Minocherhomji, I. D. Hickson, RAD52 Facilitates Mitotic DNA Synthesis Following
Replication Stress. *Mol Cell* **64** (2016).

82.   Y. Huang, *et al.*, Human telomeres that carry an integrated copy of human herpesvirus 6 are often
short and unstable, facilitating release of the viral genome from the chromosome. *Nucleic Acids Res*
42 (2014).

83.   A. Gravel, J. Gosselin, L. Flamand, Human herpesvirus 6 immediate-early 1 protein is a sumoylated
nuclear phosphoprotein colocalizing with promyelocytic leukemia protein-associated nuclear bodies.
Journal of Biological Chemistry **277**, 19679–19687 (2002).

Figure Legends

Fig. 1. HHV-6B infection and IE1 expression lead to micronuclei formation *(A)* Left panel: Representative images
of Mock and HHV-6B-infected MOLT-3 cells fixed 24 h post-infection and counterstained with DAPI. Micronuclei
are indicated by white arrows. The quantification of the micronuclei (right panel) shows the mean (n = 2, >100
micronuclei/condition). *(B)* Left panel: representative images of U2OS cell lines and clones stably expressing
Dox-inducible HHV-6B IE1 (C10 and C102). IE1 expression was induced with 1 µg/mL Dox for 48 h prior to IE1
immunofluorescence. Micronuclei are indicated by white arrows. The parental cell line (Par.) was used as a
negative control. The micronuclei (right panel) quantification shows the mean ± standard deviation (SD; n = 3).
(C) Schematic of micronuclei formation *via* DNA double-strand breaks (DSBs) (i), lagging chromosomes (ii),
and anaphase bridges (ABs) (iii). *(D-E)* Quantification of micronuclei containing centromeres (D) and telomeres
(E). Cells were treated as in B and centromeres and telomeres were detected by immunofluorescence and
FISH, respectively. Data represent the mean ± SD (n = 3) (D) and the mean (n = 2, > 100 micronuclei/condition)
(E). *(F)* A representative metaphase spread from an IE1-expressing cell. Cells were treated with 1 µg/mL Dox
for 48 h, then metaphase spreads were prepared, fixed, and counterstained with DAPI. *(G)* Quantification of
chromosomal aberrations per metaphase. Data represent the mean ± SD (n = 31). **p < 0.01, ***p < 0.001,
****p < 0.0001. Scale bars = 5 µm.

Fig. 2. H2AX phosphorylation (γ-H2AX) is inhibited in HHV-6B-infected and IE1-expressing cells *(A)*
Representative γ-H2AX immunostaining in HHV-6B-infected MOLT-3 cells irradiated with 4 Gy and
immunostained for IE1 and γ-H2AX 1 h later. Mock-infected cells were used as a negative control. *(B)*
Quantification of irradiated MOLT-3-infected and Mock cells with > 10 γ-H2AX foci. Data represent the mean ±
SD (n = 3). Statistical significance was assessed using unpaired *t*-test. *(C)* Representative γ-H2AX
immunostaining in irradiated U2OS parental (Par.) and IE1-expressing cells (Clone C10). IE1 expression was

induced as in Fig. 1B. Cells were irradiated with 1 Gy and immunostained for IE1 and γ-H2AX 1 h later. *(D)*

Quantification of irradiated U2OS Par. and IE1-expressing cells with > 10 γ-H2AX foci. Data are presented as

the mean ± SD (*n* = 3). *(E)* Quantification of cells with > 10 γ-H2AX foci in irradiated (1 Gy) U2OS *PML*^(+/+) and ^-

^(/-) cells that transiently express untagged IE1. An empty vector (EV) was used as a negative control. Data

represent the mean ± SD (*n* = 3). ****p < 0.0001. Scale bars = 5 μm.

Fig. 3. HHV-6B IE1 inhibits HDR-mediated repair *(A-D)* DNA repair reporter assays for (A) homologous

recombination (DR-GFP), (B) single-strand annealing (SA-GFP), (C) break-induced replication (BIR-GFP), and

(D) non-homologous end-joining (NHEJ-GFP (EJ7)). For each assay, a schematic is presented in the top panel

and the flow cytometry-based quantification of GFP^+ cells is presented in the bottom panel. In each replicate,

GFP^+ cells were normalized to the GFP^+ cells in the positive control (I-SceI^+ or Cas9^+, set to 1.0). Data represent

the mean ± SD (*n* = 3). ****p < 0.0001.

Fig. 4. HHV-6B IE1 colocalizes with NBS1 *(A)* Signaling events triggered by DNA DSBs. *(B)* Whole cell extracts

(WCEs) from U2OS cells (Par.) and IE1-expressing U2OS stable cell lines treated with or without 1 μg/mL Dox

were immunoblotted for RAD50, NBS1, and MRE11. GAPDH was used as a loading control. *(C, E)*

Representative images of the colocalization between IE1 and NBS1 (C) and MRE11 (E). IE1-expressing cells

were treated as described in Fig. 1B and immunostained for IE1, NBS1, or MRE11. As a positive control,

irradiated U2OS cells (+IR) were fixed 15 min post-irradiation (1 Gy) and immunostained as indicated (*SI*

Appendix, Fig. S4*A*; scale bars, 5 μm). The parental cell line (Par.) was used as a negative control. *(D)*

Percentages of IE1 foci that colocalized with NBS1 (C) and MRE11 (E). Data represent the mean ± SD of three

independent experiments. *(F)* Percentage of IE1 foci that colocalized with MRE11 in stable U2OS control cells

(shCTRL) or those depleted of NBS1 (shNBS1). Data represent the mean ± SD (*n* = 2, at least 40

nuclei/condition). Statistical significance was assessed using unpaired *t*-test. ****p < 0.0001. Scale bars = 5 μm.

Fig. 5. HHV-6B IE1 interacts with NBS1, inhibits ATM activation, and is recruited to DSBs *(A)* The integrated

LacO array-based assay used to study protein colocalization at a specific locus without DSBs (*i*) or DNA repair

protein recruitment at localized DSBs (*ii*). *LacO* array repeats, mCherry-LacRnls-fusion proteins, and preys are

shown in grey, red/orange, and green, respectively. *(B)* Quantification of the indicated DSB-signaling proteins

at localized DSBs induced by ER-mCherry-LacRnls-FOKI-DD (FOKI) or at either a mCherry-LacRnls (--,

negative control) or mCherry-LacRnls HHV-6B IE1 protein foci in U2OS 2-6-5 cells. Transfected cells were

treated with 4-OHT and Shield-1 for 6 h then immunostained for ATM, pATM (S1981), γ-H2AX, RAD50, NBS1,

and MRE11 (Representative images, *SI Appendix,* Fig. S5*A-E*). For each condition, statistical significance was

analyzed against the control protein (mCherry-LacRnls). Data represent the mean ± SD (*n* = 3). *(C)*

Representative images of NBS1 recruitment at DSBs (top panel) and its colocalization with IE1 in the absence

of DSBs (bottom panel). *(D-E)* U2OS 2-6-5 cells were treated as described in (B), immunostained for γ-H2AX

(D) or pATM (E) (*SI Appendix,* Fig. S5*F-G*) and quantified as indicated. In both experiments, cells were

transfected with vectors expressing either untagged or 3×FLAG-IE1. *(F)* Representative images of pATM

inhibition at the mCherry-LacRnls-NBS1 locus. Cells were treated as described in (B). Representative images

of the negative controls are presented in *SI Appendix,* Fig. S5*G-H*. *(G)* Representative images of NBS1-

dependent IE1 recruitment to DSBs. Cells were treated with a siCTRL or siNBS1 prior to their transfection with

an untagged IE1 or an empty vector. ER-mCherry-LacR-FOKI-DD was induced as described in (B). Cells were

processed for IE1 immunofluorescence. *(H)* Quantification of the mCherry-LacR FOKI foci colocalizing with IE1.

Data represent the mean ± SD (*n* = 3). **$p < 0.05$, **$p < 0.01$, *** $p<0.001$, ****$p < 0.0001$. Scale bars = 5 μm.

Fig. 6. IE1 interacts with NBS1 and inhibits ATM through two distinct domains *(A)* Schematic of HHV-6B IE1

and the protein fragments used in this study. NBS1-BD, NBS1-binding domain; NBS1i, NBS1 inhibitory domain,

STAT2-BD: STAT2 binding-domain (aa 270–540). *(B-C)* U2OS 2-6-5 cells transfected with plasmids expressing

the indicated mCherry-LacR fusion proteins were immunostained for NBS1 (see also *SI Appendix,* Fig. S6*A-B*).

The mCherry-LacR backbone was used as a negative control (--). *(D-E)* Quantification of cells with > 10 γ-H2AX

foci. UT, untreated. Transiently transfected cells were irradiated and immunostained for γ-H2AX 1 h later (see

also *SI Appendix,* Fig. S6*G*). Untreated cells and the mCherry-LacR backbone were used as negative controls

(--). *(F)* Schematic of NBS1 and the protein fragments used in this study. FHA, forkhead-associated domain;

BRCT, BRCA1 C-terminal domain; MRE11-BM, MRE11-binding motif; ATM-BM, ATM-binding motif; IDD,

intrinsically disordered domain. *(G-H)* U2OS 2-6-5 cells were transfected with the indicated mCherry-LacR

plasmids and immunostained for IE1 (*SI Appendix,* Fig. S6*H-M*). *(I)* U2OS cells treated with siCTRL or siNBS1

were transfected with the indicated 3×FLAG and mCherry-LacR constructs. After 24 h, WCEs were prepared

and 3×-FLAG-IE1 interactors were immunoprecipitated using anti-Flag (M2) agarose beads and immunoblotted

for FLAG, mCherry, and NBS1. β-tubulin (Tub.) was used as a loading control. Data for *(B), (D),* and *(G)*

represent the mean ± SD (*n* = 3). ** $p<0.01$, ****$p < 0.0001$. Scale bars = 5 μm.

Fig. 7. NBS1 depletion impairs viral integration in cells maintaining their telomeres by homology-directed repair

(A) HHV-6B infection in permissive and semi-permissive cells. In cells semi-permissive for HHV-6B, replication

is inefficient, and the viral genome integrates at the telomeres. *(B)* MOLT-3 cells with and without NBS1 (*SI*

Appendix, Fig. S7*A*) were infected with HHV-6B at a MOI of 1 and harvested at the indicated time points.

Following cell lysis, DNA was extracted and HHV-6B was quantified by qPCR using primers for HHV-6B *U67-*

68 and human *RPP30*. Data represent the mean ± SD (*n* = 3). *(C)* MOLT-3 cells were transduced with the

indicated shRNA and passaged 5 times prior to CellTiter-Glo® analyses. Cell viability was determined using

standard curves for each cell line and normalized to the shCTRL condition for each experiment.

Table 1. Importance of NBS1 for HHV-6B chromosomal integration in ALT[+/-] cells

Cell line	ALT status	shRNA	% cells with integrated HHV-6B[a] (n)[b]	P value[c]
HeLa	Negative	CTRL	0.96 (36,320)	<2.2e[-16]
		NBS1	6.11 (33,280)	
GM847	Positive	CTRL	0.65 (21,820)	<2.2e[-16]
		NBS1	0.01 (18,320)	
U2OS	Positive	CTRL	1.60 (20,000)	<2.2e[-16]
		NBS1	0.69 (21,520)	
U2OS *PML*[-/-]	Positive	CTRL	0.71 (28,220)	ns[d]
		NBS1	0.78 (30,460)	

[a] mean of three independent cultures

[b] total number of cells analyzed

[c] Pearson's Chi-squared test with Yates' continuity correction

[d] ns, not significant

Fig 1

A IE1 γ-H2AX Merge — Mock, HHV-6B — MOLT-3, 1h post 4Gy

C IE1 γ-H2AX Merge — Par., HHV-6B IE1 C10 — U2OS, + Dox, 1h post 1Gy

B Cells with >10 γ-H2AX foci — Mock, HHV-6B — ****

D Cells with >10 γ-H2AX foci — Par., C10, C102 — **** ****

E Cells with >10 γ-H2AX foci — EV, IE1 — PML$^{+/+}$, PML$^{-/-}$ — **** ****

Fig. 4

Fig. 5

Fig 6