**SARS-CoV-2 fears green: the chlorophyll catabolite Pheophorbide a is a potent antiviral.**

**Authors:** Guillermo H. Jimenez-Aleman[1,a], Victoria Castro[2,a], Addis Longdaitsbehere[1], Marta

Gutierrez-Rodríguez[3], Urtzi Garaigorta[2], Roberto Solano[1,*] and Pablo Gastaminza[2*].

[1] Department of Plant Molecular Genetics, National Centre for Biotechnology

[2] Department of Cell & Molecular Biology, National Centre for Biotechnology

[3] Department of Biomimetics for Drug Discovery, Medicinal Chemistry Institute (IQM-CSIC)

[a] These authors contributed equally to the work.

*Corresponding authors: Roberto Solano (rsolano@cnb.csic.es) and Pablo Gastaminza

(pgastaminza@cnb.csic.es)

**Abstract**

The SARS-CoV-2 pandemic is having devastating consequences worldwide. Although vaccination advances at good pace, effectiveness against emerging variants of the virus is unpredictable. The virus has displayed a remarkable resistance to treatments and no drugs have been proved fully effective against Covid-19. Thus, despite the international efforts, there is still an urgent need for new potent and safe antivirals against SARS-CoV-2. Here we exploited the enormous potential of plant metabolism, in particular the bryophyte *Marchantia polymorpha,* and following a bioactivity-guided fractionation and mass-spectrometry approach, identified a potent SARS-CoV-2 antiviral. We found that the chlorophyll derivative Pheophorbide a (PheoA), a natural porphyrin similar to animal Protoporphyrin IX, has an extraordinary antiviral activity against SARS-CoV-2 preventing infection of cultured monkey and human cells, without noticeable citotoxicity. We also show that PheoA prevents coronavirus entry into the cells by directly targeting the viral particle. Besides SARS-CoV-2, PheoA also displayed a broad-spectrum antiviral activity against (+)strand RNA viral pathogens such as HCV, West Nile, and other coronaviruses, but not against (-)strand RNA viruses, such as VSV. Our results indicate that PheoA displays a remarkable potency and a satisfactory therapeutic index, and suggest that it may be considered as a potential candidate for antiviral therapy against SARS-CoV-2. Moreover, PheoA adds to remdesivir's efficiency and is currently employed in photoactivable cancer therapies in humans.

## Introduction

The pandemic caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is having devastating consequences, with more than 196M infected people and over 4M deaths worldwide (July 2021; https://covid19.who.int/). Besides the humanitarian cost, this pandemic carries a tremendous negative economic impact, a huge challenge for any government to overcome. The coronavirus disease 2019 (Covid-19), the respiratory illness caused by SARS-CoV-2 (Genus betacoronavirus; Subgenus sarbecovirus), has displayed a remarkable resistance to treatments and no drugs have been proved fully effective against the virus. Moreover, the Covid-19 pandemic has made evident the need for a global strategy to fight similar situations that may appear in the future.

Current efforts to eradicate Covid-19 are focused on the development of vaccines and the search for antiviral lead compounds, mainly repurposing of existing drugs. Although the vaccination campaign seems to advance at good pace, its effectiveness against some of the present and future strains of SARS-CoV-2 is hard to predict due to the existence of different strains that could drastically reduce the vaccine efficiency[1]. In addition, the best anti-Covid-19 drugs approved so far (e.g. Remdesivir, Favipiravir or its derivative Avifavir, etc), have shown only a mild effect against the virus, slightly reducing hospitalization time of patients[2]. Other treatments, such as Chloroquine and Hydroxychloroquine appear to help at least a subgroup of patients, but, possible negative side effects of these drugs remain under investigation[3]. Although, some compounds (e.g., Aplidin, Mefloquine, Nelfinavir, Protoporphyrin IX and Verteporfin) have shown potential on *in vitro* assays, and some of them also in animal models[4–6], there is an urgent need for new potent and safe antivirals against SARS-CoV-2. Noteworthy, new pathogens, including viruses, are expected to emerge in coming decades, which puts an enormous pressure on society in order to be ready to fight back future pandemics with the proper chemical, biological, and engineering tools, including effective new antivirals.

For centuries, medical needs of society have been widely covered by plants, which have an extremely rich metabolism that provides them with a wide repertoire of chemical weapons to cope with environmental biotic stresses, including viruses[7,8]. Originally recognized by traditional medicine, plants are the main source of compounds used today in pharmacology, from Aspirin (acetyl salicilate; from Salix sp.) to current anticancer drugs (e.g. Vinblastine and Vincristine from Vinca sp., or Taxol and Paclitaxel from Taxus baccata), simply to cite a few successful examples[9,10]. Therefore, the identification of new plant sources of enzymatic variants and metabolites is essential to the discovery of new drugs and their optimization by metabolic engineering. Aromatic and exotic vascular plants are commonly studied in order to identify pharmacologically interesting compounds. In contrast, the metabolic richness of bryophytes (non-vascular plants including mosses, liverworts and hornworts) has been little explored. Bryophytes are rarely attacked by pathogens (fungi, bacteria, viruses) or herbivores (insects, snails, mammals) in their natural habitats, which indicates that they are well protected by a potent arsenal of secondary defense metabolites. However, studies on their chemical constituents have been neglected until recently[11]. Indeed, only around 5% of bryophyte species have been metabolically explored, and results have shown an enormously rich diversity of secondary metabolites, particularly in liverworts[12]. Strikingly, more than 1600 terpenoids have been reported in liverworts, whereas only about 100 terpenoids have been identified in the medicinal plant *Cannabis sativa*[11,13–16]. More importantly, several liverwort species of the order Marchantiales, including *Marchantia polymorpha*, produce terpenoids and bisbibenzyls with enormous potential for pharmaceutical applications since they show remarkable antimicrobial, antioxidant, cytotoxic, anticancer and antiviral (anti-HIV) activities[11,13,15–18]. Therefore, we made use of our vast experience in vorology and Marchantia's hormonal signaling and secondary metabolism in order to explore this plant's potential as a source of antiviral metabolites, particularly, against the SARS-CoV-2 virus.

In this study, we employed a set of Marchantia wild type plants, and signalling and metabolic

mutants to systematically study the pharmacological potential of this liverwort. We found that

total extracts from all the plants displayed a remarkable antiviral activity against SARS-CoV-2.

Using a bioactivity-guided chromatographic approach, in addition to mass-spectrometry (MS),

we identified the antiviral metabolite as Pheophorbide a (PheoA), a porphyrin chlorophyll

derivative very similar to animal Protoporphyrin IX, also described as an strong antiviral. In

contrast to Protoporphyrin IX, however, which produces prophyria in humans, PheoA is non

toxic. We also found that PheoA has a broad-range antiviral activity against positive strand RNA

(+RNA) viruses and acts as a virucidal, by directly acting on the viral particle. PheoA is additive

to remdesivir, which, together with its low toxicity, suggest its potential as candidate for antiviral

therapy against SARS-CoV-2.

**Results**

***Crude extracts of M. polymorpha show anti-SARS-CoV-2 activity***

In order to explore for the presence of anti-SARS-CoV-2 metabolites in *M. polymorpha*, we

prepared crude extracts of two different *M. polymorpha* subspecies, (subsp. *ruderalis* from Japan

and subsp. *polymorpha*, from Spain). The anti-SARS-CoV-2 activity of the resulting crude

extracts was tested in Vero-E6 cell monolayers infected with the SARS-CoV-2 NL2020 strain

(Figure 1). Infection was carried out at a multiplicity of infection (MOI) of 0.001 for 72 h. In the

absence of antiviral activity, SARS-CoV-2 infection triggers cell death of Vero E6 cells and

results in loss of cell biomass, which is readily visualized as a strong reduction in crystal violet

staining in the well (DMSO, Figure 1). Treatment of the cells during infection with serial dilutions

of remdesivir (used as positive control), the only clinically approved antiviral for treatment of

Covid-19 patients, protected the cell monolayers down to its reported $EC_{50}$ of 1.5 μM (Figure

S1). Similarly, treatment of cell cultures with Marchantia crude extracts resulted in cell protection

against virus-induced cytopathic effect without any signs of citotoxicity in a broad dilution range

(Figure 1A, Ex1 and Ex2), suggesting the presence of one or more Marchantia metabolites with

strong antiviral activity.

Given that antiviral activity had been determined by an indirect measurement (Figure 1B), we set

out to define if Marchantia extracts were indeed capable of interfering with viral spread in cell

culture. Thus, Vero E6 cells were inoculated at MOI of 0.001 in the presence of control-solvent,

remdesivir (6.25 or 25 $\mu$M) and a 1:800 (v/v) dilution of the Marchantia extracts. Viral RNA

load, which in this experimental setup represents the degree of virus propagation, was determined

72 h post infection by RT-qPCR (Figure 2). In the control, viral RNA accumulated six orders of

magnitude above the assay background levels, whereas the viral RNA was undetectable in

samples treated with the antiviral remdesivir. Importantly, in samples treated with Marchantia

extracts, the viral RNA levels were comparable to those observed upon remdesivir treatment

(Figure 2). This outcome confirms the protective activity of the extracts observed in Figure 1 and

suggest the presence of at least one antiviral compound.

***Extract bioactivity does not depend on plant's secondary metabolism and is common to several***

***plant species***

Next, we explored whether the putative antiviral metabolite could belong to plant's secondary

metabolism. Jasmonates (JAs) are a family of oxylipin-derived phytohormones regulating many

aspects of plant development and growth; as well as mediating defense responses through

transcriptional activation of the secondary metabolism, which includes several classes of

compounds such as alkaloids, terpenoids and flavonoids [19–21]. In Marchantia, secondary

metabolites accumulate in specific organelles named oil bodies (OB), which are confined to

scattered idioblastic OB cells distributed throughout the thallus [22]. Therefore, we tested extracts

from *M. polymorpha* WT, Mp*coi1-2* [impaired in dn-OPDA perception, the active JA in

Marchantia [21], thus, in defense metabolite induction], and Mp*c1hdz* plants (impaired in OB

formation; MpC1HDZ is a transcription factor required for OB cells differentiation). Mp*c1hdz*

mutants render plants defective in secondary metabolites, thus, susceptible to herbivory and

microbes [23]. To our surprise, all Marchantia extracts, WT or mutant, showed similar antiviral

activity (Figure 3), indicating that the active antiviral should not belong to the plants's secondary

metabolism. Indeed, data in Figure 3 suggests that the activity of extracts is due to the presence

of a metabolite (or metabolites) that is constitutively synthesized and/or derived from the plant's

primary metabolism. Remarkably, regulation of primary metabolism is achieved in all plants by

very similar conserved metabolic pathways [24,25]. Therefore, we tested crude extracts of several

plant species [sweet amber (*Hypericum androsaemum)*, fern (*Blechnum spicant*), nettle (*Urtica*

*dioica*), moss (*Physcomitrium patens*), tobacco (*Nicotiana benthamiana*) and thale cress

(*Arabidopsis thaliana*)] for their capability of providing protection to Vero E6 cells against the

SARS-CoV-2 virus. As shown in Figure S2, certain degree of protection was observed for most

of the tested plant species; the clearest protecting activity was observed for *Marchantia* crude

extracts.

***Identification of the antiviral metabolite***

In order to identify the bioactive metabolite(s), we followed a *bioactivity-guided*

*chromatographic fractionation* of the Marchantia's WT crude extracts, which showed strong

antiviral effect in previous assays. Chromatographic fractions were obtained via flash column

employing a solvent polarity gradient, starting at *n*-hexane (100%) up to AcOEt:MeOH (4:1, v/v).

A total of 56 fractions were obtained; fractions of a similar composition, based on their thin layer

chromatography (TLCs) profiles, were combined and evaluated as 12 new pooled fractions (1-

12). Fractions 10, 11 and 12 showed antiviral activity in the monolayer protection assay (Figure

4A-B). To directly confirm their antiviral activity, the viral antigen load reduction after

inoculation of cell cultures with SARS-CoV-2 (MOI = 0.01) was measured. In this experimental

setup, viral antigen accumulates as a consequence of virus propagation and can be quantitated

using automated immunofluorescence microscopy. Figure 4C shows a dose-dependent reduction

of viral antigen accumulation and the absence of cytotoxicity, as confirmed by normal cell

numbers, estimated by DAPI staining and image analysis, and cell viability studies performed in

parallel, uninfected cultures by MTT assays (Figure 4C).

Interestingly, TLCs of the three active fractions presented red fluorescent spots (under long wave

ultraviolet light, 365 nm; Supplementary Figure S3A), which are characteristic of plant

chlorophylls, but with a smaller retention factor [Rf = 0.36, AcOEt:MeOH (9:1, v/v)] than that

of chlorophyll (Rf = 0.92). At this point, we suspected that the active antiviral metabolite(s) could

be related to plant chlorophylls, especially because a weak antiviral activity was observed at low

dilutions of fractions 2 and 3 (Figure 4A), both containing chlorophyll. Indeed, these fractions,

when re-chromatographed, yielded spots on the TLC with an Rf consistent with that observed in

the active fractions 10 to 12, indicating that the active metabolite is a chlorophyll derivative

(Supplementary Figure S3B). It is worth mentioning that heat notably accelerated the chlorophyll

decomposition into the investigated metabolite.

Next, we employed preparative TLC in order to better isolate and characterize the red-light

emitting metabolite. Photosynthetic metabolites were extracted from *M. polymorpha* WT and

Mp*c1hdz* plants, and the extracts subjected to preparative TLC. Three spots showed fluorescence

in the vicinity of the expected Rf; these spots where isolated, analysed by HPLC-UV-MS and the

bioactivity assayed as fractions C, D and E (Figure 5A). Fraction D showed the strongest antiviral

activity (Figure 5B), which corresponded with an enrichment of compound **1** in the UV

chromatograms (Figure 5C). Careful analysis of the MS spectra (positive mode) revealed a

molecular formula of $C_{35}H_{36}N_4O_5$ for compound **1**, as deduced by HR-ESI$^+$-MS from its

monoprotonated molecular ion, [M+H]$^+$, with a *m/z* of 593.2759. The identified molecular

formula (containing four nitrogen atoms), the exact mass and the characteristic red fluorescence

of the compound helped to identify **1** as the chlorophyll catabolite *Pheophorbide a* (PheoA,

Figure 5C). The identity of PheoA was further confirmed by comparison with both a

commercially available original standard (Santa Cruz Biotechnology) and a semisynthetic

sample. PheoA was obtained semi-synthetically from Marchantia in good overall yield via a

solvent-free, thus, environmentally friendly method (Supp. Figure S4 and M&M section).

**Antiviral activity of PheoA.**

To confirm the antiviral potential of PheoA, a commercially available PheoA stock solution was

serially diluted and mixed with a virus stock to inoculate Vero E6 and Huh7-ACE2 cells (human

hepatoma cells expressing ACE2). Cells were fixed 72 h post-inoculation and stained with crystal

violet to visualize the integrity of the cell monolayer. Figure S5 shows consistent protective

capacity of PheoA at concentrations above 40 ng/ml (67 nM) in both cell lines.

PheoA antiviral activity was further confirmed by immunofluorescence microscopy, to estimate

virus propagation, and an MTT assay to evaluate compound cytotoxicity. PheoA dose-response

curves demonstrate PheoA's antiviral activity against SARS-CoV-2 in Vero E6 and human lung

epithelial cells (A549-ACE2 and Calu3); in all three models, no cytotoxicity was observed

(Figure 6). This dataset was used to determine effective concentrations ($EC_{50}$ and $EC_{90}$) and

cytotoxicity indexes. The estimated $EC_{50}$ and $EC_{90}$ values were around 14 ng/mL (25 nM) and

156 ng/mL (86 nM), with a wide therapeutic window in all tested cell lines (Table 1).

**Table 1: Potency and cytotoxicity indexes of commercially available PheoA**

| Cell line | $EC_{50}$ (nM) | $EC_{90}$ (nM) | $CC_{50}$ (nM) |
|---|---|---|---|
| Vero-E6 (green monkey) | 25 | 86 | >8420 |
| A549-ACE2 (human lung) | 52 | 263 | 2020 |
| Calu3 (human lung) | 34 | 232 | >8420 |

To fully evaluate the extent to which PheoA interferes with SARS-CoV-2 infection (i.e. on viral

replication) – given the narrow dynamic range of immunofluorescence –, we employed RT-qPCR

to determine the extracellular viral RNA load in human cells inoculated with SARS-CoV-2 (MOI

= 0.001). In this experimental setup, remdesivir (5 $\mu$M) reduced extracellular viral RNA by three

orders of magnitude. Interestingly, PheoA significantly reduced viral spread even at the lowest

tested concentration (150 ng/mL; 0.25 $\mu$M), as shown by the three times log reduction in viral

RNA accumulation after a 48 h incubation period (Figure 7). These results indicate that PheoA

displays a remarkable potency and a satisfactory therapeutic index, and suggest that it may be

considered as a potential candidate for antiviral therapy against SARS-CoV-2. Furthermore,

results also suggest that PheoA is a major determinant of the antiviral activity observed in crude

*Marchantia* extracts (Figure 1). This notion is underscored by the fact that the semisynthetic

PheoA (88-94% by HPLC-UV/Vis) showed comparable potency to crude extracts in the different

cell lines (Figure S6). Nevertheless, other related chlorophyll metabolites may also contribute

with antiviral activity. In fact, pyropheophorbide a (pPheoA), which was also found in antiviral

fractions was tested to verify its antiviral potential in Vero-E6 cells. The pPheoA showed antiviral

activity in the absence of measurable cytotoxicity with an $EC_{50}$ of 185 nM (Figure S7), suggesting

lower potency than PheoA and further underscoring a major role for PheoA in the antiviral

activity of Marchantia extracts.

**Antiviral spectrum of Pheophorbide A**

PheoA has previously been proven as an antiviral against the hepatitis C virus (HCV)[26] and

virucidal against herpes simplex virus (HSV)[27]. Thus, we determined the antiviral spectrum of

PheoA on different enveloped +RNA viruses. First, we confirmed antiviral activity against HCV,

using a surrogate model of infection by propagation-deficient, *bona fide* reporter virus bearing a

luciferase reporter gene generated by trans-complementation (HCVtcp). Dose-response curves of

the luciferase activity in HCVtcp-infected Huh7 cells indicated an $EC_{50}$ of 177 ng/mL (300 nM)

for PheoA against HCV (Figure 8A), very similar to the previously reported $EC_{50}$[26].

Next, we asked whether PheoA antiviral activity against SARS-CoV-2 could also be observed

against other human coronaviruses such as hCoV-229E (Genus alphacoronavirus; Subgenus

Duvinecovirus), which has been associated with mild respiratory infections[28]. Huh7 cells were

inoculated (MOI = 0.01) and total GFP expression in the target Huh7 cells was evaluated by

automated microscopy 48 h post-inoculation. Similar to the results with the HCV infection model,

PheoA reduced viral spread ($EC_{50}$ of 76 ng/mL; 128nM) while no cytotoxicity was observed

(Figure 8B). Strikingly, comparable results were obtained in an experimental model of infection

by the West Nile Virus, a mosquito-borne zonotic pathogen that may cause encephalitis in

infected humans. A recombinant, di-cistronic infectious molecular clone expressing GFP in the

second cistron[29] was used to inoculate Huh7 cells (MOI = 0.01). Cells were imaged 48 h post

infection and the degree of virus propagation was determined via automated microscopy; dose

response curves (Figure 8) indicated antiviral activity for PheoA with an $EC_{50}$ of 38 ng/mL (68

nM). Altogether, these results suggest that PheoA displays a broad-spectrum antiviral activity

against +RNA viral pathogens. In view of these results, we decided to test PheoA antiviral activity

against the vesicular stomatitis virus (VSV), a negative-strand RNA (-RNA) virus. Dose-response

experiments, similar to those describe above, were performed employing A549-ACE2 cells.

Interestingly, the VSV-GFP[30] was not susceptible to PheoA doses largely exceeding those for

which antiviral activity was observed against SARS-CoV-2 (Figure 8D). Collectively, these

results suggest that PheoA is a broad-spectrum antiviral and that +RNA viruses are particularly

susceptible.

**Pheophorbide a can be employed in combination with remdesivir**.

Once the broad antiviral activity of PheoA has been demonstrated, we studied, in several infection

systems, whether the addition of PheoA to remdesivir treatment could result in a synergistic effect

on viral infection. Thus, combination treatments were performed with increasing doses of PheoA

and remdesivir. Drugs were mixed in different proportions, combined with infectious SARS-

CoV-2 (MOI = 0.01) and the mixtures were used to inoculate Vero E6 cells. Twenty-four hours

later, cells were fixed and processed to determine the infection efficiency as described in Figure

6. Individual treatment with either compound resulted in the expected dose-dependent inhibition

of virus infection, achieving the $EC_{50}$ at the expected doses (2000 nM for remdesivir and 40 nM

for PheoA). Increasing concentrations of PheoA improved remdesivir efficacy and viceversa.

However, full analysis of the combinations resulted in a synergy index close to three, indicating

that the drug combination is mostly additive[31], with an area of synergy at concentrations close to

the $EC_{50}$s (Figure 9). These results suggest that combinations of PheoA with other antivirals may

result beneficial, as it was observed by its additive effect in combination with remdesivir in cell

culture infection models.

**Characterization of Pheophorbide a mode of action on SARS-CoV-2 infection.**

Next, antiviral efficacy of PheoA was compared when PheoA was (i) present at all times, (ii)

added only during virus inoculation, or (iii) added only after virions had effectively penetrated

the cells in single-cycle infection experiments (MOI = 5). Imatinib, for which antiviral activity at

the level of virus entry has previously been demonstrated[32], was employed as the reference

compound. Infection efficiency revealed the expected antiviral activity for imatinib and PheoA

when maintained at all times in the experiment (Figure 10). Imatinib showed comparable efficacy

when added during the virus entry phase and greatly lost efficacy when added after virion

internalization, as expected for an entry inhibitor. Similar results were obtained with PheoA,

inhibition was nearly identical when maintained at all times or added during viral entry, but only

ca. 6% of the maximum efficacy (statistically significantly different though) was observed when

added after virion internalization. These results suggest that PheoA is mainly acting at early

stages of the infection, potentially at the level of viral entry.

In view of these results, we directly tested this hypothesis by determining the antiviral activity of

PheoA in a surrogate model of infection recapitulating only aspects related with viral entry such

as receptor recognition, virion internalization or membrane fusion. This system is based on

reporter retroviral vectors pseudotyped with SARS-CoV-2 Spike protein (Spp) or VSV

glycoprotein as a control (VSVpp). Infection efficiency in the presence of antiviral molecules is

determined as the relative expression values of the reporter gene, in this case a Firefly luciferase[32].

Relative infection efficiency was measured in the presence of the entry inhibitor imatinib (15

$\mu$M) and antiviral doses of PheoA. As expected, imatinib selectively inhibited Spp and not

VSVpp entry (Figure 11). PheoA barely interfered with either retroviral pseudotype infection

efficiency, with a maximum reduction of 40% at the maximum dose (400 ng/ml; 678 nM) (Figure

11). These results suggest that, while time-of-addition experiments suggest that PheoA interferes

predominantly with early aspects of the infection, surrogate models of viral entry indicate that it

does not interfere substantially with molecular events leading to viral entry *per se*.

PheoA irreversibly interferes with the virion infectivity in HSV and influenza infection models[27],

therefore, we explored whether PheoA could be virucidal for SARS-CoV-2 virions, a property

that would be compatible with the observations made in the time-of-addition experiments. Thus,

a known number of infectious particles ($10^5$ TCID$_{50}$) were mixed with increasing doses of PheoA

[8 ng/mL(0.014 $\mu$M) to 5000 ng/mL (8.45 $\mu$M)] and incubated for 30 minutes before residual

infectivity titer determination by endpoint dilution and TCID$_{50}$ determination. The dose of PheoA

was kept below its effective concentrations during the titration experiment. Figure 12 shows how

pre-incubation of the infectious virions with PheoA (40 ng/mL or more) resulted in irreversible,

dose-dependent inactivation of the virus infectivity. These results suggest that PheoA is virucidal

for SARS-CoV-2 virions and that virion infectivity inactivation contributes to its overall antiviral

effect.

**Discussion**

Due to their metabolic richness plants have been traditionally used as source of medicines. The

potential of plant metabolites in pharmacology is still far from being saturated, particularly in

certain plant clades. Indeed, bryophytes (non-vascular plants) are particularly rich in specialized

metabolites that are rarely found in other plant lineages [33]. Here, we explored this richness in

order to find antiviral compounds against the SARS-CoV-2 virus by employing an activity-

guided chromatographic method; and identified PheoA as a potent antiviral, very efficient not

only against SARS-CoV-2 but also against several other enveloped viruses.

The first evidence for antiviral activity of PheoA derives from observations made on HSV

infection models[34]. In those initial reports, some degree of selectivity towards other viruses was

reported, since adenovirus (Type VI), Japanese Encephalitis virus (JEV) or poliovirus were not

affected by treatment with PheoA-enriched algal extracts[35]. Subsequent studies suggested that

PheoA and pPheoA display broad-spectrum antiviral activity against enveloped viruses,

including influenza A[27] and HIV[36], but not against non-enveloped viruses[27]. Ohta et al. reported

that PheoA-containing preparations may display virucidal activity against HSV[35], a concept that

was further supported by Bouslanu et al.[27]. Our observations support that PheoA inactivates

SARS-CoV-2 (enveloped +RNA virus) virion infectivity through a virucidal mode of action.

First, time-of-addition experiments indicate that early aspects of the infection are targeted by

PheoA. Second, the study of viral entry using retroviral pseudotypes did not reveal any

measurable antiviral activity against SARS-CoV-2 entry, suggesting that receptor recognition by

the Spike protein, particle internalization and Spike-mediated fusion are not affected by PheoA.

Similar models have been used to identify key SARS-CoV-2 entry factors as well as to study

antibody neutralizing activity. Thus, irreversible inactivation of viral infectivity (virucidal

activity) was tested as a possible mechanism reconciling these apparently contradicting

observations. Pre-incubation of infectious SARS-CoV-2 virions with PheoA rendered the virions

non-infectious even when PheoA-virus dilutions were performed below active PheoA

concentrations. The virtual lack of activity of PheoA at post-entry levels may be explained by the

fact that PheoA can only act on the viral particle, or, that PheoA requires longer incubation

periods to penetrate the cell and interfere with downstream steps of the virus lifecycle. Given that

overall effectiveness of PheoA as virucidal is substantially stronger than during multiple cycle

infection experiments, it is likely that virucidal activity is the main mechanism for interference

with SARS-CoV-2 infection.

PheoA has been shown to integrate into biological membranes[37]. Thus, it is possible that PheoA

could insert into the viral envelope lipid bilayer, altering its biophysical properties, or even

disrupting it, thusrendering the virion non-infectious. This would explain PheoA's virucidal

activity and its broad-spectrum among enveloped viruses. However, some degree of selectivity

was observed since PheoA doses that completely abolished infection by several +RNA viruses

did not interfere with VSV or retroviral pseudotype (both also enveloped) infection. It likely is

possible that the membrane's lipidic composition could play a key role for PheoA incorporation.

In this sense, VSV and retroviral pseudotypes are assembled at the plasma membrane[38,39], while

the rest of the tested virus particles are assembled in intracellular compartments[40,41], which display

a very different membrane composition from that of the plasma membrane[42].

On the other hand, PheoA is a plant derived porphyrin closely related to animal porphyrins, which

have been widely described as broad-spectrum virucidals (reviewed in Sh. Lebedeva et al.[43]).

Virion inactivation is thought to occur through incorporation of porphyrins into the viral envelope

membrane and modifying its physico-chemical properties, thus interfering with host cell

recognition and fusion processes. However, porphyrins such as protoporphyrin IX display

antiviral activity independently of their virucidal activity at post-entry steps and have been

proposed to interfere with receptor (ACE2) recognition in SARS-CoV-2 infection models[44]. The

structural resemblance between PheoA and proporphyrin IX may explain their similar antiviral

properties (broad spectrum and virucidal), but, at the same time, their differences may contribute

to PheoA's increased tolerability and *in vivo* effectiveness, an issue that has extensively been

explored for different PheoA applications as photosensitizer in photodynamic therapies against

cancer [45,46].

One huge advantage of PheoA is that it is readily available from plant and algae chlorophyll.

PheoA is the dephytylation and demetallation product of chlorophyll *a*, processes mediated by

chlorophyllase and Mg-dechelatase, respectively [47,48]. Clorophyllase activity is favored by high

temperatures (60-80 ºC)[47] and its accumulation varies throughout plant development and in stress

conditions. In this study, we also exploited stress conditions (heat) that favour PheoA

accumulation and PheoA was semisynthetically prepared from *M. polymorpha* in good overall

yield.

Another advantage of PheoA is that its combination with remdesivir has an additive effect with

no cross inhibition in their antiviral activity, and a mild synergy. This, together with its low

toxicity *in vivo,* represents an advantage that could be clinically exploited[49].

**Materials and Methods**

*Equipment and reagents*

All solvents were of ACS quality unless stated otherwise. Commercially available PheoA was

purchased from Santacruz Biotechnology (>90% by HPLC). A Geno Grinder Spex/SamplePrep

2010 was employed for tissue homogenization. Glass or aluminium supported Silica gel 60

(Merck) was used for preparative and analytical TLCs, respectively; for flash column

purification, silica gel 60 Å, 230-400 mesh, 40-63 $\mu$m was employed. HPLC-UV-MS analysis

was carried out by using a Waters Separations module Alliance e2695 system, a Waters QDa

Detector Acquity QDa and a Waters Photodiode Array Detector 2996. HPLC was performed by

using a Sunfire C18 (4.6 × 50 mm, 3.5 μm) column at 30 °C, with a flow rate of 1 mL/min and

a mobile phase gradient from 70 to 95 of A (formic acid 0.1% in $CH_3CN$) in B (0.1% of formic

acid in $H_2O$) for 10 minutes. Electrospray in positive mode was used for ionization. The HR-

MS analysis was carried out by using an Agilent 1200 Series LC system (equipped with a

binary pump, an autosampler, and a column oven) coupled to a 6520 quadrupole-time of flight

(QTOF) mass spectrometer. $CH_3CN:H_2O$ (75:25, v/v) was used as the mobile phase at 0.2

mL/min. The ionization source was an ESI interface working in the positive-ion mode. The

electrospray voltage was set at 4.5 kV, the fragmentor voltage at 150 V and the drying gas

temperature at 300 °C. Nitrogen (99.5% purity) was used as nebulizer (207 kPa) and drying gas

(6 L/min). The scan range was 50–1100 m/z.

***Preparation of crude M. polymorpha extracts***

Plant material (10 g, fresh weight) was collected and dried in an oven (60 °C) until constant

weight. The dry tissue was ground to a fine powder using a Geno/Grinder (2x 2 min at 2700

rpm) and extracted two times at room temperature with 30 mL of $CHCl_3$:MeOH (2:1, v/v) for at

least 6 h each time. Extracts were combined and concentrated under a nitrogen flow. The

remaining solid was dissolved in DMSO (1 mL) to create the stock solutions employed in the

bioassays.

***Chromatographic fractionation of extracts***

Plants extracts were prepared as described above starting from ca. 20 g of plant material and

directly subjected to silica gel flash column chromatography employing a solvent polarity

gradient starting at *n*-hexane (100%) up to AcOEt:MeOH (4:1, v/v). A total of 56 metabolite-

enriched fractions were obtained. Fractions were analyzed by TLC and those of similar

composition were combined to render 12 new pooled fractions, which were screened for

antiviral activity.

***Preparative TLCs***

Photosynthetic metabolites were extracted [two times, o/n, acetone (30 mL)] from fresh, finely

grounded *M. polymorpha* thallus (ca. 40 g). The combined extracts were concentrated to a final

volume of 10 mL, centrifuged (4000 rcf), filtrated (45 μm, Whatmann PTFE filters) and

chromatographed on preparative TLC plates employing the solvent system AcOEt:MeOH (9:1,

v/v).[1] Selected fluorescent spots (C, D and E) were scraped off, eluted (AcOEt:MeOH, 4:1, v/v)

---

[1] The TLC was heated (heat gun) vigorously at the extract application point before being developed; this procedure enhances Chlorophylls conversion into PheoA.

406 and dried under a nitrogen stream. Single components (C, D and E) were prepared at a 10

407 mg/mL and submitted to both antiviral assays, as described below, and HPLC-UV-MS analysis

408 as described above.

*Semisynthetic preparation of PheoA*

Fresh *M. polymorpha* thallus (ca. 50 g) was oven dried to produce ca. 3.5 g of dry material,

which was ground to a fine powder in the GinoGrinder as described above. The obtained

powder (3 g) was extracted (3x, Acetone, 90 mL), the extracts combined and silica gel (6 g,

ratio 2:1 by weight relative to the dried leaf powder) added. The solvent was evaporated under

reduced pressure to produce an impregnated silica, which was heated (60 °C) overnight to

further potentiate PheoA production. The obtained silica was directly loaded into a flash

column that was run as follows: column ID = 3 cm, silica (70 g), *n*-hexane:AcOEt (1:1, 300

mL), AcOEt:MeOH (9:1, 300 mL; 4:1, 600 mL; 7:3, 300 mL), fractions of ca. 35 mL were

collected. Fractions were analyzed by TLC and the obtained PheoA (3.9 mg, 0.13 % from oven

dried leaf material).

*Cell culture*

Vero E6 (ATCC) and Calu3 (ATCC) cell lines were kindly provided by Dr. Enjuanes (CNB-

CSIC). A549 cells were kindly provided by Dr. Juan Ortín (CNB-CSIC) and Huh7 cells were

kindly provided by Dr. Chisari (TSRI, La Jolla). A549 and Huh7 cells were transduced with a

retroviral vector enabling expression of ACE2 in a di-cistronic expression cassette also conferring

resistance to blasticidine. Transduced populations were selected using 2.5 $\mu$g/mL of blasticidine.

All cell cultures were kept in complete media (DMEM) supplemented with 10 mM HEPES, 1X

non-essential amino acids (gibco), 100 U/mL penicillin-streptomycin (GIBCO) and 10% fetal

bovine serum (FBS; heat-inactivated at 56 °C for 30 min). Unless otherwise stated, all infection

experiments were performed at 37°C in a $CO_2$ incubator (5% $CO_2$) the presence of 2% FBS and

in the absence of selection antibiotics.

 *Viruses*

SARS-CoV-2 (Orthocoronavirinae; Alphacoronavirus; Sarbecovirus; strain NL/2020) was

kindly provided by Dr. R. Molenkamp, Erasmus University Medical Center Rotterdam. SARS-

CoV2 stocks were produced and titrated in VeroE6 cells as described previously (PMID

33917313). VSV-GFP[30] was kindly provided by Dr. Rodriguez (CNB-CSIC). WNV-GFP

recombinant virus was rescued from cloned cDNA as reported previously[29]. Trans-complemented

defective reporter HCV virions (HCVtcp) were produced as described in (Steinmann et al.,

2008)[50]. The hCoV-229E-GFP[51] was kindly provided by Dr. Thiel (University of Basel) and

propagated in Huh7 cells at 33°C in a controlled 5% $CO_2$ environment.

*Cypopathic effect protection assays in Vero E6 and Huh7-ACE2 cells*

Vero E6 or Huh7-ACE2 cell monolayers were inoculated (MOI = 0.001) in the presence of a

wide range of two-fold dilutions of the crude, or partially purified extracts, or pure compounds

and incubated for 72 h. Cytopathic effect and lack thereof was visualized by crystal violet

staining, as previously described[32]. Untreated and solvent-treated cells were included in each plate

as controls.

*Evaluation of the antiviral activity by immunofluorescence microscopy*

Vero E6, A549-ACE2 or Calu3 were seeded onto 96-well plates as described above and infected

in the presence of the indicated compound dose (MOI = 0.01). Twenty-four hours post infection

(48 h for Calu3 cells), cells were fixed for 20 minutes at troom temperature with a 4%

formaldehyde solution in PBS, washed twice with PBS and incubated with incubation buffer (3%

BSA; 0.3% Triton X100 in PBS) for 1 h. A monoclonal antibody against the N protein was diluted

in the incubation buffer (1:2000, v/v; Genetex HL344) and incubated with the cells for 1 h; after

this time, cells were washed with PBS and subsequently incubated with a 1:500 (v/v) dilution of

a goat anti-rabbit conjugated to Alexa 488 (Invitrogen-Carlsbad, CA). Nuclei were stained with

DAPI (Life Technologies) as recommended by the manufacturer during the secondary antibody

incubation. Cells were washed with PBS and imaged using an automated multimode reader

(TECAN Spark Cyto; Austria).

All the infection experiments were performed by mixing the virus and compound dilutions 1:1

(v/v) before addition to the target cells. In the time-of-addition experiments, Vero E6 cultures

were inoculated (MOI from 0.5-1) for 1 h in the presence or absence of the compounds at 37 °C.

Subsequently, virus-compound mixtures were left at all times, or removed and replaced with fresh

2% FBS complete media containing or not the tested compounds (see experimental scheme in

Figure X for details). Cells were fixed 6 h post-inoculation.

**_Viral RNA quantitation by RT-qPCR_**

A549-ACE2 cell monolayers were inoculated at MOI = 0.001 in the presence of non-toxic

concentrations of the compound. Forty-eight hours later, cell supernatants were collected and

heat-inactivated as described in (Smyrlaki et al., 2020)[52], and processed directly for RT-qPCR.

Alternatively, cell lysates were prepared using the Trizol reagent (Thermo Scientific) and the

viral RNA content was determined by RT-qPCR using previously validated sets of primers and

probes specific for the detection of the SARS-CoV-2 E gene and the cellular 18S gene, for

normalization purposes. $\Delta$Ct method was used for relative quantitation of the intracellular viral

RNA accumulation in compound-treated cells compared to the levels in infected cells treated

with DMSO (set as 100%).

**_Cytotoxicity measurement by MTT assays_**

Cell monolayers were seeded in 96-well plates. The day after cells were treated with a wide range

of compound concentrations and forty-eight hours later they were subjected to an MTT assays

using standard procedures[53]. The $CC_{50}$ values were graphically interpolated from dose-response

curves obtained with three biological replicates.

**_Assessment of viral entry using retroviral pseudotypes_**

Retroviral particles pseudotyped with SARS-2-CoV spike envelope protein (Spp) were produced

in HEK293T cells as previously described[54] with materials kindly provided by Dr. F. L. Cosset

(INSERM, Lyon) and J. M. Casasnovas and J. G. Arriaza (CNB-CSIC) for the S protein cDNA.

Particles devoid of envelope glycoproteins were produced in parallel as controls.

*Statitistical Analysis*

Descriptive statistics were calculated using Microsoft Excel. One-way ANOVA and *post-hoc*

tests were calculated using IBM SPSS Software Package (version 26). $EC_{50}$ and $EC_{90}$ values were

obtained employing the PROBIT regression method[55] using IBM SPSS vs26.

Synergy analysis was carried out in the web-based platform Synergy Finder

(https://synergyfinder.fimm.fi/ )[31].

**Acknowledgements**

This research was funded by CSIC (PIE-RD-COVID-19 ref 202040E236 to R.S., PIE-RD-

COVID-19 ref. E202020E079 to P.G and U.G, and CSIC-CoV19-153 ref PIE-202080E221 to

M.G-R.). Funding from EVA (European Virus Archive; grant agreement No 871029) is gratefully

acknowledge. GH. J-A. was supported by the Deutsche Forschungsgemeinschaft (Individual

Research Grant JI 241/2-1).

**References**

1.    Folegatti, P. M. *et al*. Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine

against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised

controlled trial. *Lancet* 1–13 (2020) doi:10.1016/S0140-6736(20)31604-4.

2.    Grein, J. *et al*. Compassionate Use of Remdesivir for Patients with Severe Covid-19. *N.*

*Engl. J. Med.* **382**, 2327–2336 (2020).

3.    Li, X. *et al*. Is hydroxychloroquine bene fi cial for COVID-19 patients ? (2020)

doi:10.1038/s41419-020-2721-8.

4.    Newman, D. J. & Cragg, G. M. Marine natural products and related compounds in

clinical and advanced preclinical trials. *J. Nat. Prod.* **67**, 1216–1238 (2004).

5.    Jan, J. T. *et al.* Identification of existing pharmaceuticals and herbal medicines as

inhibitors of SARS-CoV-2 infection. *Proc. Natl. Acad. Sci. U. S. A.* **118**, 1–8 (2021).

6.    Gu, C. *et al.* Protoporphyrin IX and verteporfin potently inhibit SARS-CoV-2 infection

in vitro and in a mouse model expressing human ACE2. *Sci. Bull.* (2020)

doi:10.1016/j.scib.2020.12.005.

7.    Wachtel-Galor, S. & Benzie, I. F. F. *Herbal Medicine: An Introduction to Its History,*

*Usage, Regulation, Current Trends, and Research Needs. Herbal Medicine:*

*Biomolecular and Clinical Aspects* (2011).

8.    Veeresham, C. Natural products derived from plants as a source of drugs. *J. Adv. Pharm.*

*Technol. Res.* **3**, 200–201 (2012).

9.    Nicolaou, K. C., Guy, R. K. & Potier, P. Taxoids: new weapons against cancer. *Sci. Am.*

**274**, 94–98 (1996).

10.   Efferth, T., Li, P. C. H., Konkimalla, V. S. B. & Kaina, B. From traditional Chinese

medicine to rational cancer therapy. *Trends Mol. Med.* **13**, 353–361 (2007).

11.   Asakawa, Y. & Ludwiczuk, A. Chemical Constituents of Bryophytes: Structures and

Biological Activity. *J. Nat. Prod.* **81**, 641–660 (2018).

12.   Asakawa, Y., Ludwiczuk, A. & Nagashima, F. Phytochemical and biological studies of

bryophytes. *Phytochemistry* **91**, 52–80 (2013).

13.   Asakawa, Y., Ludwiczuk, A. & Nagashima, F. Phytochemical and biological studies of

bryophytes. *Phytochemistry* **91**, 52–80 (2013).

14.  Asakawa, Y., Ludwiczuk, A. & Nagashima, F. *Chemical Constituents of Bryophytes.*

*Progress in the Chemistry of Organic Natural Products* vol. 1 (2013).

15.  Chen, F. *et al.* Terpenoid Secondary Metabolites in Bryophytes: Chemical Diversity,

Biosynthesis and Biological Functions. *CRC. Crit. Rev. Plant Sci.* **37**, 210–231 (2018).

16.  Davies, K. M. *et al.* The Evolution of Flavonoid Biosynthesis: A Bryophyte Perspective.

*Front. Plant Sci.* **11**, 1–21 (2020).

17.  Komala, I., Ito, T., Nagashima, F., Yagi, Y. & Asakawa, Y. Cytotoxic, radical

scavenging and antimicrobial activities of sesquiterpenoids from the Tahitian liverwort

Mastigophora diclados (Brid.) Nees (Mastigophoraceae). *J. Nat. Med.* **64**, 417–422

(2010).

18.  Asakawa, Y., Ludwiczuk, A. & Hashimoto, T. Cytotoxic and Antiviral Compounds from

Bryophytes and Inedible Fungi. *J. Pre-Clinical Clin. Res.* **7**, 73–85 (2014).

19.  Wasternack, C. & Hause, B. Jasmonates: Biosynthesis, perception, signal transduction

and action in plant stress response, growth and development. An update to the 2007

review in Annals of Botany. *Ann. Bot.* **111**, 1021–1058 (2013).

20.  Peñuelas, M. *et al.* Jasmonate-Related MYC Transcription Factors Are Functionally

Conserved in Marchantia polymorpha. *Plant Cell* **31**, 2491–2509 (2019).

21.  Monte, I. *et al.* Ligand-receptor co-evolution shaped the jasmonate pathway in land

plants. *Nat. Chem. Biol.* **14**, 480–488 (2018).

22.  Tanaka, M. *et al.* Direct evidence of specific localization of sesquiterpenes and

marchantin A in oil body cells of Marchantia polymorpha L. *Phytochemistry* **130**, 77–84

(2016).

23.  Romani, F. *et al*. Oil Body Formation in Marchantia polymorpha Is Controlled by

MpC1HDZ and Serves as a Defense against Arthropod Herbivores. *Curr. Biol.* **30**, 2815-

2828.e8 (2020).

24.  Rolin, D. Control of primary metabolism in plants. Annual Plant Reviews, Volume 22.

*Ann. Bot.* **98**, 1331–1332 (2006).

25.  Maeda, H. A. Evolutionary diversification of primary metabolism and its contribution to

plant chemical diversity. *Front. Plant Sci.* **10**, 1–8 (2019).

26.  Ratnoglik, S. L. *et al*. Antiviral activity of extracts from Morinda citrifolia leaves and

chlorophyll catabolites, pheophorbide a and pyropheophorbide a, against hepatitis C

virus. *Microbiol. Immunol.* **58**, 188–194 (2014).

27.  Bouslama, L., Hayashi, K., Lee, J. B., Ghorbel, A. & Hayashi, T. Potent virucidal effect

of pheophorbide a and pyropheophorbide a on enveloped viruses. *J. Nat. Med.* **65**, 229–

233 (2011).

28.  Hamre, D. & Procknow, J. J. A new virus isolated from the human respiratory tract.

*Proc. Soc. Exp. Biol. Med.* **121**, 190–193 (1966).

29.  Pierson, T. C. *et al*. An infectious West Nile virus that expresses a GFP reporter gene.

*Front. Microbiol.* **334**, 28–40 (2005).

30.  Ostertag, D., Hoblitzell-Ostertag, T. M. & Perrault, J. Overproduction of double-stranded

RNA in vesicular stomatitis virus-infected cells activates a constitutive cell-type-specific

antiviral response. *J. Virol.* **81**, 503–513 (2007).

31.  Ianevski, A., Giri, A. K. & Aittokallio, T. SynergyFinder 2.0: visual analytics of multi-

drug combination synergies. *Nucleic Acids Res.* **48**, W488–W493 (2020).

32.    Ginex, T. *et al*. Host-Directed FDA-Approved Drugs with Antiviral Activity against

SARS-CoV-2 Identified by Hierarchical In Silico/In Vitro Screening Methods.

*Pharmaceuticals (Basel)*. **14**, (2021).

33.    Asakawa, Y. Biologically active compounds from bryophytes. *Pure Appl. Chem.* **79**,

557–580 (2007).

34.    Chansakaow, S., Ruangrungsi, N. & Ishikawa, T. Isolation of pyropheophorbide a from

the leaves of Atalantia monophylla (ROXB.) CORR. (Rutaceae) as a possible antiviral

active principle against herpes simplex virus type 2. *Chem. Pharm. Bull. (Tokyo)*. **44**,

1415–1417 (1996).

35.    Ohta, S. *et al*. Anti-herpes simplex virus substances produced by the marine green alga,

Dunaliella primolecta. *J. Appl. Phycol.* **10**, 349–356 (1998).

36.    Zhang, H.-J. *et al*. Natural anti-HIV agents. Part IV. Anti-HIV constituents from Vatica

cinerea. *J. Nat. Prod.* **66**, 263–268 (2003).

37.    RÖDER, B., HANKE, T. H., OELCKERS, S. T., HACKBARTH, S. T. & SYMIETZ, C.

H. Photophysical properties of pheophorbide a in solution and in model membrane

systems. *J. Porphyrins Phthalocyanines* **04**, 37–44 (2000).

38.    Drokhlyansky, E., Soh, T. K. & Cepko, C. L. Preferential Budding of Vesicular

Stomatitis Virus from the Basolateral Surface of Polarized Epithelial Cells Is Not Solely

Directed by Matrix Protein or Glycoprotein. *J. Virol.* **89**, 11718–11722 (2015).

39.    Morita, E. & Sundquist, W. I. Retrovirus budding. *Annu. Rev. Cell Dev. Biol.* **20**, 395–

425 (2004).

40.    De Wit, E., Van Doremalen, N., Falzarano, D. & Munster, V. J. SARS and MERS:

Recent insights into emerging coronaviruses. *Nat. Rev. Microbiol.* **14**, 523–534 (2016).

41.    Hirsch, A. J. *et al.* The Src family kinase c-Yes is required for maturation of West Nile

594           virus particles. *J. Virol.* **79**, 11943–11951 (2005).

595    42.    Meer, G. Membrane lipids, where they are and how they behave: Sphingolipids on the

596           move. *FASEB J.* **24**, 112–124 (2010).

597    43.    Sh Lebedeva, N., A Gubarev, Y., O Koifman, M. & I Koifman, O. The Application of

598           Porphyrins and Their Analogues for Inactivation of Viruses. *Molecules* **25**, (2020).

599    44.    Gu, C. *et al.* Protoporphyrin IX and verteporfin potently inhibit SARS-CoV-2 infection

600           in vitro and in a mouse model expressing human ACE2. *Sci. Bull.* **66**, 925–936 (2021).

601    45.    Xodo, L. E., Rapozzi, V., Zacchigna, M., Drioli, S. & Zorzet, S. The chlorophyll

602           catabolite pheophorbide a as a photosensitizer for the photodynamic therapy. *Curr. Med.*

603           *Chem.* **19**, 799–807 (2012).

604    46.    Bellnier, D. A. *et al.* Clinical pharmacokinetics of the PDT photosensitizers porfimer

605           sodium (Photofrin), 2-[1-hexyloxyethyl]-2-devinyl pyropheophorbide-a (Photochlor) and

606           5-ALA-induced protoporphyrin IX. *Lasers Surg. Med.* **38**, 439–444 (2006).

607    47.    Yilmaz, C. & Gökmen, V. Chlorophyll. *Encycl. Food Heal.* 37–41 (2016)

608           doi:10.1016/B978-0-12-384947-2.00147-1.

609    48.    Saide, A., Lauritano, C. & Ianora, A. Pheophorbide A: State of the art. *Mar. Drugs* **18**,

610           1–12 (2020).

611    49.    Hajri, A. *et al.* In Vitro and In Vivo Efficacy of Photofrin® and Pheophorbide a, a

612           Bacteriochlorin, in Photodynamic Therapy of Colonic Cancer Cells¶ *J. Photochem.*

613           *Photobiol.* **75**, 140 (2002).

50. Steinmann, E., Brohm, C., Kallis, S., Bartenschlager, R. & Pietschmann, T. Efficient

trans-encapsidation of hepatitis C virus RNAs into infectious virus-like particles. *J.*

*Virol*. **82**, 7034–7046 (2008).

51. Cervantes-Barragan, L. *et al*. Dendritic cell-specific antigen delivery by coronavirus

vaccine vectors induces long-lasting protective antiviral and antitumor immunity. *MBio*

**1**, 227 (2010).

52. Smyrlaki, I. *et al*. Massive and rapid COVID-19 testing is feasible by extraction-free

SARS-CoV-2 RT-PCR. *Nat*. *Commun*. **11**, 4812 (2020).

53. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to

proliferation and cytotoxicity assays. *J*. *Immunol*. *Methods* **65**, 55–63 (1983).

54. García-Arriaza, J. *et al*. COVID-19 vaccine candidates based on modified vaccinia virus

Ankara expressing the SARS-CoV-2 spike induce robust T- and B-cell immune

responses and full efficacy in mice. *J*. *Virol*. (2021).

55.  Cleophas, T. J. & Zwinderman, A. H. Probit Models for Estimating Effective

Pharmacological Treatment Dosages (14 Tests) BT  - Machine Learning in Medicine –

A Complete Overview. in *Machine Learning in Medicine – A Complete Overview* (eds.

Cleophas, T. J. & Zwinderman, A. H.) 347–354 (Springer International Publishing,

2020).

**FIGURES**

**Figure 1:** Marchantia extracts interfere with SARS-CoV-2-induced cytopathic effect.
Vero E6 cells were inoculated with SARS-CoV-2 at MOI = 0.001 in the presence of serial 2-fold dilutions of crude extracts from two different Marchantia ecotypes, *ruderalis* (Ex1) and *polymorpha* (Ex2) and incubated for 72 h, time after which they were fixed and stained with a crystal violet solution. Mock-infected cells were used as the control of the integrity of the cell monolayer. A) Image of an experimental plate showing dose-dependent protection of *M. polymorpha* extracts in comparison with vehicle (DMSO)-treated cells. B) Numeric expression of the protective capacity as the $\log_2$ value of the highest dilution factor capable of full-monolayer protection. Data are shown as average and mean error of two biological replicates.

**Figure S1:** Remdesivir interferes with SARS-CoV-2-induced cytopathic effect.
Vero E6 cells were inoculated with SARS-CoV-2 at MOI = 0.001 in the presence of serial 2-fold dilutions of remdesivir (50-0.78 µM) and incubated for 72 h, time after which they were fixed and stained with a crystal violet solution. Representative image showing that interference with viral multiplication by remdesivir reveals its protective activity against virus-induced cytopathic effect at the expected concentrations.

**Figure 2:** Marchantia extracts interfere with SARS-CoV-2 propagation.
Vero-E6 cells were inoculated with SARS-CoV-2 (MOI = 0.001) in the presence of vehicle (DMSO), remdesivir (RMDV; 25µM) or a 1:800 (v/v) dilution of crude extracts from Marchantia ecotypes, *ruderalis* (Ex1) and *polymorpha* (Ex2). Parallel samples remained uninfected as control (mock). Total RNA was extracted 72 h post-infection and subjected to RT-qPCR to determine viral load. Normalized viral RNA levels are shown as percentage of the viral RNA found in vehicle-treated cells. Data are shown as mean (± SD) of three biological replicates. Statistical significance was estimated using one-way ANOVA and a Dunnet´s post-hoc test (*p<0.05).

**Figure 3:** Antiviral candidates are not likely derived from Marchantia´s secondary metabolism. Vero E6 cells were inoculated with SARS-CoV-2 (MOI = 0.001) in the presence of serial 2-fold dilutions of crude extracts from WT, Mp*coi1-2* or Mp*c1hdz* Marchantia plants. Cultures were incubated for 72 h, time after which they were fixed and stained with a crystal violet solution. Mock-infected cells were used as the control of the integrity of the cell monolayer. A) Representative experiment showing dose-dependent protection of plant extracts in comparison with vehicle (DMSO)-treated cells. B) Numeric expression of the protective capacity as the $\log_2$ value of the highest dilution factor capable of full monolayer protection. Data are shown as average (± mean error) of two biological replicates.

**Figure S2:** Crude extracts of several plant species show antiviral potential against SARS-CoV-2 infection in cell culture.

Vero E6 cells were inoculated with SARS-CoV-2 at MOI of 0.001 in the presence of serial 2-fold dilutions of crude extracts from sweet amber (*Hypericum androsaemum*), fern (*Blechnum spicant*), nettle (*Urtica dioica*), moss (*Physcomitrium patens*), tocacco (*Nicotiana benthamiana*), thale cress (*Arabidopsis thaliana*). Cultures were incubated for 72 h, time after which they were fixed and stained with a crystal violet solution. Mock-infected cells were used as a control of the integrity of the cell monolayer. A) Image of an experimental plate showing the dose-dependent protection of plant extracts in comparison with vehicle (DMSO)-treated cells. B) Numeric expression of the protective capacity as the $\log_2$ value of the highest dilution factor capable of full monolayer protection. Data are shown as average and mean error of two biological replicates.

**Figure 4:** Extract fractionation and identification of antiviral fraction pools.
Vero E6 cells were inoculated with SARS-CoV-2 (MOI = 0.001) in the presence of serial 2-fold dilutions of vehicle (DMSO), a crude Marchantia and the fraction pools. Inoculated cultures were incubated for 72 h, time after which they were fixed and stained with a crystal violet solution. Mock-infected cells were used as the control of the integrity of the cell monolayer (non-infected). A) Experimental plates showing cell monolayer integrity (purple) in the presence of fractions containing the antiviral compound(s). B) Numeric expression of the protective capacity as the $\log_2$ value of the highest dilution factor capable of full monolayer protection. Data are shown as average and mean error of two biological replicates. C) Vero E6 cells were inoculated (MOI = 0.01) in the presence of the indicated fraction dilutions and incubated for 24 h before fixation and processing for immunofluorescence microscopy and cytotoxicity assays as described in the materials and methods section. Data are shown as average (± SD) of three biological replicates. Statistical significance was estimated using one-way ANOVA (Dunnet´s post-hoc test, $p < 0.05$).

**A**

**Fraction pool number**

**B**

**Fraction pool number**

**Figure S3:** A) TLC plate of chromatographic fractions under long wave ultraviolet light (365 nm). B) TLC of chlorophyll containing fractions after heating. Red fluorescent spots, characteristic of plant chlorophylls are squared. Solvent system AcOEt:MeOH (9:1, v/v). arrows heads depict the solvent front.

**Figure 5:** Identification of the main antiviral molecule present in Marchantia extracts.
A) Representative TLC of WT and Mp*c1hdz* marchantia extracts. Compounds C, D and E (box) were tested for their antiviral potential. B) Vero E6 cells were inoculated (MOI = 0.01) in the presence of the indicated compounds and incubated for 24 h before fixation and processing for immunofluorescence microscopy. Data are shown as average and SD of three biological replicates. Statistical significance was estimated using one-way ANOVA and a Dunnet´s post-hoc test (*p<0.05). C) Representative HPLC/MS analysis (shown for WT) of the C, D, E compounds, including exact mass determination of the antiviral candidate **1** and the inferred chemical structure.

**Figure S4:** Scheme of Pheophobide a preparation from plant material. KG, Silica gel 60. AcOEt, ethyl acetate. MeOH, methanol.

# Vero-E6

# Huh7-ACE2

**Figure S5:** Pheophorbide a protects Vero-E6 and human hepatoma Huh7 cells from virus-induced cytopathic effect.

Vero-E6 or Huh7 cells were inoculated with SARS-CoV-2 at MOI = 0.001 in the presence of the indicated PheoA concentrations. Cultures were incubated for 72 h, time after which they were fixed and stained with a crystal violet solution. Mock-infected (N.I.) cells were used as a control of the integrity of the cell monolayer.

**Figure 6:** Pheophorbide A shows antiviral activity against SARS-CoV-2 in Vero E6 cells and human lung epithelial A549-ACE2 and Calu3 cell lines.
Commercially available PheoA was serially diluted and mixed (1:1) with SARS-CoV-2 preparations to achieve the indicated compound concentrations and a final MOI of 0.005 for Vero E6 and Calu3 and 0.01 for A549-ACE2 cells. Cultures were incubated for 48 h, fixed and processed for automated immunofluorescence microscopy analysis. Parallel, uninfected cultures were processed for cytotoxicity evaluation using an MTT assay. Relative infection efficiency data (N=3 per dose) are shown as individual data and a PROBIT regression curve (green line) using the represented values. Cytotoxicity data (N=3 per dose) are shown as the individual data and a moving average trendline.

**Figure 7:** Pheophorbide A interferes with viral propagation in human cell lines as determined by RT-qPCR.

A549-ACE2 cells were inoculated at MOI = 0.01 in the presence of increasing concentrations of PheoA or 5000 nM remdesivir and incubated for 48 h. Samples of the supernatants were collected, heat-inactivated and directly subjected to RT-qPCR to estimate overall infection efficiency. Data are expressed as relative values compared with the vehicle (DMSO)-treated cells and are shown as mean and standard deviation of three biological replicates (N=3). Statistical significance was estimated using one-way ANOVA and a Dunnet´s post-hoc test (*p<0.05).

**Figure S6:** Semi-synthetic PheoA preparations display antiviral activity against SARS-CoV-2. Semi-synthetic PheoA was produced as described in Figure S4, serially diluted and mixed 1:1 with SARS-CoV-2 preparations to achieve the indicated compound concentrations and a final multiplicity of infection of 0.005 for Vero-E6 and Calu3, and 0.01 for A549-ACE2 cells. Cultures were incubated for 48 h, fixed and processed for automated immunofluorescence microscopy analysis. Relative infection efficiency data (N=3 per dose) are shown as individual data and a PROBIT regression curve (green line) using the represented values.

**Figure S7:** Pyropheophorbide a displays antiviral activity against SARS-CoV-2. Commercially available pPheoA was serially diluted and mixed 1:1 with SARS-CoV-2 preparations to achieve the indicated compound concentrations and a final multiplicity of infection of 0.005 for Vero E6 and Calu3, and 0.01 for A549-ACE2 cells. Cultures were incubated for 48 h, fixed and processed for automated immunofluorescence microscopy analysis. Relative infection efficiency data (N=3 per dose) are shown as individual data and a PROBIT regression curve (green line) using the represented values.

**Figure 8:** Antiviral spectrum of PheoA against different RNA viruses.
The effectiveness of PheoA was tested against four different viruses recombinant RNA viruses expressing GFP. Briefly, cells were inoculated in the presence of increasing concentrations of PheoA at MOI 0.01 and incubated to enable virus propagation. At the endpoint, cells were fixed and counter-stained with DAPI to control for unexpected cytotoxic effects. Relative infection efficiency was estimated using automated microscopy and is expressed as percentage of the infection efficiency observed in control wells. A) …… is missing. B) Huh7 cells were infected with hCoV-229E-GFP and fixed 48 h post-inoculation. C) Huh7 cells were infected with WNV-GFP and fixed 48 h post-inoculation. D) A549-ACE2 cells were inoculated with VSV-GFP and fixed 16 h post-inoculation. Individual replicate data are shown as green dots (N=3) and the PROBIT regression curve used to estimate $EC_{50}$ values is shown.

**Figure 9:** Combination treatment of Pheophorbide a with remdesivir.
Vero E6 cells were inoculated at MOI = 0.005 in the presence of increasing concentrations of PheoA in combination with increasing doses of remdesivir. Twenty-four hours post infection, cells were fixed and processed for automated immunofluorescence microscopy. Relative infection efficiency values were estimated as percentage of the values obtained in mock-treated cells.  A) Data are shown as average of two biological replicates. B) Heatmap describing the areas of synergy within the combination treatments.

**Figure 10:** Time-of-addition experiments indicate that PheoA interferes with early aspects of SARS-CoV-2 infection.

Vero E6 cells were inoculated at MOI = 5 in the presence (gray) or absence (white) of the indicated doses of PheoA or imatinib as described in both the text and the scheme. Cells were incubated for 6 h in the presence (gray) or absence (white) before chemical fixation and processing for immunofluorescence microscopy. Infection efficiency is expressed as the percentage of that observed in vehicle DMSO-treated cells and is shown as average and standard deviation of three biological replicates (N=3). Statistical significance was estimated using one-way ANOVA and a Dunnet´s post-hoc test (*p<0.05).

**Figure 11:** SARS-CoV-2-pseudotyped retroviral vectors (Spp) are not susceptible to PheoA antiviral activity.

Vero E6 cells were inoculated with retroviral pseudotypes bearing the SARS-CoV-2 Spike (SARS2pp) or the VSV-G glycoprotein (VSVpp) in the presence of the indicated compound concentrations. Forty-eight hours post-infection, cells were lysed and infection efficiency was estimated by the luciferase reporter gene activity. Relative infection values were calculated as percentage of the luciferase activity observed in vehicle (DMSO)-treated cells and is shown as average and standard deviation of three biological replicates (N=3). Statistical significance was estimated using one-way ANOVA and a Dunnet´s post-hoc test (*p<0.05).

**Figure 12:** Pheophorbide A shows virucidal activity against SARS-CoV-2.
SARS-CoV-2 virus stocks were diluted to obtain $1*10^5$ TCID50/mL and were mixed with increasing concentrations of PheoA or the vehicle (DMSO). Virus-compound mixtures were incubated at room temperature for 30 minutes and were serially diluted to determine the remaining infectivity titer using endpoint dilution and determination of virus-induced cytopatic effect by crystal violet staining. Values are expressed as LOG TCID50/mL and shown as the average and standard deviation of three independent experiments (N=3). Statistical significance was estimated using one-way ANOVA and a Dunnet´s post-hoc test (*p<0.05).