Both consumptive and non-consumptive effects of predators

impact mosquito populations and have implications for disease

transmission

Marie C. Russell[a]*, Catherine M. Herzog[b]*, Zachary Gajewski[c], Chloe Ramsay[d], Fadoua El

Moustaid[c], Michelle V. Evans[e,f] Trishna Desai[g], Nicole L. Gottdenker[e,h], Sara L. Hermann[i],

Alison G. Power[j], and Andrew C. McCall[k]

*Co-first authorship; for correspondence: Marie C. Russell, marie.clare.russell@gmail.com

[a]Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot, UK

[b]Center for Infectious Disease Dynamics, Pennsylvania State University, University Park,

PA, USA

[c]Department of Biological Sciences, Virginia Polytechnic Institute and State University,

Blacksburg, VA, USA

[d]Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA

[e]Odum School of Ecology & Center for Ecology of Infectious Diseases, University of

Georgia, Athens, GA, USA

[f]MIVEGEC, IRD, CNRS, Université Montpellier, Montpellier, France

[g]Nuffield Department of Population Health, University of Oxford, Oxford, UK

[h]Department of Veterinary Pathology, University of Georgia College of Veterinary Medicine,

Athens, GA, USA

[i]Department of Entomology, Pennsylvania State University, University Park, PA, USA

[j]Department of Ecology & Evolutionary Biology, Cornell University, Ithaca, NY, USA

[k]Biology Department, Denison University, Granville, OH, USA

Abstract

Predator-prey interactions influence prey traits through both consumptive and non-

consumptive effects, and variation in these traits can shape vector-borne disease dynamics.

Meta-analysis methods were employed to generate predation effect sizes by different

categories of predators and mosquito prey. This analysis showed that multiple families of

aquatic predators are effective in consumptively reducing mosquito survival, and that the

survival of *Aedes*, *Anopheles*, and *Culex* mosquitoes is negatively impacted by consumptive

effects of predators. Mosquito larval size was found to play a more important role in

explaining the heterogeneity of consumptive effects from predators than mosquito genus.

Mosquito survival and body size were reduced by non-consumptive effects of predators, but

development time was not significantly impacted. In addition, *Culex* vectors demonstrated

predator avoidance behavior during oviposition. The results of this meta-analysis suggest that

predators limit disease transmission by reducing both vector survival and vector size, and that

associations between drought and human West Nile virus cases could be driven by the vector

behavior of predator avoidance during oviposition. These findings are likely to be useful to

infectious disease modelers who rely on vector traits as predictors of transmission.

Introduction

While it is well-known that predation reduces vector populations through consumptive

effects, non-consumptive effects of predators can also greatly impact prey demographics

(Preisser et al., 2005). Mosquitoes are vectors of a variety of debilitating and deadly diseases,

including malaria, lymphatic filariasis, and arboviruses, such as chikungunya, Zika, and

dengue (Weaver and Reisen, 2010, World Health Organization (WHO), 2020). Consequently,

there is motivation from a public health perspective to better understand the different drivers

of variation in mosquito traits that can ultimately impact vector population growth and

disease transmission. In addition, recent work has suggested that incorporation of vector trait

variation into disease models can improve the reliability of their predictions (Cator et al.,

2020). In this study, systematic review and meta-analysis methods are used to synthesize a

clearer understanding of the consumptive and non-consumptive effects of predators on

mosquito traits, including survival, oviposition, development, and size.

Mosquito insecticide resistance is recognized as a growing problem (Hancock et al., 2018,

Hemingway and Ranson, 2000, Liu, 2015) leading some to suggest that control efforts should

rely more heavily on "non-insecticide based strategies" (Benelli et al., 2016). The

consumptive effects of predators on mosquitoes have previously been harnessed for

biocontrol purposes. Past biocontrol efforts have used predators such as cyclopoid copepods

(Kay et al., 2002, Marten, 1990, Russell et al., 1996, Veronesi et al., 2015) and mosquitofish

(Pyke, 2008, Seale, 1917) to target the mosquito's aquatic larval stage. The strength of the

consumptive effects of these predators on mosquitoes can be influenced by multiple factors,

including predator-prey size ratio and temperature. Predator-prey body size ratios tend to be

higher in freshwater habitats than other types of habitats (Brose et al., 2006), and attack rate

tends to increase with temperature (Kalinoski and DeLong, 2016, Dam and Peterson, 1988),

though other studies suggest a unimodal response to temperature (Uiterwaal and Delong,

2020, Englund et al., 2011).

Predators can also have non-consumptive effects on prey (Peacor and Werner, 2001), and

these effects are thought to be more pronounced in aquatic ecosystems than in terrestrial

ecosystems (Preisser et al., 2005). Non-consumptive effects of predators are the result of the

prey initiating anti-predator behavioral and/or physiological trait changes that can aid in

predator avoidance (Hermann and Landis, 2017, Lima and Dill, 1990). Such plasticity in

certain prey traits may also result in energetic costs (Lima, 1998). Predator detection is key

for these trait changes to occur and can be mediated by chemical, tactile, and visual cues

(Hermann and Thaler, 2014). In mosquitoes, exposure to predators is known to affect a

variety of traits including behavior, size, development, and survival (Arav and Blaustein,

2006, Bond et al., 2005, Roberts, 2012, Roux et al., 2015, Zuharah et al., 2013). Experimental

observations of predator effects on mosquito size and development are inconsistent and

results sometimes vary by mosquito sex. For example, exposure to predation was found to

increase the size of *Culex pipiens* mosquitoes (Alcalay et al., 2018) but decrease the size of

Culiseta longiareolata (Stav et al., 2005). In addition, female *Aedes triseriatus* exhibited

shorter development times when exposed to predation at high nutrient availability (Ower and

Juliano, 2019), but male *C. longiareolata* had longer development times in the presence of

predators (Stav et al., 2005). In some cases, a shared evolutionary history between predator

and prey organisms can strengthen the non-consumptive effects of predators on mosquitoes

(Buchanan et al., 2017, Sih, 1986).

This investigation assesses the consumptive and non-consumptive effects of predators on

mosquito traits and describes how these effects could impact disease transmission. The roles

of vector genus, predator family, mosquito larval instar (an indicator of prey size), and

temperature are also examined as potential moderators of predator effects. Non-consumptive

effects of predators are expected to cause a smaller reduction in mosquito survival than

consumptive effects because, in practice, measures of consumptive effects always include

both consumptive and non-consumptive effects. Based on previous findings, larger predators

are more likely to consumptively reduce mosquito survival (Kumar et al., 2008). In addition,

Aedes mosquito larvae may be more vulnerable to consumptive predation than other genera

because of the high degree of motility observed in this genus (Dieng et al., 2003, Marten and

Reid, 2007, Soumare and Cilek, 2011). The oviposition response to predation is expected to

be weakest among *Aedes* species that oviposit above the water line, due in part to their

delayed-hatching eggs (Vonesh and Blaustein, 2010). Predation is predicted to reduce

mosquito size and lengthen development time, consistent with the reduced growth response

observed in other insect systems (Hermann and Landis, 2017). Certain non-consumptive

effects of predation, particularly oviposition site selection and decreased vector size, are

likely to play important roles in the dynamics of mosquito-borne disease.

Materials and Methods

Literature Screening:

A systematic search was conducted for studies on predation of mosquitoes that were

published between 1970 and July 1, 2019 using both PubMed® and Web of Science™ search

engines, according to the PRISMA protocol (Moher et al., 2009). Mosquito vectors of the

Anopheles and *Aedes* genera were specifically highlighted in our search terms because these

genera contain the vector species that transmit malaria, yellow fever, and dengue – the three

most deadly mosquito-borne diseases worldwide (Hill et al., 2005). Searches included 18

combinations of three vector predation terms (mosquito predat*, *Anopheles* predat*, *Aedes*

predat*) and six trait terms (survival, mortality, development, fecundity, dispers*, host

preference). Abstracts from the 1,136 studies were each screened by two different co-authors,

using the "metagear" package in R (Lajeunesse, 2016, R Core Team, 2020). If either screener

thought the study had information relevant to predation of mosquitoes, or both screeners

thought the abstract was ambiguous, the study was read in full. This resulted in 306 studies

that were fully reviewed to determine if any predation data could be extracted (Fig 1).

Fig 1. Flowchart demonstrating the literature search, screening process, data exclusions, and the resulting seven different vector trait data subsets

Study Exclusion Criteria:

Data were extracted from studies that collected data on non-consumptive and/or consumptive

effects of predators on mosquitoes. Studies were required to have a mean, error measurement,

and at least two replicates for both control and predator treatments. The control treatment was

required to have all the same conditions as the predator treatment, such as prey density and

type of water, without the predators. Studies that were not published in English and studies

that did not differentiate between predators of multiple families were excluded. Studies were

also excluded if oviposition by free-flying female mosquitoes could have interfered with

observing the consumptive effects of predators on vector survival. The final database

comprised data extracted from 60 studies (Table S1). The data included observations from

laboratory experiments, as well as semi-field experiments, in which mesocosms of different

treatments were observed in outdoor settings.

Data Extraction:

Variables related to the publication, the vector, the predator, and the effect size (Table 1)

were extracted from each study. Data from tables and text were recorded as they were

published, and data from figures were extracted using WebPlotDigitizer (Rohatgi, 2020).

Error measurements that were not originally presented as standard deviations were converted

to standard deviations prior to the effect size calculation.

Table 1: Variables extracted from included studies
152

Variable	Description
Publication data:	
Title	Full study title
Journal	Name of journal that published the study
Year	Year of publication
Study environment	Environment where the experiment took place: lab or semi-field
Vector data:	
Order, Family, Genus, Species	Taxonomic identification
Trait	Outcome that was measured (e.g., survival, development, etc.)
Stage	Life stage: egg, larva, pupa, or adult
Larval instar	Early (1^{st} and 2^{nd} instars), late (3^{rd} and 4^{th} instars), both, or NA (eggs, pupae, or adults)
Sex	Male or female
Predator data:	
Phylum, Class, Order, Family, Genus, Species	Taxonomic identification
Starved	Whether the predator was starved: yes or no
Time starved	Amount of time that the predator was starved (in minutes)
Predation effect	Consumptive or non-consumptive
Effect size data:	
Units	Units of extracted data
Control mean	Average of the outcome measured among the controls
Control standard deviation	Standard deviation of the outcome measured in the controls
Control number of replicates	Number of control replicates
Predation mean	Average of the outcome measured in the predator treatment
Predation standard deviation	Standard deviation of the outcome measured in the predator treatment
Predation number of replicates	Number of predation replicates
Experiment ID	Alphabetic assignment to mark observations sharing a control group or representing the same prey individuals as originating from the same experiment
Additional data:	
Experiment time (days)	Duration of the experiment in days
Data source	Graph or text
Number of predators	Number of predators with access to prey, or "cue" if there are no predators with direct access to prey
Number of prey (vectors)	Number of mosquito prey that are exposed to predation
Arena volume (mL)	Volume of the arena where prey encounter predators
Time exposed to predator(s)	Amount of time (in days) when the predator has direct access to the mosquito prey
Temperature (°C)	Temperature during the predation interaction
Type of predator cue	Predator cues, or cues from both predator(s) and dying conspecifics; NA for observations with a consumptive predation effect

Data Exclusions:

A PRISMA plot of literature inclusion and exclusion is provided in Figure 1. Observations

where insecticide was used were excluded because insecticides are known to interfere with

consumptive and non-consumptive effects of predators (Delnat et al., 2019, Janssens and

Stoks, 2012). In addition, observations from experiments with mosquito prey of two or more

species were excluded because it was not possible to account for effects from apparent

competition or prey-switching. Observations of vector fecundity, vector competence,

behavioral traits other than oviposition, as well as observations where the vector trait was

marked as "other" were not analyzed because each of these traits were only recorded from

three or fewer studies.

Due to protandry, the earlier emergence of males to maximize their reproductive success,

mosquitoes respond to sex-specific selective forces that influence their development time and

body size (Kleckner et al., 1995). Under low resource conditions, female mosquitoes are

likely to maximize body mass by extending their development time, whereas males tend to

minimize their development time at the expense of lower body mass (Kleckner et al., 1995).

Observations of mosquito development time and body size in our database that were not sex-

specific were excluded so that these vector traits could be analyzed while controlling for sex.

In addition, among the observations of development time and body size, some predator

means did not necessarily represent an evenly weighted average of the replicates. For

example, if a total of twenty mosquitoes from three different predator replicates survived to

adulthood, the mean development time and size of those twenty individuals may have been

reported. To represent an evenly weighted average of the replicates, it is necessary to first

calculate summary statistics among multiple individuals that emerge from the same replicate,

and then report the average of the replicate-specific means. Observations that might have

been influenced by uneven representation of replicates were excluded to prevent pseudo-

replication from altering later meta-analyses.

For consumptive observations where life stage-specific survival was reported after more than

10 days of predator exposure, only data on survival marked by adult emergence were

included for analysis. Effects observed among immature vector stages after such a long

period of predator exposure were not analyzed because they could have resulted from a

combination of non-consumptive effects on development, and consumptive effects on

survival. Development time observations that were reported as the inverse of development

time (units of days^{-1}) were excluded because although their means could be converted to units

of days, their standard deviations could not be converted to match units of days. In cases

where multiple body sections of the same mosquitoes were measured to produce multiple size

observations, only the wing measurement was included in the analysis to prevent pseudo-

replication. Observations in which both the control and the predator treatments had standard

deviations of zero were excluded because the meta-analysis methods did not support non-

positive sampling variances.

Fig 2. Mosquito predator classes (bold font) and families (italicized font) included in the database and the vector traits that they may influence (in parentheses); predator images not to scale, and placed randomly with respect to the different mosquito life stages

Image sources:

phylopic.org (CC BY 3.0 or public domain): Actinopterygii (creator: Milton Tan), Arachnida (creators: Sidney Frederic Harmer & Arthur Everett Shipley, vectorized by Maxime Dahirel), Branchiopoda (creator: Africa Gomez), and Insecta (creator: Marie Russell)

BioRender.com: Amphibia, Hexanauplia, and Malacostraca class silhouettes; mosquito larval instars, pupa, and blood-feeding adult

Trishna Desai: mosquito egg raft

Exclusions and Data Substitutions for Predator Treatment Means of Zero:

One study that was included in our database reported egg survival data as the hatch rate of

field collected *Culex pervigilans* rafts (Zuharah et al., 2013). However, mosquitoes have been

shown to lay eggs independent of mating (O'Meara, 1979), and hatch rates of zero have

previously been observed in rafts laid by *Culex* females that were held separately from males

(Su and Mulla, 1997). Thus, hatch rates of zero were excluded from further analysis because

these values may represent unfertilized egg rafts, rather than a strong impact of predators on

survival. Twenty of the 187 consumptive survival observations had a predation mean of zero,

and each of these zeros resulted from experiments that began with a specified number of live

larvae. Consumptive survival zeros were each replaced with 0.5% of the starting number of

mosquito prey to avoid undefined effect sizes. In addition, there was one zero out of the 36

oviposition predation means; this value had units of "number of egg rafts laid" and was

replaced with 0.5 rafts. Similar methods for replacing zero values in the treatment mean with

small non-zero values have previously been employed (Thapa et al., 2018).

The final analysis dataset included seven subsets: consumptive effects on survival, non-

consumptive effects on survival, oviposition, development (female and male), and size

(female and male). The data included 187 observations from 34 studies of consumptive

survival, 24 observations from seven studies of non-consumptive survival, 36 observations

from 12 studies of oviposition, 14 observations from seven studies of female development, 14

observations from seven studies of male development, 27 observations from 10 studies of

female size, and 18 observations from nine studies of male size (Fig 1). These observations

covered seven different classes of predator families (Fig 2).

Data Analysis:

Measuring Effect Sizes and Heterogeneity

All analyses were conducted in R version 4.0.2 (R Core Team, 2020). For each subset of trait

data (Fig 1), the ratio of means (ROM) measure of effect size was calculated using the

"escalc" function from the "metafor" package; this effect measure is equal to a log-

transformed fraction, where predation mean is the numerator and control mean is the

denominator (Viechtbauer, 2010). Random effects models, using the "rma.uni" function,

were run with the ROM effect sizes as response variables; each model had a normal error

distribution and a restricted maximum likelihood (REML) estimator for τ^2, the variance of the

distribution of true effect sizes (Viechtbauer, 2010). Although these random effects models

could not account for multiple random effects or moderators, they provided overall estimates

of the ROM effect sizes and estimates of the I^2 statistics. Each I^2 statistic represented the

percentage of total variation across studies due to heterogeneity (Higgins et al., 2003). If the

I^2 statistic was equal to or greater than 75%, the heterogeneity was considered to be high

(Higgins et al., 2003), and high heterogeneity has previously motivated further testing of

moderators (Vincze et al., 2017).

Assessing Publication Bias

Publication bias was assessed by visually inspecting funnel plots and conducting Egger's

regression test ("regtest" function) with standard error as the predictor (Sterne and Egger,

2001, Viechtbauer, 2010). If the Egger's regression test showed significant evidence of

publication bias based on funnel plot asymmetry, the "trim and fill" method ("trimfill"

function) was used to estimate how the predation effect size might change after imputing

values from missing studies (Duval and Tweedie, 2000b, Duval and Tweedie, 2000a,

Viechtbauer, 2010). The trim and fill method has previously been recommended for testing

the robustness of conclusions related to topics in ecology and evolution (Jennions and Moller,

2002). Of the two trim and fill estimators, R_0 and L_0, that were originally recommended

(Duval and Tweedie, 2000b, Duval and Tweedie, 2000a), the L_0 estimator was used in this

study because it is more appropriate for smaller datasets (Shi and Lin, 2019).

Testing Moderators

Data subsets that had high heterogeneity, observations from at least 10 studies, and no

evidence of publication bias according to Egger's regression results were analyzed further

using multilevel mixed effects models with the "rma.mv" function (Viechtbauer, 2010,

Higgins et al., 2020). All multilevel mixed effects models had normal error distributions,

REML estimators for τ^2, and accounted for two random factors: effect size ID, and

experiment ID nested within study ID. Moderators, such as predator family, vector genus,

larval instar (directly correlated to prey size), and temperature, were tested within each data

subset to determine if they affected the observed heterogeneity in ROM effect sizes. For

categorical moderators, the intercept of the multilevel mixed effects model was removed,

allowing an analysis of variance (ANOVA) referred to as the "test of moderators" to indicate

if any of the categories had an effect size different than zero. For data subsets with

observations from 10 to 29 studies, only one moderator was tested at a time to account for

sample size constraints. For subsets with observations from a higher number of studies (30 or

more), up to two moderators were tested at once, and interaction between moderators was

also tested. The small sample corrected Akaike Information Criterion (AICc) was used to

compare multilevel mixed effects models and to select the model of best fit within each data

subset; differences in AICc greater than two were considered meaningful (Burnham and

Anderson, 2004).

Both the database and the R code file showing all analyses are publicly accessible from the

Dryad Digital Repository.

Results

Random Effects Models:

Each data subset (Fig 1) had an I^2 statistic of greater than 75%, indicating high heterogeneity

(Higgins et al., 2003). Random effects model results showed that predators consumptively

decreased mosquito survival with an effect size of -1.23 (95% CI -1.43, -1.03), p-value

<0.0001, and non-consumptively reduced survival with a smaller effect size of -0.11 (95% CI

-0.17, -0.04), p-value = 0.0016. In addition, predators non-consumptively reduced oviposition

behavior with an effect size of -0.87 (95% CI -1.31, -0.42), p-value = 0.0001, and mosquito

body size was non-consumptively reduced by predators in both males and females; the

female effect size was -0.13 (95% CI -0.19, -0.06), p-value = 0.0002, and the male effect size

was -0.03 (95% CI -0.06, -0.01), p-value = 0.0184. There was not a significant non-

consumptive effect of predators on either male or female development time; the female effect

size was -0.01 (95% CI -0.09, 0.07), p-value = 0.7901, and the male effect size was -0.04

(95% CI -0.12, 0.04), p-value = 0.3273.

The Egger's regression test results showed that the non-consumptive survival subset, both

development time subsets (male and female), and the female size subset exhibited funnel plot

asymmetry indicative of publication bias. The "trim and fill" procedure identified missing

studies in the non-consumptive survival subset and the female size subset, but the procedure

did not identify any missing studies in either of the development time subsets. Three studies

were estimated to be missing from the non-consumptive survival data, and accounting for

imputed values from missing studies resulted in a shift in the predation effect size from -0.11

(95% CI -0.17, -0.04), p-value = 0.0016, to -0.13 (95% CI -0.20, -0.07), p-value <0.0001.

Two studies were estimated to be missing from the female size data, and accounting for

imputed values from these missing studies shifted the predation effect size from -0.13 (95%

CI -0.19, -0.06), p-value = 0.0002, to -0.10 (95% CI -0.17, -0.03), p-value = 0.0083. Shifts in

effect size estimates due to the trim and fill procedure were minor and did not cause any of

the observed effects of predators to change direction or become insignificant.

Table 2: Candidate multilevel mixed effects models of consumptive effects from predators on
mosquito survival, fitted to dataset of effect sizes (n = 187 from 34 studies), and ranked by
corrected Akaike's information criterion (AICc).
309

Moderator(s)	Test of Moderators (degrees of freedom, p-value)	AICc	ΔAICc
predator family x vector genus	28, <0.0001	500.5	0
predator family	19, <0.0001	507.0	6.5
predator family + vector genus	23, <0.0001	508.1	7.6
vector genus	5, <0.0001	573.0	72.5
none	----	576.5	76.0

Fig 3. Effect sizes and 95% confidence intervals for consumptive effects of predators, for different categories of moderators (with number of studies in parentheses): a.) predator family with predator class in the right-hand column, b.) vector genus, and c.) larval instar

Table 3: Candidate multilevel mixed effects models of consumptive effects from predators, fitted to dataset of effect sizes where larval instar is not missing (n = 163 from 30 studies), and ranked by corrected Akaike's information criterion (AICc).

Moderator(s)	Test of Moderators (degrees of freedom, p-value)	AICc	ΔAICc
predator family x larval instar	25, <0.0001	429.2	0
predator family + larval instar	19, <0.0001	443.5	14.3
predator family x vector genus	25, <0.0001	455.0	25.8
predator family	17, <0.0001	456.8	27.6
predator family + vector genus	21, <0.0001	458.4	29.2
larval instar	3, <0.0001	503.1	73.9
vector genus	5, <0.0001	504.7	75.5
none	----	508.5	79.3

Multilevel Mixed Effects Models:

The consumptive survival and oviposition data subsets met the criteria of high heterogeneity, observations from at least 10 studies, and no evidence of publication bias. Therefore, these data subsets were tested for moderators using multilevel mixed effects models. Predator families that decreased mosquito survival included Cyprinidae: -3.44 (95% CI -5.79, -1.09), p-value = 0.0042; Poeciliidae: -1.42 (95% CI -2.67, -0.16), p-value = 0.0270; Ambystomatidae: -5.18 (95% CI -7.94, -2.42), p-value = 0.0002; Aeshnidae: -2.93 (95% CI -4.80, -1.07), p-value = 0.0020; and Notonectidae: -2.14 (95% CI -3.07, -1.21), p-value <0.0001 (Fig 3a). Vector genera that experienced significant decreases in survival due to consumptive effects of predators included *Aedes*: -1.23 (95% CI -1.81, -0.65), p-value <0.0001; *Anopheles*: -1.34 (95% CI -2.01, -0.66), p-value = 0.0001; and *Culex*: -1.41 (95% CI -1.96, -0.86), p-value <0.0001 (Fig 3b). Among all 187 consumptive survival observations from 34 studies, the best model fit, according to AICc value, was achieved when an interaction between predator family and vector genus was included in the model (Table 2). However, among the 163 larval stage consumptive survival observations from 30 studies, adding an interactive term between larval instar (an indicator of prey size) and predator family had a greater improvement on model fit than adding an interactive term between

vector genus and predator family (Fig 3c, Table 3). Temperature did not affect the

heterogeneity of consumptive survival data, either as a linear moderator: -0.01 (95% CI -0.10,

0.07), p-value = 0.7559, or a quadratic moderator: 0.00 (95% CI 0.00, 0.00), p-value =

0.8184. The best oviposition model fit, according to AICc value, was achieved when vector

genus was added as a moderator (Table 4). The mean oviposition effect size was not

significantly different than zero for *Aedes*: 0.32 (95% CI -2.14, 2.79), p-value = 0.7970, or

Culiseta: -0.61 (95% CI -1.83, 0.62), p-value = 0.3329, but for *Culex* mosquitoes, oviposition

was significantly decreased by predator presence: -1.69 (95% CI -2.82, -0.56), p-value =

0.0033 (Fig 4).

Table 4: Candidate multilevel mixed effects models of non-consumptive effects of predators
on mosquito oviposition behavior, fitted to dataset of effect sizes (n = 36 from 12 studies),
and ranked by corrected Akaike's information criterion (AICc).
392

Moderator(s)	Test of Moderators (degrees of freedom, p-value)	AICc	ΔAICc
vector genus	3, 0.0149	122.1	0
none	----	125.2	3.1
predator family	12, 0.8855	167.9	45.8

402   **Fig 4.** Oviposition effect sizes and 95% confidence intervals for different categories of vector
403   genus (with number of studies in parentheses)

Discussion

In this study, laboratory and semi-field empirical data were obtained through a systematic literature review and used to conduct a meta-analysis that assessed consumptive and non-consumptive effects of predators on mosquito prey. Some results agree with previously observed trends, such as greater consumptive effects from larger predators (Kumar et al., 2008, Peters, 1983) and no oviposition response to predator cues among container-breeding *Aedes* mosquitoes (Vonesh and Blaustein, 2010). However, this meta-analysis revealed additional trends. Mosquito larval instar had an important role in moderating consumptive effects of predators, likely because of its direct correlation to prey size. Furthermore, a small, but significant, decrease in mosquito survival due to non-consumptive effects of predators was observed, suggesting that mosquitoes can be "scared to death" by predators (Preisser et al., 2005). Both male and female body sizes were also reduced among mosquitoes that had been exposed to predators, and predator avoidance during oviposition was observed among female *Culex* mosquitoes. Effects of predators on different vector traits, particularly survival, body size, and oviposition behavior, have the potential to influence infectious disease dynamics.

Consumptive effects of predators on survival:
Several larger predators reduced mosquito survival, including freshwater fish (Cyprinidae and Poeciliidae), salamander larvae (Ambystomatidae), dragonfly larvae (Aeshnidae), and backswimmers (Notonectidae) (Fig 3a). This finding is consistent with a previous analysis which showed a positive linear relationship between predator body mass and ingestion rate across taxa (Peters, 1983). In addition, more effect size heterogeneity in the consumptive survival data was explained by an interaction between predator family and larval instar than

was explained by an interaction between predator family and vector genus (Table 3). This

result suggests that the relative sizes of predator and prey groups could play a more important

role in determining consumptive mosquito survival than variations in predator responses to

different behaviors of prey genera, which are likely to be shaped by the degree of shared

evolutionary history between trophic levels (Buchanan et al., 2017). Larval instar is an

indicator of mosquito size, and previous modelling work has provided evidence of prey size

selection by predators to maximize energetic gain (Mittelbach, 1981). While smaller

cyclopoid copepods are more effective against early instar mosquito larvae (Dieng et al.,

2002), larger predators including tadpoles, giant water bugs, dragonfly larvae, fish, and

backswimmers are more effective against late instar larvae (Kweka et al., 2011).

Non-consumptive effects of predators on survival:

Exposure to predation cues significantly lowered mosquito survival, and this non-

consumptive effect has also been observed in dragonfly larvae prey (*Leucorrhinia intacta*)

that were exposed to caged predators (McCauley et al., 2011). The reduction in mosquito

survival from non-consumptive effects of predators was significantly smaller than the

reduction that was observed from consumptive effects. This is partially due to the practical

constraints of most experimental designs, which cause consumptive and non-consumptive

effects of predators on survival to be grouped together and reported as consumptive effects.

The greater impact of combined consumptive and non-consumptive effects, in comparison to

only non-consumptive effects, has previously been observed in pea aphids (*Acyrthosiphon*

pisum) (Nelson et al., 2004).

Non-consumptive effects of predators on body size:

While predators did not significantly impact mosquito development time through non-

consumptive effects in either sex, mosquito body size was decreased by the non-consumptive

effects of predators in both sexes. Smaller body size is associated with lower reproductive

success in mosquitoes because smaller females lay fewer eggs (Blackmore and Lord, 2000,

Lyimo and Takken, 1993, Oliver and Howard, 2011, Styer et al., 2007, Tsunoda et al., 2010),

and smaller males produce less sperm (Hatala et al., 2018, Ponlawat and Harrington, 2007).

These effects suggest that predation could non-consumptively reduce mosquito population

growth. The smaller size of mosquitoes exposed to predators could also limit disease

transmission. Vector lifespan contributes disproportionately to disease transmission because

older vectors are more likely to have been exposed to pathogens, more likely to already be

infectious after having survived the extrinsic incubation period, and more likely to survive

long enough to bite subsequent hosts (Cator et al., 2020). It is well-established that smaller

mosquito body size is associated with shorter mosquito lifespan (Araújo et al., 2012, Hawley,

1985, Reisen et al., 1984, Reiskind and Lounibos, 2009, Xue et al., 2010). Therefore, non-

consumptive effects of predators may limit the transmission of mosquito-borne diseases.

Non-consumptive effects of predators on oviposition behavior:

Predator presence also non-consumptively reduced oviposition behavior in adult female

mosquitoes. Meta-regression results showed that *Culex* females significantly avoid

oviposition sites that contain predators or predator cues, but *Aedes* and *Culiseta* females do

not avoid these sites, despite a slight non-significant trend towards predator avoidance in

Culiseta (Fig 4). Both *Culex* and *Culiseta* mosquitoes have an "all-or-none" oviposition

strategy (Johnson and Fonseca, 2014), in which they lay hundreds of rapidly-hatching eggs in

rafts on the water's surface (Day, 2016). Such an oviposition strategy is conducive to

evolving predator avoidance behaviors, and a previous meta-analysis showed significant

predator avoidance in both *Culex* and *Culiseta* during oviposition (Vonesh and Blaustein,

2010). Conversely, it is likely that an oviposition response to predation is not particularly

advantageous for *Aedes* because the delayed hatching of their eggs (Day, 2016) can prevent

the level of predation risk at the time of oviposition from matching the level of predation risk

present in the eventual larval environment (Vonesh and Blaustein, 2010). The predator

avoidance response in *Aedes* species that lay their eggs above the water's edge in containers

has previously been described as "non-existent" (Vonesh and Blaustein, 2010). Both *Aedes*

species included in this study's oviposition data subset, *Ae. albopictus* and *Ae. aegypti*, meet

the criterion of ovipositing above water in containers (Juliano, 2009). Predator avoidance

during oviposition has previously been found to increase the mosquito population size at

equilibrium (Spencer et al., 2002). However, this study's results and those of a previous

meta-analysis (Vonesh and Blaustein, 2010) suggest that models of oviposition site selection,

such as those using parameters from Notonectidae predators and *Culiseta* prey (Kershenbaum

et al., 2012), are not generalizable to *Aedes* vectors.

Implications for West Nile virus disease dynamics:

Predator avoidance during oviposition by *Culex* mosquitoes (Fig 4) may be of particular

importance to West Nile virus (WNV) disease dynamics. Previous work has shown that *Cx.*

pipiens, *Cx. restuans*, and *Cx. tarsalis* all avoid predator habitats (Vonesh and Blaustein,

2010), and that *Cx. pipiens* is the primary bridge vector of WNV responsible for spill-over

transmission from avian reservoir hosts to humans (Fonseca et al., 2004, Hamer et al., 2008a,

Kramer et al., 2008, Andreadis, 2012). *Cx. pipiens* mosquitoes can live in permanent aquatic

environments, such as ground pools (Amini et al., 2020, Barr, 1967, Dida et al., 2018,

Sulesco et al., 2015), ponds (Lühken et al., 2015), stream edges (Amini et al., 2020), and lake

edges (Vinogradova, 2000) that are more common in rural areas, but *Cx. pipiens* are also

found in urban and suburban residential areas, where they typically breed in artificial

containers (Sulesco et al., 2015), including tires (Lühken et al., 2015, Nikookar et al., 2017,

Verna, 2015), rainwater tanks (Townroe and Callaghan, 2014), and catch basins (Gardner et

al., 2012). Small artificial containers, such as discarded tires, are generally unlikely to harbor

larger predators, including freshwater fish (Cyprinidae and Poeciliidae), salamander larvae

(Ambystomatidae), dragonfly larvae (Aeshnidae), and backswimmers (Notonectidae),

because temporary aquatic environments cannot support the relatively long development

times of these organisms. The mean dispersal distance of adult *Culex* mosquitoes is greater

than one kilometer (Ciota et al., 2012, Hamer et al., 2014), and female *Cx. pipiens* have

exhibited longer dispersal distances after developing in the presence of a fish predator

(Alcalay et al., 2018). Therefore, predator avoidance during oviposition may cause *Cx.*

pipiens populations to disperse from permanent aquatic environments in more rural areas to

artificial container environments in urbanized areas, where the risk of human WNV infection

is higher (Brown et al., 2008).

Predator cue levels may be altered by climate conditions, and these changes in cue levels can

impact WNV transmission to humans. Drought has previously been associated with human

WNV cases (Johnson and Sukhdeo, 2013, Marcantonio et al., 2015, Roehr, 2012, Shaman et

al., 2005, Epstein and Defilippo, 2001, Paull et al., 2017), but the association has thus far

lacked a clear underlying mechanism. Under drought conditions, the density of aquatic

organisms increases and predation pressures can intensify due to compressed space and high

encounter rates (Amundrud et al., 2019). A previous study of a stream ecosystem found that

impacts of fish predation are more severe during the dry season (Dudgeon, 1993). In addition,

reductions in water volume can facilitate consumption of mosquito larvae by crane fly larvae

(Tipulidae), whereas mosquito consumption by tipulids was not observed at a higher water

level (Amundrud et al., 2019). Laboratory and semi-field studies have shown that mosquitoes

respond to a gradient of predator cues (Roux et al., 2014, Silberbush and Blaustein, 2011).

The frequency of larval anti-predator behavior is correlated with the concentration of predator

cues (Roux et al., 2014), and adult female mosquitoes prefer oviposition sites with lower

predator densities (Silberbush and Blaustein, 2011). Therefore, as predator cue levels increase

due to drought, permanent aquatic habitats are likely to transition from suitable oviposition

sites for one generation of female mosquitoes, to unsuitable oviposition sites for the next

generation.

When suitable oviposition sites are absent, females retain their eggs until sites become

available (Bentley and Day, 1989). *Cx. pipiens* females can retain their eggs for up to five

weeks, allowing them enough time to find container sites with low predation risk, often

located in residential areas (Johnson and Fonseca, 2014). The movement of gravid female *Cx.*

pipiens to residential areas increases the risk of WNV spill-over to humans because these

vectors are likely to have already blood-fed at least once (Clements, 1992), suggesting that

they have a higher risk of WNV infection, relative to non-gravid mosquitoes. This is

consistent with studies that have reported associations between drought and WNV-infected

mosquitoes in urban and residential areas (Johnson and Sukhdeo, 2013, Paull et al., 2017). In

addition, vertical transmission of WNV from gravid females to their progeny may occur

during oviposition (Rosen, 1988), when the virus is transmitted by an accessory gland fluid

that attaches eggs to one another (Nelms et al., 2013). Because the rate of vertical

transmission in *Cx. pipiens* increases with the number of days following WNV infection

(Anderson et al., 2008), extended searches for oviposition sites due to drought could increase

the frequency of vertical transmission. However, the impact of vertical transmission on WNV

epidemics is thought to be minimal because when transmission to an egg raft did occur, only

4.7% of the progeny were found to be infected as adults (Anderson et al., 2008), and only

about half of those infected adults are estimated to be female. In summary, the movement of

Cx. pipiens females toward more residential areas, combined with potential limited WNV

amplification from increased vertical transmission, suggests that the vector trait of predator

avoidance during oviposition can serve as a plausible explanation for associations between

drought and human WNV cases.

Another theory for the association between drought and human WNV cases is based on the

hypothesis that increased contact between mosquito vectors and passerine reservoir hosts

occurs during drought conditions (Paull et al., 2017, Shaman et al., 2005). The proposed

aggregation of bird and mosquito populations during drought was originally thought to occur

in humid, densely vegetated hammocks – a type of habitat that is specific to southern Florida

(Shaman et al., 2005), but WNV incidence is more consistently clustered in other regions of

the US, particularly the Northern Great Plains (Centers for Disease Control and Prevention

(CDC), 2021, Sugumaran et al., 2009). Northern cardinals (*Cardinalis cardinalis*), American

robins (*Turdus migratorius*), and house sparrows (*Passer domesticus*) were among the bird

species that most frequently tested seropositive for WNV antibodies in 2005 and 2006 in

Chicago, where high numbers of human cases were reported (Hamer et al., 2008b), and these

passerine species are more abundant in residential areas, regardless of precipitation patterns

(Anderson, 2006b, Beddall, 1963, Lepczyk et al., 2008). Apart from drought, landowners'

participation in supplemental bird feeding, providing bird houses, gardening, and maintaining

vegetation can strongly influence passerine abundance in residential areas (Lepczyk et al.,

2004). Furthermore, as terrestrial foragers that can obtain hydration from their diet of insects,

fruits, and other plant material (Anderson, 2006a, Brzek et al., 2009, Malmborg and Willson,

1988, Renne et al., 2000), passerine reservoir hosts of WNV are less likely to move in

response to drought than the mosquito vectors of WNV, which have obligate aquatic life

stages.

While hatch-year birds are more vulnerable to mosquito biting, and thus contribute to the

amplification of WNV (Hamer et al., 2008b), it is illogical to expect an increased abundance

of hatch-year birds during drought conditions. However, some have argued that in cases

where drought decreases the abundance of juvenile birds, the ratio of mosquitoes to birds

increases, and this could lead to higher WNV prevalence in the mosquito population (Paull et

al., 2017). Although reductions in both hatching success (George et al., 1992) and survival of

recently fledged birds (Adams et al., 2006) have been observed during drought conditions,

the impact of drought on avian abundance varies widely by species (Verner and Purcell,

1999). In particular, synanthropic species, such as those likely to harbor WNV, are less

negatively affected by drought (Albright et al., 2010). Additionally, the droughts that impact

avian abundance often occur over much longer periods of time than the seasonal droughts

that predict WNV transmission to humans. For example, avian abundance has been modeled

based on precipitation metrics spanning 32 weeks, and house wren (*Troglodytes aedon*)

abundance has been predicted by precipitation averages spanning four years (Verner and

Purcell, 1999). Finally, birds with higher levels of stress hormones are more likely to be fed

on by mosquitoes, and certain factors associated with residential areas, such as road noise,

light pollution, and pesticide exposure, can cause avian stress (Gervasi et al., 2016).

Therefore, elevated avian stress hormones in these habitats may contribute to WNV

prevalence in the mosquito population, independent of drought conditions.

Implications for mosquito-borne disease modelling:

Although the aquatic phase of the mosquito life cycle is often overlooked in mathematical

models of mosquito-borne pathogen transmission (Reiner et al., 2013), vector survival at

immature stages plays an important role in determining mosquito population abundance,

which is an essential factor for predicting disease transmission (Beck-Johnson et al., 2013).

The results of this study show that mosquito survival decreases among the *Aedes*, *Anopheles*,

and *Culex* genera due to consumptive effects of predators (Fig 3b), and that there is also a

reduction in mosquito survival due to non-consumptive effects. Other studies have

demonstrated that aquatic predators dramatically impact mosquito survival and abundance.

For example, a biocontrol intervention relying on the application of copepod predators

eliminated *Aedes albopictus* from three communes in Nam Dinh, Vietnam, where dengue

transmission was previously detected, and reduced vector abundance by 86-98% in three

other communes (Kay et al., 2002). Conversely, the annual abundance of *Culex* and

Anopheles mosquitoes was observed to increase 15-fold in semi-permanent wetlands in the

year following a drought, likely because the drought eliminated aquatic predators from

wetlands that dried completely, and mosquitoes were able to re-colonize newly-formed

aquatic habitats more quickly than their most effective predators (Chase and Knight, 2003).

While relationships between temperature and different vector traits, such as fecundity and

lifespan, have been incorporated into models of temperature effects on mosquito population

density (El Moustaid and Johnson, 2019), models of predator effects on vector borne disease

transmission have focused primarily on the impacts of predation on vector survival. Previous

models have shown that predators of vector species can decrease or eliminate pathogen

infection in host populations as vector fecundity increases (Moore et al., 2010). The findings

of this meta-analysis suggest that predators also decrease vector fecundity through non-

consumptive effects on vector body size. In addition, the entomological inoculation rate

(EIR) is likely to be reduced by effects of predators on mosquito fecundity and lifespan, as

well as effects of predators on mosquito survival. The EIR has been defined as the product of

three variables: (m) the number of mosquitoes per host, (a) the daily rate of mosquito biting,

and (s) the proportion of mosquitoes that are infectious (Beck-Johnson et al., 2013). Based on

this study's findings, predators are likely to decrease the number of mosquitoes per host by

reducing mosquito survival through both consumptive and non-consumptive effects, and by

reducing mosquito fecundity through non-consumptive effects on body size. In addition,

predators are likely to decrease the proportion of mosquitoes that are infectious by shortening

the vector lifespan through non-consumptive effects on body size. The relationship between

mosquito body size and biting rate is unclear, with some studies showing higher biting rates

among larger mosquitoes (Araújo et al., 2012, Gunathilaka et al., 2019), and others reporting

higher biting rates among smaller mosquitoes (Farjana and Tuno, 2013, Leisnham et al.,

2008). The links between factors that influence the EIR and observed effects of predators on

mosquito prey demonstrate the necessity of including both consumptive and non-

consumptive effects of predators in models of mosquito-borne disease.

Conclusion

This meta-analysis on mosquito predation demonstrates that predators not only play an

important role in directly reducing mosquito populations, but also have non-consumptive

effects on surviving mosquitoes that may ultimately reduce further population growth and

decrease disease transmission. While families of larger sized predators were effective in

reducing mosquito survival, other factors, such as impacts on native species, as well as the

economic cost of mass-rearing and field applications (Kumar and Hwang, 2006, Pyke, 2008),

should be carefully considered before selecting a predator as a suitable biocontrol agent.

Predictive disease models are likely to be more reliable when the non-consumptive effects of

predation are incorporated. Although exposure of mosquito larvae to predators is

commonplace in outdoor field settings, it remains rare in most laboratory-based assessments

of vector traits. Therefore, mosquitoes observed in nature are likely to have smaller body

sizes than those observed under optimal laboratory conditions. It is important for disease

modelers to recognize these impacts of predation on vector traits as they can reduce mosquito

population growth and limit disease transmission due to shorter vector lifespans. Within the

WNV disease system, consideration of the oviposition behavioral response to predation cues

by *Culex* vectors can improve current understanding of the association between drought and

human cases. This study provides general estimates of the effects of predators on selected

mosquito traits for use in predictive disease models.

Future directions:

Modelling efforts that aim to optimize the application of biocontrol predators should also

consider incorporating predator effects on vector survival, fecundity, and lifespan. These

additions to predictive models of various biocontrol interventions are likely to help public

health officials choose the most cost-effective strategies for limiting disease transmission. In

the sixty-study database that was compiled, only one study was designed to directly measure

the effect of larval-stage predation on vector competence (Roux et al., 2015). Therefore,

future efforts to assess the impact of predators on mosquito-borne disease transmission

should prioritize experimental studies in which infected mosquito larvae are observed

throughout an initial period of aquatic exposure to predators, followed by a period of blood-

feeding in the adult stage.

Two studies from the compiled database examined the compatibility of predators with

Bacillus thuringiensis var. *israelensis* (*Bti*), a commonly used bacterial biocontrol agent

(Chansang et al., 2004, Op de Beeck et al., 2016). Previous studies have supported the

simultaneous application of cyclopoid copepod predators and *Bti* (Marten et al., 1993, Tietze

et al., 1994), but additional analyses are needed on the use of *Bti* with other families of

mosquito predators. Populations of other insect pests, such as the southern green stink bug

(*Nezara viridula*), are known to be regulated by both predators and parasites (Ehler, 2002).

The literature search conducted for this meta-analysis returned studies on water mite parasites

(Rajendran and Prasad, 1994) and nematode parasitoids (de Valdez, 2006) of mosquitoes, and

ascogregarine parasites have previously been evaluated as biocontrol agents against *Aedes*

mosquitoes (Tseng, 2007). A more thorough review of the impacts of parasites and

parasitoids on vector traits, such as survival, fecundity, and lifespan, is needed before

incorporating these potential biocontrol agents into integrated vector control plans.

Three studies in the sixty-study database included experiments where two mosquito prey

species were made available to the predator species (Grill and Juliano, 1996, Griswold and

Lounibos, 2005b, Micieli et al., 2002). In these cases, the effect size measurement for each

mosquito species could be influenced by interspecific competition, or a preference of the

predator species for a certain prey species. Hetero-specific prey observations were excluded

from this meta-analysis, but future analyses centered on the concepts of interspecific

competition or predator preferences might further evaluate these data. In addition, this meta-

analysis investigated consumptive and non-consumptive effects of predators separately. More

research is needed to determine how models should combine these different types of predator

effects to accurately reflect predation interactions as they occur in natural environments.

Author Contributions

Conceptualization, all authors; data curation, MCR, CMH, ZG, CR, FEM, MVE, TD, NLG, SLH, ACM; formal analysis, investigation, methodology, and validation MCR, CMH, ZG, CR, MVE, ACM; project administration, MCR, CMH, ACM; software, MCR, CMH, MVE, ZG, ACM; supervision CMH, NLG, SLH, AGP, ACM; visualization, MCR, CMH, TD, ACM; writing— original draft, MCR; writing—review and editing, all authors. None of the authors have any competing interests to declare.

Acknowledgements

We are grateful to Dr. Lauren Cator for facilitating the early stages of this project and helping to edit this manuscript. We also thank Dr. Peter Hudson for his helpful advice in early discussions about the project's aims. In addition, we thank all members of the VectorBiTE RCN (https://vectorbite.org/) for taking the initiative to forge productive collaborations between empiricists and modelers in vector ecology. This work was funded by NIH grant 1R01AI122284-01 and BBSRC grant BB/N013573/1 as part of the joint (NIH-NSF-USDA-BBSRC) Ecology and Evolution of Infectious Diseases program. It was also funded by a President's PhD Scholarship from Imperial College London awarded to Marie C. Russell.

Supplementary Information

727   Table S1: Publications included in the database and their types of vector trait data
728

Publication	Vector Trait			
	Survival	Behavior	Size	Development
(Alcalay et al., 2018)	☒	☐	☒	☒
(Alcalay et al., 2019)	☐	☒	☐	☐
(Arav and Blaustein, 2006)	☐	☒	☐	☐
(Beketov and Liess, 2007)	☒	☐	☒	☒
(Blaustein et al., 2005)	☐	☒	☐	☐
(Blaustein and Margalit, 1996)	☒	☐	☐	☐
(Bond et al., 2005)	☐	☐	☒	☒
(Bowatte et al., 2013)	☒	☒	☐	☐
(Bucciarelli et al., 2019)	☒	☐	☐	☐
(Calliari et al., 2003)	☒	☐	☐	☐
(Chandrasegaran et al., 2018)	☐	☒	☐	☐
(Chansang et al., 2004)	☒	☐	☐	☐
(Cuthbert et al., 2018)	☒	☐	☐	☐
(Cuthbert et al., 2019a)	☒	☐	☐	☐
(Cuthbert et al., 2019b)	☐	☒	☐	☐
(de Valdez, 2007)	☐	☒	☐	☐
(Dieng et al., 2002)	☒	☐	☐	☐
(Dieng et al., 2003)	☒	☐	☐	☐
(Farajollahi et al., 2009)	☒	☐	☐	☐
(Fischer et al., 2012)	☐	☐	☒	☒
(Fontana-Bria et al., 2017)	☐	☐	☒	☒
(Futami et al., 2008)	☒	☐	☐	☐
(Grill and Juliano, 1996)	☒	☒	☒	☒
(Griswold and Lounibos, 2005a)	☒	☐	☐	☐
(Irwin and Paskewitz, 2009)	☒	☐	☐	☐
(Kalimuthu et al., 2017)	☒	☐	☐	☐
(Knight et al., 2004)	☒	☐	☐	☒
(Kraus and Vonesh, 2010)	☐	☒	☐	☐
(Krol et al., 2019)	☒	☐	☐	☒
(Kweka et al., 2011)	☒	☐	☐	☐
(Marten et al., 2000a)	☒	☐	☐	☐
(Marten et al., 2000b)	☒	☐	☐	☐
(Mercer et al., 2005)	☒	☐	☒	☐
(Micieli et al., 2002)	☒	☐	☐	☐
(Minakawa et al., 2007)	☒	☐	☐	☐
(Nannini and Juliano, 1998)	☒	☐	☐	☐
(Ohba et al., 2012)	☒	☒	☒	☐
(Op de Beeck et al., 2016)	☐	☐	☒	☒
(Orr and Resh, 1989)	☒	☐	☐	☐
(Pintar et al., 2018)	☐	☒	☐	☐
(Roberts, 2012)	☐	☒	☒	☒
(Roberts, 2018)	☐	☐	☒	☒
(Roux et al., 2015)	☒	☐	☒	☒

Publication (continued)	Vector Trait (continued)			
	Survival	Behavior	Size	Development
(Rubbo et al., 2011)	☒	☒	☒	☐
(Schrama et al., 2018)	☒	☐	☐	☐
(Silberbush et al., 2015)	☒	☐	☐	☒
(Silberbush and Blaustein, 2008)	☐	☒	☐	☐
(Silberbush and Blaustein, 2011)	☐	☒	☐	☐
(Soghigian et al., 2017)	☒	☒	☐	☐
(Staats et al., 2016)	☐	☒	☐	☐
(Stav et al., 2005)	☒	☐	☒	☒
(Subramaniam et al., 2015)	☒	☐	☐	☐
(Tietze and Mulla, 1990)	☒	☐	☐	☒
(van Uitregt et al., 2012)	☒	☒	☒	☒
(van Uitregt et al., 2013)	☐	☒	☒	☐
(Veronesi et al., 2015)	☒	☐	☐	☐
(Wallace and Merritt, 1999)	☒	☐	☒	☒
(Yanoviak, 1999)	☒	☐	☒	☐
(Zuharah et al., 2013)	☒	☐	☐	☐
(Zuharah and Lester, 2010)	☒	☐	☐	☐

References

ADAMS, A. A. Y., SKAGEN, S. K. & SAVIDGE, J. A. 2006. Modeling post-fledging survival of Lark Buntings in response to ecological and biological factors. *Ecology,* 87**,** 178-188.

ALBRIGHT, T. P., PIDGEON, A. M., RITTENHOUSE, C. D., CLAYTON, M. K., FLATHER, C. H., CULBERT, P. D., WARDLOW, B. D. & RADELOFF, V. C. 2010. Effects of drought on avian community structure. *Global Change Biology,* 16**,** 2158-2170.

ALCALAY, Y., TSURIM, I. & OVADIA, O. 2018. Female mosquitoes disperse further when they develop under predation risk. *Behavioral Ecology,* 29**,** 1402-1408.

ALCALAY, Y., TSURIM, I. & OVADIA, O. 2019. Multi-scale oviposition site selection in two mosquito species. *Ecological Entomology,* 44**,** 347-356.

AMINI, M., HANAFI-BOJD, A. A., AGHAPOUR, A. A. & CHAVSHIN, A. R. 2020. Larval habitats and species diversity of mosquitoes (Diptera: Culicidae) in West Azerbaijan Province, Northwestern Iran. *Bmc Ecology,* 20.

AMUNDRUD, S. L., CLAY-SMITH, S. A., FLYNN, B. L., HIGGINS, K. E., REICH, M. S., WIENS, D. R. H. & SRIVASTAVA, D. S. 2019. Drought alters the trophic role of an opportunistic generalist in an aquatic ecosystem. *Oecologia,* 189**,** 733-744.

ANDERSON, J. F., MAIN, A. J., DELROUX, K. & FIKRIG, E. 2008. Extrinsic incubation periods for horizontal and vertical transmission of West Nile virus by Culex pipiens pipiens (Diptera: Culicidae). *J Med Entomol,* 45**,** 445-51.

ANDERSON, T. R. 2006a. Chapter 6: Foraging behavior and food. *Biology of the Ubiquitous House Sparrow: From Genes to Populations.* New York: Oxford University Press.

ANDERSON, T. R. 2006b. Chapter 10: Human commensalism and pest management. *Biology of the Ubiquitous House Sparrow: From Genes to Populations.* New York: Oxford University Press.

ANDREADIS, T. G. 2012. THE CONTRIBUTION OF CULEX PIPIENS COMPLEX MOSQUITOES TO TRANSMISSION AND PERSISTENCE OF WEST NILE VIRUS IN NORTH AMERICA. *Journal of the American Mosquito Control Association,* 28**,** 137-151.

ARAÚJO, M. D., GIL, L. H. S. & E-SILVA, A. D. 2012. Larval food quantity affects development time, survival and adult biological traits that influence the vectorial capacity of Anopheles darlingi under laboratory conditions. *Malaria Journal,* 11.

ARAV, D. & BLAUSTEIN, L. 2006. Effects of pool depth and risk of predation on oviposition habitat selection by temporary pool dipterans. *Journal of Medical Entomology,* 43**,** 493-497.

BARR, A. R. 1967. Occurrence and distribution of the Culex pipiens complex. *Bulletin of the World Health Organization,* 37**,** 293-296.

BECK-JOHNSON, L. M., NELSON, W. A., PAAIJMANS, K. P., READ, A. F., THOMAS, M. B. & BJØRNSTAD, O. N. 2013. The effect of temperature on Anopheles mosquito population dynamics and the potential for malaria transmission. *PLoS One,* 8**,** e79276.

BEDDALL, B. G. 1963. Range expansion of the cardinal and other birds in the northeastern states. *The Wilson Bulletin,* 75**,** 140-158.

BEKETOV, M. A. & LIESS, M. 2007. Predation risk perception and food scarcity induce alterations of life-cycle traits of the mosquito Culex pipens. *Ecological Entomology,* 32**,** 405-410.

BENELLI, G., JEFFRIES, C. L. & WALKER, T. 2016. Biological Control of Mosquito Vectors: Past, Present, and Future. *Insects,* 7.

BENTLEY, M. D. & DAY, J. F. 1989. Chemical ecology and behavioral aspects of mosquito oviposition. *Annu Rev Entomol,* 34**,** 401-21.

BLACKMORE, M. S. & LORD, C. C. 2000. The relationship between size and fecundity in Aedes albopictus. *Journal of Vector Ecology,* 25**,** 212-217.

BLAUSTEIN, L., BLAUSTEIN, J. & CHASE, J. 2005. Chemical detection of the predator Notonecta irrorata by ovipositing Culex mosquitoes. *Journal of Vector Ecology,* 30**,** 299-301.

BLAUSTEIN, L. & MARGALIT, J. 1996. Priority Effects in Temporary Pools: Nature and Outcome of Mosquito Larva-Toad Tadpole Interactions Depend on Order of Entrance. *Journal of Animal Ecology,* 65**,** 77-84.

BOND, J. G., ARREDONDO-JIMENEZ, J. I., RODRIGUEZ, M. H., QUIROZ-MARTINEZ, H. & WILLIAMS, T. 2005. Oviposition habitat selection for a predator refuge and food source in a mosquito. *Ecological Entomology,* 30**,** 255-263.

BOWATTE, G., PERERA, P., SENEVIRATHNE, G., MEEGASKUMBURA, S. & MEEGASKUMBURA, M. 2013. Tadpoles as dengue mosquito (Aedes aegypti) egg predators. *Biological Control,* 67**,** 469-474.

BROSE, U., JONSSON, T., BERLOW, E. L., WARREN, P., BANASEK-RICHTER, C., BERSIER, L. F., BLANCHARD, J. L., BREY, T., CARPENTER, S. R., BLANDENIER, M. F. C., CUSHING, L., DAWAH, H. A., DELL, T., EDWARDS, F., HARPER-SMITH, S., JACOB, U., LEDGER, M. E., MARTINEZ, N. D., MEMMOTT, J., MINTENBECK, K., PINNEGAR, J. K., RALL, B. C., RAYNER, T. S., REUMAN, D. C., RUESS, L., ULRICH, W., WILLIAMS, R. J., WOODWARD, G. & COHEN, J. E. 2006. Consumer-resource body-size relationships in natural food webs. *Ecology,* 87**,** 2411-2417.

BROWN, H. E., CHILDS, J. E., DIUK-WASSER, M. A. & FISH, D. 2008. Ecological
factors associated with West Nile virus transmission, northeastern United States.
Emerg Infect Dis, 14**,** 1539-45.
BRZEK, P., KOHL, K., CAVIEDES-VIDAL, E. & KARASOV, W. H. 2009. Developmental
adjustments of house sparrow (Passer domesticus) nestlings to diet composition.
Journal of Experimental Biology, 212**,** 1284-1293.
BUCCIARELLI, G. M., SUH, D., LAMB, A. D., ROBERTS, D., SHARPTON, D.,
SHAFFER, H. B., FISHER, R. N. & KATS, L. B. 2019. Assessing effects of non-
native crayfish on mosquito survival. *Conservation Biology,* 33**,** 122-131.
BUCHANAN, A. L., HERMANN, S. L., LUND, M. & SZENDREI, Z. 2017. A meta-
analysis of non-consumptive predator effects in arthropods: the influence of
organismal and environmental characteristics. *Oikos,* 126**,** 1233-1240.
BURNHAM, K. P. & ANDERSON, D. R. 2004. Multimodel Inference: Understanding AIC
and BIC in Model Selection. *Sociological Methods & Research,* 33**,** 261-304.
CALLIARI, D., SANZ, K., MARTINEZ, M., CERVETTO, G., GOMEZ, M. & BASSO, C.
2003. Comparison of the predation rate of freshwater cyclopoid copepod species on
larvae of the mosquito Culex pipiens. *Medical and veterinary entomology,* 17**,** 339-
342.
CATOR, L. J., JOHNSON, L. R., MORDECAI, E. A., EL MOUSTAID, F., SMALLWOOD,
T. R. C., LADEAU, S. L., JOHANSSON, M. A., HUDSON, P. J., BOOTS, M.,
THOMAS, M. B., POWER, A. G. & PAWAR, S. 2020. The Role of Vector Trait
Variation in Vector-Borne Disease Dynamics. *Frontiers in Ecology and Evolution,* 8.
CENTERS FOR DISEASE CONTROL AND PREVENTION (CDC). 2021. *West Nile virus:*
Final Cumulative Maps & Data for 1999-2019 [Online]. Available:
https://www.cdc.gov/westnile/statsmaps/cumMapsData.html [Accessed 02/07/2021].
CHANDRASEGARAN, K., SINGH, A., LAHA, M. & QUADER, S. 2018. Playing it safe?
Behavioural responses of mosquito larvae encountering a fish predator. *Ethology*
Ecology & Evolution, 30**,** 70-87.
CHANSANG, U. R., BHUMIRATANA, A. & KITTAYAPONG, P. 2004. Combination of
Mesocyclops thermocyclopoides and Bacillus thuringiensis var. israelensis: A better
approach for the control of Aedes aegypti larvae in water containers. *Journal of*
Vector Ecology, 29**,** 218-226.
CHASE, J. M. & KNIGHT, T. M. 2003. Drought-induced mosquito outbreaks in wetlands.
Ecology Letters, 6**,** 1017-1024.
CIOTA, A. T., DRUMMOND, C. L., RUBY, M. A., DROBNACK, J., EBEL, G. D. &
KRAMER, L. D. 2012. Dispersal of Culex mosquitoes (Diptera: Culicidae) from a
wastewater treatment facility. *J Med Entomol,* 49**,** 35-42.
CLEMENTS, A. N. 1992. *The Biology of Mosquitoes: Development, Nutrition and*
Reproduction, London, Chapman & Hall.
CUTHBERT, R. N., DALU, T., WASSERMAN, R. J., COUGHLAN, N. E., CALLAGHAN,
A., WEYL, O. L. F. & DICK, J. T. A. 2018. Muddy waters: Efficacious predation of
container-breeding mosquitoes by a newly-described calanoid copepod across
differential water clarities. *Biological Control,* 127**,** 25-30.
CUTHBERT, R. N., DALU, T., WASSERMAN, R. J., WEYL, O. L. F., CALLAGHAN, A.,
FRONEMAN, W. & DICK, J. T. A. 2019a. Sex-skewed trophic impacts in ephemeral
wetlands. *Freshwater Biology,* 64**,** 359-366.
CUTHBERT, R. N., ORTIZ-PEREA, N., DICK, J. T. A. & CALLAGHAN, A. 2019b.
Elusive enemies: Consumptive and ovipositional effects on mosquitoes by predatory
midge larvae are enhanced in dyed environments. *Biological Control,* 132**,** 116-121.

DAM, H. G. & PETERSON, W. T. 1988. THE EFFECT OF TEMPERATURE ON THE GUT CLEARANCE RATE-CONSTANT OF PLANKTONIC COPEPODS. *Journal of Experimental Marine Biology and Ecology,* 123**,** 1-14.

DAY, J. F. 2016. Mosquito Oviposition Behavior and Vector Control. *Insects,* 7.

DE VALDEZ, M. R. 2007. Predator avoidance behavior of Aedes aegypti mosquito larvae infected with mermithid nematodes (Nematoda: Mermithidae). *J Vector Ecol,* 32**,** 150-3.

DE VALDEZ, M. R. W. 2006. Parasitoid-induced behavioral alterations of Aedes aegypti mosquito larvae infected with mermithid nematodes (Nematoda : Mermithidae). *Journal of Vector Ecology,* 31**,** 344-354.

DELNAT, V., TRAN, T. T., JANSSENS, L. & STOKS, R. 2019. Resistance to a chemical pesticide increases vulnerability to a biopesticide: Effects on direct mortality and mortality by predation. *Aquat Toxicol,* 216**,** 105310.

DIDA, G. O., ANYONA, D. N., ABUOM, P. O., AKOKO, D., ADOKA, S. O., MATANO, A. S., OWUOR, P. O. & OUMA, C. 2018. Spatial distribution and habitat characterization of mosquito species during the dry season along the Mara River and its tributaries, in Kenya and Tanzania. *Infectious Diseases of Poverty,* 7.

DIENG, H., BOOTS, M., TUNO, N., TSUDA, Y. & TAKAGI, M. 2002. A laboratory and field evaluation of Macrocyclops distinctus, Megacyclops viridis and Mesocyclops pehpeiensis as control agents of the dengue vector Aedes albopictus in a peridomestic area in Nagasaki, Japan. *Medical and Veterinary Entomology,* 16**,** 285-291.

DIENG, H., BOOTS, M., TUNO, N., TSUDA, Y. & TAKAGI, M. 2003. Life history effects of prey choice by copepods: Implications for biocontrol of vector mosquitoes. *Journal of the American Mosquito Control Association,* 19**,** 67-73.

DUDGEON, D. 1993. THE EFFECTS OF SPATE-INDUCED DISTURBANCE, PREDATION AND ENVIRONMENTAL COMPLEXITY ON MACROINVERTEBRATES IN A TROPICAL STREAM. *Freshwater Biology,* 30**,** 189-197.

DUVAL, S. & TWEEDIE, R. 2000a. A nonparametric "trim and fill" method of accounting for publication bias in meta-analysis. *Journal of the American Statistical Association,* 95**,** 89-98.

DUVAL, S. & TWEEDIE, R. 2000b. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics,* 56**,** 455-463.

EHLER, L. E. 2002. An evaluation of some natural enemies of Nezara viridula in northern California. *BioControl,* 47**,** 309-325.

EL MOUSTAID, F. & JOHNSON, L. R. 2019. Modeling Temperature Effects on Population Density of the Dengue Mosquito Aedes aegypti. *Insects,* 10**,** 393.

ENGLUND, G., OHLUND, G., HEIN, C. L. & DIEHL, S. 2011. Temperature dependence of the functional response. *Ecology Letters,* 14**,** 914-921.

EPSTEIN, P. R. & DEFILIPPO, C. 2001. West Nile Virus and Drought. *Global Change and Human Health,* 2**,** 105-107.

FARAJOLLAHI, A., KESAVARAJU, B., NELDER, M. P., CRANS, S. C. & GAUGLER, R. 2009. An unusual larval collection and survival of Orthopodomyia signifera in the presence of the predator Toxorhynchites rutilus septentrionalis. *J Am Mosq Control Assoc,* 25**,** 370-3.

FARJANA, T. & TUNO, N. 2013. Multiple Blood Feeding and Host-Seeking Behavior in Aedes aegypti and Aedes albopictus (Diptera: Culicidae). *Journal of Medical Entomology,* 50**,** 838-846.

FISCHER, S., PEREYRA, D. & FERNÁNDEZ, L. 2012. Predation ability and non-consumptive effects of Notonecta sellata (Heteroptera: Notonectidae) on immature stages of Culex pipiens (Diptera: Culicidae). *Journal of Vector Ecology,* 37**,** 245-251.

FONSECA, D. M., KEYGHOBADI, N., MALCOLM, C. A., MEHMET, C., SCHAFFNER, F., MOGI, M., FLEISCHER, R. C. & WILKERSON, R. C. 2004. Emerging vectors in the Culex pipiens complex. *Science,* 303**,** 1535-1538.

FONTANA-BRIA, L., SELFA, J., TUR, C. & FRAGO, E. 2017. Early exposure to predation risk carries over metamorphosis in two distantly related freshwater insects. *Ecological Entomology,* 42**,** 255-262.

FUTAMI, K., SONYE, G., AKWEYWA, P., KANEKO, S. & MINAKAWA, N. 2008. Diving behavior in Anopheles gambiae (Diptera: Culicidae): avoidance of a predacious wolf spider (Araneae: Lycosidae) in relation to life stage and water depth. *J Med Entomol,* 45**,** 1050-6.

GARDNER, A. M., HAMER, G. L., HINES, A. M., NEWMAN, C. M., WALKER, E. D. & RUIZ, M. O. 2012. Weather variability affects abundance of larval Culex (Diptera: Culicidae) in storm water catch basins in suburban Chicago. *Journal of medical entomology,* 49**,** 270-276.

GEORGE, T. L., FOWLER, A. C., KNIGHT, R. L. & MCEWEN, L. C. 1992. Impacts of a Severe Drought on Grassland Birds in Western North Dakota. *Ecol Appl,* 2**,** 275-284.

GERVASI, S. S., BURKETT-CADENA, N., BURGAN, S. C., SCHREY, A. W., HASSAN, H. K., UNNASCH, T. R. & MARTIN, L. B. 2016. Host stress hormones alter vector feeding preferences, success, and productivity. *Proceedings of the Royal Society B-Biological Sciences,* 283.

GRILL, C. P. & JULIANO, S. A. 1996. Predicting species interactions based on behaviour: Predation and competition in container-dwelling mosquitoes. *Journal of Animal Ecology,* 65**,** 63-76.

GRISWOLD, M. W. & LOUNIBOS, L. P. 2005a. Competitive Outcomes of Aquatic Container Diptera Depend on Predation and Resource Levels. *Annals of the Entomological Society of America,* 98**,** 673-681.

GRISWOLD, M. W. & LOUNIBOS, L. P. 2005b. Does differential predation permit invasive and native mosquito larvae to coexist in Florida? *Ecol Entomol,* 30**,** 122-127.

GUNATHILAKA, H. N., UPILIKA, H., UDAYANGA, N. W. B. A. & AMARASINGHE, D. 2019. Effect of Larval Nutritional Regimes on Morphometry and Vectorial Capacity of Aedes aegypti for Dengue Transmission. *BioMed Research International,* 2019**,** 1-11.

HAMER, G. L., ANDERSON, T. K., DONOVAN, D. J., BRAWN, J. D., KREBS, B. L., GARDNER, A. M., RUIZ, M. O., BROWN, W. M., KITRON, U. D., NEWMAN, C. M., GOLDBERG, T. L. & WALKER, E. D. 2014. Dispersal of adult culex mosquitoes in an urban west nile virus hotspot: a mark-capture study incorporating stable isotope enrichment of natural larval habitats. *PLoS neglected tropical diseases,* 8**,** e2768-e2768.

HAMER, G. L., KITRON, U. D., BRAWN, J. D., LOSS, S. R., RUIZ, M. O., GOLDBERG, T. L. & WALKER, E. D. 2008a. Culex pipiens (Diptera : Culicidae): A bridge vector of West Nile virus to humans. *Journal of Medical Entomology,* 45**,** 125-128.

HAMER, G. L., WALKER, E. D., BRAWN, J. D., LOSS, S. R., RUIZ, M. O., GOLDBERG, T. L., SCHOTTHOEFER, A. M., BROWN, W. M., WHEELER, E. & KITRON, U. D. 2008b. Rapid amplification of West Nile virus: The role of hatch-year birds. *Vector-Borne and Zoonotic Diseases,* 8**,** 57-67.

HANCOCK, P. A., WIEBE, A., GLEAVE, K. A., BHATT, S., CAMERON, E., TRETT, A., WEETMAN, D., SMITH, D. L., HEMINGWAY, J., COLEMAN, M., GETHING, P.

W. & MOYES, C. L. 2018. Associated patterns of insecticide resistance in field populations of malaria vectors across Africa. *Proceedings of the National Academy of Sciences of the United States of America,* 115**,** 5938-5943.

HATALA, A. J., HARRINGTON, L. C. & DEGNER, E. C. 2018. Age and Body Size Influence Sperm Quantity in Male Aedes albopictus (Diptera: Culicidae) Mosquitoes. *Journal of Medical Entomology,* 55**,** 1051-1054.

HAWLEY, W. A. 1985. THE EFFECT OF LARVAL DENSITY ON ADULT LONGEVITY OF A MOSQUITO, AEDES-SIERRENSIS - EPIDEMIOLOGICAL CONSEQUENCES. *Journal of Animal Ecology,* 54**,** 955-964.

HEMINGWAY, J. & RANSON, H. 2000. Insecticide resistance in insect vectors of human disease. *Annual Review of Entomology,* 45**,** 371-391.

HERMANN, S. L. & LANDIS, D. A. 2017. Scaling up our understanding of non-consumptive effects in insect systems. *Current Opinion in Insect Science,* 20**,** 54-60.

HERMANN, S. L. & THALER, J. S. 2014. Prey perception of predation risk: volatile chemical cues mediate non-consumptive effects of a predator on a herbivorous insect. *Oecologia,* 176**,** 669-676.

HIGGINS, J. P. T., THOMAS, J., CHANDLER, J., CUMPSTON, M., LI, T., PAGE, M. J. & WELCH, V. A. 2020. Cochrane Handbook for Systematic Reviews of Interventions. Cochrane.

HIGGINS, J. P. T., THOMPSON, S. G., DEEKS, J. J. & ALTMAN, D. G. 2003. Measuring inconsistency in meta-analyses. *BMJ (Clinical research ed.),* 327**,** 557-560.

HILL, C. A., KAFATOS, F. C., STANSFIELD, S. K. & COLLINS, F. H. 2005. Arthropod-borne diseases: vector control in the genomics era. *Nature Reviews Microbiology,* 3**,** 262-268.

IRWIN, P. & PASKEWITZ, S. 2009. Investigation of fathead minnows (Pimephales promelas) as a biological control agent of Culex mosquitoes under laboratory and field conditions. *J Am Mosq Control Assoc,* 25**,** 301-9.

JANSSENS, L. & STOKS, R. 2012. How does a pesticide pulse increase vulnerability to predation? Combined effects on behavioral antipredator traits and escape swimming. *Aquat Toxicol,* 110-111**,** 91-8.

JENNIONS, M. D. & MOLLER, A. P. 2002. Publication bias in ecology and evolution: an empirical assessment using the 'trim and fill' method. *Biological Reviews,* 77**,** 211-222.

JOHNSON, B. J. & FONSECA, D. M. 2014. The effects of forced-egg retention on the blood-feeding behavior and reproductive potential of Culex pipiens (Diptera: Culicidae). *Journal of Insect Physiology,* 66**,** 53-58.

JOHNSON, B. J. & SUKHDEO, M. V. K. 2013. Drought-Induced Amplification of Local and Regional West Nile Virus Infection Rates in New Jersey. *Journal of Medical Entomology,* 50**,** 195-204.

JULIANO, S. A. 2009. Species Interactions Among Larval Mosquitoes: Context Dependence Across Habitat Gradients. *Annual Review of Entomology,* 54**,** 37-56.

KALIMUTHU, K., PANNEERSELVAM, D. C., CHOU, C., LIN, S.-M., TSENG, L.-C., TSAI, K.-H., MURUGAN, K. & HWANG, J.-S. 2017. Predatory efficiency of the copepod Megacyclops formosanus and toxic effect of the red alga Gracilaria firma-synthesized silver nanoparticles against the dengue vector Aedes aegypti. *Hydrobiologia,* 785.

KALINOSKI, R. M. & DELONG, J. P. 2016. Beyond body mass: how prey traits improve predictions of functional response parameters. *Oecologia,* 180**,** 543-550.

KAY, B. H., NAM, V. S., VAN TIEN, T., YEN, N. T., PHONG, T. V., DIEP, V. T. B., NINH, T. U., BEKTAS, A. & AASKOV, J. G. 2002. Control of Aedes vectors of

dengue in three provinces of Vietnam by use of Mesocyclops (Copepoda) and community-based methods validated by entomologic, clinical, and serological surveillance. *American Journal of Tropical Medicine and Hygiene,* 66**,** 40-48.

KERSHENBAUM, A., SPENCER, M., BLAUSTEIN, L. & COHEN, J. E. 2012. Modelling evolutionarily stable strategies in oviposition site selection, with varying risks of predation and intraspecific competition. *Evolutionary Ecology,* 26**,** 955-974.

KLECKNER, C. A., HAWLEY, W. A., BRADSHAW, W. E., HOLZAPFEL, C. M. & FISHER, I. J. 1995. PROTANDRY IN AEDES SIERRENSIS - THE SIGNIFICANCE OF TEMPORAL VARIATION IN FEMALE FECUNDITY. *Ecology,* 76**,** 1242-1250.

KNIGHT, T. M., CHASE, J. M., GOSS, C. W. & KNIGHT, J. J. 2004. Effects of interspecific competition, predation, and their interaction on survival and development time of immature Anopheles quadrimaculatus. *J Vector Ecol,* 29**,** 277-84.

KRAMER, L. D., STYER, L. M. & EBEL, G. D. 2008. A global perspective on the epidemiology of West Nile virus. *Annual Review of Entomology,* 53**,** 61-81.

KRAUS, J. M. & VONESH, J. R. 2010. Feedbacks between community assembly and habitat selection shape variation in local colonization. *Journal of Animal Ecology,* 79**,** 795-802.

KROL, L., GORSICH, E. E., HUNTING, E. R., GOVENDER, D., VAN BODEGOM, P. M. & SCHRAMA, M. 2019. Eutrophication governs predator-prey interactions and temperature effects in Aedes aegypti populations. *Parasites & Vectors,* 12.

KUMAR, R. & HWANG, J. S. 2006. Larvicidal efficiency of aquatic predators: A perspective for mosquito biocontrol. *Zoological Studies,* 45**,** 447-466.

KUMAR, R., MUHID, P., DAHMS, H. U., TSENG, L. C. & HWANG, J. S. 2008. Potential of three aquatic predators to control mosquitoes in the presence of alternative prey: a comparative experimental assessment. *Marine and Freshwater Research,* 59**,** 817-835.

KWEKA, E. J., ZHOU, G. F., GILBREATH, T. M., AFRANE, Y., NYINDO, M., GITHEKO, A. K. & YAN, G. Y. 2011. Predation efficiency of Anopheles gambiae larvae by aquatic predators in western Kenya highlands. *Parasites & Vectors,* 4.

LAJEUNESSE, M. J. 2016. Facilitating systematic reviews, data extraction and meta-analysis with the metagear package for r. *Methods in Ecology and Evolution,* 7**,** 323-330.

LEISNHAM, P. T., SALA, L. M. & JULIANO, S. A. 2008. Geographic variation in adult survival and reproductive tactics of the mosquito Aedes albopictus. *Journal of Medical Entomology,* 45**,** 210-221.

LEPCZYK, C. A., FLATHER, C. H., RADELOFF, V. C., PIDGEON, A. M., HAMMER, R. B. & LIU, J. G. 2008. Human impacts on regional avian diversity and abundance. *Conservation Biology,* 22**,** 405-416.

LEPCZYK, C. A., MERTIG, A. G. & LIU, J. G. 2004. Assessing landowner activities related to birds across rural-to-urban landscapes. *Environmental Management,* 33**,** 110-125.

LIMA, S. & DILL, L. 1990. Behavioral Decisions Made under the Risk of Predation: A Review and Prospectus. *Canadian Journal of Zoology-revue Canadienne De Zoologie - CAN J ZOOL,* 68**,** 619-640.

LIMA, S. L. 1998. Nonlethal Effects in the Ecology of Predator-Prey Interactions: What are the ecological effects of anti-predator decision-making? *BioScience,* 48**,** 25-34.

LIU, N. N. 2015. Insecticide Resistance in Mosquitoes: Impact, Mechanisms, and Research Directions. *In:* BERENBAUM, M. R. (ed.) *Annual Review of Entomology, Vol 60.*

LÜHKEN, R., STEINKE, S., LEGGEWIE, M., TANNICH, E., KRÜGER, A., BECKER, S. & KIEL, E. 2015. Physico-Chemical Characteristics of Culex pipiens sensu lato and Culex torrentium (Diptera: Culicidae) Breeding Sites in Germany. *Journal of Medical Entomology,* 52**,** 932-936.

LYIMO, E. O. & TAKKEN, W. 1993. EFFECTS OF ADULT BODY-SIZE ON FECUNDITY AND THE PRE-GRAVID RATE OF ANOPHELES-GAMBIAE FEMALES IN TANZANIA. *Medical and Veterinary Entomology,* 7**,** 328-332.

MALMBORG, P. K. & WILLSON, M. F. 1988. FORAGING ECOLOGY OF AVIAN FRUGIVORES AND SOME CONSEQUENCES FOR SEED DISPERSAL IN AN ILLINOIS WOODLOT. *Condor,* 90**,** 173-186.

MARCANTONIO, M., RIZZOLI, A., METZ, M., ROSA, R., MARINI, G., CHADWICK, E. & NETELER, M. 2015. Identifying the Environmental Conditions Favouring West Nile Virus Outbreaks in Europe. *Plos One,* 10.

MARTEN, G. G. 1990. Elimination of Aedes-Albopictus from Tire Piles by Introducing Macrocyclops-Albidus (Copepoda, Cyclopidae). *Journal of the American Mosquito Control Association,* 6**,** 689-693.

MARTEN, G. G., CHE, W. Y. & BORDES, E. S. 1993. Compatibility of Cyclopoid Copepods with Mosquito Insecticides. *Journal of the American Mosquito Control Association,* 9**,** 150-154.

MARTEN, G. G., NGUYEN, M., MASON, B. J. & NGO, G. 2000a. Natural control of Culex quinquefasciatus larvae in residential ditches by the copepod Macrocyclops albidus. *Journal of Vector Ecology,* 25**,** 7-15.

MARTEN, G. G., NGUYEN, M. & NGO, G. 2000b. Copepod predation on Anopheles quadrimaculatus larvae in rice fields. *Journal of Vector Ecology,* 25**,** 1-6.

MARTEN, G. G. & REID, J. W. 2007. Cyclopoid copepods. *Journal of the American Mosquito Control Association,* 23**,** 65-92.

MCCAULEY, S. J., ROWE, L. & FORTIN, M. J. 2011. The deadly effects of "nonlethal" predators. *Ecology,* 92**,** 2043-8.

MERCER, D. R., WETTACH, G. R. & SMITH, J. L. 2005. Effects of larval density and predation by Toxorhynchites amboinensis on Aedes polynesiensis (Diptera : culicidae) developing in coconuts. *Journal of the American Mosquito Control Association,* 21**,** 425-431.

MICIELI, M. V., MARTI, G. & GARCÍA, J. J. 2002. Laboratory evaluation of Mesocyclops annulatus (Wierzejski, 1892) (Copepoda: Cyclopidea) as a predator of container-breeding mosquitoes in Argentina. *Mem Inst Oswaldo Cruz,* 97**,** 835-8.

MINAKAWA, N., FUTAMI, K., SONYE, G., AKWEYWA, P. & KANEKO, S. 2007. Predatory capacity of a shorefly, Ochthera chalybescens, on malaria vectors. *Malaria Journal,* 6**,** 104.

MITTELBACH, G. G. 1981. FORAGING EFFICIENCY AND BODY SIZE - A STUDY OF OPTIMAL DIET AND HABITAT USE BY BLUEGILLS. *Ecology,* 62**,** 1370-1386.

MOHER, D., LIBERATI, A., TETZLAFF, J. & ALTMAN, D. G. 2009. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ,* 339**,** b2535.

MOORE, S. M., BORER, E. T. & HOSSEINI, P. R. 2010. Predators indirectly control vector-borne disease: linking predator-prey and host-pathogen models. *J R Soc Interface,* 7**,** 161-76.

NANNINI, M. A. & JULIANO, S. A. 1998. Effects of the Facultative Predator Anopheles barberi on Population Performance of its Prey Aedes triseriatus (Diptera Culicidae). *Annals of the Entomological Society of America,* 91**,** 33-42.

NELMS, B. M., FECHTER-LEGGETT, E., CARROLL, B. D., MACEDO, P., KLUH, S. & REISEN, W. K. 2013. Experimental and natural vertical transmission of West Nile virus by California Culex (Diptera: Culicidae) mosquitoes. *J Med Entomol,* 50**,** 371-8.

NELSON, E. H., MATTHEWS, C. E. & ROSENHEIM, J. A. 2004. PREDATORS REDUCE PREY POPULATION GROWTH BY INDUCING CHANGES IN PREY BEHAVIOR. *Ecology,* 85**,** 1853-1858.

NIKOOKAR, S. H., FAZELI-DINAN, M., AZARI-HAMIDIAN, S., MOUSAVINASAB, S. N., ARABI, M., ZIAPOUR, S. P., SHOJAEE, J. & ENAYATI, A. 2017. Species composition and abundance of mosquito larvae in relation with their habitat characteristics in Mazandaran Province, northern Iran. *Bulletin of Entomological Research,* 107**,** 598-610.

O'MEARA, G. F. 1979. Variable expressions of autogeny in three mosquito species. *International Journal of Invertebrate Reproduction,* 1**,** 253-261.

OHBA, S.-Y., OHTSUKA, M., SUNAHARA, T., SONODA, Y., KAWASHIMA, E. & TAKAGI, M. 2012. Differential responses to predator cues between two mosquito species breeding in different habitats. *Ecological Entomology,* 37**,** 410-418.

OLIVER, J. & HOWARD, J. J. 2011. Fecundity of Wild-Caught Gravid Culiseta morsitans (Diptera: Culicidae). *Journal of Medical Entomology,* 48**,** 196-201.

OP DE BEECK, L., JANSSENS, L. & STOKS, R. 2016. Synthetic predator cues impair immune function and make the biological pesticide Bti more lethal for vector mosquitoes. *Ecological Applications,* 26**,** 355-366.

ORR, B. K. & RESH, V. H. 1989. Experimental test of the influence of aquatic macrophyte cover on the survival of Anopheles larvae. *J Am Mosq Control Assoc,* 5**,** 579-85.

OWER, G. D. & JULIANO, S. A. 2019. The demographic and life-history costs of fear: Trait-mediated effects of threat of predation on Aedes triseriatus. *Ecology and Evolution,* 9**,** 3794-3806.

PAULL, S. H., HORTON, D. E., ASHFAQ, M., RASTOGI, D., KRAMER, L. D., DIFFENBAUGH, N. S. & KILPATRICK, A. M. 2017. Drought and immunity determine the intensity of West Nile virus epidemics and climate change impacts. *Proceedings of the Royal Society B-Biological Sciences,* 284.

PEACOR, S. D. & WERNER, E. E. 2001. The contribution of trait-mediated indirect effects to the net effects of a predator. *Proceedings of the National Academy of Sciences,* 98**,** 3904.

PETERS, R. H. 1983. Ingestion. *In:* BECK, E., BIRKS, H. J. B. & CONNOR, E. F. (eds.) *The ecological implications of body size.* Cambridge: Cambridge University Press.

PINTAR, M. R., BOHENEK, J. R., EVELAND, L. L. & RESETARITS, W. J. 2018. Colonization across gradients of risk and reward: Nutrients and predators generate species-specific responses among aquatic insects. *Functional Ecology,* 32**,** 1589-1598.

PONLAWAT, A. & HARRINGTON, L. C. 2007. Age and body size influence male sperm capacity of the dengue vector Aedes aegypti (Diptera : Culicidae). *Journal of Medical Entomology,* 44**,** 422-426.

PREISSER, E. L., BOLNICK, D. I. & BENARD, M. F. 2005. Scared to death? The effects of intimidation and consumption in predator-prey interactions. *Ecology,* 86**,** 501-509.

PYKE, G. H. 2008. Plague Minnow or Mosquito Fish? A Review of the Biology and Impacts of Introduced Gambusia Species. *Annual Review of Ecology Evolution and Systematics,* 39**,** 171-191.

R CORE TEAM 2020. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.

RAJENDRAN, R. & PRASAD, R. S. 1994. A laboratory study on the life cycle and feeding behaviour of Arrenurus madaraszi (Acari: Arrenuridae) parasitizing Anopheles mosquitoes. *Ann Trop Med Parasitol,* 88**,** 169-74.

REINER, R. C., JR., PERKINS, T. A., BARKER, C. M., NIU, T., CHAVES, L. F., ELLIS, A. M., GEORGE, D. B., LE MENACH, A., PULLIAM, J. R., BISANZIO, D., BUCKEE, C., CHIYAKA, C., CUMMINGS, D. A., GARCIA, A. J., GATTON, M. L., GETHING, P. W., HARTLEY, D. M., JOHNSTON, G., KLEIN, E. Y., MICHAEL, E., LINDSAY, S. W., LLOYD, A. L., PIGOTT, D. M., REISEN, W. K., RUKTANONCHAI, N., SINGH, B. K., TATEM, A. J., KITRON, U., HAY, S. I., SCOTT, T. W. & SMITH, D. L. 2013. A systematic review of mathematical models of mosquito-borne pathogen transmission: 1970-2010. *J R Soc Interface,* 10**,** 20120921.

REISEN, W. K., MILBY, M. M. & BOCK, M. E. 1984. THE EFFECTS OF IMMATURE STRESS ON SELECTED EVENTS IN THE LIFE-HISTORY OF CULEX-TARSALIS. *Mosquito News,* 44**,** 385-395.

REISKIND, M. H. & LOUNIBOS, L. P. 2009. Effects of intraspecific larval competition on adult longevity in the mosquitoes Aedes aegypti and Aedes albopictus. *Medical and Veterinary Entomology,* 23**,** 62-68.

RENNE, I. J., GAUTHREAUX, S. A. & GRESHAM, C. A. 2000. Seed dispersal of the Chinese tallow tree (Sapium sebiferum (L.) Roxb.) by birds in coastal South Carolina. *American Midland Naturalist,* 144**,** 202-215.

ROBERTS, D. 2012. Responses of three species of mosquito larvae to the presence of predatory dragonfly and damselfly larvae. *Entomologia Experimentalis Et Applicata,* 145**,** 23-29.

ROBERTS, D. 2018. Predator feeding vibrations encourage mosquito larvae to shorten their development and so become smaller adults: Predator feeding vibrations. *Ecological Entomology*.

ROEHR, B. 2012. US hit by massive West Nile virus outbreak centred around Texas. *British Medical Journal,* 345.

ROHATGI, A. 2020. *WebPlotDigitizer* [Online]. Available: https://automeris.io/WebPlotDigitizer [Accessed 07/14/2020].

ROSEN, L. 1988. Further observations on the mechanism of vertical transmission of flaviviruses by Aedes mosquitoes. *Am J Trop Med Hyg,* 39**,** 123-6.

ROUX, O., DIABATE, A. & SIMARD, F. 2014. Divergence in threat sensitivity among aquatic larvae of cryptic mosquito species. *Journal of Animal Ecology,* 83**,** 702-711.

ROUX, O., VANTAUX, A., ROCHE, B., YAMEOGO, K. B., DABIRE, K. R., DIABATE, A., SIMARD, F. & LEFEVRE, T. 2015. Evidence for carry-over effects of predator exposure on pathogen transmission potential. *Proceedings of the Royal Society B-Biological Sciences,* 282**,** 20152430.

RUBBO, M. J., LANTERMAN, J. L., FALCO, R. C. & DANIELS, T. J. 2011. The Influence of Amphibians on Mosquitoes in Seasonal Pools: Can Wetlands Protection Help to Minimize Disease Risk? *Wetlands,* 31**,** 799-804.

RUSSELL, B. M., MUIR, L. E., WEINSTEIN, P. & KAY, B. H. 1996. Surveillance of the mosquito Aedes aegypti and its biocontrol with the copepod Mesocyclops aspericornis in Australian wells and gold mines. *Medical and Veterinary Entomology,* 10**,** 155-160.

SCHRAMA, M., GORSICH, E. E., HUNTING, E., BARMENTLO, S. H., BEECHLER, B. & VAN BODEGOM, P. M. 2018. Eutrophication and predator presence overrule the effects of temperature on mosquito survival and development. *PLoS Neglected Tropical Diseases,* 3.

SEALE, A. 1917. The Mosquito Fish, *Gambusia affinis* (Baird and Girard), in the Philippine Islands. *Philippine Journal of Science,* 12**,** 177-187.

SHAMAN, J., DAY, J. F. & STIEGLITZ, M. 2005. Drought-induced amplification and epidemic transmission of West Nile Virus in southern Florida. *Journal of Medical Entomology,* 42**,** 134-141.

SHI, L. Y. & LIN, L. F. 2019. The trim-and-fill method for publication bias: practical guidelines and recommendations based on a large database of meta-analyses. *Medicine,* 98.

SIH, A. 1986. ANTIPREDATOR RESPONSES AND THE PERCEPTION OF DANGER BY MOSQUITO LARVAE. *Ecology,* 67**,** 434-441.

SILBERBUSH, A., ABRAMSKY, Z. & TSURIM, I. 2015. Effects of fish cues on mosquito larvae development. *Acta Tropica,* 150**,** 196-199.

SILBERBUSH, A. & BLAUSTEIN, L. 2008. Oviposition habitat selection by a mosquito in response to a predator: are predator-released kairomones air-borne cues? *J Vector Ecol,* 33**,** 208-11.

SILBERBUSH, A. & BLAUSTEIN, L. 2011. Mosquito females quantify risk of predation to their progeny when selecting an oviposition site. *Functional Ecology,* 25**,** 1091-1095.

SOGHIGIAN, J., VALSDOTTIR, L. R. & LIVDAHL, T. P. 2017. A parasite's modification of host behavior reduces predation on its host. *Ecology and Evolution,* 7**,** 1453-1461.

SOUMARE, M. K. F. & CILEK, J. E. 2011. The Effectiveness of Mesocyclops Longisetus (Copepoda) for the Control of Container-Inhabiting Mosquitoes in Residential Environments. *Journal of the American Mosquito Control Association,* 27**,** 376-383.

SPENCER, M., BLAUSTEIN, L. & COHEN, J. E. 2002. OVIPOSITION HABITAT SELECTION BY MOSQUITOES (CULISETA LONGIAREOLATA) AND CONSEQUENCES FOR POPULATION SIZE. *Ecology,* 83**,** 669-679.

STAATS, E. G., AGOSTA, S. J. & VONESH, J. R. 2016. Predator diversity reduces habitat colonization by mosquitoes and midges. *Biology Letters,* 12**,** 20160580.

STAV, G., BLAUSTEIN, L. & MARGALIT, Y. 2005. INDIVIDUAL AND INTERACTIVE EFFECTS OF A PREDATOR AND CONTROPHIC SPECIES ON MOSQUITO POPULATIONS. *Ecological Applications,* 15**,** 587-598.

STERNE, J. A. C. & EGGER, M. 2001. Funnel plots for detecting bias in meta-analysis: Guidelines on choice of axis. *Journal of Clinical Epidemiology,* 54**,** 1046-1055.

STYER, L. M., MEOLA, M. A. & KRAMER, L. D. 2007. West Nile virus infection decreases fecundity of Culex tarsalis females. *Journal of Medical Entomology,* 44**,** 1074-1085.

SU, T. & MULLA, M. S. 1997. Physiological aspects of autogeny in Culex tarsalis (Diptera: Culicidae): influences of sugar-feeding, mating, body weight, and wing length. *J Vector Ecol,* 22**,** 115-21.

SUBRAMANIAM, J., MURUGAN, K., PANNEERSELVAM, C., KOVENDAN, K., MADHIYAZHAGAN, P., KUMAR, P. M., DINESH, D., CHANDRAMOHAN, B., SURESH, U., NICOLETTI, M., HIGUCHI, A., HWANG, J. S., KUMAR, S., ALARFAJ, A. A., MUNUSAMY, M. A., MESSING, R. H. & BENELLI, G. 2015. Eco-friendly control of malaria and arbovirus vectors using the mosquitofish Gambusia affinis and ultra-low dosages of Mimusops elengi-synthesized silver nanoparticles: towards an integrative approach? *Environ Sci Pollut Res Int,* 22**,** 20067-83.

SUGUMARAN, R., LARSON, S. R. & DEGROOTE, J. P. 2009. Spatio-temporal cluster analysis of county-based human West Nile virus incidence in the continental United States. *International Journal of Health Geographics,* 8**,** 43.

SULESCO, T. M., TODERAS, L. G., USPENSKAIA, I. G. & TODERAS, I. K. 2015. Larval Habitats Diversity and Distribution of the Mosquito (Diptera: Culicidae) Species in the Republic of Moldova. *Journal of Medical Entomology,* 52**,** 1299-1308.

THAPA, R., MIRSKY, S. B. & TULLY, K. L. 2018. Cover Crops Reduce Nitrate Leaching in Agroecosystems: A Global Meta-Analysis. *Journal of Environmental Quality,* 47**,** 1400-1411.

TIETZE, N. S., HESTER, P. G., SHAFFER, K. R., PRESCOTT, S. J. & SCHREIBER, E. T. 1994. INTEGRATED MANAGEMENT OF WASTE TIRE MOSQUITOS UTILIZING MESOCYCLOPS-LONGISETUS (COPEPODA, CYCLOPIDAE), BACILLUS-THURINGIENSIS VAR ISRAELENSIS, BACILLUS-SPHAERICUS, AND METHOPRENE. *Journal of the American Mosquito Control Association,* 10**,** 363-373.

TIETZE, N. S. & MULLA, M. S. 1990. Influence of tadpole shrimp, Triops longicaudatus (Notostraca:Triopsidae), stocking rate on Culex tarsalis development in experimental field microcosms. *J Am Mosq Control Assoc,* 6**,** 265-9.

TOWNROE, S. & CALLAGHAN, A. 2014. British Container Breeding Mosquitoes: The Impact of Urbanisation and Climate Change on Community Composition and Phenology. *PLOS ONE,* 9**,** e95325.

TSENG, M. 2007. Ascogregarine parasites as possible biocontrol agents of mosquitoes. *Journal of the American Mosquito Control Association,* 23**,** 30-4.

TSUNODA, T., FUKUCHI, A., NANBARA, S. & TAKAGI, M. 2010. Effect of body size and sugar meals on oviposition of the yellow fever mosquito, Aedes aegypti (Diptera: Culicidae). *J Vector Ecol,* 35**,** 56-60.

UITERWAAL, S. F. & DELONG, J. P. 2020. Functional responses are maximized at intermediate temperatures. *Ecology*.

VAN UITREGT, V. O., HURST, T. P. & WILSON, R. S. 2012. Reduced size and starvation resistance in adult mosquitoes, Aedes notoscriptus, exposed to predation cues as larvae. *Journal of Animal Ecology,* 81**,** 108-115.

VAN UITREGT, V. O., HURST, T. P. & WILSON, R. S. 2013. Greater costs of inducible behavioural defences at cooler temperatures in larvae of the mosquito, Aedes notoscriptus. *Evolutionary Ecology,* 27**,** 13-26.

VERNA, T. N. 2015. Species Composition and Seasonal Distribution of Mosquito Larvae (Diptera: Culicidae) in Southern New Jersey, Burlington County. *Journal of Medical Entomology,* 52**,** 1165-1169.

VERNER, J. & PURCELL, K. L. 1999. Fluctuating populations of House Wrens and Bewick's Wrens in foothills of the western Sierra Nevada of California. *Condor,* 101**,** 219-229.

VERONESI, R., CARRIERI, M., MACCAGNANI, B., MAINI, S. & BELLINI, R. 2015. Macrocyclops Albidus (Copepoda: Cyclopidae) for the Biocontrol of Aedes Albopictus and Culex Pipiens in Italy. *Journal of the American Mosquito Control Association,* 31**,** 32-43.

VIECHTBAUER, W. 2010. Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software,* 36**,** 1-48.

VINCZE, E., SERESS, G., LAGISZ, M., DINGEMANSE, N. & SPRAU, P. 2017. Does Urbanization Affect Predation of Bird Nests? A Meta-Analysis. *Frontiers in Ecology and Evolution,* 5.

VINOGRADOVA, E. B. 2000. *Culex pipiens pipiens mosquitoes: taxonomy, distribution, ecology, physiology, genetics, applied importance and control,* Sofia, Bulgaria, Pensoft.

VONESH, J. R. & BLAUSTEIN, L. 2010. PREDATOR-INDUCED SHIFTS IN
MOSQUITO OVIPOSITION SITE SELECTION: A META-ANALYSIS and
IMPLICATIONS FOR VECTOR CONTROL. *Israel Journal of Ecology &*
Evolution, 56**,** 263-279.
WALLACE, J. R. & MERRITT, R. W. 1999. Influence of microclimate, food, and predation
on Anopheles quadrimaculatus (Diptera : Culicidae) growth and development rates,
survivorship, and adult size in a Michigan pond. *Environmental Entomology,* 28**,** 233-
239.
WEAVER, S. C. & REISEN, W. K. 2010. Present and future arboviral threats. *Antiviral*
Research, 85**,** 328-345.
WORLD HEALTH ORGANIZATION (WHO). 2020. *Vector-borne diseases* [Online].
Available: https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases
[Accessed 02/07/2021].
XUE, R. D., BARNARD, D. R. & MULLER, G. C. 2010. Effects of Body Size and
Nutritional Regimen on Survival in Adult Aedes albopictus (Diptera: Culicidae).
Journal of Medical Entomology, 47**,** 778-782.
YANOVIAK, S. P. 1999. Effects of Mecistogaster spp. (Odonata: Pseudostigmatidae) and
Culex mollis (Diptera: Culicidae) on Litter Decomposition in Neotropical Treehole
Microcosms. *The Florida Entomologist,* 82**,** 462-468.
ZUHARAH, W. F., FADZLY, N. & LESTER, P. J. 2013. Lethal and sublethal impacts of
predaceous backswimmer Anisops wakefieldi (Hemiptera: Notonectidae) on the life-
history traits of the New Zealand mosquito Culex pervigilans (Diptera: Culicidae). *J*
Med Entomol, 50**,** 1014-24.
ZUHARAH, W. F. & LESTER, P. J. 2010. The influence of aquatic predators on mosquito
abundance in animal drinking troughs in New Zealand. *Journal of Vector Ecology,*
35**,** 347-353.

1322