Regulatory perturbations of ribosome allocation results in a trade-off between fast growth and adaptation capacity

Authors:

David Hidalgo[1], César A. Martínez-Ortiz[1], Bernhard O. Palsson[2,3], José I. Jiménez[4], José Utrilla[1*]

[1]Centre for Genomic Sciences, Universidad Nacional Autónoma de México, Campus Morelos, Cuernavaca, México

[2]Department of Bioengineering, University of California San Diego, La Jolla, California, United States of America

[3]Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Lyngby, Denmark

[4]Department of Life Sciences, Imperial College London, South Kensington Campus, London, SW7 2AZ, United Kingdom

*Corresponding author:

Name: José Utrilla

Address: UNAM-Centro de Ciencias Genómicas. Av. Universidad s/n Col. Chamilpa. Cuernavaca, Morelos, México. 62210

utrilla@ccg.unam.mx

Highlights

Strains with the 3 stronger ribosomal operons grow faster than the wild-type

Fast-growing strains show higher ribosomal content

Comparison among *rrn* and RNAP mutants shows commonalities

Fast-growing strains show adaptation trade-offs

Mutant's proteome is larger and optimized for growth

Abstract

Bacteria regulate their cellular resource allocation to enable their fast growth-adaptation to a variety of environmental niches. We studied the ribosomal allocation, growth and expression profile of two sets of fast-growing mutants of *Escherichia coli* K-12 MG1655. Mutants with 3 copies of the stronger ribosomal RNA operons grew faster than the wild-type strain in minimal media and show similar phenotype to previously studied *rpoB* mutants. All of them displayed increased ribosomal content, a longer diauxic shift and a reduced activity of the *aceBAK* operon, indicative of repressed gluconeogenic pathways. Transcriptomic profiles of fast-growing mutants showed common downregulation of hedging functions and upregulated growth functions. Proteome allocation estimations showed an increase in the growth-related proteome for fast-growing strains, but not an increased cellular budget for recombinant protein production. These results show that two different regulatory perturbations (rRNA promoters or *rpoB* mutations) increasing ribosomal allocation optimize the proteome for growth with a concomitant fitness cost.

Keywords: ribosomes, rRNA, resource allocation, fitness, RNA polymerase.

Introduction

Cellular growth is the result of a highly regulated process that demands the allocation of cellular resources towards different tasks. Bacteria have evolved to regulate resource allocation to maximize survival under harsh and changing environments. During fast exponential growth ribosomal RNA represents ~86% of the total RNA, ribosomes constitute over 30% of the biomass (Dennis and Bremer, 2008), and translation uses ~40% of the cellular energy (Wilson and Nierhaus, 2007). Therefore, the regulation of ribosomal content is central to bacterial physiology.

The main control step in ribosome biogenesis is the transcription of ribosomal RNA (rRNA) (Dennis et al., 2004), which hoards much of the available RNA polymerase (RNAP). RNAP is central to the regulation of the ribosomal allocation, it is the target of the (p)ppGpp stringent response and fast-growing strains selected by Adaptive Laboratory Evolution (ALE) often present mutations in the *rpoB* or *rpoC* genes that code for the core RNAP (Phaneuf et al., 2019). ALE selects for RNAP mutants that grow faster than the parental strain in minimal media, display adaptation trade-offs (such as the glucose-acetate diauxic shift) and slower growth in rich media. The effects of these mutations have been attributed to the reprogramming of the whole transcriptome, and the reallocation of resources from downregulated hedging functions to growth, however their effects on the regulation of the ribosome allocation have not been studied (Cheng et al., 2014; Conrad et al., 2010; Utrilla et al., 2016).

A clear correlation between copy number of ribosomal RNA operons, growth rate and growth efficiency has been established for many bacterial species (Roller et al., 2016). In the model bacterium *Escherichia coli* rRNA is transcribed from seven ribosomal operons (*rrn*A, B, C, D, E, G and H) that provide the necessary amount of ribosomal transcription for fast growth. However, it has been shown through progressive deletions that *E. coli* can survive with only one of the seven (Asai et al., 1999a). The resulting ribosomal operon elimination mutants have been studied in rich medium, showing reduced growth rate, increased adaptation time in glucose-LB shift-up assays and reduced RNA content (Asai et al., 1999b; Condon et al., 1995). Previous studies in *E. coli* have demonstrated that not all the ribosomal operons have the same transcription strength in different conditions, even though the structure of the *rrn* promoters is similar, the genomic location and slight differences in the promoter sequences result in different promoter strength and sensitivity to stringent response (Kolmsee et al., 2011; Maeda et al., 2015). Also, there is evidence showing that poor nutrient conditions produce heterogeneous populations of ribosomes that regulate the expression of stress response genes (Kurylo et al., 2018).

It has been recently shown that the cost of the translationally inactive (unused) ribosomes, present in *E. coli*, may reduce the growth capacity in a static environment but can be useful in a dynamic or changing environment (Dai et al., 2016). A negative correlation between growth rate and adaptation rate has been recently observed (Basan et al., 2020). This evidence suggests that a fraction of inactive ribosomes may confer fitness advantages in changing conditions, such as those found in nature, however, under constant environments the inactive ribosomes may constitute a burden that constrains the maximum achievable growth rate. An open question regarding the mechanism of growth rate increase of ALE selected mutants in constant minimal media conditions, is if they reduce the amount of inactive ribosomes. The reduction of costly inactive ribosomes could increase growth rate in constant environments but would result in a loss of capacity to resume growth under changing conditions.

Here we studied the allocation of ribosomal resources in two sets of strains with regulatory perturbations that increase growth rates in minimal media. Two mutants with the three stronger ribosomal operon (SQ53: *rrn*CEH, SQ78: *rrn*BCH) expression resulted in higher resource allocation to growth and decreased adaptation capacity. These results are similar to the optimized growth strategies displayed by fast-growing RNAP mutants obtained by ALE (*rpoB*E546V and *rpoB*E672K), which highlights common and specific mechanisms of regulatory reprograming of ribosomal allocation.

Results

The effect of *rrn* operon number on the growth phenotype in different environments

Mutants in ribosomal operons have been constructed previously and partially studied (Quan et al., 2015). However, the implications of the reduction of ribosomal operons in the general cellular physiology has not been studied. Even less from the perspective of analyzing the regulation of growth phenotypes of the mutants under different environments. To this end we grew the Wild Type (WT: *E. coli* MG1655) strain and the mutants lacking one to six *rrn* operon copies (Table 1, Quan et al., 2015) in rich and minimal media supplemented with different carbon sources. The results showed that while there is a positive correlation between growth rate and number of *rrn* operons in rich medium (LB with glucose and M9 with glucose supplemented with casamino acids, CAA) (Figure 1A), in minimal media with different carbon sources there is an optimal amount of *rrn* operons that maximizes the growth rate (3 *rrn* operons; SQ53: *rrn*CEH, SQ78: *rrn*BCH) and this number is different from that of the WT strain (7 *rrn*) (Figure 1B).

During our experiments in glucose M9 medium, where acetate is typically produced as an overflow metabolite and latter consumed in a second growth phase, we observed that faster *rrn* mutants showed a diminished ability to resume exponential growth once glucose had been exhausted (Fig 1D), while the WT strain transitioned swiftly through the glucose-acetate shift. At this point we recognized similarities between the phenotype exhibited by *rrn* mutants and two previously reported strains (*rpoB*E546V and *rpoB*E672K, Utrilla et al., 2016) each carrying a point mutation in the *rpoB* gene of the RNAP. Both types of mutants grew faster in glucose minimal media, showed glucose-acetate growth shift defects, and grew slower than the WT in rich media, such as glucose + CAA and LB with glucose (Figures 1C, D and E).

Figure 1. Growth and growth rates. A, the effect of *rrn* operon copy number on the growth rate in rich media (LB with glucose and glucose supplemented with casamino acids). B, the effect of *rrn* operon copy number on the growth rate in minimal media (glucose, arabinose and galactose). C, Triplicate

growth dynamics of selected *rrn* strains and two *rpoB* mutants in LB with glucose. D, Triplicate growth dynamics of selected fast strains in glucose. E, Growth rates of selected strains in LB supplemented with glucose. F, Growth rates of selected strains in glucose minimal medium. Error bars represent standard deviations from nine replicates; some are smaller than the size of the marker.

The ribosome production and the growth rate change proportionally in all mutants

Recently many studies have pointed to the fact that reserve ribosomes and translation efficiency may play a fundamental role in the observed maximum growth rate in a given condition (Dai et al., 2016). Since we studied mutants with a reduced ribosomal operon copy number that grow faster than the WT strain, we hypothesize that a reduced amount of ribosomal content plus a higher translation efficiency may be a possible mechanism for the increased growth rate of the mutants in minimal media. Therefore, one of the main questions we addressed in this work was whether the ribosome production in the studied mutants changes at the same magnitude as the WT in different growth environments. To measure ribosome content in the studied strains we used a previously reported *in vivo* system to indirectly monitor ribosome concentration. A ribosomal protein-GFP fusion was used to replace the native *rplI* gene (Fig. 2A), whose protein product is a component of the large subunit of the ribosome (Kim et al., 2021).

We measured ribosomal content (via GFP fluorescence) and maximum growth rates (μ) in 11 different conditions which rendered μ ranging from 0.24 to 1.53 h^{-1}. We plotted the normalized *rplI*:msfGFP fluorescence during exponential growth as a function of growth rate (Fig 2B); we observed that regardless of the number of rRNA deletions, the slope of the regression of ribosomal content and the and μ is maintained (Fig. 2C). The results also showed that the mutants that grew faster than the WT strain in the same environment did so by increasing the ribosomal content at a similar magnitude than the WT strain for different environments (data points fell on the same linear trend as the WT). These observations show that the increase in growth rate was derived from an increase in ribosomal content and not from a reduction of inactive ribosomes or increased in ribosomal efficiency. We conclude this since the reduction of inactive ribosomes or the increase of ribosomal efficiency would result in higher growth rate with similar o less ribosomal content shifting the trend from what was shown by the WT. Previous evidence (Kolmsee et al., 2011, Maeda et al., 2015) showed that the promoters of each rRNA operon have different strengths and sensitivity to environmental and regulatory signals such as (p)ppGpp (Fig. S5). Utrilla et al, showed that even though the *rpoB*E546V and E672K mutations are not in the specific (p)ppGpp binding site, they belong to the same "structural community" where molecular motions of residues are strongly correlated. They observed that these mutations perturb the stability of the RNAP open complex in the presence

of (p)ppGpp perturbing the response to stringent response with implications in its affinity for sigma factors, ribosomal promoters and other regulatory effects at the transcriptional level. Since both set of mutants increase ribosomal content proportional to the growth increase in 11 evaluated media (Figs S1 and S2), we postulate that the mechanism for doing so is the stronger expression of ribosomal RNA resulted from two different molecular mechanisms: the use of the stronger set of ribosomal promoters in SQ53 and SQ78 strains and the perturbed stability of the RNAP open complex in *rpoB* mutants.

Figure 2. *in vivo* ribosome content measurements. A, The ribosomal protein-GFP fusion in the chromosome. An 18 bp linker (coding for amino acids SGGGGS) is placed between the genes (Not to scale). The fused genes encode the ribosomal L9-msfGFP chimeric protein. TAA denotes end of translation. B, Linear regression of the normalized *rplI*:msfGFP fluorescence as a function of growth rate for all strains in several conditions. The yellow line represents the regression for all strains data combined. Each strain's corresponding points are categorized by color (see legend). Error bars are the standard deviation from triplicates. The regression line's shadow is the 95% confidence. C, Comparison of each strain's individual normalized *rplI*:msfGFP slopes obtained by linear regression. No significant differences between any mutant and MG1655 (95% confidence) were found.

Faster growth phenotypes show a reduced adaptation capacity

In *E. coli* aerobic cultures, growth in glucose minimal medium is accompanied by acetate production (Enjalbert et al., 2015). Two well-defined growth phases can be differentiated in these cultures with a maximum growth rate in glucose and another in acetate (Fig. 3A). We observed that mutants displayed a reduced capacity to adapt and grow in the acetate growth phase. Figure 3B shows the maximum growth rates (µ) before (lilac) and after (green) the diauxic shift. While fast growing mutants (SQ53: *rrn*CEH, SQ78: *rrn*BCH and both in *rpoB*) had a higher growth rate during the glucose phase, they were slower than the WT after the shift (only 28% of the growth rate achieved by the WT for SQ53 and *rpoB*E546V). SQ110, with only one *rrn* copy, grows so poorly that it did not even have a diauxic shift. Regarding the time required to reach the post-shift maximum growth rate, our data showed a delay correlated with the number of rrn deletions (Fig. 3C). However, this may depend on the precise deletions as exemplified by SQ53 which was particularly slow in contrast to SQ78 despite both containing 3 *rrn* copies (see discussion below). Additionally, both *rpoB* mutants showed a similar diauxic behavior to that of SQ78, though with slight differences in the post-shift growth rate which decreased by 28% in *rpoB*E546V and 68% in *rpoB*E672K. These observations show an important phenotypic trait in the fast-growing mutants: they took longer to adapt to a new carbon source and resume exponential growth in acetate and their growth rates were lower than MG1655 in the post-shift phase.

Figure 3. Growth dynamics and PaceB activity. A, Typical bacterial diauxic growth diagram. First, a single carbon source is consumed, reaching a growth rate (μ1). After a pause, another carbon source is consumed and exponential growth is resumed, reaching a second growth rate (μ2) lower than the first one. B, Maximum growth rates (μ) in glucose before (lilac) and after (green) the growth shift. C, Time at maximum growth rates (μ) in glucose before (lilac) and after (green) the growth shift. D, Dynamics of the normalized activity of the *aceB* promoter (dGFP/dt/OD) in MG1655. Dashed line represents the time of growth shift. Red lines represent activity slopes analyzed in panels E and F. E, Comparison of normalized *PaceB* activity slope towards maximum between mutants and MG1655. *

denotes significant difference with 95% confidence. F, Comparison of normalized *PaceB* activity slope after maximum between mutants and MG1655. * denotes significant difference with 95% confidence.

aceB promoter activity dynamics

The genes *aceA* and *aceB* are essential for growth in acetate. They are responsible for the glyoxylate shunt in the tricarboxylic acid cycle and show a strong transcriptional response in the glucose-acetate shift (Enjalbert et al 2015). In order to test the regulatory response of the *aceBAK* operon in the fast-growing mutants, we used a reporter plasmid containing the *aceB* promoter controlling the expression of GFP (Zaslaver et al., 2006). The normalized Promoter Activity (PA), a measure of RNAP transcription at a certain promoter, was monitored *in vivo* over time in M9-glucose 4 g/L cultures (dGFP/dt/OD, Fig. 3D). In those conditions, two clear phases were observed: a) a gradual increase in *aceB* PA to reach a maximum value of activity hours before the growth shift, and b) a decrease in *aceB* PA towards the transition to acetate growth. The slopes for each phase of this activity dynamics (blue and yellow lines) were determined by a linear regression (Fig. 3E and F). There was a 12-20% reduction in the SQ53 *aceB* PA slope towards its maximum and 54% of the WT PA for the *rpoB* E672K mutant. After this point, the PA slope fell abruptly for the studied mutants with a reduction ranging from 2.47 to 1.37-fold of the WT *aceB* PA. Since the concentration of acetate in each culture may be different and could affect the regulatory response, we also measured the *aceB* PA dynamics in media with glucose-acetate mixtures showing results consistent with those mentioned above: a lower slope of the aceB PA during the first phase and a higher negative slope during the second growth phase (Figures S3 and S4). These results showed a reduced transcription of the *aceBAK* operon in both the rRNA and RNAP mutants and suggested a reduced availability of the RNAP to transcribe the genes needed to efficiently adapt to a new carbon source that resulted in a loss of adaptation capacity in the fast-growing mutants.

The transcriptional profiles of *rrn* and *rpoB* mutants show common differentially expressed functions

After assessing similarities during growth associated with the RNAP in both types of mutants, we compared the transcriptomes in glucose minimal medium of two selected fast strains: SQ53 and the previously reported *rpoB* E672K mutant. Increased growth rate of *rpoB* mutants in glucose minimal medium has been attributed to the reduction of hedging functions, having their associated cellular resources being channeled to growth functions (Kim et al., 2020; Utrilla et al., 2016).   Results for SQ53 showed 104 Differentially Expressed Genes (DEGs), of which 74 are downregulated (Figure 4A). Broadly, the downregulated genes belong to

hedging functions such as oxidative and osmotic stress, acid resistance, general stress responses and a portion of genes of unknown function (Figure 4B). For the 30 upregulated genes we observed the majority of growth functions such as carbohydrate transport and metabolism, amino acid uptake, and metabolism and protein synthesis. When we compared the expression profile changes among SQ53 and *rpoB* mutants we found similarities in terms of the broad categories of DEGs, such as osmotic and oxidative stress, acid resistance, ion transport and homeostasis and genes with unknown function.

Figure 4. Transcriptomic profile comparison between fast *rrn* and *rpoB* mutants. A, Number of upregulated (red) and downregulated (blue) genes for SQ53 and *rpoB*. Intersections denote the number

of shared genes between mutants. B, Heatmap showing expression levels of hedging and growth functions for SQ53 and rpoB672. Genes were categorized by the functions shown.

We further analyzed the data by performing a proteome balance calculation to estimate the contribution in mass by each DEG (Lastiri-Pancardo et al., 2020, see methods). While a given gene may show differences in its transcription, it is the size and copy number of the corresponding protein that has the largest impact on the cellular resource and energy budget (Lynch and Marinov, 2015). From these calculations we observed that most of the impact on the cellular budget was associated with a few proteins showing a Pareto-shape response regardless of the mutant. This means that 80% of the change in cellular budget associated with gene expression comes from just a few genes (Figure 5). For upregulated genes, most categories corresponded to amino acid uptake, protein synthesis, carbohydrate transport and metabolism and ion homeostasis. For downregulated genes, the associated categories are more heterogeneous. Within the genes with more impact, functional categories corresponded to oxidative and osmotic stress, DNA damage and metabolic process, ion homeostasis and defense mechanisms. Another important observation derived from these calculations is that the estimated proteome change was higher for upregulated functions compared to the down regulated ones. Upregulated genes for SQ53 contributed three times the mass of those downregulated (6.27 vs 2.09 fg., respectively). For the *rpoB*E672K mutant the difference is more prominent, as upregulated genes contributed 5.97 times more than downregulated genes (52.43 vs 8.76 fg., respectively). This means that the estimated increase in the growth proteome for both studied mutants is higher than the reduction of the hedging functions. This may explain the better capacity to grow in minimal media but the reduced ability to shift conditions. The mutants show a larger and more specialized proteome that prevents them from being prepared for environmental adaptations.

Figure 5. Proteome mass contribution in femtograms for DEGs. A, B Proteome mass contribution in femtograms for up and downregulated genes for each strain, respectively. The yellow line represents the cumulative percent (right axis) for the 20 genes with the largest contribution.

Greed versus fear response and the proteomic budget of fast-growing mutants

In order to get a deeper understanding of the transcriptomic changes in fast growing strains we used the iModulon classification of the DEG data sets. iModulons are a set of genes obtained by independent component analysis of transcriptomic data sets that reflect the transcriptional regulatory network of *E. coli* (Sastry et al., 2019).We found significant and conserved changes in few iModulons: there was a general increase of the iModulons related to growth (pyruvate, BCAA-2), and a general reduction to the "fear" iModulons (*rpoS*) (Figs. 6A and S6). Tan et al 2020 reported the expression patterns in different heterologous protein production strains and observed that those proteins that elicited a weaker expression of fear iModulon were more productive for heterologous protein production. We therefore assessed the availability of cellular resources for recombinant gene expression using the isocost lines methodology that measures the protein budget on the basis of the relationship of two co-expressed fluorescent reporters (Gyorgy et al., 2015). Even though the fast-growing strains show a higher expression of the translation machinery iModulons ("greed") and reduced the expression of the *rpoS* iModulons (fear), we found a reduced proteomic budget in these strains (Figure 6B). Recently, it has been observed that in a defined environment fast growth and unnecessary protein production are inversely correlated (Kim et al., 2021; Scott et al., 2010). Therefore, the increased allocation of resources to growth obtained in these strains is not correlated with a higher budget for a synthetic function. Taken all together, our results suggest that the similar phenotype of the RNAP and rRNA mutants is the result of different expression profiles with similar categories of genes being differentially expressed.

Figure 6. iModulons and isocostlines. A, DEGs categorized by iModulons for SQ53 and *rpoB*E672K. Color bar indicates fold change expression level. B, Isocost lines indicating differences in cellular budget between mutants and the WT.

Discussion

The bacterial growth rate is the result of a complex regulatory interplay that fine tunes the allocation of cellular resources such as the ribosomes. Several works have proven that the growth rate in a determined environment can be increased if the proteome is optimized, by regulatory and other mutations, to that specific environment (Cheng et al., 2014; Heckmann et al., 2020; LaCroix et al., 2015). Indeed, rewiring *E. coli* transcription networks with fusions of promoters and master regulators (e.g. *arcA*, *crp*, *rpoD*) upregulated ribosomal-associated genes and increased growth rates (Baumstark et al., 2015). Here, we show that different regulatory perturbations that increase ribosomal synthesis result in higher growth rates but are also associated with phenotypic and molecular trade-offs.

In this work, we studied *E. coli* mutants lacking one to six copies of the seven ribosomal operons. We found that, while in rich media ribosomal deletion results in a reduced growth, there is a combination of three ribosomal operons that results in higher growth in minimal media. We compared the fast-growing ribosomal mutants to the ALE selected RNAP mutants (*rpoB*E546V and *rpoB*E672K) due to their striking growth phenotype similarity: these strains exhibit increased growth rate in minimal media with longer diauxic shifts and decreased growth rate in rich media. To explain the observed phenomenon, we postulate two hypotheses A) a reduced amount of ribosomal expression accompanied with increased translation efficiency, and B) a regulatory effect on RNAP distribution, being able to increase the expression of rRNA in very specific conditions with faster growth and a reprogramming of the whole transcriptome. Our *in vivo* measurements of ribosomal content show that these fast-growing strains did so by increasing ribosome synthesis, their data points fell into the same trendline of ribosomal content versus growth rate as the WT, ruling out the first hypothesis. Even though some studies point to an increase in ribosomal efficiency or the reduction of reserve ribosomes as a possible mechanism of increase in growth rate (Dai et al., 2016), here we show that the increase in growth rate in the studied mutants was due to the increase in ribosomal content. We attribute this to two different mechanisms with similar consequences: the previously shown differential regulation and strengths of ribosomal operons (Kolmsee et al., 2011; Maeda et al., 2015) or mutations in RNAP that may perturb the sensibility to the (p)ppGpp stringent response. Both mechanisms can perturb the (p)ppGpp - ribosomal allocation regulatory circuit. Therefore, in wild type *E. coli* the (p)ppGpp - ribosomal allocation regulatory circuit is fine tuned to constrain the maximum growth rate with a better adaptation response.

We found that mutants that showed faster growth rate in glucose also showed the slowest growth rate in the acetate consumption phase. This agrees with a study (Basan et al., 2020)

where cultures with higher pre-shift growth rates show the longest times to resume exponential growth in shift-down experiments due to reduced metabolite pools and enzyme distribution. Furthermore, the *aceB* PA results show that these fast-growing strains allocate less resources to anticipate the carbon source shift. This is very likely due to the sensitivity of the *rrn* promoters to environmental and regulatory signals, the SQ53 (*rrn*CEH) and the SQ78 (*rrn*BCH) strains both harbor the *rrn*H ribosomal operon that has been shown to be upregulated in minimal media (Kurylo et al., 2018). Furthermore, as it has been previously shown, the P1 promoters of *rrn* operons have heterogeneous strength and sensitivity to (p)ppGpp (Kolmsee et al., 2011, Maeda et al., 2015, Fig.S5). The SQ53 mutant has the strongest and least (p)ppGpp sensitive P1 promoter (*rrn*E). The SQ78 mutant, differing only in having *rrn*B instead of *rrn*E, has a more responsive P1 promoter to reduce rRNA transcription. Our results suggest that the sensitivity of the P1 promoters to regulatory signals allows the RNAP to be allocated to the transcription of genes required for shifting growth to a gluconeogenic carbon source in a more timely fashion. In summary, the WT strain with higher aceB PA had a shorter time to resume exponential growth compared to the fast-growing mutants with lower aceB PA. The fast-growing *rpoB* mutants showed a similar phenotype, and while the detailed molecular mechanism of the regulatory reprograming in these mutants needs to be further studied, overall these results point to a reduced sensitivity to (p)ppGpp that drives RNAP to highly express ribosomal operons with a lower capacity to shift to alternative promoters.

The transcriptome of the fastest growing strain, SQ53, showed a general profile similar to what was previously observed for the *rpoB*E672K strain. There was a conserved response of increased growth functions and a reduction of hedging functions, however the specific genes differentially expressed in the strains were not the same (Utrilla et al. 2016). In this study, we conducted a detailed assessment of the proteomic contribution of each DEG in both mutants compared to the WT strain. We observed that there is a greater increase in proteome related to growth functions than the estimated proteome reduction from downregulation of hedging functions. These results provide new insights into the molecular phenotype of the fast-growing mutants: they produce a larger and more specialized proteome resulting in a trade-off in adaptation capacity. Here we show that the fast growth phenotype may be associated to a more efficient recruiting of the RNA polymerase to the *rrn* promoters with a concomitant reduced expression of bet-hedging genes such as those of metabolic adaptation to gluconeogenic carbon sources, thus highlighting its evolutionary role for selecting fast growth phenotypes.

Analyzing the transcriptomic changes with the iModulon methodology we identified a general shift in the greed versus fear response; however, we did not observe a larger proteomic budget for a synthetic circuit expression as measured by isocost lines. Even though previous works show that less fear iModulon expression, when expressing unnecessary proteins, results in higher proteomic budget (Ceroni et al., 2015; Tan et al., 2020), our results show the growth rate related greed versus fear balance do not result in increased proteomic budget. Recent evidence shows that a faster growth rate in a given environment results in a reduced budget for a genetic circuit (Kim et al., 2021). From a resource allocation and a cellular minimization point of view (Hidalgo and Utrilla, 2019), the detailed study of these strains is an important effort to understand and harness design principles for tailored cellular chassis construction.

In this work, we took a systems-based approach to study perturbations on rRNA synthesis at a molecular level. We show that increasing ribosomal content resulting from regulatory perturbations increases growth rate in minimal media but has an adaptation capacity cost and less resource availability for engineered functions. Our findings show that the *E. coli* proteome can naturally achieve faster growth rates in minimal media but its regulatory program limits the ribosomal allocation to channel resources for hedging functions. The understanding of the correlation between growth, gene expression machinery, and its regulation, are of great interest for those studying the general physiology of bacteria and can help to guide successful bioengineering efforts.

Acknowledgements

We thank H. King for rRNA quantification equipment support. We acknowledge the funding provided by UNAM–DGAPA-PAPIIT project IN213420 and Newton advanced Fellowship Project NA160328. D.H. is a doctoral student from Programa de Doctorado en Ciencias Biomédicas, Universidad Nacional Autónoma de México (UNAM), and received a fellowship 508351 from CONACYT. C.M. acknowledges the Programa de Maestría y Doctorado en Ciencias Bioquímicas, UNAM and the masters scholarship 825745 from CONACyT.

Author contributions

Experiments, data processing, statistical analyses and figures were conducted by D.H. RNA extraction, library construction and initial RNAseq data processing and analysis was carried out by C.M. Critical analysis of experimental data and RNAseq categorization was done by D.H and J.U. *In vivo* ribosomal measurements analysis and isocost lines were carried by D.H. and J.J. The project was conceived by D.H. and J.U. Supervision and guidance by J.U. The manuscript was written by D.H., J.U., J.J. and B.P.

Declaration of interests

B.P. is inventor in a US patent filed by UCSD concerning the *rpoB* mutant strains.

Accession numbers

The accession number for the RNAseq data of SQ53 and processed files of SQ53 and rpoBE672K reported here is GEO:GSE180830. The accession number for the RNAseq data of rpoBE672K reported in Utrilla et al., 2016 is GEO:GSE59377.

Methods

Strains

The WT strain used in this work was *E. coli* K12 MG1655.

The *rrn* deletion strains are derived from *E. coli* K12 MG1655 and were generated by Quan et al. 2015. and obtained through the Coli Genetic Stock Center (CGSC, http://cgsc2.biology.yale.edu/)

Table 1 summarizes the strain's genotype.

Table 1. *rrn* deletion strains

Strain	Genotype	*rrn* in chromosome
E. coli MG1655	*ilvG rfb-50 rph-1*	7: ABCDEGH
SQ37	Δ*rrn*E	6: ABCDGH
SQ40	Δ*rrn*EG	5: ABCDH
SQ49	Δ*rrn*GBA	4: CDEH
SQ53	Δ*rrn*GBAD	3: CEH
SQ78	Δ*rrn*GADE	3: BCH
SQ88	Δ*rrn*GADEH(ptRNA67)	2: BC
SQ110	Δ*rrn*GADBHC(ptRNA67)	1: E

Table 1. Strain genotypes. Mutants derived from the K12 MG1655 strain contain deletions of one or several *rrn* operons. Each *rrn* operon contains a copy of the *rrn*, *rrl* and *rrf* genes. In the intergenic and distal positions of these operons there are tRNA genes for different amino acids. Strains may be complemented with a plasmid containing tRNA genes when needed (ptRNA67, SpcR).

The *rpoB* mutants *rpoB*E546V and *rpoB*E672K were obtained from Utrilla et al., 2016.

Growth media

Rich complex media used in this work were LB (Sigma), LB with 1 g/L glucose and M9 minimal media with 0.2% casamino acids (cAA) (Sigma) and 4 g/L glucose. The minimal medium used was M9 (33.7 mM, Na2HPO4; 22.0 mM, KH2PO4; 8.55 mM, NaCl; 9.35 mM, NH4Cl; 1 mM, MgSO4; 0.3 mM, CaCl2) supplemented with trace nutrients (13.4 µM, EDTA; 3.1 µM, FeCl3 - 6H2O; 0.62 µM, ZnCl2; 76 nM, CuCl2 - 2H2O; 42 nM, CoCl2 - 2H2O; 162 nM, H3BO3; 8.1 nM, MnCl2 - 4H2O).

All carbon sources (glucose, glycerol, arabinose, galactose, sodium gluconate, xylose, fructose) were at 4 g/L except for acetate and succinate (2 g/L) unless otherwise stated (diauxic shift experiments). Antibiotic concentrations were Gentamicin 25 µg/mL, Spectinomycin (Spc) 50 µg/mL and Kanamycin 50 µg/mL unless otherwise stated.

Cell growth

Frozen strains in glycerol at -80°C were grown in solid medium and incubated overnight to 24h at optimal temperature. A colony was put into liquid medium with the corresponding antibiotics were needed and incubated overnight to 24h in agitation (37°C)

On the day of the experiment, either plates (50 µL) of flasks (70 mL) were inoculated at 0.05 OD 600 nm.

Growth-associated calculations and determinations

Growth rates

The growth rates for the plate experiments (3 replicates per plate for 3 independent plates) were determined computationally using an algorithm based on gaussian processes (Swain *et al.*, 2016) running on Python. From this analysis, the magnitude and time of the maximum growth rate and of the maximum OD were obtained, as well as the time and magnitude of local growth rate maxima. Additionally, for selected growth media, the time and OD at diauxic changes was analyzed manually.

To corroborate the validity of the determinations, growth rates from these experiments were also calculated computationally by linear regression.

For the flask experiments, growth rates were calculated by linear regression of the data during exponential growth. The average and standard deviation of three independent experiments is reported.

When needed, fluorescence intensity in plate experiments was also measured.

rplI:msfGFP-Tagged strains

Strains were electroporated with the cpEMG *rplI*:msfGFP plasmid and recovered in rich medium overnight. 200 µL of the recovery were plated in LB Km (25 µg/mL). Cointegration in green fluorescent colonies was verified by PCR using primers emg rpli Fwd (5'-agggataacagggtaatctgcgctcgctacctgtccctgct-3') and yifz emg Rev (5'-gcctgcaggtcgactctagagtatttattgcaagatgtcgaat-3'). Km resistance was not removed since it does not interfere with the rplI:msfGFP fusion. Experiments were run in the absence of the antibiotic.

pPaceB-GFP

The Km resistant plasmid containing the aceB promoter fused with GFP was constructed by Zaslaver et al., 2006). Cells were transformed by electroporation following a 1-hour recovery and plated on LB with Km. Fluorescent colonies were selected for the plate assays.

RNA sequencing

SQ53 and MG1655 cells were cultured at 37°C in 25 mL of glucose minimal medium. During early exponential growth, 3 mL were sampled into a tube and 3mL of RNA protect reagent were added and mixed. After a 5-minute incubation, samples were lysed enzymatically (Lysozyme from chicken egg white, Sigma). RNA from samples was purified using a modified version of the Quick-RNA miniprep kit from Zymo research. RNA integrity of purified samples was verified by gel electrophoresis (1% agarose) using TAE buffer. Then, 10 µL of each replicate at a 500 ng/µL concentration was used as the input material to be separated from ribosomal RNA using the Zymo-Seq RiboFree Total RNA Library kit from Zymo research. The resulting material was used to construct the libraries using the TruSeq Stranded mRNA kit from Illumina and sequenced using the Illumina Next-Seq 500 platform. Libraries were sequenced to 10 million paired reads. These reads were then verified for quality and the adapter sequences were cut. Sequence alignment using the *E. coli* genome (GCF_000005845.2_ASM584v2) as reference was performed using R packages from bioconductor. Tables containing the raw transcripts were used as input for the normalization algorithm and significance tests from the DeSeq2 package, comparing replicates from the WT and mutant strain in the same experiment. Cut values were set to $p < 0.1$ with a base 2 logarithmic fold change different from 0. Genes passing these criteria were recognized as

differentially expressed. The accession number for the RNAseq data reported in this paper is GEO: GSE180830 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE180830).

Proteome estimation

The proteome estimation calculation, as reported by Lastiri-Pancardo et al. 2020, is based on the number of transcripts (r_i) and the corresponding protein copy number per cell ($C\,\text{cell}\,i$). It is assumed that the transcripts yield the proteome reported by Schmidt et al. in glucose. The translation efficiency rate (s_i) is calculated by

$$s_i = \frac{C\,cell\,i}{r_i}$$

The rate for each protein is then used to estimate the protein copies per cell (P_i) for the mutants and transform that to mass with

$$P_i = \frac{r_i \times s_i \times MW}{NA}$$

where P_i is a given protein i contribution in mass, r_i is the mutant's transcript abundance in FPKMs, MW is the protein's molecular weight and NA is Avogadro's number (6.022×10^{23}). For each protein, the difference between mutant and WT Pi is its contribution in mass.

Gene categorization

Differentially expressed genes from SQ53 and rpoB672 were categorized using available data from ecocyc (https://ecocyc.org/) and uniprot (https://www.uniprot.org/) and compared with previous categorization by Schmidt, 2016. Final categories were manually cured based on the most frequent annotations from available data sources. Genes with poor or unavailable information regarding function or associated sigma factor were considered in the hedging function group.

References

Asai, T., Zaporojets, D., Squires, C., and Squires, C.L. (1999a). An escherichia coli strain with all chromosomal rRNA operons inactivated: Complete exchange of rRNA genes between bacteria. Proc. Natl. Acad. Sci. U. S. A. *96*, 1971–1976.

Asai, T., Condon, C., Voulgaris, J., Zaporojets, D., Shen, B., Al-Omar, M., Squires, C., and Squires, C.L. (1999b). Construction and initial characterization of Escherichia coli strains with few or no intact chromosomal rRNA operons. J. Bacteriol. *181*, 3803–3809.

Basan, M., Honda, T., Christodoulou, D., Hörl, M., Chang, Y.F., Leoncini, E., Mukherjee, A., Okano, H., Taylor, B.R., Silverman, J.M., et al. (2020). A universal trade-off between growth and lag in fluctuating environments. Nature *584*, 470–474.

Baumstark, R., Hänzelmann, S., Tsuru, S., Schaerli, Y., Francesconi, M., Mancuso, F.M., Castelo, R., and Isalan, M. (2015). The propagation of perturbations in rewired bacterial gene networks. Nat. Commun. 2015 61 *6*, 1–11.

Ceroni, F., Algar, R., Stan, G.-B., and Ellis, T. (2015). Quantifying cellular capacity identifies gene expression designs with reduced burden. Nat. Methods *12*, 415–418.

Cheng, K.-K., Lee, B.-S., Masuda, T., Ito, T., Ikeda, K., Hirayama, A., Deng, L., Dong, J., Shimizu, K., Soga, T., et al. (2014a). Global metabolic network reorganization by adaptive mutations allows fast growth of Escherichia coli on glycerol. Nat. Commun. 2014 51 *5*, 1–9.

Condon, C.N., Liveris, D., Squires, C., Schwartz, I., and Squires, C.L. (1995). rRNA Operon Multiplicity in Escherichia coli and the Physiological Implications of rrn Inactivation. J. Bacteriol. *177*, 4152–4156.

Conrad, T.M., Frazier, M., Joyce, A.R., Cho, B.-K., Knight, E.M., Lewis, N.E., Landick, R., and Palsson, B.Ø. (2010). RNA polymerase mutants found through adaptive evolution reprogram Escherichia coli for optimal growth in minimal media. Proc. Natl. Acad. Sci. *107*, 20500–20505.

Dai, X., Zhu, M., Warren, M., Balakrishnan, R., Patsalo, V., Okano, H., Williamson, J.R., Fredrick, K., Wang, Y.-P., and Hwa, T. (2016). Reduction of translating ribosomes enables Escherichia coli to maintain elongation rates during slow growth.

Dennis, P.P., and Bremer, H. (2008). Modulation of Chemical Composition and Other

Parameters of the Cell at Different Exponential Growth Rates. EcoSal Plus *3*.

Dennis, P.P., Ehrenberg, M., and Bremer, H. (2004). Control of rRNA Synthesis in Escherichia coli: a Systems Biology Approach. Microbiol. Mol. Biol. Rev. *68*, 639–668.

Enjalbert, B., Cocaign-Bousquet, M., Portais, J.C., and Letisse, F. (2015). Acetate exposure determines the diauxic behavior of Escherichia coli during the glucose-acetate transition. J. Bacteriol. *197*, 3173–3181.

Gyorgy, A., Jiménez, J.I., Yazbek, J., Huang, H.-H., Chung, H., Weiss, R., and Del Vecchio, D. (2015). Isocost Lines Describe the Cellular Economy of Genetic Circuits. Biophys. J. *109*, 639–646.

Heckmann, D., Campeau, A., Lloyd, C.J., Phaneuf, P. V., Hefner, Y., Carrillo-Terrazas, M., Feist, A.M., Gonzalez, D.J., and Palsson, B.O. (2020). Kinetic profiling of metabolic specialists demonstrates stability and consistency of in vivo enzyme turnover numbers. Proc. Natl. Acad. Sci. U. S. A. *117*, 23182–23190.

Hidalgo, D., and Utrilla, J. (2019). Resource allocation principles and minimal cell design. In Minimal Cells: Design, Construction, Biotechnological Applications, (Springer International Publishing), pp. 211–230.

Kim, J., Darlington, A., Salvador, M., Utrilla, J., and Jiménez, J.I. (2020). Trade-offs between gene expression, growth and phenotypic diversity in microbial populations. Curr. Opin. Biotechnol. *62*, 29–37.

Kim, J., Darlington, A.P.S., Bates, D.G., and Jimenez, J.I. (2021). The interplay between growth rate and nutrient quality defines gene expression capacity Running Title: Growth rate and nutrient capacity define expression capacity. BioRxiv 2021.04.02.438188.

Kolmsee, T., Delic, D., Agyenim, T., Calles, C., and Wagner, R. (2011). Differential stringent control of Escherichia coli rRNA promoters: Effects of ppGpp, DksA and the initiating nucleotides. Microbiology *157*, 2871–2879.

Kurylo, C.M., Parks, M.M., Juette, M.F., Zinshteyn, B., Altman, R.B., Thibado, J.K., Vincent, C.T., and Blanchard, S.C. (2018). Endogenous rRNA Sequence Variation Can Regulate Stress Response Gene Expression and Phenotype. Cell Rep. *25*, 236-248.e6.

LaCroix, R. a, Sandberg, T.E., O'Brien, E.J., Utrilla, J., Ebrahim, A., Guzman, G.I., Szubin, R., Palsson, B.O., and Feist, A.M. (2015). Use of Adaptive Laboratory Evolution To Discover

Key Mutations Enabling Rapid Growth of Escherichia coli K-12 MG1655 on Glucose Minimal Medium. Appl. Environ. Microbiol. *81*, 17–30.

Lastiri-Pancardo, G., Mercado-Hernández, J.S., Kim, J., Jiménez, J.I., and Utrilla, J. (2020). A quantitative method for proteome reallocation using minimal regulatory interventions. Nat. Chem. Biol. *16*, 1026–1033.

Lynch, M., and Marinov, G.K. (2015). The bioenergetic costs of a gene. Proc. Natl. Acad. Sci. U. S. A. *112*, 15690–15695.

Maeda, M., Shimada, T., and Ishihama, A. (2015). Strength and Regulation of Seven rRNA Promoters in Escherichia coli. PLoS One *10*, e0144697.

Phaneuf, P. V, Gosting, D., Palsson, B.O., and Feist, A.M. (2019). ALEdb 1.0: a database of mutations from adaptive laboratory evolution experimentation. Nucleic Acids Res. *47*, D1164–D1171.

Quan, S., Skovgaard, O., McLaughlin, R.E., Buurman, E.T., and Squires, C.L. (2015). Markerless Escherichia coli rrn deletion strains for genetic determination of ribosomal binding sites. G3 Genes, Genomes, Genet. *5*, 2555–2557.

Roller, B.R.K., Stoddard, S.F., and Schmidt, T.M. (2016). Exploiting rRNA operon copy number to investigate bacterial reproductive strategies. Nat. Microbiol. *1*, 16160.

Sastry, A. V., Gao, Y., Szubin, R., Hefner, Y., Xu, S., Kim, D., Choudhary, K.S., Yang, L., King, Z.A., and Palsson, B.O. (2019). The Escherichia coli transcriptome mostly consists of independently regulated modules. Nat. Commun. *10*, 1–14.

Schmidt, A., Kochanowski, K., Vedelaar, S., Ahrné, E., Volkmer, B., Callipo, L., Knoops, K., Bauer, M., Aebersold, R., and Heinemann, M. (2016). The quantitative and condition-dependent Escherichia coli proteome. Nat. Biotechnol. *34*, 104–110.

Scott, M., Gunderson, C.W., Mateescu, E.M., Zhang, Z., and Hwa, T. (2010). Interdependence of Cell Growth and Gene Expression: Origins and Consequences. Science (80-.). *330*, 1099–1102.

Swain, P.S., Stevenson, K., Leary, A., Montano-Gutierrez, L.F., Clark, I.B.N., Vogel, J., and Pilizota, T. (2016). Inferring time derivatives including cell growth rates using Gaussian processes. Nat. Commun. *7*, 1–8.

Tan, J., Sastry, A. V, Fremming, K.S., Bjørn, S.P., Seo, S., Voldborg, B.G., and Palsson,

B.O. (2020). Independent component analysis of E. coli's transcriptome reveals the cellular processes that respond to heterologous gene expression. Metab. Eng.

Utrilla, J., O'Brien, E.J., Chen, K., McCloskey, D., Cheung, J., Wang, H., Armenta-Medina, D., Feist, A.M., and Palsson, B.O. (2016). Global Rebalancing of Cellular Resources by Pleiotropic Point Mutations Illustrates a Multi-scale Mechanism of Adaptive Evolution. Cell Syst. *2*, 260–271.

Wilson, D.N., and Nierhaus, K.H. (2007). The Weird and Wonderful World of Bacterial Ribosome Regulation. Crit. Rev. Biochem. Mol. Biol. *42*, 187–219.

Zaslaver, A., Bren, A., Ronen, M., Itzkovitz, S., Kikoin, I., Shavit, S., Liebermeister, W., Surette, M.G., and Alon, U. (2006). A comprehensive library of fluorescent transcriptional reporters for Escherichia coli. Nat. Methods *3*, 623–628.