EP300 (p300) mediated histone butyrylation is critical for adipogenesis

Aditya Bhattacharya [1], Sourav Chatterjee [1], Utsa Bhaduri [2], Akash Kumar Singh [1], Madavan Vasudevan [3], Koneni V Sashidhara [5], Rajdeep Guha [6], Aamir Nazir [7], Srikanta Kumar Rath [7], Nagashayana Natesh [8], Tapas K Kundu [1,4]

1. Transcription and Disease Laboratory, Molecular Biology and Genetics Unit, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore, India.
2. Chromatin Biology Laboratory, Molecular Biology and Genetics Unit, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore, India.
3. Theomics International Pvt Ltd, Bangalore
4. Division of Neuroscience and Aging Biology,CSIR-Central Drug Research Institute, Lucknow
5. Division of Medicinal and Process Chemistry, Central Drug Research Institute, Lucknow, India
6. Division of Laboratory Animal Facility, Central Drug Research Institute, Lucknow, India
7. Division of Toxicology and Experimental Medicine, Central Drug Research Institute, Lucknow, India
8. Central Government Health Scheme Wellness Centre Number 8, Domlur, Bangalore, India

E-mail ID of the corresponding author: tapas@jncasr.ac.in

Affiliations:

Aditya Bhattacharya – Ph.D. graduate student

Sourav Chatterjee – Post-doctoral researcher

Utsa Bhaduri – Research assistant

Madavan Vasudevan – Founder and CEO, Theomics International Pvt Ltd

Akash Kumar Singh - Integrated PhD student

Koneni V Sashidhara - Professor

Rajdeep Guha – Assistant Professor

Aamir Nazir- Principal Scientist

Srikanta Kumar Rath- Chief Scientist

Nagashayana Natesh – Senior physician

Tapas K Kundu - Professor

Abstract

Objective

The master epigenetic enzyme EP300 (p300) besides having lysine acetyltransferase activity can also catalyse other acylation modifications (propionylation, butyrylation, crotonylation etc.), the physiological implications of which are yet to be established fully. We hypothesized that p300 catalysed histone butyrylation may have a causal relationship with adipogenesis and the consequent obesity.

Methods

Histone butyrylation pattern was investigated in 3T3L1 cells upon adipogenesis by immunoblotting and chromatin immunoprecipitation experiments. A small molecule modulator that could specifically inhibit p300 catalysed butyrylation without affecting its canonical acetyltransferase activity was screened from a series of compounds and then administered in differentiating 3T3L1 adipocytes as well as high fat diet-induced and genetically obese mice to validate the importance of butyrylation in adipogenesis.

Results

Histone butyrylation was increased upon adipogenesis both globally and locally in the promoters of pro-adipogenic genes along with an upregulation in the expression of acyl CoA generating enzyme Acss2, knockdown of which led to reduced butyrylation. Treatment of differentiating 3T3L1 cells with the p300 specific butyrylation inhibitor LTK-14A led to

abrogation of adipogenesis with reduced expression of pro-adipogenic genes and inhibition of

H4K5 butyrylation. LTK-14A administration could also attenuate weight gain in both mice

models of obesity by preventing adipocyte hypertrophy via H4K5 butyrylation inhibition.

Conclusion

Our results indicate that p300 catalysed histone butyrylation may have a causal relationship

with the process of adipogenesis. Site specific inhibition of butyrylation could lead to

adipogenesis repression and hence this epigenetic modification could be targeted for obesity

treatment.

Keywords: Acetylation; adipogenesis; butyrylation; garcinol; p300

Highlights

●Histone butyrylation has been established as a new epigenetic signature in the context of

adipogenesis.

•To the best of our knowledge, this is the first report of a selective inhibitor of p300 catalysed histone

acylation (butyrylation) without affecting its canonical acetyltransferase activity.

•Specific inhibition of H4K5 butyrylation could be a possible mechanism for inhibiting adipogenesis and

hepatic steatosis leading to better control of obesity.

•LTK-14A class of molecule could be developed as anti-obesity therapeutics.

Introduction

Chromatin function is spatially and temporally regulated by several processes like incorporation of histone variants, remodeling of nucleosomes by ATP-dependent remodelers and histone post-translational modifications. Interestingly, epigenetics and the metabolic state of the cell are intimately intertwined because the enzymes catalyzing the modifications use co-factors derived from several metabolic reactions (1). In recent times, advancements made in high sensitivity mass spectrometry have led to the discovery of new acylation modifications that structurally resemble acetylation, but differ in hydrocarbon chain length, hydrophobicity and charge (2). Emerging evidence indicates that these modifications occur dynamically on histones and could either have distinct functions from acetylation, or, have a similar but more profound impact on chromatin de-compaction, thereby amplifying the transcriptional output (3). EP300 (p300) is the master acyltransferase that has been found to promiscuously acylate histones using diverse CoA derivatives (2).

Due to the rising prevalence of obesity in the modern-day society, there has been increased research on the molecular mechanisms behind its manifestation. At the core of obesity lies the adipose tissue which regulates energy homeostasis and metabolism. Hence, the developmental process of adipose tissue i.e. adipogenesis and its dysregulation needs to be studied for clinical management of obesity-related disorders. The process of adipogenesis is accompanied by a significant increase in global histone acetylation levels, along with a rise in expression of p300 and a concomitant decrease in the levels of several histone deacetylases (4). Regulation of gene expression is dictated by the fluctuating activities of histone acetyltransferases and histone deacetylases. Moreover, the importance of histone acetylation for adipogenesis has been further underlined by the finding that knockdown of ATP citrate lyase, an enzyme responsible for the regeneration of acetyl CoA in the cytoplasm and nucleus, leads to decrease in histone acetylation and also suppression of adipogenesis (5). Therefore, it can be assumed that p300

plays a crucial role in promoting the expression of adipogenesis related genes by acetylating the

histones in the corresponding gene regulatory regions, especially the promoters. Amongst the different

coenzyme A derivatives, butyric acid is an important short chain fatty acid and an intermediate of fatty

acid synthesis/breakdown processes. p300 has been known to carry out butyrylation of histones which

has been causally linked to the process of spermatogenesis (3).

In this study we report that histone butyrylation is critical in adipoogenesis, probed by the first known

acylation-specific small molecule inhibitor of p300. The newly discovered importance of histone

butyrylation in adipogenesis makes it an attractive target for small molecule compound mediated

inhibition of adipogenesis, thereby controlling obesity with lesser collateral damage.

Materials and Methods

Animals and treatment for obesity-related studies: Male C57BL/6J mice (8 weeks old) were bred in -

house and acclimatized for 1 week prior to experiments. Mice were maintained on a 12 hour light/dark

cycle and given ad libitum access to food and water. Experiments were carried out in accordance with

the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA)

guidelines and were approved by the Institutional Animal Ethics Committee of JNCASR

(201/GO/Re/S/2000/CPCSEA dated 1st June, 2000 Renewed in Aug 2015). The age-matched animals

were maintained on a standard laboratory chow diet (AMRUT   M/s Krishna Valley Agrotech ,

Maharashtra) or high fat diet (60 kcal% fat) (Research Diets, Cat. No. D12492) with or without LTK-14A

(50 mg/Kg body weight) and water. The compound was mixed with the high fat diet according to dosage

requirements in the laboratory every week. Body weight and food intake were measured twice every

week.

For the experiment on genetically obese mice, male db/db mice and db+/- mice (8-10 weeks old) bred

in-house were obtained and acclimatized for 1 week prior to experiments. Mice were maintained on a

12-hour light/dark cycle at 22+/-3 °C and a relative humidity of 55% and given ad libitum access to food

and water. The experiments were carried out in accordance with the Committee for the Purpose of

Control and Supervision of Experiments on Animals (CPCSEA) guidelines, Government of India and were

approved by the Institutional Animal Ethics Committee of CSIR-CDRI, Lucknow (IAEC/2020/115/Renew-

0/Sl.No.09). The age-matched animals were maintained on a standard laboratory pelleted diet (Altromin

International, Cat. No. 1320) and water. db/db mice were oral gavaged (50 mg/kg) with LTK14A in 0.5%

carboxymethyl cellulose, an equal volume of the vehicle was orally gavaged to control db/db mice for a

total period of 30 days. db+/- animals were kept as an untreated normal control group. The body

weights and feed intake was recorded every week throughout the experimental period.

Cell culture: The mammalian cell line 3T3L1 was obtained from ATCC (American Type Culture Collection)

and was grown in DMEM (Dulbecco's Modified Eagle's Medium). The growth medium was

supplemented with 10% FBS (Fetal Bovine Serum). The cell line was grown in 37°C incubator with 5%

CO_2 and 90% relative humidity.

Adipogenesis and Oil Red-O staining: 3T3L1 preadipocytes were grown to confluency and 24 hours

post-confluence, cells were incubated with a differentiation medium (growth medium containing 1μM

Dexamethasone, 0.5 mM IBMX and 1 μg/ml insulin)(Sigma Aldrich) for 2 days in the presence/absence

of LTK-14A. Then the cells were supplemented with a maintenance medium with/without LTK-14A

(growth medium with 1 μg/ml insulin only) every alternate day. On the 6th day post-induction of

differentiation the cells were fixed with 3.7% formaldehyde solution followed by staining with Oil Red-O

dye (Sigma-Aldrich) (prepared in 60% isopropanol) for 1 hour at room temperature. The incorporated

dye was extracted using isopropanol and absorbance values were determined spectrophotometrically at

510 nm to estimate total amount of lipid accumulation under different experimental conditions.

Acid extraction of histones and immunoblotting: 3T3L1 cells were washed with ice-cold PBS and then

resuspended in Triton extraction buffer (PBS containing 0.5% Triton X-100 (v/v) and 2 mM PMSF). The

cells were allowed to lyse on ice for 10 minutes, spun down, washed with PBS and the process was

repeated another time. Then the proteins were acid extracted with 0.2 N HCl for two hours on ice. The

debris was spun down and then histones were precipitated by incubating the supernatant with 33%

trichloroacetic acid (Sigma-Aldrich) at 4^{o}C for 30 minutes. The histones were then pelleted down and

then given two washes with acetone. The pellet was allowed to air-dry for 10 minutes and then

resuspended with 50 mM Tris-Cl, pH 7.4.   Extraction of histones from epididymal fat pads was

performed by mechanically homogenizing the adipose tissues in Triton extraction buffer and then

following the same subsequent steps as mentioned before, except for the step of acid extraction with

HCl, in which the incubation was performed for four hours on ice.

The samples were then run in 15% SDS-PAGE, transferred onto PVDF membrane (Milipore) in transfer

buffer (25 mM Tris, 192 mM glycine, 0.036% SDS in 20% methanol), blocked with 5% skimmed milk

solution to prevent non-specific antibody binding and finally probed with antibodies against acetylation

and butyrylation marks prepared in 1% skimmed milk or FBS solution (5% BSA, 1x TBS, 0.1% Tween-20).

HRP-conjugated secondary antibodies (Abcam) were used against the primary antibodies to obtain a

chemiluminiscent signal in the presence of SuperSignal West Pico Chemiluminiscent Substrate (Thermo

Scientific) which was developed using the VersaDoc imaging system (Biorad).

Synthesis of LTK14A: Isolation of garcinol from Garcinia indica and synthesis of isogarcinol from it were

carried out following the protocols as mentioned in Mantelingu et al, (2007) (11). For LTK-14A synthesis,

anhydrous potassium carbonate (0.41 mmol) was added to a stirred solution of isogarcinol (100 mg, 0.16

mmol) in dry acetone. The mixture was stirred for 1 hour at room temperature followed by dropwise

addition of dimethyl sulfate (0.41 mmol) under a nitrogen gas atmosphere. The resulting reaction

mixture was further stirred at room temperature for 14 hours. Progress of the reaction was monitored

through thin layer chromatography. Acetone was evaporated from the reaction mass, ice was added and

acidified with 5% aqueous HCl (pH ~ 5-6). A pale brown precipitate was observed, which was filtered and

washed with cold water. The dry mass was then chromatographed on silica gel using 5-6% ethyl acetate

in hexane as an eluting solvent mixture. 720 mg (69%) LTK14A was isolated as a white solid. NMR

spectral analysis was consistent with the desired product (Sup. Fig.3A). A 20 mg sample was dissolved in

hot methanol and placed in a flat surface glass vial with access to very slow solvent evaporation. Crystals

as colorless needles were isolated. The purity of the synthesized LTK-14A was found to be more than

98% as verified by HPLC (Sup. Fig. 3C) and only the protonated species of LTK-14A were observed in the

ESI spectra of the corresponding fraction (Sup. Fig. 3B).

X-ray crystal structure determination for LTK14A (CCDC: 1969185): Suitable single crystal with

approximate dimensions of 0.35 × 0.18 × 0.15 mm3 was used for X-ray diffraction analyses by mounting

on the tip of a glass fibre in air. Data were collected on a Bruker kappa apex 2 with Mo Kα (λ=0.71073 Å)

at 296 (2) K. The structure was solved by direct method using program SHELXL-97 and subsequent Fast

Fourier Transform technique. Crystallographic data and experimental details for LTK14A are summarized

in Table 1 and Sup. Fig. 3D.

Chromatin Immunoprecipitation: Nuclear ChIP was performed using intact nuclei from cells after

removal of lipids. 3T3L1 that were either undifferentiated or differentiated for 3 to 6 days were cross-

linked using 1% formaldehyde. Nuclear fractions were obtained by sequentially lysing the cells in two

buffers- lysis buffer 1 (140 mM NaCl, 1 mM EDTA, 50 mM HEPES, 10% Glycerol, 0.5% NP-40, 0.25%

Triton X-100) and lysis buffer 2 (10 mM Tris pH 8, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA). This was

followed by lysis in SDS lysis buffer (1% SDS, 10 mM EDTA, 50 mM Tris-HCl pH 8). Lysates were sonicated

in a Diagenode Bioruptor (Liège, Belgium) to produce 200 to 1000 bp DNA fragments. The cell lysates

were incubated with H4K5 Bu antibody (PTM Biolabs) or preimmune IgG per sample and 25 µl of

magnetic protein G Dynabeads (catalog no. 10003D; Novex, Waltham, MA) overnight at 4°C. Beads were

then washed successively with low-salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl

pH 8, and 150 mM NaCl), high-salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8,

and 500 mM NaCl), LiCl buffer (250 mM LiCl, 1% NP-40, 1% NaDOC, 1 mM EDTA, and 10 mM Tris-HCl, pH

8), and TE (10 mM Tris-HCl pH 8 and 1 mM EDTA). DNA-protein complexes were recovered from beads

in elution buffer (0.1% SDS and 100 mM NaHCO3). The ChIP eluates and input samples in the elution

buffer were then reverse cross-linked by adding 200 mM NaCl and 20 µg proteinase K (Sigma) and

incubating at 65°C for 4 h. Subsequently, 20 µg of RNase A (Sigma) were added and the samples were

further incubated for 15 min at 37°C. The immunoprecipitated DNA was extracted using phenol-

chloroform, ethanol precipitated, and used for quantitative PCR. The region-specific primer sets used for

the ChIP-qPCR analysis have been mentioned in Table 3. These primers were designed based on the

genomic coordinates with enriched H3K27 acetylation in the vicinity of promoters of candidate genes in

brown adipose tissue, the information for which was obtained from NCBI37/mm9 assembly

(ENCODE/LICR).

Immunohistochemistry: Collected liver tissue samples were stored in 4% paraformaldehyde for 24

hours after which they were cryoprotected in 30% sucrose solution for two weeks. Cryosections were

performed at 7 µm sections using Cryostat LeicaCM1850 UV. For staining, the tissue sections were

washed with PBS followed by antigen retrieval with 0.01 M citrate buffer (pH 6). The tissues were then

permeabilised with 0.3% Triton X-100/PBS (PBST) and blocked with 2% serum followed by incubation

with primary antibodies overnight at 4°C. The next day, secondary antibody incubation was carried out

for one hour at room temperature followed by staining of the nuclei with Hoechst 33342 and mounting

with 70% glycerol. Images were acquired by confocal microscopy. For quantitation, the intensity of each

modification-specific staining was normalized with respect to the Hoechst staining intensity.

Reagents and antibodies: The media used for cell culture was purchased from Sigma-Aldrich; DMEM

(Ref: 1152). Several other cell culture reagents were obtained from HIMEDIA, such as PBS (Ref: TL1006),

Trypsin-EDTA solution 10X (Ref: TCL070) and Antibiotic antimyotic solution 100X (Ref: A002A). Multiple

reagents used for cell culture were obtained from Sigma-Aldrich: Insulin (Ref: I6634), Dexamethasone

(Ref: D4902), IBMX (Ref: I5879). TRIzol reagent was obtained from Ambion life technologies (Ref:

15596018). The commercial antibodies used in this study are Pan anti-butyryllysine (PTM Biolabs, SKU:

PTM 301), Butyryl-Histone H4 (Lys 12) rabbit pAb (PTM Biolabs, SKU: PTM 308), Butyryl-Histone H3 (Lys

9) rabbit pAb (PTM Biolabs, SKU: PTM 305), Butyryl-Histone H3 (Lys 23) mouse mAb (PTM Biolabs, SKU:

PTM 307), Butyryl-Histone H4 (Lys 8) rabbit pAb (PTM Biolabs, SKU: PTM 311), Butyryl-Histone H4 (Lys 5)

rabbit pAb (PTM Biolabs, SKU: PTM 313), Butyryl-Histone H3 (Lys 27) rabbit pAb (PTM Biolabs, SKU: PTM

315).

Statistical analysis: All statistical analyses were performed using GraphPad Prism 7 Software (California,

USA). Data obtained from two or three individual experiments as mentioned in figure legends, were

expressed as mean +/- SD. Either two-tailed unpaired Student's t-test or one-way ANOVA with

Dunnett's/Bonferroni's multiple comparision was used to determine the statistical significance values.

The mutagenecity analysis data were analyzed by comparing the groups using one-way ANOVA followed

by Neman-Keuls test. Body weight data for acute toxicity test were summarized in Mean ± SD and

compared by two factor (Group and Period) repeated measure analysis of variance (RM ANOVA) and the

significance of mean difference within (intra) and between (inter) the groups was done by Newman-

Keuls post hoc test. A P-value of equal to or less than 0.05 was considered statistically significant.

Acknowledgements: TKK is a recipient of JC Bose fellowship from the Department of Science and

Technology, Govt. of India (SR/S2/JCB-28/2010). AB is a senior research fellow of UGC. This work was

supported by Life Science Education & Training at JNCASR grant (DBT/INF/22/SP27679/2018). We

acknowledge Dr. Rahul Gajbhiye from NIPER, Kolkata for the assistance in mass spectrometry-related

experiments in their central instrument facility. We thank Dr. R.G. Prakash for providing the necessary

facilities to perform the mice experiment in Animal Facility, JNCASR and the confocal facility, JNCASR for

the immunofluorescence imaging analysis. We also thank Kruthi HT for helping in the preparation of the

figures and graphical abstract. We further acknowledge the Sophisticated Analytical Instrument Facility

and Research (SAIFR, Lucknow) for providing spectral data. This manuscript bears CDRI communication

number 2021/TK.

Conflict of interest: Authors declare no conflict of interest.

Author Contributions: AB and TKK conceived the project. AB designed and performed all the biological

experiments. SC performed the initial synthesis of all compounds and molecular docking analysis. KVS

performed bulk scale synthesis of LTK-14A for animal experiments. AKS and RG assisted in the animal

experiments. UB and MV performed *in silico* analysis of high throughput RNA sequencing data. AN did

the toxicity evaluation and SKR evaluated the mutagenecity. NN provided the raw material *Garcinia*

indica for compound isolation. AB and TKK analyzed the data and wrote the manuscript.

Results

Physiological levels of butyryl CoA and histone butyrylation increase during adipogenesis:

Under normal circumstances, the cellular levels of specific coenzyme A derivatives are much less than

acetyl CoA (6).We surmised that the situation might alter in particular cases where metabolic pathways

promoting the production of other acyl CoA derivatives become more predominant. One such possible

scenario could be the process of adipogenesis in which fatty acid synthesis and breakdown can give rise

to several intermediate products like butyryl CoA.

The causal relationship between butyryl CoA and adipogenesis was investigated using the 3T3L1 pre-

adipocyte cell line as a model system (fig 1A). Our RT-qPCR data showed a significant increase in the

expression of *Acss2* (acyl-CoA synthetase short-chain family member 2) upon differentiation (fig 1B).

Previously, ACSS2 had been established as the enzyme responsible for the generation of acyl-CoA

derivatives from their corresponding short chain fatty acids (7). Indeed, siRNA mediated knockdown of

Acss2 (fig 1C) led to a drastic reduction in histone butyrylation with a relatively lesser effect on

acetylation (fig 1D). This indicated that the ACSS2 generated butyryl CoA might be playing an important

physiological role in adipogenesis. These findings prompted us to investigate whether adipocyte

differentiation also led to an increased butyrylation of proteins. We observed that in accordance with

the previous report (4), there was an increase in acetylation level at histone H3 lysine 9 (H3K9 Ac) in the

differentiated cells as compared to the undifferentiated cells (Fig. 1C, panel i). Interestingly

immunoblotting with pan butyryllysine and site-specific antibodies (H3K9, H3K23, H3K27, H4K5, H4K8

and H4K12) against butyrylation (Fig 1C, panel ii and iii) revealed that butyrylation at each of these sites

also increased upon differentiation. Chromatin immunoprecipitation with antibody against H4K5

butyrylation revealed that the modification enrichment levels increased in promoters of different pro-

adipogenic genes in a time dependant manner. *Pparg*, the master regulator of adipogenesis, exhibited a

steady increase in butyrylation level in its promoter with the progress of differentiation (fig 1F, panel i)

while butyrylation increased only at the terminal stage of adipogenesis in the promoter of *Lep*, encoding

the adipocyte marker leptin which gets expressed in mature adipocytes (fig 1F, panel ii). Interestingly,

the promoter of *Cebpd*, which regulates *Pparg* expression did not show any enhanced butyrylation in its

promoter on day 3 and instead, exhibited increased butyrylation on day 6 of adipogenesis (fig 1F, panel

iii).   Collectively these results showed that during adipogenesis, EP300 catalysed butyrylation of

histones could be an important epigenetic signature.

Identification of a small molecule that selectively inhibits p300 catalysed histone butyrylation *in vitro*:

In order to investigate the contribution of p300 catalyzed butyrylation per se in adipogenesis, we

adopted a chemical biology-based approach. For this purpose, a series of derivatives of garcinol, a

polyisoprenylated benzophenone isolated from *Garcinia indica*, were synthesized (8). These derivatives,

unlike garcinol, were found to be non-toxic to cells and showed a high specificity toward the catalytic

activity inhibition of EP300. The derivatives were generated by internal cyclisation of garcinol with

further addition of one or two functional groups to the remaining free -OH groups- giving rise to mono

and disubstituted derivatives (Sup. fig. 1A). Amongst these, a few monosubstituted derivatives could

potently inhibit EP300 catalysed acetylation as well as butyrylation, while their disubstituted

counterparts were unable to inhibit acetylation (Sup. fig. 2 B, C and D). This was probably due to

increased steric hindrance brought about by the addition of an extra functional group that interfered

with the interaction between the compounds and EP300.

An inverse correlation between catalytic rates of acylation reactions by EP300 and increasing acyl chain

length had already been established previously (9). Based on this, we hypothesized since butyrylation is

catalysed by EP300 at a much slower rate than acetylation, one of the compounds that could not inhibit

EP300 catalysed acetylation might be able to inhibit the slower butyrylation reaction. Our *in vitro*

butyrylation assay using recombinant p300 and histone H3 as the substrate revealed a few potential

candidates (Sup. Fig. 2 E) as butyrylation inhibitors. Amongst them, LTK-14A, which is 13, 14 dimethoxy

isogarcinol, was capable of inhibiting full-length EP300 catalysed butyrylation at a concentration of 25

µM (fig. 2 C and D), with minimal effect on EP300 catalysed acetylation in similar reaction conditions (fig.

2 A and B).

In order to understand the underlying mechanism of the selective inhibition by LTK-14A, we performed

molecular docking analysis of LTK-14A (Fig. 2 G) and the catalytic domain. To delineate this, we solved

the crystal structure of LTK-14A (Table 1, Sup. Fig.2A). LTK-14 (fig. 2E), a specific inhibitor of EP300

catalysed acetylation and butyrylation (Sup. Fig. 2 B and C) was used for further comparative analysis.

The energy minimized analysis revealed that both LTK-14 and LTK-14A could potentially bind with the

active site of the EP300 catalytic domain forming several hydrophobic interactions (Fig.2 F and H). LTK-

14, due to the presence of a free phenolic –OH group, could hydrogen bond with a carbonyl oxygen

atom present in the peptide backbone of S1396 residue. However, LTK-14A has all of its phenolic –OH

groups blocked and so it is incapable of forming such an interaction. This hydrogen bonding by LTK-14

could imply a stronger binding to the enzyme, while LTK-14A, by interacting with fewer residues

presumably inhibits only the slower butyrylation reaction and the interaction may not be strong enough

to deter the faster acetylation reaction.

Some of the residues with which LTK-14A could potentially form hydrophobic interaction have already

been reported before as being critical for targeting inhibition of the enzymatic activity of EP300 by the

first reported EP300 inhibitor lysyl CoA (10). These residues include C1438, Y1446, Y1397 and W1436. To

verify this observation we generated a point mutant of p300 catalytic domain in which cysteine 1438,

showing the highest propensity to interact with LTK-14A, was mutated to alanine. We found that the

C1438A catalytic domain is equally active as the wild-type catalytic domain in an *in vitro* lysine

acetyltransferase assay (Sup. Fig. 2B). We observed that the wild-type EP300 catalytic domain catalysed

butyrylation was inhibited by LTK-14A most prominently at 25 µM concentration while acetylation was

not inhibited (compare lane 6 in each of fig.2 I-L). In contrast, neither acetylation nor butyrylation

modification catalysed by the mutant EP300 catalytic domain could be inhibited by LTK-14A even at 25

µM concentration (compare lane 6 in each of fig. 2 M-P). Altogether, these data suggest that LTK-14A

indeed interacts with C1438 within the active site of the EP300 catalytic domain, and this residue is

critical for its interaction to bring about butyrylation reaction inhibition.

LTK-14A is a metabolically stable, cell-permeable compound that significantly inhibits adipogenesis:

Upon establishing that LTK-14A is an EP300 acylation-specific inhibitor *in vitro*, its cellular effects were

tested using 3T3L1 cells. The quality of our compound used in all biological experiments was acceptably

good with more than 97% purity (Sup. Fig. 2 C, D and E). We found the LTK14A is highly cell-permeable

and metabolically stable as it could be detected by mass spectrometry analysis from metabolites

extracted from LTK-14A treated cells after six days (Fig. 3A). MTT assay showed that LTK-14A at 25 µM

concentration did not have any adverse effect on the cellular oxidoreductase activity, a signature of

metabolic output of cells, (Fig. 3B) suggesting that LTK-14A is non-toxic to 3T3L1 even after 6 days of

treatment. Oil Red-O staining of 3T3L1 cells differentiated in the presence of LTK-14A followed by

spectrophotometric quantitaion revealed significant inhibition of adipogenesis by the compound (Fig.

3C, D).

In order to understand the transcriptional mechanism of LTK14A-mediated butyrylation inhibition on

adipogenesis, adipogenic cells were treated with LTK-14A and subjected to transcriptome analysis. We

used untreated and solvent treated cells as control. Volcano plot (Fig. 3E) and heat map analysis (Fig. 3F)

revealed that upon treatment with the compound, a large number of genes were differentially

expressed compared to DMSO treated and untreated conditions. To better understand the LTK-14A

affected genes, we focused on the common genes that were differentially expressed in the two cases-

(1) LTK-14A versus no inhibitor and (2) LTK-14A versus DMSO. Among these common genes, 2200 genes

were found to be downregulated and 1309 genes were upregulated (Fig. 4A) with a majority of them to

be protein-coding genes (Fig. 4B). Gene ontology analysis revealed that LTK14A treatment

predominantly affected the genes involved in lipid metabolism pathways (Fig. 4C, E) which were

minimally affected by DMSO alone (Fig. 4D). The gene regulatory networks and the respective pathways

were constructed with the major genes that were affected by LTK-14A treatment compared to DMSO

treated and untreated conditions (Fig. 4F, Sup. Fig.3A). They collectively depict how LTK-14A, by

inhibiting the expression of a large number of genes that are interconnected via several pathways

associated with lipid metabolism, brings about a collapse in the adipogenesis process as a whole.

A few of the pro-adipogenic genes downregulated by LTK-14A, as indicated by RNA sequencing, were

selected for further validation (Sup. Fig. 3B). RT-qPCR analysis corroborated that except for *Cebpd*, most

of the other pro-adipogenic genes (*Pparg, Cebpa, Adipoq, Lep, Slc2a4, Fasn* and *Fabp4*) showed a

marked downregulation upon LTK-14A treatment (Fig. 4G). *Acss2* was also downregulated by LTK-14A

(Sup. Fig.4C). The upregulation of anti-adipogenic genes was also verified by performing RT-qPCR

analysis for two such candidate genes- *Kdm4c* and *Tead4* (Sup. Fig.3C). Collectively, the RNA sequencing

analysis and the validation of gene targets further convincingly suggested that LTK-14A can inhibit

adipogenesis in 3T3L1 cells, mainly by repressing the expression of important pro-adipogenic genes.

LTK-14A inhibits histone butyrylation in 3T3L1 cells, without affecting acetylation or other post-

translational modifications:

Furthermore we wanted to find out whether LTK-14A mediated downregulation of adipogenesis is

causally linked to the inhibition of histone butyrylation in the 3T3L1 cells. We observed that treatment

with LTK-14A inhibited overall butyrylation of histones (Fig. 5 A, B). Among the specific modification sites

tested, H3K23 and H4K5 butyrylation showed a significant decrease upon treatment with LTK-14A (Fig. 5

A, B) while butyrylation at other tested sites was affected minimally (Sup. Fig. 3D). Acetylation of

histones (H3K9, H3K18 and H4K12) was also found to be unaffected (Fig. 5 C, D) upon LTK-14A

treatment. To establish the specificity of the LTK14A, the effect of LTK-14A on lysine and arginine

methylation along with phosphorylation of histones was also investigated. We observed (Fig. 5 E, F) no

significant changes in these marks upon the compound treatment, thereby indicating that in the 3T3L1

cells, LTK-14A specifically inhibits butyrylation.

Inhibition of butyrylation attenuates weight gain of high fat diet fed C57BL6/J mice:

After observing the anti-adipogenic effect of LTK-14A on 3T3L1 cells, we investigated whether this

compound could also affect adipogenesis and obesity at an organismal level. For this purpose, C57BL6/J

mice were maintained either on a normal diet or a high fat diet with or without LTK-14A. In due course

of time it was observed that the mice maintained on a high fat diet along with LTK-14A had a weight

gaining trend intermediate between the normal diet and high fat diet fed mice (Fig.6A,B). Dissection of

the mice revealed that the fat accumulation in high fat diet-fed mice was the highest, followed by those

on high fat diet and LTK-14A, and then by those on normal diet (Fig. 6C). The epididymal fat pads were

separately weighed and again it was observed that the weights of the fat pads from compound treated

mice were much less than that of high fat diet-fed mice (Fig. 6 D, E). The diet consumption pattern of the

mice showed that the two groups on high fat diet with or without LTK-14A, had a similar consumption

rate indicating that the reduced weight gain of the compound treated mice is not due to discrepancy in

diet consumption (Table 2). Hematoxylin and eosin staining of adipose tissue and liver (Fig. 6 F,G)

showed signs of adipocyte hypertrophy and hepatic steatosis in high fat diet-fed mice, which were

absent in the mice maintained on normal diet or high fat diet with LTK-14A.

Immunofluorescence staining of the liver tissue samples revealed that there was a significant increase in

H4K5 butyrylation levels in high fat diet-fed mice compared to normal diet-fed mice in which the

modification was barely detectable and it was almost completely inhibited upon LTK-14A treatment (Fig.

6H panel iii, Fig. 6I panel iii). On the other hand, no significant difference in H4K12 (Fig. 6H panel i, Fig. 6I

panel i) and H3K14 acetylation (Fig. 6H panel ii, Fig. 6I panel ii) levels could be observed across all the

three groups. Furthermore, histone modification pattern in the adipose tissues was studied with acid

extracted histones from the epididymal fat pads. H4K5 butyrylation was reduced in the LTK-14A treated

samples compared to those of high fat diet-fed mice (Fig. 6 J, N) while H3K14, H3K18 and H4K12

acetylation levels were relatively unchanged across all the three groups (Fig. 6 J, K, L, M). The metabolic

stability of LTK-14A in mice was verified by its detection in ESI spectra of analytes extracted from the

liver of compound treated mice (Sup. Fig 3 E).

The butyrylation specific inhibitor reduces the weight of genetically obese db/db mice:

The previous model system involved the use of LTK-14A as a prophylactic. To test whether LTK-14A

could also reduce obesity by inhibiting histone butyrylation, genetically obese homozygous mutant

db/db mice were used. Heterozygous mutant db+/- mice were taken as control. The hyperphagic db/db

mice were orally administered LTK-14A daily at the same dosage as before (50 mg/Kg body weight). We

observed that upon oral gavaging of LTK-14A, the overweight mice not only showed an arrest in weight

gain, but their body weights were significantly reduced by the end of one month of treatment (Fig. 7 A,

B, C). The food intake of LTK-14A treated mice was marginally less than that of the vehicle-treated mice

(Table 2, Supplementary Information), although both groups consumed more food compared to the

db+/- mice during the treatment regime. The average weight of the liver (Fig. 7 D,E) and epididymal fat

pads (Fig. 7 F,G) was the highest for db/db mice given vehicle control and least for the db+/- mice, while

that for the compound treated db/db mice was intermediate. Hematoxylin and eosin staining of liver

and adipose tissue(Fig. 7 H,I) showed signs of adipocyte hypertrophy and hepatocyte ballooning in

vehicle-treated db/db mice, which were absent in the db+/- and LTK-14A treated db/db mice.

Similar to the high fat diet-induced obesity model, the more obese homozygous leptin receptor mutant

mice (db/db) liver had a greater level of H4K5 butyrylation compared to leaner heterozygous mutant

mice (db +/-) (Fig 7J panel iii, Fig. 7K panel iii). Moreover, LTK-14A treated db/db mice liver showed a

decrease in H4K5 butyrylation compared to vehicle control-treated mice liver. In contrast, H4K12 (Fig. 7J

panel i, Fig. 7K panel i) and H3K14 acetylation (Fig. 7J panel ii, Fig. 7K panel ii) were only partially

upregulated in the db/db mice and did not get significantly reduced upon LTK-14A treatment.

Immunoblotting with acid extracted histones from epididymal fat pads showed that H4K5 butyrylation

was reduced in the adipose tissues of LTK-14A treated mice (Fig. 7 L,P) compared to vehicle control

treated obese mice. H3K14, H3K18 and H4K12 acetylation were not significantly altered in any of the

three groups (Fig. 7 L,M,N,O). Finally, RT-qPCR analysis showed that just like in cellular model of

adipogenesis, LTK-14A treatment led to robust inhibition of important pro-adipogenic genes *Pparg*, *Lep*,

Slc2a4 and *Adipoq* in the fat pads of compound treated db/db mice (Fig. 7Q). Collectively these data

establish that obesity can not only be prevented but also reverted by specific inhibition of H4K5

butyrylation, thereby highlighting the importance of this modification in the context of adipogenesis.

LTK-14A is non-mutagenic and non-toxic in nature for biological applications: Based on the

observations of obesity attenuating effects of LTK-14A in two different mice models, we performed

some preclinical tests to ensure that the obesity arrest was not a result of any deleterious side effect of

toxicity from the compound. Plate incorporation assay was carried out in four different nutritional

auxotrophic mutants of Salmonella typhimurium (TA97a, TA98, TA100 & TA102) in the absence or

presence of rat liver extract (S9 enzyme fraction). It was observed that the number of revertants

obtained from LTK-14A was similar to that from vehicle control and far less compared to that from the

positive control in all four strains, both in the absence and presence of liver extract (Sup. Fig 4 A-H).

Furthermore, acute toxicity test was performed on rats for checking whether a single administration of

excessively high dose of LTK-14A would alter the body weight and food/water consumption pattern of

the rats. The body weight changes in the LTK-14A administered rats were similar to the vehicle control-

treated ones (Table 6) with no gross morphological changes in any organ. Moreover, there was also no

significant treatment-related effect on the food and water consumption of rats of both gender for all

three doses compared to vehicle control (Table 6).

Collectively these data imply that LTK-14A does not have deleterious toxic effects on the test animals

and could have a potential for downstream applications for tackling obesity-related complications.

Discussion

Reports on the physiological relevance of acylation modifications are scanty. Breakthroughs have been

made in establishing the importance of butyrylation in spermatogenesis (5), β-hydroxybutyrylation in

starvation response (11) and crotonylation in inflammatory response, digestive system, cancer and HIV

latency (7,12,13,14). Lysine propionylation has also been reported as an activation mark on histones

that could be useful in various physiological processes (15). The list of these modifications seems to be

increasing continuously, with the recent additions of histone serotonylation (16), lactylation (17) and

dopaminylation (18). The fact that serotonin and dopamine are chemical neurotransmitters while

lactate is derived from pyruvate, a glycolytic end product, and all these molecules bring about a

modification of histones thereby mediating gene expression in several tissues, bear testimony to the

hypothesis that the abundance levels of cellular metabolites in different physiological contexts can lead

to modification of histones with a plethora of diverse chemical moieties that can selectively modulate

gene expression patterns.

Our initial experiments in 3T3L1 adipogenesis indicated an important role of ACSS2 in the differentiation

process by generating butyryl CoA for histone butyrylation. Inside the cells ACL (ATP citrate lyase) and

ACSS2 have been implicated in the generation of nuclear acetyl CoA. Both the enzymes are known for

regulating gene expression in different contexts by acetylating histones in specific genomic loci (19-21).

Knockdown of *Acss2* had a much more pronounced effect on H4K5 butyrylation compared to H3K18

acetylation. This is probably due to the fact that ACSS2 is the sole/major producer of acyl CoA species

while acetyl CoA can be generated by diverse means and the knockdown of a single acetyl CoA producer

was not enough to have a robust effect on global acetylation level of histones. Similar observations have

already been made for histone crotonylation in HeLa cells (7).

Due to redundancy in the functionality of p300 catalyzed butyrylation with acetylation, the significance

of butyrylation in adipogenesis was demonstrated using a pharmacological approach. LTK-14A could

selectively and specifically inhibit histone butyrylation and still inhibit adipogenesis by downregulating

the expression of many pro-adipogenic genes. However amongst these genes *CEBPd* expression did not

get affected by LTK-14A possibly because the expression of this gene is less prone to be regulated by

histone butyrylation in its promoter. Indeed, our chromatin immunoprecipitation study indicated that in

the initial stage of induction of differentiation in pre-adipocytes, there was no enrichment of H4K5

butyrylation in the promoter of *Cebpd*. This could explain why the butyrylation inhibitor LTK-14A

treatment at the induction stage did not inhibit *Cebpd* expression, since there was no buyrylation mark

enrichment for inhibition in its promoter to begin with. An increased butyrylation in its promoter at the

terminal stage could be either through p300 catalysed modification or non-enzymatic means with

greater availability of butyryl CoA, or both. Currently, we have not been able to verify this. However our

preliminary observations with promoter-specific studies indicate that H4K5 butyrylation plays a role in

regulating the expression of the critical transcription factor *Pparg*. Inhibition of butyrylation by LTK-14A

repressed the expression of the master regulator which further led to the inhibition of downstream

genes like *Lep*, whose expression is dependent on PPARγ. While cell line-based studies indicate that

C/EBPδ regulates the expression of a cascade of transcription factors like C/EBPα and PPARγ, this fact

has been questioned in knockout mice which have poorly differentiated adipose tissue but still express

Pparg and *Cebpa* (22). This indicates that there could be other factors playing a redundant function like

C/EBPδ in the initiation stage of adipogenesis. Moreover, an additional function of C/EBPδ may seem to

be downstream of C/EBPα and PPARγ at the terminal stage of adipogenesis. It has been speculated that

this could involve inducing the expression of ligands that activate PPARγ (23). Therefore, the increased

butyrylation in the promoter of *Cebpd* on day 6 of adipogenesis might be responsible for its role in the

terminal stage. The expression of *Cebpd* is independent of *Pparg* and hence repression of *Pparg* by

butyrylation inhibition with LTK-14A did not repress *Cebpd*.

Acss2, which encodes the enzyme responsible for butyryl CoA biosynthesis was also downregulated by

LTK-14A which might have further contributed to reduced butyrylation of histones during inhibition of

adipogenesis. Amongst the different histone modification patterns investigated, H4K5 butyrylation was

consistently affected by LTK-14A in different cell line and mice models, while the other site specific

butyrylation marks were relatively unaltered, indicating that there may be a hierarchy in the importance

of these modifications in the regulation of adipogenesis. While our work was in progress, a new report

was published in which it was demonstrated that inhibition of fatty acid synthesis as well as oxidation

leads to a reduction in H4K5 butyrylation while acetylation remains unaffected (24). This further

reinforces our own hypothesis that fatty acid metabolism is an important source for certain acyl CoAs

such as butyryl CoA, derived from the short chain fatty acid butyric acid. Another work published during

this period revealed that HBO1, a member of the MYST family of acetyltransferase, also possesses the

ability to catalyse acylation reactions including butyrylation (25). Mass spectrometry studies of histone

modifications altered by knockdown of HBO1 in HeLa cells showed that H4K5 butyrylation does not get

affected by depletion of HBO1 indicating that it is a p300 specific mark. LTK-14, the monomethoxy

derivative of isogarcinol was found to be a specific inhibitor of p300 acetyltransferase activity (8)

without affecting PCAF. Since its disubstituted counterpart LTK-14A was found to be relatively inert with

respect to LTK-14, it seems plausible that LTK-14A would be unlikely to affect other epigenetic

modifications besides p300 catalysed acylation.

Global histone acetylation has previously been reported to remain unchanged in liver and adipose

tissues of mice even after high fat diet consumption (26), which was consistent with the observations we

made in both mice models studied. Since acetyl CoA is highly abundant, fluctuations in its levels did not

lead to major changes in histone acetylation patterns. Because the usual butyryl CoA levels are far less

than their corresponding Michaelis Menten constant for p300, an increase in butyryl CoA levels due to

excess lipid consumption through high fat diet or *de novo* lipid synthesis through excess food

consumption, resulted in a concomitant increase in butyrylation by p300 which served as the nutritional

sensor for butyryl CoA. Enhanced H4K5 butyrylation was not observed in mature adipose tissues of

obese mice possibly due to the predominance of adipocyte population in the tissues of mice from all the

groups (27).

Conclusion

In the 21st century, obesity has become a major epidemic leading to several other disorders such as type 2 diabetes and arteriosclerosis. Histone acetylation inhibition could be regarded as a possible method for attenuating adipogenesis and obesity. But acetylation is a widespread phenomenon that is required in several other physiological processes besides adipogenesis. Hence, treatment with an acetyltransferase inhibitor at a particular range of concentration could have undesirable side effects and toxicity. In contrast, butyrylation is a rarer event, repressing which may be less harmful to other processes. Indeed, our preclinical tests are indicative of the fact that LTK-14A may be suitable for development as a possible anti-obesity drug due to its negligible toxicity side effects. The present finding thus establishes the importance of histone butyrylation in adipogenesis, making it an attractive target for pharmacological inhibition of adipogenesis and thereby controlling obesity (Graphical abstract).

References

1. Haws, S. A., Leech, C. M., Denu, J. M. Metabolism and the Epigenome: A Dynamic Relationship. Trends Biochem 2020; 45, 731-747

2. Chen, Y., Sprung, R., Tang, Y., Ball, H., Sangras, B., Kim et al Lysine propionylation and butyrylation are novel post-translational modifications in histones. Mol Cell Proteomics 2007; **6**: 812-819

3. Goudarzi, A., Zhang, D., Huang, H., Barral, S., Kwon, O. K., Qi, S. et al Dynamic Competing Histone H4 K5K8 Acetylation and Butyrylation Are Hallmarks of Highly Active Gene Promoters. Mol Cell 2016;**62**:169-180

4. Yoo, E. J., Chung, J. J., Choe, S. S., Kim, K. H., and Kim, J. B. Down-regulation of histone deacetylases stimulates adipocyte differentiation. J Biol Chem 2006; **281**:6608-6615

5. Wellen, K. E., Hatzivassiliou, G., Sachdeva, U. M., Bui, T. V., Cross, J. R., and Thompson, C. B. ATP-

citrate lyase links cellular metabolism to histone acetylation. Science 2009;**324**:1076-1080

6. King, M. T., and Reiss, P. D. Separation and measurement of short-chain coenzyme-A

compounds in rat liver by reversed-phase high-performance liquid chromatography. Anal

Biochem 1985;**146**:173-179

7. Sabari, B. R., Tang, Z., Huang, H., Yong-Gonzalez, V., Molina, H., Kong, H. E. et al Intracellular

crotonyl-CoA stimulates transcription through p300-catalyzed histone crotonylation. Mol Cell

2015;**58**:203-215

8. Mantelingu, K., Reddy, B. A., Swaminathan, V., Kishore, A. H., Siddappa, N. B., Kumar, G. V. et al

Specific inhibition of p300-HAT alters global gene expression and represses HIV replication.

Chem Biol 2007;**14**:645-657

9. Kaczmarska, Z., Ortega, E., Goudarzi, A., Huang, H., Kim, S., Márquez et al Structure of p300 in

complex with acyl-CoA variants. Nat Chem Biol 2017; **13**:21-29

10. Liu, X., Wang, L., Zhao, K., Thompson, P. R., Hwang, Y., Marmorstein, R. et al The structural basis

of protein acetylation by the p300/CBP transcriptional coactivator. Nature 2008;**451**:846-850

11. Xie, Z., Zhang, D., Chung, D., Tang, Z., Huang, H., Dai, L. et al Metabolic Regulation of Gene

Expression by Histone Lysine β-Hydroxybutyrylation. Mol Cell 2016;**62**:194-206

12. Fellows, R., Denizot, J., Stellato, C., Cuomo, A., Jain, P., Stoyanova, E. et al Microbiota derived

short chain fatty acids promote histone crotonylation in the colon through histone deacetylases.

Nat Commun 2018;**9**:105

13. Wan, J., Liu, H., and Ming, L. Lysine crotonylation is involved in hepatocellular carcinoma

progression. Biomed Pharmacother 2019;**111**:976-982

14. Jiang, G., Nguyen, D., Archin, N. M., Yukl, S. A., Méndez-Lagares, G., Tang, Y. et al HIV latency is

reversed by ACSS2-driven histone crotonylation. J Clin Invest 2018;**128**:1190-1198

15. Kebede, A. F., Nieborak, A., Shahidian, L. Z., Le Gras, S., Richter, F., Gómez, D. A. et al Histone

propionylation is a mark of active chromatin. Nat Struct Mol Biol 2017;**24**:1048-1056

16. Farrelly, L. A., Thompson, R. E., Zhao, S., Lepack, A. E., Lyu, Y., Bhanu, N. V. et al   Histone

serotonylation is a permissive modification that enhances TFIID binding to H3K4me3. Nature

2019;**567**:535-539

17. Zhang, D., Tang, Z., Huang, H., Zhou, G., Cui, C., Weng, Y. et al. Metabolic regulation of gene

expression by histone lactylation. Nature **574**,575-580 (2019)

18. Lepack, A.E., Werner, C.T., Stewart, A.F., Fulton, S.L., Zhong, P., Farrelly, L.A. et al.

Dopaminylation of histone H3 in ventral tegmental area regulates cocaine seeking. Science **368**,

197-201 (2020)

19. Mews, P., Donahue, G., Drake, A.M., Luczak, V., Abel, T., Berger, S.L. Acetyl-CoA synthetase

regulates histone acetylation and hippocampal memory. Nature 2017; 546, 381–386

20. Wellen, K.E., Hatzivassiliou, G., Sachdeva, U.M., Bui, T.V., Cross, J.R., Thompson, C.B. ATP-

citrate lyase links cellular metabolism to histone acetylation. Science2009; 324, 1076–1080

21. Mews, P., Egervari, G., Nativio, R., Sidoli, S., Donahue, G., Lombroso, S.I. Alcohol metabolism

contributes to brain histone acetylation. Nature 2019; 574, 717–721

22. Tanaka, T., Yoshida, N., Kishimoto, T., Akira, S. Defective adipocyte differentiation in mice

lacking the C/EBPbeta and/or C/EBPdelta gene. EMBO J. 1997; 16, 7432–7443.

23. Hamm, J.K., Park, B.H., Farmer, S.R. A role for C/EBPbeta in regulating peroxisome proliferator-

activated receptor gamma activity during adipogenesis in 3T3-L1 preadipocytes. J. Biol. Chem.

2001; 276, 18464–18471.

24. Gao, M. et al. Metabolically controlled histone H4K5 acylation/acetylation ratio drives BRD4

genomic distribution. Cell Rep 2021; 36(4):109460.

25. Xiao, Y. et al. HBO1 is a versatile histone acyltransferase critical for promoter histone acylations.

Nucleic Acids Res 2021; 49(14):8037-8059.

26. Carrer, A., Parris, J. L., Trefely, S., Henry, R. A., Montgomery, D. C., Torres, A. et al Impact of a

High-fat Diet on Tissue Acyl-CoA and Histone Acetylation Levels. J Biol Chem 2017;**292**:3312-

3322

27. Burgeois, C., Gorwood, J., Barrail-Tran, A., Lagathu, C., Capeau, J., Desjardins, D. et al. Specific

Biological Features of Adipose Tissue, and their impact on HIV persistence. Front Microbiol **17**

(2019)

28. Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley et al Differential gene and transcript

expression analysis of RNA-seq experiments with TopHat and Cufflinks. Nat Protoc 2012;**7**:562-

578

29. Thomas, P. D., Campbell, M. J., Kejariwal, A., Mi, H., Karlak, B., Daverman, R. et al PANTHER: a

library of protein families and subfamilies indexed by function. Genome Res 2003;**13**:2129-2141

30. de Hoon, M. J. L., Imoto, S., Nolan, J.,Miyano, S. Open source clustering software. Bioinformatics

2004;**9**:1453-1454

Figure 1

Fig.1 Histone butyrylation increases during adipogenesis (A) Phase contrast images of 3T3L1 cells
exhibiting fibroblast like morphology in their pre-adipocyte state and accumulation of lipid droplets in
their adipocyte state. The lipid droplets can be stained using Oil Red-O dye. (B) RT-qPCR analysis was
performed to check for transcript levels of Acss2 in undifferentiated and differentiated 3T3L1 cells. Error
bars denote mean +/- SEM of three biological replicates; two-tailed unpaired Student's t-test: * $P < 0.05$,
**$P < 0.01$, ***$P < 0.001$, ns: not significant. (C) Knockdown of Acss2 in differentiated 3T3L1 cells was
confirmed by RT-qPCR analysis. Error bars denote mean +/- SEM of three biological replicates; two-tailed
unpaired Student's t-test: * $P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ns: not significant. (D) Histone
acetylation and butyrylation pattern in 3T3L1 cells upon Acss2 knockdown was checked using antibodies
against acetylated histone H3K18 and butyrylated H4K5. (E) Histone acetylation and butyrylation levels
in 3T3L1 cells in undifferentiated and differentiated state were analysed by immunoblotting with
antibodies specific for acetylated H3K9, butyrylated H3K9, H3K23, H3K27, H4K5, H4K8, H4K12 and pan-
butyryl lysine antibodies . Antibodies against core histones H3 and H4 were used as loading controls
along with Direct Blue staining of all four core histones. Lane 1- undifferentiated pre-adipocyte (Day 0)
and Lane 2 - differentiated adipocyte (Day 7) in each of the panels i, ii and iii. (F) Enrichment of H4K5
butyrylation in the promoters of (i) Pparg, (ii) Lep and (iii) Cebpd represented in the form of fold change
over IgG. Error bars denote mean +/- SEM of three biological replicates; one-way ANOVA with Dunnett's
multiple comparision was performed: * $P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ns: not significant.

Figure 2

Figure 2: LTK-14A selectively inhibits p300 catalysed butyrylation without affecting its acetyltransferase activity: Immunoblotting with antibodies against acetylated H3K18 (A) and pan-butyryl lysine (C) to check for the relative inhibition of acetyltransferase and butyryltransferase activities of full length p300 by LTK-14A at different concentrations. Recombinant histone H3 was used as substrate. Lane 1, enzyme alone; Lane 2, enzyme + DMSO; Lane 3-6, enzyme + LTK-14A at concentrations 10, 15, 20 and 25 µM respectively. Images of Western blotting are given in panels F and H and their quantitations are given in panels B and D. Molecular structures of mono- (LTK-14) and di- (LTK-14A) substituted isogarcinol (E and G). Molecular docking studies for studying comparative interaction of LTK-14 and LTK-14A with the amino acid residues of p300 catalytic domain (F and H). The predicted interactions are denoted by tick lines. Hydrogen bonding is shown in red and hydrophobic interaction is shown in green. Immunoblotting with antibodies against acetylated H3K18 (I) and pan-butyryl lysine (K) to check for the relative inhibition of acetyltransferase and butyryltransferase activities of wild type p300 catalytic domain by LTK-14A at different concentrations. Lane 1, enzyme alone; Lane 2, enzyme + DMSO; Lane 3-6, enzyme + LTK-14A at concentrations 10, 15, 20 and 25 µM respectively. Images of Western blotting are given in panels I and K and their quantitations are given in panels J and L. Immunoblotting with antibodies against acetylated H3K18 (M) and pan-butyryl lysine (O) to check for the relative inhibition of acetyltransferase and butyryltransferase activities of catalytically active p300 catalytic domain with point mutation C1438A by LTK-14A at different concentrations. Lane 1, enzyme alone; Lane 2, enzyme + DMSO; Lane 3-6, enzyme + LTK-14A at concentrations 10, 15, 20 and 25 µM respectively. Images of Western blotting are given in panels M and O and their quantitation are given in panels N and P. Error bars denote mean +/- SEM of three biological replicates, one-way ANOVA with Bonferroni's multiple comparison; * $P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ns: not significant.

Figure 3

Fig. 3: The butyrylation inhibitor represses adipogenesis in 3T3L1 cells: (A) The permeability and
stability of LTK-14A in 3T3L1 was tested by LC-ESI-MS analysis of metabolites, extracted from the
compound treated 3T3L1 cells to detect the protonated LTK-14A in the ESI spectra of the eluted fraction.
(B) MTT assay was performed to check for any possible cellular toxicity effect of LTK-14A, using 25 µM
concentration of the compound on 3T3L1 cells upon incubation for varying lengths of time (2, 4 and 6
days). Comparison was done with cells that were left untreated or treated with DMSO as solvent
control. The effect of LTK-1a4A (25 µM) treatment on adipogenesis was tested by Oil Red-O staining of
the adipocytes (C) and the quantitation of staining is represented graphically in panel D. Error bars
denote mean +/- SEM of three biological replicates,one-way ANOVA with Bonferroni's multiple
comparison; * P < 0.05, **P < 0.01, ***P < 0.001, ns: not significant. (E) The effect of LTK-14A on the
global transcription network inside 3T3L1 cells was investigated by high throughput transcriptome
analysis with RNA isolated from compound treated, DMSO treated and untreated 3T3L1 cells (n=2 in
each case). Differential gene expression pattern of three conditions compared with each other, depicted
in volcano plots- No Inhibitor versus DMSO , No Inhibitor versus LTK-14A and DMSO versus LTK-14A. The
downregulated and upregulated genes are shown in blue and red respectively.(F) Heat maps showing
the differential gene expression pattern between LTK-14A treated cells and DMSO treated or untreated
cells.

Figure 4

Figure 4: LTK-14A represses expression of several pro-adipogenic genes and inhibits lipid metabolism:
(A) Venn diagram showing the number of differentially expressed genes upon LTK-14A treatment with
respect to untreated and DMSO treated conditions. (B) Classification of the common differentially
expressed genes according to their type. Gene ontology analysis was performed to study the different
biological pathways that were affected by LTK-14A in 3T3L1 compared to DMSO treated as well as
untreated conditions, as predicted by Panther software. The pathways were ranked in the order of
those with maximum number of genes affected (E). A different ranking order was created on the basis of
fold enrichment i.e. the pathways in which the ratio of number of observed affected genes over number
of expected genes to be affected was the highest (C). A similar analysis was also performed for genes
differentially expressed in DMSO versus untreated conditions (D). (F) A molecular network was
constructed to depict the interconnectivity between the lipid metabolism related genes and pathways
that were affected upon their dysregulation by LTK-14A compared to DMSO treated conditions. (G) RT-
qPCR analysis was performed to validate the putative target genes of LTK-14A. Error bars denote mean
+/- SEM of three biological replicates; one-way ANOVA with Bonferroni's multiple comparison: * P <
0.05, **P < 0.01, ***P < 0.001, ns: not significant.

Figure 5

Fig.5: LTK-14A inhibits histone butyrylation in 3T3L1 cells, without affecting acetylation or other post-translational modifications: Histone butyrylation levels in 3T3L1 cells that were treated with LTK-14A were compared with that in DMSO treated and untreated conditions by immunoblotting with antibodies against butyrylated histone H3K23 , H4K5 and pan-butyryl lysine. Histones H3 and H4 were used as loading controls (A). The fold change of quantified band intensity is graphically represented in panel (B) in which the error bars denote mean +/- SEM of three biological replicates. Under similar conditions histone acetylation levels were compared by immunoblotting using antibodies against acetylated histones H3K18, H3K9 and H4K12 (C). The result is graphically represented in panel (D) in which error bars denote mean +/- SEM of three biological replicates. Immunoblotting was also performed with antibodies against histone H3R17 dimethylation, H3S10 Phosphorylation and H3K9 dimethylation (E). The result is graphically represented in panel F in which error bars denote mean +/- SEM of three biological replicates. For all data one-way ANOVA with Bonferroni's multiple comparision was performed: * $P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ns: not significant .

Figure 6

Fig. 6: The weight gaining of high fat diet fed mice (C57BL6/J) could be significantly reduced by the butyrylation specific inhibitor of p300: Trend of weight gaining (A) and average body weight gained after 16 weeks (B) of C57BL6/J mice maintained on normal chow diet, high fat diet and high fat diet mixed with LTK-14A was plotted; n=4 in each group. Representative images of dissected bodies showing fat accumulation (C) and epididymal fat pads (D) of the mice from three groups. The average weight of the fat pads is represented graphically (E); n=4 in each group. Representative images of morphology of adipose tissue (F) and liver (G) from the mice of the three groups, as studied by hematoxylin and eosin staining. For all data one-way ANOVA with Dunnett's multiple comparision was performed: $* P < 0.05$, $**P < 0.01$, $***P < 0.001$, ns: not significant. (H) Immunofluorescence microscopy of liver sections stained with antibodies against histone H4K12 acetylation (panel i), histone H3K14 acetylation (panel ii) and histone H4K5 butyrylation (panel iii). The fluorescence intensity of the modified histone marks was normalised against DAPI staining of whole nuclei for quantitation which is graphically depicted in (I) panel i, ii and iii ;n=4 in each group; one-way ANOVA with Dunnett's multiple comparision: $* P < 0.05$, $**P < 0.01$, $***P < 0.001$, ns: not significant. (J) Histone H3K14, H3K18, H4K12 acetylation and H4K5 butyrylation levels in epididymal fat pads of mice were estimated by immunoblotting . Histone H3 was used as loading control The fold change of quantified band intensity is graphically represented in panel K,L, M and N ;n=3 in each group; one-way ANOVA with Dunnett's multiple comparision: $* P < 0.05$, $**P < 0.01$, $***P < 0.001$, ns: not significant.

Figure 7

Fig. 7: Oral gavaging of the butyrylation inhibitor reduces weight of genetically obese db/db mice:
Trend of weight gaining (A) and average body weight gained after 30 days (B) of db +/- and db/db mice
orally administered with either LTK-14A or vehicle solution ; n=6 in each group. (C) Representative
images of dissected bodies showing fat accumulation (panel iii) of the mice from three groups.
Representative images of liver (D) from the mice of three groups and average weight of liver (E)
depicted graphically; n= 6 in each group. Representative images of epididymal fat pads (F) from the mice
of three groups and average weight of fat pads (G) depicted graphically; n= 6 in each group.
Representative images of morphology of liver (H) and adipose tissue (I) from the mice of the three
groups, as studied by hematoxylin and eosin staining. (J) Immunofluorescence microscopy of liver
sections stained with antibodies against histone H4K12 acetylation (panel i), histone H3K14 acetylation
(panel ii) and histone H4K5 butyrylation (panel iii). The fluorescence intensity of the modified histone
marks was normalised against DAPI staining of whole nuclei for quantitation which is graphically
depicted in (K) panel i, ii and iii ;n=4 in each group; one-way ANOVA with Dunnett's multiple
comparision: * P < 0.05, **P < 0.01, ***P < 0.001, ns: not significant. (L) Histone H3K14, H3K18, H4K12
acetylation and H4K5 butyrylation levels in epididymal fat pads of mice were estimated by
immunoblotting. Histone H3 was used as loading control The fold change of quantified band intensity is
graphically represented in panel M,N, O and P; n=3 in each group; one-way ANOVA with Dunnett's
multiple comparision: * P < 0.05, **P < 0.01, ***P < 0.001, ns: not significant. (Q) qRT-PCR with RNA
isolated from epididymal fat pads to check the expression pattern of lipogenesis related pro-adipogenic
genes upon LTK-14A treatment; n=3/4 in each group; one-way ANOVA with Dunnett's multiple
comparision: * P < 0.05, **P < 0.01, ***P < 0.001, ns: not significant

Graphical abstract

Graphical abstract:

Proposed model for the role of p300-mediated histone butyrylation in adipogenesis:

In pre-adipocytes, there exists a basal level of histone acetylation while butyrylation is present to a
much lesser extent owing to low stoichiometric levels of butyryl CoA. Induction of adipogenesis causes a
simultaneous upregulation of histone acetylation and butyrylation marks leading to increased rate of
adipogenesis and concomitant transcriptional activation of pro-adipogeneic genes.

Onset of obesity in mice, either due to excess energy intake through high fat diet consumption or
increased de novo synthesis of fatty acids due to leptin receptor gene mutation leading to hyperphagic
behavior, is accompanied by adipocyte hyperplasia and hypertrophy. Both the organs of adipose tissue
and liver were found to have enhanced levels of H4K5 butyrylation during obesity. LTK-14A, a
butyrylation specific inhibitor could efficiently prevent the processes of adipogenesis and adipocyte
hypertrophy due to inhibition of H4K5 butyrylation in these organs. Thus the compound could attenuate
weight gain by selective inhibition of butyrylation without affecting acetylation, thereby highlighting the
importance of histone butyrylation in adipogenesis and obesity.

Supplementary Section:

Supplementary Materials and Methods:

siRNA experiments: ON-TARGETplus siRNA SMART pools against mouse Acss2 (catalog no.L-065412-01-
0005) was purchased from Dharmacon (Dharmacon, Lafayette, CO, USA). 3T3L1 cells were transfected
with Acss2 siRNA at 25 nM concentration, using RNAiMAX (catalog no. 13778-150, Thermo Fisher
Scientific) according to manufacturer's protocols. Transfection was carried out for 48 hours after which
cells were induced for differentiation for 6 days followed by RNA isolation and histone extraction.

Molecular docking analysis: Glide module of Schrodinger suite was used to dock LTK-14 and LTK-14A to
p300 catalytic domain (PDB: 3BIY). Chimera was used as visualizing software. The protein was prepared
using the wizard tool in Maestro version 10.2.The crystal structure of LTK-14A (CCDC 1969185) and LTK-
14 (CCDC 645420) were used as the most optimized and energy minimized ligand structures. The
receptor grid for p300 catalytic domain was generated by specifying the binding (active) site residues
using Receptor grid generation in the Schrodinger Glide application. Once the receptor grid was
generated, the ligands LTK-14 and LTK-14A were docked to the p300 catalytic domain using Glide
version 5.8 molecular docking.

**Untargeted analysis of intracellular metabolites by Ultra-performance liquid chromatography coupled
with time-of-flight mass spectrometer (Q-TOF LC/MS):** Profiling of intracellular metabolites was
performed on an agilent 1290 Infinity LC system coupled to Agilent 6545 Accurate-Mass Quadrupole
Time-of- Flight (QTOF) with Agilent Jet Stream Thermal Gradient Technology. The UPLC system was
assembled with a Diode array detector (DAD) and autosampler. The Chromatographic separation was
achieved on Agilent ZORBAX SB- C18 column (2.1 × 100 mm, 1.8 μm) as stationary phase. The mobile
phase consisted of a linear gradient of 100 mM ammonium formate (A) and Acetonitrile (B): 0–10.0 min,
30-80% B (v/v); 10.0–15.0 min, 80–100% B (v/v); 15.0–20.0 min, 100% B (v/v); 20.0–21.0 min, 100–30%
B (v/v); 21.0–25.0 min, 30% B. The sample was dissolved in 1 ml methanol (LCMS Grade), centrifuged
and supernatant was taken for UPLC-QTOF-MS analysis. The column was reconditioned for 5 minutes
prior to the next injection. The flow rate was 0.5 ml/min, and the injected volume was 20 μL. The UPLC
was connected to the MS analysis was performed on an Agilent 6545 Accurate-Mass Q-TOF/MS system
with an electrospray ionization (ESI) source. Considering the MS conditions, positive ion mode was used
to obtain high-resolution mass spectra. The ESI source parameters were: drying gas (N2) flow, 8 L/min;
drying gas temperature, 200°C. Other parameters were set as nebuliser gas, 35 psig; capillary voltage,
3000 V; skimmer voltage, 65 V; nozzle voltage 1000 V and fragmentor voltage 150 V. The data
acquisition on the LC-QTOF was performed using Agilent Mass Hunter Acquisition software (Agilent
Technologies, Santa Clara, CA, USA). The data were deconvoluted into individual chemical peaks with
Agilent MassHunter Qualitative Analysis (MassHunterQual, Agilent Technologies).

Site directed mutagenesis: Point mutation was introduced into the p300 catalytic domain sequence
cloned within pET-28b plasmid by using the Quikchange site directed mutagenesis kit (Agilent). Briefly,
the template was mixed with Pfuturbo DNA polymerase, designed forward and reverse primers and
dNTPs for PCR (Initial denaturation for 30 seconds at 95°C; Cycles: 95°C 30'', 55°C for 60'', 68°C for 7';

final elongation at 68°C for 5'.) The mixture was then incubated with DpnI enzyme for 2.5 hours for
removal of the parent wild type sequence followed by transformation of competent XL10Gold cells and
kanamycin selection of positive transformants. The inserted mutation was verified by sequencing and
alignment with the wild type sequence using ClustalOmega software. The sequences of the primers used
are as follows:

Forward primer- 5'- CCTCACTTGGTGGAGCTGCCCAAAATATGCCCTGTTGTG – 3'

Reverse primer- 5'- CACAACAGGGCATATTTGGGCAGCTCCACCAAGTGAGG -3'

Purification of recombinant full-length p300 and p300 catalytic domain: Polyhistidine tagged
recombinant p300 was purified from Sf21 insect cells by Ni-NTA based affinity chromatography. These
cells were infected by the baculovirus containing p300 for 72 hours. Once the cells showed signs of
infection, they were scraped off and centrifuged to separate the spent media. The cell pellets were
resuspended in ice cold homogenisation buffer (20 mM Tris, pH 7.5; 20 mM imidazole, 10% glycerol, 0.2
mM EDTA, 300 mM KCl, 0.1%NP-40, 2 mM PMSF, 2 mM βmercaptoethanol) and then homogenised
using a Dounce homogeniser (5 cycles, 6 strokes per cycle with 5 minutes interval between each cycle).
The cell debris was separated by centrifugation and then the lysate was incubated with pre-equilibriated
Ni-NTA slurry for 2.5 hours at 4°C on an end-to-end rotor. Then the beads were washed with wash
buffer (20 mM Tris, pH 7.5; 40 mM imidazole) nine times to remove unwanted contaminants. Finally
p300 was eluted using elution buffer (20 mM Tris, pH 7.5; 250 mM imidazole).

Polyhistidine tagged p300 catalytic domain cloned in pET28b plasmid and Sirt2 were used for co-
transforming BL21(DE3) cells. At first, 50 ml primary culture was prepared in Luria broth containing
kanamycin (50 μg/ml) and ampicillin (100 μg/ml) (HIMEDIA) and then it was scaled up 10 times to
prepare secondary culture. Incubation was done at 37°C, 180 rpm till O.D. reached 0.37 followed by
induction with 0.2 mM IPTG and a further incubation at 30°C for 4 hours. Cells were pelleted at 6000
rpm, 4°C for 10 minutes and stored at -80°C. Resuspension and homogenisation of the pellet was
carried out in homogenisation buffer (20 mM Tris, pH 7.5; 20 mM imidazole, 10% glycerol, 0.2 mM
EDTA, 300 mM KCl, 0.1%NP-40, 2 mM PMSF, 2 mM βmercaptoethanol). 3 cycles of sonication were done
for 3 minutes at 35% amplitude with 5 minutes interval between each cycle. The cell debris were
separated by pelleting at 12000 rpm for 30 minutes at 4°C. The supernatant was equilibriated with Ni-
NTA beads slurry (Novagen) in an end-to-end rotor for 3 hours at 4 C. Eight washes were given with
wash buffer (20 mM Tris, pH 7.5; 40 mM imidazole). Elution was carried out in batches with elution
buffer (20 mM Tris, pH 7.5; 250 mM imidazole).

The enzymatic activity of the proteins was estimated by filter binding assay.

In vitro **filter binding assay:** Different dilutions of purified enzyme were incubated in reaction buffer (50
mM Tris-HCl pH 7.5, 1 mM DTT, 10 % Glycerol, 2 mM PMSF) with 10 mM NaBu, tritiated acetyl CoA (50

μCi) (PerkinElmer, Part No: NET290050UC) and Xenopus histone H3 (500 ng) for 30 minutes at 30^{o}C. The
reaction mixture was then spotted on P81 filter paper and the paper was allowed to dry at room
temperature. The paper was washed in wash buffer (50 mM sodium bicarbonate and 50 mM sodium
carbonate), dried at 90^{o}C and immersed in enhancing solution (2.5% PPO and 0.25% POPOP in toluene)
for taking radioactive counts using liquid scintillation counter (Perkin Elmer, MicroBeta2 system).

In vitro **acylation (acetylation/butyrylation) assay:** Acylation reactions were performed in reaction
buffer (25 mM Tris-HCl pH 7.5, 100 mM NaCl, 0.1 mM EDTA, 1 mM DTT, 10 % Glycerol, 1x PIC) with 100
ng/mL TSA, and 50 μM Butyryl-CoA/Acetyl CoA. Xenopus histone H3 (1 μg) was used as the substrate
and full length p300 or p300 catalytic domain (10,000 cpm activity) was used as the enzyme. Reactions
were incubated in presence or absence of butyrylation inhibitor for 10 minutes at 30 °C, followed by
initiation of the acetylation/butyrylation reaction by the addition of acetyl CoA/butyryl CoA. After a
further 10 minutes, reactions were stopped by addition of Laemmli buffer and samples were used for
immunoblotting.

Transcriptomics analysis by RNA-seq: RNA integrity was checked by Agilent Bioanalyzer 2100, only
samples with clean rRNA peaks were used. Libraries for RNA-seq were prepared according to KAPA
Stranded RNA-Seq Kit with RiboErase (KAPA Biosystems, Wilmington, MA) system. Final library quality
and quantity were analyzed by Agilent Bioanalyzer 2100 and Life Technologies Qubit3.0 Fluorometer,
respectively. 150 bp Paired-end sequencing was performed on Illumina HiSeq 4000 (Illumnia Inc., San
Diego, CA).

House mouse (Mus musculus, strain C57BL/6J) genome (mm10) was downloaded from GENCODE and
indexed using Bowtie2-build with default parameters. Adapter removal was done using Trim Galore (v
0.4.4) and each of the raw Fastq files were passed through a quality check using FastQC. PCR duplicates
were removed using the Samtools 1.3.1 with the help of 'rmdup' option. Each of the raw files was then
aligned to mm10 genome assembly using TopHat2 with default parameters for paired-end sequencing
as described in (28). After aligning, quantification of transcripts was performed using Cufflinks and then
Cuffmerge was used to create merged transcriptome annotation. Finally differentially expressed (DE)
genes were identified using Cuffdiff. The threshold for DE genes was log2 (fold change) >1.5 for up
regulated genes and log2 (fold change) <1.5 for down regulated genes with p-value <0.05.

GO enrichment analysis

Gene Ontology (GO) analysis was performed in PANTHER (29). Significant enrichment test was
performed with the set of differentially expressed genes in PANTHER and Bonferroni correction method
was applied to get the best result of significantly enriched biological processes.

Fisher's exact test

Fisher's exact test was performed in PANTHER Gene Ontology (GO) where p-value significance was
calculated based on the ratio of obtained number of genes to the expected number of genes (O/E)
considering the total number of genes for the respective pathway in Mus musculus with a FDR of <0.05.

Heatmap and clustering of genes

Unsupervised hierarchical clustering method was performed using Cluster 3.0 (30) with Pearson
Correlation and average linkage rule. Gene expression data (FPKM of all samples) was taken and log2
transformed. Low expressed (FPKM<0.05) and invariant genes were removed. Then genes were
centered and clustering was performed based on differential expression pattern of genes and fold
change. Finally, the heatmap was visualized in Java TreeView 3.0.

Biological analysis of differentially expressed transcripts and pathway regulatory network modeling

Statistically significant differentially expressed transcripts were subjected to GO and Pathway
enrichment using DAVID tool. Only those GO and pathways with a FDR score of <=0.05 was considered
for further downstream analysis. Key biologically dysregulated GO and Pathways along with the
differentially expressed genes was provided as an input to Pathreg algorithm from Theomics
International Pvt Ltd, Bangalore, India for gene regulatory network modeling. The result (nodes and
edges) of the Pathreg algorithm was provided as an input to Cytoscape v2.8.2 to identify key nodes and
edges that could be representative of the gene regulatory changes upon treatment.

Quantitative real time PCR: Total RNA was isolated from the 3T3L1 cells and epididymal fat pads by
phenol-chloroform method. In case of LTK-14A treatment, 3T3L1 cells were allowed to differentiate for
7 days following the standard adipogenesis protocol, in presence of the compound at 25 μM
concentration. The control experiments were 3T3L1 cells differentiated with or without DMSO as
solvent control and differentiation was done only with the addition of only the cocktail of chemicals for
adipogenesis. RNA was isolated from the cells after the period of 7 days in all three cases. For isolation
of RNA from mice fat pads, the adiose tissue samples were lysed in Trizol using a mechanical
homogenizer followed by phenol-chloroform method of extraction. The isolated RNA samples were
treated with 2 units/μL DNase I to remove any residual DNA followed by ethanol precipitation at -80°C.
2μg RNA was used for cDNA synthesis using MMLV reverse transcriptase (Sigma) in presence of 0.5 mM
dNTP and 3.5 μM oligo dT (Sigma). RT-qPCR amplification was done in kappa SYBR green reagent
(Biosystems) with the help of Real time PCR instrument (Applied Biosystems by Life Technologies).For
estimation of relative fold change in RNA expression, the following calculation was done: Relative fold
change = 2-ΔΔCt, where ΔΔCt=ΔCt value of sample – ΔCt value of control and ΔCt= Ct value of target – Ct
value of internal control actin for individual sample. The sequences of primers are provided in Table 4.

Hematoxylin and eosin staining of liver and adipose tissues: Staining was performed on paraffin
embedded sections of liver and epididymal fat pads having the thickness of 5 μm and 10 μm
respectively. Briefly, the sections were deparaffinized in xylene, followed by immersion in absolute
alcohol, 90% ethanol, 70% ethanol, 50% ethanol and water before staining with hematoxylin for 5-10
minutes. The sections were immersed in above mentioned solutions in reverse order before counter-

staining in eosin for 15 seconds. Then the samples were dipped in absolute ethanol and xylene before
mounting with DPX.

MTT assay: 12500 cells/ml were seeded in presence/absence of LTK-14A in 96-well plate and at
different intervals of time (2 days, 4 days, 6 days) , 1/10th volume of 5mg/ml MTT solution was added to
it followed by incubation at 37 °C for 3 hours. The formed crystals were dissolved in DMSO and
spectrometric estimation of colour intensity was done at 570 nm.

1052

Evaluation of mutagenecity by plate incorporation assay: Five doses of LTK-14A (10.0㎍, 33.0㎍,
100.0㎍, 333.0㎍ and 1000.0㎍/plate) were selected to check for its possible mutagenecity. Seven
groups decided as Group 1- Control, Group 2 Positive Control and 5 doses of LTK-14A (Group 3 to Group
7: 10.0㎍, 33.0㎍, 100.0㎍, 333.0㎍ and 1000.0㎍/plate) were tested. Four tester strains of Salmonella
typhimurium (TE 97a, TA 98, TA 100, TA102) were used and their respective positive controls were used
in absence of S9 fraction (Table 5); 2-aminoanthracene was used as the positive control in presence of
S9 fraction (Table 5). The tester strains were grown overnight in Luria Broth antibiotic selection medium
at 37㎕C. LTK-14A prepared in DMSO was mixed with the bacterial culture, 0.5mM Histidine-Biotin and
top agar in a culture tube and the content was overlaid on to minimal glucose plate. In case of plate
incorporation assay with S9 mix, LTK-14A was incubated with S9 mix in cold condition for 15 to 20
minutes followed by the same procedure as the plate incorporation assay without S9. The plates were
incubated in an inverted position in 37㎕C incubators for 48 hours following which the number of
revertants was counted using automated colony counter.

Single dose toxicity study: The single dose toxicity study on LTK-14A was conducted in accordance with
Good Laboratory Practice Principles as published by the OECD in 1998, in accordance with the Drugs and
Cosmetics Rule (New Drugs and Clinical trials, 2019/ DCGI). LTK-14A (formulated in vehicle i.e. 0.5%
sodium salt of carboxymethyl cellulose) was administered to Sprague Dawley rats once by oral route on
the day of scheduled treatment. The three treatment groups were 250 mg, 500 mg, and 1000 mg/kg
body weight, while vehicle control was administered to control group. The rat equivalent doses were
calculated from the dosage used in mice experiments following FDA guidelines for drug administration.
The doses finally administered were 10, 20 and 40 times the corresponding mice equivalent dose. Food,
but not water, was removed thus letting the animals fast over night prior to dosing. After administration
of LTK-14A, food was withheld for further 2-3 hours and the animals were observed at 30 minutes, 1
hour, 2 hours and 4 hours from the time of dosing. Animals were subsequently observed once daily for
any signs of toxicity, mortality, or any other observations including conditions of skin, hair coat, eyes,
mucous membrane, movement, activity, tremors, convulsions, salivation, diarrhea, posture, piloerection
and lethargy. Food intake, water intake and body weight of animals was recorded weekly. Terminal
sacrifice of surviving animals was performed after 14 days using CO2 euthanasia technique following
which animals were necropsied. Animals were fasted over night before necropsy. Gross observations
were recorded.

1083

Supplementary Figure 1

1084

1085

1086

1087

1088

1089

1090

Sup.Fig.1: (A) Structures of garcinol and its cyclised and substituted derivatives. The site of internal
cyclisation and mono/di-substitutions are shown in brown colour. Initial screening of the different
monno-substituted derivatives for their differential effect on p300 catalysed acetylation (B) and
butyrylation (C) of histone H3 at a concentration of 10 µM. Immunoblotting with antibodies against
acetylated H3K18 (B) and pan-butyryl lysine (C). Similar screening was performed with di-substituted
derivatives for their effect on p300 catalysed acetylation (D) and (E) of histone H3 at a concentration of
25 µM.

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

Supplementary Figure 2

A

B

C

D

E

F

1119

1120

1121

1122

1123

1124

1125

1126

1127

Sup. Fig.2: (A) Crystal structure of LTK-14A (CCDC: 1969185) (B) Bar graphs depicting the
acetyltransferase activity of wild type an mutant C1438A p300 catalytic domain as meeasured by filter
binding assay. Error bars denote mean +/- SEM of four technical replicates; two-tailed unpaired
Student's t-test: * P < 0.05, **P < 0.01, ***P < 0.001, ns: not significant. (C) Proton NMR, (D) HR-MS
spectra and (D) HPLC chromatogram of LTK-14A. (F) The presence of LTK-14A in the physiological
system upon administering the compound was verified by LC-ESI-MS analysis of the metabolite extracts
of liver.

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

Supplementary Figure 3

Sup. Fig. 3: (A) Molecular pathway analysis for the expression status of major lipid metabolism related genes and their interrelationship with the pathways that were affected in LTK-14A versus untreated conditions. (B) Pro-adipogenic genes that could be putative targets for LTK-14A were identified and RPKM values for these genes have been depicted graphically.(C) RT-qPCR analysis was performed to validate the putative target genes of LTK-14A. Error bars denote mean +/- SEM of three biological replicates. (D) Histone butyrylation levels in 3T3L1 cells that were treated with LTK-14A were compared with that in DMSO treated and untreated conditions by immunoblotting with antibodies against butyrylated histone H3K9, H3K27, H4K8 and H4K12. Histones H3 and H4 were used as loading controls.

Supplementary Figure 4

1191

1192

1193

1194

1195

1196

Sup. Fig.4: Evaluation of mutagenecity of LTK-14A at different doses was tested by performing plate incorporation assay with four different tester strains of Salmonella typhimurium (TE 97a, TA 98, TA 100, TA102). Panel (A) to (D) – tests without S9 fraction. Panel (E) to (H) – tests with S9 fractions. Error bars denote mean +/- SEM of three biological replicates; One-way ANOVA followed by Neman-Keuls test.

1224   **Table 1: Crystal data and structure refinement for LTK14A (CCDC: 1969185)**

1225

Empirical formula	$C_{40}H_{54}O_6$
Formula weight	630.86
Temperature/K	296 (2)
Crystal system	monoclinic
Space group	C2
a/Å	19.1677(11)
b/Å	11.5860(7)
c/Å	16.8399(11)
α/°	90
β/°	90
γ/°	90
Volume/Å^3	3739.8(4)
Z	110
ρ_{calc}g/cm^3	1.417
μ/mm^{-1}	0.131
F(000)	1650
Crystal size/mm^3	0.35 × 0.18 × 0.15
Radiation	MoKα (λ = 0.71073)
2Θ range for data collection/°	0.995 to 25.75
Index ranges	-23 ≤ h ≤ 23, -14 ≤ k ≤ 14, -20 ≤ l ≤ 20
Reflections collected	108274
Independent reflections	108274 [R_{int} = 0.0811, R_{sigma} = 0.0325]
Data/restraints/parameters	7123/3/415
Goodness-of-fit on F^2	1.045
Final R indexes [I>=2σ (I)]	R_1 = 0.1038, wR_2 = 0.2797
Final R indexes [all data]	R_1 = 0.1362, wR_2 = 0.3157

1226

1227

1228

1229

1230   **Table 2: Food consumption pattern of mice used in obesity studies**

Set A			
Type of diet	**Chow diet**	**High fat diet**	**High fat diet and LTK-14A**
Average food consumed per mice per day (gram)	8.0157	1.95547	2.04051
Standard error of mean	1.31838	0.23142	0.24589
Set B			
Type of mice	**db +/-**	**db/db and vehicle control**	**db/db and LTK-14A**
Average food consumed per week per mouse (gram)	25.8	37.8	34
Standard error of mean	1.04	1.64	2.08

1231

1232   **Table 3: List of primers used in ChIP-qRTPCR**

Gene	Primer sequence 5' - > 3'
Pparg Fwd	AGCTCGATGACCATAAGCCT
Pparg Rev	ATAGGCAGTGCATCAGCGAA
Lep Fwd	AGGTCATGTGGACAGCTTGG
Lep Rev	CGTCCAGCCTTGCTTTTTCC
Cebpd Fwd	CGCTTTTCAGCCTGGACAG
Cebpd Rev	TGGAGTCAATGTAGGCGCTG

1233

1234   Table 4: List of primers used in gene expression analysis by qRTPCR

Gene	Primer sequence 5' - > 3'
Actin Fwd	**CCACTGTCGAGTCGCGTCC**
Actin Rev	**GTCATCCATGGCGAACTGGTG**
Pparg Fwd	**TCCGTGATGGAAGACCACTC**
Pparg Rev	**CCCTTGCATCCTTCACAAGC**
Cebpd Fwd	**ATCGACTTCAGCGCCTACAT**
Cebpd Rev	**GCTTTGTGGTTGCTGTTGAA**
aP2 Fwd	**TCTCCAGTGAAAACTTCGAT**
aP2 Rev	**TTACGCTGATGATCAATGTT**
Slc2a4 Fwd	**ACGACGGACACTCCATCTGTTG**
Slc2a4 Rev	**GGAGACATAGCTCATGGCTGGAA**
Lep Fwd	**GAGCGGGATCAGGTTTTGTGGT**
Lep Rev	**TGTCACTCTTTCCCGGTCTCTTCA**

Adipoq Fwd	CAGTGGATCTGACGACACCAA
Adipoq Rev	GAACAGGAGAGCTTGCAACAGT
Cebpa Fwd	TACCGAGTAGGGGGAGCAAA
Cebpa Rev	TCATTTTTCTCACGGGGCCA
TEAD4 Fwd	GCCACAGCCTTCCACAGTAA
TEAD4 Rev	AAGCTCCTTGCCAAAACCCT
Kdm4c Fwd	CAGTGCGGAACAAGTCTTCCAA
Kdm4c Rev	CAGATGTTTCTGCGCCGCT
Fasn Fwd	GGGTGTGAGTGGTTCAGAGG
Fasn Rev	AGGGCAATGCTTGGTCCTTT
Acss2 Fwd	TCGGAACTTGATGCGATGGG
Acss2 Rev	TTTCCCCAAAACTCCCGTGG

1235

1236   **Table 5: List of mutagenic agents used as positive control in Ames test**

Tester Strains	Without S9	With S9
TA-97a	9-aminoacridine (50µg/plate)	2-aminoanthracene (5µg/plate)
TA-98	N-nitro-o-phenylene diamine (2.5µg/plate)	2-aminoanthracene (5µg/plate)
TA-100	Sodiumazide (5µg/plate)	2-aminoanthracene (5µg/plate)
TA-102	Mitomycin C (0.5µg/plate)	2-aminoanthracene (5µg/plate)

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246   **Table 6: Details of single dose toxicity test**

1247   **Body weight (gram, mean +/- SD of animals in all groups**

Group		Male				Female		
		Initial	Mid	Final		Initial	Mid	Final
Vehicle control	Mean	174.75	217.38	232.13	Mean	152.21	169.81	171.5
	±SD	7.61	11.38	18.38	±SD	6.58	6.57	6.65
LTK-14A (250 mg/Kg bw)	Mean	171.94	216.62	244.21	Mean	151.99	174.54	176.35
	±SD	6.87	13.63	29.81	±SD	6.43	6.06	9.5
LTK-14A (500 mg/Kg bw)	Mean	176.3	229.75	243.14	Mean	153.32	176.55	175.8
	±SD	5.53	17.08	12.25	±SD	6.44	10.86	6.48
LTK-14A (1000 mg/Kg bw)	Mean	176.24	217.65	226.19	Mean	153.81	178.58	179.52
	±SD	7.51	12.31	23.57	±SD	7.29	7.46	8.45

1248

1249   **Average food intake (gram/day/animal) in each group**

Group	Male			Female		
	Initial	Middle	Final	Initial	Middle	Final
Vehicle control	20.38	20.24	21.24	17.84	15.65	15.95
LTK-14A (250 mg/Kg bw)	20.03	20.79	22.62	20	16.32	16.65
LTK-14A (500 mg/Kg bw)	20.99	21.14	23.29	18.22	18.13	17.09
LTK-14A (1000 mg/Kg bw)	20.76	20.74	19.95	18.17	17.82	16.97

1250

1251   **Average water intake (ml/day/animal) in each group**

Group	Male			Female		
	Initial	Middle	Final	Initial	Middle	Final
Vehicle control	32	30	34	25	21	26
LTK-14A (250 mg/Kg bw)	32	27	34	26	21	24
LTK-14A (500 mg/Kg bw)	36	33	40	25	23	26
LTK-14A (1000 mg/Kg bw)	34	31	38	26	22	32

1252

1253