EP300 (p300) mediated histone butyrylation is critical for adipogenesis

Running title: Lysine butyrylation inhibition blocks adipogenesis

Aditya Bhattacharya [1], Sourav Chatterjee [1], Utsa Bhaduri [2], Akash Kumar Singh [1], Koneni V
Sashidhara [4], Rajdeep Guha [5], Aamir Nazir [6], Srikanta Kumar Rath [6], Nagashayana Natesh , Tapas K
Kundu [1,3]

1   Transcription and Disease Laboratory, Molecular Biology and Genetics Unit, Jawaharlal Nehru Centre for
Advanced Scientific Research, Bangalore, India.
2   Chromatin Biology Laboratory, Molecular Biology and Genetics Unit, Jawaharlal Nehru Centre for
Advanced Scientific Research, Bangalore, India.
3   Division of Neuroscience and Aging Biology,CSIR-Central Drug Research Institute, Lucknow
4   Division of Medicinal and Process Chemistry, Central Drug Research Institute, Lucknow, India
5   Division of Laboratory Animal Facility, Central Drug Research Institute, Lucknow, India
6   Division of Toxicology and Experimental Medicine, Central Drug Research Institute, Lucknow, India
7   Central Government Health Scheme Wellness Centre Number 8, Domlur, Bangalore, India

E-mail ID of corresponding author: tapas@jncasr.ac.in

Affiliations:

Aditya Bhattacharya – PhD graduate student

Sourav Chatterjee – Post-doctoral researcher

Utsa Bhaduri – Research assistant

Akash Kumar Singh - Integrated PhD student

Koneni V Sashidhara - Professor

Rajdeep Guha – Assistant Professor

Aamir Nazir- Principal Scientist

Srikanta Kumar Rath- Chief Scientist

Nagashayana Natesh – Senior physician

Tapas K Kundu - Professor

Abstract

The master epigenetic enzyme EP300 (p300) besides having lysine acetyltransferase activity can also

catalyse other acylation modifications (propionylation, butyrylation, crotonylation etc.), the

physiological implications of which are yet to be established fully. Here, by performing immunoblotting

and chromatin immunoprecipitation assays, we report that histone butyrylation increases both globally

as well as locally in the promoters of pro-adipogenic genes upon adipogenesis. Moreover, the enzyme

ACSS2 which is responsible for generating acyl CoA for being used as a cofactor in acylation reactions is

upregulated in adipogenesis and its genetic ablation leads to reduction of butyrylation. To delineate the

role of p300 catalysed butyrylation from acetylation in adipogenesis, we have identified a semi-synthetic

derivative (LTK-14A) of garcinol which specifically inhibited histone butyrylation without affecting

acetylation mediated by p300. Treatment of 3T3L1 cells with LTK-14A significantly abolished

adipogenesis with downregulation of pro-adipogenic genes along with the inhibition of H4K5

butyrylation. Administering the specific inhibitor to high fat diet fed C57BL6/J mice as well as genetically

obese db/db mice led to an attenuation/decrease in their weight gain respectively. The reduced obesity

could be at least partially attributed to the targeted inhibition of H4K5 butyrylation, as observed by

immunofluorescence staining of the inhibitor treated mice liver sections and immunoblotting with

histones extracted from epididymal fat pads. This report therefore not only for the first time causally

links histone butyrylation with adipogenesis but also presents a novel small molecule modulator that

could be developed for anti-obesity therapeutics. Initial preclinical studies indicate that LTK-14A can be

a safe candidate drug for treatment of obesity.

Keywords: Acetylation; adipogenesis; butyrylation; garcinol; p300

Introduction

Chromatin function is spatially and temporally regulated by several processes like incorporation of histone variants, remodeling of nucleosomes by ATP-dependent remodelers and histone post-translational modifications. Interestingly, epigenetics and the metabolic state of the cell are intimately intertwined because the enzymes catalyzing the modifications use co-factors derived from several metabolic reactions (1). In recent times, advancements made in high sensitivity mass spectrometry have led to the discovery of new acylation modifications that structurally resemble acetylation, but differ in hydrocarbon chain length, hydrophobicity and charge (2). Emerging evidence indicates that these modifications occur dynamically on histones and could either have distinct functions from acetylation, or, have a similar but more profound impact on chromatin de-compaction, thereby amplifying the transcriptional output (3). EP300 (p300) is the master acyltransferase that has been found to promiscuously acylate histones using diverse CoA derivatives (2).

Due to the rising prevalence of obesity in the modern day society, there has been increased research on the molecular mechanisms behind its manifestation. At the core of obesity lies the adipose tissue which regulates energy homeostasis and metabolism. Hence, the developmental process of adipose tissue i.e. adipogenesis and its dysregulation needs to be studied for clinical management of obesity related disorders. The process of adipogenesis is accompanied by a significant increase in global histone acetylation levels, along with a rise in expression of p300 and a concomitant decrease in the levels of several histone deacetylases (4). Regulation of gene expression is dictated by the fluctuating activities of histone acetyltransferases and histone deacetylases. Moreover, the importance of histone acetylation for adipogenesis has been further underlined by the finding that knockdown of ATP citrate lyase, an enzyme responsible for the regeneration of acetyl CoA in the cytoplasm and nucleus, leads to decrease in histone acetylation and also suppression of adipogenesis (5). Therefore, it can be assumed that p300

plays a crucial role in promoting the expression of adipogenesis related genes by acetylating the

histones in the corresponding gene regulatory regions, especially the promoters. Amongst the different

coenzyme A derivatives, butyric acid is an important short chain fatty acid and an intermediate of fatty

acid synthesis/breakdown processes. p300 has been known to carry out butyrylation of histones which

has been causally linked to the process of spermatogenesis (3).

In this study we report that histone butyrylation is critical in adipoogenesis, probed by the first known

acylation specific small molecule inhibitor of p300. The newly discovered importance of histone

butyrylation in adipogenesis makes it an attractive target for small molecule compound mediated

inhibition of adipogenesis, thereby controlling obesity with lesser collateral damage.

Materials and Methods

Animals and treatment for obesity related studies: Male C57BL/6J mice (8 weeks old) were bred in

house and acclimatized for 1 week prior to experiments. Mice were maintained on a 12 hour light/dark

cycle and given ad libitum access to food and water. Experiments were carried out in accordance with

the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA)

guidelines and were approved by the Institutional Animal Ethics Committee of JNCASR

(201/GO/Re/S/2000/CPCSEA dated 1st June, 2000 Renewed in Aug 2015). The age-matched animals

were maintained on a standard laboratory chow diet (AMRUT   M/s Krishna Valley Agrotech ,

Maharashtra) or high fat diet (60 kcal% fat) (Research Diets, Cat. No. D12492) with or without LTK-14A

(50 mg/Kg body weight) and water. The compound was mixed with the high fat diet according to dosage

requirements in the laboratory every week. Body weight and food intake was measured twice every

week.

For the experiment on genetically obese mice, male db/db mice and db+/- mice (8-10 weeks old) bred in

house were obtained and acclimatized for 1 week prior to experiments. Mice were maintained on a 12-

hour light/dark cycle at 22+/-3 ⬚C and a relative humidity of 55% and given ad libitum access to food and

water. The experiments were carried out in accordance with the Committee for the Purpose of Control

and Supervision of Experiments on Animals (CPCSEA) guidelines, Government of India and was approved

by the Institutional Animal Ethics Committee of CSIR-CDRI, Lucknow (IAEC/2020/115/Renew-

0/SI.No.09). The age-matched animals were maintained on a standard laboratory pelleted diet (Altromin

International, Cat. No. 1320) and water. db/db mice were oral gavaged (50 mg/kg) with LTK14A in 0.5%

carboxymethyl cellulose, equal volume of the vehicle was orally gavaged to control db/db mice for a

total period of 30 days. db+/- animals were kept as untreated normal control group. The body weights

and feed intake was recorded every week throughout the experimental period. The lean and fat mass of

the mice were determined using Echo MRI in live mice on day 15th and day 30th.

Cell culture: The mammalian cell line 3T3L1 was obtained from ATCC (American Type Culture Collection)

and was grown in DMEM (Dulbecco's Modified Eagle's Medium). The growth medium was

supplemented with 10% FBS (Fetal Bovine Serum). The cell line was grown in 37°C incubator with 5%

CO_2 and 90% relative humidity.

siRNA experiments: ON-TARGETplus siRNA SMART pools against mouse Acss2 (catalog no.L-065412-01-

0005) was purchased from Dharmacon (Dharmacon, Lafayette, CO, USA). 3T3L1 cells were transfected

with Acss2 siRNA at 25 nM concentration, using RNAiMAX (catalog no. 13778-150, Thermo Fisher

Scientific) according to manufacturer's protocols. Transfection was carried out for 48 hours after which

cells were induced for differentiation for 6 days followed by RNA isolation and histone extraction.

Adipogenesis and Oil Red-O staining: 3T3L1 preadipocytes were grown to confluency and 24 hours

post-confluence, cells were incubated with differentiation medium (growth medium containing 1µM

Dexamethasone, 0.5 mM IBMX and 1 μg/ml insulin)(Sigma Aldrich) for 2 days in the presence/absence

of LTK-14A. Then the cells were supplemented with maintenance medium with/without LTK-14A

(growth medium with 1 μg/ml insulin only) every alternate day. On the 6th day post induction of

differentiation the cells were fixed with 3.7% formaldehyde solution followed by staining with Oil Red-O

dye (Sigma-Aldrich) (prepared in 60% isopropanol) for 1 hour at room temperature. The incorporated

dye was extracted using isopropanol and absorbance values were determined spectrophotometrically at

510 nm to estimate total amount of lipid accumulation under different experimental conditions.

Acid extraction of histones and immunoblotting: 3T3L1 cells were washed with ice-cold PBS and then

resuspended in Triton extraction buffer (PBS containing 0.5% Triton X-100 (v/v) and 2 mM PMSF). The

cells were allowed to lyse on ice for 10 minutes, spun down, washed with PBS and the process was

repeated another time. Then the proteins were acid extracted with 0.2 N HCl for two hours on ice. The

debris was spun down and then histones were precipitated by incubating the supernatant with 33%

trichloroacetic acid (Sigma-Aldrich) at 4⬚C for 30 minutes. The histones were then pelleted down and

then given two washes with acetone. The pellet was allowed to air-dry for 10 minutes and then

resuspended with 50 mM Tris-Cl, pH 7.4. Extraction of histones from epididymal fat pads was

performed by mechanically homogenizing the adipose tissues in Triton extraction buffer and then

following the same subsequent steps as mentioned before, except for the step of acid extraction with

HCl, in which the incubation was performed for four hours on ice.

The samples were then run in 15% SDS-PAGE, transferred onto PVDF membrane (Milipore) in transfer

buffer (25 mM Tris, 192 mM glycine, 0.036% SDS in 20% methanol), blocked with 5% skimmed milk

solution to prevent non-specific antibody binding and finally probed with antibodies against acetylation

and butyrylation marks prepared in 1% skimmed milk or FBS solution (5% BSA, 1x TBS, 0.1% Tween-20).

HRP-conjugated secondary antibodies (Abcam) were used against the primary antibodies to obtain a

chemiluminiscent signal in the presence of SuperSignal West Pico Chemiluminiscent Substrate (Thermo

Scientific) which was developed using the VersaDoc imaging system (Biorad).

Synthesis of LTK14A: Isolation of garcinol from Garcinia indica and synthesis of isogarcinol from it were

carried out following the protocols as mentioned in Mantelingu et al, (2007) (11). For LTK-14A synthesis,

anhydrous potassium carbonate (0.41 mmol) was added to a stirred solution of isogarcinol (100 mg, 0.16

mmol) in dry acetone. The mixture was stirred for 1 hour at room temperature followed by dropwise

addition of dimethyl sulfate (0.41 mmol) under nitrogen gas atmosphere. The resulting reaction mixture

was further stirred at room temperature for 14 hours. Progress of the reaction was monitored through

thin layer chromatography. Acetone was evaporated from the reaction mass, ice was added and

acidified with 5% aqueous HCl (pH ~ 5-6). A pale brown precipitate was observed, which was filtered and

washed with cold water. The dry mass was then chromatographed on silica gel using 5-6% ethyl acetate

in hexane as eluting solvent mixture. 720 mg (69%) LTK14A was isolated as white solid. NMR spectral

analysis was consistent with the desired product (Sup. Fig. 3A). A 20 mg sample was dissolved in hot

methanol and placed in a flat surface glass vial with access to very slow solvent evaporation. Crystals as

colorless needles were isolated. The purity of the synthesized LTK-14A was found to be more than 98%

as verified by HPLC (Sup. Fig. 3C) and only the protonated species of LTK-14A were observed in the ESI

spectra of the corresponding fraction (Sup. Fig. 3B).

X-ray crystal structure determination for LTK14A (CCDC: 1969185): Suitable single crystal with

approximate dimensions of 0.35 × 0.18 × 0.15 mm3 was used for X-ray diffraction analyses by mounting

on the tip of a glass fibre in air. Data were collected on a Bruker kappa apex 2 with Mo Kα (λ=0.71073 Å)

at 296 (2) K. The structure was solved by direct method using program SHELXL-97 and subsequent Fast

Fourier Transform technique. Crystallographic data and experimental details for LTK14A are summarized

in Table 1 and Sup. Fig. 3D.

Untargeted analysis of intracellular metabolites by Ultra-performance liquid chromatography coupled

with time-of-flight mass spectrometer (Q-TOF LC/MS): Profiling of intracellular metabolites was

performed on an agilent 1290 Infinity LC system coupled to Agilent 6545 Accurate-Mass Quadrupole

Time-of- Flight (QTOF) with Agilent Jet Stream Thermal Gradient Technology. The UPLC system was

assembled with a Diode array detector (DAD) and autosampler. The Chromatographic separation was

achieved on Agilent ZORBAX SB- C18 column (2.1 × 100 mm, 1.8 μm) as stationary phase. The mobile

phase consisted of a linear gradient of 100 mM ammonium formate (A) and Acetonitrile (B): 0–10.0 min,

30-80% B (v/v); 10.0–15.0 min, 80–100% B (v/v); 15.0–20.0 min, 100% B (v/v); 20.0–21.0 min, 100–30%

B (v/v); 21.0–25.0 min, 30% B. The sample was dissolved in 1 ml methanol (LCMS Grade), centrifuged

and supernatant was taken for UPLC-QTOF-MS analysis. The column was reconditioned for 5 minutes

prior to the next injection. The flow rate was 0.5 ml/min, and the injected volume was 20 μL. The UPLC

was connected to the MS analysis was performed on an Agilent 6545 Accurate-Mass Q-TOF/MS system

with an electrospray ionization (ESI) source. Considering the MS conditions, positive ion mode was used

to obtain high-resolution mass spectra. The ESI source parameters were: drying gas (N2) flow, 8 L/min;

drying gas temperature, 200°C. Other parameters were set as nebuliser gas, 35 psig; capillary voltage,

3000 V; skimmer voltage, 65 V; nozzle voltage 1000 V and fragmentor voltage 150 V. The data

acquisition on the LC-QTOF was performed using Agilent Mass Hunter Acquisition software (Agilent

Technologies, Santa Clara, CA, USA). The data were deconvoluted into individual chemical peaks with

Agilent MassHunter Qualitative Analysis (MassHunterQual, Agilent Technologies).

Site directed mutagenesis: Point mutation was introduced into the p300 catalytic domain sequence

cloned within pET-28b plasmid by using the Quikchange site directed mutagenesis kit (Agilent). Briefly,

the template was mixed with Pfuturbo DNA polymerase, designed forward and reverse primers and

dNTPs for PCR (Initial denaturation for 30 seconds at 95°C; Cycles: 95°C 30'', 55°C for 60'', 68°C for 7';

final elongation at 68°C for 5'.) The mixture was then incubated with DpnI enzyme for 2.5 hours for

removal of the parent wild type sequence followed by transformation of competent XL10Gold cells and

kanamycin selection of positive transformants. The inserted mutation was verified by sequencing and

alignment with the wild type sequence using ClustalOmega software. The sequences of the primers used

are as follows:

Forward primer- 5'- CCTCACTTGGTGGAGCTGCCCAAAATATGCCCTGTTGTG – 3'

Reverse primer- 5'- CACAACAGGGCATATTTGGGCAGCTCCACCAAGTGAGG -3'

Purification of recombinant full-length p300 and p300 catalytic domain: Polyhistidine tagged

recombinant p300 was purified from Sf21 insect cells by Ni-NTA based affinity chromatography. These

cells were infected by the baculovirus containing p300 for 72 hours. Once the cells showed signs of

infection, they were scraped off and centrifuged to separate the spent media. The cell pellets were

resuspended in ice cold homogenisation buffer (20 mM Tris, pH 7.5; 20 mM imidazole, 10% glycerol, 0.2

mM EDTA, 300 mM KCl, 0.1%NP-40, 2 mM PMSF, 2 mM βmercaptoethanol) and then homogenised

using a Dounce homogeniser (5 cycles, 6 strokes per cycle with 5 minutes interval between each cycle).

The cell debris was separated by centrifugation and then the lysate was incubated with pre-equilibriated

Ni-NTA slurry for 2.5 hours at 4°C on an end-to-end rotor. Then the beads were washed with wash

buffer (20 mM Tris, pH 7.5; 40 mM imidazole) nine times to remove unwanted contaminants. Finally

p300 was eluted using elution buffer (20 mM Tris, pH 7.5; 250 mM imidazole).

Polyhistidine tagged p300 catalytic domain cloned in pET28b plasmid and Sirt2 were used for co-

transforming BL21(DE3) cells. At first, 50 ml primary culture was prepared in Luria broth containing

kanamycin (50 μg/ml) and ampicillin (100 μg/ml) (HIMEDIA) and then it was scaled up 10 times to

prepare secondary culture. Incubation was done at 37°C, 180 rpm till O.D. reached 0.37 followed by

induction with 0.2 mM IPTG and a further incubation at 30°C for 4 hours. Cells were pelleted at 6000

rpm, 4°C for 10 minutes and stored at -80°C. Resuspension and homogenisation of the pellet was

carried out in homogenisation buffer (20 mM Tris, pH 7.5; 20 mM imidazole, 10% glycerol, 0.2 mM

EDTA, 300 mM KCl, 0.1%NP-40, 2 mM PMSF, 2 mM βmercaptoethanol). 3 cycles of sonication were done

for 3 minutes at 35% amplitude with 5 minutes interval between each cycle. The cell debris were

separated by pelleting at 12000 rpm for 30 minutes at 4°C. The supernatant was equilibriated with Ni-

NTA beads slurry (Novagen) in an end-to-end rotor for 3 hours at 4 C. Eight washes were given with

wash buffer (20 mM Tris, pH 7.5; 40 mM imidazole). Elution was carried out in batches with elution

buffer (20 mM Tris, pH 7.5; 250 mM imidazole).

The enzymatic activity of the proteins was estimated by filter binding assay.

In vitro **filter binding assay:** Different dilutions of purified enzyme were incubated in reaction buffer (50

mM Tris-HCl pH 7.5, 1 mM DTT, 10 % Glycerol, 2 mM PMSF) with 10 mM NaBu, tritiated acetyl CoA (50

μCi) (PerkinElmer, Part No: NET290050UC) and Xenopus histone H3 (500 ng) for 30 minutes at 30°C. The

reaction mixture was then spotted on P81 filter paper and the paper was allowed to dry at room

temperature. The paper was washed in wash buffer (50 mM sodium bicarbonate and 50 mM sodium

carbonate), dried at 90⬚C and immersed in enhancing solution (2.5% PPO and 0.25% POPOP in toluene)

for taking radioactive counts using liquid scintillation counter (Perkin Elmer, MicroBeta2 system).

***In vitro* acylation (acetylation/butyrylation) assay**: Acylation reactions were performed in reaction

buffer (25 mM Tris-HCl pH 7.5, 100 mM NaCl, 0.1 mM EDTA, 1 mM DTT, 10 % Glycerol, 1x PIC) with 100

ng/mL TSA, and 50 μM Butyryl-CoA/Acetyl CoA. Xenopus histone H3 (1 μg) was used as the substrate

and full length p300 or p300 catalytic domain (10,000 cpm activity) was used as the enzyme. Reactions

were incubated in presence or absence of butyrylation inhibitor for 10 minutes at 30 °C, followed by

initiation of the acetylation/butyrylation reaction by the addition of acetyl CoA/butyryl CoA. After a

further 10 minutes, reactions were stopped by addition of Laemmli buffer and samples were used for

immunoblotting.

Molecular docking analysis: Glide module of Schrodinger suite was used to dock LTK-14 and LTK-14A to

p300 catalytic domain (PDB: 3BIY). Chimera was used as visualizing software. The protein was prepared

using the wizard tool in Maestro version 10.2.The crystal structure of LTK-14A (CCDC 1969185) and LTK-

14 (CCDC 645420) were used as the most optimized and energy minimized ligand structures. The

receptor grid for p300 catalytic domain was generated by specifying the binding (active) site residues

using Receptor grid generation in the Schrodinger Glide application. Once the receptor grid was

generated, the ligands LTK-14 and LTK-14A were docked to the p300 catalytic domain using Glide

version 5.8 molecular docking.

Chromatin Immunoprecipitation: Nuclear ChIP was performed using intact nuclei from cells after

removal of lipids. 3T3L1 that were either undifferentiated or differentiated for 3 to 6 days were cross-

linked using 1% formaldehyde. Nuclear fractions were obtained by sequentially lysing the cells in two

buffers- lysis buffer 1 (140 mM NaCl, 1 mM EDTA, 50 mM HEPES, 10% Glycerol, 0.5% NP-40, 0.25%

Triton X-100) and lysis buffer 2 (10 mM Tris pH 8, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA). This was

followed by lysis in SDS lysis buffer (1% SDS, 10 mM EDTA, 50 mM Tris-HCl pH 8). Lysates were

sonicated in a Diagenode Bioruptor (Liège, Belgium) to produce 200 to 1000 bp DNA fragments. The cell

lysates were incubated with H4K5 Bu antibody (PTM Biolabs) or preimmune IgG per sample and 25 µl of

magnetic protein G Dynabeads (catalog no. 10003D; Novex, Waltham, MA) overnight at 4°C. Beads were

then washed successively with low-salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl

pH 8, and 150 mM NaCl), high-salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH

8, and 500 mM NaCl), LiCl buffer (250 mM LiCl, 1% NP-40, 1% NaDOC, 1 mM EDTA, and 10 mM Tris-

HCl, pH 8), and TE (10 mM Tris-HCl pH 8 and 1 mM EDTA). DNA-protein complexes were recovered

from beads in elution buffer (0.1% SDS and 100 mM NaHCO3). The ChIP eluates and input samples in

the elution buffer were then reverse cross-linked by adding 200 mM NaCl and 20 µg proteinase K

(Sigma) and incubating at 65°C for 4 h. Subsequently, 20 µg of RNase A (Sigma) were added and the

samples were further incubated for 15 min at 37°C. The immunoprecipitated DNA was extracted using

phenol-chloroform, ethanol precipitated, and used for quantitative PCR. The region-specific primer sets

used for the ChIP-qPCR analysis have been mentioned in Table 3. These primers were designed based on

the genomic coordinates with enriched H3K27 acetylation in the vicinity of promoters of candidate

genes in brown adipose tissue, the information for which was obtained from NCBI37/mm9 assembly

(ENCODE/LICR).

Quantitative real time PCR: Total RNA was isolated from the 3T3L1 cells and epididymal fat pads by

phenol-chloroform method. In case of LTK-14A treatment, 3T3L1 cells were allowed to differentiate for

7 days following the standard adipogenesis protocol, in presence of the compound at 25 µM

concentration. The control experiments were 3T3L1 cells differentiated with or without DMSO as

solvent control and differentiation was done only with the addition of only the cocktail of chemicals for

adipogenesis. RNA was isolated from the cells after the period of 7 days in all three cases. For isolation

of RNA from mice fat pads, the adiose tissue samples were lysed in Trizol using a mechanical

homogenizer followed by phenol-chloroform method of extraction. The isolated RNA samples were

treated with 2 units/μL DNase I to remove any residual DNA followed by ethanol precipitation at -80□C.

2μg RNA was used for cDNA synthesis using MMLV reverse transcriptase (Sigma) in presence of 0.5 mM

dNTP and 3.5 μM oligo dT (Sigma). RT-qPCR amplification was done in kappa SYBR green reagent

(Biosystems) with the help of Real time PCR instrument (Applied Biosystems by Life Technologies).For

estimation of relative fold change in RNA expression, the following calculation was done: Relative fold

change = $2^{-\Delta\Delta Ct}$, where ΔΔCt=ΔCt value of sample − ΔCt value of control and ΔCt= Ct value of target − Ct

value of internal control actin for individual sample. The sequences of primers are provided in Table 4.

Transcriptomics analysis by RNA-seq: RNA integrity was checked by Agilent Bioanalyzer 2100, only

samples with clean rRNA peaks were used. Libraries for RNA-seq were prepared according to KAPA

Stranded RNA-Seq Kit with RiboErase (KAPA Biosystems, Wilmington, MA) system. Final library quality

and quantity were analyzed by Agilent Bioanalyzer 2100 and Life Technologies Qubit3.0 Fluorometer,

respectively. 150 bp Paired-end sequencing was performed on Illumina HiSeq 4000 (Illumnia Inc., San

Diego, CA).

House mouse (Mus musculus, strain C57BL/6J) genome (mm10) was downloaded from GENCODE and

indexed using Bowtie2-build with default parameters. Adapter removal was done using Trim Galore (v

0.4.4) and each of the raw Fastq files were passed through a quality check using FastQC. PCR duplicates

were removed using the Samtools 1.3.1 with the help of 'rmdup' option. Each of the raw files was then

aligned to mm10 genome assembly using TopHat2 with default parameters for paired-end sequencing

as described in (28). After aligning, quantification of transcripts was performed using Cufflinks and then

Cuffmerge was used to create merged transcriptome annotation. Finally differentially expressed (DE)

genes were identified using Cuffdiff. The threshold for DE genes was log2 (fold change) >1.5 for up

regulated genes and log2 (fold change) <1.5 for down regulated genes with p-value <0.05.

GO enrichment analysis

Gene Ontology (GO) analysis was performed in PANTHER (29). Significant enrichment test was

performed with the set of differentially expressed genes in PANTHER and Bonferroni correction method

was applied to get the best result of significantly enriched biological processes.

Fisher's exact test

Fisher's exact test was performed in PANTHER Gene Ontology (GO) where p-value significance was

calculated based on the ratio of obtained number of genes to the expected number of genes (O/E)

considering the total number of genes for the respective pathway in Mus musculus with a FDR of <0.05.

Heatmap and clustering of genes

Unsupervised hierarchical clustering method was performed using Cluster 3.0 (30) with Pearson

Correlation and average linkage rule. Gene expression data (FPKM of all samples) was taken and log2

transformed. Low expressed (FPKM<0.05) and invariant genes were removed. Then genes were

centered and clustering was performed based on differential expression pattern of genes and fold

change. Finally, the heatmap was visualized in Java TreeView 3.0.

Biological analysis of differentially expressed transcripts and pathway regulatory network modeling

Statistically significant differentially expressed transcripts were subjected to GO and Pathway

enrichment using DAVID tool. Only those GO and pathways with a FDR score of <=0.05 was considered

for further downstream analysis. Key biologically dysregulated GO and Pathways along with the

differentially expressed genes was provided as an input to Pathreg algorithm from Theomics

International Pvt Ltd, Bangalore, India for gene regulatory network modeling. The result (nodes and

edges) of the Pathreg algorithm was provided as an input to Cytoscape v2.8.2 to identify key nodes and

edges that could be representative of the gene regulatory changes upon treatment.

Hematoxylin and eosin staining of liver and adipose tissues: Staining was performed on paraffin

embedded sections of liver and epididymal fat pads having the thickness of 5 μm and 10 μm

respectively. Briefly, the sections were deparaffinized in xylene, followed by immersion in absolute

alcohol, 90% ethanol, 70% ethanol, 50% ethanol and water before staining with hematoxylin for 5-10

minutes. The sections were immersed in above mentioned solutions in reverse order before counter-

staining in eosin for 15 seconds. Then the samples were dipped in absolute ethanol and xylene before

mounting with DPX.

Immunohistochemistry: Collected liver tissue samples were stored in 4% paraformaldehyde for 24

hours after which they were cryoprotected in 30% sucrose solution for two weeks. Cryosections were

performed at 7 μm sections using Cryostat LeicaCM1850 UV. For staining, the tissue sections were

washed with PBS followed by antigen retrieval with 0.01 M citrate buffer (pH 6). The tissues were then

permeabilised with 0.3% Triton X-100/PBS (PBST) and blocked with 2% serum followed by incubation

with primary antibodies overnight at 4⬚C. The next day, secondary antibody incubation was carried out

for one hour at room temperature followed by staining of the nuclei with Hoechst 33342 and mounting

with 70% glycerol. Images were acquired by confocal microscopy. For quantitation, intensity of each

modification specific staining was normalized with respect to the Hoechst staining intensity.

MTT assay: 12500 cells/ml were seeded in presence/absence of LTK-14A in 96-well plate and at different intervals of time (2 days, 4 days, 6 days) , 1/10th volume of 5mg/ml MTT solution was added to it followed by incubation at 37 °C for 3 hours. The formed crystals were dissolved in DMSO and spectrometric estimation of colour intensity was done at 570 nm.

Evaluation of mutagenecity by plate incorporation assay: Five doses of LTK-14A (10.0μg, 33.0μg, 100.0μg, 333.0μg and 1000.0μg/plate) were selected to check for its possible mutagenecity. Seven groups decided as Group 1- Control, Group 2 Positive Control and 5 doses of LTK-14A (Group 3 to Group 7: 10.0μg, 33.0μg, 100.0μg, 333.0μg and 1000.0μg/plate) were tested. Four tester strains of *Salmonella typhimurium* (TE 97a, TA 98, TA 100, TA102) were used and their respective positive controls were used in absence of S9 fraction (Table 5); 2-aminoanthracene was used as the positive control in presence of S9 fraction (Table 5). The tester strains were grown overnight in Luria Broth antibiotic selection medium at 37°C. LTK-14A prepared in DMSO was mixed with the bacterial culture, 0.5mM Histidine-Biotin and top agar in a culture tube and the content was overlaid on to minimal glucose plate. In case of plate incorporation assay with S9 mix, LTK-14A was incubated with S9 mix in cold condition for 15 to 20 minutes followed by the same procedure as the plate incorporation assay without S9. The plates were incubated in an inverted position in 37°C incubators for 48 hours following which the number of revertants was counted using automated colony counter.

Single dose toxicity study: The single dose toxicity study on LTK-14A was conducted in accordance with Good Laboratory Practice Principles as published by the OECD in 1998, in accordance with the Drugs and Cosmetics Rule (New Drugs and Clinical trials, 2019/ DCGI). LTK-14A (formulated in vehicle i.e. 0.5%

sodium salt of carboxymethyl cellulose) was administered to Sprague Dawley rats once by oral route on

the day of scheduled treatment. The three treatment groups were 250 mg, 500 mg, and 1000 mg/kg

body weight, while vehicle control was administered to control group. The rat equivalent doses were

calculated from the dosage used in mice experiments following FDA guidelines for drug administration.

The doses finally administered were 10, 20 and 40 times the corresponding mice equivalent dose. Food,

but not water, was removed thus letting the animals fast over night prior to dosing. After administration

of LTK-14A, food was withheld for further 2-3 hours and the animals were observed at 30 minutes, 1

hour, 2 hours and 4 hours from the time of dosing. Animals were subsequently observed once daily for

any signs of toxicity, mortality, or any other observations including conditions of skin, hair coat, eyes,

mucous membrane, movement, activity, tremors, convulsions, salivation, diarrhea, posture, piloerection

and lethargy. Food intake, water intake and body weight of animals was recorded weekly. Terminal

sacrifice of surviving animals was performed after 14 days using CO_2 euthanasia technique following

which animals were necropsied. Animals were fasted over night before necropsy. Gross observations

were recorded.

Reagents and antibodies: The media used for cell culture was purchased from Sigma-Aldrich; DMEM

(Ref: 1152). Several other cell culture reagents were obtained from HIMEDIA, such as PBS (Ref: TL1006),

Trypsin-EDTA solution 10X (Ref: TCL070) and Antibiotic antimyotic solution 100X (Ref: A002A). Multiple

reagents used for cell culture were obtained from Sigma-Aldrich: Insulin (Ref: I6634), Dexamethasone

(Ref: D4902), IBMX (Ref: I5879). TRIzol reagent was obtained from ambion life technologies (Ref:

15596018). The commercial antibodies used in this study are Pan anti-butyryllysine (PTM Biolabs, SKU:

PTM 301), Butyryl-Histone H4 (Lys 12) rabbit pAb (PTM Biolabs, SKU: PTM 308), Butyryl-Histone H3 (Lys

9) rabbit pAb (PTM Biolabs, SKU: PTM 305), Butyryl-Histone H3 (Lys 23) mouse mAb (PTM Biolabs, SKU:

PTM 307), Butyryl-Histone H4 (Lys 8) rabbit pAb (PTM Biolabs, SKU: PTM 311), Butyryl-Histone H4 (Lys 5)

rabbit pAb (PTM Biolabs, SKU: PTM 313), Butyryl-Histone H3 (Lys 27) rabbit pAb (PTM Biolabs, SKU: PTM

315).

Statistical analysis: All statistical analyses were performed using GraphPad Prism 7 Software (California,

USA). Data obtained from two or three individual experiments as mentioned in figure legends, were

expressed as mean +/- SD. Either two-tailed unpaired Student's t-test or one-way ANOVA with

Dunnett's/Bonferroni's multiple comparision was used to determine the statistical significance values.

The mutagenecity analysis data was analyzed by comparing the groups using One-way ANOVA followed

by Neman-Keuls test. Body weight data for acute toxicity test were summarized in Mean ± SD and

compared by two factor (Group and Period) repeated measure analysis of variance (RM ANOVA) and the

significance of mean difference within (intra) and between (inter) the groups was done by Newman-

Keuls post hoc test. A P-value of equal to or less than 0.05 was considered statistically significant.

Acknowledgements: TKK is a recipient of JC Bose fellowship from Dept. of Science and Technology,

Govt. of India (SR/S2/JCB-28/2010). AB is a senior research fellow of UGC. This work was supported by

Life Science Education & Training at JNCASR grant (DBT/INF/22/SP27679/2018). We acknowledge Dr.

Rahul Gajbhiye from NIPER, Kolkata for the assistance in mass spectrometry related experiments in their

central instrument facility. We thank Dr. R.G. Prakash for providing the necessary facilities to perform

the mice experiment in Animal Facility, JNCASR and the confocal facility, JNCASR for the

immunofluorescence imaging analysis. We also thank Kruthi HT for helping in preparation of the figures

and graphical abstract. We further acknowledge the Sophisticated Analytical Instrument Facility and

Research (SAIFR, Lucknow) for providing spectral data. This manuscript bears CDRI communication

number 2021/TK.

Conflict of interest: Authors declare no conflict of interest.

Author Contributions: AB and TKK conceived the project. AB designed and performed all the biological

experiments. SC performed initial synthesis of all compounds and molecular docking analysis. KVS

performed bulk scale synthesis of LTK-14A for animal experiments. AKS and RG assisted in the animal

experiments. UB performed *in silico* analysis of high throughput RNA sequencing data. AN did the

toxicity evaluation and SKR evaluated the mutagenecity. NN provided the raw material *Garcinia indica*

for compound isolation. AB and TKK analyzed the data and wrote the manuscript.

Results

Physiological levels of butyryl CoA and histone butyrylation increases during adipogenesis:

Under normal circumstances, the cellular levels of specific coenzyme A derivatives are much less than

acetyl CoA (6).We surmised that the situation might alter in particular cases where metabolic pathways

promoting the production of other acyl CoA derivatives become more predominant. One such possible

scenario could be the process of adipogenesis in which fatty acid synthesis and breakdown can give rise

to several intermediate products like butyryl CoA.

The causal relationship of butyryl CoA and adipogenesis was investigated using 3T3L1 pre-adipocyte cell

line as model system (fig 1A). Our RT-qPCR data showed a significant increase in the expression of *Acss2*

(acyl-CoA synthetase short-chain family member 2) upon differentiation (fig 1B). Previously, ACSS2 had

been established as the enzyme responsible for generation of acyl-CoA derivatives from their

corresponding short chain fatty acids (7). Indeed, siRNA mediated knockdown of *Acss2* (fig 1C) led to a

drastic reduction in histone butyrylation with relatively lesser effect on acetylation (fig 1D). This

indicated that the ACSS2 generated butyryl CoA might be playing an important physiological role in

adipogenesis. These findings prompted us to investigate whether adipocyte differentiation also led to an

increased butyrylation of proteins. We observed that in accordance with the previous report (4), there

was an increase in acetylation level at histone H3 lysine 9 (H3K9 Ac) in the differentiated cells as

compared to the undifferentiated cells (Fig. 1C, panel i). Interestingly immunoblotting with pan

butyryllysine and site specific antibodies (H3K9, H3K23, H3K27, H4K5, H4K8 and H4K12) against

butyrylation (Fig 1C, panel ii and iii) revealed that butyrylation at each of these sites also increased upon

differentiation. Chromatin immunoprecipitation with antibody against H4K5 butyrylation revealed that

the modification enrichment levels increased in promoters of different pro-adipogenic genes in a time

dependant manner. *Pparg*, the master regulator of adipogenesis, exhibited a steady increase in

butyrylation level in its promoter with the progress of differentiation (fig 1F, panel i) while butyrylation

increased only at the terminal stage of adipogenesis in the promoter of *Lep*, encoding the adipocyte

marker leptin which gets expressed in mature adipocytes (fig 1F, panel ii). Interestingly, the promoter of

Cebpd, which regulates *Pparg* expression did not show any enhanced butyrylation in its promoter on day

3 and instead, exhibited increased butyrylation on day 6 of adipogenesis (fig 1F, panel iii).   Collectively

these results showed that during adipogenesis, EP300 catalysed butyrylation of histones could be an

important epigenetic signature.

Identification of a small molecule that selectively inhibits p300 catalysed histone butyrylation *in vitro*:

In order to investigate the contribution of p300 catalyzed butyrylation per se in adipogenesis, we

adopted a chemical biology based approach. For this purpose, a series of derivatives of garcinol, a

polyisoprenylated benzophenone isolated from *Garcinia indica*, were synthesized (8). These derivatives,

unlike garcinol, were found to be non-toxic to cells and showed a high specificity towards the catalytic

activity inhibition of EP300. The derivatives were generated by internal cyclisation of garcinol with

further addition of one or two functional groups to the remaining free -OH groups- giving rise to mono

and disubstituted derivatives (Sup. fig. 1A). Amongst these, a few monosubstituted derivatives could

potently inhibit EP300 catalysed acetylation as well as butyrylation, while their disubstituted

counterparts were unable to inhibit acetylation (Sup. fig. 2 B, C and D). This was probably due to

increased steric hindrance brought about by the addition of an extra functional group that interfered

with the interaction between the compounds and EP300.

An inverse correlation between catalytic rates of acylation reactions by EP300 and increasing acyl chain

length had already been established previously (9). Based on this, we hypothesized since butyrylation is

catalysed by EP300 at a much slower rate than acetylation, one of the compounds that could not inhibit

EP300 catalysed acetylation might be able to inhibit the slower butyrylation reaction. Our *in vitro*

butyrylation assay using recombinant p300 and histone H3 as the substrate revealed a few potential

candidates (Sup. Fig. 2 E) as butyrylation inhibitors. Amongst them, LTK-14A, which is 13, 14 dimethoxy

isogarcinol, was capable of inhibiting full length EP300 catalysed butyrylation at a concentration of 25

µM (fig. 2 C and D), with minimal effect on EP300 catalysed acetylation in similar reaction conditions (fig.

2 A and B).

In order to understand the underlying mechanism of the selective inhibition by LTK-14A, we performed

molecular docking analysis of LTK-14A (Fig. 2 G) and the catalytic domain. To delineate this, we solved

the crystal structure of LTK-14A (Table 1, Sup. Fig.2A). LTK-14 (fig. 2E), a specific inhibitor of EP300

catalysed acetylation and butyrylation (Sup. Fig. 2 B and C) was used for further comparative analysis.

The energy minimized analysis revealed that both LTK-14 and LTK-14A could potentially bind with the

active site of the EP300 catalytic domain forming several hydrophobic interactions (Fig.2 F and H). LTK-

14, due to the presence of a free phenolic –OH group, could hydrogen bond with a carbonyl oxygen

atom present in the peptide backbone of S1396 residue. However, LTK-14A has all of its phenolic –OH

groups blocked and so it is incapable of forming such an interaction. This hydrogen bonding by LTK-14

could imply a stronger binding to the enzyme, while LTK-14A, by interacting with fewer residues

presumably inhibits only the slower butyrylation reaction and the interaction may not be strong enough

to deter the faster acetylation reaction.

Some of the residues with which LTK-14A could potentially form hydrophobic interaction have already

been reported before as being critical for targeting inhibition of the enzymatic activity of EP300 by the

first reported EP300 inhibitor lysyl CoA (10). These residues include C1438, Y1446, Y1397 and W1436. To

verify this observation we generated a point mutant of p300 catalytic domain in which cysteine 1438,

showing the highest propensity to interact with LTK-14A, was mutated to alanine. We found that the

C1438A catalytic domain is equally active as the wild-type catalytic domain in an *in vitro* lysine

acetyltransferase assay (Sup. Fig. 2B). We observed that the wild-type EP300 catalytic domain catalysed

butyrylation was inhibited by LTK-14A most prominently at 25 µM concentration while acetylation was

not inhibited (compare lane 6 in each of fig.2 I-L). In contrast, neither acetylation nor butyrylation

modification catalysed by the mutant EP300 catalytic domain could be inhibited by LTK-14A even at 25

µM concentration (compare lane 6 in each of fig. 2 M-P). Altogether, these data suggest that LTK-14A

indeed interacts with C1438 within the active site of the EP300 catalytic domain, and this residue is

critical for its interaction to bring about butyrylation reaction inhibition.

LTK-14A is a metabolically stable, cell permeable compound that significantly inhibits adipogenesis:

Upon establishing that LTK-14A is an EP300 acylation-specific inhibitor *in vitro*, its cellular effects were

tested using 3T3L1 cells. The quality of our compound used in all biological experiments was acceptably

good with more than 97% purity (Sup. Fig. 2 C, D and E). We found the LTK14A is highly cell-permeable

and metabolically stable as it could be detected by mass spectrometry analysis from metabolites

extracted from LTK-14A treated cells after six days (Fig. 3A). MTT assay showed that LTK-14A at 25 µM

concentration did not have any adverse effect on the cellular oxidoreductase activity, a signature of

metabolic output of cells, (Fig. 3B) suggesting that LTK-14A is non-toxic to 3T3L1 even after 6 days of

treatment. Oil Red-O staining of 3T3L1 cells differentiated in the presence of LTK-14A followed by

spectrophotometric quantitaion revealed significant inhibition of adipogenesis by the compound (Fig.

3C, D).

In order to understand the transcriptional mechanism of LTK14A-mediated butyrylation inhibition on

adipogenesis, adipogenic cells were treated with LTK-14A and subjected to transcriptome analysis. We

used untreated and solvent treated cells as control. Volcano plot (Fig. 3E) and heat map analysis (Fig. 3F)

revealed that upon treatment with the compound, a large number of genes were differentially

expressed compared to DMSO treated and untreated conditions. To better understand the LTK-14A

affected genes, we focused on the common genes that were differentially expressed in the two cases-

(1) LTK-14A versus no inhibitor and (2) LTK-14A versus DMSO. Among these common genes, 2200 genes

were found to be downregulated and 1309 genes were upregulated (Fig. 4A) with a majority of them to

be protein coding genes (Fig. 4B). Gene ontology analysis revealed that LTK14A treatment

predominantly affected the genes involved in lipid metabolism pathways (Fig. 4C, E) which were

minimally affected by DMSO alone (Fig. 4D). The gene regulatory networks and the respective pathways

were constructed with the major genes that were affected by LTK-14A treatment compared to DMSO

treated and untreated conditions (Fig. 4F, Sup. Fig.3A). They collectively depict how LTK-14A, by

inhibiting expression of a large number of genes that are interconnected via several pathways

associated with lipid metabolism, brings about a collapse in the adipogenesis process as a whole.

A few of the pro-adipogenic genes downregulated by LTK-14A, as indicated by RNA sequencing, were

selected for further validation (Sup. Fig. 3B). RT-qPCR analysis corroborated that except for *Cebpd*, most

of the other pro-adipogenic genes (*Pparg, Cebpa, Adipoq, Lep, Slc2a4, Fasn* and *Fabp4*) showed a

marked downregulation upon LTK-14A treatment (Fig. 4G). *Acss2* was also downregulated by LTK-14A

(Sup. Fig.4C). The upregulation of anti-adipogenic genes was also verified by performing RT-qPCR

analysis for two such candidate genes- *Kdm4c* and *Tead4* (Sup. Fig.3C). Collectively, the RNA sequencing

analysis and the validation of gene targets further convincingly suggested that LTK-14A can inhibit

adipogenesis in 3T3L1 cells, mainly by repressing the expression of important pro-adipogenic genes.

LTK-14A inhibits histone butyrylation in 3T3L1 cells, without affecting acetylation or other post-

translational modifications:

Furthermore we wanted to find out whether LTK-14A mediated downregulation of adipogenesis is

causally linked to the inhibition of histone butyrylation in the 3T3L1 cells. We observed that treatment

with LTK-14A inhibited overall butyrylation of histones (Fig. 5 A, B). Among the specific modification sites

tested, H3K23 and H4K5 butyrylation showed a significant decrease upon treatment with LTK-14A (Fig. 5

A, B) while butyrylation at other tested sites were affected minimally (Sup. Fig. 3D). Acetylation of

histones (H3K9, H3K18 and H4K12) was also found to be unaffected (Fig. 5 C, D) upon LTK-14A

treatment. To establish the specificity of the LTK14A, the effect of LTK-14A on lysine and arginine

methylation along with phosphorylation of histones were also investigated. We observed (Fig. 5 E, F) no

significant changes in these marks upon the compound treatment, thereby indicating that in the 3T3L1

cells, LTK-14A specifically inhibits butyrylation.

Inhibition of butyrylation attenuates weight gain of high fat diet fed C57BL6/J mice:

After observing the anti-adipogenic effect of LTK-14A on 3T3L1 cells, we investigated whether this

compound could also affect adipogenesis and obesity at an organismal level. For this purpose, C57BL6/J

mice were maintained either on a normal diet or a high fat diet with or without LTK-14A. In due course

of time it was observed that the mice maintained on a high fat diet along with LTK-14A had a weight

gaining trend intermediate between the normal diet and high fat diet fed mice (Fig.6A,B). Dissection of

the mice revealed that the fat accumulation in high fat diet maintained mice was the highest, followed

by those on high fat diet and LTK-14A, and then by those on normal diet (Fig. 6C). The epididymal fat

pads were separately weighed and again it was observed that weights of the fat pads from compound

treated mice were much less than that of high fat diet fed mice (Fig. 6 D,E). The diet consumption

pattern of the mice showed that the two groups on high fat diet with or without LTK-14A, had a similar

consumption rate indicating that the reduced weight gain of the compound treated mice is not due to

discrepancy in diet consumption (Table 2). Hematoxylin and eosin staining of adipose tissue and liver

(Fig. 6 F,G) showed signs of adipocyte hypertrophy and hepatic steatosis in high fat diet fed mice, which

were absent in the mice maintained on normal diet or high fat diet with LTK-14A.

Immunofluorescence staining of the liver tissue samples revealed that there was a significant increase in

H4K5 butyrylation levels in high fat diet fed mice compared to normal diet fed mice in which the

modification was barely detectable and it was almost completely inhibited upon LTK-14A treatment (Fig.

6H panel iii, Fig. 6I panel iii). On the other hand, no significant difference in H4K12 (Fig. 6H panel i, Fig. 6I

panel i) and H3K14 acetylation (Fig. 6H panel ii, Fig. 6I panel ii) levels could be observed across all the

three groups. Furthermore, histone modification pattern in the adipose tissues was studied with acid

extracted histones from the epididymal fat pads. H4K5 butyrylation was reduced in the LTK-14A treated

samples compared to those of high fat diet fed mice (Fig. 6 J, N) while H3K14, H3K18 and H4K12

acetylation levels were relatively unchanged across all the three groups (Fig. 6 J, K, L, M). The metabolic

stability of LTK-14A in mice was verified by its detection in ESI spectra of analytes extracted from the

liver of compound treated mice (Sup. Fig 3 E).

The butyrylation specific inhibitor reduces weight of genetically obese db/db mice:

The previous model system involved the use of LTK-14A as a prophylactic. To test whether LTK-14A

could also reduce obesity by inhibiting histone butyrylation, genetically obese homozygous mutant

db/db mice were used. Heterozygous mutant db+/- mice were taken as control. The hyperphagic db/db

mice were orally administered LTK-14A daily at the same dosage as before (50 mg/Kg body weight). We

observed that upon oral gavaging of LTK-14A, the overweight mice not only showed an arrest in weight

gain, but their body weights were significantly reduced by the end of one month of treatment (Fig. 7

A,B,C). The food intake of LTK-14A treated mice was marginally less than that of the vehicle treated mice

(Table 2, Supplementary Information), although both groups consumed more food compared to the

db+/- mice during the treatment regime. Average weight of the liver (Fig. 7 D,E) and epididymal fat pads

(Fig. 7 F,G) was the highest for db/db mice given vehicle control and least for the db+/- mice, while that

for the compound treated db/db mice was intermediate. *In vivo* Echo MRI of live mice revealed that

total adiposity i.e. fat mass of the mice was significantly lowered upon LTK-14A treatment (Fig. 7 H) and

lean mass was marginally affected (Fig. 7 I). Hematoxylin and eosin staining of adipose tissue and liver

(Fig. 7 J,K) showed signs of adipocyte hypertrophy and hepatocyte ballooning in vehicle treated db/db

mice, which were absent in the db+/- and LTK-14A treated db/db mice.

Similar to the high fat diet induced obesity model, the more obese homozygous leptin receptor mutant

mice (db/db) liver had greater level of H4K5 butyrylation compared to leaner heterozygous mutant mice

(db +/-) (Fig 7L panel iii, Fig. 7M panel iii). Moreover, LTK-14A treated db/db mice liver showed a

decrease in H4K5 butyrylation compared to vehicle control treated mice liver. In contrast, H4K12 (Fig. 7L

panel i, Fig. 7M panel i) and H3K14 acetylation (Fig. 7L panel ii, Fig. 7M panel ii) were only partially

upregulated in the db/db mice and did not get significantly reduced upon LTK-14A treatment.

Immunoblotting with acid extracted histones from epididymal fat pads showed that H4K5 butyrylation

was reduced in the adipose tissues of LTK-14A treated mice (Fig. 7 N,R) compared to vehicle control

treated obese mice. H3K14, H3K18 and H4K12 acetylation were not significantly altered in any of the

three groups (Fig. 7 N,O,P,Q). Finally, RT-qPCR analysis showed that just like in cellular model of

adipogenesis, LTK-14A treatment led to robust inhibition of important pro-adipogenic genes *Pparg*, *Lep*,

Slc2a4 and *Adipoq* in the fat pads of compound treated db/db mice (Fig. 7S). Collectively these data

establish that obesity can not only be prevented but also reverted by specific inhibition of H4K5

butyrylation, thereby highlighting the importance of this modification in the context of adipogenesis.

LTK-14A is non-mutagenic and non-toxic in nature for biological applications: Based on the

observations of obesity attenuating effects of LTK-14A in two different mice models, we performed

some preclinical tests to ensure that the obesity arrest was not as a result of any deleterious side effect

of toxicity from the compound. Plate incorporation assay was carried out in four different nutritional

auxotrophic mutants of Salmonella typhimurium (TA97a, TA98, TA100 & TA102)in absence or presence

of rat liver extract (S9 enzyme fraction). It was observed that the number of revertants obtained from

LTK-14A was similar to that from vehicle control and far less compared to that from the positive control

in all four strains, both in the absence and presence of liver extract (Sup. Fig 4 A-H). Furthermore, acute

toxicity test was performed in rats for checking whether a single administration of excessively high dose

of LTK-14A would alter the body weight and food/water consumption pattern of the rats. The body

weight changes in the LTK-14A administered rats were similar to the vehicle control treated ones (Table

6) with no gross morphological changes in any organ. Moreover, there was also no significant treatment

related effect on the food and water consumption of rats of both gender for all three doses compared to

vehicle control (Table 6).

Collectively these data imply that LTK-14A does not have deleterious toxic effects on the test animals

and could have a potential for downstream applications for tackling obesity related complications.

Discussion

Reports on the physiological relevance of acylation modifications are scanty. Breakthroughs have been made in establishing the importance of butyrylation in spermatogenesis (5), β-hydroxybutyrylation in starvation response (11) and crotonylation in inflammatory response, digestive system, cancer and HIV latency (7,12,13,14). Lysine propionylation has also been reported as an activation mark on histones that could be useful in various physiological processes (15). The list of these modifications seems to be increasing continuously, with the recent additions of histone serotonylation (16), lactylation (17) and dopaminylation (18). The fact that serotonin and dopamine are chemical neurotransmitters while lactate is derived from pyrvate, a glycolytic end product, and all these molecules bring about modification of histones thereby mediating gene expression in several tissues, bear testimony to the hypothesis that the abundance levels of cellular metabolites in different physiological contexts can lead to modification of histones with a plethora of diverse chemical moieties that can selectively modulate gene expression patterns.

Our initial experiments in 3T3L1 adipogenesis indicated an important role of ACSS2 in the differentiation process by generating butyryl CoA for histone butyrylation. Inside the cells ACL (ATP citrate lyase) and ACSS2 that have been implicated in generation of nuclear acetyl CoA. Both the enzymes are known for regulating gene expression in different contexts by acetylating histones in specific genomic loci (19-21). Knockdown of *Acss2* had a much more pronounced effect on H4K5 butyrylation compared to H3K18 acetylation. This is probably due to the fact that ACSS2 is the sole/major producer of acyl CoA species while acetyl CoA can be generated by diverse means and knockdown of a single acetyl CoA producer was not enough to have a robust effect on global acetylation level of histones. Similar observations have already been made for histone crotonylation in HeLa cells (7).

Due to redundancy in the functionality of p300 catalyzed butyrylation with acetylation, the significance

of butyrylation in adipogenesis was demonstrated using a pharmacological approach. LTK-14A could

selectively and specifically inhibit histone butyrylation and still inhibit adipogenesis by downregulating

expression of many pro-adipogenic genes. However amongst these genes *CEBPd* expression did not get

affected by LTK-14A possibly because the expression of this gene is less prone to be regulated by histone

butyrylation in its promoter. Indeed, chromatin immunoprecipitation study indicated that in the initial

stage of induction of differentiation in pre-adipocytes, there was no enrichment of H4K5 butyrylation in

the promoter of *Cebpd*. This could explain why the butyrylation inhibitor LTK-14A treatment at the

induction stage did not inhibit *Cebpd* expression, since there was no buyrylation mark enrichment for

inhibition in its promoter to begin with. An increased butyrylation in its promoter at the terminal stage

could be either through p300 catalysed modification or non-enzymatic means with greater availability of

butyryl CoA, or both. Currently, we have not been able to verify this. However our preliminary

observations with promoter specific studies indicate that H4K5 butyrylation plays a role in regulating the

expression of the critical transcription factor *Pparg*. Inhibition of butyrylation by LTK-14A repressed the

expression of the master regulator which further led to the inhibition of downstream genes like *Lep*,

whose expression is dependent on PPARγ. While cell line based studies indicate that C/EBPδ regulates

the expression of a cascade of transcription factors like C/EBPα and PPARγ, this fact has been

questioned in knockout mice which have poorly differentiated adipose tissue but still express *Pparg* and

Cebpa (22). This indicates that there could be other factors playing a redundant function like C/EBPδ in

the initiation stage of adipogenesis. Moreover, an additional function of C/EBPδ may seem to be

downstream of C/EBPα and PPARγ at the terminal stage of adipogenesis. It has been speculated that this

could involve inducing the expression of ligands that activate PPARγ (23). Therefore, the increased

butyrylation in the promoter of *Cebpd* on day 6 of adipogenesis might be responsible for its role in the

terminal stage. The expression of *Cebpd* is independent of *Pparg* and hence repression of *Pparg* by

butyrylation inhibition with LTK-14A did not repress *Cebpd*.

Acss2, which encodes the enzyme responsible for butyryl CoA biosynthesis was also downregulated by

LTK-14A which might have further contributed to reduced butyrylation of histones during inhibition of

adipogenesis. Amongst the different histone modification patterns investigated, H4K5 butyrylation was

consistently affected by LTK-14A in different cell line and mice models, while the other site specific

butyrylation marks were relatively unaltered, indicating that there may be a hierarchy in the importance

of these modifications in regulation of adipogenesis. While our work was in progress, a new report was

published in which it was demonstrated that inhibition of fatty acid synthesis as well as oxidation leads

to reduction in H4K5 butyrylation while acetylation remains unaffected (24). This further reinforces our

own hypothesis that fatty acid metabolism is an important source for certain acyl CoAs such as butyryl

CoA, derived from the short chain fatty acid butyric acid. Another work published during this period

revealed HBO1, a member of MYST family of acetyltransferase, also possesses the ability to catalyse

acylation reactions including butyrylation (25). Mass spectrometry studies of histone modifications

altered by knockdown of HBO1 in HeLa cells showed that H4K5 butyrylation does not get affected by

depletion of HBO1 indicating that it is a p300 specific mark.

Global histone acetylation has previously been reported to remain unchanged in liver and adipose

tissues of mice even after high fat diet consumption (26), which was consistent with the observations we

made in both mice models studied. Since acetyl CoA is highly abundant, fluctuations in its levels did not

lead to major changes in histone acetylation pattern. Because the usual butyryl CoA levels are far less

than their corresponding Michaelis Menten constant for p300, an increase in butyryl CoA levels due to

excess lipid consumption through high fat diet or *de novo* lipid synthesis through excess food

consumption, resulted in a concomitant increase in butyrylation by p300 which served as the nutritional

sensor for butyryl CoA. Enhanced H4K5 butyrylation was not observed in mature adipose tissues of

obese mice possibly due to predominance of adipocyte population in the tissues of mice from all the

groups (27).

In the 21st century, obesity has become a major epidemic leading to several other disorders such as

type 2 diabetes and arteriosclerosis. Histone acetylation inhibition could be regarded as a possible

method for attenuating adipogenesis and obesity. But acetylation is a widespread phenomenon that is

required in several other physiological processes besides adipogenesis. Hence, treatment with an

acetyltransferase inhibitor at a particular range of concentration could have undesirable side effects and

toxicity. In contrast, butyrylation is a rarer event, repressing which may be less harmful to other

processes. Indeed, our preclinical tests are indicative of the fact that LTK-14A may be suitable for

development as a possible anti-obesity drug due to its negligible toxicity side effects. The present finding

thus establishes an importance of histone butyrylation in adipogenesis, making it an attractive target for

pharmacological inhibition of adipogenesis and thereby controlling obesity (Graphical abstract).

References

1. Haws, S. A., Leech, C. M., Denu, J. M. Metabolism and the Epigenome: A Dynamic Relationship.

Trends Biochem 2020; 45, 731-747

2. Chen, Y., Sprung, R., Tang, Y., Ball, H., Sangras, B., Kim et al Lysine propionylation and

butyrylation are novel post-translational modifications in histones. Mol Cell Proteomics 2007; **6**:

812-819

3. Goudarzi, A., Zhang, D., Huang, H., Barral, S., Kwon, O. K., Qi, S. et al Dynamic Competing

Histone H4 K5K8 Acetylation and Butyrylation Are Hallmarks of Highly Active Gene Promoters.

Mol Cell 2016;**62**:169-180

4. Yoo, E. J., Chung, J. J., Choe, S. S., Kim, K. H., and Kim, J. B. Down-regulation of histone
deacetylases stimulates adipocyte differentiation. J Biol Chem 2006; **281**:6608-6615

5. Wellen, K. E., Hatzivassiliou, G., Sachdeva, U. M., Bui, T. V., Cross, J. R., and Thompson, C. B. ATP-
citrate lyase links cellular metabolism to histone acetylation. Science 2009;**324**:1076-1080

6. King, M. T., and Reiss, P. D. Separation and measurement of short-chain coenzyme-A
compounds in rat liver by reversed-phase high-performance liquid chromatography. Anal
Biochem 1985;**146**:173-179

7. Sabari, B. R., Tang, Z., Huang, H., Yong-Gonzalez, V., Molina, H., Kong, H. E. et al Intracellular
crotonyl-CoA stimulates transcription through p300-catalyzed histone crotonylation. Mol Cell
2015;**58**:203-215

8. Mantelingu, K., Reddy, B. A., Swaminathan, V., Kishore, A. H., Siddappa, N. B., Kumar, G. V. et al
Specific inhibition of p300-HAT alters global gene expression and represses HIV replication.
Chem Biol 2007;**14**:645-657

9. Kaczmarska, Z., Ortega, E., Goudarzi, A., Huang, H., Kim, S., Márquez et al Structure of p300 in
complex with acyl-CoA variants. Nat Chem Biol 2017; **13**:21-29

10. Liu, X., Wang, L., Zhao, K., Thompson, P. R., Hwang, Y., Marmorstein, R. et al The structural basis
of protein acetylation by the p300/CBP transcriptional coactivator. Nature 2008;**451**:846-850

11. Xie, Z., Zhang, D., Chung, D., Tang, Z., Huang, H., Dai, L. et al Metabolic Regulation of Gene
Expression by Histone Lysine β-Hydroxybutyrylation. Mol Cell 2016;**62**:194-206

12. Fellows, R., Denizot, J., Stellato, C., Cuomo, A., Jain, P., Stoyanova, E. et al Microbiota derived
short chain fatty acids promote histone crotonylation in the colon through histone deacetylases.
Nat Commun 2018;**9**:105

13. Wan, J., Liu, H., and Ming, L. Lysine crotonylation is involved in hepatocellular carcinoma
progression. Biomed Pharmacother 2019;**111**:976-982

14. Jiang, G., Nguyen, D., Archin, N. M., Yukl, S. A., Méndez-Lagares, G., Tang, Y. et al HIV latency is

reversed by ACSS2-driven histone crotonylation. J Clin Invest 2018;**128**:1190-1198

15. Kebede, A. F., Nieborak, A., Shahidian, L. Z., Le Gras, S., Richter, F., Gómez, D. A. et al Histone

propionylation is a mark of active chromatin. Nat Struct Mol Biol 2017;**24**:1048-1056

16. Farrelly, L. A., Thompson, R. E., Zhao, S., Lepack, A. E., Lyu, Y., Bhanu, N. V. et al   Histone

serotonylation is a permissive modification that enhances TFIID binding to H3K4me3. Nature

2019;**567**:535-539

17. Zhang, D., Tang, Z., Huang, H., Zhou, G., Cui, C., Weng, Y. et al. Metabolic regulation of gene

expression by histone lactylation. Nature **574**,575-580 (2019)

18. Lepack, A.E., Werner, C.T., Stewart, A.F., Fulton, S.L., Zhong, P., Farrelly, L.A. et al.

Dopaminylation of histone H3 in ventral tegmental area regulates cocaine seeking. Science **368**,

197-201 (2020)

19. Mews, P., Donahue, G., Drake, A.M., Luczak, V., Abel, T., Berger, S.L. Acetyl-CoA synthetase

regulates histone acetylation and hippocampal memory. Nature 2017; 546, 381–386

20. Wellen, K.E., Hatzivassiliou, G., Sachdeva, U.M., Bui, T.V., Cross, J.R., Thompson, C.B. ATP-

citrate lyase links cellular metabolism to histone acetylation. Science2009; 324, 1076–1080

21. Mews, P., Egervari, G., Nativio, R., Sidoli, S., Donahue, G., Lombroso, S.I. Alcohol metabolism

contributes to brain histone acetylation. Nature 2019; 574, 717–721

22. Tanaka, T., Yoshida, N., Kishimoto, T., Akira, S. Defective adipocyte differentiation in mice

lacking the C/EBPbeta and/or C/EBPdelta gene. EMBO J. 1997; 16, 7432–7443.

23. Hamm, J.K., Park, B.H., Farmer, S.R. A role for C/EBPbeta in regulating peroxisome proliferator-

activated receptor gamma activity during adipogenesis in 3T3-L1 preadipocytes. J. Biol. Chem.

2001; 276, 18464–18471.

24. Gao, M. et al. Metabolically controlled histone H4K5 acylation/acetylation ratio drives BRD4

genomic distribution. Cell Rep 2021; 36(4):109460.

25. Xiao, Y. et al. HBO1 is a versatile histone acyltransferase critical for promoter histone acylations.

Nucleic Acids Res 2021; 49(14):8037-8059.

26. Carrer, A., Parris, J. L., Trefely, S., Henry, R. A., Montgomery, D. C., Torres, A. et al Impact of a

High-fat Diet on Tissue Acyl-CoA and Histone Acetylation Levels. J Biol Chem 2017;**292**:3312-

3322

27. Burgeois, C., Gorwood, J., Barrail-Tran, A., Lagathu, C., Capeau, J., Desjardins, D. et al. Specific

Biological Features of Adipose Tissue, and their impact on HIV persistence. Front Microbiol **17**

(2019)

28. Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley et al Differential gene and transcript

expression analysis of RNA-seq experiments with TopHat and Cufflinks. Nat Protoc 2012;**7**:562-

578

29. Thomas, P. D., Campbell, M. J., Kejariwal, A., Mi, H., Karlak, B., Daverman, R. et al PANTHER: a

library of protein families and subfamilies indexed by function. Genome Res 2003;**13**:2129-2141

30. de Hoon, M. J. L., Imoto, S., Nolan, J.,Miyano, S. Open source clustering software. Bioinformatics

2004;**9**:1453-1454

Figure legends:

Fig.1 Histone butyrylation increases during adipogenesis (A) Phase contrast images of 3T3L1 cells

exhibiting fibroblast like morphology in their pre-adipocyte state and accumulation of lipid droplets

780 in their adipocyte state. The lipid droplets can be stained using Oil Red-O dye. (B) RT-qPCR analysis

781 was performed to check for transcript levels of Acss2 in undifferentiated and differentiated 3T3L1

782 cells. Error bars denote mean +/- SEM of three biological replicates; two-tailed unpaired Student's t-

783 test: * $P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ns: not significant. (C) Knockdown of Acss2 in

784 differentiated 3T3L1 cells was confirmed by RT-qPCR analysis. Error bars denote mean +/- SEM of

785 three biological replicates; two-tailed unpaired Student's t-test: * $P < 0.05$, **$P < 0.01$, ***$P < 0.001$,

786 ns: not significant. (D) Histone acetylation and butyrylation pattern in 3T3L1 cells upon Acss2

787 knockdown was checked using antibodies against acetylated histone H3K18 and butyrylated H4K5.

788 (E) Histone acetylation and butyrylation levels in 3T3L1 cells in undifferentiated and differentiated

789 state were analysed by immunoblotting with antibodies specific for acetylated H3K9, butyrylated

790 H3K9, H3K23, H3K27, H4K5, H4K8, H4K12 and pan-butyryl lysine antibodies . Antibodies against core

791 histones H3 and H4 were used as loading controls along with Direct Blue staining of all four core

792 histones. Lane 1- undifferentiated pre-adipocyte (Day 0) and Lane 2 - differentiated adipocyte (Day

793 7) in each of the panels i, ii and iii. (F) Enrichment of H4K5 butyrylation in the promoters of (i) Pparg,

794 (ii) Lep and (iii) Cebpd represented in the form of fold change over IgG. Error bars denote mean +/-

795 SEM of three biological replicates; one-way ANOVA with Dunnett's multiple comparision was

796 performed: * $P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ns: not significant.

797 **Figure 2: LTK-14A selectively inhibits p300 catalysed butyrylation without affecting its**

798 **acetyltransferase activity:** Immunoblotting with antibodies against acetylated H3K18 (A) and pan-

799 butyryl lysine (C) to check for the relative inhibition of acetyltransferase and butyryltransferase

800 activities of full length p300 by LTK-14A at different concentrations. Recombinant histone H3 was

801 used as substrate. Lane 1, enzyme alone; Lane 2, enzyme + DMSO; Lane 3-6, enzyme + LTK-14A at

802 concentrations 10, 15, 20 and 25 µM respectively. Images of Western blotting are given in panels F

803 and H and their quantitations are given in panels B and D. Molecular structures of mono- (LTK-14)

and di-(LTK-14A) substituted isogarcinol (E and G). Molecular docking studies for studying

comparative interaction of LTK-14 and LTK-14A with the amino acid residues of p300 catalytic

domain (F and H). The predicted interactions are denoted by tick lines. Hydrogen bonding is shown

in red and hydrophobic interaction is shown in green. Immunoblotting with antibodies against

acetylated H3K18 (I) and pan-butyryl lysine (K) to check for the relative inhibition of

acetyltransferase and butyryltransferase activities of wild type p300 catalytic domain by LTK-14A at

different concentrations. Lane 1, enzyme alone; Lane 2, enzyme + DMSO; Lane 3-6, enzyme + LTK-

14A at concentrations 10, 15, 20 and 25 µM respectively. Images of Western blotting are given in

panels I and K and their quantitations are given in panels J and L. Immunoblotting with antibodies

against acetylated H3K18 (M) and pan-butyryl lysine (O) to check for the relative inhibition of

acetyltransferase and butyryltransferase activities of catalytically active p300 catalytic domain with

point mutation C1438A by LTK-14A at different concentrations. Lane 1, enzyme alone; Lane 2,

enzyme + DMSO; Lane 3-6, enzyme + LTK-14A at concentrations 10, 15, 20 and 25 µM respectively.

Images of Western blotting are given in panels M and O and their quantitation are given in panels N

and P. Error bars denote mean +/- SEM of three biological replicates, one-way ANOVA with

Bonferroni's multiple comparison; * $P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ns: not significant.

Fig. 3: The butyrylation inhibitor represses adipogenesis in 3T3L1 cells: (A) The permeability and

stability of LTK-14A in 3T3L1 was tested by LC-ESI-MS analysis of metabolites, extracted from the

compound treated 3T3L1 cells to detect the protonated LTK-14A in the ESI spectra of the eluted

fraction. (B) MTT assay was performed to check for any possible cellular toxicity effect of LTK-14A,

using 25 µM concentration of the compound on 3T3L1 cells upon incubation for varying lengths of

time (2, 4 and 6 days). Comparison was done with cells that were left untreated or treated with

DMSO as solvent control. The effect of LTK-14A (25 µM) treatment on adipogenesis was tested by

Oil Red-O staining of the adipocytes (C) and the quantitation of staining is represented graphically in

panel D. Error bars denote mean +/- SEM of three biological replicates,one-way ANOVA with

Bonferroni's multiple comparison; * P < 0.05, **P < 0.01, ***P < 0.001, ns: not significant. (E) The

effect of LTK-14A on the global transcription network inside 3T3L1 cells was investigated by high

throughput transcriptome analysis with RNA isolated from compound treated, DMSO treated and

untreated 3T3L1 cells (n=2 in each case). Differential gene expression pattern of three conditions

compared with each other, depicted in volcano plots- No Inhibitor versus DMSO , No Inhibitor

versus LTK-14A and DMSO versus LTK-14A. The downregulated and upregulated genes are shown in

blue and red respectively.(F) Heat maps showing the differential gene expression pattern between

LTK-14A treated cells and DMSO treated or untreated cells.

Figure 4: LTK-14A represses expression of several pro-adipogenic genes and inhibits lipid

metabolism: (A) Venn diagram showing the number of differentially expressed genes upon LTK-14A

treatment with respect to untreated and DMSO treated conditions. (B) Classification of the common

differentially expressed genes according to their type. Gene ontology analysis was performed to

study the different biological pathways that were affected by LTK-14A in 3T3L1 compared to DMSO

treated as well as untreated conditions, as predicted by Panther software. The pathways were

ranked in the order of those with maximum number of genes affected (E). A different ranking order

was created on the basis of fold enrichment i.e. the pathways in which the ratio of number of

observed affected genes over number of expected genes to be affected was the highest (C). A

similar analysis was also performed for genes differentially expressed in DMSO versus untreated

conditions (D). (F) A molecular network was constructed to depict the interconnectivity between the

lipid metabolism related genes and pathways that were affected upon their dysregulation by LTK-

14A compared to DMSO treated conditions. (G) RT-qPCR analysis was performed to validate the

putative target genes of LTK-14A. Error bars denote mean +/- SEM of three biological replicates;

one-way ANOVA with Bonferroni's multiple comparison: * P < 0.05, **P < 0.01, ***P < 0.001, ns:

not significant.

Fig.5: LTK-14A inhibits histone butyrylation in 3T3L1 cells, without affecting acetylation or other

post-translational modifications: Histone butyrylation levels in 3T3L1 cells that were treated with

LTK-14A were compared with that in DMSO treated and untreated conditions by immunoblotting

with antibodies against butyrylated histone H3K23 , H4K5 and pan-butyryl lysine. Histones H3 and

H4 were used as loading controls (A). The fold change of quantified band intensity is graphically

represented in panel (B) in which the error bars denote mean +/- SEM of three biological replicates.

Under similar conditions histone acetylation levels were compared by immunoblotting using

antibodies against acetylated histones H3K18, H3K9 and H4K12 (C). The result is graphically

represented in panel (D) in which error bars denote mean +/- SEM of three biological replicates.

Immunoblotting was also performed with antibodies against histone H3R17 dimethylation, H3S10

Phosphorylation and H3K9 dimethylation (E). The result is graphically represented in panel F in

which error bars denote mean +/- SEM of three biological replicates. For all data one-way ANOVA

with Bonferroni's multiple comparision was performed: * P < 0.05, **P < 0.01, ***P < 0.001, ns: not

significant .

Fig. 6: The weight gaining of high fat diet fed mice (C57BL6/J) could be significantly reduced by the

butyrylation specific inhibitor of p300: Trend of weight gaining (A) and average body weight gained

after 16 weeks (B) of C57BL6/J mice maintained on normal chow diet, high fat diet and high fat diet

mixed with LTK-14A was plotted; n=4 in each group. Representative images of dissected bodies

showing fat accumulation (C) and epididymal fat pads (D) of the mice from three groups. The

average weight of the fat pads is represented graphically (E); n=4 in each group. Representative

images of morphology of adipose tissue (F) and liver (G) from the mice of the three groups, as

studied by hematoxylin and eosin staining. For all data one-way ANOVA with Dunnett's multiple

comparision was performed: * P < 0.05, **P < 0.01, ***P < 0.001, ns: not significant. (H)

Immunofluorescence microscopy of liver sections stained with antibodies against histone H4K12

acetylation (panel i), histone H3K14 acetylation (panel ii) and histone H4K5 butyrylation (panel iii).

The fluorescence intensity of the modified histone marks was normalised against DAPI staining of

whole nuclei for quantitation which is graphically depicted in (I) panel i, ii and iii ;n=4 in each group;

one-way ANOVA with Dunnett's multiple comparision: * P < 0.05, **P < 0.01, ***P < 0.001, ns: not

significant. (J) Histone H3K14, H3K18, H4K12 acetylation and H4K5 butyrylation levels in epididymal

fat pads of mice were estimated by immunoblotting . Histone H3 was used as loading control The

fold change of quantified band intensity is graphically represented in panel K,L, M and N ;n=3 in each

group; one-way ANOVA with Dunnett's multiple comparision: * P < 0.05, **P < 0.01, ***P < 0.001,

ns: not significant.

Fig. 7: Oral gavaging of the butyrylation inhibitor reduces weight of genetically obese db/db mice:

Trend of weight gaining (A) and average body weight gained after 30 days (B) of db +/- and db/db

mice orally administered with either LTK-14A or vehicle solution ; n=6 in each group. (C)

Representative images of dissected bodies showing fat accumulation (panel iii) of the mice from

three groups. Representative images of liver (D) from the mice of three groups and average weight

of liver (E) depicted graphically; n= 6 in each group. Representative images of epididymal fat pads (F)

from the mice of three groups and average weight of fat pads (G) depicted graphically; n= 6 in each

group. Total fat mass (H) and total lean mass (I) of mice from three groups are depicted graphically;

n= 6 in each group. Representative images of morphology of liver (J) and adipose tissue (K) from the

mice of the three groups, as studied by hematoxylin and eosin staining. (L) Immunofluorescence

microscopy of liver sections stained with antibodies against histone H4K12 acetylation (panel i),

histone H3K14 acetylation (panel ii) and histone H4K5 butyrylation (panel iii). The fluorescence

intensity of the modified histone marks was normalised against DAPI staining of whole nuclei for

quantitation which is graphically depicted in (M) panel i, ii and iii ;n=4 in each group; one-way

ANOVA with Dunnett's multiple comparision: * $P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ns: not

significant. (N) Histone H3K14, H3K18, H4K12 acetylation and H4K5 butyrylation levels in epididymal

fat pads of mice were estimated by immunoblotting. Histone H3 was used as loading control The

fold change of quantified band intensity is graphically represented in panel O,P, Q and R; n=3 in

each group; one-way ANOVA with Dunnett's multiple comparision: * $P < 0.05$, **$P < 0.01$, ***$P <$

0.001, ns: not significant. (S) qRT-PCR with RNA isolated from epididymal fat pads to check the

expression pattern of lipogenesis related pro-adipogenic genes upon LTK-14A treatment;

n=3/4 in each group; one-way ANOVA with Dunnett's multiple comparision: * $P < 0.05$, **P

< 0.01, ***$P < 0.001$, ns: not significant

Graphical abstract:

Proposed model for the role of p300-mediated histone butyrylation in adipogenesis:

In pre-adipocytes, there exists a basal level of histone acetylation while butyrylation is present to a

much lesser extent owing to low stoichiometric levels of butyryl CoA. Induction of adipogenesis

causes a simultaneous upregulation of histone acetylation and butyrylation marks leading to

increased rate of adipogenesis and concomitant transcriptional activation of pro-adipogeneic genes.

Onset of obesity in mice, either due to excess energy intake through high fat diet consumption or

increased de novo synthesis of fatty acids due to leptin receptor gene mutation leading to

hyperphagic behavior, is accompanied by adipocyte hyperplasia and hypertrophy. Both the organs

of adipose tissue and liver were found to have enhanced levels of H4K5 butyrylation during obesity.

LTK-14A, a butyrylation specific inhibitor could efficiently prevent the processes of adipogenesis and

adipocyte hypertrophy due to inhibition of H4K5 butyrylation in these organs. Thus the compound

could attenuate weight gain by selective inhibition of butyrylation without affecting acetylation,

thereby highlighting the importance of histone butyrylation in adipogenesis and obesity.

Figure 1

Figure 2

Figure 3

A

B

C

3T3L1
Pre-adipocytes

8 Days
No inhibitor/DMSO/LTK-14A

Adipocytes

No inhibitor   DMSO   LTK14A

D

F

LTK-14A
downregulated
genes

LTK-14A
upregulated
genes

E

No inhibitor vs DMSO   No inhibitor vs LTK-14A   DMSO vs LTK-14A

A

LTK-14A vs No Inhibitor
4879

Common Genes
(affected by LTK-14A)
3509

1370

4334
LTK-14A vs DMSO

Upregulated (1309)
Downregulated (2200)

B

91%

- Protein coding
- lncRNA
- Processed pseudogene
- Transcribed unprocessed pseudogene
- transcribed_unitary_pseudogene
- Unclassified

C

Biological processes	Fold enrichment	P-value	FDR
fatty acid beta-oxidation using acyl-CoA dehydrogenase	6.93	0.0000556	0.00304
positive regulation of cholesterol biosynthetic process	6.06	0.00129	0.0442
positive regulation of cholesterol metabolic process	6.06	0.00129	0.0441
positive regulation of sterol biosynthetic process	6.06	0.00129	0.044
acetyl-CoA biosynthetic process from pyruvate	6.06	0.00129	0.0439

D

Biological process	Fold enrichment	P-value	FDR
receptor localization to synapse	4.78	0.0000625	0.0448
protein localization to synapse	4.37	0.0000078	0.00724
regulation of synaptic plasticity	3.55	1.25E-09	0.00000393
signal release	2.97	0.0000189	0.0166
exocytosis	2.83	0.00000493	0.00519

E / **F** / **G**

No Inhibtor
DMSO
LTK-14A

Figure 5

A

B

C

D

E

F

H3R17me2
H3S10Ph
H3K9me2
H3

Figure 6

Figure 7

Graphical abstract