1 **An abundant myeloid progenitor shapes neonatal hematopoiesis of naked mole-rats**

2 Stephan Emmrich[1], Andrei Seluanov[1], Vera Gorbunova[1,2]

3   [1] Department of Biology, University of Rochester, Rochester NY 14627, USA
4   [2] Corresponding authors
5
6
7 ***Correspondence:**
8
Vera Gorbunova
University of Rochester
434 Hutchison Hall, River Campus
Rochester NY 14627-0211
Phone: 585-275-7740
Email: vera.gorbunova@rochester.edu

Abstract

The naked mole-rat became a significant animal model for its exceptional longevity and

resistance mechanisms to cancers, however the developmental properties of their native stem

and progenitor compartments are poorly understood. Here we report a high frequency bone

marrow progenitor exclusively present in neonatal animals. The heterotypic developmental stage

was marked by an immunophenotype resembling adult naked mole-rat CMP-like progenitors with

mature neutrophilic Thy1.1-antigen levels and showed a clear differentiation bias and

transcriptome signature towards myelopoiesis. Importantly, myeloid progenitors overexpressed

both CXCL12 and its alternative receptor CXCR7, thereby abrogating their homing capacity and

produced multiple secreted matrix components related to neonatal bone marrow niche formation.

These results indicate a transient ontogenic cell stage during postnatal hematopoiesis unique to

naked mole-rats and further advances our understanding of the developmental steps involved in

creating the regenerative cell pool and its environment in a long-lived species.

Introduction

The naked mole-rat (*Heterocephalus glaber*) has become an attractive animal model in aging research due to its longevity and resistance to disease. Naked mole-rats are subterranean eusocial rodents found in grasslands of East Africa. This remarkable rodent with the size of a mouse has a maximum lifespan of over 39 years in captivity and does not display increased mortality during aging[1,2]. Previously we discovered that high molecular weight hyaluronic acid is responsible for the resistance of naked mole-rats to solid tumors[3], while further unique molecular adaptations such as a split 28S rRNA and a fusion isoform in the INK4A/ARF locus confer improved cellular responses towards senescence and transformation[4,5]. We created a naked mole-rat hematlas by employing cross-reactive antibodies for fluorescence-activated cell sorting (FACS), purifying hematopoietic stem and progenitor cell (HSPC) subsets and using colony formation, xenotransplantation and Cellular Indexing of Transcriptomes and Epitopes by Sequencing (CITE-Seq) to link single-cell transcriptomic cell type annotation with FACS immunophenotypes[6]. Hallmarks of naked mole-rat hematopoiesis are high basal levels of neutrophil granulocytes (GC) accompanied by reduced B-lymphopoiesis, splenic erythropoiesis as major route of red blood cell production, prolonged cell cycle duration of hematopoietic cells and a primitive HSPC compartment marked by CD34, THY1 and TM4SF1. Moreover, we revealed the presence of an ectopic cervical thymus along with a delay of immunosenescence and thymic involution in naked mole-rats, and showed that naked mole-rat T-lymphopoiesis results in strongly reduced CD8+ cytotoxic T cell frequencies[7].

Every second, an adult human produces nearly 2 million red blood cells[8]. This enormous turnover rate, required to sustain life in all vertebrates, is accomplished unilaterally by hematopoietic stem cells (HSCs). As a paradigm of stem cell biology, HSCs are self-renewing cells capable of differentiating into the wide variety of mature blood lineages (LIN) in our bodies, ranging from red blood cells to the entire spectrum of immune cells[9]. Vertebrate hematopoiesis

starts within the yolk sac to succinctly continue over aorta-gonads-mesonephros, placenta and

fetal liver (FL), whereby each ontogenic HSC stage is marked by distinct immunophenotypes and

transcriptional signatures[10]. FL HSCs have higher regenerative potential than young adult marrow

HSCs[11], and predominantly produce myeloid cell outputs in contrast to balanced adult HSCs[12].

Moreover the FL phenotype is retained up to the 3rd week of life in mice[13]. The SLAM family

members CD48, CD150 and CD244 are differentially expressed among primitive progenitors in

adult mouse bone marrow (BM) and currently allow the highest purification grade of LIN$^-$/Sca-

1$^+$/Kit$^+$/CD48$^-$/CD150$^+$/CD41$^-$ long-term LT-HSCs[14]. Similarly, highest purity of long-term

multilineage reconstituting fetal liver HSCs was obtained from the LIN$^-$/Sca-1$^+$/Mac-1$^+$/CD48$^-$

/CD150$^+$ immunophenotype[15]. The most prevalent HSPC population within FL LIN$^-$/Sca-1$^+$/Kit$^+$

(LSK) during E15.5-E18.5 were Flt3$^-$/CD48$^+$/CD150$^+$ cells comprising megakaryocyte-biased

MPP2, a multipotent progenitor (MPP) subset[16]. FL-MPP2 frequencies declined to adult levels

around post-natal day 0 (P0), until which these MPP2 showed a strong decline in colony formation

capacity and in vivo repopulation, a paradigm of a functionally distinct but immunophenotypically

identical pre-natal HSPC population as compared to its adult counterpart[17].

Here we characterized the post-natal HSPC landscape in naked mole-rats, and found a

myeloid-biased progenitor stage specific to newborn BM and to a lesser extent the spleen. The

progenitors concomitantly overexpressed CXCL12 and CXCR7, leading to impaired BM niche

homing. Our results provide an unprecedented finding of stage-specific neonatal myeloid

progenitors without homing capacity, improving our current understanding of stem cell biology in

the naked mole-rat as a model of exceptional longevity.

Results

An abundant progenitor population specific to naked mole-rat neonatal hematopoiesis

Previously we functionally characterized purified naked mole-rat HSPC subsets by cross-

reactive monoclonal antibodies[6]. Using cross-reactive clones against the human stem cell

markers CD34 and THY1 created a complex staining pattern characteristic for adult naked mole-

rat bone marrow (BM, Figure 1a). The transcriptional identity of neutrophil granulocytes (GC),

erythroid (ERY) and T cells (TC) as well as megakaryocytic erythroid progenitors (MEP) and

HSPCs gated from the flow cytometry (FACS) staining was confirmed by CITE-Seq[18]. We then

harvested hematopoietic tissues from P2-3 newborns to be profiled by our naked mole-rat HSPC

FACS panel. Neonatal BM featured a distinctive Thy1.1/CD34 staining pattern, with less mature

GCs than adult BM (Figure 1b). Notably, an abundant neonate Thy1.1^{hi}/CD34^{hi} cell population

extending from the Thy1.1^{int}/CD34^{hi} HSPC fraction was absent in adult animals. A 4.8-fold

increase of Thy1.1/CD34 double-positive cells in neonate compared to adult livers indicates fetal

hematopoiesis as conserved origin of later developmental steps (Supplementary Figure 1).

Reasoning from the Thy1.1^{hi} levels seen exclusively in neutrophils[6], we referred to

Thy1.1^{hi}/CD34^{hi} cells as neonate myeloid progenitors (NMP).

We previously showed that CD11b and CD18 label naked mole-rat myeloid peripheral

blood cells, whereas high CD90-antigen marks neutrophils and dim CD90-antigen T cells[6]. This

labelling pattern was confirmed in adult marrow (Supplementary Figure S1). Together with

CD125, which labels B cells[7], we combined a lineage depletion cocktail (LIN) used to further purify

the HSPC fraction (Figure 1c). The gating limit for LIN$^-$ was positioned to exclude 2 orders of

magnitude of positively LIN-stained cells and in parallel to exclude any high complexity side

scatter cells. Compared to adult BM neonate marrow featured a higher LIN$^-$ fraction, which is

expected given the higher amount of erythropoiesis and stem and progenitors required for

postnatal development[19]. Specific to neonates however was a strongly expanded

LIN$^+$/Thy1.1$^{int/hi}$/CD34^{hi} population containing cells in some animals with the highest CD34 and

Thy1.1 staining intensities observed in any tissue (Figure 1d). Although this immunophenotype

resembled adult LIN$^+$/Thy1.1^{int}/CD34^{hi} CP1 (candidate population), neonate

LIN$^+$/Thy1.1$^{int/hi}$/CD34^{hi} cells expressed CD90-antigen at high levels and did not express CD11b

(Supplementary Figure 1). Hence we coined this fraction TC90 (Thy1.1$^{int/hi}$/CD34^{hi}/CD90^{lo}), which

was predominantly found in neonatal marrow as a 4.6-fold increase compared to spleen (Figure

1e). While there was ~5-fold increase of CP1 (0.66±0.22%) over LIN$^-$/Thy1.1^{int}/CD34^{hi} LTCs

(0.13±0.07) in adult BM[6], the NMP population specific to neonate whole BM was exclusively

detected in the LIN$^+$ TC90 fraction (8.1±3.8%), which comprised 38-fold more cells than neonate

LIN$^-$ CP2 (0.21±0.21%; Figure 1f).

Next we immunophenotyped murine adult and P3 HSPC fractions, which showed an

increased LIN$^-$/Sca-1$^+$/Kit$^+$ (LSK) stem and progenitor compartment in neonates across

hematopoietic tissues (Figure 2a). Expansion of LSKs relative to adult tissues was highest in

neonatal liver, resembling fetal liver hematopoiesis progressing towards the neonatal-to-adult

stage transition (Figure 2b). Cd150$^+$ LSK long-term hematopoietic stem cells (LT-HSCs) are

slightly elevated in 6 month-old adult marrow, however multipotent progenitors (MPP) are 4.2-fold

more abundant in neonatal mice (Supplementary Figure 1). Myeloid progenitor populations within

the LIN$^-$/Sca-1$^-$/Kit$^+$ (LK) compartment were unaltered. Naked mole-rat LTCs, which equivalently

to murine LSKs have lympho-myelo-erythroid potential and harbor the primitive HSC pool[6], remain

at neonatal levels in adult animals (Figure 2c-d). This is a further neotenic trait, the retention of

juvenile features in adult animals, of the naked mole-rat[20]. Consistent with a prevalence of

erythropoiesis around birth neonate CP5, comprised of hyperproliferative late erythroblasts[6],

expanded 3.5-fold in BM and 4.7-fold in spleen (Supplementary Figure 2).

Conclusively, we found a large expansion of CD90-antigen positive cells co-expressing

naked mole-rat stem cell markers Thy1.1-antigen and CD34 specifically in neonates, whereas

primitive stem and progenitor LTCs remained at neonatal levels in adult hematopoietic tissues.

TC90 cells feature unique transcriptome and splicing signatures

To address molecular changes between neonate and adult HSPCs we integrated RNA-

Seq profiles of TC90s in our previously reported dataset from adult CP1/LTC/CP3/CP4[6]. This

combined to 9863 expressed orthologues across 24 normalized samples (Supplementary Figure

2). Using t-distributed stochastic neighborhood embedding (t-SNE)[21] neonate TC90s clearly

formed a separate cluster, however adult CP1 and LTCs could not be resolved into their separate

immunophenotypes but instead clustered together as CP1/LTC partition, same for CP3/4 (Figure

3a). Differential gene expression analysis found most upregulated genes for TC90s (Figure 3b).

Differentially expressed genes (DEGs) were fed into single sample gene set enrichment analysis

(ssGSEA) using a curated collection of expression signatures from human, mouse and naked

mole-rat HSPCs for cell type annotation[6]. The adult partitions CP1/LTC and CP3/4 showed

consensus annotation patterns towards their combined FACS population identity: Human HSC,

human CD34[+] and naked mole-rat LTC and lymphomyeloid signatures enriched while naked

mole-rat common erythroid signatures depleted for CP1/2; human MEP, naked mole-rat CP4 and

common erythroid signatures enriched but lymphoid and myeloid signatures depleted in CP3/4

(Figure 3c). TC90 cells shared a human myeloid gene signature with CP1/2 and were enriched

for human and mouse HSC gene expression, resembling a myeloid-biased early progenitor.

Next we checked whether this profile was also present as differential transcript usage

(DTU). Using the same dataset as in Figure 3a we obtained 15038 unique transcripts assigned

through the FRAMA BM transcriptome[6,22], leading to virtually the same clustering (Supplementary

Figure 2). Likewise, TC90 cells had highest DTU, thus we checked for inversely expressed

transcripts (IETs; see Methods) from the same gene locus in TC90s (Supplementary Figure 2).

Out of 12 total IETs we found an isoform i6 of POLR2B strongly upregulated exclusively in TC90

(1260-fold p<10^{-11}; Figure 3d). BLAST alignments of the mRNA coding regions (ORF) revealed

an ubiquitously expressed POLR2B isoform i7 downregulated in TC90 (3.6-fold p<10^{-6}) giving rise

to the full-length RNA-Polymerase-2β subunit protein. The truncated i6 lacks protrusion and lobe

domains (Figure 6e), which contribute to maintenance of the transcription bubble but are known

as dispensible region in bacterial homologues[23]. Another IET pair was found at the OGG1 locus,

a DNA repair enzyme excising 8-oxoguanine which can be formed upon reactive oxygen species

(ROS) exposure[24]. TC90s overexpressed the i25 full-length isoform (5.5-fold p<10^{-7}), while

expression of a truncated i29 lacking both the mitochondrial targeting signal and an N-terminal

domain contributing to substrate recognition was completely abolished (Figure 3f-g). The IET

isoform with the strongest depletion for TC90s was i12 of the TANK gene (890-fold, p<10^{-4}; Figure

3h), an adaptor binding to TRAF1-3 and thereby modulating TNFR:NF-κB signaling[25].

Interestingly, the two isoforms solely differed in their 5'UTR, with only the i21 isoform harboring a

9bp TOP motif associated with mTORC1-mediated translational induction[26] (Figure 3i).

Overall neonate TC90s differ from adult naked mole-rat LTCs and progenitors by very

high abundances in spleen and BM, high CD90-antigen levels and specific alternative splicing

patterns.

The SDF-1α/CXCR4-axis is altered in TC90 cells

The ontogeny-associated differences between murine fetal and adult hematopoietic

tissues are conserved in humans[27]. To segregate the inherent properties of TC90s as an earlier

progenitor pool in naked mole-rats, we compared them to LTCs along with human cord blood

(CB) and marrow HSCs (Figure 4a, Supplementary Figure 3). Using the MSigDB hallmark

geneset collection[28] we classified the stem cell types according to their signatures, which yielded

Epithelial-Mesenchymal-Transition (EMT) as top ssGSEA category for TC90 cells (Figure 4b).

Importantly, the top leading edge gene for EMT in TC90s was CXCL12 (Figure 4c). To our

surprise we found the cognate receptor CXCR4 as one of the top downregulated genes in TC90s.

Apart from its primary target CXCL12 can also bind to the alternate chemokine receptor homolog

ACKR3[29], leading to ligand internalization and β-arrestin recruitment. Strikingly, TC90 cells

specifically overexpressed all three ACKR interceptors (Figure 4c). The arrestin family of adaptor

proteins can induce MAPKs, AKT but also NF-κB signaling[30], the latter could be further modulated

by i21 TANK activation (Figure 3h-i). Consistent with a fetal myeloid predominance TC90s

overexpressed CD14 and CD38, along with the stem cell master transcription factor EBF1 (Figure

4c). On the contrary there are key marker genes of the naked mole-rat stem cell state emerging

from population and single cell RNA-Seq such as ICAM2, HOXA9, GATA2, and HHEX, MEP

markers MDK, NFE2 and conserved HSC regulators TAL1, BMI1, MEIS1 and HMGA2, appear

strongly decreased in TC90s (Figure 4c, Supplementary Table X). An intriguing finding are the

coordinated upregulation of antiproliferative p21, BTG1 and 4 IGF-binding proteins together with

corresponding reduction of 4 centromer proteins, CEP85, POLR1A and MYB. Despite elevated

frequencies of TC90s in neonatal BM, spleen and liver but in line with a depleted G2M-Core Set

cell cycle signature (Figure 3c), this suggests the neonate-specific LIN+/Thy1.1hi/CD34hi extension

of the CP1-like LIN+/Thy1.1int/CD34hi subset within TC90 cells as a granulocytic precursor with

little self-renewal and proliferative activity.

Whereas an absence of HSC maintenance genes would be expected on a committed

progenitor such as TC90, we aimed to confirm the inversed CXCR4/ACKR expression pattern by

extending the comparison to human HSCs and committed lymphomyeloid progenitors. Thus we

integrated expression profiles of common lymphoid progenitors (CLP), multipotent lymphoid

progenitors (MLP), granulocytic monocytic progenitors (GMP) and HSCs from 6 different RNA-

Seq datasets [31-35] across cord blood and adult BM stages with LTC, CP1 and TC90 transcriptomes

(Figure 5a). This compound dataset comprised 51 samples and 10291 common orthologs

(Supplementary Figure 3). Unsupervised hierarchical clustering faithfully recapitulated each of the

two ontogenic stages and species groupings (Figure 5b). Remarkably, CXCL12 and ACKR3, also

known as CXCR7, were specifically overexpressed in TC90s as compared to all other cell types

(Figure 3c). Concordantly, CXCR4 levels were lowest in naked mole-rat TC90 cells. While ACKR2

was also overexpressed in TC90, the alternate CXCR7 ligand CXCL11 was absent in naked mole-

rat HSPCs (Supplementary Figure 3). Interestingly, HMGA2 which maintains the fetal stem cell

status in concert with LIN28B[36], was higher expressed throughout human neonatal stem and

progenitor populations, whereas naked mole-rat TC90 showed less HMGA2 transcription than

adult LTCs and CP1 (Supplementary Figure 3). The same pattern was seen for MLLT3, the fusion

partner AF9 of t(9;11) MLL-rearranged translocations, which has a role in early mouse

hematopoiesis[37,38], whereas KMT2A was generally lower expressed in naked mole-rats and

MLLT1 showed no significant changes across species, cell types or ontogenies (Supplementary

Figure 3). The ssGSEA profile of TC90 cells compared to neonate and adult HSPCs reproduced

the EMT signature as top MSigDb hallmark term ($q<10^{-29}$; Supplementary Table Y). Using the

fGSEA tool to retrieve the leading edge genes also scored EMT as top term ($q=0.00055$;

Supplementary Table Z), which were enriched with multiple secreted proteins responsible for

organization of or interacting with the extracellular matrix (ECM), such as Collagen Type I Alpha

2 Chain, PCOLCE, Fibronectin 1, Fibulin 1/2/5, Elastin, Transgelin, Thrombospondin 1, Syndecan

1/4, Periostin, Nidogen 2, Versican, Decorin, ECM1 and MGP (Figure 5d). In conjunction with

elevated skeletal muscle-related TPM1, GEM, SGCB transcripts TC90s overexpressed several

factors implicated in cell migration and motility. Strikingly, THY1 mRNA expression across species

and ontogenic stage correlated with the TC90 Thy1.1[hi] subset immunophenotype (Figure 1d, 5d).

Here we report the first developmental stage-specific gene expression pattern for naked

mole-rat neonatal committed progenitor TC90 cells, marked by a conserved EMT signature,

CXCL12 signaling through ACKRs and absence of multiple HSC maintenance genes. Moreover,

high expression of a broad ECM component spectrum suggests a role in remodeling of the bone

marrow microenvironment.

Neonate TC90 are myeloid-biased CMP-like progenitors without homing capacity

The unique transcriptome signature of neonatal TC90 was further interrogated using

colony formation and xenotransplantations (TX) of sorted naked mole-rat HSPCs. Multipotency

as the capacity to form multiple distinct lineages via differentiation can be assayed through

quantitation of progenitor frequencies from colony assays in methylcellulose[39]. We previously

validated the scoring of colony-forming unit (CFU) types by cytochemistry of single colonies[6].

Here we show representative morphologies for each colony type derived from naked mole-rat

adult marrow LTCs (Supplementary Figure 2). CFU-M (Monocytic/Macrophage) colonies

displayed colorless, large cells as scattered aggregations, CFU-GMs (Granulocytic-Monocytic)

had high amounts of scattered single neutrophils, whereas BFU-Es (Burst-forming unit Erythroid)

were larger clusters of condensed small cells with brownish coloration indicating heme production,

and CFU-Es (Colony-forming unit Erythroid) had smaller clusters of condensed very small cells

with brown-to-red coloration. Upon comparison between neonate and adult sorted HSPC fractions

we found that all neonate populations produced significantly more CFU-GM (Granulocytic-

Monocytic) colonies (Figure 6a). Neonatal CFU-GM colonies were generally of larger diameters

than any other colony type (Supplementary Figure 4). Given the myeloid bias of ontogenetically

younger fetal liver cells in mice this was expected[12]. Moreover, we saw neonate CP3 MEPs

yielding less CFU-E colonies (2.8-fold, p=0.0017), which derive from more differentiated erythroid

precursors than BFU-E clones, indicating a higher frequency of early progenitors within the

erythroid lineage pool of neonate MEPs.

When adult LTCs or CP1 were transplanted via intravenous injection (i.v.) into

immunodeficient mice (NSGS), both produced similar peak engraftment rates at 12-16% host BM

chimerism between 2 and 4 weeks post TX[6]. Thus we compared CP1 with neonate TC90

xenografts administered either i.v or through intrafemoral injection (i.f.) to account for potential

engraftment failures due to impaired BM homing (Figure 6b). At 4 weeks post TX through i.v.

there was clearly no engraftment of TC90s, whereas adult CP1 reached 12.6±4% BM chimerism

(Figure 6c). Next we quantified i.f. xenografts by separating each femur and tibia from recipients

to distinguish the direct marrow injection site of the left femur from the left tibia and contralateral

bones (Figure 6d-e). Adult CP1 produced robust multilineage engraftment of 20±2.4% at the

injected femur. Reconstitution of non-injected bones requires HSPC homing through the

circulation, resulting in diminished engraftment distal the i.f. injection site[40]. Conversely, CP1

xenografts displayed decreasing chimerism in contralateral right femori with 14.8±3.3%, right

tibiae with 10.9±2.5% and left tibiae with 5.3±2.1%. Strikingly however, TC90 chimerism yielded

4.4±1.1% at the i.f. site, whereas non-injected bones had no engraftment (Figure 6d).

Conclusively, HSPC homing in neonate TC90 is abrogated through downregulated CXCR4

expression accompanied by high ACKR levels acting as decoys for SDF-1α signaling. Previously

we applied the Thy1.1/CD34 staining pattern to differentiate between the major xenograft

lineages[6]. The TC90 grafts of left femori have very little CD34$^+$ engraftment compared to CP1

(Figure 6e). Instead Thy1.1^{hi}/CD34$^-$ GCs form the bulk of TC90 progeny, which is in line with their

high frequency of CFU-GM colony formation and naked mole-rat neutrophilic Thy1.1-antigen and

CD90-antigen levels at the cell surface.

Thus we showed that TC90 cells have common myeloid progenitor (CMP) potential

with a strong bias towards neutrophil cell fate and no capacity of HSPC homing.

Discussion

268   Naked mole-rats are the longest-lived rodents that remain healthy until the end of their

269 lives and are resistant to age-related diseases including cancer. Fetal and neonatal stem cells

270 are programmed in a developmental stage-specific manner to establish hematopoiesis and

271 become the progeny of the adult HSC pool, which is crucial for lifelong blood and immune cell

272 production. Thus characterization of neonatal HSPCs will improve understanding of the biology

273 that enables the exceptional lifespan of naked mole-rats.

274   Here we show that naked mole-rat neonatal BM is enriched with a specific progenitor

275 stage we termed TC90. Mice showed a substantial expansion of neonatal MPPs responsible for

276 higher HSPC frequencies throughout hematopoietic tissues, while their more committed

277 progenitors remained unchanged towards the adult stage. By contrast, the primitive HSPC

278 compartment marked by LTCs in naked mole-rats remained unchanged, while expanded TC90s

279 are LIN$^+$ and their adult counterpart CP1 is a myeloid-biased CMP-like progenitor with durable

280 self-renewal as show by CITE-Seq and FACS[6]. Moreover, the majority of TC90 cells are CD11b$^-$

281 /CD90^{hi}/Thy1.1^{hi}/CD34^{hi} extending from the CP1 LIN$^+$/Thy1.1^{int}/CD34^{hi} immunophenotype,

282 thereby forming an intermediate differentiation stage linked to CD11b$^+$/CD90^{hi}/Thy1.1^{hi}/CD34$^-$

283 mature GCs. Whole BM of naked mole-rats comprises ~74% neutrophil GCs and their precursors,

284 whereas mice have ~52% and humans ~61% GCs[6,41]. Hence the accumulation of a neonatal

285 myeloid progenitor (Figure 1a), dedicated to increase innate immune effector GCs appears to be

286 an adaptation to protect the newborn and sustain ample blood and marrow GC frequencies in

287 adult animals. It is important to note that the TC90 sorting and quantitation gate subsumes a

288 heterogenous population of neonatal CP1 and committed granulocytic precursors, based on the

289 CD11b$^-$/Thy1.1^{hi}/CD90^{hi} NMP immunophenotype. This explains the shift towards granulocytic

290 differentiation and accounts for the decreased erythroid commitment seen for TC90 in colony

291 assays and i.f. xenografts.

TC90 RNA-Seq profiles revealed downregulation of CXCR4 and concurrent

overexpression of several ACKR orthologs and CXCL12, suggesting alternate SDF-1α/CXCR4-

axis signaling. Indeed TC90s produced xenografts upon direct marrow transfer at the injection

site, yet host marrow distal from the i.f. site was not repopulated, and i.v. delivered transplants

failed to engraft. This points towards a BM homing deficiency in TC90 due to competitive inhibition

of CXCR4 by CXCR7[42]. CXCR7 can heterodimerize with CXCR4 and interfere with CXCR4-

mediated integrin activation but not chemotaxis[43,44], CXCL11 triggers CXCR7-induced cell

adhesion in myeloid leukemia cells[45]. In T cells CXCR7/CXCL12 interaction promotes survival

and chemotaxis in a G-protein independent manner[46], instead signaling occurs through β-

arrestins[47]. Conceivably, their exceptionally high transcript levels render TC90 cells as a potential

key source of CXCL12 in naked mole-rat neonatal BM. Furthermore, TC90 overexpress multiple

secreted factors for ECM organization, pointing towards a role in modeling of the bone marrow

niche. If responsible for generating the chemotactic SDF-1α gradient, TC90 cells employing

CXCR7 to abolish CXCR4-mediated migratory signaling could remain in and contribute to the

developing niche. This would explain both the TC90 insensitivity to homing signals in a

transplantation setting, as well as TC90 frequencies of 8.1±3.8% in neonatal whole BM (Figure

1f), enabling sustained production of ECM components.

In conclusion, we found an abundant neonatal myeloid progenitor stage that supplies the

newborn with an essential reservoir of neutrophils to support a myeloid-based immune system

and potentially acts as source of the SDF-1α homing chemokine and multiple ECM components.

This reveals novel aspects of myelopoiesis in the developing blood system of naked mole-rats

and adds transient developmental stages promoting GC accumulation and producing CXCL12

and CXCR7 as further unique traits of these long-lived rodents.

Materials and Methods

Quantification and Statistical Analysis

Data are presented as the mean ± SD. Statistical tests performed can be found in the figure legends. P values of less than 0.05 were considered statistically significant. Statistical analyses were carried out using Prism 9 software (GraphPad) unless otherwise stated.

Supplementary Material

Supplementals can be found within this manuscript file.

Acknowledgements

The authors thank the Center for Integrated Research Computing (CIRC) at the University of Rochester for providing computational resources and technical support and the URMC Flow Core for assistance with sorting. This work was supported by the US National Institutes of Health grants to V.G. and A.S. S.E. is a fellow of HFSP.

Author Contributions

S.E. designed and supervised research, performed experiments and analyzed data; S.E., A.S. and V.G. wrote the manuscript.

References

Buffenstein, R. & Ruby, J. G. Opportunities for new insight into aging from the naked mole-rat and other non-traditional models. *Nature Aging* **1**, 3-4, doi:10.1038/s43587-020-00012-4 (2021).

Ruby, J. G., Smith, M. & Buffenstein, R. Naked Mole-Rat mortality rates defy gompertzian laws by not increasing with age. *Elife* **7**, doi:10.7554/eLife.31157 (2018).

Tian, X. *et al.* High-molecular-mass hyaluronan mediates the cancer resistance of the naked mole rat. *Nature* **499**, 346-349, doi:10.1038/nature12234 (2013).

Tian, X. *et al.* INK4 locus of the tumor-resistant rodent, the naked mole rat, expresses a functional p15/p16 hybrid isoform. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 1053-1058, doi:10.1073/pnas.1418203112 (2015).

Azpurua, J. *et al.* Naked mole-rat has increased translational fidelity compared with the mouse, as well as a unique 28S ribosomal RNA cleavage. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 17350-17355, doi:10.1073/pnas.1313473110 (2013).

Emmrich, S. *et al.* Naked mole-rat hematopoietic stem and progenitors are highly quiescent with an inherent myeloid bias. *Nature communications* **submitted** (2021).

Emmrich, S. *et al.* An ectopic cervical thymus and no thymic involution until midlife in naked mole-rats. *Aging Cell* **submitted** (2021).

Higgins, J. M. Red blood cell population dynamics. *Clin Lab Med* **35**, 43-57, doi:10.1016/j.cll.2014.10.002 (2015).

Orkin, S. H. & Zon, L. I. Hematopoiesis: an evolving paradigm for stem cell biology. *Cell* **132**, 631-644, doi:10.1016/j.cell.2008.01.025 (2008).

McKinney-Freeman, S. *et al.* The transcriptional landscape of hematopoietic stem cell ontogeny. *Cell stem cell* **11**, 701-714, doi:10.1016/j.stem.2012.07.018 [doi] (2012).

Rebel, V. I., Miller, C. L., Eaves, C. J. & Lansdorp, P. M. The repopulation potential of fetal liver hematopoietic stem cells in mice exceeds that of their liver adult bone marrow counterparts. *Blood* **87**, 3500-3507 (1996).

Benz, C. *et al.* Hematopoietic stem cell subtypes expand differentially during development and display distinct lymphopoietic programs. *Cell stem cell* **10**, 273-283, doi:10.1016/j.stem.2012.02.007 (2012).

Bowie, M. B. *et al.* Identification of a new intrinsically timed developmental checkpoint that reprograms key hematopoietic stem cell properties. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 5878-5882, doi:0700460104 (2007).

Kiel, M. J. *et al.* SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. *Cell* **121**, 1109-1121, doi:10.1016/j.cell.2005.05.026 (2005).

Kim, I., He, S., Yilmaz, O. H., Kiel, M. J. & Morrison, S. J. Enhanced purification of fetal liver hematopoietic stem cells using SLAM family receptors. *Blood* **108**, 737-744, doi:10.1182/blood-2005-10-4135 (2006).

Hall, T., Sriram, P. & McKinney-Freeman, S. Murine Fetal Bone Marrow HSPCs Undergo a Dramatic Shift in Frequency at Birth. *Blood* **134**, 2471-2471, doi:10.1182/blood-2019-129073 (2019).

Hall, T. *et al.* 3083 – THE MURINE BONE MARROW NICHE DOES NOT SUPPORT A DIVERSE ARRAY OF HEMATOPOIETIC STEM AND PROGENITOR CELLS UNTIL

BIRTH. *Experimental Hematology* **88**, S63,
doi:https://doi.org/10.1016/j.exphem.2020.09.098 (2020).

18   Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single
cells. *Nature methods* **14**, 865-868, doi:10.1038/nmeth.4380 (2017).

19   Palis, J. Ontogeny of erythropoiesis. *Curr Opin Hematol* **15**, 155-161,
doi:10.1097/MOH.0b013e3282f97ae1 (2008).

20   Skulachev, V. P. *et al.* Neoteny, Prolongation of Youth: From Naked Mole Rats to
"Naked Apes" (Humans). *Physiological Reviews* **97**, 699-720,
doi:10.1152/physrev.00040.2015 (2017).

21   van der Maaten, L. & Hinton, G. Visualizing Data using t-SNE. *Journal of machine*
learning research : JMLR **9(Nov)**, 2579-2605 (2008).

22   Bens, M. *et al.* FRAMA: from RNA-seq data to annotated mRNA assemblies. *BMC*
genomics **17**, 54-58, doi:10.1186/s12864-015-2349-8 (2016).

23   Cramer, P., Bushnell, D. A. & Kornberg, R. D. Structural basis of transcription: RNA
polymerase II at 2.8 angstrom resolution. *Science (New York, N.Y.)* **292**, 1863-1876,
doi:10.1126/science.1059493 (2001).

24   Nishioka, K. *et al.* Expression and differential intracellular localization of two major forms
of human 8-oxoguanine DNA glycosylase encoded by alternatively spliced OGG1
mRNAs. *Molecular biology of the cell* **10**, 1637-1652, doi:10.1091/mbc.10.5.1637 (1999).

25   Li, C. *et al.* Downstream regulator TANK binds to the CD40 recognition site on TRAF3.
Structure (London, England : 1993) **10**, 403-411, doi:S0969-2126(02)00733-5 (2002).

26   Liu, X. *et al.* Regulation of mitochondrial biogenesis in erythropoiesis by mTORC1-
mediated protein translation. *Nature cell biology* **19**, 626-638, doi:10.1038/ncb3527
(2017).

27   Holyoake, T. L., Nicolini, F. E. & Eaves, C. J. Functional differences between
transplantable human hematopoietic stem cells from fetal liver, cord blood, and adult
marrow. *Experimental hematology* **27**, 1418-1427, doi:S0301-472X(99)00078-8 (1999).

28   Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set
collection. *Cell systems* **1**, 417-425, doi:10.1016/j.cels.2015.12.004 (2015).

29   Balabanian, K. *et al.* The chemokine SDF-1/CXCL12 binds to and signals through the
orphan receptor RDC1 in T lymphocytes. *The Journal of biological chemistry* **280**,
35760-35766, doi:M508234200 (2005).

30   Smith, J. S. & Rajagopal, S. The beta-Arrestins: Multifunctional Regulators of G Protein-
coupled Receptors. *The Journal of biological chemistry* **291**, 8969-8977,
doi:10.1074/jbc.R115.713313 (2016).

31   Chen, L. *et al.* Transcriptional diversity during lineage commitment of human blood
progenitors. *Science (New York, N.Y.)* **345**, 1251033, doi:10.1126/science.1251033
(2014).

32   Casero, D. *et al.* Long non-coding RNA profiling of human lymphoid progenitor cells
reveals transcriptional divergence of B cell and T cell lineages. *Nature immunology* **16**,
1282-1291, doi:10.1038/ni.3299 (2015).

33   Corces, M. R. *et al.* Lineage-specific and single-cell chromatin accessibility charts
human hematopoiesis and leukemia evolution. *Nature genetics* **48**, 1193-1203,
doi:10.1038/ng.3646 (2016).

34   Notta, F. *et al.* Distinct routes of lineage development reshape the human blood
hierarchy across ontogeny. *Science (New York, N.Y.)* **351**, aab2116,
doi:10.1126/science.aab2116 (2016).

35   Drissen, R., Thongjuea, S., Theilgaard-Monch, K. & Nerlov, C. Identification of two
distinct pathways of human myelopoiesis. *Science immunology* **4**,
10.1126/sciimmunol.aau7148, doi:eaau7148 (2019).

36 Copley, M. R. *et al.* The Lin28b-let-7-Hmga2 axis determines the higher self-renewal
potential of fetal haematopoietic stem cells. *Nature cell biology* **15**, 916-925,
doi:10.1038/ncb2783 (2013).
37 McKinney-Freeman, S. *et al.* The transcriptional landscape of hematopoietic stem cell
ontogeny. *Cell Stem Cell* **11**, 701-714, doi:10.1016/j.stem.2012.07.018 (2012).
38 Collins, E. C. *et al.* Mouse Af9 is a controller of embryo patterning, like Mll, whose
human homologue fuses with Af9 after chromosomal translocation in leukemia. *Mol Cell*
Biol **22**, 7313-7324, doi:10.1128/MCB.22.20.7313-7324.2002 (2002).
39 Broxmeyer, H. E. Colony assays of hematopoietic progenitor cells and correlations to
clinical situations. *Critical reviews in oncology/hematology* **1**, 227-257, doi:S1040-
8428(84)80013-X (1984).
40 McKenzie, J. L., Gan, O. I., Doedens, M. & Dick, J. E. Human short-term repopulating
stem cells are efficiently detected following intrafemoral transplantation into NOD/SCID
recipients depleted of CD122+ cells. *Blood* **106**, 1259-1261, doi:10.1182/blood-2005-03-
1081 (2005).
41 Greer, J. P. *et al. Wintrobe's Clinical Hematology*. 14th edn, (Lippincott Williams &
Wilkins (LWW), 2019).
42 Naumann, U. *et al.* CXCR7 functions as a scavenger for CXCL12 and CXCL11. *PLoS*
One **5**, e9175, doi:10.1371/journal.pone.0009175 (2010).
43 Levoye, A., Balabanian, K., Baleux, F., Bachelerie, F. & Lagane, B. CXCR7
heterodimerizes with CXCR4 and regulates CXCL12-mediated G protein signaling.
Blood **113**, 6085-6093, doi:10.1182/blood-2008-12-196618 (2009).
44 Hartmann, T. N. *et al.* A crosstalk between intracellular CXCR7 and CXCR4 involved in
rapid CXCL12-triggered integrin activation but not in chemokine-triggered motility of
human T lymphocytes and CD34+ cells. *J Leukoc Biol* **84**, 1130-1140,
doi:10.1189/jlb.0208088 (2008).
45 Tarnowski, M. *et al.* CXCR7: a new SDF-1-binding receptor in contrast to normal
CD34(+) progenitors is functional and is expressed at higher level in human malignant
hematopoietic cells. *Eur J Haematol* **85**, 472-483, doi:10.1111/j.1600-0609.2010.01531.x
(2010).
46 Kumar, R. *et al.* CXCR7 mediated Gialpha independent activation of ERK and Akt
promotes cell survival and chemotaxis in T cells. *Cell Immunol* **272**, 230-241,
doi:10.1016/j.cellimm.2011.09.015 (2012).
47 Rajagopal, S. *et al.* Beta-arrestin- but not G protein-mediated signaling by the "decoy"
receptor CXCR7. *Proc Natl Acad Sci U S A* **107**, 628-632,
doi:10.1073/pnas.0912852107 (2010).

Figures and Legends

Figure 1

Fig. 1. Marrow immunophenotyping reveals a specific neonatal progenitor stage.

a, Representative FACS gating of viable naked mole-rat adult [left] and neonate [right] whole BM cells. GC, neutrophil granulocytes; TC, T cells; HSPC, hematopoietic stem and progenitor cells; MEP, megakaryocytic erythroid progenitor; ERY, erythroid cells; NMP, neonatal myeloid progenitor. **b**, Frequencies of $LIN^+/Thy1.1^{hi}/CD34^-$ GCs in adult (N=46) and neonatal (N=23) BM; p-values were determined by unpaired Student's t-test. Representative FACS gating of viable naked mole-rat adult [left] and neonate [right] **c**, whole BM or **d**, LIN^+ BM cells. LIN, lineage cocktail CD11b/CD18/CD90/CD125; TC90, $CD11b^-/CD90^+/Thy1.1^{int/hi}/CD34^{hi}$. **e**, Frequencies of TC90s in NBM (N=23) and neonatal spleens (N=22); p-values were determined by unpaired Welch's t-test. **f**, Frequencies of neonatal BM cell fractions. p-value determined by Brown-Forsythe's One-way ANOVA; N=23.

Figure 2

Fig. 2. Neonatal HSPC compartment frequencies in mice and naked mole-rats.

a, Representative FACS gating of viable mouse neonate [left] and adult [right] LIN$^-$ BM cells. LK,

LIN$^-$/Sca-1$^-$/Kit$^+$; LSK, LIN$^-$/Sca-1$^+$/Kit$^+$. **b**, Frequencies of LSKs across mouse hematopoietic

tissues. Adult BM (N=12), neonate BM (N=18), adult spleen (N=10), neonate spleen (N=16), adult

liver (N=10), neonate liver (N=16); p-value determined by Sidak's Two-way ANOVA. **c**,

Representative FACS gating of viable naked mole-rat neonate [left] and adult [right] LIN$^-$ BM cells.

CP3, LIN$^-$/Thy1.1^{lo}/CD34^{hi}; CP4, LIN$^-$/Thy1.1$^-$/CD34^{hi}; CP5, LIN$^-$/Thy1.1$^-$/CD34^{lo}; CP6, LIN$^-$

/Thy1.1^{lo}/CD34$^-$; CP7, LIN$^-$/Thy1.1$^-$/CD34$^-$; LTC, LIN$^-$/Thy1.1^{int}/CD34^{hi}. **d**, Frequencies of LTCs

across naked mole-rat hematopoietic tissues. Adult BM (N=47), NBM (N=23), adult spleen

(N=29), neonate spleen (N=23), adult liver (N=13), neonate liver (N=18); p-value determined by

Sidak's Two-way ANOVA.

Figure 3

Fig. 3. Distinct expression and splicing patterns in neonatal TC90.

a, Unsupervised t-SNE clustering of neonate TC90 (N=5) and adult LTC (N=5), CP1 (N=5), CP3

(N=4), CP4 (N=5) transcriptomes. LTC/CP1 and CP3/4 form overlapping partitions, TC90s cluster

as a distinct group. **b**, Heat- [top panel] and Coldmap [bottom panel] of differentially expressed

genes (DEG) for each of the three groupings TC90, CP1/LTC, CP3/4. **c**, GSEA of naked mole-

rat neonate and adult BM HSPCs using each DEG signature genes. GeneRatio, (signature ∩

term) / (signature ∩ all terms); NES, normalized enrichment score; displayed the top genesets

filtered for GeneRatio > 0.041 and NES > 0.1 | NES < 0.1; each $q < 10^{-4}$. Note that the HSC

genesets enriched in TC90s are distinct from those enriched in CP1/LTC. **d**, POLR2B; **f**, OGG1;

g, TANK. Isoform downregulated in TC90, green; Isoform upregulated in TC90, black. **e**, POLR2B

domains in amino acids (AA): Rpb2_1 protrusion, black; Rpb2_2 lobe, red; Rpb2_6 hybrid binding

& wall, blue; β chain signature, purple; Rbp2_7 anchor & clamp, orange. **g**, OGG1 domains: MTS,

orange; OGG_N N-terminal, blue; ENDO3c endonuclease, yellow; HhH-GPD motif, black. **i**,

TANK transcripts in nucleotides (nt): TOP motif, red; ORF, white box.

Figure 4

Fig. 4. TC90 cells inversely expressed the CXCR4/CXCL12 ratio.

a, Unsupervised t-SNE clustering, color code used on heatmaps throughout this Figure. CB-HSC,

human neonatal cord-blood HSCs; Adult HSC, human adult HSC; LTC, adult naked mole-rat

HSPCs. **b**, Heatmap showing upregulated [top] and downregulated [bottom] orthologs with framed

TC90 DEG signature; z-score derived from voom-transformed transcripts per million (TPMs). **c**,

GSEA of TC90 DEG signature displaying the top 10 q-value terms filtered for NES > 0.1 and

GeneRatio > 0.02; each q < 10^{-6}. Circle size, GeneRatio; Square color, -log10(q-value).

Figure 5

Fig. 5. Extended cross-species comparison confirms CXCR7/CXCL12 overexpression.

Unsupervised t-SNE clustering with **a**, Colors for cell types, symbols for datasets; **b**, Colors for

ontogeny, symbols for species. hHSC, human HSCs; hMLP, human multipotent lymphoid

progenitor; hCLP, human common lymphoid progenitor; hGMP, human granulocytic monocytic

progenitor; hsa, *homo sapiens*; hgl, *heterocephalus glaber*. **c**, Log-transformed TPM expression

across cell types, species and ontogenies for orthologs CXCL12 [top], CXCR7 [middle] and

CXCR4 [bottom]. **d**, Heatmap the leading edge genes of the top GSEA term

EPITHELIAL_MESENCHYMAL_TRANSITION ($q < 5.5 \times 10^{-4}$; NES = 2.54); z-score derived from

voom-transformed TPMs.

Figure 6

Fig. 6. TC90s are myeloid-biased progenitors without homing capacity *in vivo*.

a, Colony assay of sorted naked mole-rat adult (N=4) and neonate (N=3) BM HSPCs, grown at 32°C for 3 weeks. Error bars denote s.d., p-value determined by Two-way ANOVA, corrected for multiple testing with Two-stage linear step-up procedure of Benjamini, Krieger and Yekutieli. **b**, Naked mole-rat adult CP1 or neonate TC90 cells were FACS-purified and xenotransplanted (TX) into NSGS hosts via **c**, Intravenous (*i.v.*) injection, host whole BM from all combined long bones at 4 weeks post TX. Chimerism gated as viable cells positive for either or both naked mole-rat cross-reactive HSPC markers CD34 and Thy1.1-antigen. **d**, Intrafemoral (*i.f.*) injection, host whole BM from each single long bone at 4 weeks post TX. e, Representative FACS gating for chimerism on *i.f.* injected hosts with adult CP1 [left panel] or neonate TC90 [right panel].

532   **Supplementary Figures and Legends**

Supplementary Figure 1

Supplementary Fig. 1. TC90 immunophenotyping, Mouse stem and progenitor FACS.

a, Frequencies of Thy1.1$^+$/CD34$^+$ cells in adult (N=13) and neonatal (N=18) livers; p-values were determined by unpaired Welch's t-test. **b**, Single color flow cytometry histograms of neonatal BM co-stained for CD90-antigen [left panel] and CD11b [right panel] and the LIN/Thy1.1/CD34 labelling; blue line, LIN$^+$/Thy1.1^{hi}/CD34$^-$ GCs; green line, Thy1.1^{int}/CD34$^-$ TCs; red line, LIN$^-$/Thy1.1$^-$/CD34$^-$ CP7 as negative controls; shaded AUC, LIN$^+$/Thy1.1$^{int/hi}$/CD34^{hi} TC90s. Representative FACS gating of neonate [left panel] or adult [right panel] BM **c**, LSKs or **d**, LKs; Frequencies of adult (N=6) and neonatal (N=9) populations see scatter respective plots, p-values were determined by Sidak's Two-way ANOVA.

Supplementary Figure 2

Supplementary Fig. 2. RNA-Seq TC90 vs adult naked mole-rat HSPCs.

a, Frequencies of CP4 and CP5 across adult BM (N=48), neonate BM (N=23), adult spleen (N=30)

and neonatal spleen (N=22); p-value determined by Tukey's Two-way ANOVA. **b**, Filtered,

normalized and log2-transformed naked mole-rat transcriptomes, color code see **c**, Unsupervised

t-SNE clustering of neonate TC90 (N=5) and adult LTC (N=5), CP1 (N=5), CP3 (N=4), CP4 (N=5)

isoform counts. The clustering pattern of gene counts in Figure 3a is reproduced: LTC/CP1 and

CP3/4 form overlapping partitions, TC90s cluster as a distinct group. **d**, Heatmap of 12 inversely

expressed transcript pairs from the same gene locus, whereof each single isoform is differentially

expressed in TC90 vs other CPs.

Supplementary Figure 3

 Supplementary Fig. 3. RNA-Seq cross-species, cross-ontogenies.

 a, RLE-plot of filtered, size factor-normalized, log2-transformed and median-centered TPMs

 [transcripts per million] of human and naked mole-rat transcriptomes. **b**, Log-transformed TPM

 expression across cell types, species and ontogenies for indicated orthologs.

Supplementary Figure 4

Supplementary Fig. 4. Naked mole-rat colony assays.

a, Photomicrographs of representative CFU morphologies from an adult BM LTC sample, grown for 3 weeks at 32°C; Scale bar 250µm. Hematoxylin & Eosin staining of spleen sections; Scale bar 200µm [bottom]. **b**, Merged Scan of representative colony assays for adult CP1 [left] and neonate TC90 [right]; Scale bar 400mm. Note that TC90 features larger scattered colonies, pointing towards granulocytic precursors with higher regenerative potential than adult CFU-GMs.