Title: Artificial mimicry of seasonal transcriptome dynamics in *Arabidopsis thaliana* reveals short-

and long-term responses to environmental conditions

Authors / Affiliations

Yuko Kurita[1], Hironori Takimoto[2], Mari Kamitani[1], Yoichi Hashida[3], Makoto Kashima[1], Ayumi

Tezuka[1], Takanari Tanabata[4], Atsushi J. Nagano[1]

[1]Faculty of Agriculture, Ryukoku University, 1-5 Yokotani, Ooe-cho, Seta, Otsu, Shiga 520-2194

Japan

[2]Faculty of Computer Science and Systems Engineering, Okayama Prefectural University, Soja-shi,

Okayama 719−1197 Japan

[3]Faculty of Agriculture, Takasaki University of Health and Welfare, 54 Nakaorui-machi,

Takasaki-shi, Gunma 370-0033, Japan

[4]Kazusa DNA Research Institute, 2-6-7 Kazusa-kamatari, Kisarazu, Chiba 292-0818, Japan

Author for correspondence: Atsushi J. Nagano

17   e-mail: anagano@agr.ryukoku.ac.jp

Total wordcount: 4940

Introduction: 401

Materials and methods: 1510

Results: 2189

Discussion: 840

Number of figures: 7

Color figures: All figures

Number of tables: 0

Supporting information: 11 figures, 11 tables

Summary

- Plants must respond to various environmental factors that change seasonally. In a
previous study, seasonally oscillating genes were identified by a massive
time-series transcriptome analysis in a wild population of *Arabidopsis halleri* ssp.
gemmifera, a sister species of *Arabidopsis thaliana*.

- To analyze the function of these seasonally oscillating genes, we established an
experimental system to mimic seasonal expression trends using *A. thaliana*.
Arabidopsis thaliana plants were cultured under conditions that mimicked
average monthly temperatures and daylengths in a "smart growth chamber mini,"
a hand-made low-cost small chamber.

- Under different short-term incubations, the seasonal trends of 1627 seasonally
oscillating genes were mimicked. These seasonally oscillating genes had varying
temporal responsiveness (constant, transient, and incremental).

- Our findings suggest that plants perceive and integrate information about
environmental stimuli in the field by combining seasonally oscillating genes with
temporal responsiveness.

Key words

Introduction

Introduction

Plants responses to seasonal changes in environmental factors are essential mechanisms.

The analysis of gene expression patterns is a useful approach to understand how plants respond to

environmental stimuli and adapt physiologically and morphologically. Over the past few decades, a

number of studies have analyzed seasonal gene expression patterns (Andersson *et al.*, 2004;

Galindo González *et al.*, 2012; Nagano *et al.*, 2012, 2019; Lu *et al.*, 2020). Vigorous research has

uncovered the mechanisms of several seasonal phenomena, such as vernalization in

flowering, growth cessation, and bud dormancy (Song *et al.*, 2013; Maurya & Bhalerao, 2017;

Falavigna *et al.*, 2019; Luo & He, 2020). However, there are many seasonal phenomena whose

mechanisms have not yet been elucidated, or genes that show seasonally oscillating expression

patterns but whose functions have not yet been clarified. While the seasonal phenomena of plants

are fascinating, they take a lot of time to investigate and these investigations are difficult. In

addition to field studies, multifaceted approaches are needed to improve the efficiency and speed of

this type of research.

Several genes that show seasonal oscillation patterns in gene expression in *Arabidopsis*

halleri ssp. *gemmifera* (*Ahg*), a perennial herbaceous plant closely related to *Arabidopsis thaliana*

(*Ath*), have been identified by seasonal transcriptome analysis (Nagano *et al.*, 2019). The seasonal

expression patterns of many of these seasonally oscillating genes (SO genes) were defined by

annual temperature changes rather than annual changes in daylength. However, the molecular

mechanisms regulating the seasonal expression of a large number of SO genes are still unknown.

The expression patterns of these genes may be affected by long-term environmental stimuli. For

example, *FLC* is associated with vernalization in *Ath* (Jean Finnegan, 2015; Bouché *et al.*, 2017).

However, it is also expected that there are SO genes whose seasonal expression patterns are formed

by transient responses to environmental stimuli. Functional analyses are needed to distinguish the

temporal responsiveness of SO genes to environmental stimuli.

We aimed to mimic seasonal expression trends in the field using model plants and a

simple experimental system in the laboratory. In this study, we cultured *Ath* plants under average

temperature and daylength conditions of each month for a short period (7, 3, or 1-day) to test how

many genes can mimic the seasonal expression of *Ahg*. To culture plants in parallel under different

temperature and daylength conditions, we developed a hand-made low-cost small chamber, called a

smart growth chamber mini (SGCmini).

Materials and Methods

Smart growth chamber mini (SGCmini)

The system configuration of the developed smart growth chamber mini (SGCmini) is shown in Supporting Information Fig. **S1a**. The proposed system consisted of an incubator body, Peltier unit, circuit unit, and control unit. Supporting Information Fig. **S1b** depicts the proposed system, excluding the control unit. The incubator body was made of acrylic; the bottom surface was made of black acrylic and the rest was transparent acrylic (thickness: 10 mm). The transparent acrylic parts were all integrated and had holes into which the Peltier unit could be attached (described below). The outer dimensions of the transparent acrylic part were 80 mm × 200 mm × 200 mm.

The appearance of the Peltier unit is shown in Supporting Information Fig. **S1c**. This unit comprised a 40 × 40 mm Peltier device, two heat sinks, and two fans. The heat sinks were adhered on both sides of the Peltier device and the black insulation sheets enclosed them. The side with the small heat sink was inserted through the hole in the incubator body and housed in the body. A small fan was installed to circulate the air inside the incubator. The side with the larger heat sink was for heat dissipation, which enabled air cooling with a large fan.

The circuit unit consisted of an Arduino Uno R3, a DC brush motor driver shield for Arduino (SHIELD-MD10, Cytron Technologies, Malaysia), and two temperature sensors (MCP9808, Microchip Technology, USA). Two temperature sensors were installed in the incubator. The circuit unit sends information from the temperature sensors to the control unit (PC) via serial communication.

The control unit was equipped with a GUI, which determined the extent of temperature control for heating and cooling to achieve the "target temperature set by the GUI" in the incubator. The proportional integral (PI) control determined the temperature control according to the current temperature in the incubator, and it was sent to the circuit unit as the control signal. According to the signal received, the circuit unit controlled the current sent to the Peltier unit by pulse-width

modulation (PWM).

The GUI of the control unit is illustrated in Supporting Information Fig. **S1d**. In this GUI,

the temperature in the incubator could be set in increments of 0.5 °C. In addition to the current

temperature, the temperature profile for the past 48 h could be displayed on a chart. The GUI also

provided other functions, such as the output of a temperature control log and the capability to

change the set temperature. The developed incubator could control the temperature with an error

range of 0.2 °C within a range of −9 to +20 °C from the standard room temperature. For colder

conditions, such as that for February, the SGCmini was operated in a growth chamber (LH-241SP,

NK system Osaka, Japan) set below 15 °C.

For plant cultivation, the incubator body was painted black on four sides to block external

light (Fig. **1a**). An LED light unit (25 pcs; color: red and blue; size: 30.8 × 30.8 cm; PPFD:

approximately 35 μmol m^{-2} s^{-1}) was placed on the top of the incubator body. The lighting of the

LED light unit was controlled using a timer.

Setting of culture condition in SGCmini

To mimic the seasonal expression of *Arabidopsis halleri* ssp. *gemmifera* (*Ahg*) in the field,

simplified conditions of the environment of the *Ahg* sampling site (the Omoide River, Taka-cho,

35° 06′ N, 134° 55′ E, altitude 190–230 m above sea level) (Nagano *et al.*, 2019) were set as culture

conditions (Fig. **2a**). For daylength settings, sunrise and sunset times on the 15th of each month in

Nishiwaki city were obtained from the National Astronomical Observatory of Japan. For

temperature settings, hourly average temperatures at Nishiwaki (averaged over 2006–2016) were

obtained from the Japan Meteorological Agency. The average value of the temperature from sunrise

to sunset was set as the daytime temperature, and the average value of the temperature from sunset

to sunrise was set as the nighttime temperature. It was difficult to mimic temperatures below 5 °C,

so when the temperature was below 5 °C, such as in January, the experiment was conducted with a

5 °C setting.

We also conducted a high-temperature culture experiment, in which the daytime

temperature of the August condition was changed to 32 °C or 35 °C (Fig. **4a**).

Plant materials

Seeds of *A. thaliana* (*Ath*) (Col-0, CS70000) were sown on 1/2 MS medium with 0.5%

sucrose and 0.6% gellan gum in square Petri dishes and cultured at 20 °C with 8 h light for 20 d.

Plants were transferred to 1/2 MS medium with 0.25% gellan gum in round plates and cultivated for

3, 7, or 9 d under the same conditions as the pre-culture to recover from the stress of transplanting.

After the pre-culture, the round plates were placed in the SGCmini. Plants were cultivated under the

conditions of each month for 7, 3, or 1 d (Fig. **2b** and **5a**). To equalize the age of the plants at the

sampling date, we extended the pre-culture period to shorten the culture period. However, for a

1-day culture with short pre-culture conditions, the pre-culture period was set to 3 d in order to

evaluate the effect of plant age at the start date of treatment (Supporting Information Fig. **S8a**).

For morphometry, all leaves were cut off and scanned using a scanner (GT-S650, EPSON,

Japan). Leaf area and leaf length were measured from the scanned images using ImageJ software.

ANOVA and Tukey HSD were performed in R software to test for differences between conditions

($P < 0.05$).

RNA-Seq

For RNA extraction, leaves larger than 1 mm without cotyledons and juvenile leaves

were harvested from one plant as one sample between 11:00 and 13:00. The leaves were

immediately frozen in liquid nitrogen and stored at −80 °C until extraction.

Samples were ground with two zirconia beads using a TissueLyser II (QIAGEN, MD,

USA) with pre-chilled adapters at −80 °C. Total RNA from leaves was extracted using a Maxwell

16 LEV Plant RNA Kit (Promega, WI, USA). The amount of RNA was determined using Quant-iT

RNA Assay Kit, broad range (Thermo Fisher Scientific, Waltham, MA, USA) and Tecan plate

reader Infinite 200 PRO (Tecan, Männedorf, Switzerland). For RNA-Seq library preparation, 400 or

500 ng of total RNA per sample was used. The library was prepared using the Lasy-Seq v1.0 or

v1.1 protocol (Kamitani *et al.*, 2019) (https://sites.google.com/view/lasy-seq/). The quality of the

library was assessed using a Bioanalyzer (Agilent Technologies). The libraries were sequenced with

a HiSeq 2500 or Hiseq X (Illumina) to produce single reads of 50 bp (Library 1 and 2) or 150bp

(Library 3).

Preprocessing and quality filtering of RNA-Seq data were performed using

Trimmomatic-0.33 (Bolger *et al.*, 2014). Preprocessed reads were mapped on transcript sequences

in TAIR 10 with Bowtie1(v1.1.1) (Langmead *et al.*, 2009) and quantified using RSEM-1.2.21(Li &

Dewey, 2011). The output of the RSEM was analyzed using R (R Core Team, 2019).

RNA-Seq data of *A. halleri* were obtained from previous studies (Nagano *et al.*, 2019).

RNA-Seq data of *A. thaliana* and *A. halleri* were analyzed as follows. Samples with fewer than

$10^{5.5}$ reads were excluded from the analysis. The attributes of the samples used in the analysis are

listed in Supporting Information Table **S1**. *Ahg* genes with an average $\log_2(\text{rpm}+1) > 2$ were

designated as expressed genes (16158 genes) (Supporting Information Fig. **S2**). A total of 90.3% of

the expressed genes in *Ahg* (14587 genes) were annotated as orthologs of *Ath* genes by reciprocal

BLAST. Subsequent seasonal expression analysis was performed using these annotated

Ahg-expressed genes (14587 genes) and their orthologous genes in *Ath*.

Analysis of seasonal expression

To compare the seasonal trends of *Ahg* and *Ath*, the expressions of the *Ahg*-expressed

genes and their orthologous genes in *Ath* were fitted to a cosine curve using the *nls* function in R

(Fig. **3a**). Amplitude and phase were obtained from each fitted cosine curve over a 1-y period.

Genes with an amplitude > 1 were defined as seasonally oscillating genes (SO genes). Since the

value of $\log_2(\text{rpm} + 1)$ was fitted with a cosine curve, an amplitude > 1 indicates that the difference

in seasonal oscillation is more than two-fold. We compared the amplitude and phase of *Ahg* and *Ath*

genes to determine "seasonality-induced genes (SI genes)" that could mimic the seasonal

expression trends of *Ahg* in the field. A gene whose amplitude difference was smaller than 1 and

whose phase difference was smaller than 45 d was defined as a SI gene.

Gene ontology enrichment analysis

For gene ontology (GO) enrichment analysis, GO annotations for *Ath* genes were

obtained from the TAIR database. Statistical tests of enrichment analysis were performed using the

Fisher's exact test function in R. A total of 14,472 expressed genes with GO annotations were

included in the test. Multiple testing corrections were performed using FDR (Benjamini &

Hochberg, 1995). Representative GO terms were summarized based on the REVIGO output

(http://revigo.irb.hr) (Supek *et al.*, 2011) (allowed similarity = 0.5) and visualized with reference to

a previous protocol (Bonnot *et al.*, 2019).

Results

Smart growth chamber mini

To cultivate plants under various conditions, we developed a space- and cost-saving small

incubator named as "smart growth chamber mini" (SGCmini) (Fig. **1a**; Supporting Information Fig.

S1). The SGCmini can control the inside temperature within a range of –9 to +20 °C from the

outside temperature. The Peltier unit was controlled to equalize the average of the measured

temperatures of the two sensors installed at the diagonal corners of the incubator body with the set

temperature (Fig. **1b–g**). It took a few minutes to 30 min for the average temperature in the

SGCmini to reach the set temperature (Fig. **1c,d**). The temperature records of the SGCmini under

the March and August conditions are shown in Fig. **1e** and **1f**. Multiple SGCminis can be controlled

using a single PC. This allows us to save space and conduct multiple experiments under various

conditions simultaneously in one room.

Pseudo-seasonal change of *Ath* plant growth in SGCmini for 7 d

To mimic the seasonal trend of gene expression in *A. halleri* (*Ahg*) in the natural

environment, *A. thaliana* (*Ath*) plants were cultivated in the SGCmini for 7 d under simplified

temperature and daylength settings based on the conditions of the natural habitat for each month

(Supporting Information Table **S2**; Fig. **2a,b**). Sunrise and sunset times on the 15th of each month

were set as light-on and light-off times. The average temperature from sunrise to sunset for each

month was set as the daytime temperature, and the average temperature from sunset to sunrise for

each month was set as the nighttime temperature.

The total leaf area and maximum leaf blade length increased from February to May,

remained constant from May to September, and decreased after October (Fig. **2c–e,h**). The number

of leaves increased from February to May, remained constant from May to June, decreased slightly

in July and August, and then increased again in September before decreasing after October (Fig. **2f**).

The petiole length increased from April to July and then decreased after August (Fig. **2d,g**). Thus,

the 7-d culture induced morphological differences under each month's condition.

Comparison of seasonal trends of *Ahg* in field and pseudo-seasonal trends of *Ath* in SGCmini

for 7 d

To analyze how many genes mimic the seasonal expression trend of *Ahg* genes, gene

expression data of *Ath* plants grown in the SGCmini for 7 d were obtained by RNA-Seq. Samples

with more than $10^{5.5}$ reads were used for analysis (*Ahg*: 474 samples; *Ath*: 72 samples; Supporting

Information Fig. **S2a**). *Ahg*-expressed genes (14587 genes) ($\log_2$(mean rpm +1) > 2) and their

orthologous genes in *Ath* were used for analysis (Supporting Information Fig. **S2**). The seasonal

trend of gene expression was summarized as the amplitude (α) and phase (φ) by cosine curve fitting

(Fig. **3a**). The phase was defined as the time point of the highest gene expression in a year. We

defined genes whose amplitude was greater than 1 as SO genes (Fig. **3b**).

Based on this criterion, 4312 and 4016 genes were identified as SO genes in *Ahg* and *Ath*,

respectively. Although phases of these SO genes were concentrated in summer (June to August) or

winter (December to February) in both *Ahg* and *Ath*, the phase of *Ahg* SO genes was wider than that

of *Ath* SO genes (Fig. **3c**). In *Ahg*, 48% (2099 genes) and 33.5% (1445 genes) of the 4312 SO genes

had phases distributed in summer and winter, respectively. In *Ath*, 52% (2125 genes) and 42%

(1693 genes) of the 4016 SO genes had phases distributed in summer and winter, respectively.

Among 4312 *Ahg* SO genes and 4016 *Ath* SO genes, 1757 genes overlapped (Fig. **3d**). By

comparing the amplitude and phase in *Ath* and *Ahg*, we defined genes whose amplitude difference

was < ± 1 and whose phase difference was < ± 45 days as seasonality-induced genes (SI genes) (Fig.

3e−g). Between *Ahg* plants grown in the field and *Ath* plants cultured in the SGCmini for 7 d, 946

genes were found to be SI genes.

To elucidate the functional aspects of the SI genes, we tested the enrichment of genes

with a functional annotation. We found that 63 GOs were significantly enriched (adjusted *P* < 0.05,

Supporting Information Table **S3**). For example, genes with annotations of 'response to temperature

stimulus' (GO: 0009266, adjusted $P = 0.009$) were enriched. One of these genes, *AhgLHY* (*LATE*

ELONGATED HYPOCOTYL 1), which is a core clock gene, has been reported in a previous study

as a gene that showed seasonally oscillating expression at noon. *AthLHY* (AT1G01060.1) had

almost the same seasonal trend as *AhgLHY* (Fig. **3h**). It has been reported that the amplitude of the

circadian rhythm of *AhgLHY* changes seasonally, and that sampling at noon results in a seasonal

trend of high expression in winter and low expression in summer (Nagano *et al.*, 2019).

Mimicry of gene expression in response to higher temperature

Several of the genes annotated as "response to temperature stimuli" were SI genes, such

as *LHY*, and others were not. One possible reason for the non-mimicry is the mild temperature

setting of the SGCmini. For example, although we chose the average temperatures, in summer,

plants transiently face higher temperatures in the field. To mimic this condition, a high-temperature

experiment was performed. In the experiment, daytime temperature in the August condition was

changed to 32 °C or 35 °C (Fig. **4a,b**).

GAMMA TONOPLAST INTRINSIC PROTEIN 1 (*TIP1;1*, AT2G36830) and *VACUOLAR*

INVERTASE 1 (*VI1*, AT1G62660) showed pseudo-seasonal oscillations in *Ath* (Fig. **4c,d**).

Although their phases were almost the same as that of *Ahg*, their amplitudes were smaller than the

threshold (amplitude difference < 1). When cultivated under higher temperatures, the expressions of

these genes were successfully elevated to a level similar to that in *Ahg*.

In contrast, in the field, *Ahg* orthologs of *HEAT SHOCK PROTEIN 18.2* (*HSP18.2*,

AT5G59720) and *HSP22* (AT4G10250) showed heterogeneous high gene expression levels among

plant individuals in summer (Fig. **4e,f**). Heterogeneous expression is speculated to be a response to

the heterogeneous microenvironment around individual plants, that is, locally elevated temperatures

(Nagano *et al.*, 2019). Although the expressions of *HSP18.2* and *HSP22* were not high under

August conditions (daytime temperature: 28.3 °C), these expressions were notably elevated at 32 °C

and 35 °C (Fig. **4e,f**). Thus, for some genes, non-average conditions, such as the highest or lowest

temperatures in a day, are necessary to mimic the seasonal trends in gene expression.

Pseudo-seasonal change of plant growth in SGCmini for 1 or 3 d

To investigate the effect of culture period length, plants were cultured in the SGCmini for

1 or 3 d under the condition of each month (Fig. **5**; Supporting Information Fig. **S3** and **S4**). As

shown in Fig. **5a**, pre-culture periods were changed depending on the culture period in each month

condition to make the days after sowing uniform when morphology was measured. As the culture

period in each month became shorter, the pseudo-seasonal change in morphology became more

moderate (Fig. **5b,c**). In the 3-d culture, total leaf area increased under July and August conditions

(Fig. **5b**; Supporting Information Fig. **S3d**). Petiole length increased under warmer conditions (June

to September). The number of leaves slightly increased under the July, August, and September

conditions (Fig. **5b**; Supporting Information Fig. **S3f**). The leaf blade length slightly increased

under the July condition (Fig. **5b**; Supporting Information Fig. **S3g**). In the 1-d culture, there was no

significant difference in plant growth (Fig. **5b**; Supporting Information Fig. **S4**). Morphological

observations showed that the differences among the conditions of each month were smaller in the

shorter culture period lengths.

Differences between culture periods (7, 3, and 1-d) at $P < 0.001$ using ANOVA and

Tukey's HSD test are shown in Fig. **5b**. Taking the number of tests into account, we chose a

conservative threshold (results of differences at $P < 0.05$ are shown in Supporting Information

Tables **S4–7**). Significant differences in total leaf area were detected between 7- and 3-d cultures in

March and between 7- and both 3- and 1 d cultures in November. Significant differences in the

number of leaves were detected between 7- and both 3- and 1-d cultures in January and between 7-

and 3-d cultures in February and August, indicating that development of visible leaves tended to be

suppressed under these conditions in the 7-d culture. Significant differences in maximum petiole

length were detected between 7- and 1-d cultures from June to September, indicating that petiole

elongation tended to be promoted under these conditions as the culture period increased. A

significant difference in maximum petiole length was also detected between the 7- and 3-d cultures

in March. No significant difference was detected in the leaf blade length between the culture

periods.

Comparison of seasonal expression trends of *Ahg* and *Ath* cultured in SGCmini for

a shorter period

RNA-Seq was performed to clarify how the length of the culture period affected the

pseudo-seasonal trend of gene expression in *Ath*. The SO genes and SI genes were defined in the

same manner as the analysis of 7-d culture data (Fig. **6**; Supporting Information Fig. **S5** and **S6**).

The number of SO genes was similar in the 7-d culture (4016 genes) and 3-d culture (3975 genes)

(Fig. **6a**). The number of SO genes in the 1-d culture was greater than that in the 3- and 7-d cultures

(4575 genes). The number of SO genes that were annotated as orthologs of *Ahg* SO genes showed

the same tendency (7-d culture: 1757 genes, 3-d culture: 1766 genes, 1-d culture: 1886 genes). As

the culture period became shorter, the phase distribution became more concentrated in summer and

winter (Fig. **6b**). A total of 412 SI genes were shared with three conditions (Fig. **6c**). Examples of

the shared genes were *LHY* and S*UCROSE SYNTHASE 1* (*SUS1*, AT5G20830). These genes

showed almost the same seasonal trends regardless of the length of the culture period (Fig. **7a,b**).

Interestingly, hundreds of genes mimicked the *Ahg* seasonal trend only when the culture

period was longer or shorter (SI genes in 7-d culture but not in 1-d culture: 436 genes, SI genes in

1-d culture but not in 7-d culture: 508 genes). *ALCOHOL DEHYDROGENASE 1* (*ADH1*,

AT1G77120) and *JACALIN-RELATED LECTINS 5* (*JAL5*, AT1G52000) were classified as SI

genes in the 1-d culture but not in the 7-d culture (Fig. **6d, 7c,d**). As the culture period increased,

the amplitude of these genes decreased. *SWEET17* (AT4G15920) and *TRANSPARENT TESTA 8*

(*TT8*, AT4G09820) were classified as SI genes in the 7-d culture but not in the 1-d culture (Fig. **6c,**

7e,f). The amplitude of these genes decreased as the culture period became shorter. TT8 is a bHLH

transcription factor that regulates flavonoid biosynthesis with other factors (Li, 2014; Xu *et al.*, 2015). Anthocyanin regulatory and biosynthetic pathways and predicted seasonal expression plots of *Ahg* and *Ath* are shown in Supporting Information Fig. **S7–10**. Among these genes, the amplitudes of the pseudo-seasonal trend of *DIHYDROFLAVONOL 4-REDUCTASE* (*DFR*, AT5G42800), *LEUCOANTHOCYANIDIN DIOXYGENASE* (*LDOX*, AT5G42800), *PRODUCTION OF ANTHOCYANIN PIGMENT 1* (*PAP1*, AT1G56650), and *PAP2* (AT1G66390) became larger as the culture period increased, similar to *TT8* (Supporting Information Fig. **S7a,c, S8**, and **S10**). In our criteria, *F3'H*, *MYB11*, *MYB111*, *PAP1*, *PAP2*, *TT8*, *GL3*, and *EGL3* were defined as SO genes in *Ahg*. Among them, only *TT8* and *PAP1* were defined as SI genes.

The reason for the difference in the amplitude of the seasonal trend among the three conditions in *Ath* could be the difference in the length of the culture period (7, 3, and 1 d) and/or the difference in the plant age at the start of the treatment in the SGCmini (23, 27, and 29 d). To investigate the effect of plant age on the amplitude of SO genes, plants at 23 d after seeding were cultured in the SGCmini for 1 d under the January and August conditions (Supporting Information Fig. **S11a**), and RNA-Seq was performed. The expression levels of SO genes of younger plants tended to have slightly smaller differences between the January and August conditions (Supporting Information Fig. **S11b**). However, the variation due to the difference in the length of the culture period was greater than that due to plant age (Supporting Information Fig. **S11b**–d).

In total, 1627 genes were identified as SI genes in at least one of the three conditions (Fig. **6c**; Supporting Information Fig. **S2b,** Table **S8**). To elucidate the functional aspects of the SI and non-SI genes, we performed GO enrichment analysis (adjusted $P < 0.05$, Supporting Information Tables –**S9–11**). In all SI genes, the GO terms "gene expression," "response to abiotic stimulus," and "response to temperature stimulus" were enriched (Fig. **6e**). In all non-SI genes, the GO terms "secondary metabolic process," "response to biotic stimulus," and "response to abiotic stimulus" were enriched. The GO terms "secondary metabolic process" and "glucosinolate metabolic process" were enriched in genes that passed the amplitude criteria but were not in phase with *Ahg*. This

353 indicates that several SO genes associated with these GO terms require other factors (other

354 environmental stimuli, etc.) to mimic the seasonal trend of *Ahg* in the appropriate phase, even

355 though they were induced under mimic conditions in the present experiment.

Discussion

Short-term incubation of *Ath* plants in the SGCmini under conditions representing each month could mimic the seasonal trends of 1627 SO genes in *Ahg*. The results of the 7-, 3-, and 1-d culture periods revealed that the culture period required for mimicry differed depending on the SO gene (Fig. **6c, 7;** Supporting Information Fig. **S8**). For some SO genes, the length of the culture period seems to be unrelated to the mimicking of seasonal trends of *Ahg* gene expression in the field because 412 SI genes were observed under all three conditions. These SO genes may ensure robustness to environmental changes by responding accurately to the current environment, regardless of the past environment. *LHY* was one of these SO genes (Fig. **7a**) and is known as a core clock gene that contributes to temperature compensation of the circadian clock (Gould *et al.*, 2006; Salomé *et al.*, 2010; Gil & Park, 2019). In field environments, temperatures can change drastically, even within a few days. The constant responsiveness of *LHY* may contribute to maintaining the circadian rhythm under fluctuating environmental conditions in the field.

Conversely, there were several SO genes that mimicked the *Ahg* seasonal trend only when the culture period was shorter (Fig. **7c,d**). *ADH1* was one of these SO genes. Song *et al*. showed that *ADH1* transcription was induced at 3 h following 4 °C stress treatment, reached the highest expression level at 24 h, then declined (Song *et al.*, 2017). Since its amplitude decreased under longer incubations, *ADH1* seems to respond to environmental changes (in this experiment, changes in temperature and daylength from pre-culture conditions to conditions of each month in the SGCmini), rather than to absolute values at the time, as LHY does. *ADH1* is not only involved in the cold stress response (Song *et al.*, 2017), but its transcription is also known to be induced by various other stresses (Winter *et al.*, 2007; Shi *et al.*, 2017). SO genes such as *ADH1*, which mimicked field seasonal trends only in shorter culture periods, may play a role in responding to short-term environmental changes due to their transient expression.

Several SO genes mimicked field seasonal trends only in longer culture periods, such as *SWEET17* and *TT8* (Fig. **7e,f**). In *Ath*, late biosynthetic genes (LBGs) of the anthocyanin

biosynthetic pathway, *DFR* and *LDOX*, and their transcriptional regulators, *TT8*, *PAP1*, and *PAP2*,

showed an increase in the amplitude of the seasonal trend with the extension of the culture period,

suggesting a regulatory mechanism that responds to continuous environmental stimuli such as low

temperature (Supporting Information Fig. **S7–10**). One possible mechanism driving this is

epigenetic regulation. Recent studies have revealed that epigenetic regulation is involved in the

regulation of anthocyanin biosynthesis gene expression (Cai *et al.*, 2019; Zheng *et al.*, 2019). In the

field, analysis of seasonal histone modifications of *Ahg* suggests that histone H3 lysine 27

trimethylation is involved in gene expressions that depend on memory of longer-term (more than a

few weeks) environmental conditions (Nishio *et al.*, 2020). In our study, although the plants only

experienced the treatment once, it is known that the memory of past intermittent stimuli such as hot

or cold temperatures can also affect plant response (Leuendorf *et al.*, 2020; Yamaguchi *et al.*, 2021).

Future studies will need to include these findings to clarify the response to fluctuating field

environments. The differences in the temporal responsiveness of SO genes may be one factor that

contributes to the variation in the phase of the seasonal trend (Fig. **6b**).

Early biosynthetic genes (EBGs) and LBGs, except *F3'H*, did not show seasonal

oscillations in *Ahg* in the field (Supporting Information Fig. **S7** and **S8**). However, their

transcriptional regulators showed seasonal oscillations in various phases (Supporting Information

Fig. **S7, S9,** and **S10**), suggesting that the constitutive expression of EBGs and LBGs in the field is

the outcome of the integration of multiple environmental stimuli by transcriptional regulators.

Flavonoid biosynthesis is affected not only by low temperature, but also by light quality, drought,

nutrition, and virus infection (Cominelli *et al.*, 2008; Lillo *et al.*, 2008; Cai *et al.*, 2019; Honjo *et al.*,

2020). While highly oscillating genes are eye-catching, it is also important to remember that

seemingly non-oscillating genes may be involved in plant responses to seasonal changes. Such a

complex response mechanism of secondary metabolites may be one of the reasons why the GO

term "secondary metabolic process" was detected in the GO analysis of SO genes that could not be

mimicked (Fig. **6e**).

In conclusion, we successfully mimicked the seasonal trends of approximately 38% of

Ahg SO genes using *Ath* plants in an experimental system set at the average temperature and

daylength of each month. This is the first study to comprehensively analyze the response of SO

genes to relatively short-term environmental stimuli. The results of this study and the experimental

system will be useful for investigating the function of unknown SO genes. To adapt to complex

environmental changes in the field, plants probably interpret seasonal changes by converting

environmental stimuli into gene expression at different temporal resolutions.

Acknowledgements

We greatly appreciate Hiroshi Kudoh and his laboratory members for field data

acquisition and discussions, Tetsuro Mimura for his advice and discussions on the experimental

system, Kyoko Mogami for her help with RNA-Seq library preparation, and Motohiro Mihara and

Hitoshi Ooshima of Dynacom for their support in RNA-Seq analysis, Daisuke Kyogoku for his

advice on English language editing, Editage (www.editage.com) for English language editing. This

work was supported by JST CREST (JPMJCR15O2) and ACCEL (JPMJAC1403).

Author Contribution

A.J.N. and Y.K. designed the research. H.T. and T.T. developed SGCmini. M. Kamitani,

Y.H., M. Kashima, A.T. contributed to library preparation and data analysis. Y.K. performed

experiments and data analysis. Y.K., H.T. and A.J.N. wrote the paper.

Data Availability

The RNA-seq data were submitted to the NCBI Sequence Read Archive repository under

the BioProject number PRJNA739047 (https://www.ncbi.nlm.nih.gov/bioproject/ PRJNA739047).

R scripts and data required for analysis (count data, etc.) are available via the GitHub repository

(https://github.com/naganolab/AthSGCmini_pseudo-seasonal_RNA-Seq).

Reference

Andersson A, Keskitalo J, Sjödin A, Bhalerao R, Sterky F, Wissel K, Tandre K, Aspeborg H,

Moyle R, Ohmiya Y, *et al.* **2004**. A transcriptional timetable of autumn senescence. *Genome biology*

5: R24.

Benjamini Y, Hochberg Y. **1995**. Controlling the False Discovery Rate : A Practical and Powerful

Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**:

289–300.

Bolger AM, Lohse M, Usadel B. **2014**. Trimmomatic: A flexible trimmer for Illumina sequence data.

Bioinformatics **30**: 2114–2120.

Bonnot T, Gillard M, Nagel D. **2019**. A Simple Protocol for Informative Visualization of Enriched

Gene Ontology Terms. *Bio-Protocol* **e3429**.

Bouché F, Woods DP, Amasino RM. **2017**. Winter memory throughout the plant kingdom:

Different paths to flowering. *Plant Physiology* **173**: 27–35.

Cai H, Zhang M, Chai M, He Q, Huang X, Zhao L, Qin Y. **2019**. Epigenetic regulation of

anthocyanin biosynthesis by an antagonistic interaction between H2A.Z and H3K4me3. *New*

Phytologist **221**: 295–308.

Cominelli E, Gusmaroli G, Allegra D, Galbiati M, Wade HK, Jenkins GI, Tonelli C. **2008**.

Expression analysis of anthocyanin regulatory genes in response to different light qualities in

Arabidopsis thaliana. *Journal of Plant Physiology* **165**: 886–894.

Falavigna V da S, Guitton B, Costes E, Andrés F. **2019**. I want to (Bud) break free: The potential

role of DAM and SVP-like genes in regulating dormancy cycle in temperate fruit trees. *Frontiers in*

Plant Science **9**: 1990.

Galindo González LM, El Kayal W, Ju CJT, Allen CCG, King-Jones S, Cooke JEK. **2012**.

Integrated transcriptomic and proteomic profiling of white spruce stems during the transition from

active growth to dormancy. *Plant, Cell and Environment* **35**: 682–701.

Gil KE, Park CM. **2019**. Thermal adaptation and plasticity of the plant circadian clock. *New*

Phytologist **221**: 1215–1229.

Gould PD, Locke JCW, Larue C, Southern MM, Davis SJ, Hanano S, Moyle R, Milich R,

Putterill J, Millar AJ, *et al.* 2006. The molecular basis of temperature compensation in the

Arabidopsis circadian clock. *Plant Cell* **18**: 1177–1187.

Honjo MN, Emura N, Kawagoe T, Sugisaka J, Kamitani M, Nagano AJ, Kudoh H. **2020**.

Seasonality of interactions between a plant virus and its host during persistent infection in a natural

environment. *ISME Journal* **14**: 506–518.

Jean Finnegan E. **2015**. Time-dependent stabilization of the +1 nucleosome is an early step in the

transition to stable cold-induced repression of FLC. *Plant Journal* **84**: 875–885.

Kamitani M, Kashima M, Tezuka A, Nagano AJ. **2019**. Lasy-Seq: a high-throughput library

preparation method for RNA-Seq and its application in the analysis of plant responses to fluctuating

temperatures. *Scientific Reports* **9**: 1–14.

Langmead B, Trapnell C, Pop M, Salzberg SL. **2009**. Ultrafast and memory-efficient alignment of

short DNA sequences to the human genome. *Genome Biology* **10**: R25.

Leuendorf JE, Frank M, Schmülling T. **2020**. Acclimation, priming and memory in the response of

Arabidopsis thaliana seedlings to cold stress. *Scientific Reports* **10**: 1–11.

Li S. **2014**. Transcriptional control of flavonoid biosynthesis: Fine-tuning of the MYB-bHLH-WD40

(MBW) complex. *Plant Signaling and Behavior* **9**: e27522.

Li B, Dewey CN. **2011**. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC Bioinformatics* **12**: 323.

Lillo C, Lea US, Ruoff P. **2008**. Nutrient depletion as a key factor for manipulating gene expression

and product formation in different branches of the flavonoid pathway. *Plant, Cell and Environment*

31: 587–601.

Lu H, Gordon MI, Amarasinghe V, Strauss SH. **2020**. Extensive transcriptome changes during

seasonal leaf senescence in field-grown black cottonwood (Populus trichocarpa Nisqually-1).

Scientific Reports **10**: 1–14.

Luo X, He Y. **2020**. Experiencing winter for spring flowering: A molecular epigenetic perspective on

vernalization. *Journal of Integrative Plant Biology* **62**: 104–117.

Maurya JP, Bhalerao RP. **2017**. Photoperiod- and temperature-mediated control of growth cessation

and dormancy in trees: A molecular perspective. *Annals of Botany* **120**: 351–360.

Nagano AJ, Kawagoe T, Sugisaka J, Honjo MN, Iwayama K, Kudoh H. **2019**. Annual

transcriptome dynamics in natural environments reveals plant seasonal adaptation. *Nature Plants* **5**:

74–83.

Nagano AJ, Sato Y, Mihara M, Antonio BA, Motoyama R, Itoh H, Nagamura Y, Izawa T. **2012**.

Deciphering and prediction of transcriptome dynamics under fluctuating field conditions. *Cell* **151**:

1358–1369.

Nishio H, Nagano AJ, Ito T, Suzuki Y, Kudoh H. **2020**. Seasonal plasticity and diel stability of

H3K27me3 in natural fluctuating environments. *Nature Plants* **6**: 1091–1097.

R Core Team. **2019**. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria.

Salomé PA, Weigel D, McClunga CR. **2010**. The role of the Arabidopsis morning loop components

CCA1, LHY, PRR7, and PRR9 in temperature compensation. *Plant Cell* **22**: 3650–3661.

Shi H, Liu W, Yao Y, Wei Y, Chan Z. **2017**. Alcohol dehydrogenase 1 (ADH1) confers both abiotic

and biotic stress resistance in Arabidopsis. *Plant Science* **262**: 24–31.

Song YH, Ito S, Imaizumi T. **2013**. Flowering time regulation: Photoperiod- and

temperature-sensing in leaves. *Trends in Plant Science* **18**: 575–583.

Song Y, Liu L, Wei Y, Li G, Yue X, An L. **2017**. Metabolite profiling of adh1 mutant response to

cold stress in Arabidopsis. *Frontiers in Plant Science* **7**: 2072.

Supek F, Bošnjak M, Škunca N, Šmuc T. **2011**. Revigo summarizes and visualizes long lists of

gene ontology terms. *PLoS ONE* **6**: e21800.

Winter D, Vinegar B, Nahal H, Ammar R, Wilson G V., Provart NJ. **2007**. An 'electronic

fluorescent pictograph' Browser for exploring and analyzing large-scale biological data sets. *PLoS*

ONE **2**: e718.

Xu W, Dubos C, Lepiniec L. **2015**. Transcriptional control of flavonoid biosynthesis by

MYB-bHLH-WDR complexes. *Trends in Plant Science* **20**: 176–185.

Yamaguchi N, Matsubara S, Yoshimizu K, Seki M, Hamada K, Kamitani M, Kurita Y,

Nomura Y, Nagashima K, Inagaki S, *et al.* **2021**. H3K27me3 demethylases alter HSP22 and

HSP17.6C expression in response to recurring heat in Arabidopsis. *Nature Communications* **12**: 1–16.

Zheng T, Tan W, Yang H, Zhang L, Li T, Liu B, Id DZ, Id HL. **2019**. Regulation of anthocyanin

accumulation via MYB75/HAT1/TPL-mediated transcriptional repression. *PLoS Genetics* **15**:

e1007993.

 Figure legends

Fig. 1 Performance of the smart growth chamber mini (SGCmini). (**a**) Picture of the SGCmini. The

walls of the incubator body were painted for shading. (**b**) Temperature plot of a test run. The

temperature was changed stepwise from 16 °C to 26 °C every 2 h. Light blue line: outside temperature;

pink lines: inside temperature measured by the sensors; black line: preset temperature; red line: average

temperature of sensor 1 and sensor 2. (**c**) Enlarged plot of (**b**) at the lowest temperature setting. (**d**)

Enlarged plot of (**b**) at the highest temperature setting. (**e**) Temperature plot for 3 d under the March

condition. (**f**) Temperature plot for 3 d under the August condition.

Fig. 2 Experimental settings and morphological traits of *Arabidopsis thaliana* (*Ath*) plants grown in

the SGCmini for 7 d. (**a**) Setting of each condition in the SGCmini. Yellow bars: daylength; red circles:

daytime temperatures; blue triangles; nighttime temperatures. (**b**) Schematic diagram of the culture

condition. (**c**) Whole images of plants grown under February, May, and July conditions. (**d**) Leaves of

plants grown under February, May, and July conditions. The youngest leaves are on the right. Bar = 10

mm. (**e**) Total leaf area. (**f**) Number of leaves. (**g**) Maximum petiole length. (**h**) Maximum leaf blade

length. Bar: SD. n = 5–9. Different letters indicate differences between conditions at $P < 0.05$ using

ANOVA and Tukey's HSD test.

Fig. 3 Comparison of seasonal trends of gene expression in *Arabidopsis halleri* ssp. *gemmifera*

(*Ahg*) in the field and *Arabidopsis thaliana* (*Ath*) in the SGCmini for 7 d. (**a**) Schematic diagram of

comparison method of seasonal trends of *Ahg* and *Ath*. Black circles indicate observed value by

RNA-Seq. The solid orange (*Ahg*) and blue (*Ath*) lines show the fitted cosine curve. The orange and

blue dashed lines indicate the phases of *Ahg* and *Ath*. "α" and "φ" indicate the amplitude and the

phase of the fitted cosine curve. (**b**) Histograms of amplitude (α) of expressed genes. Left: *Ahg*.

Right: *Ath*. The regions with $\alpha > 1$ are colored. (**c**) Histograms of phase (φ) of expressed genes.

Left: *Ahg*. Right: *Ath*. The regions with $\alpha > 1$ are colored. (**d**) Venn diagram of *Ahg* and *Ath* SO

genes. Left circle: *Ahg*. Right circle: *Ath*. (**e**) Scatter plot of amplitudes in *Ahg* and *Ath*. Only genes

with $\alpha > 1$ in both species are colored according to density using the *densCols* function in R. (**f**)

Scatter plot of phases in *Ahg* and *Ath*. Only genes with $\alpha > 1$ in both species are colored according

to density using the *densCols* function in R. (**g**) Venn diagram of genes that passed amplitude and

phase criteria. Left circle: genes that passed amplitude criteria ($\Delta\alpha < \pm 1$). Right circle: genes that

passed phase criteria ($\Delta\varphi < \pm 45$ days). (**h**) Seasonal expression plots of *LHY*. From left, fitted

cosine curve plot of *Ahg* and *Ath*, observed value and fitted line plot of *Ahg* and *Ath*, respectively.

Fig. 4 Effect of higher temperature on gene expression. (**a**) Schematic diagram of culture conditions.

(**b**) Conditions of the high temperature experiment. Yellow bars: daylength; red circles: daytime

temperatures; blue triangles: nighttime temperatures. (**c**) Gene expression plot of *TIP1;1*. From left,

fitted cosine curve plot of *Arabidopsis halleri* ssp. *gemmifera* (*Ahg*) and *Arabidopsis thaliana* (*Ath*),

observed value and fitted line plot of *Ahg* and *Ath*, respectively, and a plot of temperature and

expression levels. Black circles indicate observed values. Orange (*Ahg*) and blue (*Ath*) lines show

the fitted cosine curve. Orange and blue dashed lines indicate the phases of *Ahg* and *Ath*,

respectively. Pink triangles and red squares indicate observed values when plants were grown at

32 °C and 35 °C. (**d**) *VI1*. (E) *HSP18.2*. (F) *HSP22.0*.

Fig. 5 Comparison of morphological traits of *Arabidopsis thaliana* (*Ath*) plants grown in the

SGCmini for 1, 3, or 7 d. (**a**) Schematic diagram of culture conditions. (**b**) Morphological traits.

From left, total leaf area, number of leaves, maximum petiole length, and maximum leaf blade

length. Dark blue squares, middle blue triangles, and light blue circles indicate data for plants

grown in the SGCmini for 7, 3, and 1 d, respectively. n = 4–9. Different letters below the plots

indicate differences between culture periods at $P < 0.001$ using ANOVA and Tukey's HSD test. (**c**)

Leaves of plants grown under the August condition. From the top, 7-, 3- and 1-d cultures. The

youngest leaves are on the right. Bar = 10 mm.

Fig. 6 Comparison of seasonal trends of gene expression between different culture periods. (**a**)

Comparison of the number of seasonally oscillating (SO) genes. Darker regions indicate the number

of *Arabidopsis thaliana* (*Ath*) SO genes common with the *Arabidopsis halleri* ssp. *gemmifera* (*Ahg*)

SO genes. (**b**) Comparison of the distribution of phases of SO genes. Orange circles: *Ahg*; dark blue

rhombuses: *Ath* 7-d culture; blue squares: *Ath* 3-d culture; light blue triangles: *Ath* 1-d culture. (**c**)

Venn diagram of seasonality-induced genes (SI genes). Left circle: *Ath* 7-d culture; right circle: *Ath*

3-d culture; bottom circle: *Ath* 1-d culture. (**d**) Scatter plot of the alpha of SI genes in the 7-d culture

(vertical axis) and 1-d culture (horizontal axis). Each point is colored based on the local density of

points. Pink circles indicate genes shown in Fig. **7**. (**e**) Enriched biological process gene ontology

(GO) terms. "total SI gene" indicates total SI genes that mimicked the seasonal trend of *Ahg* in at

least one of the three conditions (7-, 3-, and 1-d culture). "total non-SI gene" indicates non-SI genes

that did not mimic the seasonal trend of *Ahg*. "non-SI gene (phase)" indicates genes that passed the

amplitude criteria but were not in phase with *Ahg*. Circle size indicates the number of genes. Color

scale indicates the adjusted *P* values in the GO enrichment analysis.

Fig. 7 Seasonal expression plots of seasonality-induced genes. From left, fitted cosine curve plot

of *Arabidopsis halleri* ssp. *gemmifera* (*Ahg*) and *Arabidopsis thaliana* (*Ath*), observed value and

fitted line plot of *Ahg* and *Ath* in the 7-, 3-, and 1-d cultures, respectively. (**a**) *LHY*. (**b**) *SUS1*. (**c**)

ADH1. (**d**) *JAL5*. (**e**) *SWEET17*. (**f**) *TT8*.

Supporting Information

Fig. S1 Design of the smart growth chamber mini (SGCmini).

Fig. S2 Summary of sample sets and gene sets in this study.

Fig. S3 Morphological traits of *Arabidopsis thaliana* (*Ath*) plants grown in the SGCmini for 3 d.

Fig. S4 Morphological traits of *Arabidopsis thaliana* (*Ath*) plants grown in the SGCmini for 1 d.

Fig. S5 Comparison of seasonal trends of gene expression in *Arabidopsis halleri* ssp. *gemmifera*

(*Ahg*) and *Arabidopsis thaliana* (*Ath*) grown in the SGCmini for 3 d.

Fig. S6 Comparison of seasonal trends of gene expression in *Arabidopsis halleri* ssp. *gemmifera*

(*Ahg*) and *Arabidopsis thaliana* (*Ath*) grown in the SGCmini for 1 d.

Fig. S7 Anthocyanin regulatory and biosynthetic pathways and seasonal trend plots of

Arabidopsis halleri ssp. *gemmifera* (*Ahg*) and *Arabidopsis thaliana* (*Ath*).

Fig. S8 Seasonal expression plots of early biosynthetic genes (EBGs) and late biosynthetic genes

(LBGs) in the anthocyanin biosynthetic pathway.

Fig. S9 Seasonal expression plots of transcriptional regulators that regulate early biosynthetic

genes (EBGs) in the anthocyanin biosynthetic pathway.

Fig. S10 Seasonal expression plots of transcriptional regulators that regulate late biosynthetic

genes (LBGs) in the anthocyanin biosynthetic pathway.

Fig. S11 Comparison of gene expression patterns by length of pre-culture period.

(a)

	Seeding	Trans planting	Start of treatment	Sampling
		Pre-culture		Culture in SGCmini
7-day culture	20 days	3 days		7 days
3-day culture	20 days	7 days		3 days
1-day culture	20 days	9 days		1 day

20°C, 8h light · each month condition

(b)

Total leaf area (mm²) — Number of leaves — Petiole length (mm) — Leaf blade length (mm)

day: 1, 3, 7

(c)

Aug condition

7-day

3-day

1-day