Title: Integration of short- and long-term responses to environmental stimuli shape seasonal

transcriptome dynamics

Running Head: Artificial mimicry of seasonal transcriptome

Corresponding Author: A. J. Nagano

Faculty of Agriculture, Ryukoku University, 1-5 Yokotani, Ooe-cho, Seta, Otsu, Shiga 520-2194

Japan

Tel: +81-77-599-5656

e-mail: anagano@agr.ryukoku.ac.jp

Subject Areas: (2) environmental and stress responses, (3) regulation of gene expression

Number of black and white figures: 0

Color figures: 7

Number of tables: 0

supplementary material: 12 figures, 14 tables

Title: Integration of short- and long-term responses to environmental stimuli shape seasonal transcriptome dynamics

Running Head: Artificial mimicry of seasonal transcriptome

Authors / **Affiliations**

Yuko Kurita[1], Hironori Takimoto[2], Mari Kamitani[1], Yoichi Hashida[3], Makoto Kashima[1], Ayumi Tezuka[1], Takanari Tanabata[4], Atsushi J. Nagano[1,5,*]

[1]Faculty of Agriculture, Ryukoku University, 1-5 Yokotani, Ooe-cho, Seta, Otsu, Shiga 520-2194 Japan

[2]Faculty of Computer Science and Systems Engineering, Okayama Prefectural University, Soja-shi, Okayama 719−1197 Japan

[3]Faculty of Agriculture, Takasaki University of Health and Welfare, 54 Nakaorui-machi, Takasaki-shi, Gunma 370-0033, Japan

[4]Kazusa DNA Research Institute, Department of Frontier Research and Development, 2-6-7 Kazusa-kamatari, Kisarazu, Chiba 292-0818, Japan

[5]Institute for Advanced Biosciences, Keio University, 403-1 Nipponkoku, Daihouji, Tsuruoka, Yamagata, 997-0017, Japan

Author for correspondence: Atsushi J. Nagano

e-mail: anagano@agr.ryukoku.ac.jp

Abstract

Plants must respond to various seasonally changing environmental stimuli. In a previous

study, seasonally oscillating genes were identified by a massive time-series transcriptome analysis

for the wild population of *Arabidopsis halleri* ssp. *gemmifera*, a sister species to *Arabidopsis*

thaliana. It was not clear how environmental stimuli shaped the seasonal expression pattern of these

seasonally oscillating genes. In this study we show that responses in different time-scales

contributed to the formation of seasonal expression patterns for several genes. To analyze the

seasonally oscillating genes, we established an experimental system to mimic seasonal expression

trends using *A. thaliana* and a "smart growth chamber mini," a hand-made low-cost small chamber.

Arabidopsis thaliana plants were cultured under conditions that mimicked the average monthly

temperatures and daylengths under different day-scale incubation. In total, the seasonal trends of

1627 seasonally oscillating genes were mimicked, and they showed varying temporal responses

(constant, transient, and incremental) to environmental stimuli. Our results suggest that plants

perceive and integrate information regarding environmental stimuli in the field by combining

seasonally oscillating genes with different temporal responsiveness.

Introduction

Plants responses to seasonal changes in environmental factors are essential mechanisms. The analysis of gene expression patterns is a useful approach to understand how plants respond to environmental stimuli and adapt physiologically and morphologically. Over the past few decades, a number of studies have analyzed seasonal gene expression patterns (Andersson et al., 2004; Galindo González et al., 2012; Lu et al., 2020; Nagano et al., 2019, 2012). Vigorous research has uncovered the mechanisms of several seasonal phenomena, such as vernalization in flowering, growth cessation, and bud dormancy (Falavigna et al., 2019; Luo and He, 2020; Maurya and Bhalerao, 2017; Song et al., 2013). However, there are many seasonal phenomena whose mechanisms have not yet been elucidated, or genes that show seasonally oscillating expression patterns but whose functions have not yet been clarified.

The previous study have identified several genes that show seasonal oscillation patterns in gene expression in *Arabidopsis halleri* ssp. *gemmifera* (*Ahg*), a perennial herbaceous plant closely related to *Arabidopsis thaliana* (*Ath*) by seasonal transcriptome analysis (Nagano et al., 2019). The seasonal expression patterns of many of these seasonally oscillating genes (SO genes) were defined by annual temperature changes rather than annual changes in daylength. However, the molecular mechanisms regulating the seasonal expression of a large number of SO genes are still unknown. Expression patterns of several SO genes are thought to be affected by long-term environmental stimuli (month-scale response: weeks to months) such as *FLC,* which is associated with vernalization in *Ath* (Bouché et al., 2017; Jean Finnegan, 2015). On the contrary, it is thought that other SO genes show a minute-scale response (minutes to hours) to the environmental stimuli, such as *CBF/DREB1*s for the cold stress response (Gong et al., 2020; Ye et al., 2019). In addition, the mechanism of day-scale response (1 day to 1 week) to repeated temperature stimulation has recently attracted attention (Leuendorf et al., 2020; Yamaguchi et al., 2021). Nagano et al. revealed

87   that there are numerous genes whose expression can be explained by meteorological data from the

88   past one to several days (Nagano et al., 2012).

89   The seasonal phenomena of plants are fascinating; however, field studies are

time-consuming and challenging. Moreover, in the field, plants are simultaneously exposed to many

biotic and abiotic environmental stimuli. To understand the molecular mechanisms of plant

responses in the field, effective experimental systems which can separate and control the

environmental stimuli, as well as field studies, are essential. Recently, combination studies of the

field and experimental systems, such as ecotrons (Roy et al., 2021) and the smart growth chamber

(SGC) (Hashida et al., 2022), which can reproduce microclimatic conditions, have been reported. In

the study of the function of SO genes, understanding their properties by using controlled

environments will be helpful. For example, what stimuli shape the seasonal trends of the SO gene

and the length of time required for stimulus acceptance.

99   In this study, we focused on the responses to temperature and day-length occurring within

several days (day-scale responses). We cultured *Ath* plants under simplified monthly conditions

with average temperature and daylength of each month for 7, 3, or 1-day(s) to test how many genes

can mimic the seasonal expression of *Ahg*. Compared to *Ahg* plants, which are difficult to culture in

large quantities in the laboratory, *Ath* plants can be cultured easily and quickly, and abundant

genomic information of such plants is available. In addition, *Ath* allows for further research on SO

genes using a variety of molecular genetic tools. To culture plants in parallel under different

temperature and daylength conditions, we developed a hand-made low-cost small chamber, called a

smart growth chamber mini (SGCmini).

Results

Development of a space- and cost-saving incubator with parallel control

To cultivate plants under various conditions, we developed a space- and cost-saving small incubator named as "smart growth chamber mini" (SGCmini) (Fig. **1A**; Supplementary Fig. **S1**). The SGCmini can control the inside temperature within a range of –9 to +20 °C from the outside temperature. The Peltier unit was controlled to equalize the average of the measured temperatures of the two sensors installed at the diagonal corners of the incubator body with the set temperature (Fig. **1B–G**). It took a few minutes to 30 min for the average temperature in the SGCmini to reach the set temperature (Fig. **1C,D**). The temperature records of the SGCmini under the March and August conditions are shown in Fig. **1E** and **1F**. Multiple SGCminis can be controlled using a single PC. This allows us to save space and conduct multiple experiments under various conditions simultaneously in one room.

The 7-d culture in smart growth chamber mini induced morphological differences under conditions of each month

To mimic the seasonal trend of gene expression in *A. halleri* (*Ahg*) in the natural environment, *A. thaliana* (*Ath*) plants were cultivated in the SGCmini for 7 d under simplified temperature and daylength settings based on the conditions of the natural habitat for each month (Supplementary Table **S2**; Fig. **2A,B**). In the Northern Hemisphere, summer occurs from June through August, and winter occurs from December through February. Sunrise and sunset times on the 15[th] of each month were set as light-on and light-off times. The average temperature from sunrise to sunset for each month was set as the daytime temperature, and the average temperature from sunset to sunrise for each month was set as the nighttime temperature.

The total leaf area and maximum leaf blade length increased from February to May, remained constant from May to September, and decreased after October (Fig. **2C–E,H**). The

number of leaves increased from February to May, remained constant from May to June, decreased

slightly in July and August, and then increased again in September before decreasing after October

(Fig. **2F**). The petiole length increased from April to July and then decreased after August (Fig.

2D,G). Thus, the 7-d culture induced morphological differences under each month's condition.

Seasonal expression of *Ahg* was partially reproduced in *Ath* plants cultured for 7 days under

the condition of each month

To analyze how many genes mimic the seasonal expression trend of *Ahg* genes, gene

expression data of *Ath* plants grown in the SGCmini for 7 d were obtained by RNA-Seq. Samples

with more than $10^{5.5}$ reads were used for analysis (*Ahg*: 474 samples; *Ath*: 72 samples;

Supplementary Fig. **S2A**). *Ahg*-expressed genes (14587 genes) ($\log_2$(mean rpm +1) > 2) and their

orthologous genes in *Ath* were used for analysis (Supplementary Fig. **S2**). The seasonal trend of

gene expression was summarized as the amplitude (α) and phase (φ) by cosine curve fitting (Fig.

3A). The phase was defined as the time point of the highest gene expression in a year. We defined

genes whose amplitude was greater than 1 as SO gene (Fig. **3B**).

Based on this criterion, 4312 and 4016 genes were identified as SO genes in *Ahg* and *Ath*,

respectively. Although phases of these SO genes were concentrated in summer (June to August) or

winter (December to February) in both *Ahg* and *Ath*, the phase of *Ahg* SO genes was wider than that

of *Ath* SO genes (Fig. **3C**). In *Ahg*, 48% (2099 genes) and 33.5% (1445 genes) of the 4312 SO

genes had phases distributed in summer and winter, respectively. In *Ath*, 52% (2125 genes) and

42% (1693 genes) of the 4016 SO genes had phases distributed in summer and winter, respectively.

Among 4312 *Ahg* SO genes and 4016 *Ath* SO genes, 1757 genes overlapped (Fig. **3D**). By

comparing the amplitude and phase in *Ath* and *Ahg*, we defined genes whose amplitude difference

was < ± 1 and whose phase difference was < ± 45 days as seasonality reproduced genes (SR genes)

(Fig. **3E−G**). Between *Ahg* plants grown in the field and *Ath* plants cultured in the SGCmini for 7 d,

946 genes were found to be SR genes.

To elucidate the functional aspects of the SR genes, we tested the enrichment of genes

with a functional annotation. We found that 63 GOs were significantly enriched (adjusted $P < 0.05$,

Supplementary Table **S3**). For example, genes with annotations of 'response to temperature

stimulus' (GO: 0009266, adjusted $P = 0.009$) were enriched. One of these genes, *AhgLHY* (*LATE*

ELONGATED HYPOCOTYL 1), which is a core clock gene, has been reported in a previous study

as a gene that showed seasonally oscillating expression at noon. *AthLHY* (AT1G01060.1) had

almost the same seasonal trend as *AhgLHY* (Fig. **3H**). It has been reported that the amplitude of the

circadian rhythm of *AhgLHY* changes seasonally, and that sampling at noon results in a seasonal

trend of high expression in winter and low expression in summer (Nagano et al., 2019).

Several SO genes required extreme high temperatures to reproduce their seasonal trends

Several of the genes annotated as "response to temperature stimuli" were SR genes, such

as *LHY*, and others were not. One possible reason for the non-mimicry is the mild temperature

setting of the SGCmini. For example, although we chose the average temperatures, in summer,

plants transiently face higher temperatures in the field. To mimic this condition, a high-temperature

experiment was performed. In the experiment, daytime temperature in the August condition was

changed to 32 °C or 35 °C (Fig. **4A,B**).

GAMMA TONOPLAST INTRINSIC PROTEIN 1 (*TIP1;1*, AT2G36830) and *VACUOLAR*

INVERTASE 1 (*VI1*, AT1G62660) showed pseudo-seasonal oscillations in *Ath* (Fig. **4C,D**).

Although their phases were almost the same as that of *Ahg*, their amplitudes were smaller than the

threshold (amplitude difference < 1). When cultivated under higher temperatures, the expressions of

these genes were successfully elevated to a level similar to that in *Ahg*.

In contrast, in the field, *Ahg* orthologs of *HEAT SHOCK PROTEIN 18.2* (*HSP18.2*,

AT5G59720) and *HSP22* (AT4G10250) showed heterogeneous high gene expression levels among

plant individuals in summer (Fig. **4E,F**). Heterogeneous expression is speculated to be a response to

the heterogeneous microenvironment around individual plants, that is, locally elevated temperatures

(Nagano et al., 2019). Although the expressions of *HSP18.2* and *HSP22* were not high under

August conditions (daytime temperature: 28.3 °C), these expressions were notably elevated at 32 °C

and 35 °C (Fig. **4E,F**). Thus, for some genes, non-average conditions, such as the highest or lowest

temperatures in a day, are necessary to mimic the seasonal trends in gene expression.

Shorter cultivation periods resulted in smaller morphological seasonal changes

To investigate the effect of culture period length, plants were cultured in the SGCmini for

1 or 3 d under the condition of each month (Fig. **5**; Supplementary Fig. **S3** and **S4**). As shown in Fig.

5A, pre-culture periods were changed depending on the culture period in each month condition to

make the days after sowing uniform when morphology was measured. As the culture period in each

month became shorter, the pseudo-seasonal change in morphology became more moderate (Fig.

5B,C). In the 3-d culture, total leaf area increased under July and August conditions (Fig. **5B**;

Supplementary Fig. **S3D**). Petiole length increased under warmer conditions (June to September).

The number of leaves slightly increased under the July, August, and September conditions (Fig. **5B**;

Supplementary Fig. **S3F**). The leaf blade length slightly increased under the July condition (Fig.

5B; Supplementary Fig. **S3G**). In the 1-d culture, there was no significant difference in plant growth

(Fig. **5B**; Supplementary Fig. **S4**). Morphological observations showed that the differences among

the conditions of each month were smaller in the shorter culture period lengths.

Differences between culture periods (7, 3, and 1-d) at $P < 0.001$ using ANOVA and

Tukey's HSD test are shown in Fig. **5B**. Taking the number of tests into account, we chose a

conservative threshold (results of differences at $P < 0.05$ are shown in Supplementary Tables **S4–7**).

Significant differences in total leaf area were detected between 7- and 3-d cultures in March and

between 7- and both 3- and 1 d cultures in November. Significant differences in the number of

leaves were detected between 7- and both 3- and 1-d cultures in January and between 7- and 3-d

cultures in February and August, indicating that development of visible leaves tended to be

suppressed under these conditions in the 7-d culture. Significant differences in maximum petiole

length were detected between 7- and 1-d cultures from June to September, indicating that petiole

elongation tended to be promoted under these conditions as the culture period increased. A

significant difference in maximum petiole length was also detected between the 7- and 3-d cultures

in March. No significant difference was detected in the leaf blade length between the culture

periods.

Each seasonally oscillating gene required a different length of cultivation period to reproduce

its seasonal trend

RNA-Seq was performed to clarify how the length of the culture period affected the

pseudo-seasonal trend of gene expression in *Ath*. The SO genes and SR genes were defined in the

same manner as the analysis of 7-d culture data (Fig. **6**; Supplementary Fig. **S5** and **S6**). The

number of SO genes was similar in the 7-d culture (4016 genes) and 3-d culture (3975 genes) (Fig.

6A). The number of SO genes in the 1-d culture was greater than that in the 3- and 7-d cultures

(4575 genes). The number of SO genes that were annotated as orthologs of *Ahg* SO genes showed

the same tendency (7-d culture: 1757 genes, 3-d culture: 1766 genes, 1-d culture: 1886 genes). As

the culture period became shorter, the phase distribution became more concentrated in summer and

winter (Fig. **6B**). A total of 412 SR genes were shared with three conditions (Fig. **6C**). Examples of

the shared genes were *LHY* and S*UCROSE SYNTHASE 1* (*SUS1*, AT5G20830). These genes

showed almost the same seasonal trends regardless of the length of the culture period (Fig. **7A,B**).

Interestingly, hundreds of genes mimicked the *Ahg* seasonal trend only when the culture

period was longer or shorter (SR genes in 7-d culture but not in 1-d culture: 436 genes, SR genes in

1-d culture but not in 7-d culture: 508 genes). *ALCOHOL DEHYDROGENASE 1* (*ADH1*,

AT1G77120) and *JACALIN-RELATED LECTINS 5* (*JAL5*, AT1G52000) were classified as SR

genes in the 1-d culture but not in the 7-d culture (Fig. **6D, 7C,D**). As the culture period increased,

the amplitude of these genes decreased. *SWEET17* (AT4G15920) and *TRANSPARENT TESTA 8*

(*TT8*, AT4G09820) were classified as SR genes in the 7-d culture but not in the 1-d culture (Fig. **6C,**

7E,F). The amplitude of these genes decreased as the culture period became shorter. TT8 is a bHLH

transcription factor that regulates flavonoid biosynthesis with other factors (Li, 2014; Xu et al.,

2015). Anthocyanin regulatory and biosynthetic pathways and predicted seasonal expression plots

of *Ahg* and *Ath* are shown in Supplementary Fig. **S7–10**. Among these genes, the amplitudes of the

pseudo-seasonal trend of *DIHYDROFLAVONOL 4-REDUCTASE* (*DFR*, AT5G42800),

LEUCOANTHOCYANIDIN DIOXYGENASE (*LDOX*, AT5G42800), *PRODUCTION OF*

ANTHOCYANIN PIGMENT 1 (*PAP1*, AT1G56650), and *PAP2* (AT1G66390) became larger as the

culture period increased, similar to *TT8* (Supplementary Fig. **S7A,C, S8**, and **S10**). In our criteria,

F3'H, *MYB11*, *MYB111*, *PAP1*, *PAP2*, *TT8*, *GL3*, and *EGL3* were defined as SO genes in *Ahg*.

Among them, only *TT8* and *PAP2* were defined as SR genes in *Ath*.

The reason for the difference in the amplitude of the seasonal trend among the three

conditions in *Ath* could be the difference in the length of the culture period (7, 3, and 1 d) and/or the

difference in the plant age at the start of the treatment in the SGCmini (23, 27, and 29 d). To

investigate the effect of plant age on the amplitude of SO genes, plants at 23 d after seeding were

cultured in the SGCmini for 1 d under the January and August conditions (Supplementary Fig.

S11A), and RNA-Seq was performed. The expression levels of SO genes of younger plants tended

to have slightly smaller differences between the January and August conditions (Supplementary Fig.

S11B). However, the variation due to the difference in the length of the culture period was greater

than that due to plant age (Supplementary Fig. **S11B**–**D**).

In total, 1627 genes were identified as SR genes in at least one of the three conditions

(Fig. **6C**; Supplementary Fig. **S2B,** Table **S8**). To elucidate the functional aspects of the SR and

non-SR genes, we performed GO enrichment analysis (adjusted $P < 0.05$, Supplementary Tables

S9–11). In all SR genes, the GO terms "gene expression," "response to abiotic stimulus," and

"response to temperature stimulus" were enriched (Fig. **6E**). The GO term "response to abiotic

stimulus" was enriched only in the SR gene in 7-d culture (Fig. **S12,** Supplementary Tables **S12-14**).

In all non-SR genes, the GO terms "secondary metabolic process," "response to biotic stimulus,"

and "response to abiotic stimulus" were enriched. The GO terms "secondary metabolic process"

and "glucosinolate metabolic process" were enriched in genes that passed the amplitude criteria but

were not in phase with *Ahg*. This indicates that several SO genes associated with these GO terms

require other factors (other environmental stimuli, etc.) to mimic the seasonal trend of *Ahg* in the

appropriate phase, even though they were induced under mimic conditions in the present

experiment.

Discussion

Short-term incubation of *Ath* plants in the SGCmini under conditions representing each month could mimic the seasonal trends of 1627 SO genes in *Ahg*. The results of the 7-, 3-, and 1-d culture periods revealed that the culture period required for mimicry differed depending on the SO gene (Fig. **6C, 7;** Supplementary Fig. **S8**). For some SO genes, the length of the culture period seems to be unrelated to the mimicking of seasonal trends of *Ahg* gene expression in the field because 412 SR genes were observed under all three conditions. These SO genes may ensure robustness to environmental changes by responding accurately to the current environment, regardless of the past environment. *LHY* is one of these SO genes (Fig. **7A**); it is known as a core clock gene that contributes to temperature compensation of the circadian clock (Gil and Park, 2019; Gould et al., 2006; Salomé et al., 2010). In field environments, temperatures can change drastically, even within a few days. The constant responsiveness of *LHY* may contribute to maintaining the circadian rhythm under fluctuating environmental conditions of the field. In our previous study, we found that 1792 SO genes had both seasonal and diurnal oscillations (62.2% of SO genes)(Nagano et al., 2019). Recently, the complex effects of clock genes in the natural environment have received increasing attention (Oravec and Greenham, 2022) because rising temperatures due to global warming disrupt existing plant responses to day length and temperature during seasonal events. Accordingly, we have now established a convenient experimental system using SGCmini. We will be able to assess the contribution of each gene (such as clock genes) to the seasonal transcriptome from pseudo-seasonal experiments using mutants of each gene in *Ath*.

Conversely, there were several SO genes that mimicked the *Ahg* seasonal trend only when the culture period was shorter (Fig. **7C,D**). *ADH1* was one of these SO genes. Song *et al*. showed that *ADH1* transcription was induced at 3 h following 4 °C stress treatment, reached the highest expression level at 24 h, then declined (Song et al., 2017). Since its amplitude decreased under longer incubations, *ADH1* seems to respond to environmental changes (in this experiment, changes in temperature and daylength from pre-culture conditions to conditions of each month in

the SGCmini), rather than to absolute values at the time, as LHY does. *ADH1* is not only involved

in the cold stress response (Song et al., 2017), but its transcription is also known to be induced by

various other stresses (Shi et al., 2017; Winter et al., 2007). SO genes such as *ADH1*, which

mimicked field seasonal trends only in shorter culture periods, may play a role in responding to

rapid environmental changes (within 1 day) due to their transient expression.

Several SO genes mimicked field seasonal trends only in longer culture periods, such as

SWEET17 and *TT8* (Fig. **7E,F**). In *Ath*, late biosynthetic genes (LBGs) of the anthocyanin

biosynthetic pathway, *DFR* and *LDOX*, and their transcriptional regulators, *TT8*, *PAP1*, and *PAP2*,

showed an increase in the amplitude of the seasonal trend with the extension of the culture period,

suggesting a regulatory mechanism that responds to continuous environmental stimuli such as low

temperature (Supplementary Fig. **S7–10**). One possible mechanism driving this is epigenetic

regulation. Recent studies have revealed that epigenetic regulation is involved in the regulation of

anthocyanin biosynthesis gene expression (Cai et al., 2019; Zheng et al., 2019). In the field, analysis

of seasonal histone modifications of *Ahg* suggests that histone H3 lysine 27 trimethylation is

involved in gene expressions that depend on memory of longer-term (more than a few weeks)

environmental conditions (Nishio et al., 2020). In our study, although the plants only experienced

the treatment once, the memory of past intermittent stimuli, such as hot or cold temperatures, can

also affect plant seasonal response (Leuendorf et al., 2020; Yamaguchi et al., 2021). Evaluating the

contribution of longer-term (month-scale) and intermittent environmental stimulus inputs to

seasonal trends is one of the next challenges in the study of SO genes.

The differences in the temporal responsiveness of SO genes may contribute to the

variation in the phase of the seasonal trend (Fig. **6B**). In addition, integrated response to multiple

stimulus information may also be responsible for the wide distribution of phases in field *Ahg*. The

phase of pseudo-seasonal trends of *PAP1* in *Ath* did not match that in *Ahg* (Fig. **S7C**). In SGCmini,

PAP1 expression is up-regulated in winter conditions, but in the field, its expression is higher in

summer than in winter. Although previous studies have shown that *PAP1* and *PAP2* are

up-regulated at low temperatures (Petridis et al., 2016), *PAP2*, rather than *PAP1* is suggested to

play a more specific role in the accumulation of flavonoids at cold temperatures (Hannah et al.,

2006). *PAP1* expression is also induced by other stimuli, such as high light intensity (Li et al., 2016;

Shi and Xie, 2010). An integrated response to multiple stimulus information may also be the reason

for the enrichment of GO terms like "response to stress" or "response to abiotic stimuli" in non-SR

genes (Fig.**6E**). For some SO genes, multiple stimulus inputs in the field may form seasonal trends

that are not correlated with a single environmental stimulus, such as temperature.

Alternatively, multiple seasonal stimuli may mask seasonal oscillations. Early

biosynthetic genes (EBGs) and LBGs, except *F3'H*, did not show seasonal oscillations in *Ahg* in the

field (Supplementary Fig. **S7** and **S8**). However, their transcriptional regulators showed seasonal

oscillations in various phases (Supplementary Fig. **S7, S9,** and **S10**), suggesting that the constitutive

expression of EBGs and LBGs in the field is the outcome of the integration of multiple

environmental stimuli by transcriptional regulators. Flavonoid biosynthesis is affected not only by

low temperature, but also by light quality, drought, nutrition, and virus infection (Cai et al., 2019;

Cominelli et al., 2008; Honjo et al., 2020; Lillo et al., 2008). While highly oscillating genes are

eye-catching, it is also important to remember that seemingly non-oscillating genes may be

involved in plant responses to seasonal changes. Such a complex response mechanism of secondary

metabolites may be one of the reasons why the GO term "secondary metabolic process" was

detected in the GO analysis of SO genes that could not be mimicked (Fig. **6E**).

In conclusion, we successfully mimicked the seasonal trends of approximately 38% of

Ahg SO genes using *Ath* plants by incubating them for 1 to 7 days in an experimental system set at

the average temperature and daylength of each month. The results indicate that the field seasonal

trend of these genes is shaped by day-scale responses to environmental stimuli rather than

month-scale responses to long-term inputs. Conversely, the seasonal trends of 62% of the SO genes

could not be mimicked. Although one of the reasons could be the lack of long-term inputs of

environmental stimuli, there are many other possibilities. For example, species differences between

Ahg and *Ath* and environmental stimuli which were not replicated in the SGCmini (fluctuation of

environmental stimuli, biotic stress, etc.) may also be responsible. We will next clarify which of

these possibilities is important for seasonal trends of non-mimicked SO genes in *Ahg*. This is the

first comprehensive study of the day-scale response of SO genes to environmental stimuli. In order

to adapt to complex environmental changes in the field, plants likely interpret seasonal changes by

converting environmental stimuli into gene expression at different temporal resolutions (minute-,

day- and month-scale). Although the functions of many SO genes are still unknown, the results of

this study and the experimental system can be useful for investigating these functions.

Materials and Methods

Smart growth chamber mini (SGCmini)

The system configuration of the developed smart growth chamber mini (SGCmini) is shown in Supplementary Fig. **S1A**. The proposed system consisted of an incubator body, Peltier unit, circuit unit, and control unit. Supplementary Fig. **S1B** depicts the proposed system, excluding the control unit. The incubator body was made of acrylic; the bottom surface was made of black acrylic and the rest was transparent acrylic (thickness: 10 mm). The transparent acrylic parts were all integrated and had holes into which the Peltier unit could be attached (described below). The outer dimensions of the transparent acrylic part were 80 mm × 200 mm × 200 mm.

The appearance of the Peltier unit is shown in Supplementary Fig. **S1C**. This unit comprised a 40 × 40 mm Peltier device, two heat sinks, and two fans. The heat sinks were adhered on both sides of the Peltier device and the black insulation sheets enclosed them. The side with the small heat sink was inserted through the hole in the incubator body and housed in the body. A small fan was installed to circulate the air inside the incubator. The side with the larger heat sink was for heat dissipation, which enabled air cooling with a large fan.

The circuit unit consisted of an Arduino Uno R3, a DC brush motor driver shield for Arduino (SHIELD-MD10, Cytron Technologies, Malaysia), and two temperature sensors (MCP9808, Microchip Technology, USA). Two temperature sensors were installed in the incubator. The circuit unit sends information from the temperature sensors to the control unit (PC) via serial communication.

The control unit was equipped with a GUI, which determined the extent of temperature control for heating and cooling to achieve the "target temperature set by the GUI" in the incubator. The proportional integral (PI) control determined the temperature control according to the current temperature in the incubator, and it was sent to the circuit unit as the control signal. According to the signal received, the circuit unit controlled the current sent to the Peltier unit by pulse-width modulation (PWM).

The GUI of the control unit is illustrated in Supplementary Fig. **S1D**. In this GUI, the

temperature in the incubator could be set in increments of 0.5 °C. In addition to the current

temperature, the temperature profile for the past 48 h could be displayed on a chart. The GUI also

provided other functions, such as the output of a temperature control log and the capability to

change the set temperature. The developed incubator could control the temperature with an error

range of 0.2 °C within a range of −9 to +20 °C from the standard room temperature. For colder

conditions, such as that for February, the SGCmini was operated in a growth chamber (LH-241SP,

NK system Osaka, Japan) set below 15 °C.

For plant cultivation, the incubator body was painted black on four sides to block external

light (Fig. **1A**). An LED light unit (25 pcs; color: red and blue; size: 30.8 × 30.8 cm; PPFD:

approximately 35 µmol m^{-2} s^{-1}) was placed on the top of the incubator body. The lighting of the

LED light unit was controlled using a timer.

Setting of culture condition in SGCmini

To mimic the seasonal expression of *Arabidopsis halleri* ssp. *gemmifera* (*Ahg*) in the field,

simplified conditions of the environment of the *Ahg* sampling site (the Omoide River, Taka-cho,

35° 06′ N, 134° 55′ E, altitude 190–230 m above sea level) (Nagano et al., 2019) were set as culture

conditions (Fig. **2A**). For daylength settings, sunrise and sunset times on the 15th of each month in

Nishiwaki city were obtained from the National Astronomical Observatory of Japan. For

temperature settings, hourly average temperatures at Nishiwaki (averaged over 2006–2016) were

obtained from the Japan Meteorological Agency. The average value of the temperature from sunrise

to sunset was set as the daytime temperature, and the average value of the temperature from sunset

to sunrise was set as the nighttime temperature. It was difficult to mimic temperatures below 5 °C,

so when the temperature was below 5 °C, such as in January, the experiment was conducted with a

5 °C setting.

We also conducted a high-temperature culture experiment, in which the daytime

temperature of the August condition was changed to 32 °C or 35 °C (Fig. **4A**).

Plant materials

Seeds of *A. thaliana* (*Ath*) (Col-0, CS70000) were sown on 1/2 MS medium with 0.5%

sucrose and 0.6% gellan gum in square Petri dishes and cultured at 20 °C with 8 h light for 20 d.

Plants were transferred to 1/2 MS medium with 0.25% gellan gum in round plates and cultivated for

3, 7, or 9 d under the same conditions as the pre-culture to recover from the stress of transplanting.

After the pre-culture, the round plates were placed in the SGCmini. Plants were cultivated under the

conditions of each month for 7, 3, or 1 d (Fig. **2B** and **5B**). To equalize the age of the plants at the

sampling date, we extended the pre-culture period to shorten the culture period. However, for a

1-day culture with short pre-culture conditions, the pre-culture period was set to 3 d in order to

evaluate the effect of plant age at the start date of treatment (Supplementary Fig. **S12A**).

For morphometry, all leaves were cut off and scanned using a scanner (GT-S650, EPSON,

Japan). Leaf area and leaf length were measured from the scanned images using ImageJ software.

ANOVA and Tukey HSD were performed in R software to test for differences between conditions

($P < 0.05$).

RNA-Seq

For RNA extraction, leaves larger than 1 mm without cotyledons and juvenile leaves

were harvested from one plant as one sample between 11:00 and 13:00. The leaves were

immediately frozen in liquid nitrogen and stored at −80 °C until extraction.

Samples were ground with two zirconia beads using a TissueLyser II (QIAGEN, MD,

USA) with pre-chilled adapters at −80 °C. Total RNA from leaves was extracted using a Maxwell

16 LEV Plant RNA Kit (Promega, WI, USA). The amount of RNA was determined using Quant-iT

RNA Assay Kit, broad range (Thermo Fisher Scientific, Waltham, MA, USA) and Tecan plate

reader Infinite 200 PRO (Tecan, Männedorf, Switzerland). For RNA-Seq library preparation, 400 or

500 ng of total RNA per sample was used. The library was prepared using the Lasy-Seq v1.0 or

v1.1 protocol (Kamitani et al., 2019) (https://sites.google.com/view/lasy-seq/). The quality of the

library was assessed using a Bioanalyzer (Agilent Technologies). The libraries were sequenced with

a HiSeq 2500 or Hiseq X (Illumina) to produce single reads of 50 bp (Library 1 and 2) or 150bp

(Library 3).

440   Preprocessing and quality filtering of RNA-Seq data were performed using

441   Trimmomatic-0.33 (Bolger et al., 2014). Preprocessed reads were mapped on transcript sequences

442   in TAIR 10 with Bowtie1(v1.1.1) (Langmead et al., 2009) and quantified using RSEM-1.2.21(Li

443   and Dewey, 2011). The output of the RSEM was analyzed using R (R Core Team, 2019).

444   RNA-Seq data of *A. halleri* were obtained from previous studies (Nagano et al., 2019).

RNA-Seq data of *A. thaliana* and *A. halleri* were analyzed as follows. Samples with fewer than

$10^{5.5}$ reads were excluded from the analysis. The attributes of the samples used in the analysis are

listed in Supplementary Table **S1**. *Ahg* genes with an average $\log_2(\text{rpm}+1) > 2$ were designated as

expressed genes (16158 genes) (Supplementary Fig. **S2**). A total of 90.3% of the expressed genes in

Ahg (14587 genes) were annotated as orthologs of *Ath* genes by reciprocal BLAST. Subsequent

seasonal expression analysis was performed using these annotated *Ahg*-expressed genes (14587

genes) and their orthologous genes in *Ath*.

Analysis of seasonal expression

454   To compare the seasonal trends of *Ahg* and *Ath*, the expressions of the *Ahg*-expressed

genes and their orthologous genes in *Ath* were fitted to a cosine curve using the *nls* function in R

(Fig. **3A**). Amplitude and phase were obtained from each fitted cosine curve over a 1-y period.

Genes with an amplitude > 1 were defined as seasonally oscillating genes (SO genes). Since the

value of $\log_2(\text{rpm} + 1)$ was fitted with a cosine curve, an amplitude > 1 indicates that the difference

in seasonal oscillation is more than two-fold. We compared the amplitude and phase of *Ahg* and *Ath*

genes to determine "seasonality reproduced genes (SR genes)" that could mimic the seasonal

expression trends of *Ahg* in the field. A gene whose amplitude difference was smaller than 1 and

whose phase difference was smaller than 45 d was defined as a SR gene.

Gene ontology enrichment analysis

For gene ontology (GO) enrichment analysis, GO annotations for *Ath* genes were

obtained from the TAIR database. Statistical tests of enrichment analysis were performed using the

Fisher's exact test function in R. A total of 14,472 expressed genes with GO annotations were

included in the test. Multiple testing corrections were performed using FDR (Benjamini and

Hochberg, 1995). Enriched GO terms are those with adjusted $p < 0.05$. Representative GO terms

were summarized based on the REVIGO output (http://revigo.irb.hr) (Supek et al., 2011) (allowed

similarity = 0.5) and visualized with reference to a previous protocol (Bonnot et al., 2019).

Data Availability Statement

The RNA-seq data were submitted to the NCBI Sequence Read Archive repository under

the BioProject number PRJNA739047 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA739047).

R scripts and data required for analysis (count data, etc.) are available via the GitHub repository

(https://github.com/naganolab/AthSGCmini_pseudo-seasonal_RNA-Seq).

Funding information

This work was supported by Japan Science and Technology Agency CREST

[JPMJCR15O2], ACCEL [JPMJAC1403], FOREST [JPMJFR210B] and a project [JPNP18016] of

the New Energy and Industrial Technology Development Organization (NEDO).

Acknowledgements

We greatly appreciate Hiroshi Kudoh and his laboratory members for field data

acquisition and discussions, Tetsuro Mimura for his advice and discussions on the experimental

system, Kyoko Mogami for her help with RNA-Seq library preparation, and Motohiro Mihara and

Hitoshi Ooshima of Dynacom Co., Ltd. (Chiba, Japan) for their support in RNA-Seq analysis,

Daisuke Kyogoku for his advice on English language editing.

Author Contribution

A.J.N. and Y.K. designed the research. H.T. and T.T. developed SGCmini. M. Kamitani,

Y.H., M. Kashima, A.T. contributed to library preparation and data analysis. Y.K. performed

experiments and data analysis. Y.K., H.T. and A.J.N. wrote the paper.

Reference

Andersson, A., Keskitalo, J., Sjödin, A., Bhalerao, R., Sterky, F., Wissel, K., et al. (2004) A

transcriptional timetable of autumn senescence. *Genome Biol*. 5: R24.

Benjamini, Y., and Hochberg, Y. (1995) Controlling the False Discovery Rate□: A Practical and

Powerful Approach to Multiple Testing. *J R Stat Soc Ser B*. 57: 289–300.

Bolger, A.M., Lohse, M., and Usadel, B. (2014) Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics*. 30: 2114–2120.

Bonnot, T., Gillard, M., and Nagel, D. (2019) A Simple Protocol for Informative Visualization of

Enriched Gene Ontology Terms. *Bio-Protocol*. e3429.

Bouché, F., Woods, D.P., and Amasino, R.M. (2017) Winter memory throughout the plant kingdom:

Different paths to flowering. *Plant Physiol*. 173: 27–35.

Cai, H., Zhang, M., Chai, M., He, Q., Huang, X., Zhao, L., et al. (2019) Epigenetic regulation of

anthocyanin biosynthesis by an antagonistic interaction between H2A.Z and H3K4me3. *New*

Phytol. 221: 295–308.

Cominelli, E., Gusmaroli, G., Allegra, D., Galbiati, M., Wade, H.K., Jenkins, G.I., et al. (2008)

Expression analysis of anthocyanin regulatory genes in response to different light qualities in

Arabidopsis thaliana. *J Plant Physiol*. 165: 886–894.

Falavigna, V. da S., Guitton, B., Costes, E., and Andrés, F. (2019) I want to (Bud) break free: The

potential role of DAM and SVP-like genes in regulating dormancy cycle in temperate fruit trees.

Front Plant Sci. 9: 1990.

Galindo González, L.M., El Kayal, W., Ju, C.J.T., Allen, C.C.G., King-Jones, S., and Cooke, J.E.K.

(2012) Integrated transcriptomic and proteomic profiling of white spruce stems during the

transition from active growth to dormancy. *Plant, Cell Environ*. 35: 682–701.

Gil, K.E., and Park, C.M. (2019) Thermal adaptation and plasticity of the plant circadian clock. *New*

Phytol. 221: 1215–1229.

Gong, Z., Xiong, L., Shi, H., Yang, S., Herrera-Estrella, L.R., Xu, G., et al. (2020) Plant abiotic stress

response and nutrient use efficiency. *Sci China Life Sci*. 63: 635–674.

Gould, P.D., Locke, J.C.W., Larue, C., Southern, M.M., Davis, S.J., Hanano, S., et al. (2006) The

molecular basis of temperature compensation in the Arabidopsis circadian clock. *Plant Cell*. 18:

1177–1187.

Hannah, M.A., Wiese, D., Freund, S., Fiehn, O., Heyer, A.G., and Hincha, D.K. (2006) Natural

genetic variation of freezing tolerance in arabidopsis. *Plant Physiol*. 142: 98–112.

Hashida, Y., Tezuka, A., Nomura, Y., Kamitani, M., Kashima, M., Kurita, Y., et al. (2022) Fillable

and unfillable gaps in plant transcriptome under field and controlled environments. *Plant Cell*

Environ. 45: 2410–2427.

Honjo, M.N., Emura, N., Kawagoe, T., Sugisaka, J., Kamitani, M., Nagano, A.J., et al. (2020)

Seasonality of interactions between a plant virus and its host during persistent infection in a

natural environment. *ISME J*. 14: 506–518.

Jean Finnegan, E. (2015) Time-dependent stabilization of the +1 nucleosome is an early step in the

transition to stable cold-induced repression of FLC. *Plant J*. 84: 875–885.

Kamitani, M., Kashima, M., Tezuka, A., and Nagano, A.J. (2019) Lasy-Seq: a high-throughput

library preparation method for RNA-Seq and its application in the analysis of plant responses to

fluctuating temperatures. *Sci Rep*. 9: 1–14.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009) Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biol*. 10: R25.

Leuendorf, J.E., Frank, M., and Schmülling, T. (2020) Acclimation, priming and memory in the

response of Arabidopsis thaliana seedlings to cold stress. *Sci Rep*. 10: 1–11.

Li, B., and Dewey, C.N. (2011) RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC Bioinformatics*. 12: 323.

Li, S. (2014) Transcriptional control of flavonoid biosynthesis: Fine-tuning of the

MYB-bHLH-WD40 (MBW) complex. *Plant Signal Behav*. 9: e27522.

Li, S., Wang, W., Gao, J., Yin, K., Wang, R., Wang, C., et al. (2016) MYB75 phosphorylation by

MPK4 is required for light-induced anthocyanin accumulation in arabidopsis. *Plant Cell*. 28:

2866–2883.

Lillo, C., Lea, U.S., and Ruoff, P. (2008) Nutrient depletion as a key factor for manipulating gene

expression and product formation in different branches of the flavonoid pathway. *Plant, Cell*

Environ. 31: 587–601.

Lu, H., Gordon, M.I., Amarasinghe, V., and Strauss, S.H. (2020) Extensive transcriptome changes

during seasonal leaf senescence in field-grown black cottonwood (Populus trichocarpa

Nisqually-1). *Sci Rep*. 10: 1–14.

Luo, X., and He, Y. (2020) Experiencing winter for spring flowering: A molecular epigenetic

perspective on vernalization. *J Integr Plant Biol*. 62: 104–117.

Maurya, J.P., and Bhalerao, R.P. (2017) Photoperiod- and temperature-mediated control of growth

cessation and dormancy in trees: A molecular perspective. *Ann Bot*. 120: 351–360.

Nagano, A.J., Kawagoe, T., Sugisaka, J., Honjo, M.N., Iwayama, K., and Kudoh, H. (2019) Annual

transcriptome dynamics in natural environments reveals plant seasonal adaptation. *Nat Plants*. 5:

74–83.

Nagano, A.J., Sato, Y., Mihara, M., Antonio, B.A., Motoyama, R., Itoh, H., et al. (2012) Deciphering

and prediction of transcriptome dynamics under fluctuating field conditions. *Cell*. 151:

1358–1369.

Nishio, H., Nagano, A.J., Ito, T., Suzuki, Y., and Kudoh, H. (2020) Seasonal plasticity and diel

stability of H3K27me3 in natural fluctuating environments. *Nat Plants*. 6: 1091–1097.

Oravec, M.W., and Greenham, K. (2022) The adaptive nature of the plant circadian clock in natural

environments. *Plant Physiol*. 968–980.

Petridis, A., Döll, S., Nichelmann, L., Bilger, W., and Mock, H.P. (2016) Arabidopsis thaliana

G2-LIKE FLAVONOID REGULATOR and BRASSINOSTEROID ENHANCED

EXPRESSION1 are low-temperature regulators of flavonoid accumulation. *New Phytol*. 211:

577  912–925.

578 R Core Team (2019) R: A language and environment for statistical computing. R Foundation for

579  Statistical Computing, Vienna, Austria. .

580 Roy, J., Rineau, F., De Boeck, H.J., Nijs, I., Pütz, T., Abiven, S., et al. (2021) Ecotrons: Powerful and

581  versatile ecosystem analysers for ecology, agronomy and environmental science. *Glob Chang*

582  *Biol*. 27: 1387–1407.

583 Salomé, P.A., Weigel, D., and McClunga, C.R. (2010) The role of the Arabidopsis morning loop

584  components CCA1, LHY, PRR7, and PRR9 in temperature compensation. *Plant Cell*. 22:

585  3650–3661.

586 Shi, H., Liu, W., Yao, Y., Wei, Y., and Chan, Z. (2017) Alcohol dehydrogenase 1 (ADH1) confers

587  both abiotic and biotic stress resistance in Arabidopsis. *Plant Sci*. 262: 24–31.

588 Shi, M.Z., and Xie, D.Y. (2010) Features of anthocyanin biosynthesis in pap1-D and wild-type

589  Arabidopsis thaliana plants grown in different light intensity and culture media conditions.

590  *Planta*. 231: 1385–1400.

591 Song, Y., Liu, L., Wei, Y., Li, G., Yue, X., and An, L. (2017) Metabolite profiling of adh1 mutant

592  response to cold stress in Arabidopsis. *Front Plant Sci*. 7: 2072.

593 Song, Y.H., Ito, S., and Imaizumi, T. (2013) Flowering time regulation: Photoperiod- and

594  temperature-sensing in leaves. *Trends Plant Sci*. 18: 575–583.

595 Supek, F., Bošnjak, M., Škunca, N., and Šmuc, T. (2011) Revigo summarizes and visualizes long lists

596  of gene ontology terms. *PLoS One*. 6: e21800.

597 Winter, D., Vinegar, B., Nahal, H., Ammar, R., Wilson, G. V., and Provart, N.J. (2007) An 'electronic

598  fluorescent pictograph' Browser for exploring and analyzing large-scale biological data sets.

599  *PLoS One*. 2: e718.

600 Xu, W., Dubos, C., and Lepiniec, L. (2015) Transcriptional control of flavonoid biosynthesis by

601  MYB-bHLH-WDR complexes. *Trends Plant Sci*. 20: 176–185.

602 Yamaguchi, N., Matsubara, S., Yoshimizu, K., Seki, M., Hamada, K., Kamitani, M., et al. (2021)

H3K27me3 demethylases alter HSP22 and HSP17.6C expression in response to recurring heat in

Arabidopsis. *Nat Commun*. 12: 1–16.

Ye, K., Li, H., Ding, Y., Shi, Y., Song, C., Gong, Z., et al. (2019)

BRASSINOSTEROID-INSENSITIVE2 Negatively Regulates the Stability of Transcription

Factor ICE1 in Response to Cold Stress in Arabidopsis. *Plant Cell*. 31: 2682–2696.

Zheng, T., Tan, W., Yang, H., Zhang, L., Li, T., Liu, B., et al. (2019) Regulation of anthocyanin

accumulation via MYB75/HAT1/TPL-mediated transcriptional repression. *PLoS Genet*. 15:

e1007993.

Fig. 1 Performance of the smart growth chamber mini (SGCmini). (A) Picture of the SGCmini. The

walls of the incubator body were painted for shading. (B) Temperature plot of a test run. The

temperature was changed stepwise from 16 °C to 26 °C every 2 h. Light blue line: outside temperature;

pink lines: inside temperature measured by the sensors; black line: preset temperature; red line: average

temperature of sensor 1 and sensor 2. (C) Enlarged plot of (B) at the lowest temperature setting. (D)

Enlarged plot of (B) at the highest temperature setting. (E) Temperature plot for 3 d under the March

condition. (F) Temperature plot for 3 d under the August condition.

Fig. 2 Experimental settings and morphological traits of *Arabidopsis thaliana* (*Ath*) plants grown in

the SGCmini for 7 d. (A) Setting of each condition in the SGCmini. Yellow bars: daylength; red circles:

daytime temperatures; blue triangles; nighttime temperatures. (B) Schematic diagram of the culture

condition. (C) Whole images of plants grown under February, May, and July conditions. (D) Leaves of

plants grown under February, May, and July conditions. The youngest leaves are on the right. Bar = 10

mm. (E) Total leaf area. (F) Number of leaves. (G) Maximum petiole length. (H) Maximum leaf blade

length. Bar: SD. n = 5–9. Different letters indicate differences between conditions at $P < 0.05$ using

ANOVA and Tukey's HSD test.

Fig. 3 Comparison of seasonal trends of gene expression in *Arabidopsis halleri* ssp. *gemmifera*

(*Ahg*) in the field and *Arabidopsis thaliana* (*Ath*) in the SGCmini for 7 d. (A) Schematic diagram of

comparison method of seasonal trends of *Ahg* and *Ath*. Black circles indicate observed value by

RNA-Seq. The solid orange (*Ahg*) and blue (*Ath*) lines show the fitted cosine curve. The orange and

blue dashed lines indicate the phases of *Ahg* and *Ath*. "α" and "φ" indicate the amplitude and the

phase of the fitted cosine curve. (B) Histograms of amplitude (α) of expressed genes. Left: *Ahg*.

Right: *Ath*. The regions with $\alpha > 1$ are colored. (C) Histograms of phase (φ) of expressed genes.

Left: *Ahg*. Right: *Ath*. The regions with $\alpha > 1$ are colored. (D) Venn diagram of *Ahg* and *Ath* SO

genes. Left circle: *Ahg*. Right circle: *Ath*. (E) Scatter plot of amplitudes in *Ahg* and *Ath*. Only genes

with $\alpha > 1$ in both species are colored according to density using the *densCols* function in R. (F)

Scatter plot of phases in *Ahg* and *Ath*. Only genes with $\alpha > 1$ in both species are colored according

to density using the *densCols* function in R. (G) Venn diagram of genes that passed amplitude and

phase criteria. Left circle: genes that passed amplitude criteria ($\Delta\alpha < \pm 1$). Right circle: genes that

passed phase criteria ($\Delta\varphi < \pm 45$ days). (H) Seasonal expression plots of *LHY*. From left, fitted

cosine curve plot of *Ahg* and *Ath*, observed value and fitted line plot of *Ahg* and *Ath*, respectively.

Fig. 4 Effect of higher temperature on gene expression. (A) Schematic diagram of culture

conditions. (B) Conditions of the high temperature experiment. Yellow bars: daylength; red circles:

daytime temperatures; blue triangles: nighttime temperatures. (C) Gene expression plot of *TIP1;1*.

From left, fitted cosine curve plot of *Arabidopsis halleri* ssp. *gemmifera* (*Ahg*) and *Arabidopsis*

thaliana (*Ath*), observed value and fitted line plot of *Ahg* and *Ath*, respectively, and a plot of

temperature and expression levels. Black circles indicate observed values. Orange (*Ahg*) and blue

(*Ath*) lines show the fitted cosine curve. Orange and blue dashed lines indicate the phases of *Ahg*

and *Ath*, respectively. Pink triangles and red squares indicate observed values when plants were

grown at 32 °C and 35 °C. (D) *VI1*. (E) *HSP18.2*. (F) *HSP22.0*.

Fig. 5 Comparison of morphological traits of *Arabidopsis thaliana* (*Ath*) plants grown in the

SGCmini for 1, 3, or 7 d. (A) Schematic diagram of culture conditions. (B) Morphological traits.

From left, total leaf area, number of leaves, maximum petiole length, and maximum leaf blade

length. Dark blue squares, middle blue triangles, and light blue circles indicate data for plants

grown in the SGCmini for 7, 3, and 1 d, respectively. n = 4–9. Different letters below the plots

indicate differences between culture periods at $P < 0.001$ using ANOVA and Tukey's HSD test. (C)

Leaves of plants grown under the August condition. From the top, 7-, 3- and 1-d cultures. The

youngest leaves are on the right. Bar = 10 mm.

Fig. 6 Comparison of seasonal trends of gene expression between different culture periods. (A) Comparison of the number of seasonally oscillating (SO) genes. Darker regions indicate the number of *Arabidopsis thaliana* (*Ath*) SO genes common with those of *Arabidopsis halleri* ssp. *gemmifera* (*Ahg*) defined in the field experiment data. The x-axis indicates the number of days of the length of the incubation period in SGCmini. (B) Comparison of the distribution of phases of SO genes. Orange circles: *Ahg*; dark blue rhombuses: *Ath* 7-d culture; blue squares: *Ath* 3-d culture; light blue triangles: *Ath* 1-d culture. (C) Venn diagram of seasonality reproduced genes (SR genes). Left circle: *Ath* 7-d culture; right circle: *Ath* 3-d culture; bottom circle: *Ath* 1-d culture. (D) Scatter plot of the alpha of SR genes in the 7-d culture (vertical axis) and 1-d culture (horizontal axis). Each point is colored based on the local density of points. Pink circles indicate genes shown in Fig. **7**. (E) Enriched biological process gene ontology (GO) terms. The "total SR gene" indicates the total SR number of genes that mimicked the seasonal trend of *Ahg* in at least one of the three conditions (7-, 3-, and 1-d culture) and corresponds to the colored region in Fig. 6C. The "total non-SR gene" indicates non-SR genes that did not mimic the seasonal trend of *Ahg* and corresponds to the non-colored region in Fig. 6C. Among the "total non-SR genes," "non-SR gene (phase)" indicates genes that passed the amplitude criteria but were not in phase with *Ahg*. Circle size indicates the number of genes. Color scale indicates the adjusted *P* values in the GO enrichment analysis.

Fig. 7 Seasonal expression plots of seasonality reproduced genes. From left, fitted cosine curve plot of *Arabidopsis halleri* ssp. *gemmifera* (*Ahg*) and *Arabidopsis thaliana* (*Ath*), observed value and fitted line plot of *Ahg* and *Ath* in the 7-, 3-, and 1-d cultures, respectively. (A) *LHY*. (B) *SUS1*. (C) *ADH1*. (D) *JAL5*. (E) *SWEET17*. (F) *TT8*.

A

LED light unit

Temperature sensor

Circuit unit

Incubator body

Peltier unit

A

Seeding | Trans planting | Start of treatment | Sampling

20 days | Pre-culture 3 days | Culture in SGCmini 7 days

20°C, 8h light | Aug

High temperature, 32°C

High temperature, 35°C

B

C AT2G36830.1, TIP1;1, gamma tonoplast intrinsic protein

Fitted curve plot | Ahg482663 $\alpha = 4.3$, $\varphi = 193$ | AT2G36830.1 (Ath d7) $\alpha = 2.11$, $\varphi = 204$

D AT1G62660.3, VI1, Glycosyl hydrolases family 32 protein

Fitted curve plot | Ahg907175 $\alpha = 5.37$, $\varphi = 205$ | AT1G62660.3 (Ath d7) $\alpha = 3.71$, $\varphi = 217$

E AT5G59720.1, HSP18.2, heat shock protein 18.2

Fitted curve plot | Ahg496104 $\alpha = 2.72$, $\varphi = 203$ | AT5G59720.1 (Ath d7) $\alpha = 1.02$, $\varphi = 192$

F AT4G10250.1, HSP22.0, HSP20-like chaperones superfamily protein

Fitted curve plot | Ahg489869 $\alpha = 2.61$, $\varphi = 208$ | AT4G10250.1 (Ath d7) $\alpha = 0.07$, $\varphi = 347$

A

B

C