Biomorphogenic feedbacks and the spatial organisation of a dominant grass steer dune
development

Dries Bonte[1*], Femke Batsleer[1], Sam Provoost[3], Valérie Reijers[5], Martijn L. Vandegehuchte[1,6], Ruben Van De
Walle[1], Sebastian Dan[2], Hans Matheve[1], Pieter Rauwoens[4], Glenn Strypsteen[4], Tomohiro Suzuki [2], Toon
Verwaest[2], Jasmijn Hillaert[1,3]

1. Ghent University, Dept. Biology, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium
2. Flanders Hydraulics Research, Berchemlei 115, 2140 Antwerp, Belgium
3. Institute for Nature and Forest research, Herman Teirlinckgebouw, Havenlaan 88 bus 73, 1000 Brussel
4. KU Leuven, Campus Brugge, Department of Civil Engineering, Spoorwegstraat 12
8200 Brugge
5. Department of Physical Geography, Faculty of Geosciences, Utrecht University, Utrecht, 3508 TC the
Netherlands
6. Department of Biology, Norwegian University of Science and Technology, Høgskoleringen 5, 7491
Trondheim, Norway

*Corresponding author: Dries Bonte (dries.bonte@ugent.be)

Abstract

Nature-based solutions to mitigate the impact of future climate change depend on restoring biological
diversity and natural processes. Coastal foredunes represent the most important natural flood barriers
along coastlines worldwide, but their area has been squeezed dramatically because of a continuing
urbanisation of coastlines, especially in Europe. Dune development is steered by the development of
vegetation in interaction with sand fluxes from the beach. Marram grass (*Calamagrostis arenaria*,
formerly *Ammophila arenaria*) is the main dune building species along most European coasts, but also
in other continents where the species was introduced. Engineering of coastal dunes, for instance by
building dunes in front of dikes, needs to be based on a solid understanding of the species' interactions
with the environment. Only quantitative approaches enable the further development of mechanistic
models and coastal management strategies that encapsulate these biomorphogenic interactions. We
here provide a quantitative review of the main biotic and physical interactions that affect marram grass
performance, their interactions with sand fluxes and how they eventually shape dune development.
Our review highlights that the species' spatial organisation is central to dune development. We further
demonstrate this importance by means of remote sensing and a mechanistic model and provide an
outlook for further research on the use of coastal dunes as a nature-based solution for coastal
protection.

Keywords: Nature-based solution, dune, coastal safety, ecological feedbacks, cover, spatial configuration,
synthesis, model, remote sensing

I. Introduction

As climate change induces sea level rise and possibly heavier storms, coastal protection is in a transition phase from hard structural engineering towards soft measures that can adapt dynamically to a changing environment (Borsje et al., 2011; IPCC, 2014, 2018; Vousdoukas et al., 2018). Ecosystem-based approaches complementing engineering with functional parts of the natural system provide such an alternative to conventional coastal defense ('hard engineering'). Indeed, estuarine and coastal soft sediment systems are dynamic by nature and their inherent ecological processes may be exploited to enhance resilience (Temmerman et al., 2013). Coastal foredunes represent the most important natural flood barrier for much of the European coastline and 30% of all shorelines worldwide (Martinez and Psuty, 2004; Reijers et al., 2019). In contrast to urban and other infrastructure, coastal dunes have the capacity to grow with rising sea level due to interactions between plant growth (Duarte et al., 2013) and aeolian sediment supply (de Vries et al., 2012; Strypsteen et al., 2019a). Therefore, they are currently considered as an important nature-based solution for coastal protection (Borsje et al., 2011; Duarte et al., 2013; Temmerman et al., 2013).

The use of foredunes as an engineering tool cannot be achieved without a deep understanding of the organizational properties of the natural dune system. Coastal dunes develop in first instance by sand accretion at the upper beach. In regions with predominant onshore winds, the magnitude of aeolian sand flux can primarily be described as a function of wind speed and grain size, but it also depends on soil moisture content, fetch length, beach and dune morphology (Delgado-Fernandez 2010, de Vries et al. 2012, Strypsteen et al. 2019). These aeolian fluxes impact the performance of a keystone species in foredunes from the European Atlantic coast : marram grass (*Calamagrostis arenaria (L.) Roth*, formerly *Ammophila arenaria;* Huiskes 1977). The species is the dominant species from white dunes, as protected within the directive 92/43 EEC [Shifting dunes along the shoreline with *Ammophila arenaria*, code 2120] (Perrino et al. 2014, European Commission 2007).

Biomorphogenesis refers to the process where biota like plants but also animals induce changes in the form of the environment they live in. Marram grass is such as an engineering species (Bakker, 1976; Puijenbroek et al., 2017) as its growth and performance depends on, and in turn influences, aeolian sand fluxes and hence, dune development (Hesp, 2002; Zarnetske et al., 2015; Strypsteen et al., 2019). Phenomena where the value of one state variable directly or indirectly affects the sign, direction and rate at which that variable changes, is defined as a feedback (Maxwell et al., 2017). These feedbacks can be positive (self-amplifying) or negative (self-dampening). As soon as sand dynamics cease, marram grass starts to lose its vigour and declines in abundance, making way for the development of grey dunes [called "Fixed coastal dunes with herbaceous vegetation (grey dunes)" code 2130* (European Commission 2007)] the next stage in the vegetation succession, dominated by drought tolerant mosses (Fig. 1). The degeneration of marram grass by sand stabilization was already noted by Marshall (1965), who called this phenomenon "The *Ammophila* problem" (not to be confused with the "Ammophila problem" referring to the invasion of the species outside its natural range as mentioned by e.g., Wiedemann & Pickart (1996).

The extraordinary sand fixing capacity of marram grass has been recognized in northwestern Europe for many centuries. *C. arenaria* was introduced for dune fixation in different parts of the world such as North America (Buell et al. 1995), Chile (Castro 1988), South Africa (Hertling & Lubke 1999), New Zealand (Hilton et al. 2004) and Australia (Webb et al. 2000). We here review the current state of the art with respect to the species' biotic and abiotic drivers of performance (Part II). We subsequently review the quantitative evidence of feedbacks with sand dynamics and demonstrate by both a new model and remote sensing how marram grass spatial configuration affects dune development (Part III). We end this review by an outlook towards further research.

Figure 1. Conceptual figure on the role of marram grass as an engineer in foredune formation. A. Once established, the species' sand capture ability shapes local sand accumulation, leading to an increase in dune volume. This sand accumulation will promote the species' growth, unless burial is too severe. If sand accumulation ceases, either due to decreased input from the sea, or to sheltering effects from surrounding vegetation, the plant performance will decrease due to pathogen accumulation in the roots, after which marram grass will degenerate. B. These dynamic feedbacks depend on the species' spatial configuration and external environmental conditions and will eventually shape the development of its volume and form, and, hence, its stability and resilience against storm surges under climate change.

II.   Abiotic and biotic drivers of marram grass performance

II.1 Abiotic drivers of marram grass performance

II.1.1. marram establishment

Marram grass establishes through seed germination or shooting of rhizome fragments detached from tussocks by coastal erosion. Konlecher & Hilton (2009) showed the potential for marine dispersal of such rhizome fragments over hundreds of kilometres, depending on regional sea currents. Seeds are mainly dispersed by wind, although the species only shows week morphological adaptations to wind dispersal (Huiskes 1979). Dispersal experiments by Pope (2006) and McLachlan (2014) suggest that a large majority of the seeds end up within a distance of less than 1 meter from the parent plant and wind dispersal abilities are probably restricted to several tens of meters. The potential for seed establishment of *Calamagrostis arenaria* is very high. First, this is due to a substantial seed production. Salisbury (1952) estimated over 20 000 caryopses are formed yearly per plant tussock. Second, marram grass has a long-lived seedbank. Viable seeds of up to 21 years old were recovered (Hilton et al. 2019). Third, the germination potential is high. Experiments under optimal laboratory conditions yielded germination percentages between 82 and 94% (Huiskes et al. 1979; Van Der Putten 1990, Bendimered et al. 2007, Lim 2011). In the field however, the establishment success of *C. arenaria* from seed is reputed to be very low on average (Huiskes 1977), although locally frequent germination (has been)/was observed in coastal dunes in The Netherlands (Van Der Putten 1990), New Zealand (Esler 1974) and North-America (Wiedemann 1987). Frequent establishment of marram is observed in embryonic foredunes and damp dune slacks (authors' personal observations). Seed germination strongly decreases with sand burial (Van der Putten 1990, Lim 2011, McLachlan 2014). Seedling emergence decreases linearly with burial depth, with a 3 cm burial already resulting in a germination reduction of 60% and no more seedlings emerge when seeds are buried under 9cm of sand; Fig. 2). The results we obtained from a burial experiment (See Supplementary material 1) are very similar to the findings of Lim (2011).

Figure 2. Seedling emergence in relation to sand burial (see supplement1)

Huiskes (1977) and Van Der Putten (1990) showed that the highest germination rates are obtained
with a fluctuating (day/night) temperature regime and a day temperature exceeding about 20°C. These
results are enhanced by stratification (cold pre-treatment). Optimal germination was obtained with a
20/30°C night/day temperature, with germination inhibited at lower temperatures of 10/20°C
night/day. These germination requirements retrieved in the lab correspond well with observations of
seeds germination in spring, when the temperature has risen sufficiently. Germination occurs only
under moist conditions (Huiskes 1979) and is inhibited when salinity exceeds 9 g/l (Chergui et al. 2013).
132

II.1.2. Marram growth and survival
Once established, marram grass growth and survival depends largely on the exposure of its local
environment for the physical forces of wind and water, that can directly dislodge plants or indirectly
affect growth and survival by transporting sediment. Partial burial of seedlings resulted in a 50-60%
increase of shoots length and root dry mass, but this vertical growth increases at the expense of lateral
growth and overall shoot biomass (number of tillers, which was maximal at 0-40% burial of the shoot
height) (Ievinsh and Andersone-Ozola, 2020). *C. arenaria* biomass increase showed a parabolic
response to burial with optimal growing performance at burial rates of 31 cm of sand per growing
season. (Nolet et al. 2018). The tolerance for burying was estimated to 78 to 96 cm burial/year. Reijers
et al. (2021) found more mature tussocks (clonal fragments containing ± 8 shoots) to perform equally
well under 0 or 2 cm burial every two weeks, but high mortality when burial reached 4 cm. Sediment
burial also indirectly influence marram grass growth by protecting the plants against the detrimental
effects of coastal flooding. Higher and larger embryo dunes are less susceptible to erosion during the
winter storm season, which positively influences marram grass growth during summer (Van
Puyenbroek et al. 2017).

Besides exposure to physical forces, soil nutrient levels can have a large influence on marram grass
performance as well. In general, sandy coastal systems are nutrient-limited and *C. arenaria* can cope
with these nutrient-poor conditions through symbiosis with arbuscular mycorrhizal fungi and by
recycling its own plant material through slow decomposition (Kowalchuk et al. 2002, Reijers et al.
2020).

Despite its occurrence in nutrient-limited conditions, *C. arenaria* requires substantial
levels of nitrogen, phosphorus and potassium for good growth (Willis 1965). A higher availability of N
and P in lime- and iron-poor dunes, due to atmospheric deposition, has been proposed as a mechanism
of the species' local expansion in coastal dunes (Kooijman et al., 1998; Kooijman and Besse, 2002).

Increases in temperature, nutrients and precipitation stimulate vegetation growth and lead to a global
greening of coastal dunes (Jackson et al., 2019). This global greening affects the natural sediment-
sharing capacity of coastal dunes, by hampering sediment transport to the hinterland (Gao et al. 2020).
Reduced sediment mobility and dune stabilization are thought to threaten several ecological functions,
while it can increase the protective function of coastal dunes by lowering erosion susceptibility
(Delgado-Fernandez et al. 2019, Gao et al. 2020, Pye & Blott 2020).

II.2 Biotic constraints on marram grass performance

II.2.1. Negative plant-soil feedback

Marram grass was found to perform worse in its own rhizosphere soil than in either sand from the sea
floor or in sterilized soil from its own rhizosphere (van der Putten et al. 1988, 1993), demonstrating
that a biotic factor in the soil causes a decline in marram grass performance. The exact cause of this
biotic control is to date unclear. The first studies attempting to pinpoint the soil organisms causing the
decline of marram grass implicated root-feeding nematodes as well as pathogenic fungi (Van der
Putten et al. 1990, De Rooij and van der Goes 1995, van der Putten and van der Stoel 1998, van der
Stoel and van der Putten 2002). However, the exact species causing a performance reduction could
not be identified across these studies. Competitive and facilitating interactions among these co-
infecting belowground parasites (Brinkman 2005&a,b,c) but also more complex trophic interactions,
including those with microbes within the rhizophere (Piśkiewicz et al. 2008, 2009; Costa et al. 2012)
were found to be mediators of marram performance under experimental conditions. Furthermore, it
has been shown that the negative effect of certain nematode species can be mitigated by the positive
effect of mycorrhizal and endophytic fungi (Little and Maun 1996, de la Peña et al. 2006, Hol et al.
2007). Overall, the net effect on marram grass performance of all naturally occurring members of the
soil community is generally negative. Although the exact mechanism is difficult to identify, evidence
for the "escape hypothesis" remains strong, i.e., marram grass needs regular burial by wind-blown
sand free of soil organisms so that it can grow new roots into an –at least temporarily– enemy-free
space. Plant-soil feedbacks caused by other plant species also play a role. Conditioning of soils by
Carpobrotus edulis (L.) N.E.Br., a species originating from South Africa and one of the most invasive
plant species in the Mediterranean, suppresses marram grass biomass and in some cases survival rate
(de la Peña et al. 2010). The increase of *Carpobrotus* in the dunes of central Italy (Sperandii et al. 2018)
has therefore been linked to large-scale decreases in marram grass. Marram grass also shows a
reduced germination on soil invaded by *Acacia longifolia* (Andrews) Willd., yet it performed better on
invaded than native soil after 12 weeks of growth (Morais et al. 2019).

II.2.2. Aboveground biotic interactions
The aboveground organisms associated with marram grass are in general well known (i.e., Huiskes
1979, Heie 1982, 1986, Holman 2009, Vandegehuchte et al. 2010a), but very little is known about their
effects on plant performance. The associated herbivore species have the potential to induce serious
reductions in aboveground performance in a controlled environment (Balachowsky and Mesnil 1935,
Nye 1958, Heie 1986, Vandegehuchte et al. 2010a), but so far no experiments were conducted in
nature. Marram grass does not seem to be controlled to any significant extent by mammalian grazers
either (Badhresa 1977, Huiskes 1979), except for some feeding on young shoots (Rowan 1913). Seed
predation has been observed (Huiskes 1979) but its magnitude and/or impact on marram grass
demography is unknown.

II.2.3. Control of above- and belowground communities by marram grass intraspecific variation
Intraspecific variation among marram grass populations can have strong effects on the abundance and
community composition of both above- and belowground invertebrate species (Vandegehuchte et al.
2011). This variation is linked to genetic variation in plant growth, which likely explains higher

abundances of aboveground invertebrates on local than on non-local marram grass populations.
Contrasting effects were found for root herbivores as their abundance and species richness negatively
covaried with the aboveground ones (Vandegehuchte et al. 2011, 2012). Additionally, it has to be
noted that a full soil biota community can have stronger effects on marram grass performance than
local abiotic soil properties (Vandegehuchte et al. 2010c), although performance can differ significantly
among soils differing substantially in abiotic properties. The relationships between marram grass and
its aboveground invertebrates can therefore not be understood independently of its belowground
invertebrates and the abiotic conditions of the soil.
214
215

II.2.4. Learning from elsewhere: marram grass as invasive species
Explanations for success of marram grass in its novel range have been sought in the popular "enemy
release hypothesis" (Keane and Crawley, 2002), mainly focusing on belowground enemies. Growth of
marram grass was significantly less reduced on soils from South African sites than on soils from the
Netherlands, indicating a weakened negative plant-soil feedback and thus potential role for enemy
release in South African soils (Knevel et al. 2004). However, this contrasts with findings from coastal
dunes of California, where soil sterilization experiments have shown that the performance of marram
grass is reduced to similar extents as in Europe when grown on non-sterilized soil (Beckstead and
Parker 2003), suggesting there is no enemy release. Furthermore, soil biota from three native South
African plant species did not suppress marram grass growth, but biota from soils beneath the tropical
cosmopolitan dropseed *Sporobolus virginicus* (L.) Kunth did, suggesting that this plant species may
confer biotic resistance against invasion by marram grass (Knevel et al. 2004). A large sampling
campaign of soil and roots from Tasmania, New Zealand, South Africa and the west coast of the USA
revealed that marram grass did not have fewer root-feeding nematode taxa in these regions than in
its native range. However, native plants in the novel range had more specialist root-feeding nematode
taxa than marram grass, while specialists such as cyst and root-knot nematodes, which are common in
the native range of marram grass, were not found in the southern hemisphere (van der Putten et al.
2005). Invasiveness of marram grass thus seems correlated with an escape from specialized root-
feeding nematodes.

II.3. Dynamic feedbacks between aeolian fluxes and vegetation development
The capture rates of sand by vegetation and its effect on dune topography have been intensely studied
(e.g., Hesse & Simpson 2006). There is also abundant literature on how obstacles that represent
vegetation obstruct or facilitate sand fluxes, with strong analogies to research on fluid dynamics.
Typically, multiple configuration of height and density of the obstacles are used (e.g., reed stems (Arens
et al., 2001); see (Mayaud and Webb, 2017) for a comprehensive review on aeolian sand transport in
drylands). These studies quantify how much of the total force of the wind by drag is reduced by the
vegetation, also referred to as shear stress partitioning and expressed as drag coefficients (Raupach,
1992). All studies show this drag coefficient to be positively related to the roughness induced by the
density and impermeability of the set of obstacles, and their height (Hesp et al., 2019). Since these
experiments use marram-grass surrogates like artificial cylinders, stem bundles or even dead plant
material, they do not represent the realised morphology of dune vegetation, which precludes further
progress in understanding the feedbacks between sediment capture and plant growth. Clusters of
tillers enhance sand deposition by lowering wind speed and associated shear stress within the
vegetation canopy (Charbonneau & Casper 2018). Larger tussocks are able to capture more sand,
thereby imposing a positive feedback on their own development and vigour. The plants react to burial
by rapid production of elongated stem internodes, but the exact extent of this growth response is
unknown except for young plants under lab conditions (Levinsh and Andersone-Ozola 2020). As sand
burial induces the production of high-density vertical tillers and horizontally expanding rhizomes
(Reijers et al., 2021), marram grass steers dune morphology (van der Putten et al. 2005, Hart et al.

2012, Darke et al. 2016). *C. arenaria* is, because of this growth strategy, associated with the
development of higher and steeper dunes compared to those formed by its North-American sister
species *C. breviligulata*, making dunes build by the former potentially more resistant to erosion
(Zarnetske et al., 2012; Seabloom et al., 2013; Charbonneau et al., 2016).

Both vertical and horizontal growth responses influence the size and shape of *C. arenaria* tussocks, but
also directly determine remaining sand drift at the rear side of these vegetated patches (Reijers et al.,
2021). With increasing densities and cover, *C. arenaria* subsequently stabilizes the mobile sand
(Huiskens 1979). At least in European coastal dunes, the ceasing sand fluxes mediated by the species'
increasing densities, and the resulting increases in dune height and slope, then induce on longer time
frames a negative feedback on the species' vigour in the long run, causing the species to slowly die off
(e.g., Van der Putten 1994, De Rooij and Van der Goes 1995, van der Putten and van der Stoel 1998,
van der Stoel and van der Putten 2002) as the resource (fresh sand) becomes limiting. The spatial
configuration and morphology of the vegetation is therefore dynamically coupled to shear stress. Sand
capture directly alters potential density, growth and lateral expansion of the vegetation, which
feedbacks to patterns in flow parameters (velocity, turbulence, intermittency) because of sheltering
effects by vegetation and dune topography. The qualitative importance of these feedbacks for the
large-scale geomorphology of coastal dunes is well-appreciated (Durán and Herrmann, 2006; Hesse
and Simpson, 2006; Durán et al., 2009; Durán and Moore, 2013), but very few data on the feedbacks
between sand fluxes and vigour of the foredune vegetation are available. So far, different vegetation
states rather than vegetation dynamics have been linked to dune height potential and subsequent risks
of overtopping events and flooding (Seabloom et al., 2013).

III Towards an integrated model of the plant-sand feedbacks
III.1 The integration of vegetation-dune feedbacks in existing process-based models
The importance of the vegetation-dune feedbacks is still not well understood, let alone quantified and
incorporated into predictive models for coastal dune dynamics (Puijenbroek et al., 2017). Current
state-of-the-art 3D models for coastal dune development (e.g., DUBEVEG (Keijsers et al., 2015), CDM
(Durán and Moore, 2013), AeoLIS (Hoonhout and de Vries, 2016)) are able to simulate topographic
development of coastal dunes and sediment transport at spatial scales relevant for coastal managers
as a function of sediment supply, probabilities of vegetation development, descriptions of flow field,
and dune erosion by waves. These (coupled) 3D coastal dune models are a product of the basic physical
principles and sediment transport models, and they are essential for the prediction of dune
development. Furthermore, they need validation from field experiments containing high-quality
datasets relevant for dune development. With exception of the DUBEVEG model (De Groot et al.,
2011), which has coarse vegetation dynamics incorporated, recent coastal functioning models have
ignored ecological interactions across scales (e.g., Duran and Moore 2013, Van Westen et al. 2019). As
important engineer species from coastal dunes differ in physical features and life history, they
differently affect dune development, with for instance sand couch grass (*Elymus farctus* (Viv.)
Runemark ex Melderis) giving rise to 'lower broader dunes', and marram grass enabling dunes to
develop into a 'higher, hummocky peaked topography' (Hesp, 2002; Puijenbroek et al., 2017; Reijers
et al., 2019; Schwarz et al., 2019).

All existing dune erosion models treat the processes at the dune face in a simplified way. No process-
based description is implemented to describe the formation of vertical cliffs at the dune foot
undermining the dune slope with subsequent geotechnical failures of the dune slope that results in
slumps of sand on the beach that can be taken away by the waves. For this marshes, Bendoni et al.
(2019) implemented a hydro-morphodynamic interaction model in XBeach (Roelvink et al., 2009) to
evaluate erosion of marsh boundaries due to wave impact. Although this study is limited to the
cohesive sediments' environment, soil reinforcement due to roots has been modelled, which might be
extended to other environments in the future. Physical scale model experiments, with dunes,

vegetation and disturbances scaled towards lab conditions, have demonstrated that roots, which geotechnically strengthen a sand volume, significantly reduce the dune erosion compared to bare sand (Feagin et al. 2019, Bryant et al. 2019). Presently, only indirect implementations are possible namely by tuning calibration parameters influencing the morphodynamics.

III.2. Insights from a new simulation model

III.2.1. The geography of marram spatial configuration

From II.3, it is clear that feedbacks between the environment and the spatial distribution of marram grass impact dune development. We mapped marram cover and spatial contingency in 20x20 m² grid cells along the coastlines of northern France, Belgium, Netherlands and South-England (see supplement S2), to identify realistic ranges in nature. Marram grass is –as predicted from the species' biology- predominantly showing a clustered distribution with JC (join-count; an established method that quantitatively determines the degree of clustering or dispersion, see S2) values between 20 and 80, so ranging from random (values close to zero) to highly clustered patterns (Figure 3). A mean clustering pattern with JC values around 50 is stable over the four studies regions. No underdispersed (so regular) patterns were observed. Interestingly, marram grass spatial cover at these spatial scales is strongly country-specific with UK and France being represented by well-vegetated dunes. Dunes in Belgium and the Netherlands appear to be in more dynamics states with quite a substantial presence of areas with a low cover (see IV).

Figure 3. Density distribution plots of observed cover (P) and spatial correlation (JC) of marram grass in 20x20 m² grid cells in dunes from the Isles of Scilly to Norfolk in England (UK) and from Somme (Fr) to Texel (NL) on the continental side along the North Sea and the Flemish West coast (B) (right panel). Note that for visualisation, a subsampling of 1000 points (2%) was performed for (A).

III.2.2. Simulating dune growth in relation to marram grass spatial configuration

We developed a grid-based dune simulation model that computes aeolian transport processes and changes in vegetation growth and dune morphology based on their dynamic feedbacks and marram spatial organisation. The landscape is grid-based with cells having dimensions of 0.20x0.20 m². The 100x100 cell matrix therefore corresponds with a dune area of 20x20m². We refer to supplementary material S2 for a detailed process overview, references to the code. We simulated changes in aeolian processes and wind dynamics at a day-resolution. We used averages over 7 years (2010-2017) received from the Royal Meteorological Institute at Koksijde, and scaled them to the four main different wind directions as used in the model by (Nolet et al. 2018b), each corresponding to a side of the landscape. Wind speed and direction is drawn daily from a normal distribution, based on monthly average wind speed and its standard deviation. *Sand input*, the material blown into the system from the beach (N-direction here), is expressed as a relative percentage of the maximum sand saturation flux, i.e. the

maximal amount of sand that can be carried by the wind. Lateral winds (E and W-directions in the grid)
have an influx which corresponds with the most recent outflux of a lateral wind (thus, we represent
the landscape as tube to avoid edge artefacts). This amount is constantly updated during a simulation.

Sand deposition is directly dependent on shear velocity which is a function of wind velocity Hoonhout
and de Vries 2016; Durán et al., 2010), vegetation density (Durán et al. 2010) and its roughness (Durán
and Moore 2013). Increases in shear stress due to funnel effects are included, as are gravity and shelter
effects. Maximum angles of repose are set to 34° (Durán et al., 2010)when vegetation is absent (Durán
et al., 2010). These angles increase with vegetation density. As such, avalanches are less prevalent
when plant density is high. Moreover, erosion is inhibited in locations sheltered by lee slopes at an
angle of maximal 14° (Kroy et al., 2002). *Marram grass* dynamics are seasonal (only growth in spring
and summer), with local growth modelled as outlined in this review. Vertical as well as lateral growth
during the growing season is modelled as a logistic function up to maximal heights, and directly
depending on sand deposition (Nolet et al., 2018), leading to positive growth under intermediate sand
accretion and complete burial leading to marram grass die-off. Lateral growth follows Lévy-patterns as
determined by (Reijers et al., 2019), and are here modelled by simplified neighbour expansion
processes. No growth occurs during autumn and winter but sand accumulation continuous. The net
height after winter burial determines the starting conditions for vertical growth in the next season. No
germination events were modelled as these are to date not (or only rarely) witnessed in foredunes the
last decade.

To validate the model predictions, we compared outcomes from the model with those from a statistical
model linking changes in topography over five years as derived from LIDAR in relation to the initial
marram spatial configuration as determined from aerial photographs in 2015 in Belgium (Fig. 4;
detailed methods in supplementary material S3 and S4). Our model simulations (Fig 5, upper panels)
were run for initial marram cover in the range 0.1-0.9 (as without marram cover, only erosion of the
initialised sand volume is occurring without establishment) and join counts between 20 and 70).

A. The RGB, NIR and DTM data are retrieved for the sites under study

B. All input data are used to generate vegetation maps (marram cover, 20x20 cm²)

C. Differential DTM's of changes in dune height over 5 years are produced (20x20 cm²)

D. Marram spatial distribution statistics are caluculated at grids of 20x20 m² (darker colours are higher values)

Cover (P)

Spatial correlation (JC)

E. Differential heights are integrated into grids of 20x20 m² (red: erosion; blue: accretion)

Figure 4. Workflow for connecting observed differences in dune height over a period of 5 years to marram spatial
configuration in coastal foredunes (case presented: Schipgatduinen, Koksijde, Belgium)

The modelled height changes agree in general terms with those observed. The observed larger effects
in the field therefore suggest slightly larger sand input, either due to sand availability or changes in
wind strengths, from the beach as sand input initiated in the model based on rough estimates from
the Belgian coast (unpub. data Rauwoens & Strypsteen). Alternatively, the small scale of the
mechanistic model may underestimate wind saturation and therefore sand displacements (see IV.
outlook). Dune height increases at intermediate cover of marram grass, so *P~0.5*. However, the
simulation model predicts increases to be maximal under low cover and more random (i.e., less
clustered) distributions of the marram grass tussocks (low *P* and low *JC*). According to the simulation
model, local changes in dune topography, estimated as the coefficient of variation (CV) of grid cell-
level differences in height, show most changes occurring in dunes with clustered marram patches or
patches with a low amount of vegetation, but more random patterns. Predicted changes from LIDAR
follow the same pattern as the ones generated by the simulation model. Only under low cover and
intermediate clumping, a more homogenous increase of the dune is predicted by the computer model.

386
**Figure 5. Output of the simulations computing changes in dune height and topography (CV of height changes over all grid
cells) over a period of 5 years (upper panel), and similar metrics as observed from LiDAR data from coastal dunes along
the Belgian west coast (lower panels).**

Analysis of the LIDAR data also showed decreasing dynamics with increasing distance from the sea (see
supplement S4). The obtained effect sizes (supplement S4) and accompanying visualisations of the
modelled effects (figure 5) indicate that the observed changes in integrated dune height and form
differ from those of the simulation in the sign and strength of the cover x spatial clustering (P:JC and
the interaction between P² and JC). The most prominent difference lies in the predicted erosion
dynamics under low marram cover and a strong clustering. This suggests that the sand accretion
capacities of marram grass under these conditions needs to be re-evaluated.

IV.   Discussion and outlook
A mechanistic understanding of the vegetation-sedimentation feedbacks that steer the natural
development of coastal dunes is essential for conserving and restoring the function of coastal dunes

as natural flood barriers. Climate change, and its impact on feedbacks between marram grass and sand fluxes, is anticipated to strongly alter dune formation and dune resilience (Pakeman et al., 2015; van Puijenbroek et al., 2017). We here reviewed the state of knowledge on the ecology of marram grass in relation to dune formation, flow attenuation, sediment deposition and plant growth. Our model and LIDAR analysis showed that the joint increase of volume and variability under low cover and less clustered spatial configurations have the highest impact on local sand accretion and dune morphology. Such conditions steer impose a positive feedback on vertical growth. Strong erosion dynamics are conversely anticipated to preclude establishment at further distances from existing tussocks. Scale-dependent feedbacks lead to patterns of self-organisation (Rietkerk & van de Koppel 2008) and need to be quantified and further integrated into mechanistic models to forecast coastal dune formation in relation to climatic conditions. Earlier research showed considerable variation in marram growth (Vandegehuchte et al. 2010a) and expansion strategies (Reijers et al. 2021), and changes here-in can be expected with respect to future climatic conditions. The relevance of this intraspecific variation remains to be understood, also from the perspective of planting actions to actively build resilient dunes in the light of climate change.

A resilient coastal dune system is anticipated to be one where vegetation and bare sand coexist in a stable equilibrium, hence a state to which the system should bounce back after any disturbance, e.g., by erosion. The permanent loss of sand dynamics by changing sand input, fragmentation or anthropogenic dune stabilization are expected to lead to catastrophic shifts causing dunes to become hyperstatically fixed by plantation and succession (Jackson et al. 2019, Gao et al. 2020). On the other hand, at too low initial densities, the vegetation may be disrupted by strong sand drifts, also following intense trampling by people, leading to a hyperdynamic and unvegetated state. A resilient dune should balance between both extremes (Borsje et al. 2011) and this resilience will thus largely be determined by the current vegetation density and configuration, local conditions of sand supply, connectivity with the beach, and erosion. The state of the marram dune can be expected to impact further inland sand drift. Narrow stretches thereby have the potential to determine dune stability at larger spatial scales by affecting the total dune system volume, and the further vegetation succession dynamics (e.g., Olff et al. 1993, Fenu et al. 2013). These are less relevant from a coastal protection perspective but of major importance for biodiversity conservation (European Commission on Habitat of Directive 92/43 EEC)

Coastal dunes along the coast of the North-Sea and Channel show a remarkable convergence in the spatial clustering across the four studied countries, and this clustering seems to be preserved across the range of vegetation cover. This finding suggests an optimal clustering in European dunes, which is anticipated to result from the species' self- organizing capacity. At intermediate cover, this clustering leads to largest changes in dune growth. We anticipate that the availability of sufficient aeolian dynamics at small scales drives this overall increase in dune volume. This review also shows these conditions to facilitate marram grass performance because of the steady supply of fresh sand. Although more research is needed, this finding suggests that such a spatial configuration can optimize both marram grass performance and dune resilience by maximizing growth. Deviations from this state, especially in terms of cover –note that the clustering metric becomes less relevant with increasing cover– are then likely disturbed states resulting from either ceasing sand dynamics or vegetation development. As we only documented patterns in marram grass spatial configuration, we lack insights into the underlying causes. Are they due to sediment transport potential, or correlated to region-specific variables such as tidal amplitude, wave height, or beach width? Alternatively, it is not unlikely that variation in both dune management –especially planting campaigns– and the differences in recreational pressure are at the basis for this variation. Since we showed marram grass' spatial configuration to affect both dune growth and topography and therefore sand fluxes further inland, this variation is anticipated to have strong implications for coastal protection. Pending on the state, recreational pressures may constrain dune stabilisation and keep the system in a dynamic, and presumed optimal state with respect to resilience, or facilitate erosion and the transition to

hypermobile states. Clearly, the negative and positive contributions of such recreational pressures
need to be assessed case by case, and in direct connection to the local environmental (boundary)
conditions (Nunes et al. 2020).

Incorporating the available information allowed us to mechanistically build models that support
generic predictions of dune volume and topography change at short spatial and temporal scales. The
model does, however, still contain gaps in terms of parameterisation and validation (both
observational and experimental), especially with regard to very dynamic conditions (low, clustered
cover by marram grass). While any prediction in this specific parameter range can be questioned for
its relevance ("How natural are these configurations, if we do not observe them?"), we argue that this
is of the utmost relevance in the light of dune-building campaigns where marram grass needs to be
planted, for instance in front of existing dikes. The presented simulation model also operates at
relatively small scales relevant for vegetation dynamics, but potentially underestimating realised wind
saturation in barely vegetated dunes. Deviations between the observed and predicted changes in dune
volume likely result from such scaling issues. Upscaling of dune-vegetation dynamics can be achieved
by linking sand-output conditions from the most seaward-oriented dunes as input conditions for those
more inland. Under such conditions, sand-vegetation dynamics need to be extended towards other
species occurring along the expected succession gradient. It remains to be studied whether
simplifications using vegetation height and biomass (Duran *et al.* 2009), as the mediator of such
interactions suffice. A major driver of dune development is the amount of sand input from the beach.
While difficult to measure, the joint analysis of vegetation and dune development may be used as a
reliable predictor of such sea-land sediment fluxes by means of inverse modelling.

More research is needed on potential regional changes in the interactions among vegetation
development, dune growth and sand fluxes, for example caused by differences in climatic conditions
and marram grass genetic variation. Dune volumes are prime determinants of their functioning in
coastal protection, with large bodies of sand providing a larger safe zone against inundation risks from
storm surges. Vegetation dynamics are expected to have a strong impact on the healing capacity of
foredunes, i.e., how fast they recover to earlier states after erosion by such storm events, or in the
longer term resulting from sea level rise. Current models enable us to predict changes in dune volume
and form, but it remains unknown how erosion is affected by the vegetated state of the dune. For
example, to which degree erosion resistance is determined by the marram grass root network still
needs to be elucidated. Ultimately, the addition of such information on the mechanistic underpinnings
of the patterns generated by our models would reduce the uncertainty of their predictions to the
benefit of all stakeholders involved.

Funding
This research was done with financial support by the Interreg 2seas project Endure. DB is funded by
Ugent-BOF-grant BOF.24Y.2021.0012.01. FB is supported by Research Foundation – Flanders (FWO)

Contribution to the Field Statement
Coastal dunes provide safety against future climate change induced sea-level rises and storm surges.
Contrary to hard protections, they have the advantage to grow with the sea-level. Coastal dunes thus
provide important safety services, especially in highly urbanised lowland areas. Because these coastal
landscapes are formed by the interplay between the dune building marram grass and sand dynamics,
a quantitative study of these interactions is necessary to understand and forecast how climate change
will impact dune development and its resilience. We here provide a synthesis on the available insights,
and complement these insights with new experiments and remote sensing data from coastal dunes
along the North Sea. This information is integrated into a mechanistic model to demonstrate the
importance of marram grass ecology, and its emerging spatial configuration for dune development.

We build on these insights to identify important caveats and future research directions to effectively
manage and construct dunes as a solid nature-based solution from the fundamental ecological and
geomorphological building blocks.

Reference list

Arens, S. M., Baas, A. C. W., Van Boxel, J. H., and Kalkman, C. (2001). Influence of reed stem density
on foredune development. Earth Surf. Process. Landforms 26, 1161–1176. doi:10.1002/esp.257.

Bakker, J. P. (1976). Phytogeographical Aspects of the Vegetation of the Outer Dunes in the Atlantic
Province of Europe. Journal of Biogeography 3, 85. doi:10.2307/3038138.

Balachowsky, A., Mesnil, L. (1935). Les Insectes Nuisibles aux Plantes Cultivées. Paris.Beckstead, J.
and Parker, I. M. 2003. Invasiveness of Ammophila arenaria: Release from soil-borne pathogens? -
Ecology 84: 2824-2831.

Bendimered, F. Z., Mehdadi, Z. & Benhassaini, H. (2007). Etude de la germination et de croissance
foliaire de l'oyat (Ammophila arenaria (L) Link. en conditions contrôlées. Acta Botanica Gallica 154(1):
129–140.

Borsje, B. W., van Wesenbeeck, B. K., Dekker, F., Paalvast, P., Bouma, T. J., van Katwijk, M. M., et al.
(2011). How ecological engineering can serve in coastal protection. Ecological Engineering 37, 113–
122. doi:10.1016/j.ecoleng.2010.11.027.

Brinkman EP, Duyts H, Van Der Putten WH (2005a) Consequences of variation in species diversity in a
community of root-feeding herbivores for nematode dynamics and host plant biomass. Oikos 110:
417–427.

Brinkman EP, Duyts H, Van Der Putten WH (2005b) Competition between endoparasitic nematodes
and effect on biomass of Ammophila arenaria (marram grass) as affected by timing of inoculation
and plant age. Nematology 7: 169–178

Brinkman EP, Troelstra SR, van der Putten WH (2005c) Soil feedback effects to the foredune grass
Ammophila arenaria by endoparasitic root-feeding nematodes and whole soil communities. Soil
Biology & Biochemistry 37: 2077–2087.

Brinkman, E. P., et al. (2015). Plant-feeding nematodes in coastal sand dunes: occurrence, host
specificity and effects on plant growth. - Plant Soil 397: 17-30.

Castro, C. A. (1988). The artificial construction of foredunes and the interference of dune-beach
interaction, Chile. Journal of Coastal Research, Special Issue 3: 103–107.

Charbonneau, B. & Casper, B. B. (2018). Wind tunnel tests inform Ammophila planting spacing for
dune management. Shore & Beach, 86: 37-46.

Charbonneau, B. R., Wnek, J. P., Langley, J. A., Lee, G., and Balsamo, R. A. (2016). Above vs.
belowground plant biomass along a barrier island: Implications for dune stabilization. Journal of
Environmental Management 182, 126–133. doi:10.1016/j.jenvman.2016.06.032.

Chergui, A., Latifa, E. H. & Mohammed, M. (2013). The effects of temperature, hydric & saline stress
on the germination of marram grass seeds (Ammophila arenaria L.) of the SIBE of Moulouya
embouchure (Mediterranean - North-eastern Morocco). Research Journal of Pharmaceutical,
Biological and Chemical Sciences 4(1): 1333-1339.

Costa, S. R., Kerry, B.R., Bardgett, R.D., & Davies, K.G. (2012). Interactions between nematodes and
their microbial enemies in coastal sand dunes. - Oecologia 170: 1053-1066.

Darke, I. B Walker, I.J. & Hesp, P.A. (2016). Beach-dune sediment budgets and dune morphodynamics
following coastal dune restoration, Wickaninnish Dunes, Canada. - Earth Surf. Process. Landf. 41:
1370-1385.

De Groot, A. V., Berendse, F., Riksen, M., Baas, A., Slim, P., and van Dobben, H., Stroosnijder, L.
(2011). Modelling coastal dune formation and associated vegetation development. Geophysical
Research.

de la Peña E, Echeverría Rodríguez S, van der Putten WH, Freitas H, Moens M (2006) Mechanism of
control of root-feeding nematodes by mycorrhizal fungi in the dune grass Ammophila arenaria. New
Phytologist 169: 829–840.

de la Peña, E., de Clercq, N., Bonte, D. et al. (2010) Plant-soil feedback as a mechanism of invasion by
Carpobrotus edulis . Biol Invasions 12, 3637–3648.

De Rooij - Van der Goes PCEM (1995) The role of plant-parasitic nematodes and soil-borne fungi in
the decline of Ammophila-arenaria (L) Link. New Phytologist 129: 661–669.

de Vries, S., Southgate, H. N., Kanning, W., and Ranasinghe, R. (2012). Dune behavior and aeolian
transport on decadal timescales. Coastal Engineering 67, 41–53.
doi:10.1016/j.coastaleng.2012.04.002.

Delgado-Fernandez, I. (2010). A review of the application of the fetch effect to modelling sand supply
to coastal foredunes. Aeolian Research 2, 61–70. doi:10.1016/j.aeolia.2010.04.001.

Delgado-Fernandez, I., Davidson-Arnott, R. G., & Hesp, P. A. (2019). Is 'remobilisation'nature
restoration or nature destruction? A commentary. Journal of Coastal Conservation, 23(6), 1093-1103.

Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I., and Marbà, N. (2013). The role of coastal
plant communities for climate change mitigation and adaptation. Nature Climate Change 3, 961–968.
doi:10.1038/nclimate1970.

Durán, O., and Herrmann, H. J. (2006). Vegetation Against Dune Mobility. Phys. Rev. Lett. 97, 188001.
doi:10.1103/PhysRevLett.97.188001.

Durán, O., and Moore, L. J. (2013). Vegetation controls on the maximum size of coastal dunes.
Proceedings of the National Academy of Sciences 110, 17217–17222. doi:10.1073/pnas.1307580110.

Durán, O., Parteli, E. J. R., and Herrmann, H. J. (2010). A continuous model for sand dunes: Review,
new developments and application to barchan dunes and barchan dune fields. Earth Surface
Processes and Landforms 35, 1591–1600. doi:10.1002/esp.2070.

Durán, O., Schwämmle, V., Lind, P. G., and Herrmann, H. J. (2009). The dune size distribution and
scaling relations of barchan dune fields. Granular Matter 11, 7–11. doi:10.1007/s10035-008-0120-4.

Esler, A. E. (1974). Vegetation of the sand country bordering the Waitakere Range, Auckland: the
southern beaches. Proceedings of the New Zealand Ecological Society 21: 72–77.

European Commission DG Environment (2007) Interpretation manual of European Union habitats
(version EUR27). Brussels: European Commission DG Environment.

Fenu, G., Carboni, M., Acosta, A.T.R. & Bachetta G. (2013). Environmental Factors Influencing Coastal
Vegetation Pattern: New Insights from the Mediterranean Basin. Folia Geobot 48, 493–508

Gao, J., Kennedy, D. M., & Konlechner, T. M. (2020). Coastal dune mobility over the past century: A
global review. Progress in Physical Geography: Earth and Environment, 44(6), 814-836.

Hart, A. T., et al. (2012). The Impact of Ammophila arenaria Foredune Development on Downwind
Aerodynamics and Parabolic Dune Development. - J. Coast. Res. 28: 112-122.

Heie O.E. (1982). The Aphidoidea (Hemiptera) of Fennoscandia and Denmark. II. The family
Drepanosiphidae. Fauna Entomologica Scandinavica, vol. 11. Scandinavian Science Press Ltd.,
Klampenborg.

Heie O.E. (1986). The Aphidoidea (Hemiptera) of Fennoscan- dia and Denmark. III. Family Aphididae:
subfamily Pterocommatinae & tribe Aphidini of subfamily Aphidinae.

Hesp, P. (2002). Foredunes and blowouts: initiation, geomorphology and dynamics. Geomorphology
48, 245–268. doi:10.1016/S0169-555X(02)00184-8.

Hesp, P. A., Dong, Y., Cheng, H., and Booth, J. L. (2019). Wind flow and sedimentation in artificial
vegetation: Field and wind tunnel experiments. Geomorphology 337, 165–182.
doi:10.1016/j.geomorph.2019.03.020.

Hertling, U.M. & Lubke, R.A. (1999). Indigenous and Ammophila arenaria-dominated dune vegetation
on the South African Cape coast. Appl. Veg. Sci. 2: 157–168

Hesse, P. P., and Simpson, R. L. (2006). Variable vegetation cover and episodic sand movement on
longitudinal desert sand dunes. Geomorphology 81, 276–291. doi:10.1016/j.geomorph.2006.04.012.

Hilton, M. Konlechner, T., McLachlan, K., Lim, D. & Lord, J. (2019). Long-lived seed banks of
Ammophila arenaria prolong dune restoration programs. Journal of Coastal Conservation 23: 461–
471.

Hilton, M., Duncan, M. & Jul, A. (2004). Processes of Ammophila arenaria (marram grass) invasion
and indigenous species displacement, Stewart Island, New Zealand. Journal of Coastal Research
21(1): 175–185.

Hol WHG, de la Pena E, Moens M, Cook R (2007) Interaction between a fungal endophyte and root
herbivores of Ammophila arenaria. Basic and Applied Ecology 8: 500–509.

Holman J. (2009). Host Plant Catalog of Aphids, Palaearctic Region. Springer.

Hoonhout, B., and de Vries, S. (2016). A process-based model for aeolian sediment transport and
spatiotemporal varying sediment availability. Journal of Geophysical Research: Earth Surface 121,
1555–1575. doi:10.1002/2015JF003692.

Huiskes, A. H. L. (1977). The natural establishment of ammophila arenaria from seed. Oikos 29 (1):
133-138.

Huiskes, A. H. L. (1979). Biological flora of the British Isles: Ammophila arenaria (L.) Link (Psamma
arenaria (L.) Roem. et Schult; Calamagrostis arenaria (L.) Roth). Journal of Ecology 67(1): 363-382.

Ievinsh, G., and Andersone-Ozola, U. (2020). Variation in Growth Response of Coastal Dune-Building
Grass Species Ammophila Arenaria and Leymus Arenarius to Sand Burial. Botanica 26, 116–125.
doi:10.2478/botlit-2020-0013.

IPCC (2014). Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to
the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. , eds. Team, R. K.
Pachauri, and L. A. Meyer Geneva, Switserland.

IPCC (2018). Special Report on Global Warming of 1.5°C.

Jackson, D. W. T., Costas, S., González-Villanueva, R., and Cooper, A. (2019). A global 'greening' of
coastal dunes: An integrated consequence of climate change? Global and Planetary Change 182,
103026. doi:10.1016/j.gloplacha.2019.103026.

Keane, R. M., and Crawley, M. J. (2002). Exotic plant invasions and the enemy release hypothesis.
Trends in Ecology & Evolution 17, 164–170. doi:10.1016/S0169-5347(02)02499-0.

Keijsers, J. G. S., De Groot, A. V., and Riksen, M. J. P. M. (2015). Vegetation and sedimentation on
coastal foredunes. Geomorphology 228, 723–734. doi:10.1016/j.geomorph.2014.10.027.

Knevel, I. C., Lans, T., Menting, F.B., Hertling, U.M. & Van der Putten, W. (2004). Release from native
root herbivores and biotic resistance by soil pathogens in a new habitat both affect the alien
Ammophila arenaria in South Africa. - Oecologia 141: 502-510.

Konlechner, T. M. & Hilton, M. J. (2009). The potential for marine dispersal of Ammophila arenaria
(marram grass) rhizome. Journal of Coastal Research 56: 434–437.

Kooijman, A. M., and Besse, M. (2002). The higher availability of N and P in lime-poor than in lime-
rich coastal dunes in the Netherlands. Journal of Ecology 90, 394–403.
doi:https://doi.org/10.1046/j.1365-2745.2001.00661.x.

Kooijman, A. M., Dopheide, J. C. R., Sevink, J., Takken, I., and Verstraten, J. M. (1998). Nutrient
limitations and their implications on the effects of atmospheric deposition in coastal dunes; lime-
poor and lime-rich sites in the Netherlands. Journal of Ecology 86, 511–526.
doi:https://doi.org/10.1046/j.1365-2745.1998.00273.x.

Kowalchuk, G. A., De Souza, F. A., & Van Veen, J. A. (2002). Community analysis of arbuscular
mycorrhizal fungi associated with Ammophila arenaria in Dutch coastal sand dunes. Molecular
Ecology, 11(3), 571-581.

Kroy, K., Sauermann, G., and Herrmann, H. J. (2002). Minimal model for sand dunes. Physical Review
Letters 88, 543011–543014. doi:10.1103/PhysRevLett.88.054301.

Lim, D. (2011). Marram grass seed ecology: the nature of the seed bank and secondary dispersal. MSc
thesis, University of Otago, Dunedin, 177 p.

Little LR, Maun MA (1996) The 'Ammophila problem' revisited: A role for mycorrhizal fungi. Journal
of Ecology 84: 1–7.

Marshall JK (1965) Corynephorus-Canescens (L) P Beauv as a model for the Ammophila problem.
Journal of Ecology 53: 447–463

Martinez, M. L., and Psuty, N. P. (2004). Coastal Dunes Ecology and Conversation. Springer Berlin
Heidelberg doi:10.1017/CBO9781107415324.004.

Maun, M. A. (1998). Adaptations of plants to burial in coastal sand dunes. Canadian Journal of Botany
76(5): 713-738.

Maxwell, P. S., Eklöf, J. S., Katwijk, M. M. van, O'Brien, K. R., Torre-Castro, M. de la, Boström, C., et al.
(2017). The fundamental role of ecological feedback mechanisms for the adaptive management of
seagrass ecosystems – a review. Biological Reviews 92, 1521–1538. doi:10.1111/brv.12294.

Mayaud, J. R., and Webb, N. P. (2017). Vegetation in Drylands: Effects on Wind Flow and Aeolian
Sediment Transport. Land 6, 64. doi:10.3390/land6030064.

McLachlan, K. (2014). The dispersal, establishment and management of Ammophila arenaria seed in
coastal foredunes. MSc Thesis, University of Otago, 179 p.

Morais, M. C., Oliveira, P., Marchante, H., Freitas, H. & Marchante, E. (2019). Is richer always better?
Consequences of invaded N-rich soils for the early growth of a native and an invasive species. - Flora
260: 8.

Nolet, C., van Puijenbroek, M., Suomalainen, J., Limpens, J., and Riksen, M. (2018). UAV-imaging to
model growth response of marram grass to sand burial: Implications for coastal dune development.
Aeolian Research 31, 50–61. doi:10.1016/j.aeolia.2017.08.006.

Nunes, LJ.R., Raposo, M. & Gomes, C.J.P. (2020). The Impact of Tourism Activity on Coastal Biodiversity: A Case Study at Praia da Cova Redonda (Algarve—Portugal). Environments 7: 88.

Nye, I. W. B. (1958). The external morphology of some dipterous larvae living in the Gramineae of Britain. Transactions of the Royal Entomological Society 110: 411–487.

Olff H, Huisman J, Van Tooren BF (1993) Species dynamics and nutrient accumulation during early primary succession in coastal sand dunes. Journal Ecology 81:693–706

Pakeman, R. J., Alexander, J., Beaton, J., Brooker, R., Cummins, R., Eastwood, A., et al. (2015). Species composition of coastal dune vegetation in Scotland has proved resistant to climate change over a third of a century. Global Change Biology 21, 3738–3747. doi:10.1111/gcb.12999.

Perrino, E.V., Tomaselli, V., Costa, R., Pavone, P. (2013) Conservation status of habitats (Directive 92/43 EEC) of coastal and low hill belts in a mediterranean biodiversity hot spot (Gargano - Italy). Plant Biosystems 147 (4), 1006-1028. doi:10.1080/11263504.2013.860052.

Piśkiewicz AM, Duyts H, van der Putten WH (2008) Multiple species-specific controls of root-feeding nematodes in natural soils. Soil Biology & Biochemistry 40: 2729–2735.

Piskiewicz AM, Duyts H, van der Putten WH (2009) Soil microorganisms in coastal foredunes control the ectoparasitic root-feeding nematode Tylenchorhynchus ventralis by local interactions. Functional Ecology 23: 621–626.

Puijenbroek, M. E. B., Teichmann, C., Meijdam, N., Oliveras, I., Berendse, F., and Limpens, J. (2017). Does salt stress constrain spatial distribution of dune building grasses Ammophila arenaria and Elytrichia juncea on the beach? Ecology and Evolution 7, 7290–7303. doi:10.1002/ece3.3244.

Pye, K., & Blott, S. J. (2020). Is 're-mobilisation'nature restoration or nature destruction? A commentary. Discussion. Journal of Coastal Conservation, 24(1), 1-4.

Raupach, M. R. (1992). Drag and drag partition on rough surfaces. Boundary-Layer Meteorol 60, 375–395. doi:10.1007/BF00155203.

Reijers, V. C., Hoeks, S., Belzen, J. van, Siteur, K., Rond, A. J. A. de, Ven, C. N. van de, et al. (2021). Sediment availability provokes a shift from Brownian to Lévy-like clonal expansion in a dune building grass. Ecology Letters 24, 258–268. doi:10.1111/ele.13638.

Reijers, V. C., Lammers, C., de Rond, A. J., Hoetjes, S. C., Lamers, L. P., & van der Heide, T. (2020). Resilience of beach grasses along a biogeomorphic successive gradient: resource availability vs. clonal integration. Oecologia, 192(1), 201-212.

Reijers, V. C., Siteur, K., Hoeks, S., van Belzen, J., Borst, A. C. W., Heusinkveld, J. H. T., et al. (2019). A Lévy expansion strategy optimizes early dune building by beach grasses. Nature Communications 10, 2656. doi:10.1038/s41467-019-10699-8.

Roelvink, D., Reniers, A., van Dongeren, A., van Thiel de Vries, J., McCall, R., and Lescinski, J. (2009). Modelling storm impacts on beaches, dunes and barrier islands. Coastal Engineering 56, 1133–1152. doi:10.1016/j.coastaleng.2009.08.006.

Schwarz, C., Brinkkemper, J., and Ruessink, G. (2019). Feedbacks between Biotic and Abiotic Processes Governing the Development of Foredune Blowouts: A Review. Journal of Marine Science and Engineering 7, 2. doi:10.3390/jmse7010002.

Seabloom, E. W., Ruggiero, P., Hacker, S. D., Mull, J., and Zarnetske, P. (2013). Invasive grasses,
climate change, and exposure to storm-wave overtopping in coastal dune ecosystems. Global Change
Biology 19, 824–832. doi:https://doi.org/10.1111/gcb.12078.

Sperandii, M. G., et al. (2018). Hard times for Italian coastal dunes: insights from a diachronic analysis
based on random plots. - Biodivers. Conserv. 27: 633-646.

Strypsteen, G., Houthuys, R., and Rauwoens, P. (2019). Dune Volume Changes at Decadal Timescales
and Its Relation with Potential Aeolian Transport. Journal of Marine Science and Engineering 7, 357.
doi:10.3390/jmse7100357.

Temmerman, S., Meire, P., Bouma, T. J., Herman, P. M. J., Ysebaert, T., and De Vriend, H. J. (2013).
Ecosystem-based coastal defence in the face of global change. Nature 504, 79–83.
doi:10.1038/nature12859.

Van der Putten, W.H., Maas, P.W., Van Gulik, W.J.M. & brinckman, H. (1990). Characterization of soil
organisms involved in the degeneration of Ammophila arenari. Soil Biology and Biochemistry 22: 845-
852

van der Putten WH, van der Stoel CD (1998) Plant parasitic nematodes and spatio-temporal variation
in natural vegetation. Applied Soil Ecology 10: 253–262

van der Putten WH, Vandijk C, Peters BAM (1993) Plant-specific soil-borne diseases contribute to
succession in foredune vegetation. Nature 362: 53–56.

van der Putten WH, Vandijk C, Troelstra SR (1988) Biotic soil factors affecting the growth and
development of Ammophila arenaria. Oecologia 76: 313–320.

Van der Putten, W. H. (1990). Establishment of Ammophila arenaria (marram grass) from culms,
seeds and rhizomes. J Appl Ecol 27: 188–199

van der Putten, W. H., et al. (2005). Invasive plants and their escape from root herbivory: a
worldwide comparison of the root-feeding nematode communities of the dune grass Ammophila
arenaria in natural and introduced ranges. - Biol. Invasions 7: 733-746.

van der Stoel CD, van der Putten WH, Duyts H (2002) Development of a negative plant-soil feedback
in the expansion zone of the clonal grass Ammophila arenaria following root formation and
nematode colonization. Journal of Ecology 90: 978–988.

van Puijenbroek, M. E. B., Limpens, J., de Groot, A. V., Riksen, M. J. P. M., Gleichman, M., Slim, P. A.,
et al. (2017). Embryo dune development drivers: beach morphology, growing season precipitation,
and storms. Earth Surface Processes and Landforms 42, 1733–1744. doi:10.1002/esp.4144.

Van Puijenbroek, M. E. B., Nolet, C., de Groot, A. V., Suomalainen, J. M., Riksen, M. J. P. M., Berendse,
F., and Limpens, J. (2017). Exploring the contributions of vegetation and dune size to early dune
development using unmanned aerial vehicle (UAV) imaging, Biogeosciences, 14, 5533–5549,
https://doi.org/10.5194/bg-14-5533-2017

Van Westen, B., De Vries, S., Reniers, A. J. H. M., Den Bieman, J. P., Hoonhout, B. M., Rauwoens, P., Et
Al. (2019). Aeolian Modelling Of Coastal Landform Development. In Coastal Sediments 2019 (World
Scientific), 1354–1364. Doi:10.1142/9789811204487_0118.

Vandegehuchte, M.L., de la Peña, E. & Bonte, D. (2010a) Aphids on Ammophila arenaria in Belgium:
first reports, phenology and host range expansion. Belgian Journal of Zoology 140: 77-88.

Vandegehuchte, M.L., de la Peña, E. & Bonte, D. (2010b) Interactions between root and shoot
herbivores of Ammophila arenaria in the laboratory do not translate into correlated abundances in
the field. Oikos, 119, 1011–1019.

Vandegehuchte, M.L., de la Peña, E. & Bonte, D. (2010c) Relative Importance of Biotic and Abiotic Soil
Components to Plant Growth and Insect Herbivore Population Dynamics. PLoS ONE 5: e12937

Vandegehuchte, M.L., de la Peña, E. & Bonte, D. (2011) Contrasting covariation of above- and
belowground invertebrate species across plant genotypes. Journal of Animal Ecology, 80, 148–158.

Vandegehuchte, M.L., de la Peña, E., Breyne, P. & Bonte, D. (2012) Non-local genotypes of a resident
grass species reduce invertebrate species richness. Insect Conservation and Diversity 5: 453–460.

Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., Jevrejeva, S., Jackson, L. P., et al.
(2018). Global probabilistic projections of extreme sea levels show intensification of coastal flood
hazard. Nature Communications 9, 2360. doi:10.1038/s41467-018-04692-w.

Webb, C. E., Oliver, I. & Pik, A. J. (2000). Does coastal foredune stabilisation with Ammophila arenaria
restore plant and arthropod communities in Southeastern Australia? Restoration Ecology 8(3): 283-
288.

Wiedemann, A. M. (1987). The ecology of European beachgrass (Ammophila arenaria (L.) Link): a
review of the literature. Technical Report 87-1-01, Oregon Department of Fish and Wildlife, Corvallis,
18p.

Wiedemann, A. M. and Pickart, A. (1996). The Ammophila problem on the Northwest Coast of North
America. - Landsc. Urban Plann. 34: 287-299.

Willis AJ (1965) The influence of mineral nutrients on the growth of Ammophila arenaria. Journal of
Ecology 53: 735–745.

Zarnetske, P. L., Ruggiero, P., Seabloom, E. W., and Hacker, S. D. (2015). Coastal foredune evolution:
the relative influence of vegetation and sand supply in the US Pacific Northwest. Journal of The Royal
Society Interface 12, 20150017. doi:10.1098/rsif.2015.0017.

Zarnetske, P. L., Skelly, D. K., and Urban, M. C. (2012). Biotic Multipliers of Climate Change. Science
336, 1516–1518. doi:10.1126/science.1222732.