Completion of neural crest cell production and emigration is regulated by retinoic acid-dependent inhibition of BMP signaling

Dina Rekler and Chaya Kalcheim·*

Department of Medical Neurobiology, Institute of Medical Research Israel-Canada (IMRIC) and the Edmond and Lily Safra Center for Brain Sciences (ELSC).

Hebrew University of Jerusalem-Hadassah Medical School

Jerusalem 9112102, P.O.Box 12272, Israel

Emails: Dina Rekler, dina.rekler@mail.huji.ac.il; Chaya Kalcheim, kalcheim@cc.huji.ac.il

*To whom correspondence should be addressed.

Running title: Local RA signals the end of neural crest production

Key words: BAMBI, cell specification, dorsal interneurons, Foxd3, Hairy, neural tube, Raldh2, Roof plate, Snai2, Sox9.

Abstract

Production and emigration of neural crest cells is a transient process followed by the emergence

of the definitive roof plate. The mechanisms regulating the end of neural crest ontogeny are

poorly understood. Whereas early crest development is stimulated by mesoderm-derived retinoic

acid, we report that the end of the neural crest period is regulated by retinoic acid synthesized in

the dorsal neural tube. Inhibition of retinoic acid signaling in the neural tube prevents the normal

upregulation of BMP inhibitors in the nascent roof plate and prolongs the period of BMP

responsiveness which otherwise ceases close to roof plate establishment. Consequently, neural

crest production and emigration are extended well into the roof plate stage. Although several roof

plate-specific genes are normally expressed in the absence of retinoic acid signaling, roof plate

and crest markers are co-expressed in single cells and this domain also contains dorsal

interneurons. Hence, the cellular and molecular architecture of the roof plate is compromised.

Collectively, our results demonstrate that neural tube-derived retinoic acid, via inhibition of BMP

signaling, is an essential factor responsible for the end of neural crest generation and the proper

segregation of dorsal neural lineages.

Introduction

To ensure normal development, embryogenesis takes place in a stereotypic sequence and a spatially regulated manner. An example of such a precision is the development of the central and peripheral subdivisions of the nervous system (CNS and PNS), that arise from a common domain in the dorsal neural tube (NT). This domain sequentially generates neural crest cells (NC), progenitors of the peripheral nervous system (PNS)(Le Douarin and Kalcheim, 1999), followed by the definitive roof plate (RP) of the CNS, which is flanked ventrally by dorsal interneuron populations (Andrews et al., 2019; Kalcheim and Kumar, 2017; Krispin et al., 2010).

At trunk levels of the avian axis, NC progenitors sequentially emigrate from the NT for a period of about two days while actively proliferating before and after delamination and largely synchronizing to the S-phase of the cell cycle during the emigration event (Burstyn-Cohen and Kalcheim, 2002). Lineage tracing experiments showed that RP progenitors originate ventral to the cohort of premigratory NC (Krispin et al., 2010b). As a result of continuous cell emigration, RP precursors undergo a ventro-dorsal shift and, upon completion of NC departure, they reach the dorsal midline of the NT and form the definitive RP (Krispin et al., 2010b). RP progenitors gradually exit the cell cycle to become post-mitotic (Kahane and Kalcheim, 1998), and function primarily as a dorsal organizer, providing a gradient of Bone Morphogenetic Protein (BMP) and Wnt activity which is crucial for development of dorsal interneurons (Chizhikov and Millen, 2005; Le Dréau and Martí, 2013; Muroyama et al., 2002; Zechner et al., 2007) and for proliferation of ependymal cells at later stages (Shinozuka et al., 2019; Xing et al., 2018).

Many aspects of early NC ontogeny were extensively studied, from induction to specification, epithelial to mesenchymal transition (EMT) and emigration. Yet, virtually nothing is known about the mechanisms underlying the end of NC production and transition to definitive RP (Rekler and Kalcheim, 2021).

BMP has been found to regulate several aspects of NC development. In the early neurula, BMP

is derived from the non-neural ectoderm and participates in NC induction at the neural plate

border (Endo et al., 2003). Upon closure of the NT, cells of the dorsal domain synthesize and

secrete BMP, which acts in a graded rostro-caudal fashion to activate Wnt signaling and a

network of downstream genes that altogether promote NC EMT and emigration [Reviewed in

(Kalcheim, 2018)].

A first step towards understanding the segregation between NC and RP, was the finding that

approaching the end of the NC production in the avian embryo, the dorsal NT becomes

insensitive to BMP, even though this factor continues to be produced by nascent RP cells. This

suggested that local downregulation of BMP signaling is necessary for the end of NC stage, a

conclusion further illustrated by a premature cessation of NC proliferation and delamination

upon early inhibition of BMP signaling (Nitzan et al., 2016). These events were mediated by

the transcription factor HES1/Hairy1, as early misexpression of *hes1/hairy1* in the dorsal NT

led to a downregulation of the *Alk3* BMP receptor and of pSmad activity (Nitzan et al., 2016).

However, the mechanism that orchestrates these BMP-dependent processes, leading to

cessation of NC production, segregation of NC and RP lineages, and formation of the definitive

RP, remains unknown. Along this line, whether cessation of NC production and RP

development are regulated by the same or by independent mechanisms is yet to be elucidated.

A recent RNA-seq analysis revealed transcriptional heterogeneity between premigratory NC

and definitive RP cells, uncovering a set of genes expressed specifically at the onset of the RP

stage (Ofek et al., 2021). One of these genes is *Raldh2/Aldh1a2*, that encodes an enzyme

responsible for the biosynthesis of the morphogen retinoic acid (RA) (Haselbeck et al., 1999),

central to multiple neurodevelopmental processes (Diez del Corral and Morales, 2014; Lara-

Ramírez et al., 2013; Wilson et al., 2004). RA derived from the *Raldh2*-expressing somitic

mesoderm has been described as a key factor in NT development, including control of

motoneuron specification (Novitch et al., 2003; Wilson et al., 2004) and initiation of NC EMT

and emigration (Martinez-Morales et al., 2011). Although expression of *Raldh2* in the dorsal

NT was previously reported (Berggren et al., 1999; Blentic et al., 2003), its precise temporal

dynamics vis-à-vis NC/RP phases and function/s have not been characterized. Hence, the

discovery of local RA production in the dorsal NT only by the end of the NC stage prompted

us to hypothesize a role for RA in events concerning the late NC.

In this study, we unravel a fundamental role of dorsal NT-derived RA in controlling the end of

NC production. Using two separate tools to inhibit RA signaling in the quail embryo, we

demonstrate that RA is responsible for downregulation of BMP signaling in the dorsal NT. In

the absence of RA signaling, cells in the dorsal NT retain expression of NC markers well into

the RP stage which also displays additional NC traits such as a high mitotic rate and impaired

epithelial properties. Furthermore, we show that lack of RA signaling prolongs cell

delamination into the RP stage. The latter effect is abrogated by concomitant inhibition of BMP

signaling, suggesting that, epistatically, RA acts upstream of BMP signaling. Remarkably,

maintenance of NC traits in the dorsal NT does not prevent the advent of RP traits, or RP-

dependent specification of dorsal interneurons. Rather, in the absence of RA activity, the spatial

and temporal segregation of dorsal lineages, as well as the structural integrity of this signaling

center, are compromised.

Results

RA is both produced and active in the nascent RP

The spatio-temporal dynamics of RA, including synthesis, degradation and activity, was examined in the trunk of quail embryos from stages corresponding to NC production until formation of the nascent RP. RA is produced from retinol via two consecutive reactions: first, retinol is converted to retinaldehyde by RDH enzymes, then it is oxidized to RA by RALDH/ALDH1; additionally, CYP1B1 catalyzes both reactions [Fig. 1A and (Chambers et al., 2007; Kam et al., 2012; Reijntjes et al., 2005; Rhinn and Dollé, 2012). Of the three known RALDH-encoding genes, only *Raldh2* is expressed in the avian trunk at these developmental stages (Blentic et al., 2003). At an early NC stage, *Raldh2* was expressed in the intermediate and paraxial somitic mesoderm but not in the neuroepithelium (Fig. 1B). At an advanced NC stage, *Raldh2* continued to be transcribed in subdomains of the somite-derived sclerotome and dermomyotome (Fig. 1C) and was weakly detected in the dorsal NT (Fig. 1C, arrowhead). Subsequently, *Raldh2* was strongly expressed in the nascent RP itself (Fig. 1D, arrowhead). In contrast, no expression of *Cyp1B1* was apparent in the NT at any stage, although robust transcription was evident in the paraxial mesoderm throughout development (Fig. 1E-G).

RA is degraded by CYP26 enzymes of the cytochrome P450 family. *Cyp26A1* is the only member of the family to be expressed in the avian trunk at early stages of nervous system development (Blentic et al., 2003). At an early NC stage, *Cyp26A1* was absent from the NT and adjacent mesoderm, it was later detected in a salt and pepper pattern in dorsal interneuron progenitors, but was excluded from the dorsal midline (Fig. 1H-I). As the definitive RP formed, it was also occasionally detected in RP cells (Fig. 1J).

CRABP-I is a carrier protein of RA, which participates in regulation of RA function and/or metabolism (Maden et al., 1989; Venepally et al., 1996). At the early NC stage there was no

prominent expression of *CrabpI* in either the NT or paraxial mesoderm (Fig. 1K); at an advanced NC stage it appeared in the basal domain of the NT, corresponding to the positions adopted by differentiating neurons, yet absent from the dorsal and ventral midline regions (Fig. 1L). At the RP stage *CrabpI* was additionally expressed in two stripes at the lateral domain of the RP [(Fig. 1M, and see (Ofek et al., 2021)].

Next, we examined expression of RA receptors. RA acts by binding to a heterodimer of nuclear receptors, RAR and RXR, each of which is encoded by three genetic variants. Expression of all RA receptors was predominantly ubiquitous with at least one variant of each subunit expressed in the dorsal NT at both NC and RP stages (RARα, RARβ and RXRγ, respectively, but not RARγ or RXRα) (Supplem. Fig. S1). The dynamic expression of various components of the RA pathway, suggests that RA acts not only during early NC development, but also during the transition into the nascent RP where a tight regulation of local RA levels may take place.

To directly examine the activity of RA in the dorsal NT, we electroporated a distable GFP plasmid driven by a retinoic acid response element (RARE-d2EGFP) at E2.5 and analyzed GFP fluorescence 12 and 48 hours later, at NC and RP stages, respectively. Whereas control RFP was apparent throughout the dorso-ventral extent of the NTs, RARE activity was primarily present within the dorsal half of the NT at both stages (Fig. 1N-O').

Together, these data show that whereas NC cells respond to mesoderm-derived RA (see Introduction for refs.), but do not synthesize it, the emerging RP both synthesizes and responds to this factor. Hence during the transition from NC to definitive RP, the latter becomes a local source of RA. This prompted us to hypothesize that locally produced RA plays a role in either the end of NC production and emigration, in the ensuing formation of the RP, or in both processes.

Figure 1- RA is both produced and active in the nascent RP

(A) Schematic illustration of the metabolic and signaling pathways of RA.

(B-M) Expression patterns of genes participating in RA synthesis and metabolism, both at early and advanced NC stages (E2 and E3, respectively), and at the RP stage (E4). ISH for *Raldh2/Aldh1a2* (B-D), *Cyp1B1* (E-G), *Cyp26A1* (H-J) and *Crabp1* (K-M).

(N-O') RA activity in the NT at NC and RP stages, indicated by the presence of distable GFP (d2EGFP) driven under the control of a RARE element. Embryos were electroporated at E2.5 along with control RFP to monitor efficiency of transfection, and analyzed 12 hours (N-N') or 48 hours (O-O') later. RARE activity was evident in the dorsal NT of NC-stage embryos (N=8/8), and RP-stage embryos (N=7/7).

Abbreviations, EpS, epithelial somite, No, notochord, NT, neural tube, DM, dermomyotome, Scl, sclerotome, DRG, dorsal root ganglion. Scale bar, 50 μm.

RA is responsible for the loss of sensitivity of dorsal NT progenitors to BMP during the transition from NC to RP

In previous studies, we showed that BMP signaling is essential for NC EMT and emigration (Sela-Donenfeld and Kalcheim, 1999). Subsequently, when approaching the RP stage, although BMP is still produced by the nascent RP, the latter becomes refractory to it, likely via downregulation of BMP receptor 1A (*Alk3*), and upregulation of BMP inhibitory genes (Nitzan et al., 2016; Ofek et al., 2021). The onset of *Raldh2* synthesis close to RP development prompted us to hypothesize that RA, locally produced in the dorsal NT, regulates this dynamic cellular behavior vis-a-vis BMP. To this end, we implemented two independent means to inhibit RA activity *in vivo*: 1) RARα403 - a truncated human RA receptor that lacks the transactivation domain and therefore acts in a dominant-negative fashion (Gupta and Sen, 2015) , and 2) Cyp26A1 which encodes a RA-degrading enzyme.

First, the efficacy of these constructs was tested by co-electroporation with RARE-RFP. RFP intensity was significantly reduced by 79% and 88% in Cyp26A1- and RARα403- electroporated embryos, respectively, when compared to GFP-only controls. Similarly, RARα403 almost completely abolished the RARE-AP signal observed in controls (Supplem. Fig. S2). Thus, both RARα403 and Cyp26A1 can be effectively used to examine the consequences of RA signaling inhibition.

Next, we investigated the effect of RA on the dynamics of BMP signaling. NTs were electroporated at E2.5 (25-27 somite pairs at the level of epithelial somites, NC stage) with either RARα403 or Cyp26A1 and analyzed 2 days later (E4, RP stage) for pSmad1/5/8 expression, a readout of BMP activity. As expected (Nitzan et al., 2016), the RP of control embryos was already devoid of BMP activity at this stage (Fig.2A,A', arrows). In contrast, the RP domain of embryos treated with either RARα403 or Cyp26A1 was pSmad-positive,

indicating a failure to normally downregulate BMP signaling (Fig. 2B-C', arrowheads). The

dI1 progenitor population of dorsal interneurons, located ventral to the RP, was also pSmad-

positive, consistent with its dependency on BMP for proper development (Duval et al., 2019;

Lee et al., 1998; Tozer et al., 2013; Wine-Lee et al., 2004). Quantification of pSmad intensity

confirmed a significant increase in Cyp26A1 and RARα403-treated embryos compared to

controls. This was apparent even if measurement of pSmad intensity comprised the dorsal NT

from its midline to the ventral border of pSmad+ staining, that included dI1 progenitors (Fig.

2D). These results show that endogenous RA activity accounts for the downregulation of BMP

signaling in the dorsal NT during the transition from NC to RP.

We next examined whether the observed RA-dependent downregulation of BMP activity is

mediated by a corresponding upregulation of BMP inhibitors at the RP stage. In this context,

Hairy1 was previously reported to act as a specific RP-derived BMP inhibitor thus participating

in the end of NC production (Nitzan et al., 2016). Furthermore *hairy1/qHes4*, and the known

BMP inhibitors *BAMBI* (Brazil et al., 2015) and *Grem1* (Sun et al., 2006), were unraveled as

specific RP markers in a recent RNA-seq analysis comparing dorsal NT cells at both

premigratory NC and RP stages (Ofek et al., 2021). Embryos electroporated with either

Cyp26A1 or RARα403 at E2.5 exhibited a substantially decreased intensity of *BAMBI* mRNA

staining in the RP relative to controls (Fig. 2E-H). Similarly, RARα403 significantly affected

the upregulation of both *hairy1* and *Gem1* in RP when compared to control embryos (Fig. 2I-

N). These results show that the downregulation of BMP signaling by RA is accounted for, at

least partially, by upregulation of BMP inhibitory genes.

In a previous study, we reported that BMP-dependent emigration of NC cells is mediated by

Wnt activity (Burstyn-Cohen et al., 2004), raising the possibility that, in the absence of RA,

enhanced Wnt activity is associated with extended BMP signaling at the RP stage. To this end,

we assessed the effect of RA inhibition on Wnt signaling in RP by electroporating a distable

Wnt reporter, 12XTopFlashd2GFP, together with control RFP or RARα403/RFP. Whereas

very low 12XTopFlashd2GFP expression was apparent under control conditions, significantly

enhanced expression in the RP was monitored in the RARα403-transfected NTs (Supplem.

Fig. S3). Together, these results suggest an involvement of RA activity in the downregulation

of BMP and of downstream Wnt signaling towards ending NC production and/or emigration.

Figure 2- <u>RA is responsible for the loss of sensitivity of dorsal NT progenitors to BMP during the transition from NC to RP.</u>

(A-C') Immunostaining for pSmad1/5/8. Embryos at E2.5 were co-electroporated with RARα403, Cyp26A1 or with a control PCAGG vector, along with GFP, and analyzed at E4. Note absence of BMP activity in the RP of control embryos (N=8, A-A') whereas only ventrally located dorsal interneurons are positive, while the RP domain itself is positive in Cyp26A1 (N=8) and RARα403-electroporated embryos (N=7, B-B',C-C'). (D) Quantification of pSmad staining intensity (measured area includes dorsal interneurons). Four-to-21 sections per embryo were analyzed. *p < 0.05, **p < 0.005 , one-way ANOVA with post-hoc Tukey's tests.

(E-N) In-situ hybridization of embryos electroporated with Cyp26a1 or RARα403 at E2.5 and analyzed at E4. Asterisks mark the RP domain.

(E-H) ISH for *BAMBI*, showing downregulation in the RP of Cyp26A1- and RARα403-treated embryos. (H) For quantification of the intensity of *BAMBI*, 6-to-21 sections per embryo were analyzed. N = 7,7 and 6 embryos for control, Cyp26A1 and RARα403, respectively. *p < 0.05, ****p < 0.0001, one-way ANOVA with post-hoc Tukey's tests.

(I-J, M) ISH for *hairy1*, showing downregulation in the RP of RARα403-treated embryos. (M) For quantification of the intensity of *hairy1,* 7-18 sections per embryo were analyzed. N = 7 embryos for each group. **p < 0.005, Student's unpaired t-test.

(K-L,N) ISH for *Grem1*, showing downregulation in the RP of RARα403-treated embryos. For quantification of the intensity of *Grem1*, 11-27 sections per embryo were analyzed. N= 8 embryos for each group. **p < 0.005 via Student's unpaired t-test. Abbreviations, dI1, dorsal interneurons 1, RP, roof plate. Scale bar, 50 μm.

RA downregulates NC markers during the transition to definitive RP

Our results prompted us to investigate whether the observed persistence of BMP and Wnt

activity into the RP stage is associated with a retention of NC identity.

Foxd3, *snai2* and *sox9* are known premigratory markers of trunk NC cells (Nieto et al., 1994;

Sela-Donenfeld and Kalcheim, 1999), and are downregulated in the dorsal NT upon emigration

of neural progenitors, being absent from prospective melanoblasts and RP (Krispin et al.,

2010b; Nitzan et al., 2012; Ofek et al., 2021). Expression of *foxd3*, *snai2* and *sox9* transcripts

was examined at the RP stage in embryos electroporated with RARα403. Strikingly, whereas

no mRNA expression was detected in controls, all three genes persisted in the dorsal NT of

treated embryos (Fig. 3A-F', arrowheads). Moreover, using immunohistochemistry to assess

Sox9 protein, we confirmed its maintenance in the dorsal NT at the RP stage both in Cyp26A1-

and RARα403-electroporated embryos (Fig. 3G-J).

These findings suggest that endogenous RA signaling underlies the loss of NC identity towards

the end of the NC period.

Figure 3- RA downregulates NC markers during the transition to definitive RP

(A-F') ISH for neural crest markers on embryos electroporated with GFP along with control PCAGG or RARα403 at E2.5 and analyzed at E4. Arrowheads point to the RP in RARα403-treated embryos. (A-B') ISH for *foxd3*, showing no expression in the RP of control embryos (N= 7) compared to strong expression in the RP of all RARα403-treated embryos (N = 7). (C-D') ISH for *snai2*, showing no expression in the RP of control embryos (N = 11) and clear expression in the RP of all RARα403-treated embryos (N = 11). (E-F) ISH for *sox9*, showing no expression in the RP of control embryos (N = 9) and strong expression in the RP of all RARα403-treated embryos (N = 9). Note as well expression of *sox9* mRNA in more ventral regions of the NT and in adjacent sclerotome.

(G-J) Immunostaining for Sox9 in control and experimental embryos, showing the presence of Sox9 protein in the RP of Cyp26A1- and RARα403-treated embryos (arrowheads), compared to absence of Sox9 in the RP of control embryos. RFP was co-electroporated in both groups. Dashed lines delineate the RP domain. (J) Measurement of Sox9 staining intensity. For quantification, 15 sections per embryo were analyzed. N = 10, 7, and 11 embryos for control, Cyp26A1 and RARα403 groups, respectively. $*p < 0.05$, $****p < 0.0001$, one-way ANOVA with post-hoc Tukey's tests. Abbreviations, NT, neural tube, DRG, dorsal root ganglion. Scale bar, 50 μm.

RA signaling promotes cell cycle exit and epithelial traits in dorsal NT cells

NC cells proliferate extensively while residing in the NT and also during migration. Upon the end of NC production, dorsal NT progenitors gradually exit from the cell cycle to become post-mitotic RP cells (Kahane and Kalcheim, 1998; Nitzan et al., 2016). We asked whether RA inhibition affects cell cycle withdrawal of nascent RP cells. To this end, we immunostained embryos at E4 (RP stage) for the mitotic marker phosphohistone H3 (pH3), following electroporation of RARα403 or Cyp26A1 at E2.5 (NC stage). Remarkably, a significantly higher number of mitotic nuclei was observed in the RP of embryos treated with either Cyp26A1 or RARα403 compared to controls which exhibited a low extent of mitosis (Fig. 4A-D). These results suggest that endogenous RA inhibits cell proliferation, consistent with previous data obtained in forebrain and optic tectum of avian embryos (Gupta and Sen, 2015; Kukreja et al., 2020).

Furthermore, characteristic of post-mitotic, differentiating progenitors is the localization of nuclei to the basal domain of the neuroepithelium, as observed in control embryos at the RP stage (Fig. 4E). In contrast, in RARα403-electroporated embryos, nuclei were spread throughout its apico-basal extent. Quantification showed that their localization to the apical half of the epithelium significantly exceeds that seen in control conditions (Fig. 4E-G). This is consistent with the higher extent of cell proliferation monitored in the absence of local RA signaling, reflecting the retention of interkinetic nuclear migration with presence of mitotic nuclei in the ventricular zone, and/or a disorganization of the nascent RP under experimental conditions.

Another critical process that occurs following the end of NC emigration is the restoration of epithelial cell traits in the nascent RP (Nitzan et al., 2016). One of these traits is the deposition of a continuous and prominent laminin-containing basal lamina covering the normal RP (Fig.

4H,H'). This basal lamina is incomplete at the NC stage due to continuous cell delamination. Inhibition of RA signaling resulted in a weaker and discontinuous pattern of laminin immunostaining dorsal to the RP (Fig. 4I-J), indicative of the maintenance of NC characteristics at the later RP stage. Consistently, upon electroporation with membrane-tethered GFP, it was apparent in controls that cells extended along the apico-basal domain of the RP (Fig. 4H). In contrast, in the absence of RA activity, many cells were devoid of epithelial integrity, including labeled cells detaching from the epithelium (Fig.4I, arrows). Combined, these results show that inhibition of RA signaling in the dorsal NT prevents cells from exiting the cell cycle and from adopting cellular traits that characterize the normal ontogeny of a RP.

Figure 4- RA signaling promotes cell cycle exit and epithelial traits in dorsal NT cells

(A-D) Immunostaining for phospho-Histone H3 of embryos electroporated at E2.5 with either control PCAGG, RARα403 or Cyp26A1, and analyzed at E4. Arrowheads in B-C' point to pH3$^+$ cells in the RP domain of RARα403- and Cyp26A1- electroporated embryos. The quantification of pH3$^+$ cells in the RP is represented in (D). 120 sections were analyzed per embryo. N = 20, 9 and 9 embryos for control, RARα403 and Cyp26A1, respectively. **p < 0.005, ****p < 0.0001, one-way ANOVA with post-hoc Tukey's tests.

(E-G) Apico-basal localization of Hoechst-stained nuclei in embryos electroporated as in A-D. Whereas in control embryos most nuclei are restricted to the basal domain of the RP, in the RP of RARα403-treated embryos, nuclei are apparent across the entire thickness of the tissue. Confocal images were z-stacked and nuclei in the apical and basal halves of the RP domain were counted. The proportion of apically-localized nuclei is represented in (G). Thirteen-to-17 sections were analyzed per embryo. N = 10 and 12 embryos for control and RARα403 groups, respectively, ****p < 0.0001, exact p-value Mann-Whitney test.

(H-J) Immunostaining for Laminin of embryos electroporated as in A-D. Arrows and arrowheads in insets of H' and I' indicate the presence of either a continuous or a discontinuous RP basal lamina, respectively. Arrows in I point to delaminating cells. (J) Quantification of Laminin intensity along the basal aspect of the RP. Six-to-18 sections were analyzed per embryo. N = 9 embryos both for control and RARα403 treatments. *p < 0.05, Student's unpaired t-test. Abbreviations, NT, neural tube. Scale bar, (A-C), 50 μm, (E-F, H-I), 10 μm.

RA is responsible for the end of NC delamination

As described above, inhibiting RA activity in the dorsal NT extends the period of BMP activity and preserves NC properties such as premigratory genes and cell proliferation into the RP stage. We therefore asked whether the most discernible difference between NC cells and definitive RP cells, the process of EMT and cell delamination, is prolonged as well by manipulation of RA signaling.

To this end, we implemented a double electroporation protocol: first, we electroporated RARα403 along with RFP only to one hemi-NT at E2 and at E3.5 both sides were labeled with GFP-DNA (Fig.5A). By E4, RFP-positive NC cells exited the NTs under both control and treated conditions primarily towards the side ipsilateral to the transfection and also to some extent towards the contralateral side, as previously described (Burstyn-Cohen and Kalcheim, 2002; George et al., 2007). However, the number of emigrated, late GFP-labeled cells was significantly higher on the RARα403-electroporated side when compared to the control contralateral side and also to both sides of control embryos (Fig. 5B-D, arrowheads in C'). These data show that RA signaling promotes the end of NC cell delamination.

Next, we examined the relationship between RA and BMP activities in the context of NC EMT. We hypothesized that, if the end of NC EMT is mediated by RA-dependent inhibition of BMP signaling, the continuous NC emigration observed at the RP stage in response to RARα403, should be prevented by inhibiting BMP with Smad6 (Nitzan et al., 2016).

Embryos aged E2 were bilaterally electroporated (Fig. 5A, bottom) with either control GFP, RARα403 to induce late NC delamination (Fig. 5B-D), with Smad6, or with both RARα403 and Smad6. At E3.5, control GFP was bilaterally delivered to all embryos and the outcome was analyzed at E4 (Fig. 5A). In all cases, RFP-labeled cells that indicate the efficiency of early transfections, successfully delaminated from the NT. In contrast, under control conditions, few

or no GFP-labeled cells delaminated from the NT (Fig.5E-E',I), consistent with the second electroporation being done following the end of NC emigration. As also shown above, in RARα403-treated embryos, a substantial number of GFP-labeled cells had emigrated (Fig. 5F-F',I and see also D). Treatment with Smad6 alone resembled the control situation with few delaminated GFP+ cells at the RP stage (Fig. 5G-G',I), and this effect was not different upon co-transfection of Smad6 and RARα403 (Fig. 5H-H',I). Together, these results show that RA acts upstream of BMP.

Next, we examined whether the late-delaminating cells exhibit a NC identity. For this purpose, embryos double electroporated as described in Fig 5A were immunostained for the NC marker HNK1. While in control embryos no GFP-labeled cells were apparent outside the NT, and HNK1 stained only NC derivatives such as dorsal ganglia and ventral roots, in RARα403-treated embryos a considerable number of GFP+HNK1+ cells was also detected in the mesenchyme near the NT (Supplem. Fig. S4A-B", arrowheads in B-B"). Remarkably, $95 \pm 0.8\%$ of the late-delaminating GFP-labeled cells were HNK1$^+$ (Fig. S4C), confirming their NC nature.

In addition, we examined the pattern of *foxd3*, which is expressed both in premigratory and migrating NC cells. Embryos were electroporated with RARα403 along with GFP at E2.5 and analyzed at E4. Control NTs were devoid of *foxd3* mRNA in the RP and mRNA signal was only detected in NC derivatives and dI2/V1 interneurons. In RARα403-treated embryos *foxd3* expression was evident in the dorsal NT (see also Fig. 3), and also in NC cells that presumably emigrated recently (Fig S4D-E', arrowheads), further confirming their NC identity.

Thus, local RA inhibition in the NT extends the period of NC production and further delamination into the RP stage. Collectively, these results point to RA being a major regulator of the end of the NC stage.

Figure 5- RA is responsible for the end of NC delamination

(A)Schematic representation of the experimental design for detection of late-emigrating cells (see text for details). (B-D) Embryos electroporated unilaterally at E2 with either control or RARα403 plasmids, followed by a second bilateral electroporation at E3.5 with control GFP. Emigration of GFP[+] cells was compared between electroporated and non-electroporated sides of control and experimental embryos (D). RFP was delivered with the early electroporation to monitor transfection efficiency. Images presented are adjacent to the rostral somitic domain. Whereas few GFP[+] cells delaminated at the RP stage in controls, C' and inset show late-delaminating GFP[+] cells ipsilateral to the early RARα403 electroporation (arrowheads). (D) Quantification of GFP[+] cells that emigrated from either intact or electroporated (EP) sides of control and experimental embryos. Seventy two sections per embryo were analyzed at an

equivalent axial level. N = 10 and 11 embryos for control and RARα403 groups, respectively.
Statistical tests applied were exact P-value Wilcoxon signed ranks paired test to compare
between EP and non-EP sides of embryos inside each experimental group, and exact P-value
Mann-Whitney test to compare between non-EP sides of control and RARα403 groups. ***p
< 0.001.

(E-I) Embryos were electroporated bilaterally at E2 with control, RARα403, Smad6 or
RARα403+Smad6 constructs along with RFP. At E3.5, bilateral electroporations with GFP
were performed and embryos were analyzed for emigration of GFP$^+$ cells at E4. Images
presented are opposite a rostral somitic domain. Note the presence of emigrated RFP+ cells in
all treatments. Notice as well the presence of GFP$^+$ delaminated cells in F,F', while virtually
no GFP$^+$ cells are seen outside of the NT in E,E', G,G' and H,H'. (I) Quantification of
delaminated GFP$^+$ cells. Seventy sections per embryo were analyzed at an equivalent axial
level. N = 21, 19, 6 and 12 embryos were monitored for control, RARα403, Smad6 and
Smad6+RARα403 groups, respectively. *p < 0.05, ****p < 0.0001, approximate P-value
Kruskal-Wallis with post-hoc Dunn's test. Abbreviations, ns, not significant; NT, neural tube,
DRG, dorsal root ganglion. Scale bar, 50μm.

Inhibition of RA signaling downregulates only a subset of RP markers

A major question that remains to be addressed is whether the end of the NC stage is sufficient for signaling the formation of the definitive RP, or whether these are two independently regulated processes. Given that RA deficiency delays the end of the NC stage, we asked whether this occurs at the expense of timely RP development.

As recently shown (Ofek et al., 2021), there are differences in the gene expression profile between premigratory NC cells and definitive RP cells. We selected a number of RP-specific genes that are not expressed in the dorsal NT during the NC stage and examined the effect of RA downregulation on their expression. As described in Fig. 2, expression of RP-specific BMP inhibitors was strongly reduced. Likewise, mRNA encoding *slit1*, a chemorepellent of post-floor plate commissural axons (Kidd et al., 1999) was also decreased under experimental conditions (Supplem. Fig. S5). In contrast, additional RP markers, such as *Rspo1, NDP (norrin)* or *draxin*, were normally expressed (Supplem. Fig. S5), suggesting that development of a number of RP traits is independent of RA signaling.

Since both NC and RP markers are present at the RP stage upon inhibition of RA signaling (Figs.3, Supplem Fig. S5), we proceeded to clarify whether NC and RP markers are expressed in the same cells, or, alternatively, whether there is a mix of different cell types in the RP of RARα403-treated embryos. First, we performed ISH of *Rspo1* (RP marker) and *foxd3* (NC marker) on adjacent sections of RP-stage embryos electroporated at E2.5 with either RARα403 or control GFP (Fig. 6). Whereas in control NTs, only *Rspo1* was apparent in the RP, both *Rspo1* and *foxd3* were detected in an overlapping pattern in the RP of the treated embryos (Fig. 6A-B''').

In addition, we combined a fluorescent ISH of *Rspo1* with immunostaining for SNAI2 or Sox9 in the same sections. Control NTs exhibited SNAI2 and Sox9$^+$ nuclei only outside the *Rspo1$^+$*

RP (Fig.6C-C'', E-E", see also Fig. 3G-I). In striking contrast, both markers were detected in

the same dorsal NT cells in the absence of local RA signaling (Fig.6D-D", F-G", H,I). In some

instances, $Rspo1^+$Sox9$^+$ cells were also noticed undergoing delamination from the NT (Fig.

6G-G", arrowhead).

Together, these data suggest that the loss of RA signaling does not totally prevent the advent

of RP markers implying that loss of NC identity and gain of RP fate are independently-

regulated processes. Nevertheless, timely RA activity in the dorsal NT is required for the proper

segregation of NC from RP fates.

Figure 6- Overlapping expression in RP of both NC and RP markers upon inhibition of RA signaling

(A-B''') Embryos co-electroporated at E2.5 with either control PCAGG or RARα403 along with a GFP plasmid. Embryos were fixed at E4, and adjacent sections were in-situ hybridized for *foxd3* or *Rspo1* and then superimposed to reveal overlapping domains of gene expression. Dotted lines delineate the *Rspo1* expression domain. In control embryos, no *foxd3* expression was detected in the RP (N = 2, and see also Fig. 3). In RARα403-treated embryos, overlapping expression was detected in all 4 embryos examined.

(C-D") Immunostaining for SNAI2 and fluorescent ISH for *Rspo1* were combined on the same sections of embryos electroporated at E2.5 with either control PCAGG or RARα403 and analyzed at E4. Dashed lines mark the *Rspo1* expression domain. Note the presence of SNAI2$^+$*Rspo1*$^+$ cells in the RP in D". (E-G") Immunostaining for Sox9 and fluorescent ISH for *Rspo1* were performed as above. Note the presence of Sox9$^+$*Rspo1*$^+$

cells in the RP in F" and G''. Additionally, delaminating Sox9$^+$*Rspo1*$^+$ cells (arrowheads in G-G") were apparent under experimental conditions. (H,I) Quantification of SNAI2$^+$*Rspo1*$^+$ and Sox9$^+$*Rspo1*$^+$ cells in the RP. N = 8 and 8 for SNAI2 in controls and RARα403. N = 9 and 10 embryos for control and RARα403 groups stained with Sox9, respectively. ****p < 0.0001, Welch's t-test. Abbreviations, NT, neural tube. Scale bar, 20 μm.

RP-derived dI1 interneuron development only partly depends on RA signaling

We further investigated whether depletion of RA signaling had any effect on the role of the RP as a signaling center for dorsal interneuron development. We immunostained RARα403 and control-electroporated embryos with the dI1 interneuron marker BarHL1, and analyzed the number of BarHL1$^+$ cells and their spatial distribution. Whereas no significant difference in the number of BarHL1$^+$ interneurons could be monitored between control and treated embryos (Fig. 7A), there was a collective dorsal shift in their position in the RARα403-treated NTs, with some of BarHL1$^+$ cells residing in the RP domain itself (Fig. 7B-E', arrowheads).

Thus, whereas inhibition of RA signaling does not affect RP-mediated formation of dI1 interneurons, it interferes with the spatial information conveyed onto dorsal interneurons from the RP, likely by impairing the formation of the boundary between the RP and more ventral domains. These findings further indicate that RA signaling not necessarily influences development of RP properties but rather impairs temporal and spatial segregation between dorsal cell lineages.

Figure 7- RP-derived dI1 interneuron development only partly depends on RA signaling

(A) Quantification of BarHL1[+] dI1 interneurons in E4 embryos, electroporated at E2.5 with either control PCAGG or RARα403. No significant difference in the number of BarHL1[+] neurons was monitored. Ten-to-14 sections were analyzed per embryo; N= 9 and 8 embryos for control and RARα403 groups, respectively. Student's unpaired t-test was applied.

(B, C) Spatial distribution of BarHL1[+] cells measured as the distance from the dorsal midline. 5126 cells were counted in the control group (green), and 3886 in the RARα403 group (purple). The area of overlap of control and experimental cells is in blue. Note the dorsal shift of cells in the RARα403-treated group when compared to controls. In (B) all the cells are included, while in (C) only cells included in a 30 μm distance from the dorsal midline, comprising the RP region, are presented.

(D-E') Confocal images of BarHL1-stained sections. Nuclear GFP (nucGFP) marks the electroporated domain. Note in E and E' the presence of BarHL1[+] interneurons inside the RP domain (arrowheads) compared to controls in which these neurons are localized ventral to the RP. Nuclei are visualized with Hoechst. Abbreviation, NT, neural tube. Scale bar, 50 μm.

The dorsal NT as the source of RA responsible for NC to RP transition

As shown in Fig. 1, the dorsal NT becomes a source of RA during the period marking the end of NC emigration and transition to the RP stage, suggesting this is the local source of RA that underlies the effects demonstrated here. Yet, at the same time, RA is still produced in the adjacent paraxial mesoderm. To begin addressing whether the latter tissue accounts for the effects observed, we separated the paraxial mesoderm from the NT with an impermeable membrane and assessed the development of the RP. A unilateral slit was performed at the epithelial somite stage into which a piece of aluminum foil was inserted (Fig. 8A). We analyzed RP formation using *BAMBI* expression as a sensitive readout for normal RP ontogeny that is downregulated in the absence of RA signaling (Fig.2). Notably, *BAMBI* remained strongly expressed in the RP of treated embryos to the same extent as the control sides, despite separation from the mesoderm and virtual disappearance of the somitic tissue that lost contact with the NT (Fig. 8A-C'). To exclude the possibility that RA from the mesoderm of the non-separated side is sufficient to induce normal RP development, a slit was made along the dorsal midline of the NT following insertion of the barrier, thus opening the NT and creating a physical gap between both halves of the neuroepithelium (Fig. 8D). A comparable expression of *BAMBI* mRNA on both hemi-RPs was still apparent under these conditions (Fig.8E-F"), suggesting that the paraxial mesoderm is not the source of RA important for NC to RP transition.

Figure 8- Somitic RA is dispensable for NC-to-RP transition

(A, D) Schematic representations of the experimental design to mechanically separate the NT from somitic mesoderm using an impermeable aluminum foil (see Methods for details).

(B-C', E-F') ISH for *BAMBI* on sections of embryos with intact (B-C') or split (E-F") dorsal NTs following grafting of a unilateral barrier between NT and somites. Representative images from two different embryos are presented for each treatment. *BAMBI* expression remained unchanged in the RP opposite to the operated sides in 7/7 embryos with an intact RP, and in 4/4 embryos with a split RP. Note as well that distal to the barriers, most of the somitic mesoderm disappeared (B,C,E). In some instances, the somatic layer of the lateral plate mesoderm approached the barrier (F). Abbreviations, DM, dermomyotome, DRG, dorsal root ganglion, NT, neural tube. Scale bar, 50 μm.

Figure 9- Schematic diagram summarizing the role of RA in the transition from NC to RP

(A) At the NC stage,
premigratory progenitors
residing in the dorsal NT
(green) exhibit BMP
signaling activity, express
specific NC markers (e.g.
foxd3, *snai2* and *sox9*,
green in D) and proliferate
extensively. Following a
process of EMT, NC cells
delaminate from the NT
through a discontinuous
basement membrane
(dashed line) to migrate
away. At the NC stage, RA
is supplied to the NT from
the paraxial mesoderm,
where it is produced by
Raldh2 (blue).

(B) Under normal
conditions, upon cessation
of NC production and
emigration, a definitive
RP forms. RP cells (purple) are refractory to BMP signaling, and exhibit a different gene
expression profile when compared to NC cells: they downregulate *foxd3*, *snai2* and *sox9* and
upregulate RP-specific genes (purple in D), among which are known inhibitors of BMP
signaling. RP cells gradually exit the cell cycle and do not delaminate, regenerating a
continuous basement membrane and epithelial traits. During the transition between NC and RP
stages, *Raldh2* expression begins in the dorsal NT (blue in RP), in addition to RA still produced
in somites. The RP functions as a signaling center, regulating specification and/or
differentiation of dorsal interneurons (red, dI1 progenitors) that localize ventral to the RP.

(C) In the present study we demonstrate that inhibition of RA signaling in the dorsal NT,
impairs the end of NC production and consequently, the normal transition between NC and RP
stages. First, under experimental conditions, dorsal NT cells at the RP stage retain active BMP
signaling likely via downregulation of BMP antagonists (*BAMBI, hairy1, Grem1*) otherwise
selectively expressed in RP. Second, they express both NC and RP markers (illustrated as
green/purple cells), Third, they functionally behave as NC cells, maintaining mitotic activity,
impaired epithelial traits (such as a discontinuous basement membrane, dashed line), and
continuous emigration from the NT. Fourth, although the advent of several RP-specific markers
is normal (e.g; *Rspo1, NDP, draxin*) and the number of dI1 progenitors is unchanged, the spatial
segregation between NC, RP and dI1 interneuron lineages is compromised.

(D) A selection of differential traits displayed by NC progenitors (green), normal RP cells
(purple), as well as of "RP" cells formed upon inhibition of RA signaling (green/purple).
Selected NC genes appear in green and RP-specific genes in purple. (+) and (-) depict
expression or lack of expression of relevant genes, respectively. Light blue color indicates
sites of RA production (i.e. *Raldh2* expression). Red circles represent dI1 progenitors.

Discussion

Here we show that dorsal NT-derived RA controls the end of NC production and emigration (Fig.9). This effect is mediated through BMP signaling, at least via the upregulation of BMP inhibitors in the nascent RP, which prevent persistent BMP activity and ensures the timely cessation of NC production (Fig.9). We previously reported that BMP produced in the dorsal NT is an essential inducer of NC EMT and emigration; a graded activity of BMP is created along the rostro-caudal axis of the embryo thanks to a counter-gradient of its inhibitor noggin. By the time of initial somite dissociation, noggin expression is downregulated by somite derived factors (Sela-Donenfeld and Kalcheim, 1999; Sela-Donenfeld and Kalcheim, 2002), including an interplay between mesodermal RA and Fgf8, that affect the onset but not the maintenance of NC delamination (Martínez-Morales et al., 2011). Our findings are consistent with previous results reporting that RA repressed BMP signal in P19 embryonic carcinoma cells; in vivo, RA antagonized BMP-regulated differentiation and proliferation of neural progenitor cells in the NT (Sheng et al., 2010). Likewise, RA was also found to inhibit Wnt signaling (Shaker et al., 2020), that in the present context acts downstream of BMP and whose activity was enhanced by loss of RA signaling.

In normal development, the end of NC emigration is followed by the advent of the definitive RP of the CNS (Rekler and Kalcheim, 2021). This is associated with a progressive withdrawal from the cell cycle, the regeneration of a complete basal lamina overlying the dorsal NT, of apico-basal properties of the RP epithelium and the corresponding downregulation and upregulation, respectively, of NC and RP-specific genes (Nitzan et al., 2016; Ofek et al., 2021). We report that, in the absence of RA activity, there is a continuity of several molecular and cellular properties of the NC well into the RP stage, including the presence of a subset of RP

cells that co-expresses NC markers. Together, lack of local RA activity blurs the spatial and

temporal segregation between NC and RP (Fig.9).

Despite these phenotypes, it is notable that RP-specific markers such as *Rspo1*, *NDP* or *draxin*,

were not affected, suggesting that a RP is able to develop even if RA activity is inhibited. This

raises the fundamental question whether the end of NC formation and the ensuing development

of a RP are regulated by the same or by independent mechanisms. Our results would suggest

that the mechanisms accounting for the cessation of NC phase are necessary but not sufficient

to explain all aspects of RP development. Consistent with this notion, spatio-temporal lineage

tracings revealed that RP segregate from neural progenitors of the NC before the latter

complete emigration (Krispin et al., 2010b; Nitzan et al., 2012). In addition, gain and loss of

Notch function in avians and mice, respectively, resulted in changes in RP formation with no

apparent effects on NC cell behavior (Ofek et al., 2021). This would indicate that although NC

and RP lineages arise from common progenitors (Nitzan et al., 2012) by different exposure

times and/or levels of BMP signaling (Nitzan et al., 2016; Timmer et al., 2002; Tozer et al.,

2013), they segregate rather early from each other.

Among the RP genes whose expression is perturbed in the absence of RA signaling, we found

BAMBI, hairy1 and *Gremlin*, none of which is expressed in the dorsal NT at NC stages. These

genes, whose transcription was prevented by treatment with RARα403 and/or Cyp26, encode

inhibitors of BMP signaling. Hence, RA-sensitive aspects of RP development are linked to its

regulation of BMP activity. Moreover, since RA-dependent downregulation of BMP activity

accounts for only selected facets of RP ontogeny, we posit that RP development comprises

both RA-dependent and RA-independent mechanisms.

We observed that maintenance of NC traits in RARα403-treated NTs did not interfere with

differentiation of a normal number of RP-dependent dorsal interneurons of type 1, suggesting

that the role of the RP as an interneuron inducer center was not altered. However, their

topographical dorsoventral distribution was abnormal as instead of organizing into discrete

nuclei, interneurons redistributed along the basal surface of the NT comprising the RP itself,

further demonstrating that the structural integrity of this signaling center and/or of its

boundaries is compromised; this is consistent with the observed local changes in epithelial

architecture, cell cycle properties and basal lamina deposition.

Since dI1 interneurons are also specified by BMP signaling (Chesnutt et al., 2004; Hegarty et

al., 2013; Timmer et al., 2002), it is conceivable to speculate that maintenance of BMP

responsiveness in the RP of RARα403-treated NTs affects the normal dorso-ventral gradient

of BMP activity (Tozer et al., 2013), perhaps eliciting indirectly an attractive function that

enables dI1 interneurons to migrate into the RP domain. In this context, RP-derived BMP

exhibits at a later stage chemorepellent properties vis-à-vis commissural axons (Augsburger et

al., 1999; Butler and Dodd, 2003), putting forward the additional possibility of guidance

functions for this morphogen under the present experimental conditions. One of the genes

downregulated upon abrogation of RA function is *Slit1*, a known repellent of axonal guidance

(Kidd et al., 1999), as well as of cell migration (Halperin-Barlev and Kalcheim, 2011;

Theveneau and Mayor, 2012). Loss of *Slit1* could also account for the dorsal shift of dI1

interneurons into the RP domain, and possible interactions with BMP remain to be examined,

It is worth mentioning that the NT is sensitive to RA throughout development, with RA playing

various functions at different stages [this study, (Diez del Corral and Morales, 2014; Martinez-

Morales et al., 2011; Wilson et al., 2004)]. Hence, it is plausible that the mechanisms mediating

these effects differ as a function of time and context. Here we report that precisely before the

end of NC delamination, RA begins to be synthesized in the dorsal NT (e.g; *Raldh2* expression)

in addition to the paraxial mesoderm, raising the notion that this new, local source of RA

accounts for the end of NC development. Notably, *Raldh2* is also synthesized in the RP of mouse embryos (Ofek et al., 2021). Unfortunately, available mutants of *Raldh2* are embryonic lethal at about E8.75 (Niederreither et al., 1999), precluding a focal analysis of the transition from NC to RP. In this sense, the avian embryo provides an excellent model to address events restricted both in time and space. The two methods we implemented to abrogate RA signaling in the dorsal NT act primarily in a cell autonomous fashion. However, it is still possible that given the secreted nature of the ligand, Cyp26A1 and/or RARα403 missexpression in the dorsal NT also provide a sink for somitic RA. Despite robust separation of the mesoderm from the NT, expression of *BAMBI*, a most sensitive and significant readout of RA activity, was not altered, suggesting that mesodermal RA is dispensable. What accounts for this shift in responsiveness from a mesodermal towards a neural source of factor? First, local RA could provide a stronger and more efficient signal to target progenitors given the increasing size of the embryo and the development of a continuous basement membrane dorsal to the NT. Second, cofactors of RA such as *Crabp1*, could change the nature of the cellular response upon late appearance at the RP stage. Together, changes in the molecular landscape between NC and RP stages (Ofek et al., 2021), that partly depend on local RA activity, may alter the nature of the interactions between RA and BMP/Wnt to signal a timely cessation of NC production. Along this line, the present study provides a mechanistic explanation for the separation between PNS (NC) and CNS (RP and dI1) lineages during development.

 Materials and Methods

 Embryos

Fertilized quail (*Coturnix coturnix Japonica*) eggs were obtained from commercial sources

(Moshav Mata), kept at 15°C and then incubated at 38 °C to desired stages. Embryos were

staged by the number of somite pairs formed. All experiments were conducted at the flank level

of the axis.

Expression vectors and *in ovo* electroporation

The following expression vectors were used as controls: pCAGG, pCAGGS-EGFP (Krispin et

al., 2010b), pCAG-mGFP (Addgene #14757), pCAGGS-RFP (Ofek et al., 2021). PCAGGS-

RARα403 and PCAGGS-Cyp26A1 (subcloned from PCAGGS-RARα403-IRES-eGFP and

PCAGGS-Cyp26A1-IRES-eGFP, respectively, from S. Sockanathan (Novitch et al.,

2003), were applied to inhibit RA activity. pCAB-cSmad6 was implemented to inhibit BMP

activity (Nitzan et al., 2016).

To monitor RA activity, we used the pGL3-RARE-SV40-AP reporter [from J. Sen, (Gupta and

Sen, 2015)]. Fluorescent versions of this reporter were prepared by replacing the AP tag with

either RFP or with a destabilized version of GFP (pGL3-RARE-SV40-RFP and pGL3-RARE-

SV40-d2EGFP, respectively (Clontech). Wnt activity was measured with 12XTOPFLASH-

d2EGFP (Rios et al., 2010).

Unilateral, bilateral and/or double electroporations to the NT were performed as detailed under

Results. To this end, DNA (4 ug/ul) was mixed with Fast Green and microinjected into the

lumen of the NT at the flank level of the axis. Five mm tungsten electrodes were placed on

either side of the embryo. A square wave electroporator (BTX, San Diego, CA, USA) was used

to deliver one pulse of current at 17–25 V for 6 ms.

Mechanical separation of the paraxial mesoderm from the NT

A unilateral slit between epithelial somites and NT comprising an approximate length of 8

segments was performed. A piece of aluminum foil was then inserted that extended beyond the

dorsal border of the tissues to provide a continuous barrier despite embryo growth. In some

cases, the dorsal NT was additionally opened to separate between both halves of the

neuroepithelium. Following microsurgery, embryos were reincubated for 48 hours prior to

fixation.

Immunohistochemistry

For immunostaining, embryos were fixed overnight at 4 °C with 4% formaldehyde (PFA) in

phosphate-buffered saline (PBS) (pH 7.4), embedded in paraffin wax and serially sectioned at

8 μm. Immunostaining was performed either on whole mount embryos or on paraffin sections,

as previously described (Burstyn-Cohen and Kalcheim, 2002; Kahane and Kalcheim, 2020).

Antibodies were diluted in PBS containing 5% fetal bovine serum (Biological Industries Israel,

04-007-1A) and 1% or 0.1% Triton X-100 (Sigma Aldrich Israel, X-100), respectively.

Antibodies used were: rabbit anti-GFP (1:1000, Invitrogen, Thermo-Fisher Scientific, A-

6455), mouse anti-GFP (1:100, Abcam, Ab38689), rabbit anti-RFP (1:1000, OriGene,

AP09229PU-N), guinea pig anti-pSmad1/5/8 (1:300, a gift from E. Laufer), mouse anti-H3-

pS10 (1:400, Abcam, Ab14955), rabbit anti-Laminin (1:100, Sigma Israel, L9393), rabbit anti-

Sox9 (1:150, Millipore, AB5535), rabbit anti-SNAI2 (1:500; Cell Signaling Technologies,

9585), mouse anti-CD57 (anti-HNK1, 1:500, BD Biosciences, 559048), rabbit anti-BarHL1

(1:300, Sigma Israel, HPA004809). Cell nuclei were visualized with 125 ng/ml Hoechst 33258

(Sigma Israel, 14530) diluted in PBS.

<u>In-situ hybridization</u>

For *in-situ* hybridization (ISH), embryos were fixed with Fornoy (60% ethanol, 30%

formaldehyde and 10% acetic acid) for 1 hour at room temperature, embedded in paraffin wax

and serially sectioned at 10 μm. ISH was performed on sections. In brief, sections were treated

with 1 ug/ml proteinase K, refixed in 4% PFA, then hybridized overnight at 65 °C with

digoxigenin-labeled RNA probes (Roche, 11277073910). The probes were detected with AP

coupled with anti-digoxigenin Fab fragments (Roche, 11093274910). AP reaction was

developed with 4-Nitro blue tetrazolium chloride (NBT, Roche, 11383213001) and 5-bromo-

4-chloro-3'-indolyphosphate p-toluidine salt (BCIP, Sigma Israel, B8503). In the *foxd3/Rspo1*

ISH experiment, the AP reaction of *Rspo1* was developed with Fast Red (Sigma Israel, F4648)

for 2 hours at room temperature. Importantly, all hybridizations, whether done on intact

embryos at NC vs. RP stages, or in control vs. experimental embryos, were always developed

for the same time per given probe and experiment. The RNA probes used are described in Table

S1. Several probes were produced by PCR amplification (KAPA2G Fast ReadyMix PCR kit,

Sigma Israel, KK5101). cDNA templates were synthesized by RNA precipitation followed by

reverse transcription PCR. RNAs were produced from 20ss-E4 quail embryos. Tissue samples

were homogenized with TriFast reagent, and RNA was separated with chloroform and

isopropanol. The primers used are depicted in Table S1.

 Data acquisition and statistical analysis

Measurement of the intensity of immunofluorescence and ISH signals: Intensities were

measured using FIJI (Schindelin et al., 2009). In most cases, the RP was selected as the region

of interest (ROI); mean intensity was measured and background intensity was then subtracted.

Average intensity of all sections was calculated per embryo, and values were normalized with

a control mean set as 1.

For measurement of phospho-Smad1/5/8 intensity, the ROI was selected as the dorsal domain

of the NT from the dorsal midline to the ventral border of dI1 progenitors.

To monitor intensity of laminin immunostaining in the RP basal lamina, a segmented line of

width 4 was drawn along the basal aspect of the RP, and mean intensity was measured within

that area. Average intensity was calculated for each embryo and expressed relative to a control

value set as 1.

To measure the relative position of cell nuclei across the apico-basal extent of the RP, the latter

was divided into basal and apical halves, and Hoechst-stained nuclei were manually counted

in both domains.

The distance of BarHL1[+] cells from the dorsal midline of the NT was measured for each cell

manually using FIJI, and histograms were generated using R (https://www.R-project.org/) in

Rstudio (http://www.rstudio.com/).

Images were photographed using a DP73 (Olympus) cooled CCD digital camera mounted on

a BX51 microscope (Olympus) with Uplan FL-N 20x/0.5 and 40x/0.75 dry objectives

(Olympus). Confocal sections encompassing their entire thickness were photographed using a

Nikon Eclipse 90i microscope with a Plan Apo 40x/1.3 or 100x/1.4 objectives (Nikon) and a

D-Eclipse c1 confocal system (Nikon) at 1.0 μm increments through the z-axis. Images were

z-stacked with FIJI software.

For quantification, images of control and treated sections were photographed under the same

conditions. For figure preparation, images were exported into Photoshop CS6 (Adobe). If

necessary, the levels of brightness and contrast were adjusted to the entire image and images

were cropped without color correction adjustments or γ adjustments. Graphics were generated

using Graphpad Prism 7.0, and figures were prepared using Photoshop and InDesign CS6.

Results were processed with Graphpad Prism 7 and presented as scatter plots with mean±SEM.

Data were subjected to statistical analysis using either of the following tests, as described in

the respective legends to Figures: Student's t-test, Welch's t-test, Mann-Whitney test, 1-way

ANOVA with post-hoc Tukey test, Kruskal-Wallis with post-hoc Dunn's test and Wilcoxon

signed ranks test. All tests applied were two-tailed and a P-value ≤ 0.05 was considered

significant.

Acknowledgments

We are grateful to all the researchers whose names are mentioned along the text for generously

providing reagents. We thank Avihu Klar and Joel Yisraeli for critical reading of the

manuscript. This study was supported by grants from the Israel Science Foundation (ISF

#209/18) and from the Ines Mandl Research Fund (IMRF) to CK.

Author Contributions

DR and CK conceived the article, wrote it and prepared the Figures. DR performed the

experiments. Both authors discussed and agreed on the text and approved the manuscript.

Conflicts of Interest

The authors declare no conflict of interests

References

References

Andrews, M. G., Kong, J., Novitch, B. G. and Butler, S. J. (2019). New perspectives on

the mechanisms establishing the dorsal-ventral axis of the spinal cord. *Curr Top Dev*

Biol **132**, 417–450.

Augsburger, A., Schuchardt, A., Hoskins, S., Dodd, J. and Butler, S. (1999). Bmps as

mediators of roof plate repulsion of commissural neurons. *Neuron* **24**, 127–141.

Berggren, K., McCaffery, P., Drager, U. and Forehand, C. J. (1999). Differential

distribution of retinoic acid synthesis in the chicken embryo as determined by

immunolocalization of the retinoic acid synthetic enzyme, RALDH-2. *Dev Biol* **210**,

288–304.

Blentic, A., Gale, E. and Maden, M. (2003). Retinoic acid signalling centres in the avian

embryo identified by sites of expression of synthesising and catabolising enzymes. *Dev*

Dyn **227**, 114–127.

Brazil, D. P., Church, R. H., Surae, S., Godson, C. and Martin, F. (2015). BMP

signalling: agony and antagony in the family. *Trends Cell Biol* **25**, 249–264.

Burstyn-Cohen, T. and Kalcheim, C. (2002). Association between the cell cycle and neural

crest delamination through specific regulation of G1/S transition. *Dev. Cell* **3**,.

Burstyn-Cohen, T., Stanleigh, J., Sela-Donenfeld, D. and Kalcheim, C. (2004). Canonical

Wnt activity regulates trunk neural crest delamination linking BMP/noggin signaling

with G1/S transition. *Development* **131**, 5327–5339.

Butler, S. J. and Dodd, J. (2003). A role for BMP heterodimers in roof plate-mediated

repulsion of commissural axons. *Neuron* **38**, 389–401.

Chambers, D., Wilson, L., Maden, M. and Lumsden, A. (2007). RALDH-independent

generation of retinoic acid during vertebrate embryogenesis by CYP1B1. *Development*

134, 1369–1383.

Chesnutt, C., Burrus, L. W., Brown, A. M. and Niswander, L. (2004). Coordinate

regulation of neural tube patterning and proliferation by TGFbeta and WNT activity.

Dev Biol **274**, 334–347.

Chizhikov, V. V and Millen, K. J. (2005). Roof plate-dependent patterning of the vertebrate

dorsal central nervous system. *Dev Biol* **277**, 287–295.

Diez del Corral, R. and Morales, A. V. (2014). Retinoic acid signaling during early spinal

cord development. *J. Dev. Biol.* **2**, 174–197.

Diez del Corral, R., Olivera-Martinez, I., Goriely, A., Gale, E., Maden, M. and Storey,

K. (2003). Opposing FGF and retinoid pathways control ventral neural pattern, neuronal

differentiation, and segmentation during body axis extension. *Neuron* **40**, 65–79.

Duval, N., Vaslin, C., Barata, T. C., Frarma, Y., Contremoulins, V., Baudin, X.,

Nedelec, S. and Ribes, V. C. (2019). BMP4 patterns Smad activity and generates

stereotyped cell fate organization in spinal organoids. *Development* **146**, doi:

10.1242/dev.175430.

Endo Y., Osumi N. and Wakamatsu Y. (2003). Bimodal functions of Notch-mediated

signaling are involved in neural crest formation during avian ectoderm development.

Development. **129**, 863–873.

George, L., Chaverra, M., Todd, V., Lansford, R. and Lefcort, F. (2007). Nociceptive

sensory neurons derive from contralaterally migrating, fate-restricted neural crest cells.

Nat Neurosci **10**, 1287–1293.

Gupta, S. and Sen, J. (2015). Retinoic acid signaling regulates development of the dorsal

forebrain midline and the choroid plexus in the chick. *Dev.* **142**, 1293–1298.

Halperin-Barlev, O. and Kalcheim, C. (2011). Sclerotome-derived slit1 drives directional

migration and differentiation of robo2-expressing pioneer myoblasts. *Development* **138**,.

Haselbeck, R. J., Hoffmann, I. and Duester, G. (1999). Distinct functions for Aldh1 and

Raldh2 in the control of ligand production for embryonic retinoid signaling pathways.

Dev. Genet. **25**, 353–364.

Hegarty, S. V., O'Keeffe, G. W. and Sullivan, A. M. (2013). BMP-Smad 1/5/8 signalling

in the development of the nervous system. *Prog. Neurobiol.* **109**, 28–41.

Kahane, N. and Kalcheim, C. (1998). Identification of early postmitotic cells in distinct

embryonic sites and their possible roles in morphogenesis. *Cell Tissue Res.* **294**, 297–

307.

Kahane, N. and Kalcheim, C. (2020). Neural tube development depends on notochord-

derived sonic hedgehog released into the sclerotome. *Development* **147**,.

Kalcheim, C. (2018). Neural crest emigration: From start to stop. *Genesis* **56**,.

Kalcheim, C. and Kumar, D. (2017). Cell fate decisions during neural crest ontogeny. *Int. J.*

Dev. Biol. **61**, 195–203.

Kam, R. K. T., Deng, Y., Chen, Y. and Zhao, H. (2012). Retinoic acid synthesis and

functions in early embryonic development. *Cell Biosci.* **2**, 11.

Kidd, T., Bland, K. S. and Goodman, C. S. (1999). Slit is the midline repellent for the robo

receptor in Drosophila. *Cell* **96**, 785–794.

Krispin, S., Nitzan, E. and Kalcheim, C. (2010a). The dorsal neural tube: A dynamic

setting for cell fate decisions. *Dev. Neurobiol.* **70**,.

Krispin, S., Nitzan, E., Kassem, Y. and Kalcheim, C. (2010b). Evidence for a dynamic
spatiotemporal fate map and early fate restrictions of premigratory avian neural crest.
Development **137**, 585–595.

Kukreja, S., Udaykumar, N., Yogesh, B. and Sen, J. (2020). Retinoic acid signaling
regulates proliferation and lamina formation in the developing chick optic tectum. *Dev.*
Biol. **467**, 95–107.

Lara-Ramírez, R., Zieger, E. and Schubert, M. (2013). Retinoic acid signaling in spinal
cord development. *Int. J. Biochem. Cell Biol.* **45**, 1302–1313.

Le Douarin, N. M. and Kalcheim, C. (1999). *The Neural Crest*. 2nd ed. New York:
Cambridge University Press.

Le Dréau, G. and Martí, E. (2013). The multiple activities of BMPs during spinal cord
development. *Cell. Mol. Life Sci.* **70**, 4293–4305.

Lee, K. J., Mendelsohn, M. and Jessell, T. M. (1998). Neuronal patterning by BMPs: a
requirement for GDF7 in the generation of a discrete class of commissural interneurons
in the mouse spinal cord. *Genes Dev.* **12**, 3394–3407.

Maden, M., Ong, D. E., Summerbell, D., Chytil, F. and Hirst, E. A. (1989). Cellular
retinoic acid-binding protein and the role of retinoic acid in the development of the chick
embryo. *Dev. Biol.* **135**, 124–132.

Martinez-Morales, P. L., Diez del Corral, R., Olivera-Martinez, I., Quiroga, A. C., Das,
R. M., Barbas, J. A., Storey, K. G. and Morales, A. V (2011). FGF and retinoic acid
activity gradients control the timing of neural crest cell emigration in the trunk. *J Cell*
Biol **194**, 489–503.

Martínez-Morales, P. L., del Corral, R. D., Olivera-Martínez, I., Quiroga, A. C., Das, R.

M., Barbas, J. A., Storey, K. G. and Morales, A. V. (2011). FGF and retinoic acid

activity gradients control the timing of neural crest cell emigration in the trunk. *J. Cell*

Biol. **194**, 489–503.

Muroyama, Y., Fujihara, M., Ikeya, M., Kondoh, H. and Takada, S. (2002). Wnt

signaling plays an essential role in neuronal specification of the dorsal spinal cord.

Genes Dev. **16**, 548–553.

Niederreither, K., Subbarayan, V., Dollé, P. and Chambon, P. (1999). Embryonic retinoic

acid synthesis is essential for early mouse post- implantation development. *Nat. Genet.*

21, 444–448.

Nieto, M. A., Sargent, M. G., Wilkinson, D. G. and Cooke, J. (1994). Control of cell

behavior during vertebrate development by slug, a zinc finger gene. *Science (80-.).* **264**,

835–839.

Nitzan, E., Krispin, S., Pfaltzgraff, E. R., Klar, A., Labosky, P. A. and Kalcheim, C.

(2012). A dynamic code of dorsal neural tube genes regulates the segregation between

neurogenic and melanogenic neural crest cells. *Dev.* **140**,.

Nitzan, E., Avraham, O., Kahane, N., Ofek, S., Kumar, D. and Kalcheim, C. (2016).

Dynamics of BMP and Hes1/Hairy1 signaling in the dorsal neural tube underlies the

transition from neural crest to definitive roof plate. *BMC Biol.* **14**, 23. DOI

10.1186/s12915-016-0245–6.

Novitch, B. G., Wichterle, H., Jessell, T. M. and Sockanathan, S. (2003). A requirement

for retinoic acid-mediated transcriptional activation in ventral neural patterning and

motor neuron specification. *Neuron* **40**, 81–95.

Ofek, S., Wiszniak, S., Kagan, S., Tondl, M., Schwarz, Q. and Kalcheim, C. (2021).

Notch signaling is a critical initiator of roof plate formation as revealed by the use of

RNA profiling of the dorsal neural tube. *BMC Biol.* **19:84**,

https://doi.org/10.1186/s12915-021-01014-3.

Reijntjes, S., Blentic, A., Gale, E. and Maden, M. (2005). The control of morphogen

signalling: regulation of the synthesis and catabolism of retinoic acid in the developing

embryo. *Dev Biol* **285**, 224–237.

Rekler, D. and Kalcheim, C. (2021). From Neural Crest to Definitive Roof Plate : The

Dynamic Behavior of the Dorsal Neural Tube. *Int. J. Mol. Sci.* **22**, 3911.

Rhinn, M. and Dollé, P. (2012). Retinoic acid signalling during development. *Development*

139, 843–858.

Rios, A. C., Denans, N. and Marcelle, C. (2010). Real-time observation of Wnt beta-catenin

signaling in the chick embryo. *Dev Dyn* **239**, 346–353.

Schindelin, J., Arganda-Carrera, I., Frise, E., Verena, K., Mark, L., Tobias, P., Stephan,

P., Curtis, R., Stephan, S., Benjamin, S., et al. (2009). Fiji - an Open platform for

biological image analysis. *Nat. Methods* **9**,.

Sela-Donenfeld, D. and Kalcheim, C. (1999). Regulation of the onset of neural crest

migration by coordinated activity of BMP4 and Noggin in the dorsal neural tube.

Development **126**, 4749–4762.

Sela-Donenfeld, D. and Kalcheim, C. (2002). Localized BMP4-noggin interactions generate

the dynamic patterning of noggin expression in somites. *Dev. Biol.* **246**,.

Shaker, M. R., Lee, J., Park, S., Kim, J. Y., Son, G. H., Son, J. W., Park, B. H., Rhyu, I.

J., Kim, H. and Sun, W. (2020). Stem Cell Reports. *Stem Cell Reports* **15**, 1–14.

Sheng, N., Xie, Z., Wang, C., Bai, G., Zhang, K., Zhu, Q., Song, J., Guillemot, F., Chen,

Y. G., Lin, A., et al. (2010). Retinoic acid regulates bone morphogenic protein signal

duration by promoting the degradation of phosphorylated Smad1. *Proc. Natl. Acad. Sci.*

U. S. A. **107**, 18886–18891.

Shinozuka, T., Takada, R., Yoshida, S., Yonemura, S. and Takada, S. (2019). Wnt

produced by stretched roof-plate cells is required for the promotion of cell proliferation

around the central canal of the spinal cord. *Development* **146**,.

Sun, J., Zhuang, F. F., Mullersman, J. E., Chen, H., Robertson, E. J., Warburton, D.,

Liu, Y. H. and Shi, W. (2006). BMP4 activation and secretion are negatively regulated

by an intracellular Gremlin-BMP4 interaction. *J. Biol. Chem.* **281**, 29349–29356.

Theveneau, E. and Mayor, R. (2012). Neural crest migration: interplay between

chemorepellents, chemoattractants, contact inhibition, epithelial-mesenchymal

transition, and collective cell migration. *Wiley Interdiscip Rev Dev Biol* **1**, 435–445.

Timmer, J. R., Wang, C. and Niswander, L. (2002). BMP signaling patterns the dorsal and

intermediate neural tube via regulation of homeobox and helix-loop-helix transcription

factors. *Development* **129**, 2459–2472.

Tozer, S., Le Dreau, G., Marti, E. and Briscoe, J. (2013). Temporal control of BMP

signalling determines neuronal subtype identity in the dorsal neural tube. *Development*

140, 1467–1474.

Venepally, P., Reddy, L. G. and Sani, B. P. (1996). Characterization of cellular retinoic

acid-binding protein I from chick embryo and its ligand binding properties.

Arch.Biochem.Biophys. **336**, 231–239.

Wilson, L., Gale, E., Chambers, D. and Maden, M. (2004). Retinoic acid and the control of

dorsoventral patterning in the avian spinal cord. *Dev Biol* **269**, 433–446.

Wine-Lee, L., Ahn, K. J., Richardson, R. D., Mishina, Y., Lyons, K. M. and Crenshaw,

E. B. (2004). Signaling through BMP type 1 receptors is required for development of

interneuron cell types in the dorsal spinal cord. *Development* **131**, 5393–5403.

Xing, L., Anbarchian, T., Tsai, J. M., Plant, G. W. and Nusse, R. (2018). Wnt/beta-

catenin signaling regulates ependymal cell development and adult homeostasis. *Proc*

Natl Acad Sci U S A **115**, E5954–E5962.

Zechner, D., Muller, T., Wende, H., Walther, I., Taketo, M. M., Crenshaw 3rd, E. B.,

Treier, M., Birchmeier, W. and Birchmeier, C. (2007). Bmp and Wnt/beta-catenin

signals control expression of the transcription factor Olig3 and the specification of spinal

cord neurons. *Dev Biol* **303**, 181–190.

1005