SNP and Haplotype Regional Heritability Mapping (SNHap-RHM): joint mapping of

common and rare variation affecting complex traits

Richard F. Oppong[1], Pau Navarro[2], Chris S. Haley[2,3] and Sara Knott[1]

[1]IEB, SBS, University of Edinburgh, Edinburgh, United Kingdom,

[2]MRC IGMM, University of Edinburgh, Edinburgh, United Kingdom,

[3]The Roslin Institute and Royal (Dick) School of Veterinary Studies, Edinburgh, United

Kingdom.

Correspondence: s.knott@ed.ac.uk

Abstract

We describe a genome-wide analytical approach, SNP and Haplotype Regional Heritability Mapping (SNHap-RHM), that provides regional estimates of the heritability across locally defined regions in the genome. This approach utilises relationship matrices that are based on sharing of SNP and haplotype alleles at local haplotype blocks delimited by recombination boundaries in the genome. We implemented the approach on simulated data and show that the haplotype-based regional GRMs capture variation that is complementary to that captured by SNP-based regional GRMs, and thus justifying the fitting of the two GRMs jointly in a single analysis (SNHap-RHM). SNHap-RHM captures regions in the genome contributing to the phenotypic variation that existing genome-wide analysis methods may fail to capture. We further demonstrate that there are real benefits to be gained from this approach by applying it to real data from about 20,000 individuals from the Generation Scotland: Scottish Family Health Study. We analysed height and major depressive disorder (MDD). We identified seven genomic regions that are genome-wide significant for height, and three regions significant at a suggestive threshold (p-value $< 1 \times 10^{-5}$) for MDD. These significant regions have genes mapped to within 400kb of them. The genes mapped for height have been reported to be associated with height in humans, whiles those mapped for MDD have been reported to be associated with major depressive disorder and other psychiatry phenotypes. The results show that SNHap-RHM presents an exciting new opportunity to analyse complex traits by allowing the joint mapping of novel genomic regions tagged by either SNPs or haplotypes, potentially leading to the recovery of some of the "missing" heritability.

Keywords: MDD; height; haplotypes; regional heritability mapping; missing heritability; rare variation; genome-wide analysis

Introduction

Estimates of the genetic component of complex trait variation using genotyped SNPs led to the conclusion that a proportion of the heritability of complex traits is still unexplained or "missing" (Maher, 2008; Manolio et al., 2009). Full sequence data will contain all the variants that account for all the heritability of complex traits (Wainschtein et al., 2019). Moreover, some of these true causal variants may be rare (Pritchard, 2001) and therefore may be in incomplete linkage disequilibrium (LD) with genotyped SNPs (Yang et al., 2010). Thus, some of the "missing" heritability may be "hidden" in rare variants whose effects are difficult to capture because of lack of statistical power. There is, therefore, some benefit to be gained in terms of improving the heritability estimates and uncovering gene variants involved in the control of traits by fitting genome-wide analytical models that adequately capture the combined effects of rare genetic variants (Cirulli and Goldstein, 2010; Gonzalez-Recio et al., 2015).

In light of this, we proposed a genome-wide analytical approach that draws its theoretical basis from the genome-based restricted maximum likelihood (GREML) approach (Clarke and Cooper, 2010; Maher, 2008; Manolio et al., 2009; Speed et al., 2012; Yang et al., 2011) which utilises both local and genome-wide relationship matrices to provide regional estimates of the heritability across locally defined regions in the genome (Nagamine et al., 2012). This regional heritability analysis can capture the combined effect of SNPs in a region, and thus small effect variants may be detectable. However, the analysis only captures effects associated with individual SNPs.

Haplotypes may provide a better strategy to capture genomic relationships amongst individuals in the presence of causal rare variants. Although rare variants are not in LD with

genotyped variants and thus are difficult to capture in conventional GWAS, these rare

variants, may be in LD with some haplotypes and thus can be captured using haplotype

methods. Compared with genotyped SNPs, capturing haplotype effects may offer an

advantage because haplotypes can be functional units (Vormfelde and Brockmöller, 2007).

Therefore, haplotype effects may reflect the combined effects of closely linked cis-acting

causal variants (Balding, 2006) and using haplotypes could provide real benefit over SNPs in

recovering some of the "missing" heritability and identifying novel trait-associated variants.

Therefore, we extended the SNP-based regional heritability analysis further by incorporating

haplotypes in addition to SNPs in the calculation of the regional GRMs used in the analysis

(Shirali et al., 2018). This approach includes two regional GRMs and divides the genome into

windows based on local haplotype blocks delimited by recombination boundaries.

This paper further explores the properties of both the SNP-based and the haplotype-

based regional heritability mapping (SNP-RHM and Hap-RHM respectively). We hypothesise

and show by simulation that the Hap-RHM complements existing SNP-RHM analytical

approaches by capturing regional effects in the genome that existing SNP-based methods fail

to capture. This leads us to propose a mapping strategy that jointly utilises SNP and haplotype

GRMs in a single analysis called SNHap-RHM. We then confirm the utility of this approach by

applying it to real data obtained from about 20,000 individuals from the Generation Scotland:

Scottish Family Health Study (GS: SFHS) (Smith et al., 2012). We analysed two phenotypes:

height and major depressive disorder (MDD). The aim was to uncover novel genetic loci that

may affect these traits and improve the estimates of the genetic components of the variation

in these traits.

Methods

The general statistical setting of a regional GREML analysis

Consider a vector $\boldsymbol{y}$ of phenotype values with length n, the linear mixed-effects model

for fitting the effects of genomic region i and background polygenic markers is given as:

93
$$y = X\beta + W_i u_i + Z u_b + e$$

where $\boldsymbol{y}$ is a vector of phenotypes, $\boldsymbol{X}$ is a design matrix of fixed effects, and $\boldsymbol{\beta}$ is a vector of

fixed effects, $\boldsymbol{W_i}$ is a design matrix relating phenotype measures to genetic markers in region

i and $\boldsymbol{u_i}$ is a vector of random genetic effects due to region i assumed to be multivariate

normal, $MVN\left(0, \sigma_{u_i}^2 \boldsymbol{L_{u_i}}\right)$. $\boldsymbol{L_{u_i}}$ is a relationship matrix calculated using markers (SNPs or

haplotypes) in region i: calculated in the subsequent sections as $\boldsymbol{G}$ for the SNP and $\boldsymbol{H}$ for the

haplotype-based models. $\boldsymbol{Z}$ is a design matrix for background polygenic effects of markers

outside the region i and $\boldsymbol{u_b}$ is a vector of random polygenic effect of genetic markers excluded

from region i, assumed to be multivariate normal, $MVN\left(0, \sigma_{u_b}^2 \boldsymbol{B_{u_b}}\right)$. $\boldsymbol{B_{u_b}}$ is a relationship

matrix calculated using the markers outside the region i: calculated in the subsequent section

in the same way as $\boldsymbol{G}$. And $\boldsymbol{e}$ is a vector of residual effects assumed to be multivariate normal,

$MVN(0, \sigma_e^2 \boldsymbol{I})$. $\boldsymbol{I}$ is an identity matrix.

Under the model, the vector of phenotypes $\boldsymbol{y}$ is assumed to be normally distributed,

$N(\boldsymbol{X\beta}, \boldsymbol{V})$ where the variance is

107
$$V = \sigma_{u_i}^2 \boldsymbol{L_{u_i}} + \sigma_{u_b}^2 \boldsymbol{B_{u_b}} + \sigma_e^2 \boldsymbol{I}$$

SNP-RHM: SNP-based regional GREML model

A SNP-based regional GREML analysis was first reported by Nagamine et al. (2012). The regional GREML analysis approach we employ here differs from the analysis done by Nagamine et al. (2012) in the way the regions are defined. That analysis defined local regions by breaking the genome into smaller user-defined windows of p SNPs, which overlapped by q SNPs. Here, however, we define regions based on recombination boundaries in the genome.

The regional GREML model fits two genetic relationship matrices (GRMs): one local GRM for the region and a whole-genome GRM for the remaining SNPs in the genome that are outside the region. The GRMs are genomic relatedness matrices calculated as the weighted proportion of the local or genome-wide autosomal SNPs shared identity by state (IBS) between pairs of individuals. The SNP IBS matrices are calculated as follows, following the second scaling factor proposed by VanRaden (2008)

$$G = \frac{MM'}{m}$$

where m is the total number of r local or b background autosomal SNPs, and $\boldsymbol{M}$ is a matrix of genotype codes for the sampled individuals centred by loci means and normalised by the standard deviation of each locus. $\boldsymbol{M}$ is calculated as follows for individual i at locus j

$$M_{ij} = \frac{(x_{ij} - 2p_j)}{\sqrt{2p_j(1 - p_j)}}$$

where x_{ij} is the genotype code at locus j for individual i and takes the values 0, 1 and 2 for AA, Aa and aa genotypes respectively, p_j is the frequency of allele 'a' at locus j. The SNP-based relationship for individuals i and k is therefore calculated as follows

$$130 \qquad G_{ik} = \frac{1}{m} \times \sum\nolimits_{j=1}^{m} \frac{(x_{ij} - 2p_j)(x_{kj} - 2p_j)}{2p_j(1 - p_j)}$$

Hap-RHM: Haplotype-based regional GREML model

The haplotype-based regional GREML model follows theoretically from the SNP-based

analysis and utilises haplotypes instead of SNPs as the genetic markers for the regional

analysis. The analysis fits two GRMs, a haplotype-based regional GRM and a SNP-based

background genome-wide GRM. The haplotype-based GRM is similar to the SNP-based GRM

defined in the previous section. For a locally defined region (haplotype block) containing h

haplotype variants, the haplotype-based kinship for individuals i and k is calculated as follows

$$138 \qquad H_{ik} = \frac{1}{h} \times \sum\nolimits_{j=1}^{h} \frac{(d_{ij} - 2p_j)(d_{kj} - 2p_j)}{2p_j(1 - p_j)}$$

where d_{ij} is the diplotype code (coded as the number of copies of haplotype j) for individual

i and takes the values 0, 1 and 2 for the $h_t h_t$, $h_t h_j$, $h_j h_j$ diplotypes respectively where

haplotype t is any haplotype other than haplotype j, i.e. $t \neq j$, p_j is the haplotype frequency

for haplotype j.

Phenotype Simulations

Five phenotypes were simulated using available genotypic information of 20,032

individuals from the Generation Scotland: Scottish Family Health Study (Smith et al., 2012). A

total of 593,932 genotyped SNPs were used, and missing genotypes were filled in by

imputation. A total of 555,091 SNPs remained after a QC that removed SNPs of MAF < 0.01

and SNPs that were out of Hardy-Weinberg equilibrium at p-value < 0.000001.

The five phenotypes were simulated to have a total variance of 1. This total is

composed of 0.6 environmental (residual) variance and genetic variance of 0.4. The genetic

variance was partitioned into two components, a polygenic variance of 0.3 and a total QTL

variance of 0.1 (20 QTLs, each explaining a variance of 0.005). A common polygenic variance

was simulated for all five phenotypes from 20,000 markers randomly selected across the

genome. The polygenic variance was simulated to be normally distributed with zero mean

and variance of 0.3.

For each phenotype, 20 regions (haplotype blocks) were randomly selected, one on

each chromosome (except chromosomes 6 and 8 because of the unusually high LD in the MHC

regions on chromosome 6 and a large inversion on chromosome 8 (Amador et al., 2015)), to

simulate quantitative trait loci (QTL). This gave a total of 20 QTLs for each phenotype. The

regions were delimited by natural boundaries: recombination hotspots where the estimated

recombination frequency exceeds ten centiMorgans per Megabase (10cM/Mb) with the

estimated recombination frequency between boundaries being less than ten centiMorgans

per Megabase (10cM/Mb) based on the Genome Reference Consortium Human Build 37

(International Human Genome Sequencing Consortium, 2004). This recombination threshold

resulted in a total 48,772 regions across the genome. The number and type of marker used

to simulate the QTL are what defined the five phenotypes. The five phenotypes are, a 1-SNP

QTL within the haplotype block, a multiple-SNP (5 SNPs) QTL within the haplotype block, two

types of 1-haplotype QTL within the haplotype block (taking either a common or a rare

haplotype as causal) and multiple (5) haplotype QTL within the haplotype block. Details of

these phenotypes are described below.

For the haplotype QTL phenotypes, a haplotype block is treated as a single genetic

locus having multiple alleles. Each haplotype variant within a block is considered as an allele

of that locus. Each study individual will carry two alleles, or have a diplotype, for each locus

or haplotype block. The genotype data used to simulate the phenotypes were phased using

SHAPEIT2 (Delaneau et al., 2013) to produce the haplotypes for study individuals. The multiple

haplotype QTL phenotypes were simulated by randomly sampling two rare haplotypes and

three common haplotypes within each haplotype block to give five haplotypes per block. The

two types of 1-haplotype QTL phenotypes were simulated by randomly sampling a rare

haplotype per haplotype block for one type and for the other type a common haplotype was

randomly sampled within each haplotype block.

The individual marker contribution to the polygenic effect and the QTL effects were

calculated as follows

183
$$\sigma_j^2 = 2p_j(1 - p_j)g_j^2$$

184
$$g_j = \sqrt{\frac{\sigma_j^2}{2p_j(1 - p_j)}}$$

where σ_j^2 is the contribution of a marker to the QTL or polygenic variance, g_j is the effect of

a SNP j or haplotype j randomly sampled to have polygenic or QTL effect, p_j is the frequency

of haplotype j or the effect allele of the SNP j. For the single marker QTL phenotypes, each

QTL explained a variance of 0.005. For the multiple marker QTL phenotypes, each causal

variant explained the same variance, with the effects scaled to account for LD in the region

so each QTL locus had a variance of 0.005. For the multiple haplotype QTL effects, the

haplotype effects were scaled relative to the inverse of their frequency to give a total variance

explained by the region of 0.005.

Common environmental effects were randomly sampled for the five phenotypes from

a normal distribution $N(0, \sigma_e^2)$ where σ_e^2 is 0.6. This, together with a genetic variance of 0.4,

gave a total variance of 1 for each phenotype. The final simulated phenotype for an

individual i was then calculated as follows

$$y(single\ markers\ per\ QTL\ region)_i = \sum_{j=1}^{20000} x_{ij}g_j + \sum_{j=1}^{20} x_{ij}g_j + e_i,$$

$$y(multiple\ markers\ per\ QTL\ region)_i = \sum_{j=1}^{20000} x_{ij}g_j + \sum_{l=1}^{20}\sum_{j=1}^{5} x_{ij}g_j + e_i,$$

where x_{ij} is the number of copies of the effect allele of SNP j for individual i (for haplotypes,

this is defined as d_{ij}; the number of copies of haplotype j for individual i) and g_j is the effect

of haplotype j or SNP j. Twenty replicates were analysed for each of the five phenotypes with

a different set of QTL markers sampled for each replicate.

Analysis of simulated data

We have shown previously that regional GREML analysis (Regional Heritability

Mapping or RHM) using fixed region sizes in the genome is a suitable mapping method for

finding local genetic effects (Nagamine et al., 2012). The conventional RHM model fits two

genomic relationship matrices (GRMs) in the analyses to map genetic loci that affect trait

variation: a local GRM (rGRM) calculated using SNPs located in the region and a genome-wide

GRM (gwGRM) calculated from SNPs outside the region. We have since extended this

conventional regional GREML analysis to incorporate haplotypes in the calculation of the local

GRM and have successfully implemented this in a simulation study (Shirali et al., 2018). This

study by Shirali et al. (2018) utilises a regional GREML model that breaks the genome into

naturally defined regions by delimiting them by recombination hotspots. Two types of

regional GREML models are then fitted in turn to the phenotypes. One model (SNP-RHM) uses

SNPs to estimate local genetic relationships between study individuals, and the other model

(Hap-RHM) estimates local genetic relationships amongst individuals using haplotypes.

In this simulation study, the five simulated phenotypes were analysed using the two

models, the SNP-based regional GREML model (SNP-RHM for the SNP QTL phenotypes) and

the haplotype-based regional GREML model (Hap-RHM for the haplotype QTL phenotypes).

To test the analytical models' specificity, we applied Hap-RHM to SNP QTL phenotypes and

SNP-RHM to the haplotype QTL phenotypes. We also performed a Hap-RHM analysis in which

the units of analysis in the haplotype blocks were restricted to regions of 20 or fewer SNPs

per haplotype block. This was because we observed that longer haplotype blocks had many

SNPs (and hence many, many haplotypes), and this impacted the estimation of the simulated

regional effect. The hybrid Hap-RHM, therefore, investigates whether the regional effect is

well captured by the haplotype-based model when shorter haplotypes are used.

We estimated the regional genetic variance and polygenic variance using restricted

maximum likelihood (REML). For each simulated phenotype, we analysed 220 regions in total

to map the 20 simulated QTLs. This involved analysing the region containing the QTL and ten

adjacent regions (five in either direction). In this way, we limit the analysis to the regions in

the genome with simulated effects, thereby reducing computation time considerably. Also,

by analysing neighbouring regions, we are able to explore the precision of estimates of the

location of regional effects. We assessed the significance of a region using the Likelihood Ratio

Test (LRT). The genome-wide significance threshold was calculated to be LRT = 23.9 (p-value

$< 1.02 \times 10^{-6}$) using a Bonferroni correction for testing 48,772 regions.

Also, we selected one replicate for each simulated phenotype and performed a

regional heritability analysis that jointly fitted the SNP and the haplotype GRM in an approach

that we termed SNP and Haplotype Regional Heritability Mapping (SNHap-RHM). An overview

of SNHap-RHM is shown in Figure 1.

GS: SFHS Data

Genotyping, quality control and phasing of Generation Scotland: Scottish Family Health

Study dataset

The data from the Generation Scotland: Scottish Family Health Study comprised

23,960 participants recruited from Scotland (Smith et al., 2006). The DNA from about 20,032

of the participants had been genotyped using the Illumina HumanOmniExpressExome8v1-2_A

chip (~700K genome-wide SNP chip) (Smith et al., 2012).

Quality control excluded SNPs and individuals with a call rate less than 98%, SNPs with

minor allele frequency (MAF) less than 1% and SNPs that were out of Hardy-Weinberg

equilibrium (p-value < 0.000001). A total of 555,091 autosomal SNPs passed quality control

for downstream analysis. Phasing of the GS: SFHS data was done using SHAPEIT2 (Delaneau

et al., 2013). Haplotype blocks were defined using recombination hotspots with a

recombination rate of 10cM/Mb inferred from the Reference Consortium Human Build 37

(International Human Genome Sequencing Consortium, 2004). Haplotypes variants within

blocks were determined using the phased data.

Phenotype definition

MDD status for GS: SFHS participants was assigned following an initial mental health

screening questionnaire with the questions: "Have you ever seen anybody for emotional or

psychiatric problems?" or "Was there ever a time when you, or someone else, thought you

should see someone because of the way you were feeling or acting?" Participants who

answered yes to one or both of the screening questions were further interviewed by the

structured clinical interview to diagnose mood disorders (SCID) (First et al., 2002). A total of

18,725 participants (2,603 MDD cases and 16,122 controls) were retained for analysis for

MDD. A total of 19,944 participants from the GS: SFHS were analysed for height.

SNHap-RHM of MDD and Height

SNHap-RHM fits jointly, the two types of regional GRMs, SNP-based and haplotype-

based, in the analysis of phenotypes (Figure 1). We pre-corrected the phenotypes with the

whole-genome GRM before performing SNHap-RHM to speed up the GREML analysis of each

block. This pre-correction has previously been shown to speed the regional heritability

analysis by Shirali et al. (2018). This step involved 22 separate GREML analyses each fitting a

whole-genome GRM that excluded SNPs from one chromosome. The residuals from the pre-

correction step were then used in the SNHap-RHM analysis. The models adjusted for sex, age,

age^2, and the first 20 principal components calculated from the study participants' genomic

relationship matrix (calculated using 555,091 autosomal SNPs).

The significance of a region was tested with a likelihood ratio test (LRT) with two

degrees of freedom which compared a model with three variance components fitted (the two

regional variances together with the residual variance) against a model with only the residual

variance component fitted. The individual regional variance components were subsequently

tested with an LRT with one degree of freedom which compared a model with three variance

components fitted against a model with two variance components fitted (one regional

variance component dropped from the model).

The p-values obtained from the LRTs were used to generate genome-wide association

plots for each phenotype (equivalent to GWAS Manhattan plots). The genome-wide

significance threshold was calculated to be LRT = 23.9 (p-value $< 1.02 \times 10^{-6}$) using a

Bonferroni correction for testing 48,772 regions. The suggestive significance threshold of a

region was set at an LRT = 19.5 (p-value $< 1 \times 10^{-5}$).

Results

Simulation: SNP-RHM, Hap-RHM and SNHap-RHM

SNP-RHM and Hap-RHM: We performed a regional GREML analysis that fits two GRMs

(one for the region and one for the rest of the genome) per region across multiple genomic

regions delimited by recombination hotspots. We tested two types of regional GREML

models, SNP-RHM and Hap-RHM, on 20 replicates of five simulated phenotypes. In SNP-RHM,

the regional matrix is derived from SNP genotypes whereas in Hap-RHM the regional matrix

is derived from haplotypes. The phenotypes were simulated to be determined by 20 regional

QTL effects and genome-wide polygenic effects. The regional QTL effects of the five

phenotypes were simulated using SNPs as causal variants for two of them and haplotypes for

the remaining three as described in the methods section.

The likelihood ratio test (LRT) was used to test the null hypothesis, H_0: that the genetic

variance explained by the region is not significant, against the alternative hypothesis, H_1: that

the region accounts for a significant proportion of the phenotypic variance. A large LRT

statistic is evidence against the null hypothesis, and therefore means the region explains a

significant proportion of the phenotypic variance.

The LRTs averaged over the 20 replicates of the five phenotypes are shown in Figure

2. The figure shows plots of average LRT for the QTL regions and ten adjacent regions (5 to

each side). The results show that both models detected the simulated regional effects at the

genome-wide significance level (LRT = 23.9) and can capture true causal loci in traits with

different genetic architectures. The LRTs were higher on average for the SNP-based model

(SNP-RHM) than the haplotype-based model (Hap-RHM). This could be because for Hap-RHM,

the genome-wide GRM which is a SNP-based GRM does not tag any of the background

haplotype effects that are outside any one particular region being analysed, and thus the

residual variance may be inflated by the other haplotype QTLs which downwardly impact the

LRTs.

We provide further investigation of the results from the simulation in the

supplementary data. For both analysis models, we have presented detailed results of the

relationships between the LRT statistics, region size, variance estimates and allele frequencies

(Supplementary Figures 3 – 10). We observed that the longer haplotype blocks had many

SNPs (and hence many, many haplotypes), and this impacted the estimation of the simulated

regional variance (Supplementary Figure 8). We, therefore, performed a hybrid-Hap-RHM

analysis that restricted the natural haplotype block sizes to 20 or fewer SNPs per haplotype

block. This hybrid-Hap-RHM was to investigate whether the regional variance is well captured

by Hap-RHM when shorter haplotypes are used. The hybrid-Hap-RHM underestimated the

regional variance for larger regions but did not offer any discernible improvement in the LRT

statistics (Supplementary Figure 9). The relationship between region size and estimated

variance was different between the Hap-RHM and hybrid-Hap-RHM, while we observed a

similar relationship between LRTs and the region size.

Both SNP-RHM and Hap-RHM fail to capture the simulated regional effects when the

simulated phenotype has a genetic architecture that does not match the analysis model, i.e.,

SNP or haplotype (Figure 3 and Supplementary Figure 1). These figures show the results for

the situation where the SNP QTL phenotypes were analysed with the haplotype-based model

(Hap-RHM) and the haplotype QTL phenotypes were analysed with the SNP-based model

(SNP-RHM). Both models fail to detect the simulated effects in such situations, therefore,

showing that the models complement each other since they capture effects due to different

types of genetic variants (i.e., tagged by SNPs or haplotypes).

To confirm that two models are complementary and thus independent of each other,

we implemented SNHap-RHM that fits the regional SNP and haplotype GRMs jointly, on a

replicate of each of the five simulated phenotypes. The significance of regional effects was

tested with an LRT with two degrees of freedom. The results are shown in Figure 4 and

confirm that the two models are complementary since even when fitted jointly, we can still

capture the simulated effects independently.

SNHap-RHM analysis of Height and MDD

The heritability estimates for height and MDD, calculated using the whole-genome

GRM, were 81.4% and 13.8% respectively. There were no overlaps between regions identified

as significant (tested with an LRT with one degree of freedom) by the haplotype and SNP-

based models for either of the two traits (Supplementary Figure 2). This reaffirms our

hypothesis shown by simulation that the Hap-RHM is complementary to SNP-RHM in mapping

associated genomic loci.

The regional heritability results for height and MDD are presented as plots of minus-

Log10 of the LRT p-values (Figures 5 and 6). The plots for the SNHap-RHM, SNP-RHM and Hap-

RHM analyses are shown.

The results for height show that nine regions passed the Bonferroni-corrected

genome-wide significance threshold in the analysis using SNP-RHM. No region was genome-

wide significant for height when analysed with Hap-RHM. Furthermore, these associations

still come up when SNPs and haplotypes in those regions are analysed jointly using SNHap-

RHM. There are GWAS reported genes that lie in or are within 400kb of these regions

(Supplementary Table 1).

For MDD, no region passed the Bonferroni-corrected genome-wide significance

threshold for the analysis done with the SNP-based and haplotype-based regional GREML

models (Figure 6). Three regions passed the suggestive significance threshold at p-value $<$

1×10^{-5} for Hap-RHM analysis of MDD. A further nine regions were significant at p-value $<$

5×10^{-5} for the haplotype-based analysis, and one region for the SNP-based analysis

(Supplementary Table 2). Figure 6 shows that when the two local GRMs are fitted jointly using

SNHap-RHM, the genomic regions associated with MDD can still be mapped. The associated

regions mapped by the haplotype-based model for MDD contains genes reported by GWAS

to be associated with several psychiatry phenotypes (Figure 6 and Supplementary Table 2).

The top associated region was within 400kb of the *DCC* gene. This gene is part of the NETRIN1

pathway, which has been reported to be associated with major depressive disorder in two

GWAS samples (Generation Scotland: Scottish Family Health Study and Psychiatric Genomics

Consortium) (Zeng et al., 2017). Zeng et al. (2017a) used a SNP-RHM guided by pathway

analysis (first uncover pathway association and then localise *DCC* within the pathway) to show

the *DCC* association with major depressive disorder.

A linear mixed effects model was used to test for association of the SNPs within the

suggestive significant region identified by the haplotype-based model on chromosome 3 for

MDD. The model tested for association of SNPs by fitting them individually in the model and

fitting a GRM to account for relatedness of individuals. The region on chromosome 3 was

chosen because there is a psychiatric phenotype associated gene, *MYRIP* (Luciano et al.,

2011), mapped to it, unlike the *DCC* region which has the gene outside the region. The results

are shown in Table 1. Five SNPs within this region are nominally significant at p-value < 0.05.

Four out of these five SNPs confer about 2% increased risk of the disease each. These four

SNPs lie within the *MYRIP* gene sequence. The *MYRIP* gene is expressed in the brain (Ganat et

al., 2012). A SNP in this gene is reported to be associated with brain processing speed in the

Lothian birth cohort (Luciano et al., 2011). Brain processing speed is an important cognitive

function that is compromised in psychiatric illness such as schizophrenia and depression, and

old age. Also, a SNP in the *MYRIP* gene region is associated with sleep duration (Gottlieb et

al., 2007). Sleep durations outside the normal range (both short sleep and long sleep) is

significantly associated with increased risk of depression (Mohan et al., 2017; Roberts and

Duong, 2014; Watson et al., 2014; Zhai et al., 2015). The *MYRIP* gene is also reported to have

a role in insulin secretion (Waselle et al., 2003) and low insulin levels have been linked to

depression (Greenwood et al., 2015; Pearson et al., 2010; Webb et al., 2017).

Comparison with published GWAS SNPs

For both traits, the SNPs in the regions that were significant at p-value $< 5 \times 10^{-5}$

were compared to SNPs reported in the GWAS catalogue (MacArthur et al., 2017) to be

significant for the two traits. The GWAS catalogue was accessed on the 15th of January 2021.

The results are presented in Table 2. The SNP-based and haplotype-based models identified

1,380 and 45 SNPs respectively for height, and 78 and 495 SNPs respectively for MDD taking

all SNPs within haplotype blocks significant at p-value $< 5 \times 10^{-5}$. Out of the 1,380 SNPs

identified for height by the SNP-based model, 57 SNPs spanning 20 haplotype regions were

in common with published GWAS results for height.

Discussion

We have proposed and implemented a genome-wide analytical method that analyses

genomic regions using a regional GREML model (Nagamine et al., 2012). We have since

extended this method to include haplotypes by fitting a regional haplotype-based GRM (Hap-

RHM) and also redefined genomic regions in our analysis to be delimited by recombination

hotspots generated using HapMap Phase II (Shirali et al., 2018). In this study, we build on our

previous regional GREML methods by exploring the properties of the SNP and haplotype-

based regional heritability mapping models by simulation and demonstrate that the two

variance components fitted are largely independent of each other (Supplementary Figure 2).

The novelty in this study shows the two models capture two different kinds of effects in terms

of genetic architecture and thus the two variance components can be fitted jointly (by fitting

the SNP and haplotype regional matrices together) in a joint marker regional heritability

mapping procedure called SNHap-RHM.

We hypothesised that the Hap-RHM would complement the SNP-RHM. We

investigated this hypothesis in a simulation study in which we simulated 20 replicates each of

two types of SNP QTL phenotypes and three types of haplotype QTL phenotypes. The results

show that the two GREML models can capture the effects of causal variants within genomic

loci associated with the phenotype analysed. The results also show that the two models are

specific about the type of causal effect they can capture, therefore, providing support for the

hypothesis that haplotype-based regional GREML models will complement SNP-based

regional GREML models. We provide further support for this hypothesis by fitting the two

GRMs jointly and showing (using a LRT with two degrees of freedom) that we can still capture

the simulated effects and real effects from real data.

We applied SNHap-RHM to height and MDD phenotypes from the Generation

Scotland: Scottish Family Health Study. Again, we draw comparisons between the effects

captured by the SNP-RHM and the Hap-RHM. The SNP-RHM identified more genome-wide

(GW) significant regions (p-value $< 1.02 \times 10^{-6}$) for height compared to MDD. Fifty-seven of

the SNPs identified for height by the SNP-RHM have been reported by other studies to be

associated with height. These SNPs spanned 20 genomic regions in the GS: SFHS cohort.

Height is a highly polygenic trait with many common genetic variants accounting for a majority

of the additive genetic variation (Yang et al., 2015). These common genetic variants may be

in LD with genotyped SNPs on SNP chips (these chips are disproportionately enriched for

common SNPs). Therefore, the SNP-based regional GREML model is better suited to capture

SNP loci in height compared to MDD. MDD is a very heterogeneous phenotype, and thus every

MDD case will have a set of genetic and non-genetic risk factors exclusive to them (Levinson

et al., 2014). These unique genetic risk factors will mean that at the population level, a lot of

the genetic variants driving the disease will be rare.

Three genomic regions were identified for MDD by the haplotype-based regional

GREML model with p-value $< 1 \times 10^{-5}$. The Hap-RHM works well for MDD because MDD is

believed to be driven by rare genetic variants, and the model can capture rare genetic

variants. The haplotype model can capture rare variants because of the LD between rare

variants (both typed and untyped) and the flanking variants that aggregate to form the

haplotypes within the genomic regions. There were no overlaps between regions identified

by the Hap-RHM and SNP-RHM for each of the two traits, which again support the hypothesis

that the two classes of models complement each other in mapping associated loci.

In both traits, the top significant regions we mapped at p-value $< 5 \times 10^{-5}$ had genes

mapped to those regions or within 400kb of those regions. For height, these genes have been

reported to be associated with height in humans (Gudbjartsson et al., 2008; Kichaev et al.,

2019; Lango Allen et al., 2010; Nagy et al., 2017; Tachmazidou et al., 2017; Weedon et al.,

2008; Wood et al., 2014). For MDD, these genes have been reported to be associated with

major depressive disorder and other psychiatry phenotypes (Arnau-Soler et al., 2019; Howard

et al., 2019; Liu et al., 2019; Luciano et al., 2011; Wray et al., 2018; Zeng et al., 2017). In one

of such regions for MDD, five SNPs within the region are individually significantly associated

with MDD at the nominal level (p-value < 0.05). Four of these SNPs lie within the gene

sequence of *MYRIP*, and they each confer 2% disease risk. A conventional GWAS analysis

would have missed these nominally associated SNPs because they will not reach genome-

wide (GW) significance. However, analysing these SNPs within the region as haplotypes, gave

us the power to detect the combined effect of these SNPs in the region at a suggestive-

significance level even with our relatively small sample size compared to recent genome-wide

association studies of MDD: 322,580 (Howard et al., 2018), and 480,359 (Wray et al., 2018).

The current study's primary strength is the ability of SNHap-RHM to incorporate SNP

and haplotype information jointly to map genomic regions that affect complex traits. This

gives SNHap-RHM a uniquely useful role to play in the future of complex traits analysis. The

plummeting costs of whole-genome resequencing (Caulfield et al., 2013) has shifted research

focus in GWA studies towards sequence data analysis (Höglund et al., 2019). Although whole-

genome sequence data analysis allows incorporating all the genetic variants that drive the

phenotypic variation, there may still be some variants whose individual effects may be too

small to be picked up in a conventional GWA analysis. However, regionally analysing sequence

information can help overcome this because multiple small-effect variants in a region can add

up to a substantial regional effect that can be captured by a regional SNP GRM or tagged by

a haplotype GRM. Moreover, by defining haplotype blocks using recombination hotspots,

whole-genome information can be summarised naturally without setting arbitrary blocks.

More so, regional heritability analysis of sequence data would be an efficient way to deal with

the burden of multiple testing which has long been a problem of conventional GWAS.

One limitation of the current study is the computation burden of the analyses which

necessitates the pre-correction of the phenotypes with the whole-genome GRM before

performing SNHap-RHM. This step involved 22 separate GREML analyses each fitting a whole-

genome GRM that excluded SNPs from one chromosome. Although this was done to speed

up the analysis, this step was used as an approximation to account for the background

polygenic effects of genetic markers outside each region; this would have been about 48,772

separate GREMLs to account for each region. Also, although this study thoroughly evaluates

the robustness of SNP and Haplotype RHM using simulation and demonstrates the utility of

SNHap-RHM in real phenotype analysis, seeking replication in other cohorts could have

improved our understanding and more importantly demonstrate that the analysis is portable

across studies and genotyping platforms.

Conclusions

We have implemented a regional GREML analysis and undertaken analyses of regions

in the genome delimited by recombination boundaries and shown by simulation that

haplotype-based GRMs can capture genetic variance that may be missed by conventional

SNP-based GRMs. We then applied this method in the analysis of real phenotype data from

GS: SFHS. Again, we show that the haplotype-based regional GREML model uncovers

associations in regions of the genome that explain genetic variance missed by the SNP-based

GREML model. In light of this, we went further to show that regional effects can still be

captured when the two regional GRMs (SNP and haplotype-based) are fitted jointly: an

analytical procedure we termed as SNHap-RHM. This SNHap-RHM presents an exciting new

opportunity to analyse complex traits by allowing the joint mapping of novel genomic regions

tagged by either SNPs or haplotypes, potentially leading to the recovery of some of the

"missing" heritability.

Acknowledgements

Generation Scotland received core support from the Chief Scientist Office of the Scottish

Government Health Directorates [CZD/16/6] and the Scottish Funding Council [HR03006].

Genotyping of the GS: SFHS samples was carried out by the Genetics Core Laboratory at the

Wellcome Trust Clinical Research Facility, Edinburgh, Scotland and was funded by the Medical

Research Council UK and the Wellcome Trust (Wellcome Trust Strategic Award "STratifying

Resilience and Depression Longitudinally" (STRADL) Reference 104036/Z/14/Z). The research

was conducted using the UK Biobank Resource under Application Number 19655. We

acknowledge the Medical Research Council (MRC) UK for funding (grants MC_UU_00007/10,

MC_PC_U127592696 and MC_PC_U127561128). We are grateful to all the families who took

part, the general practitioners and the Scottish School of Primary Care for their help in

recruiting them, and the whole Generation Scotland team, which includes interviewers,

computer and laboratory technicians, clerical workers, research scientists, volunteers,

managers, receptionists, healthcare assistants and nurses.

Conflict of interest

The authors declare no conflict of interest.

References

Amador, C., Huffman, J., Trochet, H., Campbell, A., Porteous, D., Wilson, J.F., Hastie, N.,
Vitart, V., Hayward, C., Navarro, P., Haley, C.S., 2015. Recent genomic heritage in
Scotland. BMC Genomics 16, 1–17. https://doi.org/10.1186/s12864-015-1605-2

Arnau-Soler, A., Macdonald-Dunlop, E., Adams, M.J., Clarke, T.-K., MacIntyre, D.J., Milburn,
K., Navrady, L., Generation Scotland, Major Depressive Disorder Working Group of
the Psychiatric Genomics Consortium, Hayward, C., McIntosh, A.M., Thomson, P.A.,
2019. Genome-wide by environment interaction studies of depressive symptoms and
psychosocial stress in UK Biobank and Generation Scotland. Transl Psychiatry 9, 14.
https://doi.org/10.1038/s41398-018-0360-y

Balding, D.J., 2006. A tutorial on statistical methods for population association studies.
Nature Reviews Genetics 7, 781–791. https://doi.org/10.1038/nrg1916

Caulfield, T., Evans, J., McGuire, A., McCabe, C., Bubela, T., Cook-Deegan, R., Fishman, J.,
Hogarth, S., Miller, F.A., Ravitsky, V., Biesecker, B., Borry, P., Cho, M.K., Carroll, J.C.,
Etchegary, H., Joly, Y., Kato, K., Lee, S.S.-J., Rothenberg, K., Sankar, P., Szego, M.J.,
Ossorio, P., Pullman, D., Rousseau, F., Ungar, W.J., Wilson, B., 2013. Reflections on
the Cost of "Low-Cost" Whole Genome Sequencing: Framing the Health Policy
Debate. PLOS Biology 11, e1001699. https://doi.org/10.1371/journal.pbio.1001699

Cirulli, E.T., Goldstein, D.B., 2010. Uncovering the roles of rare variants in common disease
through whole-genome sequencing. Nat Rev Genet 11, 415–425.
https://doi.org/10.1038/nrg2779

Clarke, A.J., Cooper, D.N., 2010. GWAS: heritability missing in action? Eur J Hum Genet 18,
859–861. https://doi.org/10.1038/ejhg.2010.35

Delaneau, O., Zagury, J.-F., Marchini, J., 2013. Improved whole-chromosome phasing for
disease and population genetic studies. Nature Methods 10, 5–6.
https://doi.org/10.1038/nmeth.2307

First, M.B., Spitzer, R.L., Gibbon, M., Williams, J.B.W., 2002. Structured Clinical Interview for
DSM-IV-TR Axis I Disorders, Research Version, Non-patient Edition.

Ganat, Y.M., Calder, E.L., Kriks, S., Nelander, J., Tu, E.Y., Jia, F., Battista, D., Harrison, N.,
Parmar, M., Tomishima, M.J., Rutishauser, U., Studer, L., 2012. Identification of
embryonic stem cell-derived midbrain dopaminergic neurons for engraftment. J.
Clin. Invest. 122, 2928–2939. https://doi.org/10.1172/JCI58767

Gonzalez-Recio, O., Daetwyler, H.D., MacLeod, I.M., Pryce, J.E., Bowman, P.J., Hayes, B.J.,
Goddard, M.E., 2015. Rare Variants in Transcript and Potential Regulatory Regions
Explain a Small Percentage of the Missing Heritability of Complex Traits in Cattle.
PLoS ONE 10, e0143945. https://doi.org/10.1371/journal.pone.0143945

Gottlieb, D.J., O'Connor, G.T., Wilk, J.B., 2007. Genome-wide association of sleep and
circadian phenotypes. BMC Med Genet 8, S9. https://doi.org/10.1186/1471-2350-8-
S1-S9

Greenwood, E.A., Pasch, L.A., Shinkai, K., Cedars, M.I., Huddleston, H.G., 2015. Putative role for insulin resistance in depression risk in polycystic ovary syndrome. Fertility and Sterility 104, 707-714.e1. https://doi.org/10.1016/j.fertnstert.2015.05.019

Gudbjartsson, D.F., Walters, G.B., Thorleifsson, G., Stefansson, H., Halldorsson, B.V., Zusmanovich, P., Sulem, P., Thorlacius, S., Gylfason, A., Steinberg, S., Helgadottir, A., Ingason, A., Steinthorsdottir, V., Olafsdottir, E.J., Olafsdottir, G.H., Jonsson, T., Borch-Johnsen, K., Hansen, T., Andersen, G., Jorgensen, T., Pedersen, O., Aben, K.K., Witjes, J.A., Swinkels, D.W., den Heijer, M., Franke, B., Verbeek, A.L.M., Becker, D.M., Yanek, L.R., Becker, L.C., Tryggvadottir, L., Rafnar, T., Gulcher, J., Kiemeney, L.A., Kong, A., Thorsteinsdottir, U., Stefansson, K., 2008. Many sequence variants affecting diversity of adult human height. Nat Genet 40, 609–615. https://doi.org/10.1038/ng.122

Höglund, J., Rafati, N., Rask-Andersen, M., Enroth, S., Karlsson, T., Ek, W.E., Johansson, Å., 2019. Improved power and precision with whole genome sequencing data in genome-wide association studies of inflammatory biomarkers. Scientific Reports 9, 16844. https://doi.org/10.1038/s41598-019-53111-7

Howard, D.M., Adams, M.J., Clarke, T.-K., Hafferty, J.D., Gibson, J., Shirali, M., Coleman, J.R.I., Hagenaars, S.P., Ward, J., Wigmore, E.M., Alloza, C., Shen, X., Barbu, M.C., Xu, E.Y., Whalley, H.C., Marioni, R.E., Porteous, D.J., Davies, G., Deary, I.J., Hemani, G., Berger, K., Teismann, H., Rawal, R., Arolt, V., Baune, B.T., Dannlowski, U., Domschke, K., Tian, C., Hinds, D.A., 23andMe Research Team, Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium, Trzaskowski, M., Byrne, E.M., Ripke, S., Smith, D.J., Sullivan, P.F., Wray, N.R., Breen, G., Lewis, C.M., McIntosh, A.M., 2019. Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. Nat Neurosci 22, 343–352. https://doi.org/10.1038/s41593-018-0326-7

Howard, D.M., Adams, M.J., Shirali, M., Clarke, T.-K., Marioni, R.E., Davies, G., Coleman, J.R.I., Alloza, C., Shen, X., Barbu, M.C., Wigmore, E.M., Gibson, J., Hagenaars, S.P., Lewis, C.M., Ward, J., Smith, D.J., Sullivan, P.F., Haley, C.S., Breen, G., Deary, I.J., McIntosh, A.M., 2018. Genome-wide association study of depression phenotypes in UK Biobank identifies variants in excitatory synaptic pathways. Nature Communications 9, 1470. https://doi.org/10.1038/s41467-018-03819-3

International Human Genome Sequencing Consortium, 2004. Finishing the euchromatic sequence of the human genome. Nature 431, 931–945. https://doi.org/10.1038/nature03001

Kichaev, G., Bhatia, G., Loh, P.-R., Gazal, S., Burch, K., Freund, M.K., Schoech, A., Pasaniuc, B., Price, A.L., 2019. Leveraging Polygenic Functional Enrichment to Improve GWAS Power. Am J Hum Genet 104, 65–75. https://doi.org/10.1016/j.ajhg.2018.11.008

Lango Allen, H., Estrada, K., Lettre, G., Berndt, S.I., Weedon, M.N., Rivadeneira, F., Willer, C.J., Jackson, A.U., Vedantam, S., Raychaudhuri, S., Ferreira, T., Wood, A.R., Weyant, R.J., Segrè, A.V., Speliotes, E.K., Wheeler, E., Soranzo, N., Park, J.-H., Yang, J., Gudbjartsson, D., Heard-Costa, N.L., Randall, J.C., Qi, L., Vernon Smith, A., Mägi, R., Pastinen, T., Liang, L., Heid, I.M., Luan, J., Thorleifsson, G., Winkler, T.W., Goddard, M.E., Sin Lo, K., Palmer, C., Workalemahu, T., Aulchenko, Y.S., Johansson, A., Zillikens, M.C., Feitosa, M.F., Esko, T., Johnson, T., Ketkar, S., Kraft, P., Mangino, M., Prokopenko, I., Absher, D., Albrecht, E., Ernst, F., Glazer, N.L., Hayward, C., Hottenga, J.-J., Jacobs, K.B., Knowles, J.W., Kutalik, Z., Monda, K.L., Polasek, O., Preuss, M., Rayner, N.W., Robertson, N.R., Steinthorsdottir, V., Tyrer, J.P., Voight, B.F., Wiklund,

F., Xu, J., Zhao, J.H., Nyholt, D.R., Pellikka, N., Perola, M., Perry, J.R.B., Surakka, I.,
Tammesoo, M.-L., Altmaier, E.L., Amin, N., Aspelund, T., Bhangale, T., Boucher, G.,
Chasman, D.I., Chen, C., Coin, L., Cooper, M.N., Dixon, A.L., Gibson, Q., Grundberg, E.,
Hao, K., Juhani Junttila, M., Kaplan, L.M., Kettunen, J., König, I.R., Kwan, T., Lawrence,
R.W., Levinson, D.F., Lorentzon, M., McKnight, B., Morris, A.P., Müller, M., Suh Ngwa,
J., Purcell, S., Rafelt, S., Salem, R.M., Salvi, E., Sanna, S., Shi, J., Sovio, U., Thompson,
J.R., Turchin, M.C., Vandenput, L., Verlaan, D.J., Vitart, V., White, C.C., Ziegler, A.,
Almgren, P., Balmforth, A.J., Campbell, H., Citterio, L., De Grandi, A., Dominiczak, A.,
Duan, J., Elliott, P., Elosua, R., Eriksson, J.G., Freimer, N.B., Geus, E.J.C., Glorioso, N.,
Haiqing, S., Hartikainen, A.-L., Havulinna, A.S., Hicks, A.A., Hui, J., Igl, W., Illig, T., Jula,
A., Kajantie, E., Kilpeläinen, T.O., Koiranen, M., Kolcic, I., Koskinen, S., Kovacs, P.,
Laitinen, J., Liu, J., Lokki, M.-L., Marusic, A., Maschio, A., Meitinger, T., Mulas, A.,
Paré, G., Parker, A.N., Peden, J.F., Petersmann, A., Pichler, I., Pietiläinen, K.H., Pouta,
A., Ridderstråle, M., Rotter, J.I., Sambrook, J.G., Sanders, A.R., Schmidt, C.O., Sinisalo,
J., Smit, J.H., Stringham, H.M., Bragi Walters, G., Widen, E., Wild, S.H., Willemsen, G.,
Zagato, L., Zgaga, L., Zitting, P., Alavere, H., Farrall, M., McArdle, W.L., Nelis, M.,
Peters, M.J., Ripatti, S., van Meurs, J.B.J., Aben, K.K., Ardlie, K.G., Beckmann, J.S.,
Beilby, J.P., Bergman, R.N., Bergmann, S., Collins, F.S., Cusi, D., den Heijer, M.,
Eiriksdottir, G., Gejman, P.V., Hall, A.S., Hamsten, A., Huikuri, H.V., Iribarren, C.,
Kähönen, M., Kaprio, J., Kathiresan, S., Kiemeney, L., Kocher, T., Launer, L.J.,
Lehtimäki, T., Melander, O., Mosley, T.H., Musk, A.W., Nieminen, M.S., O'Donnell,
C.J., Ohlsson, C., Oostra, B., Palmer, L.J., Raitakari, O., Ridker, P.M., Rioux, J.D.,
Rissanen, A., Rivolta, C., Schunkert, H., Shuldiner, A.R., Siscovick, D.S., Stumvoll, M.,
Tönjes, A., Tuomilehto, J., van Ommen, G.-J., Viikari, J., Heath, A.C., Martin, N.G.,
Montgomery, G.W., Province, M.A., Kayser, M., Arnold, A.M., Atwood, L.D.,
Boerwinkle, E., Chanock, S.J., Deloukas, P., Gieger, C., Grönberg, H., Hall, P.,
Hattersley, A.T., Hengstenberg, C., Hoffman, W., Lathrop, G.M., Salomaa, V.,
Schreiber, S., Uda, M., Waterworth, D., Wright, A.F., Assimes, T.L., Barroso, I.,
Hofman, A., Mohlke, K.L., Boomsma, D.I., Caulfield, M.J., Cupples, L.A., Erdmann, J.,
Fox, C.S., Gudnason, V., Gyllensten, U., Harris, T.B., Hayes, R.B., Jarvelin, M.-R.,
Mooser, V., Munroe, P.B., Ouwehand, W.H., Penninx, B.W., Pramstaller, P.P.,
Quertermous, T., Rudan, I., Samani, N.J., Spector, T.D., Völzke, H., Watkins, H.,
Wilson, J.F., Groop, L.C., Haritunians, T., Hu, F.B., Kaplan, R.C., Metspalu, A., North,
K.E., Schlessinger, D., Wareham, N.J., Hunter, D.J., O'Connell, J.R., Strachan, D.P.,
Wichmann, H.-E., Borecki, I.B., van Duijn, C.M., Schadt, E.E., Thorsteinsdottir, U.,
Peltonen, L., Uitterlinden, A.G., Visscher, P.M., Chatterjee, N., Loos, R.J.F., Boehnke,
M., McCarthy, M.I., Ingelsson, E., Lindgren, C.M., Abecasis, G.R., Stefansson, K.,
Frayling, T.M., Hirschhorn, J.N., 2010. Hundreds of variants clustered in genomic loci
and biological pathways affect human height. Nature 467, 832–838.
https://doi.org/10.1038/nature09410
Levinson, D.F., Mostafavi, S., Milaneschi, Y., Rivera, M., Ripke, S., Wray, N.R., Sullivan, P.F.,
2014. Genetic studies of major depressive disorder: Why are there no GWAS
findings, and what can we do about it? Biol Psychiatry 76, 510–512.
https://doi.org/10.1016/j.biopsych.2014.07.029
Liu, M., Jiang, Y., Wedow, R., Li, Y., Brazel, D.M., Chen, F., Datta, G., Davila-Velderrain, J.,
McGuire, D., Tian, C., Zhan, X., Choquet, H., Docherty, A.R., Faul, J.D., Foerster, J.R.,
Fritsche, L.G., Gabrielsen, M.E., Gordon, S.D., Haessler, J., Hottenga, J.-J., Huang, H.,

644 Jang, S.-K., Jansen, P.R., Ling, Y., Mägi, R., Matoba, N., McMahon, G., Mulas, A., Orrù,
645  V., Palviainen, T., Pandit, A., Reginsson, G.W., Skogholt, A.H., Smith, J.A., Taylor, A.E.,
646  Turman, C., Willemsen, G., Young, H., Young, K.A., Zajac, G.J.M., Zhao, W., Zhou, W.,
647  Bjornsdottir, G., Boardman, J.D., Boehnke, M., Boomsma, D.I., Chen, C., Cucca, F.,
648  Davies, G.E., Eaton, C.B., Ehringer, M.A., Esko, T., Fiorillo, E., Gillespie, N.A.,
649  Gudbjartsson, D.F., Haller, T., Harris, K.M., Heath, A.C., Hewitt, J.K., Hickie, I.B.,
650  Hokanson, J.E., Hopfer, C.J., Hunter, D.J., Iacono, W.G., Johnson, E.O., Kamatani, Y.,
651  Kardia, S.L.R., Keller, M.C., Kellis, M., Kooperberg, C., Kraft, P., Krauter, K.S., Laakso,
652  M., Lind, P.A., Loukola, A., Lutz, S.M., Madden, P.A.F., Martin, N.G., McGue, M.,
653  McQueen, M.B., Medland, S.E., Metspalu, A., Mohlke, K.L., Nielsen, J.B., Okada, Y.,
654  Peters, U., Polderman, T.J.C., Posthuma, D., Reiner, A.P., Rice, J.P., Rimm, E., Rose,
655  R.J., Runarsdottir, V., Stallings, M.C., Stančáková, A., Stefansson, H., Thai, K.K., Tindle,
656  H.A., Tyrfingsson, T., Wall, T.L., Weir, D.R., Weisner, C., Whitfield, J.B., Winsvold, B.S.,
657  Yin, J., Zuccolo, L., Bierut, L.J., Hveem, K., Lee, J.J., Munafo, M.R., Saccone, N.L.,
658  Willer, C.J., Cornelis, M.C., David, S.P., Hinds, D., Jorgenson, E., Kaprio, J., Stitzel, J.A.,
659  Stefansson, K., Thorgeirsson, T.E., Abecasis, G., Liu, D.J., Vrieze, S., 2019. Association
660  studies of up to 1.2 million individuals yield new insights into the genetic etiology of
661  tobacco and alcohol use. Nat Genet 51, 237–244. https://doi.org/10.1038/s41588-
662  018-0307-5
663 Luciano, M., Hansell, N.K., Lahti, J., Davies, G., Medland, S.E., Räikkönen, K., Tenesa, A.,
664  Widen, E., McGhee, K.A., Palotie, A., Liewald, D., Porteous, D.J., Starr, J.M.,
665  Montgomery, G.W., Martin, N.G., Eriksson, J.G., Wright, M.J., Deary, I.J., 2011. Whole
666  genome association scan for genetic polymorphisms influencing information
667  processing speed. Biol Psychol 86, 193–202.
668  https://doi.org/10.1016/j.biopsycho.2010.11.008
669 MacArthur, J., Bowler, E., Cerezo, M., Gil, L., Hall, P., Hastings, E., Junkins, H., McMahon, A.,
670  Milano, A., Morales, J., Pendlington, Z.M., Welter, D., Burdett, T., Hindorff, L., Flicek,
671  P., Cunningham, F., Parkinson, H., 2017. The new NHGRI-EBI Catalog of published
672  genome-wide association studies (GWAS Catalog). Nucleic Acids Res 45, D896–D901.
673  https://doi.org/10.1093/nar/gkw1133
674 Maher, B., 2008. Personal genomes: The case of the missing heritability. Nature News 456,
675  18–21. https://doi.org/10.1038/456018a
676 Manolio, T.A., Collins, F.S., Cox, N.J., Goldstein, D.B., Hindorff, L.A., Hunter, D.J., McCarthy,
677  M.I., Ramos, E.M., Cardon, L.R., Chakravarti, A., Cho, J.H., Guttmacher, A.E., Kong, A.,
678  Kruglyak, L., Mardis, E., Rotimi, C.N., Slatkin, M., Valle, D., Whittemore, A.S.,
679  Boehnke, M., Clark, A.G., Eichler, E.E., Gibson, G., Haines, J.L., Mackay, T.F.C.,
680  McCarroll, S.A., Visscher, P.M., 2009. Finding the missing heritability of complex
681  diseases. Nature 461, 747–753. https://doi.org/10.1038/nature08494
682 Mohan, J., Xiaofan, G., Yingxian, S., 2017. Association between sleep time and depression: a
683  cross-sectional study from countries in rural Northeastern China. J Int Med Res 45,
684  984–992. https://doi.org/10.1177/0300060517701034
685 Nagamine, Y., Pong-Wong, R., Navarro, P., Vitart, V., Hayward, C., Rudan, I., Campbell, H.,
686  Wilson, J., Wild, S., Hicks, A.A., Pramstaller, P.P., Hastie, N., Wright, A.F., Haley, C.S.,
687  2012. Localising Loci underlying Complex Trait Variation Using Regional Genomic
688  Relationship Mapping. PLoS ONE 7, e46501.
689  https://doi.org/10.1371/journal.pone.0046501

Nagy, R., Boutin, T.S., Marten, J., Huffman, J.E., Kerr, S.M., Campbell, A., Evenden, L., Gibson,
J., Amador, C., Howard, D.M., Navarro, P., Morris, A., Deary, I.J., Hocking, L.J.,
Padmanabhan, S., Smith, B.H., Joshi, P., Wilson, J.F., Hastie, N.D., Wright, A.F.,
McIntosh, A.M., Porteous, D.J., Haley, C.S., Vitart, V., Hayward, C., 2017. Exploration
of haplotype research consortium imputation for genome-wide association studies in
20,032 Generation Scotland participants. Genome Med 9, 23.
https://doi.org/10.1186/s13073-017-0414-4
Pearson, S., Schmidt, M., Patton, G., Dwyer, T., Blizzard, L., Otahal, P., Venn, A., 2010.
Depression and Insulin Resistance. Diabetes Care 33, 1128–1133.
https://doi.org/10.2337/dc09-1940
Pritchard, J.K., 2001. Are rare variants responsible for susceptibility to complex diseases?
Am. J. Hum. Genet. 69, 124–137. https://doi.org/10.1086/321272
Roberts, R.E., Duong, H.T., 2014. The Prospective Association between Sleep Deprivation
and Depression among Adolescents. Sleep 37, 239–244.
https://doi.org/10.5665/sleep.3388
Shirali, M., Knott, S.A., Pong-Wong, R., Navarro, P., Haley, C.S., 2018. Haplotype Heritability
Mapping Method Uncovers Missing Heritability of Complex Traits. Scientific Reports
8, 4982. https://doi.org/10.1038/s41598-018-23307-4
Smith, B.H., Campbell, A., Linksted, P., Fitzpatrick, B., Jackson, C., Kerr, S.M., 2012. Cohort
profile: Generation Scotland: Scottish Family Health Study (GS:SFHS). The study, its
participants and their potential for genetic research on health and illness. Int J
Epidemiol 42. https://doi.org/10.1093/ije/dys084
Smith, B.H., Campbell, H., Blackwood, D., Connell, J., Connor, M., Deary, I.J., 2006.
Generation Scotland: the Scottish Family Health Study; a new resource for
researching genes and heritability. BMC Med Genet 7.
https://doi.org/10.1186/1471-2350-7-74
Speed, D., Hemani, G., Johnson, M.R., Balding, D.J., 2012. Improved Heritability Estimation
from Genome-wide SNPs. The American Journal of Human Genetics 91, 1011–1021.
https://doi.org/10.1016/j.ajhg.2012.10.010
Tachmazidou, I., Süveges, D., Min, J.L., Ritchie, G.R.S., Steinberg, J., Walter, K., Iotchkova, V.,
Schwartzentruber, J., Huang, J., Memari, Y., McCarthy, S., Crawford, A.A., Bombieri,
C., Cocca, M., Farmaki, A.-E., Gaunt, T.R., Jousilahti, P., Kooijman, M.N., Lehne, B.,
Malerba, G., Männistö, S., Matchan, A., Medina-Gomez, C., Metrustry, S.J., Nag, A.,
Ntalla, I., Paternoster, L., Rayner, N.W., Sala, C., Scott, W.R., Shihab, H.A., Southam,
L., St Pourcain, B., Traglia, M., Trajanoska, K., Zaza, G., Zhang, W., Artigas, M.S.,
Bansal, N., Benn, M., Chen, Z., Danecek, P., Lin, W.-Y., Locke, A., Luan, J., Manning,
A.K., Mulas, A., Sidore, C., Tybjaerg-Hansen, A., Varbo, A., Zoledziewska, M., Finan,
C., Hatzikotoulas, K., Hendricks, A.E., Kemp, J.P., Moayyeri, A., Panoutsopoulou, K.,
Szpak, M., Wilson, S.G., Boehnke, M., Cucca, F., Di Angelantonio, E., Langenberg, C.,
Lindgren, C., McCarthy, M.I., Morris, A.P., Nordestgaard, B.G., Scott, R.A., Tobin,
M.D., Wareham, N.J., SpiroMeta Consortium, GoT2D Consortium, Burton, P.,
Chambers, J.C., Smith, G.D., Dedoussis, G., Felix, J.F., Franco, O.H., Gambaro, G.,
Gasparini, P., Hammond, C.J., Hofman, A., Jaddoe, V.W.V., Kleber, M., Kooner, J.S.,
Perola, M., Relton, C., Ring, S.M., Rivadeneira, F., Salomaa, V., Spector, T.D., Stegle,
O., Toniolo, D., Uitterlinden, A.G., arcOGEN Consortium, Understanding Society
Scientific Group, UK10K Consortium, Barroso, I., Greenwood, C.M.T., Perry, J.R.B.,
Walker, B.R., Butterworth, A.S., Xue, Y., Durbin, R., Small, K.S., Soranzo, N., Timpson,

N.J., Zeggini, E., 2017. Whole-Genome Sequencing Coupled to Imputation Discovers

Genetic Signals for Anthropometric Traits. Am J Hum Genet 100, 865–884.

https://doi.org/10.1016/j.ajhg.2017.04.014

Vormfelde, S.V., Brockmöller, J., 2007. On the value of haplotype-based genotype-

phenotype analysis and on data transformation in pharmacogenetics and -genomics.

Nat. Rev. Genet. 8. https://doi.org/10.1038/nrg1916-c1

Wainschtein, P., Jain, D.P., Yengo, L., Zheng, Z., TOPMed Anthropometry Working Group, T.-

O. for P.M.C., Cupples, L.A., Shadyab, A.H., McKnight, B., Shoemaker, B.M., Mitchell,

B.D., Psaty, B.M., Kooperberg, C., Roden, D., Darbar, D., Arnett, D.K., Regan, E.A.,

Boerwinkle, E., Rotter, J.I., Allison, M.A., McDonald, M.-L.N., Chung, M.K., Smith,

N.L., Ellinor, P.T., Vasan, R.S., Mathias, R.A., Rich, S.S., Heckbert, S.R., Redline, S.,

Guo, X., Chen, Y.-D.I., Liu, C.-T., Andrade, M. de, Yanek, L.R., Albert, C.M., Hernandez,

R.D., McGarvey, S.T., North, K.E., Lange, L.A., Weir, B.S., Laurie, C.C., Yang, J.,

Visscher, P.M., 2019. Recovery of trait heritability from whole genome sequence

data. bioRxiv 588020. https://doi.org/10.1101/588020

Waselle, L., Coppola, T., Fukuda, M., Iezzi, M., El-Amraoui, A., Petit, C., Regazzi, R., 2003.

Involvement of the Rab27 Binding Protein Slac2c/MyRIP in Insulin Exocytosis. Mol

Biol Cell 14, 4103–4113. https://doi.org/10.1091/mbc.E03-01-0022

Watson, N.F., Harden, K.P., Buchwald, D., Vitiello, M.V., Pack, A.I., Strachan, E., Goldberg, J.,

2014. Sleep Duration and Depressive Symptoms: A Gene-Environment Interaction.

Sleep 37, 351–358. https://doi.org/10.5665/sleep.3412

Webb, M., Davies, M., Ashra, N., Bodicoat, D., Brady, E., Webb, D., Moulton, C., Ismail, K.,

Khunti, K., 2017. The association between depressive symptoms and insulin

resistance, inflammation and adiposity in men and women. PLOS ONE 12, e0187448.

https://doi.org/10.1371/journal.pone.0187448

Weedon, M.N., Lango, H., Lindgren, C.M., Wallace, C., Evans, D.M., Mangino, M., Freathy,

R.M., Perry, J.R.B., Stevens, S., Hall, A.S., Samani, N.J., Shields, B., Prokopenko, I.,

Farrall, M., Dominiczak, A., Johnson, T., Bergmann, S., Beckmann, J.S., Vollenweider,

P., Waterworth, D.M., Mooser, V., Palmer, C.N.A., Morris, A.D., Ouwehand, W.H.,

Zhao, J.H., Li, S., Loos, R.J.F., Barroso, I., Deloukas, P., Sandhu, M.S., Wheeler, E.,

Soranzo, N., Inouye, M., Wareham, N.J., Caulfield, M., Munroe, P.B., Hattersley, A.T.,

McCarthy, M.I., Frayling, T.M., 2008. Genome-wide association analysis identifies 20

loci that influence adult height. Nat Genet 40, 575–583.

https://doi.org/10.1038/ng.121

Wood, A.R., Esko, T., Yang, J., Vedantam, S., Pers, T.H., Gustafsson, S., Chu, A.Y., Estrada, K.,

Luan, J., Kutalik, Z., Amin, N., Buchkovich, M.L., Croteau-Chonka, D.C., Day, F.R.,

Duan, Y., Fall, T., Fehrmann, R., Ferreira, T., Jackson, A.U., Karjalainen, J., Lo, K.S.,

Locke, A.E., Mägi, R., Mihailov, E., Porcu, E., Randall, J.C., Scherag, A., Vinkhuyzen,

A.A.E., Westra, H.-J., Winkler, T.W., Workalemahu, T., Zhao, J.H., Absher, D.,

Albrecht, E., Anderson, D., Baron, J., Beekman, M., Demirkan, A., Ehret, G.B.,

Feenstra, B., Feitosa, M.F., Fischer, K., Fraser, R.M., Goel, A., Gong, J., Justice, A.E.,

Kanoni, S., Kleber, M.E., Kristiansson, K., Lim, U., Lotay, V., Lui, J.C., Mangino, M.,

Leach, I.M., Medina-Gomez, C., Nalls, M.A., Nyholt, D.R., Palmer, C.D., Pasko, D.,

Pechlivanis, S., Prokopenko, I., Ried, J.S., Ripke, S., Shungin, D., Stancáková, A.,

Strawbridge, R.J., Sung, Y.J., Tanaka, T., Teumer, A., Trompet, S., Laan, S.W. van der,

Setten, J. van, Vliet-Ostaptchouk, J.V.V., Wang, Z., Yengo, L., Zhang, W., Afzal, U.,

Ärnlöv, J., Arscott, G.M., Bandinelli, S., Barrett, A., Bellis, C., Bennett, A.J., Berne, C.,

Blüher, M., Bolton, J.L., Böttcher, Y., Boyd, H.A., Bruinenberg, M., Buckley, B.M.,
Buyske, S., Caspersen, I.H., Chines, P.S., Clarke, R., Claudi-Boehm, S., Cooper, M.,
Daw, E.W., Jong, P.A.D., Deelen, J., Delgado, G., Denny, J.C., Dhonukshe-Rutten, R.,
Dimitriou, M., Doney, A.S.F., Dörr, M., Eklund, N., Eury, E., Folkersen, L., Garcia, M.E.,
Geller, F., Giedraitis, V., Go, A.S., Grallert, H., Grammer, T.B., Gräßler, J., Grönberg,
H., Groot, L.C.P.G.M. de, Groves, C.J., Haessler, J., Hall, P., Haller, T., Hallmans, G.,
Hannemann, A., Hartman, C.A., Hassinen, M., Hayward, C., Heard-Costa, N.L.,
Helmer, Q., Hemani, G., Henders, A.K., Hillege, H.L., Hlatky, M.A., Hoffmann, W.,
Hoffmann, P., Holmen, O., Houwing-Duistermaat, J.J., Illig, T., Isaacs, A., James, A.L.,
Jeff, J., Johansen, B., Johansson, Å., Jolley, J., Juliusdottir, T., Junttila, J., Kho, A.N.,
Kinnunen, L., Klopp, N., Kocher, T., Kratzer, W., Lichtner, P., Lind, L., Lindström, J.,
Lobbens, S., Lorentzon, M., Lu, Y., Lyssenko, V., Magnusson, P.K.E., Mahajan, A.,
Maillard, M., McArdle, W.L., McKenzie, C.A., McLachlan, S., McLaren, P.J., Menni, C.,
Merger, S., Milani, L., Moayyeri, A., Monda, K.L., Morken, M.A., Müller, G., Müller-
Nurasyid, M., Musk, A.W., Narisu, N., Nauck, M., Nolte, I.M., Nöthen, M.M.,
Oozageer, L., Pilz, S., Rayner, N.W., Renstrom, F., Robertson, N.R., Rose, L.M.,
Roussel, R., Sanna, S., Scharnagl, H., Scholtens, S., Schumacher, F.R., Schunkert, H.,
Scott, R.A., Sehmi, J., Seufferlein, T., Shi, J., Silventoinen, K., Smit, J.H., Smith, A.V.,
Smolonska, J., Stanton, A.V., Stirrups, K., Stott, D.J., Stringham, H.M., Sundström, J.,
Swertz, M.A., Syvänen, A.-C., Tayo, B.O., Thorleifsson, G., Tyrer, J.P., Dijk, S. van,
Schoor, N.M. van, Velde, N. van der, Heemst, D. van, Oort, F.V.A. van, Vermeulen,
S.H., Verweij, N., Vonk, J.M., Waite, L.L., Waldenberger, M., Wennauer, R., Wilkens,
L.R., Willenborg, C., Wilsgaard, T., Wojczynski, M.K., Wong, A., Wright, A.F., Zhang,
Q., Arveiler, D., Bakker, S.J.L., Beilby, J., Bergman, R.N., Bergmann, S., Biffar, R.,
Blangero, J., Boomsma, D.I., Bornstein, S.R., Bovet, P., Brambilla, P., Brown, M.J.,
Campbell, H., Caulfield, M.J., Chakravarti, A., Collins, R., Collins, F.S., Crawford, D.C.,
Cupples, L.A., Danesh, J., Faire, U. de, Ruijter, H.M. den, Erbel, R., Erdmann, J.,
Eriksson, J.G., Farrall, M., Ferrannini, E., Ferrières, J., Ford, I., Forouhi, N.G., Forrester,
T., Gansevoort, R.T., Gejman, P.V., Gieger, C., Golay, A., Gottesman, O., Gudnason,
V., Gyllensten, U., Haas, D.W., Hall, A.S., Harris, T.B., Hattersley, A.T., Heath, A.C.,
Hengstenberg, C., Hicks, A.A., Hindorff, L.A., Hingorani, A.D., Hofman, A., Hovingh,
G.K., Humphries, S.E., Hunt, S.C., Hypponen, E., Jacobs, K.B., Jarvelin, M.-R.,
Jousilahti, P., Jula, A.M., Kaprio, J., Kastelein, J.J.P., Kayser, M., Kee, F., Keinanen-
Kiukaanniemi, S.M., Kiemeney, L.A., Kooner, J.S., Kooperberg, C., Koskinen, S.,
Kovacs, P., Kraja, A.T., Kumari, M., Kuusisto, J., Lakka, T.A., Langenberg, C.,
Marchand, L.L., Lehtimäki, T., Lupoli, S., Madden, P.A.F., Männistö, S., Manunta, P.,
Marette, A., Matise, T.C., McKnight, B., Meitinger, T., Moll, F.L., Montgomery, G.W.,
Morris, A.D., Morris, A.P., Murray, J.C., Nelis, M., Ohlsson, C., Oldehinkel, A.J., Ong,
K.K., Ouwehand, W.H., Pasterkamp, G., Peters, A., Pramstaller, P.P., Price, J.F., Qi, L.,
Raitakari, O.T., Rankinen, T., Rao, D.C., Rice, T.K., Ritchie, M., Rudan, I., Salomaa, V.,
Samani, N.J., Saramies, J., Sarzynski, M.A., Schwarz, P.E.H., Sebert, S., Sever, P.,
Shuldiner, A.R., Sinisalo, J., Steinthorsdottir, V., Stolk, R.P., Tardif, J.-C., Tönjes, A.,
Tremblay, A., Tremoli, E., Virtamo, J., Vohl, M.-C., Consortium, T.E.M.R. and G.
(eMERGE), Consortium, T.Mig., Consortium, T.P., Study, T.L.C., Amouyel, P.,
Asselbergs, F.W., Assimes, T.L., Bochud, M., Boehm, B.O., Boerwinkle, E., Bottinger,
E.P., Bouchard, C., Cauchi, S., Chambers, J.C., Chanock, S.J., Cooper, R.S., Bakker,
P.I.W. de, Dedoussis, G., Ferrucci, L., Franks, P.W., Froguel, P., Groop, L.C., Haiman,

C.A., Hamsten, A., Hayes, M.G., Hui, J., Hunter, D.J., Hveem, K., Jukema, J.W., Kaplan,
R.C., Kivimaki, M., Kuh, D., Laakso, M., Liu, Y., Martin, N.G., März, W., Melbye, M.,
Moebus, S., Munroe, P.B., Njølstad, I., Oostra, B.A., Palmer, C.N.A., Pedersen, N.L.,
Perola, M., Pérusse, L., Peters, U., Powell, J.E., Power, C., Quertermous, T.,
Rauramaa, R., Reinmaa, E., Ridker, P.M., Rivadeneira, F., Rotter, J.I., Saaristo, T.E.,
Saleheen, D., Schlessinger, D., Slagboom, P.E., Snieder, H., Spector, T.D., Strauch, K.,
Stumvoll, M., Tuomilehto, J., Uusitupa, M., Harst, P. van der, Völzke, H., Walker, M.,
Wareham, N.J., Watkins, H., Wichmann, H.-E., Wilson, J.F., Zanen, P., Deloukas, P.,
Heid, I.M., Lindgren, C.M., Mohlke, K.L., Speliotes, E.K., Thorsteinsdottir, U., Barroso,
I., Fox, C.S., North, K.E., Strachan, D.P., Beckmann, J.S., Berndt, S.I., Boehnke, M.,
Borecki, I.B., McCarthy, M.I., Metspalu, A., Stefansson, K., Uitterlinden, A.G., Duijn,
C.M. van, Franke, L., Willer, C.J., Price, A.L., Lettre, G., Loos, R.J.F., Weedon, M.N.,
Ingelsson, E., O'Connell, J.R., Abecasis, G.R., Chasman, D.I., Goddard, M.E., Visscher,
P.M., Hirschhorn, J.N., Frayling, T.M., 2014. Defining the role of common variation in
the genomic and biological architecture of adult human height. Nature Genetics 46,
1173–1186. https://doi.org/10.1038/ng.3097
Wray, N.R., Ripke, S., Mattheisen, M., Trzaskowski, M., Byrne, E.M., Abdellaoui, A., Adams,
M.J., Agerbo, E., Air, T.M., Andlauer, T.M.F., Bacanu, S.-A., Bækvad-Hansen, M.,
Beekman, A.F.T., Bigdeli, T.B., Binder, E.B., Blackwood, D.R.H., Bryois, J.,
Buttenschøn, H.N., Bybjerg-Grauholm, J., Cai, N., Castelao, E., Christensen, J.H.,
Clarke, T.-K., Coleman, J.I.R., Colodro-Conde, L., Couvy-Duchesne, B., Craddock, N.,
Crawford, G.E., Crowley, C.A., Dashti, H.S., Davies, G., Deary, I.J., Degenhardt, F.,
Derks, E.M., Direk, N., Dolan, C.V., Dunn, E.C., Eley, T.C., Eriksson, N., Escott-Price, V.,
Kiadeh, F.H.F., Finucane, H.K., Forstner, A.J., Frank, J., Gaspar, H.A., Gill, M., Giusti-
Rodríguez, P., Goes, F.S., Gordon, S.D., Grove, J., Hall, L.S., Hannon, E., Hansen, C.S.,
Hansen, T.F., Herms, S., Hickie, I.B., Hoffmann, P., Homuth, G., Horn, C., Hottenga, J.-
J., Hougaard, D.M., Hu, M., Hyde, C.L., Ising, M., Jansen, R., Jin, F., Jorgenson, E.,
Knowles, J.A., Kohane, I.S., Kraft, J., Kretzschmar, W.W., Krogh, J., Kutalik, Z., Lane,
J.M., Li, Yihan, Li, Yun, Lind, P.A., Liu, X., Lu, L., MacIntyre, D.J., MacKinnon, D.F.,
Maier, R.M., Maier, W., Marchini, J., Mbarek, H., McGrath, P., McGuffin, P.,
Medland, S.E., Mehta, D., Middeldorp, C.M., Mihailov, E., Milaneschi, Y., Milani, L.,
Mill, J., Mondimore, F.M., Montgomery, G.W., Mostafavi, S., Mullins, N., Nauck, M.,
Ng, B., Nivard, M.G., Nyholt, D.R., O'Reilly, P.F., Oskarsson, H., Owen, M.J., Painter,
J.N., Pedersen, C.B., Pedersen, M.G., Peterson, R.E., Pettersson, E., Peyrot, W.J.,
Pistis, G., Posthuma, D., Purcell, S.M., Quiroz, J.A., Qvist, P., Rice, J.P., Riley, B.P.,
Rivera, M., Mirza, S.S., Saxena, R., Schoevers, R., Schulte, E.C., Shen, L., Shi, J., Shyn,
S.I., Sigurdsson, E., Sinnamon, G.B.C., Smit, J.H., Smith, D.J., Stefansson, H.,
Steinberg, S., Stockmeier, C.A., Streit, F., Strohmaier, J., Tansey, K.E., Teismann, H.,
Teumer, A., Thompson, W., Thomson, P.A., Thorgeirsson, T.E., Tian, C., Traylor, M.,
Treutlein, J., Trubetskoy, V., Uitterlinden, A.G., Umbricht, D., Auwera, S.V. der,
Hemert, A.M. van, Viktorin, A., Visscher, P.M., Wang, Y., Webb, B.T., Weinsheimer,
S.M., Wellmann, J., Willemsen, G., Witt, S.H., Wu, Y., Xi, H.S., Yang, J., Zhang, F.,
Arolt, V., Baune, B.T., Berger, K., Boomsma, D.I., Cichon, S., Dannlowski, U., Geus,
E.C.J. de, DePaulo, J.R., Domenici, E., Domschke, K., Esko, T., Grabe, H.J., Hamilton,
S.P., Hayward, C., Heath, A.C., Hinds, D.A., Kendler, K.S., Kloiber, S., Lewis, G., Li, Q.S.,
Lucae, S., Madden, P.F.A., Magnusson, P.K., Martin, N.G., McIntosh, A.M., Metspalu,
A., Mors, O., Mortensen, P.B., Müller-Myhsok, B., Nordentoft, M., Nöthen, M.M.,

O'Donovan, M.C., Paciga, S.A., Pedersen, N.L., Penninx, B.W.J.H., Perlis, R.H.,
Porteous, D.J., Potash, J.B., Preisig, M., Rietschel, M., Schaefer, C., Schulze, T.G.,
Smoller, J.W., Stefansson, K., Tiemeier, H., Uher, R., Völzke, H., Weissman, M.M.,
Werge, T., Winslow, A.R., Lewis, C.M., Levinson, D.F., Breen, G., Børglum, A.D.,
Sullivan, P.F., 2018. Genome-wide association analyses identify 44 risk variants and
refine the genetic architecture of major depression. Nature Genetics 50, 668–681.
https://doi.org/10.1038/s41588-018-0090-3
Yang, J., Bakshi, A., Zhu, Z., Hemani, G., Vinkhuyzen, A.A.E., Lee, S.H., Robinson, M.R., Perry,
J.R.B., Nolte, I.M., van Vliet-Ostaptchouk, J.V., Snieder, H., Esko, T., Milani, L., Mägi,
R., Metspalu, A., Hamsten, A., Magnusson, P.K.E., Pedersen, N.L., Ingelsson, E.,
Soranzo, N., Keller, M.C., Wray, N.R., Goddard, M.E., Visscher, P.M., 2015. Genetic
variance estimation with imputed variants finds negligible missing heritability for
human height and body mass index. Nature Genetics 47, 1114–1120.
https://doi.org/10.1038/ng.3390
Yang, J., Benyamin, B., McEvoy, B.P., Gordon, S., Henders, A.K., Nyholt, D.R., Madden, P.A.,
Heath, A.C., Martin, N.G., Montgomery, G.W., Goddard, M.E., Visscher, P.M., 2010.
Common SNPs explain a large proportion of the heritability for human height. Nat
Genet 42, 565–569. https://doi.org/10.1038/ng.608
Yang, J., Lee, S.H., Goddard, M.E., Visscher, P.M., 2011. GCTA: A Tool for Genome-wide
Complex Trait Analysis. The American Journal of Human Genetics 88, 76–82.
https://doi.org/10.1016/j.ajhg.2010.11.011
Zeng, Y., Navarro, P., Fernandez-Pujals, A.M., Hall, L.S., Clarke, T.-K., Thomson, P.A., Smith,
B.H., Hocking, L.J., Padmanabhan, S., Hayward, C., MacIntyre, D.J., Wray, N.R., Deary,
I.J., Porteous, D.J., Haley, C.S., McIntosh, A.M., 2017. A Combined Pathway and
Regional Heritability Analysis Indicates NETRIN1 Pathway Is Associated With Major
Depressive Disorder. Biol Psychiatry 81, 336–346.
https://doi.org/10.1016/j.biopsych.2016.04.017
Zhai, L., Zhang, H., Zhang, D., 2015. SLEEP DURATION AND DEPRESSION AMONG ADULTS: A
META-ANALYSIS OF PROSPECTIVE STUDIES. Depress Anxiety 32, 664–670.
https://doi.org/10.1002/da.22386

915 Table 1. SNP-based association test of MDD in the *MYRIP* gene region. The columns are the SNP ID,
916 chromosome, genome position of SNP, minor allele frequency, odds ratio, log of odds ratio, standard
917 error of log odds ratio and association p-value.

SNP information				Depression association			
SNP ID	Chr	Pos	MAF	OR	Log (OR)	SE (logOR)	p
rs9842160	3	39844703	0.14	0.97	-0.030	0.013	0.02
rs9858242	3	39847606	0.19	1.02	0.025	0.011	0.03
rs1599902	3	39954674	0.41	1.02	0.019	0.009	0.04
rs7618607	3	39947936	0.41	1.02	0.019	0.009	0.04
rs9860916	3	39944942	0.41	1.02	0.019	0.009	0.04

920 Table 2. Comparison of SNPs within significant regions identified by both models and published GWAS
921 results for height and MDD. The columns are the name of trait, number of SNPs in regions identified
922 by SNP-RHM and Hap-RHM with p-value $< 5 \times 10^{-5}$ and SNPs in published GWAS (pubGWAS) for the
923 traits, and the number of SNPs overlapping between the three.

Number of SNPS				Number of overlapping SNPS		
Trait	SNP-RHM	Hap-RHM	pubGWAS	SNP-RHM & Hap-RHM	SNP-RHM & pubGWAS	Hap-RHM & pubGWAS
Height	1380	45	4960	0	57	0
MDD	78	495	1815	0	0	0

Figure 1. A Schema outlying SNHap-RHM

Figure 2. Plots of Likelihood ratio test (LRT) statistics at each QTL loci and 5 regions either side averaged for the 20 simulations of each of the five QTL phenotypes. Plot (i) is SNP QTL phenotypes analysed using the SNP-RHM and plot (ii) is the haplotype QTL phenotypes analysed using the Hap-RHM. Both models can capture the simulated QTL effects for their respective SNP and haplotype phenotypes.

Figure 3. Plots of average LRT statistics over replicates of QTL loci across the chromosomes for the 20 simulations of each of the two SNP QTL phenotypes. The red dashed lines are genome-wide significance threshold (for 48,772 regions) and the black dashed lines are Bonferroni significance threshold (for 220 regions). The upper plot (i) is the 1-SNP QTL phenotype, and the lower plot (ii) is the multiple SNP QTL phenotype. The two phenotypes are analysed using both the SNP based model (SNP-RHM) (blue points) and the Haplotype based model (Hap-RHM) (red points). The Hap-RHM fails to capture the simulated effects for the SNP QTLs.

Figure 4. Joint analysis of the SNP and haplotype phenotypes using SNHap-RHM. The plot is an analysis of one replicate of each of the simulated phenotypes. The LRT statistics are plotted over QTL loci across the chromosomes. The red dashed lines are genome-wide significance threshold (for 48,772 regions) and the black dashed lines are Bonferroni significance threshold (for 220 regions).

Figure 5. The genome-wide evidence of haplotype block association for height. Analysis done with SNHap-RHM, SNP-RHM and Hap-RHM. The points are plots of -log10 of the p-values of regions tested with the LRT for the regional GREML analyses. The green lines are the Bonferroni-corrected genome-wide significance threshold and the red lines are the suggestive significance threshold calculated to be p-value $< 1 \times 10^{-5}$. The top association hits at p-value $< 5 \times 10^{-5}$ with genes located within the region are highlighted in blue for SNP-RHM and red for the Hap-RHM.

Figure 6. The genome-wide evidence of haplotype block association for Major Depressive Disorder. Analysis done with SNHap-RHM, SNP-RHM and Hap-RHM. The points are plots of -log10 of the p-values of regions tested with the LRT for the regional GREML analyses. The green lines are the Bonferroni-corrected genome-wide significance threshold and the red lines are the suggestive significance threshold calculated to be p-value $< 1 \times 10^{-5}$. The top association hits at p-value $< 5 \times 10^{-5}$ with genes located within the region are highlighted in blue for SNP-RHM and red for the Hap-RHM.