Microbial byproducts determine reproductive fitness of free-living and parasitic nematodes

Mericien Venzon[*1,2], Ritika Das[*2,4], Daniel J. Luciano[2,5], Julia Burnett[2], Hyun Shin Park[6], Joseph Cooper Devlin[5], Eric T. Kool[7], Joel G. Belasco[2,5], E. Jane Albert Hubbard[2,3,4,†], Ken Cadwell[2,5,8,†]

1. Vilcek Institute of Graduate Biomedical Sciences, New York University Grossman School of Medicine, New York, NY, USA
2. Kimmel Center for Biology and Medicine at the Skirball Institute, New York University Grossman School of Medicine, New York, NY, USA
3. Department of Pathology, New York University Grossman School of Medicine, New York, NY, USA
4. Department of Cell Biology, New York University Grossman School of Medicine, New York, NY, USA
5. Department of Microbiology, New York University Grossman School of Medicine, New York, NY, USA
6. Seegene Inc., Ogeum-ro, Songpa-Gu, Seoul, Republic of Korea
7. Department of Chemistry, Stanford Cancer Institute and ChEM-H Institute, Stanford University, Stanford, CA, USA
8. Division of Gastroenterology, Department of Medicine, New York University Grossman School of Medicine, New York, NY, USA

* These authors contributed equally

†Co-corresponding, equal contribution

Correspondence:

jane.hubbard@nyulangone.org

ken.cadwell@nyulangone.org

Summary

A distinguishing feature of *Trichuris* nematodes is that these parasitic worms reproduce within the digestive tracts of humans and other mammalian hosts shedding thousands of eggs daily, facilitating their sustained presence in the environment and hampering eradication efforts. Although this aspect of the lifecycle places *Trichuris* in a microbiota-rich environment, metabolic byproducts of bacteria that facilitate the reproductive development of parasites are unknown. Here, we employ a pipeline using the well-characterized free-living nematode *C. elegans* to identify microbial factors with conserved roles in the reproduction of nematodes. A screen for *E. coli* mutants that impair *C. elegans* fertility identified genes in fatty acid biosynthesis and ethanolamine utilization pathways, including *fabH* and *eutN*. *Trichuris muris* eggs displayed defective hatching in the presence of *E. coli* deficient in *fabH* or *eutN* due to reduction in arginine or elevated levels of aldehydes, respectively. Remarkably, *T. muris* reared in gnotobiotic mice colonized with these *E. coli* mutants displayed profound abnormalities including morphological defects and a failure to lay viable eggs. These findings indicate that microbial byproducts mediate evolutionarily conserved transkingdom interactions that impact the reproductive fitness of distantly-related nematodes.

Introduction

The reproductive success of the parasitic nematode *Trichuris trichiura* is evidenced by

the over 400 million individuals colonized by this soil-transmitted helminth (Vos et al.,

2016). Its enormous reproductive capacity contributes to its infectious spread and

hampers parasite eradication efforts. A distinct feature of *Trichuris* species is that

reproductive development, from egg hatching to egg laying, is completed exclusively

within the host digestive tract (Fahmy, 1954; Panesar, 1989). A new infection begins

when embryonated eggs ingested from the environment hatch in the host cecum, a

region within the gastrointestinal tract populated by a dense community of bacteria.

Although bacteria mediate egg hatching for *Trichuris muris* (Hayes et al., 2010;

Koyama, 2016; White et al., 2018), the *Trichuris* species that infects mice, a role for

bacteria at later stages of *Trichuris* development remains unclear. Our group and others

have shown that *Trichuris* infections in mice and humans affect the bacterial

composition of the gut microbiota (Holm et al., 2015; Houlden et al., 2015; Jenkins et

al., 2017; Lee et al., 2014; Ramanan et al., 2016; Rosa et al., 2021; Schachter et al.,

2020; White et al., 2018). Additionally, the *T. muris* microbiota harbors taxa similar to

those observed in the murine host microbiota, with a notable enrichment of

Proteobacteria, including Gram-negative commensals like *E. coli* (White et al., 2018).

For the free-living nematode *C. elegans*, reducing the quantity or quality of the

bacterial diet, or interfering with nutrient uptake, can impair germline development with

consequences for fertility (Dalfo et al., 2012; Gracida and Eckmann, 2013; Korta et al.,

2012; Pekar et al., 2017; Spanier et al., 2018; Starich et al., 2020). Because of the high

degree of conservation of body plan and neuromuscular system organization within the

phylum Nematoda, *C. elegans* has been used as a model for the discovery of

anthelmintics for more than four decades (Salinas and Risi, 2018; Sepulveda-Crespo et

al., 2020). Therefore, we hypothesized that bacteria-derived essential requirements for

reproductive development in the well-characterized, model organism *C. elegans* are

conserved in *Trichuris*.

Results

High-throughput screen identifies *E. coli* mutants that delay *C. elegans* fertility

To identify bacteria-derived factors that impact nematode reproduction, we

assessed fertility of young adult *C. elegans* raised in 96-well format on a library of viable

E. coli mutants suspended in liquid media. We used *C. elegans* bearing a temperature-

sensitive mutation in the gene encoding the GLP-1 Notch receptor, *glp-1(e2141),* to

facilitate quantification of fertility. At the semi-permissive temperature of 20ºC, the

number of germline progenitor cells is roughly half of the wild type (Michaelson et al.,

2010), thereby sensitizing the worms to fertility defects when signaling pathways

conveying nutritional sufficiency are altered (Dalfo et al., 2012; Korta et al., 2012). The

C. elegans strain also carried fluorescent markers for the pharynx and for embryos (Roy

et al., 2018) to facilitate counting and staging the worms, and to determine whether

individual worms contained embryos (gravid) (Fig. 1, A-B). Wells were excluded from

subsequent analyses if *E. coli* failed to grow, or if *C. elegans* exhibited developmental

arrest or severely delayed growth. Using a series of selection criteria that considered

both plate-by-plate statistical comparisons and penetrance, our primary screen of ~3000

of the 3985 *E. coli* mutants in the Keio library identified 315 bacterial mutants that

reduced the percentage of gravid worms at the established time point (Fig. 1C).

After these 315 *E. coli* mutants were re-screened in biological triplicate, ten

mutants significantly reduced the penetrance of embryo-bearing *C. elegans* compared

to our wild-type (WT) *E. coli* control strain (GC1547) in at least two out of three

biological replicates (Fig. 1C and S1, A-C). Seven of the ten *E. coli* genes fell into three

functional groups related to ethanolamine utilization, fatty acid biosynthesis or

lipopolysaccharide synthesis (Table S1). We focused on the *E. coli* mutants associated

with the two functional groups fatty acid biosynthesis and ethanolamine utilization:

ΔfabF, ΔfabH, ΔeutD, and *ΔeutN*. We pursued these functional groups because there

were two mutants each that affected these processes and these mutants displayed

consistent results across experiments, increasing our confidence that they represented

bona fide hits (Fig. 1D). Complementation of the *ΔfabH*, *ΔeutN,* and *ΔeutD* mutations

with a plasmid-borne copy of the corresponding gene confirmed their functional

relevance in the *C. elegans* fertility delay phenotype (Fig. S1D-E). For *ΔfabF*, we

rederived the *E. coli* deletion mutant by using P1 phage to transduce the original

ΔfabF::*kan* allele from the Keio library strain into the Keio progenitor strain BW25113

(Fig. S1, F-G). Similar to the original mutant, the transductant caused a significant

reduction in the percentage of embryo-bearing *C. elegans*.

To assess whether these *E. coli* mutants caused delays in somatic and/or

germline development of *C. elegans*, we performed two different assays. First, we

assayed somatic development using a time-course analysis of vulval development (Mok

et al., 2015). Among the four mutants, only worms fed *ΔeutN* exhibited delayed somatic

development (Fig. 1E). We also assayed for a delay in germline relative to somatic

development (Fig. 1F) using as an assay enhancement of the temperature-sensitive Pro

phenotype seen in a weak *glp-1(gf)* mutant (Pepper et al., 2003). Enhancement of Pro

at the permissive temperature often occurs as a result of delayed germline development

relative to somatic development (Hubbard and Schedl, 2019). Taken together, within the

limits of these two assays we infer that the soma and germline of worms fed *ΔfabH*

developed comparably with worms fed WT *E. coli, ΔfabF* and *ΔeutD* displayed delayed

germline development without a somatic delay, and *ΔeutN* delayed somatic

development as well as germline development relative to the (already delayed) somatic

development.

Because our screen assessed the proportion of gravid *C. elegans* at a single

time point, we asked whether non-gravid worms were permanently sterile or were

delayed in achieving fertility. Exact time points corresponding to larval development

differ when *C. elegans* is grown in liquid versus on solid media. Our screening time

point ("Day 1" 62-65 hours post seeding; see Methods) is an early adult timepoint when

worms fed on the WT bacteria are virtually all gravid (excluding a low background

sterility of GC1474). When we prolonged the assay by scoring the worms on Day 2—24

hours after the initial screening time point—the percentage of gravid worms significantly

increased, indicating that feeding on these *E. coli* mutants delayed *C. elegans* fertility

rather than induced permanent sterility (Fig. 1G). We also found no significant

differences when comparing the number of germline progenitor zone (PZ) nuclei in *C.*

elegans raised on each mutant or WT *E. coli* at the L4-to-adult molt (Fig. S1, H-I),

indicating that the delay in fertility is not likely due to slower accumulation of the PZ

during larval stages (Agarwal et al., 2018; Kocsisova et al., 2019; Korta et al., 2012).

DAF-2 insulin/IGF-like and DAF-7 TGFß signaling pathways promote germline

progenitor expansion and fertility in response to adequate nutrition, and this effect is

also observed in the *glp-1* mutant background used in our screen (Dalfo et al., 2012;

Michaelson et al., 2010; Pekar et al., 2017; Starich et al., 2020). Activation of these

pathways ultimately interferes with the activity of downstream transcriptional regulators,

daf-16 and *daf-5*, respectively; loss of *daf-16* or *daf-5* therefore mimics constitutive

activation of these pathways. We wished to determine whether loss of *daf-16* or *daf-5,*

neither of which alter overall growth rate, would circumvent the fertility delay caused by

each of the *E. coli* mutant diets. We fed WT or mutant *E. coli* to *daf-16* and *daf-5* mutant

C. elegans and determined the percentage of gravid worms. Loss of *daf-16* actually

exacerbated instead of rescuing the fertility defect of *C. elegans* raised on each mutant

E. coli, as well as on WT *E. coli* (Fig. S1J). Although constitutive activation of the DAF-7

TGFß pathway via loss of *daf-5* partially and variably suppressed the fertility defect of *C.*

elegans raised on *ΔeutN, ΔfabF* and *ΔfabH* (Fig. S1K), it also elevated the penetrance

of gravid *C. elegans* raised on WT *E. coli* to a similar extent, making the suppression

difficult to interpret. In short, we do not observe a simple causal relationship between

either the insulin or TGFß signaling pathways as assayed by loss of *daf-16* or *daf-5*, and

the alteration in the timing of *C. elegans* fertility caused by these four *E. coli* mutants.

E. coli mutants with altered metabolic byproducts delay _C. elegans_ fertility and

inhibit _T. muris_ egg hatching

Having identified *E. coli* mutants that delay fertility of a free-living nematode, we

examined whether these bacteria impact a distantly related parasitic nematode. *T. muris*

egg hatching was shown to be the reproductive stage sensitive to the presence of

bacteria (Hayes et al., 2010; White et al., 2018). Thus, to begin investigating possible

effects of the mutants identified in our *C. elegans* screen on the *Trichuris* lifecycle, we

adapted a previously described method to quantify *E. coli*-mediated hatching kinetics of

embryonated *T. muris* eggs *in vitro* (Hayes et al., 2010) (Fig. 2A). All *E. coli* mutants

were confirmed to have similar growth kinetics to WT *E. coli* (Fig. S2A). Two of the

mutants tested, *ΔeutN* and *ΔfabH,* elicited significantly reduced hatching compared to

rates mediated by WT *E. coli* (Fig. 2B). Complementation of *ΔeutN* or *ΔfabH* with a

plasmid-borne copy of the corresponding gene rescued hatching (Fig. S2, B-C).

To determine whether reduced egg hatching is due to the production of a toxic

factor, we performed the hatching assay utilizing WT *E. coli* mixed with each mutant

(Fig. 2C). As little as 10% *ΔeutN* in the mixture significantly inhibited the ability of WT *E.*

coli to mediate hatching, whereas the presence of *ΔfabH* had no effect. Thus, the

inhibitory effect of *ΔeutN* is dominant over WT *E. coli*, implicating the production of a

toxic factor by this mutant, while the reduced hatching rate of embryonated eggs

incubated with *ΔfabH* is likely due to deficiency of a pro-hatching factor.

Hybrid metabolomics revealed an unexpectedly low relative level of arginine and

ornithine in *ΔfabH* cell lysates compared to WT *E. coli* (Fig. 2D). Although interactions

between the fatty acid and arginine biosynthesis pathways of *E. coli* have not been

previously described, *fabH* deficiency would cause a buildup of its substrate malonyl-

ACP. A possible consequence of this buildup is compensatory decreased cycling of

acetyl-CoA, particularly through the TCA cycle, and consequential slowing of the urea

cycle and arginine biosynthesis. Supplementation of *ΔfabH* cultures with arginine or

ornithine successfully rescued *T. muris* egg hatching to levels similar to WT *E. coli* (Fig.

2, E-F). Similarly, supplementation of *ΔfabH* cultures with arginine or ornithine

significantly increased *C. elegans* fertility (Fig. S2, D-E). Consistent with a role for the

arginine biosynthesis pathway, we found that *Trichuris* egg hatching was diminished in

the presence of *ΔargE* or *ΔargH E. coli*, mutants defective in ornithine and arginine

synthesis, respectively (Fig. 2, G-H).

Although optimal *T. muris* egg hatching *in vitro* requires physical contact with *E.*

coli (Hayes et al., 2010; Koyama, 2016), a reduced amount of hatching can still occur in

a transwell system that separates bacteria and eggs into different chambers (Koyama,

2016), suggesting that both contact-dependent and independent factors contribute to

the process. Similar to previous reports, eggs hatched at approximately 50% their

maximum rate when incubated separately from WT *E. coli* in a transwell system (Fig. 2,

I-J). Hatching was further reduced when *ΔfabH* was used as the bacterium.

Supplementation of ornithine restored hatching in the presence of *ΔfabH* to levels

similar to those elicited by WT *E. coli* in the transwell system. Altogether, these findings

indicate that nematode pathways regulated by ornithine and arginine contribute to

contact-independent *E. coli*-mediated hatching of *T. muris* embryonated eggs.

Toxic intermediates of the *E. coli* ethanolamine utilization pathway inhibit *T.*

***muris* egg hatching**

To understand why *T. muris* eggs incubated with *ΔeutN E. coli* displayed the

greatest reduction in hatching, we first determined whether disrupting other parts of the

ethanolamine utilization pathway (Fig. 3A) led to similar outcomes. *E. coli* mutants

lacking enzymes in the pathway—*ΔeutB*, *ΔeutC*, *ΔeutD*, *ΔeutE*, and *ΔeutG*—facilitated

egg hatching at rates similar to WT *E. coli* (Fig. 3B). In contrast to these *eut* operon

genes, *eutN* encodes a subunit of a proteinaceous shell that enables the efficient

colocalization of the *eut* enzymes with their cofactors and substrates within a

microcompartment and prevents diffusion of pathway intermediates (Yeates et al.,

2010). It is possible that disruption of the microcompartment architecture allows the

release of a soluble toxic intermediate. Therefore, we tested whether transferring the

supernatant from *ΔeutN* culture was sufficient to inhibit the ability of WT *E. coli* to

mediate hatching (Fig. 3C). Although transferring culture supernatant from *ΔeutN*

grown in isolation had no effect, supernatant isolated from *ΔeutN* that had been

incubated with eggs (*ΔeutN^{egg}*) reduced hatching mediated by WT *E. coli* (Fig. 3D). This

suggests that *ΔeutN* produces a soluble toxic factor in the presence of *T. muris*

embryonated eggs.

Acetaldehyde is a known intermediate of the ethanolamine utilization pathway

that can readily form irreversible adducts with proteins, changing their conformation and

function (Donohue et al., 1983; Sorrell and Tuma, 1987) . Additionally, deletion of *eutN*

in *Salmonella* resulted in the greatest increase in acetaldehyde volatility among all *eut*

operon mutants during the first hours of growth (Penrod and Roth, 2006). Using a

fluorogenic hydrazone transfer (DarkZone) system to label aldehydes (Yuen et al.,

2016), we detected elevated levels of aldehydes in the culture supernatant of *ΔeutN^{egg}*,

but not in the culture supernatant of the complemented *ΔeutN* strain or when the mutant

was grown in the absence of eggs (Fig. 3E and S3A). Aldehyde levels were also

elevated in *ΔeutD* (Fig. 3E)*, the other *E. coli* mutant of this pathway identified by the *C.*

elegans screen, although the increase following egg incubation was not as striking as

for *ΔeutN*, potentially explaining why deletion of *eutD* did not inhibit hatching (Fig. 2B).

The eut operon transcriptional regulator in *E. coli, eutR,* is sensitive to the

chemical environment (Penrod and Roth, 2006). We hypothesized that *eutR* expression

may be altered in the presence of *T. muris* eggs, thus explaining why aldehyde levels

are increased when bacteria are co-incubated with eggs. We found that *eutR*

expression in WT *E. coli* and *ΔeutN* significantly increased following egg incubation

(Fig. S3B). As a control, increased *eutR* expression was detected from WT *E. coli*

grown in minimal medium supplemented with ethanolamine, consistent with previous

studies (Penrod and Roth, 2006; Rowley et al., 2020).

Next, we deleted the rate limiting enzyme of the ethanolamine utilization

pathway, *eutC*, in the *ΔeutN* background (*ΔeutNΔeutC)* to prevent the formation of toxic

intermediates (Penrod and Roth, 2006; Roof and Roth, 1989). We found that aldehyde

levels of *ΔeutNΔeutC^{egg}* were similar to that of WT *E. coli* and that *T. muris* egg

hatching was completely restored when the assay was performed with this double

mutant (Fig. 3E-F). We also found that pre-treatment of *ΔeutN^{egg}* supernatant with an

aldehyde scavenger N-acetyl cysteine completely abrogated its effects on WT *E. coli*

and restored hatching (S3C-D). Additionally, we found that supplementation of

acetaldehyde alone was sufficient to inhibit hatching elicited by WT *E. coli* in a

concentration-dependent manner and confirmed that bacteria viability was unaffected

(Fig. S3, E-F).

251 ***T. muris* reared in mice colonized with *E. coli* mutants exhibit defects in worm**

252 **morphology and fertility.**

253   *In vitro T. muris* egg hatching serves as a sensitive assay to investigate

254 nematode-bacteria interactions, but the role of intestinal bacteria in subsequent steps of

255 *Trichuris* development remains obscure because they are difficult to recreate outside

256 the mammalian host. Given that the *C. elegans* screen was focused on development of

257 the reproductive tract, we established a bacterial monocolonization mouse model to

258 investigate the role of bacterial *fabH*, *eutN*, and *eutD* in *T. muris* development and

259 reproduction (Fig. 4A). After we verified that each *E. coli* mutant colonized germ-free

260 mice to levels similar to WT *E. coli* (Fig. S4A), monocolonized mice were gavaged twice

261 with 100 eggs and sacrificed two weeks after the second gavage. *E. coli* colonization

262 and cecum length were confirmed to be similar across groups on the day of sacrifice (42

263 dpi) (Fig. S4, A-B).

264   We were able to recover adult worms from the cecum of mice monocolonized

265 with WT *E. coli* but not germ-free mice (Fig. S4C), consistent with previous observations

266 (White et al., 2018), indicating that our assay allows investigation of mechanisms by

267 which *E. coli* contributes to *T. muris* development in an intact host. Our experimental

268 setup in which mice receive high doses of embryonated eggs was designed to

269 overcome the potential reduced hatching efficiency that may occur in mice lacking a

270 complex microbiota. Therefore, as expected, we did not detect any differences in worm

271 burden when comparing mice colonized with WT *E. coli* versus *ΔfabH*, *ΔeutN*, and

272 *ΔeutD* strains (Fig. S4C). However, many of the *T. muris* recovered from mice colonized

273 with these mutants displayed a striking deformity characterized by shortened length

(Fig. 4, B-C; indicated by arrowheads). Although worms from WT *E. coli*-colonized mice

were exclusively long in length, mice colonized with *ΔfabH*, *ΔeutN*, and *ΔeutD E. coli*

produced a mixture of both long and short length worms, defined as < 1cm. The defect

was most severe in mice monocolonized with *ΔeutN* from which most worms recovered

were short. To investigate if the shortened length of *T. muris* resulted from mutant *E.*

coli colonization affecting the host, we examined T cell subsets that have been

previously established as key determinants of host susceptibility (Bancroft et al., 1994;

D'Elia et al., 2009). Flow cytometric comparison of mesenteric lymph node T cells from

mice colonized with WT or *ΔeutN E. coli* revealed no significant differences in T helper

type 1 (Th1), T helper type 2 (Th2), regulatory T (T reg) cells, or IFN-γ producing T cells

(Fig. S4D-G).

The average number of freshly laid eggs per worm was not significantly different

between the conditions or when comparing long versus short worms (Figure S4H).

However, in stark contrast to worms from WT *E. coli*-colonized mice, which laid eggs of

the typical ovoid structure, worms harvested from mice colonized with *ΔfabH*, *ΔeutD*,

and *ΔeutN E. coli* laid more eggs that displayed an aberrant rounded morphology, and

in some cases lacked opercula and a distinct shell, and the inner contents were

disorganized (Fig. 4, D-E). These structurally aberrant eggs took longer to reach

embryonated morphology, and even after a prolonged incubation period, significantly

fewer eggs appeared embryonated in the *ΔeutN* and *ΔeutD* groups (Fig. 4F). Eggs laid

by worms from all three mutant-colonized groups of mice had significantly lower

hatching rates compared to eggs laid by worms from WT *E. coli*-colonized mice when

tested for bacteria-mediated hatching *in vitro* (Fig. 4G). These hatching defects

persisted even when eggs were allowed to embryonate twice the embryonation time

published in the literature for *T. muris* (Fig. S4I), suggesting disrupted and not delayed

development. In line with our *in vitro* data (Fig. 3F), monocolonization of mice with

ΔeutNΔeutC abrogated the length and egg fitness defects of *T. muris* observed in

ΔeutN-colonized mice (Fig. 4, H-L). A similar rescue was seen when mice were

monocolonized with the *ΔeutN* complemented strain (Fig S4, J-L). These findings

indicating that bacterial deficiencies decrease *T. muris* worm size and egg fitness is

consistent with recent evidence showing that another parasitic nematode

Heligmosomoides polygyrus, lays fewer eggs and exhibits similar length defects when

reared in a germ-free host (Rausch et al., 2018).

To gain insight into the cause of morphologic and reproductive aberrancies in *T.*

muris, we performed RNA-Seq analysis of individual female worms (Fig S4M, Table

S2). The transcriptomes of worms harvested from WT *E. coli*-colonized mice clustered

separately from those of short and long worms harvested from mutant-colonized mice.

The latter did not segregate further according to length, representing no detectable

significant differences between short and long worms particularly in developmental

genes. Analysis of the top 50 most differentially detected genes between the WT- and

mutant-associated worms (Table S3) included genes related to *C. elegans* genes *egg-3*,

egg-4, *egg-5*, *alg-3*, and *alg-4* all of which are expressed in the germ line of *C. elegans*

and whose biological functions include roles in oogenesis, spermatogenesis,

fertilization, the egg-to-oocyte transition, and egg shell deposition (Boeck et al., 2016;

Cheng et al., 2009; Conine et al., 2010; Maruyama et al., 2007; Parry et al., 2009).

Crosschecking our 50 genes with published lifestage- and sex-specific transcriptomics

of *T. muris* (Foth et al., 2014), all genes found to be highly expressed in the female

worms from mutant-colonized mice were previously detected as upregulated in adults

(vs. larvae), providing additional evidence that the short worms are not disrupted in

somatic development. Additionally, these same genes were also previously detected as

more highly expressed in *T. muris* males compared to females (Foth et al., 2014).

Conversely, all but one of the genes that were downregulated in the worms from

mutant-colonized mice were found to be upregulated in *T. muris* females. These

discordances provide evidence of sexual development dysregulation in worms

harvested from mutant-colonized mice.

In conclusion, these results provide evidence for a previously undescribed role

for byproducts of microbial metabolism on nematode development, a role that appears

evolutionarily conserved. Despite lacking many genes specific to parasitism, our

findings support the utility of *C. elegans* as a model for bacteria-nematode interactions

because of its high sensitivity to its bacterial milieu. In addition to identifying two critical

pathways in bacteria, fatty acid biosynthesis and ethanolamine utilization, our findings

describe a novel pipeline that leverages genetic tools available in *C. elegans* to discover

new helminth biology. An important future direction would be to determine the role of

microbial byproducts in the setting of a complex microbiota. For instance, microbiota

rich in bacteria with highly active arginine biosynthesis or another pathway favorable for

T. muris reproduction may confer greater susceptibility to infection. Identifying the

microbes and their products that affect different stages of the helminth life cycle would

allow predictions and testable hypotheses regarding the impact of interindividual

differences in microbiota composition. We propose that *C. elegans* can be used as a

system for investigating the effects of specific microbial genes and pathways in

supporting the parasitic nematode life cycle and aid in further understanding the

transkingdom interactions that sustain human disease.

Acknowledgements

We wish to thank D. Jones, R. Rose, and L. Ash from the NYULH Metabolomics

Laboratory (SCR_017935) and C. Yun, S. Cook, and D. Kahler from the High

Throughput Biology Core (now Microscopy Laboratory, NIH S10 RR023704) for their

help in acquiring and analyzing the data presented. Both Core Laboratories are funded

partially by NIH/NCI P30CA016087. Some strains were built using strains provided by

the CGC which is funded by the NIH Office Research Infrastructure Programs (P40

OD010440), and we thank WormBase. We also thank H. Darwin and M. Pacold for

advice on microbial metabolism; S. Marion, C. Linton, M. Xu, A. Niranjan, D. Stokes, M.

Lopez-Redondo, I. Irnov, V. Torres, S. Dyzenhaus, M. Hui, J. Richards (NYU), and E.

Tait-Wojno (Univ. of Washington) for technical assistance and reagents; members of

the Hubbard Lab, Cadwell Lab, and NYU Medical Scientist Training Program for

constructive comments and technical assistance; and M. Alva, J. Carrasquillo, and D.

Basnight for assistance with gnotobiotics.

Funding

NIH grant DK093668 (K.C.)

NIH grant AI121244 (K.C.)

NIH grant HL123340 (K.C.)

NIH grant AI130945 (K.C.)

NIH grant AI140754 (K.C.)

NIH grant DK124336 (K.C.)

NIH grant R01GM130152 (E.J.A.H.)

NIH grant R35GM134876 (E.J.A.H.)

NIH grant R01GM035769 (J.G.B.)

NIH grant 5T32AI100853 (M.V.)

NIH grant T32GM136573 (M.V.)

NIH grant T32GM007308 (M.V.)

Faculty Scholar grant from the Howard Hughes Medical Institute (K.C.)

Crohn's & Colitis Foundation (K.C.)

Kenneth Rainin Foundation (K.C.)

Judith & Stewart Colton Center of Autoimmunity (K.C.)

US National Cancer Institute (CA217809) (E.T.K.)

K.C. is a Burroughs Wellcome Fund Investigator in the Pathogenesis of Infectious

Diseases.

Author contributions

M.V., R.D., E.J.A.H, and K.C. conceived the study and designed the experiments. M.V.,

R.D., and J.B. performed, analyzed, and interpreted all the experiments for *T. muris* and

C. elegans. D.J.L. performed the bacterial cloning. H.S.P and E.T.K provided DarkZone

and technical advice. J.C.D. provided technical assistance and advice for RNA

sequencing and analysis. K.C., E.J.A.H, and J.G.B oversaw analysis and interpretation

of all experiments described. M.V., R.D., E.J.A.H., and K.C. wrote the manuscript with

inputs from all authors.

Competing interests

K.C. has received research support from Pfizer, Takeda, Pacific Biosciences,

Genentech, and Abbvie. K.C. has consulted for or received honoraria from Puretech

Health, Genentech, and Abbvie. K.C. holds U.S. patent 10,722,600 and provisional

patent 62/935,035 and 63/157,225, and E.J.A.H. holds US patent 6,087,153.

Data and materials availability

All data needed to evaluate the conclusions in this study are available in the main text

and the supplementary materials or available from the corresponding authors upon

request.

Supplementary Materials

Methods

Figs. S1 to S4

Tables S1-S7

Fig. 1 Identification of *E. coli* mutants that delay *C. elegans* fertility. (A) Concept of

high throughput screening of *C. elegans* raised on *E. coli* mutants from the Keio library.

(B) Example of reduced penetrance of *C. elegans* bearing embryos at a single time point; embryos marked with GFP and pharynx with mCherry. **(C)** Detailed flowchart for schematic in (A). **(D)** Percentage of *C. elegans* bearing embryos in replicates of four *E. coli* mutants of interest from the screen. Fisher's exact test for each mutant versus controls from the same plates; total n number of worms scored are (left to right): 913, 411, 1253, 583, 1643, 550, 1561, 232. **(E)** Somatic development is delayed in *C. elegans* raised on *ΔeutN*. Percent of worms at indicated stage at 56 hours after L1 seeding to wells containing the control or mutant bacteria. $n \geq 38$ worms were scored for each *E. coli* mutant. *ΔeutN* is the only mutant that is statistically significantly different from the control GC1547 ($p \leq 0.001$). n number of worms scored (left to right): 52, 59, 61, 40, 54. **(F)** *ΔeutN* and *ΔfabF* delay germline development relative to somatic development. Percent of gonad arms displaying characteristic Pro phenotype. $p \leq 0.001$ for *ΔeutN* and *ΔfabF*. Pooled from two independent trials; n number of worms scored are (left to right): 87, 103, 79, 41, 61. **(G)** Fertility is delayed. Percent of worms with embryos on Day 1 and Day 2. $p \geq 0.05$ (not significant) for any mutant versus the control on Day 2. n number of worms scored (left to right): 547, 297, 242, 242, 60, 46, 124, 121, 253, 261. (D-G) Circles represent mean of independent biological replicates and bars show mean and SEM; pairwise Fisher's exact test versus controls (D-F) or Day 1 versus Day 2 (G). ** $p < 0.01$, **** $p < 0.0001$.

Fig. 2 *E. coli* **mutants with altered metabolic byproducts delay** *C. elegans* **fertility**

and inhibit *T. muris* **egg hatching. (A)** Experimental approach for *in vitro* hatching

assay. **(B)** Percent *in vitro* hatched *T. muris* eggs after incubation with overnight

cultures of four *E. coli* mutants identified from the *C. elegans* screen or WT *E. coli*.

Hatched eggs were checked using light microscopy at timepoints indicated (n = 4); h =

hours. **(C)** Hatching rates for mixtures of WT *E. coli* and WT *E. coli* or either Δ*fabH* or

Δ*eutN* overnight cultures in the ratios indicated. Total concentration of each mixture was

maintained at 1x 10^8 CFU/mL (n = 3). **(D)** Hierarchical clustering of metabolites that

were significantly ($p < 0.05$, Student's t-test) downregulated (blue) or upregulated (red)

in Δ*fabH* relative to WT. **(E, F)** Percent *in vitro* hatched *T. muris* eggs elicited by Δ*fabH*

and cultures supplemented with (E) 500µM arginine or (F) 100µM ornithine where

indicated (n=3). **(G)** Arginine biosynthesis pathway in *E. coli*. **(H)** Hatching rates elicited

by Δ*argE* and Δ*argH* (n = 4). **(I)** Experimental approach for hatching assay in a transwell

system. **(J)** Hatching quantified for eggs contained within a transwell chamber

separated from bacterial culture in the outer chamber by a membrane with 0.4µm pores.

Samples supplemented with ornithine as in (H) are indicated (n = 3). (B) One-way

analysis of variance (ANOVA) of area under curve compared to WT. (C, E-F, H, J) One-

way analysis of variance (ANOVA) with Dunnett's post-test compared to WT. For all

panels, measurements were taken from distinct samples. Graphs show means and

SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. ns = not significant.

Fig. 3 Toxic intermediates of the *E. coli* ethanolamine utilization pathway inhibit *T. muris* egg hatching. (A) Ethanolamine utilization pathway in *E. coli*. Modified from (Kaval and Garsin, 2018). **(B)** Hatching rates for *eut* mutants in the Keio library and WT controls (n = 3-6). **(C)** Experimental approach to assess the toxicity of soluble factors produced by *E. coli* Δ*eutN*. Filtered supernatant from Δ*eutN E. coli* or controls grown with or without *T. muris* eggs for four hours were transferred to a dish containing WT *E. coli* and *T. muris* eggs. The ability of WT *E. coli* to mediate egg hatching in the presence

of supernatant was examined four hours later. **(D)** Quantification of hatching rates for

mixtures containing WT *E. coli* cells and the indicated filtered supernatants (n = 3). **(E)**

Aldehyde concentrations in supernatants of indicated *E. coli* strains incubated with or

without *T. muris* eggs; Δ*eutN* comp. = Δ*eutN* strain where *eutN* has been complemented

on a plasmid. **(F)** Hatching rates for Δ*eutN*Δ*eutC* double mutant (n = 4). (B, D-F)

Measurements were taken from distinct samples. The average per independent

experiment for at least three technical replicates shown. Graphs show means and SEM.

One-way analysis of variance (ANOVA) with Dunnett's post-test compared to WT. * $p <$

0.05, *** $p < 0.001$, **** $p < 0.0001$. ns = not significant.

Fig. 4 *T. muris* reared in mice colonized with *E. coli* mutants exhibit defects in

worm morphology and fertility. (A) Experimental approach for *T. muris* infection of

mice monocolonized with WT or mutant *E. coli*. **(B)** Proportion of short and long worms

harvested per mouse colonized by indicated *E. coli* strains; N.D. = not detected (n = 6-9

mice per group). **(C)** Representative images of worms harvested from mice

monocolonized with WT or *ΔeutN E. coli*. A representative short worm (indicated by

arrowhead) next to a long worm is shown in the inset for *ΔeutN E. coli*. **(D)**

Representative images of eggs laid by worms from (B). Worms were incubated in RPMI

medium overnight to trigger egg laying. **(E)** Proportion of total eggs laid per worm with

normal morphology. Circles represent individual worms (WT, n = 46; *ΔfabH*, n = 68;

ΔeutD, n = 58; *ΔeutN*, n = 50). **(F)** Percentage of embryonated eggs following 8-week

incubation. **(G)** Percentage of eggs in (F) that hatched after incubation with WT *E. coli*

for 4 hours at 37°C. **(H)** Representative photos of worms harvested from WT and

ΔeutNΔeutC monocolonized C57BL/6 mice. **(I)** Representative images of eggs laid by

worms harvested from monocolonized mice groups indicated**. (J)** Proportion of total

eggs laid per worm with normal morphology**. (K)** Percentage of embryonated eggs

following 8-week incubation. **(L)** Percentage of eggs in (K) that hatched after incubation

with WT *E. coli* for 4 hours at 37°C. (B, E, F-G) Graphs show (B) means and SEM or (E-

G) medians. One-way analysis of variance (ANOVA) with Dunnett's post-test compared

to WT. (J-L) Graphs show means and SEM. Two-tailed, unpaired t-test. For all panels,

measurements were taken from distinct samples. Data shown are pooled from two

independent experiments. ** p < 0.01, **** p < 0.0001. ns = not significant.

Methods

Keio Library growth and culture

The frozen Keio collection was stamped out onto 150mm LB kanamycin (30µg/ml) plates and grown overnight at 37°C. For *C. elegans* experiments, individual colonies were inoculated into 700 µl of LB-Kan (30µg/ml) in 96 deep well plates using a compatible pin replicator. The plates were sealed with pure link air porous tape (Invitrogen cat no. 12262010) and incubated for 16-18 hours at 37°C, shaking at 250 rpm. Cultures were diluted 10X in S-media in 96 well format and OD at 600 nm was recorded for future reference to ensure that candidate positive mutants grew to an OD similar to the GC1547 wild-type control wells in the same plate. (This was done since poor bacterial growth can reduce the percentage of gravid *C. elegans*; all 315 candidates from the primary screen passed this control). Bacterial cultures were pelleted by centrifugation at 2000 rpm for 30 minutes. Supernatant was discarded and the pellet was resuspended in 100µl of S-media. 40µl of the culture was transferred into two separate plates (two technical replicates; black-walled, clear bottom 96-well microplates; Corning Cat. No. 3904) using a multi-channel pipette. The GC1547 control wild-type strain was a kanamycin resistant transformant of the Keio library starting strain BW25113, carrying the pET28a plasmid. For *T. muris in vitro* hatching assays, *E. coli* was grown overnight for 12-14 hours in Luria-Bertani broth with shaking at 225 rpm at 37 °C.

Primary screen

C. elegans GC1474 (Table S1; derived by outcrossing GC1373 (Roy et al., 2018) of the genotype *glp-1(e2141); end-1::gfp; myo-2p::mCherry*) was maintained on OP50 *E. coli* on NGM plates at 20ºC and synchronized in the L1 larval stage using hypochlorite treatment followed by overnight hatching in egg buffer (Shaham, 2006). 40-50 L1 larvae were seeded into 96-well plates in technical duplicate with 40µl of *E. coli* mutant cultures and incubated at 20°C with mild shaking in a humidified chamber for 62-65

hours. At this timepoint, control worms were fully gravid adults; control GC1547 bacteria were included on each plate in wells void of library bacteria. Prior to imaging, 40µl of 2mM levamisole was added to each well to immobilize the worms. The plates were then sealed with aluminum foil (Corning Cat. No. 6570) and imaged to detect both green (embryos) and red (pharynx) fluorescent markers using arrayscan (Thermo Scientific VTI). Images were captured at 2.5X magnification 2x2 binning in a 16 mm^2 field and digitally archived. Images captured on the red and green channels were merged in ImageJ using macros written to automate the process. Each adult animal was scored visually as "with embryos" (green embryos visible in the uterus) or "without embryos" (no green embryos visible in the uterus), and the number of worms without embryos and the total number of worms (excluding any that were arrested or any that crossed the edge of the well). Wells in which many animals had not reached adulthood were marked and were not considered further in our analysis. Ten such bacterial mutants grew well by OD but did not support worm development.

Additional culling of potential hits was performed. A variety of technical factors eliminated fewer than 10% of the total: 54 library wells were devoid of bacteria as per the plate map; 58 had a gene name in the library but the bacteria did not grow (either on solid or liquid culture); 41 were missed due to human error while scoring (noted as largely fertile but not counted individually); 5 had low n; 4 were not included due to reasons classified as 'other' (e.g., poor image) and 10 wells had poor worm growth in both replicates.

Worm counts (with/without embryos) were exported into ActivityBase (IDBS, https://www.idbs.com; a Scientific informatics software platform) and the robust Z score (a statistical measure that takes into account the median plate "without embryos" percent) was calculated for every well to minimize problems associated with plate to plate variation. Each well was analyzed with respect to percent worms without embryos, consistency (across technical replicates) and robust Z score. A cut-off of robust Z ≥ 1.5 was selected as "Z positive" (≤ 10% of the wells per plate scored above this threshold).

We also calculated the penetrance of worms without embryos for pooled technical replicates.

Criteria for selection in the primary screen weighed the robust Z and penetrance, with special consideration for cases where only one replicate was available. We included those wells for which only one technical replicate was available provided that well made the RobZ cut off (≥ 1.5) and/or percent non-gravid worms as further described below. Reasons for single-replicate included low n (62 wells), human error (49 wells), worm arrest (31 wells), poor image quality (22 wells) and 1 well marked as 'other'. Considering Z score, penetrance and replicates, the candidates were further classified as follows: High, A: both replicates Z positive and average penetrance of non-gravid worms > 20%, or B: one replicate available and it is Z positive and > 20%, or C: both replicates available but only one is Z positive and non-gravid ≥ 30%. Medium, A: both replicates Z positive and pooled non-gravid percent 13-20%, or B: only one replicate available which is Z positive and 13-20% penetrance of worms without embryos, or C: both replicates available but only one is Z positive with pooled non-gravid penetrance between 20-30%. Low: both replicates available, but only one is Z positive and non-gravid percent between 12.5-20%. This analysis yielded 315 primary screen candidates.

We then plotted the percent non-gravid in descending order for every well per plate and noted the inflection point (see (Roy et al., 2018)). We placed a subset of the mutants that passed the criteria above onto these plots and observed a strong correlation: wells with larger Z scores all appeared above the inflection point.

Secondary screen: reproducibility
The 315 candidates from the primary screen were re-selected from the original Keio library and gridded onto secondary screen plates. Two plates included only high priority candidates or a combination of randomly selected high, medium and low priority candidates to help assess whether there was a correlation between penetrance and reproducibility. No such correlation emerged, but these plates were considered as additional replicates in the analysis going forward (hence 6 plates) though 315 mutants.

Control bacteria were included in all secondary screen plates. Three biological replicates were performed in technical triplicate and scored and analyzed as described above. Technical replicate non-gravid/gravid counts were pooled and analyzed in pairwise comparison with the control on the same plates using Fisher's exact test. Candidates with p-values ≤ 0.05 in at least 2 out of 3 biological replicates were chosen for further analysis. This analysis yielded 10 candidates.

PCR validation of selected Keio mutants (completed for all ten hits)

We amplified and sequenced PCR fragments ~150 bp upstream and downstream of the predicted gene from each mutant and the control. Nested primers were used to sequence the PCR fragment and confirm the presence of the FRT and kanamycin cassette in the mutant DNA (Tables S3, S5).

P1 transduction

Mutant alleles comprising a kanamycin-resistance marker within a gene deletion were introduced into BW25113 by P1 transduction (Thomason et al., 2007) from the Keio collection (Baba et al., 2006) .The kanamycin-resistance cassette was subsequently excised to generate a markerless in-frame gene deletion by transformation with the temperature-sensitive plasmid pCP20 (Cherepanov and Wackernagel, 1995), which encodes FLP recombinase, and growth at 30°C. Finally, pCP20 was eliminated from the cells by growth at 42°C.

Complementation

For *fabH* and *eutN*, the bacterial gene coding regions plus 20 bp upstream were inserted into pBR322 downstream of a plasmid promoter by Gibson Assembly. The wild-type genomic DNA was amplified and assembled with a PCR fragment obtained from the pBR322 vector (specific primers are listed in Table S5; the fragment excluded part of the Amp^R gene between *Eco*RI and *Pst*I restriction site). PCR products were

purified using gel extraction kit (QIAEX II cat no. 20021) and 100 ng of vector was used in 1:1 molar reaction with the genomic fragment. The assembly reaction was then transformed into high efficiency DH5alpha cells (NEB C2987I). For *eutD*, the protein-coding region was amplified by PCR and cloned between the NdeI and XbaI sites downstream of the *lac*UV5 promoter of plasmid pPlacRppH6 so as to replace the *rppH* coding region (Deana et al., 2008). Plasmid preparations (Qiagen mini prep kit Cat no.27106) were sequenced using plasmid-specific primers (Table S5). Plasmids were confirmed by DNA sequence and transformed into the respective mutant (so as to create strains GC1558, GC1559, and GC1650 for *eutN* rescue, *fabH* rescue, and *eutD* rescue, respectively) and tested for reduced penetrance of the *C. elegans* non-gravid phenotype upon feeding.

Developmental timing and germline analysis

Somatic development assay
C. elegans were maintained on OP50, synchronized by hypochlorite treatment and allowed to hatch at 20ºC with shaking. 30-40 L1 larvae were seeded in 96 deep well plates with each well carrying a specific mutant clone. Worms were incubated at 20ºC with shaking for 48 hours and monitored at 400X on a Zeiss Z1 Axio Imager for vulval morphology (Mok et al., 2015) at specific time points.

Pro phenotype assay
glp-1(ar202) animals were scored for the Pro phenotype upon feeding with mutant bacteria using the same strategy as for the primary screen. Worms were maintained at 15ºC on OP50 synchronized and grown in 96 deep well format as above. Day 1 (~72 hours post seeding) worms were fixed and DAPI stained and each gonad arm was scored for the Pro phenotype (Pepper et al., 2003) at 400X.

Fertility at Day 2

GC1474 worms were maintained at 20ºC on OP50, synchronized as L1 larvae and grown in 96 deep well format as above. Worms were anesthetized and imaged as in the screen on Day 1 and Day 2 to determine changes in penetrance of fertile worms.

Progenitor zone nuclei counts at L4-Adult molt
Worms grown on GC1547 control, *eutN*, *eutD,* and *fabH* mutant bacteria were monitored for vulval development and isolated at the L4-adult molt. Worms were then stained with DAPI (Michaelson et al., 2010) and 0.5µm Z-stack images were collected and analyzed using ImageJ to determine the number of germ cells in the PZ. In approximately 30% of the worms, the transition zone was not clear. These were censored from the analysis.

Assay for dependence on DAF-7/TGFß and DAF-2 IIS pathways
C. elegans strains GC1474, GC1545 and GC1250 (Table S2) were hypochlorite-synchronized and hatched overnight at 20ºC. L1 larvae were seeded into 96 deep well plates containing specific bacterial mutants and scored for the presence of embryos in adults on Day 1 on Zeiss compound microscope at 100X.

Parasite maintenance

Stock eggs of *T. muris* E strain (Ramanan et al., 2016) were maintained in the NOD.Cg-*Prkdc^{scid}*/J (Jax) mouse strain in a specific pathogen free (SPF) facility and propagated as previously described (Antignano et al., 2011). Each egg batch was confirmed to hatch at ≥ 80% *in vitro* using method below and WT *E. coli* before use in subsequent experiments.

In vitro hatching of T. muris eggs

T. muris eggs were hatched *in vitro* by mixing 25µL of embryonated eggs at a concentration of 1 egg/1µL suspended in sterile water with 10µL of *E. coli* overnight culture and 15µL sterile LB in individual wells of a 48 well plate. Plates were incubated

at 37°C and checked every hour for a total of four hours on the Zeiss Primovert microscope to enumerate hatched eggs. Rates describe hatching after four hours of incubation unless otherwise indicated. Experiments utilizing transwell inserts (Millicell) were performed as previously described (Koyama, 2016). For experiments where cell-free supernatant was used, supernatant and cells were isolated by centrifugation and filtration through a 0.22μm syringe filter or after wash with autoclaved ddH20, respectively. Incubation with embryonated eggs was performed by mixing *T. muris* eggs at a concentration of 15 eggs/2 mL culture and keeping at 37°C for four hours.

Metabolomics

Sample preparation

Individual colonies (obtained after streaking frozen bacteria onto LB-kanamycin plates and incubating at 37ºC) were inoculated in 30 ml of LB-kanamycin (30μg/ml) liquid in a 50ml falcon tube. Cultures were grown for 16-18hr at 37ºC with 250 rpm shaking and then bacteria were pelleted by centrifugation at 1500 rpm for 20-30 minutes at 4ºC. The supernatant was transferred to a fresh tube, filtered through 0.2μm filter (Corning (431229) and frozen at -80ºC for future analysis. Pellets were resuspended in 7.5 ml of S-media (for ~250ml of S-media the recipe was 240ml of S-basal, 2.5ml of 1M potassium citrate- final concentration of 10mM, 2.5ml of 1M trace metals for final concentration of 10mM, 750μl of 1M MgS0$_4$ for final concentration of 3mM, 750μl of 1M CaCl$_2$ for final concentration of 3mM, 72μl of 100mg/ml of kanamycin for final concentration of 30μg/ml). This solution was filter sterilized using a 500ml Nalgene filter (291-3320 Fisher). Post filter sterilization, 250μl of 5mg/ml cholesterol – final concentration of 5μg/ml and 2.5ml of 250μg/ml fungizone for final concentration of 2.5μg/ml (BP264550 Fisher).

The number of CFU/ml was estimated for each culture by plating dilutions (10μl of culture was mixed with 900μl of S-media to get 10^2 or 1:100 dilution, which was then serially diluted to $1:10^5$ or $1:10^6$ on LB agar plates and counting colonies after growth at 37°C. The remaining culture was incubated at 20ºC with mild shaking overnight to

mimic *C. elegans* growth conditions. Based on CFU/ml measurements, 10^{10} cells were fast filtered (EZFIT Vacuum Manifold, Millipore EZFITLOW03 with Microfil V, Millipore MVHAWG124) and metabolite extracts were isolated as previously described (Irnov et al., 2017). Three independent biological samples were extracted on three independent days and subsequently analyzed by the NYU Langone Metabolomics Core Resource Library.

LC-MS/MS with the hybrid metabolomics method

Samples were subjected to an LCMS analysis to detect and quantify known peaks. A metabolite extraction was carried out on each sample based on a previously described method (Pacold et al., 2016). The LC column was a Millipore^{TM} ZIC-pHILIC (2.1 x150mm, 5µm) coupled to a Dionex Ultimate 3000^{TM} system and the column oven temperature was set to 25^{O}C for the gradient elution. A flow rate of 100µL/min was used with the following buffers: A) 10mM ammonium carbonate in water, pH 9.0, and B) neat acetonitrile. The gradient profile was as follows; 80-20%B (0-30 min), 20-80%B (30-31 min), 80-80%B (31-42 min). Injection volume was set to 2µL for all analyses (42 min total run time per injection).

MS analyses were carried out by coupling the LC system to a Thermo Q Exactive HF^{TM} mass spectrometer operating in heated electrospray ionization mode (HESI). Method duration was 30 min with a polarity switching data-dependent Top 5 method for both positive and negative modes. Spray voltage for both positive and negative modes was 3.5kV and capillary temperature was set to 320^{O}C with a sheath gas rate of 35, aux gas of 10, and max spray current of 100µA. The full MS scan for both polarities utilized 120,000 resolution with an AGC target of 3e6 and a maximum IT of 100ms, and the scan range was from 67-1000 *m/z*. Tandem MS spectra for both positive and negative mode used a resolution of 15,000, AGC target of 1e5, maximum IT of 50ms, isolation window of 0.4m/z, isolation offset of 0.1m/z, fixed first mass of 50m/z, and 3-way multiplexed normalized collision energies (nCE) of 10, 35, 80. The minimum AGC target was 1e4 with an intensity threshold of 2e5. All data were acquired in profile mode.

Hybrid Metabolomics Data Processing

Relative quantification of metabolites

The resulting Thermo^{TM} RAW files were converted to mzXML format using ReAdW.exe version 4.3.1 to enable peak detection and quantification. The centroided data were searched using an in-house python script Mighty_skeleton version 0.0.2 and peak heights were extracted from the mzXML files based on a previously established library of metabolite retention times and accurate masses adapted from the Whitehead Institute (Chen et al., 2016), and verified with authentic standards and/or high resolution MS/MS spectral manually curated against the NIST14MS/MS (Simon-Manso et al., 2013) and METLIN (2017)(Smith et al., 2005) tandem mass spectral libraries. Metabolite peaks were extracted based on the theoretical *m/z* of the expected ion type e.g., [M+H]$^+$, with a ±5 part-per-million (ppm) tolerance, and a ± 7.5 second peak apex retention time tolerance within an initial retention time search window of ± 0.5 min across the study samples. The resulting data matrix of metabolite intensities for all samples and blank controls was processed with an in-house statistical pipeline Metabolyze version 1.0 and final peak detection was calculated based on a signal to noise ratio (S/N) of 3X compared to blank controls, with a floor of 10,000 (arbitrary units). For samples where the peak intensity was lower than the blank threshold, metabolites were annotated as not detected, and the threshold value was imputed for any statistical comparisons to enable an estimate of the fold change as applicable. The resulting blank corrected data matrix was then used for all group-wise comparisons, and t-tests were performed with the Python SciPy (1.1.0) (Virtanen et al., 2020) library to test for differences and generate statistics for downstream analyses. Any metabolite with p-value < 0.05 was considered significantly regulated (up or down). Heatmaps were generated with hierarchical clustering performed on the imputed matrix values utilizing the R library pheatmap (1.0.12) (Kolde, 2015). Volcano plots were generated utilizing the R library, Manhattanly (0.2.0). In order to adjust for significant covariate effects (as applicable) in the experimental design the R package, DESeq2 (1.24.0) (Love et al., 2014) was used to test for significant differences. Data processing for this correction

required the blank corrected matrix to be imputed with zeroes for non-detected values instead of the blank threshold to avoid false positives. This corrected matrix was then analyzed utilizing DESeq2 to calculate the adjusted p-value in the covariate model.

Arginine and ornithine supplementation assay

fabH mutant *E. coli* or wild-type control was grown with 500µM or 100µM ornithine in LB-kanamycin overnight at 37ºC with 250 rpm shaking. For *C. elegans* experiments, worms were imaged on Day 1 as described for the primary screen. For *T. muris* experiments, hatching assay was performed as described above.

DarkZone labeling of aldehydes

Cell-free supernatants isolated from *E. coli* overnight cultures were incubated with 20µM DarkZone dye pre-dissolved in DMSO, 5mM 2,4 dimethoxyaniline catalyst (TCI America) pre-dissolved in DMSO, and buffer (100mM Tris pH 6.8, 150 mM NaCl) in a 96 well optical flat-bottom plate (Thermo-Fisher) in duplicate at 37°C for 1 hour. Adhesive plate seals were used to prevent evaporation of aldehydes. Fluorescence was measured using an EnVision 2 103 Multi-label Reader (PerkinElmer). Duplicate measurements were averaged.

Quantitative PCR

Bacterial cells grown *in vitro* were pelleted and washed with RNA Protect (Qiagen) and PBS. For all samples, RNA was was isolated using the RNeasy micro kit with DNase treatment (Qiagen). Synthesis of cDNA was conducted with the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). Samples were run without reverse transcriptase to control for genomic DNA contamination. Quantitative PCR was performed using SybrGreen (Roche) on a Roche480II Lightcycler using the following primers indicated in Table S5. Two technical replicates of each biological replicate were

included for each gene target. Relative expression of the respective genes to *16S* rRNA expression was calculated using the $\Delta\Delta C_T$ method.

N-acetyl cysteine supplementation assay

Bacterial cells and cell-free supernatants of WT *E. coli* and *eutN* mutant *E. coli* were isolated as described above. A 100mg/mL stock concentration of N-acetyl cysteine (Millipore Sigma) was dissolved in MilliQ water by stirring at 70°C. The N-acetyl cysteine was then added to supernatants at the concentrations of 8µM, 10 µM, 30 µM, and 50 µM and incubated at room temperature for one hour. WT *E. coli* cells were then resuspended in the supernatants and incubated with *T. muris* eggs to perform the hatching assay as described above.

Mice

Gnotobiotics

Previously described germ-free C57BL/6J mice (Martin et al., 2018) were maintained in flexible film isolators, and absence of faecal bacteria and fungi was confirmed by aerobic culture in brain heart infusion, sabaraud and nutrient broth (Sigma), and qPCR for bacterial 16S and eukaryotic 18S ribosomal RNA genes through sampling of stool from individual cages in each isolator on a monthly basis. Mice were transferred into individually ventilated Tecniplast ISOcages for infections to maintain sterility under positive air pressure. All animal studies were performed according to approved protocols and ethical guidelines established by the NYU Grossman School of Medicine Institutional Animal Care and Use Committee (IACUC) and Institutional Review Board.

Murine in vivo infections

Female mice were monocolonized at 6-8 weeks of age by oral gavage with 1×10^8 colony forming units per mL (CFU/mL) of indicated *E. coli* strains. Overnight *E. coli* cultures were diluted 1:100 in Luria-Bertani broth followed by 2-3 hours of growth until 1×10^8 CFU/mL was reached. Bacterial density was confirmed by dilution plating. Cultures were pelleted by centrifugation at 2437g for 10 minutes and washed once with

sterile 1x PBS. Pellets were then resuspended in sterile 1x PBS and mice were inoculated by oral gavage with $1x10^8$ CFU in a volume of 100µL. Inoculum was verified using dilution plating of aliquots.

7 and 28 days later, mice were infected by oral gavage with ~100 embryonated *T. muris* eggs. Individual worms were collected from cecal contents of all infected mice and washed in RPMI 1640 (Corning) supplemented with penicillin (100U/ml) and streptomycin (100µg/ml; Sigma). To evaluate egg laying, each worm was placed into individual wells of a 48 well plate with 200µL supplemented RPMI. Plates were then incubated at 37°C overnight in a closed tupperware (Sistema) lined with damp paper towels. The following day, eggs laid were enumerated using a Zeiss Primovert light microscope at 100X. Samples containing ~1000 eggs per condition were mounted on glass slides with glycerol mounting medium (Sigma-Aldrich) and analyzed using a Zeiss Axio Observer at 400X with oil emersion for color images or EVOS FL Auto (Life Technologies) at 200X for black and white images. Images were processed using ImageJ.

Flow Cytometry

Cells from mesenteric lymph nodes were isolated and stimulated as previously described (Ramanan et al., 2016). Stimulated cells were stained with anti-CD45 Pacific Blue, anti-CD3ε FITC, anti CD4 APC-Cy7, and anti-IFNγ AF700 from Biolegend and anti-CD19 PerCP Cy5.5 from eBioscience. Fixation and permeabilization buffers from Biolegend were used for intracellular cytokine staining, and a fixable live/dead stain from Biolegend was used to exclude dead cells. For nuclear staining, unstimulated cells were stained with anti-CD45 Pacific Blue, anti-CD3 FITC, anti-CD4 APC-Cy7, anti-FoxP3 PE-Cy7, and anti-Tbet APC from Biolegend, and anti-CD19 PerCP Cy5.5 and anti-GATA3 PE from eBioscience using the Foxp3 staining kit (eBioscience). Flow cytometric analysis was performed on a CytoFLEX analyzer (Beckman Coulter) and analyzed using FlowJo 10.0.8.

RNA isolation from harvested *T. muris* and RNA-seq

T. muris harvested from mice ceca were immediately transferred to RPMI 1640 (Invitrogen) supplemented with penicillin (100U/ml) and streptomycin (100µg/ml; Sigma) and stored at -80°C. RNA from individual worms was isolated using the RNeasy micro kit with DNase treatment (Qiagen). RNA-seq library preparation was done at the NYU School of Medicine Genome Technology Core using an automated Trio enrichment library preparation protocol (NuGEN). Libraries were then sequenced on a NovaSeq 6000 (Illumina). All sequences have been deposited at NCBI under the BioProject PRJNA803354.

RNA-seq analysis

Raw RNA-seq reads were trimmed using Trimmomatic v.036 (Bolger et al., 2014) and aligned to the reference *T. muris* genome (GenBank Assembly Accession: GCA_000612645.2) using the splice-aware STAR v2.7.3 (Dobin et al., 2013). The featureCounts command from Subread v1.6.3 (Liao et al., 2014) was used to generate counts for each gene based on how many aligned reads overlapped the exons of that gene. These counts were then normalized and used to test for differential expression using negative binomial generalized linear models implemented by DESeq2 v1.24.0 (Love et al., 2014).

Functional gene annotation of RNA-Seq hits

Annotation of the 50 most differentially detected hits from the RNA-seq analysis described above was performed using WormBase (version WS283) searches and BLAST searches against the *C. elegans* reference genome using WormBase BLAST/BLAT. The E-value threshold was set at 1E+0 and the *C. elegans* Sequencing Consortium genome project database was used. *C. elegans* genes with the highest homology were noted and their functional descriptions derived using WormBase WormMine. Stage- and sex-specific expression of hits were assigned based on previously published data (Foth et al., 201cl4).

Statistical analysis

Statistical tests and parameters used, including the definition of central value and the exact number (n) of mice or worms per group, are annotated in the corresponding figure legend. All analyses were performed with Graphpad Prism version 8.4.3 for Mac (GraphPad). The numbers of animals or biological replicates used herein were estimated on the basis of a power analysis with the following assumptions: the standard deviation will be roughly 20% of the mean; p values will be less than 0.05 when the null hypothesis is false; and the effect size (Cohen's d) is between 1.0 and 2.0. We have also carefully chosen the indicated sample size on the basis of empirical evidence of what is necessary to interpret the data and statistical significance.

Supplementary Figures

Fig. S1 Primary screen selection criteria, validation of *E. coli* mutants, and effect of mutant *E. coli* on *C. elegans*, related to Fig. 1. (A) Representative bubble plot from

secondary screen replicates. One of the six 96-well plates used in the secondary screen is shown here rendered using Vortex software data visualization tool. The size of the bubble represents percent of worms without embryos that is calculated from three technical replicates, while the color indicates the three biological trials. Boxed wells H12 and C3 indicate the negative control and a positive hit in two out of three replicates based on statistical tests. **(B)** Scatter plot showing the Robust Z (Rob Z) score for all primary screen wells scored. Scatter plot was generated using Vortex dotmatics software. Yellow dots indicate wells containing the negative control bacteria (GC1547). Dashed line indicates cut-off for Rob Z criterion. **(C)** Percentage of *C. elegans* bearing embryos in replicates of ten *E. coli* mutants identified from the secondary screen. Total n number of worms scored are (left to right): 505, 291, 418, 103, 879, 80, 586, 295, 474, 229, 1100, 501, 913, 411, 1253, 583, 1643, 550, 1561, 232. **(D)** Δ*eutN* and Δ*fabH* complementation suppresses *C. elegans* fertility defect. Total n number of worms scored is (left to right): 220, 197, 744, 1078. **(E)** Δ*eutD* complementation suppresses *C. elegans* fertility defect. Total n number of worms scored is 2276 (Δ*eutD*) and 2283 (Δ*eutD* with complementation). **(F, G)** Δ*fabF* phage transductant mimics *C. elegans* fertility defect of Δ*fabF* original clone. Circles represent mean of four independent biological replicates and bar is SEM. (F) Original *fabF* isolate from library (n = 330, 214 total worms scored, WT and Δ*fabF*, respectively), and (G) *fabF* transductant (n = 231, 66 total worms scored). **(H)** The number of progenitor zone nuclei of *glp-1(e2141)* at the L4 to Adult molt is not significantly different in worms raised on Δ*eutN*, Δ*fabH* or Δ*eutD* versus control *E. coli*. Scatter plots showing number of progenitor zone germ cell counts; each dot represents one gonad arm scored. We noted that in ~30% of *eutN* mutants (relative to 10% in control), the border of the PZ was difficult to detect; these were censored from PZ counts. The elevated penetrance of this phenotype in the Δ*eutN*-fed worms may represent a meiotic entry defect, accounting for the delay in germline development relative to somatic development. **(I)** No significant difference is observed between the number of progenitors in worms grown on BW25113 versus OP50 on standard NGM solid media. We note that the number of progenitors is markedly lower in the *glp-1(e2141)* mutant background after growth on liquid versus growth on solid media; the Keio starting strain was used in (H) since OP50 is sensitive

to kanamycin. **(J, K)** Fertility delay is not suppressed by loss of *daf-16,* nor is it unambiguously suppressed by loss of *daf-5*. Percent of worms with embryos on Day 1 in (J) *daf-16(+); daf-2(e1370) glp-1(e2141) or daf-16(m26); daf-2(e1370) glp-1(e2141)* and (K) *daf-5(+); glp-1(e2141) or daf-5(e1386); glp-1(e2141).* Loss of *daf-5* elevated the percentage of gravid worms on wild-type (WT) *E. coli*. Therefore, biological significance of a greater percentage of gravid *daf-5(e1386)* compared to *daf-5(+)* raised on the various mutant *E. coli* is unclear. Circles represent mean of independent biological replicates and bar is SEM. Total n number of worms scored are (left to right): **(J)** 269, 203, 207, 245, 179, 255, 229, 214, 200, 233 and (K) 460, 399, 386, 470, 379, 445, 388, 428, 554, 640. For (J), there is no significant suppression of the fertility delay in any case. Statistics: ANOVA and Student's t-test (H and I, respectively) and Fisher's Exact test (all other panels). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Fig. S2 *E. coli* **mutants used for** *T. muris* **egg hatching assay and effect of arginine and ornithine supplementation on** *C. elegans* **fertility, related to Fig. 2.** **(A)** Bacterial density of monoclonal cultures of each of the strains indicated measured by dilution plating; CFU = colony forming units, h = hours. **(B-C)** Hatching elicited by WT *E. coli* (WT), mutants (B) Δ*eutN* and (C) Δ*fabH,* and strains where deletions were complemented with a plasmid-borne gene (n = 3-4). One-way analysis of variance (ANOVA) with Dunnett's post-test compared with WT. **(D-E)** Percentage of *C. elegans* Day 1 adults with embryos raised on Δ*fabH E. coli* with arginine or ornithine supplementation. Fisher's exact test (Δ*fabH* alone plus or minus supplementation with 500μM arginine or 100μM ornithine); Total n number of worms scored (left to right): (D) 237, 300 and (E) 216, 149. For all panels, measurements were taken from distinct samples. Data represent the average of at least two independent experiments. Graphs show means and SEM. * p < 0.05, *** p < 0.001, **** p < 0.0001. ns = not significant.

Fig. S3 *E. coli*-derived and exogenous aldehydes reduce *T. muris* egg hatching, related to Fig. 3. (A) DarkZone standard curve for acetaldehyde. Fluorescence

intensity of samples incubated with known concentrations of acetaldehyde and 20µM DarkZone dye (n = 2). AU, arbitrary units. **(B)** Relative *eutR* expression in *E. coli* cells. eggs = pre-incubation with *T. muris* eggs for 4 hours. No RT = no reverse transcriptase control. Data pooled from 2 independent experiments. Analyzed using ANOVA with Tukey's multiple comparisons test. **(C)** Diagram of N-acetyl cysteine supplementation assay. Created using BioRender.com. **(D)** Quantification of hatching rates for mixtures containing WT *E. coli* cells and filtered supernatants supplemented with N-acetyl cysteine at the concentrations indicated (n = 2). **(E)** Quantification of hatching rates for *T. muris* eggs incubated with WT *E. coli* cultures supplemented with acetaldehyde at concentrations indicated (n = 3). Analyzed using ANOVA with Dunnett's post-test compared to 0. **(F)** Bacterial density of conditions indicated from (E) measured by dilution plating (n = 3). For all panels, measurements were taken from distinct samples. Graphs show means and SEM. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$.

Fig. S4 *T. muris* infection of germ-free and monocolonized mice, related to Fig 4.

(A) *E. coli* burden 7 dpi and 42 dpi (day of sacrifice) in stool of mice gavaged with 1x

10^8 CFU of strains indicated (n = 6-9 mice per group). **(B)** Cecum measurements taken on day of sacrifice for mice in (A). **(C)** Total worms harvested per mouse for germ-free (GF) and mice monocolonized with *E. coli* strains indicated; N.D. = not detected. **(D-G)** CD4$^+$ T cell percentage from mesenteric lymph nodes of WT and $\Delta eutN$ monocolonized mice. (D) Th1 (Tbet$^+$), (E) Th2 (GATA3$^+$), (F) Treg (FOXP3$^+$). (G) Cells were also stimulated with PMA and ionomycin and stained for intracellular IFN-γ expression, a cytokine known to be positively correlated with host susceptibility to *T. muris*. **(H)** Quantification of eggs laid per worm disaggregated by worm length. Short worms measured < 1 cm. N.D. = not detected. Harvested worms were incubated in individual wells of a 48-well plate with RPMI medium overnight and eggs laid were quantified using light microscopy. **(I)** Percentage of eggs subjected to 14 weeks of embryonation that hatched after incubation with WT *E. coli*. **(J)** Representative image of worms harvested from mice monocolonized with $\Delta eutN$ strain where *eutN* has been complemented on a plasmid = $\Delta eutN$ comp. (WT, n = 9; $\Delta eutN$ comp., n = 8). **(K)** Representative image of eggs laid by worms harvested from mice monocolonized with $\Delta eutN$ comp. **(L)** Proportion of total eggs laid per worm with normal morphology. Circles represent individual worms (WT, n = 36; $\Delta eutN$ comp., n = 29). **(M)** Heatmap of the 50 most differentially detected genes between WT- and mutant-associated worms generated using DESeq2. Colors indicate Row Z-score and clusters of genes with similar trends in expression between samples. Red indicates upregulated genes and blue indicates downregulated genes. WT = wildtype *E. coli*. L = *T. muris* larval stage. For all panels, measurements were taken from distinct samples. Data pooled from two independent experiments. Graph shows means and SEM. **** $p < 0.0001$.

Table S1 Top candidates and bacterial gene function (EcoCyc). *Fab* and *eut* genes were chosen for further characterization.

GENE	PROTEIN FUNCTION	FUNCTIONAL GROUP
fabF	β-ketoacyl-[acyl carrier protein] synthase II	Fatty acid biosynthesis
fabH	β-ketoacyl-[acyl carrier protein] synthase III	Fatty acid biosynthesis
eutD	phosphate acetyltransferase	Ethanolamine utilization
eutN	putative ethanolamine catabolic microcompartment shell protein	Ethanolamine utilization
pta	phosphate acetyltransferase	Ethanolamine utilization
rfaF	ADP-heptose—LPS heptosyltransferase 2	Lipopolysaccharide biosynthesis
rfaH	transcription antiterminator RfaH	Lipopolysaccharide biosynthesis
cmoA	carboxy-*S*-adenosyl-L-methionine synthase	Uridine modification in RNA
yhgN	putative inner membrane protein	Antibiotic resistance
holD	DNA polymerase III subunit ψ	DNA replication

Table S2 Single-worm Seq sample metadata, related to Figures 4 and S4. Data for
T. muris worms subjected to RNA extraction and sequencing analysis.

Worm ID	Sample name	run	*E. coli* associated	Worm length	Eggs laid	% normal eggs laid overnight
WT 1	MV4	1	wildtype	long	575	96
WT 2	MV5	1	wildtype	long	122	100
WT 3	MV17	2	wildtype	long	17	100
WT 4	MV18	2	wildtype	long	66	100
WT 5	MV19	2	wildtype	long	46	100
WT 6	MV1	1	wildtype	long	10	100
WT 7	MV3	1	wildtype	long	759	96
Mut 1	MV24	2	⊿eutN	short	59	2
Mut 2	MV40	2	⊿eutD	long	1	0
Mut 3	MV30	2	⊿eutD	short	2	0
Mut 4	MV33	2	⊿eutD	short	14	7
Mut 5	MV35	2	⊿eutD	long	1	0
Mut 6	MV14	1	⊿eutN	long	2	0
Mut 7	MV32	2	⊿eutD	short	6	0

Table S3 List of top 50 differentially detected genes, related to Figure S4E.
Information was collated from WormBase (version WS283) and Foth et al. (2014) for
stage specific expression data. *C. elegans* homologs were found using WormBase
BLAST/BLAT.

7. muris gene WormBase ID (increased expression in mutant) table

T. muris gene WormBase ID (increased expression in mutant)	WormBase Annotation	Gene Name	Other name	Seq Length (AA)	Stage specific exp (L2 vs L3, L3 vs adult, male vs female) (From: C. elegans Basidisijpment et al. 2014)	E-value if < 1		C. elegans gene(s) brief descriptions from WormBase/WormMine
WBGene00254232	Is predicted to enable calcium ion binding activity.	TMUE_100000 9514	TMUE_s01730	501 male	up in adults, male	2.00E-43 vyp-1	WBGene00018008	vyp-1 encodes the C. elegans VAP (VAMP-associated protein) ortholog, a transmembrane protein containing an N-terminal extracellular major sperm protein (MSP) domain; loss of vyp-1 function during development indicates that vyp-1 is required, both in vab-1-dependent and vab-1-independent pathways, for proper distal tip cell migration during somatic gonad development, ventral hypodermal cell migrations during embryonic enclosure, and anteriorly directed amphid neuron migration; like C. elegans MSPs, the VPR-1 MSP domain into Ig2; females induces oocyte maturation and gonadal sheath cell contraction; the VPR-1 MSP domain directly binds the VAB-1/Eph receptor extracellular domain in vitro and in vivo, binds the oocyte and gonadal sheath cell plasma membranes
WBGene00288269	n/a, protein coding	TMUE_200000 6815	TMUE_s00080	446 male	up in adults,			
WBGene00291903	n/a, protein coding	TMUE_200000 6496	TMUE_s00880	168 male	up in adults,	2.00E-08 vyp-1	WBGene00018008	Serpentine receptor class delta-26
WBGene00298538	n/a, protein coding	TMUE_200000 6270		113	up in adults,			
WBGene00288055	n/a, protein coding	TMUE_200000 0094	TMUE_s00390	456 male	up in adults,			
WBGene00288797	Is predicted to encode a protein with the following domain: Rho GTPase activation protein.	TMUE_200000	TMUE_s00380		up in adults,			
WBGene00298721	n/a, protein coding	TMUE_100000 6914	TMUE_s00140	175	up in adults,	0.37 nas-39	WBGene00003355	Zinc metalloprotease nas-39
WBGene00288099	n/a, protein coding	TMUE_300001 4904	TMUE_s00230	151 male	up in adults,	0.77 srd-25	WBGene00005103	
WBGene00290442	n/a, protein coding	TMUE_100000 5106	TMUE_s00580	237 male	up in adults, 26% ID, 52% +	0.19 F4SG6.8	WBGene00009960	RING-type domain-containing protein
WBGene00288079	n/a, protein coding	TMUE_200000 3015	TMUE_s00320	421 male	up in adults, 27% ID, 49% +	0.19 F4SG6.8	WBGene00003355	Zinc metalloprotease nas-39; nas-39 encodes an astacin family zinc metalloprotease and the only BMP-1/Tolloid homolog in C. elegans; in Drosophila and vertebrates nas-39 homologues activate TGF-beta signals by cleaving chordin/SOG, an inhibitor keeping the TGF-beta in inactive form; however developmental processes known to be affected by TGF-beta signaling are unaffected in nas-39 mutants, suggesting that nas-39 is not required for TGF-beta signaling; nas-39 is expressed in the pharynx, intestine, vulval and body wall muscles, and many neurons
WBGene00298863	Is predicted to encode a protein with the following domains: Zinc finger, Zinc finger C2H2-type, and TMUE_100000 4114	TMUE_100000 4114		539		4.00E-06 Y23H5A.4a	WBGene00021271	major sperm protein
WBGene00288514	Is predicted to encode a protein with the following domains: MSP (Major sperm protein) domain; PapD-like superfamily; and Immunoglobulin-like fold.	TMUE_300001 2031	TMUE_s00420	217 male	up in adults, 29% ID, 52% +	4.00E-06 Y23H5A.4a	WBGene00021271	major sperm protein
WBGene00288284	Is predicted to encode a protein with the following domains: MSP (Major sperm protein) domain; PapD-like superfamily; and Immunoglobulin-like fold.	TMUE_300001 7953	TMUE_s00390	183 male	up in adults, 47% ID, 75% +	M03E7.1	WBGene00019753	major sperm protein
WBGene00294631	Is predicted to encode a protein with the following domains: MSP (Major sperm protein) domain; Major sperm protein (MSP) domain; PapD-like superfamily; and Immunoglobulin-like fold.	TMUE_100000 3930	TMUE_s02040	239 male	up in L3, adult, 29% ID, 48% +	1.00E-13 pid-1	WBGene00016197	pid-1 encodes protein containing N-terminal LD motifs and C-terminal LM domains that is the C. elegans ortholog of vertebrate parafilin; pid-1 is required for organized myofilament structure in the pharynx and pharyngeal muscle contraction, and hence feeding and viability
WBGene00298580	Is predicted to encode a protein with the following domains: LIM domain and Zinc finger, LIM-type.	TMUE_100000 4317	TMUE_s00020	139 male	up in adults, 29% ID, 48% +	1.00E-13 pid-1	WBGene00016197	pid-1 encodes protein containing N-terminal LD motifs and C-terminal LM domains that is the C. elegans ortholog of vertebrate parafilin; pid-1 is required for organized myofilament structure in the pharynx and pharyngeal muscle contraction, and hence feeding and viability; past the first larval stage; PXL-1 is expressed in body wall and pharyngeal muscle, as well as pharyngeal epithelial cells; PXL-1 localizes to sites of actin attachment, namely dense bodies, adhesion plaques, and M-lines in body wall muscle, and podosome-like structures in pharyngeal muscle; pxl-1 expression in pharyngeal muscle and marginal cells is sufficient to rescue the pharyngeal muscle contraction defects and lethality of a pxl-1 mutant; PXL-1 physically interacts with muscle proteins DEB-1, UIG-1, LIM-8, UNC-96, UNC-98, and HUM-6.
WBGene00288184	n/a, protein coding	TMUE_100000 3466	TMUE_s00030	528 male	up in adults, 38% ID, 62% +	3.00E-51 ttx-7	WBGene00011283	Protein kinase domain-containing protein
WBGene00288793	Is predicted to enable ATP binding activity and protein kinase activity.	TMUE_100000 3767		339		3.00E-51 ttx-7	WBGene00011283	Protein kinase domain-containing protein
WBGene00296208	n/a, protein coding	TMUE_100000 5194		96				
WBGene00295526	Is predicted to enable phosphatase activity, is an ortholog of C. elegans egg-3, egg-4, and egg-5.	TMUE_300001 4972	TMUE_s03230	366 male	up in adults, egg-3, egg-4, egg-5	egg-3, egg-4, egg-5	WBGene00009701, WBGene00020682, WBGene00020685	egg-3 encodes a member of the protein tyrosine phosphatase-like (PTPL) family; EGG-3 activity is required for proper regulation of the cell surface and cortical rearrangements that occur during egg activation in response to spermmeiosis; EGG-3 physically interacts with CHS-1 and MBK-2 in the oocyte prior to meiosis, thus preventing MBK-2 from phosphorylating its substrates, such as cytoskeletal MEI-1/katanin; EGG-3 is required for normal embryonic and larval viability, embryonic polarity, eggshell formation, and the block to polyspermy; EGG-3 initially localizes to the oocyte cortex, but during anaphase of meiosis I moves into the cytoplasm where it is degraded allowing for activation of MBK-2; egg-4 encodes a pseudotyrosine phosphatase; egg-4 along with egg-5 which also encodes a pseudotyrosine phosphatase, are required for the egg-to-oocyte transition and embryonic development; EGG-4 and EGG-5 function to negatively regulate the activity of the MBK-2 DYRK family kinase by sequestering MBK-2 at the oocyte cortex and, in the case of EGG-4, by inhibiting its enzymatic activity; EGG-4 physically interacts with MBK-2 and 2 in vitro assays, appears to function as a mixed-type inhibitor of MBK-2 kinase activity; EGG-4 localizes to the oocyte cortex and nuclei; egg-5 encodes a pseudotyrosine phosphatase; egg-5, along with egg-4 which also encodes a pseudotyrosine phosphatase, are required for the egg-to-oocyte transition and embryonic development; EGG-4 and EGG-5 function to negatively regulate the activity of the MBK-2 DYRK family kinase by sequestering MBK-2 at the oocyte cortex and, in the case of EGG-4 by inhibiting its enzymatic activity; EGG-5 physically interacts with MBK-2; EGG-5 localizes to the oocyte cortex and nuclei
WBGene00291963	Is predicted to encode a protein with the following domains: Dynein light chain type 1; Dynein light chain, type 1/2; and Dynein light chain superfamily.	TMUE_100000 4981	TMUE_s00760	275 male	up in adults, 32% ID, 59% +	5.00E-33 pmp-1		dlc-1 encodes a dynein light chain 1 required for viability, that inhibits DHC-1 in vivo; DLC-1 is 92-95% identical to human dyneins b DLC-2/c-6; DLC-1 is expressed in adult body wall muscle, and in larval and adult pharynx, posterior intestine, and nervous system (e.g., ventral nerve cord, head neurons, and tail neurons); dlc-1 is probably transcriptionally activated by DAF-19; DLC-1 is required for normal embryonic and larval viability, embryonic pronuclear migration, growth speed, fertility, brood size, body shape, and oocyte integrity; anterograde movements of DLC-1 in neurons are regulated by PTL-1/Tau.
WBGene00293961	n/a, protein coding	TMUE_200000 8700	TMUE_s01440	72 male	up in adults,			
WBGene00299157	n/a, protein coding	TMUE_200000 2749	TMUE_s02550	221 male	up in adults, 41% ID, 64% +	4.00E-18 dlc-1	WBGene00001005	dlc-1 encodes a dynein light chain 1 required for viability, that inhibits DHC-1 in vivo; DLC-1 is 92-95% identical to human dyneins b DLC-2/c-6; DLC-1 is expressed in adult body wall muscle, and in larval and adult pharynx, posterior intestine, and nervous system (e.g., ventral nerve cord, head neurons, and tail neurons); dlc-1 is probably transcriptionally activated by DAF-19; DLC-1 is required for normal embryonic and larval viability, embryonic pronuclear migration, growth speed, fertility, brood size, body shape, and oocyte integrity; anterograde movements of DLC-1 in neurons are regulated by PTL-1/Tau.
WBGene00302146	Is predicted to encode a protein with the following domains: Dynein light chain type 1; Dynein light chain, type 1/2 and Dynein light chain superfamily.	TMUE_300001 4161		92		4.00E-18 dlc-1	WBGene00001005	NDZ07:1 is orthologous to the human gene PURINE NUCLEOSIDE PHOSPHORYLASE (NP; OMIM:164050), which when mutated leads to disease.
WBGene00287425	Is predicted to encode a protein tyrosine phosphatase activity; is an ortholog of C. elegans egg-3, egg-4 and egg-5.	TMUE_200000 6534	TMUE_s00160 11300	445 male	up in adults,	egg-3, egg-4, egg-5	WBGene00009701, WBGene00020682, WBGene00020685	egg-3 encodes a member of the protein tyrosine phosphatase-like (PTPL) family; EGG-3 activity is required for proper regulation of the cell surface and cortical rearrangements that occur during egg activation in response to spermmeiosis; EGG-3 physically interacts with CHS-1 and MBK-2 in the oocyte prior to meiosis, thus preventing MBK-2 from phosphorylating its substrates, such as cytoskeletal MEI-1/katanin; EGG-3 initially localizes to the oocyte cortex, but during anaphase of meiosis I moves into the cytoplasm where it is degraded allowing for activation of MBK-2.
WBGene00302548	Is predicted to encode a protein with the following domains: Histidine phosphatase superfamily (branch 1); Histidine phosphatase superfamily, clade-1; and Histidine phosphatase superfamily.	TMUE_100000 4607	TMUE_010500 2600	229				

WBGene ID	Description	TMUE code 1	TMUE code 2	Count/sex	Expression	% ID	E-value	Ortholog	WBGene ref	Protein / long description
WBGene00028471	Is predicted to enable ATP binding activity and protein kinase activity; Is an ortholog of C. elegans M7.7.	TMUE_300001 _G5300	TMUE_300001 _G0280	361 male	up in L3, adults,	28% ID, 49% +	4.00E-38	ttbk-7	WBGene00011283	Protein kinase domain-containing protein
WBGene00029062	n/a, protein coding	TMUE_100000 _G0700	TMUE_300001 _01900	163 male	up in adults,					
WBGene00034130	n/a, n/a	TMUE_010000 _0892	n/a							
WBGene00297748	Is predicted to enable peptidase inhibitor activity; Is predicted to encode a protein with the following domain: Cystine-knot cytokine.	TMUE_000000 1881	TMUE_300001 _G0800	94		41% ID, 52% +	0.87	R12A1.3a	WBGene00020017	BPTI/Kunitz inhibitor domain-containing protein
WBGene00029614	Is predicted to encode a protein with the following domain: Cystine-knot cytokine.	TMUE_300001 _0896	TMUE_300001 _G0820	286 male	up in L3, adults,					
WBGene00029613	n/a, protein coding	TMUE_300001 _0907	TMUE_300001 _02700	297 male	up in L3, adults,					
WBGene00301102	n/a, protein coding	TMUE_300001 _0868		286						
WBGene00028974	n/a, protein coding	TMUE_300001 _3984	TMUE_300001 _05900	84 male	up in adults,					
WBGene00028479	Is predicted to enable nucleic acid binding activity; Is an ortholog of C. elegans alg-3 and alg-4. In C. elegans alg-4 is involved in spermatogenesis.	TMUE_300001 _2942	TMUE_300001 _04300	918 male	up in L3, adults,			alg-3, alg-4	WBGene00011910, WBGene00006649	alg-3 encodes an Argonaute (AGO) protein that is a member of the AGO clade of Argonaute small RNA-binding proteins; together with its paralog, alg-4, alg-3 is required for normal brood sizes at elevated temperatures, specifically for the process of spermatogenesis in which spermatids are converted to mobile amoeboid sperm; alg-3/4 is also required for accumulation of male germline-specific 26G-RNAs, suggesting that alg-3/4 mediate thermotolerant fertility via a 26G-RNA-mediated pathway; a GFP::ALG-3 rescuing fusion protein is expressed in germ cells during spermatogenesis and localizes to the cytoplasm and to P granules.
WBGene00301801	n/a, protein coding	TMUE_300001 _3156		145						
WBGene00028504	Is predicted to enable nucleic acid binding activity; Is an ortholog of C. elegans alg-3 and alg-4. In C. elegans alg-4 is involved in spermatogenesis.	TMUE_100000 _3405	TMUE_300001 _G0020 13100	880 male	up in L3, adults,			alg-3, alg-4	WBGene00011910, WBGene00006649	alg-4 encodes an Argonaute (AGO) protein that is a member of the AGO clade of Argonaute small RNA-binding proteins; together with its paralog, alg-3, alg-4 is required for normal brood sizes at elevated temperatures, specifically for the process of spermatogenesis in which spermatids are converted to mobile amoeboid sperm; alg-3/4 is also required for accumulation of male germline-specific 26G-RNAs, suggesting that alg-3/4 mediate thermotolerant fertility via a 26G-RNA-mediated pathway.
WBGene00287204	Is predicted to enable alpha-amylase activity and cation binding activity; Is an ortholog of C. elegans C59B6.7	TMUE_200000 _9275	TMUE_200000 _9275	695				C59B6.7	WBGene00009820	predicted to enable alpha-amylase activity, maltose(ase), and metal ion binding activity; Is expressed in intestine; Is an ortholog of human AMY2A (amylase alpha 2A) and AMY2B (amylase alpha 2B).
WBGene00032877	n/a, protein coding	TMUE_300001 _2979 4200	TMUE_300001 _001000	313						
WBGene00030047	n/a, protein coding	TMUE_200000 _8654		142						
WBGene00301660	n/a, protein coding	TMUE_300001 _3344		781						
WBGene00024606	Is predicted to encode a protein with the following domains: Tsp2/MMR family; Tsp2/MMR-related protein, Tsp2/MMR-related superfamily.	TMUE_100000 _4599 03800	TMUE_100000 _008900 03800	238 female	up in L3, adults,					
WBGene00302075	n/a, protein coding	TMUE_100000 _3330 4500		174						
WBGene00301915	n/a, protein coding	TMUE_300001 _3424		277						
WBGene00298627	n/a, protein coding	TMUE_200000 _6954		69						
WBGene00296619	Is predicted to encode a protein with the following domain:	TMUE_000000 _0071		136						
WBGene00029071	Is predicted to enable chitin binding activity; Is an ortholog of C. elegans C39D10.7	TMUE_300001 _4378 05000	TMUE_300001 _G0920 05000	842 female	up in adults,	23% ID, 34% +	8.00E-20	C39D10.7	WBGene00016934	C39D10.7 encodes a large (1185-residue) protein with 11 chitin-binding-type-A domains; C39D10.7 transcripts are enriched in/enriched in germcells; like CEJ-1 and CPG-2, C39D10.7 might participate in/eggshell synthesis and early embryonic development, and in which spermatids are converted; however, C39D10.7 alone has no visible function in male RNAi assays.
WBGene00029893	n/a, protein coding	TMUE_300001 _4020 02000	TMUE_300001 _G0690 02000	166 female	up in adults,					
WBGene00029894	n/a, protein coding	TMUE_100000 _1882 02100	TMUE_300001 _G0020 02100	947 female	up in adults,					
WBGene00028558	Is predicted to enable lipid binding activity.	TMUE_100000 _3367	TMUE_300001 _19100	2006 female	up in L3, adults,					
WBGene00029875	n/a, protein coding	TMUE_300001 _1875	TMUE_300001 _x0680 02200	90 male	up in adults,	33% ID, 51% +	4.00E-16	dlc-1	WBGene00001005	dlc-1 encodes a dynein light chain 1, required for viability, that inhibits DHC-1 in vivo; DLC-1 is 92-95% identical to its orthologs human DYNLL1 (OMIM:601562) and DYNLL2 (OMIM:608942); DLC-1 is paralogous to DLC2A-6; DLC-1 is expressed in adult body wall muscle, and in larval and adult pharynx, posterior intestine, and nervous system (e.g., ventral nerve cord, head neurons, and tail neurons); dlc-1 is probably transcriptionally activated by DAF-19; DLC-1 is required for normal embryonic and larval viability, embryonic promotora migration, growth/speed, and fertility, brood size, normal cuticular integrity; anterograde movements of DLC-1 in neurons are regulated by PTL-1/Tau.

Table S4 Strains used in this study.

STRAIN	SPECIES	GENOTYPE	REFERENCE
BW25113	*E. coli*	*F-, Δ(araD-araB)567, ΔlacZ4787*(::rrnB-3), *λ-, rph-1, Δ(rhaD-rhaB)568, hsdR514*	(Baba et al., 2006)
GC833	*C. elegans*	*glp-1(ar202)* III	(Pepper et al., 2003)
GC1250	*C. elegans*	*daf-16(m26) I; daf-2(e1370) glp-1(e2141)* III	(Michaelson et al., 2010)
GC1474	*C. elegans*	*glp-1(e2141)* III; hjSi20 [*Pmyo-2P::mCherry::unc-54* 3'UTR] IV; *zuIs70* [pJN152: *end-1P::gfp::caax; unc-119*(+)] V	This study
GC1545	*C. elegans*	*daf-5(e1386) II; glp-1(e2141)* III	This study
GC1547	*E. coli*	BW25113 transformed with pET28	This study
GC1549	*E. coli*	BW25113 Δf*abF*::*kan*	This study
GC1558	*E. coli*	BW25113 Δ*eutN*::kan, pGC735	This study
GC1559	*E. coli*	BW25113 Δf*abH*::kan, pGC738	This study
GC1643	*E. coli*	BW25113 Δ*eutC*::*kan*	This study
GC1644	*E. coli*	BW25113 Δ*eutC*	This study
GC1645	*E. coli*	BW25113 Δ*eutN*::*kan*	This study
GC1646	*E. coli*	BW25113 Δ*eutN*	This study
GC1647	*E. coli*	BW25113 Δ*eutC* Δ*eutN*::*kan*	This study
GC1648	*E. coli*	BW25113 Δ*eutC*::*kan* Δ*eutN*	This study
GC1649	*E. coli*	BW25113 Δ*eutD*	This study
GC1650	*E. coli*	BW25113 Δ*eutD* pPlac-eutD	This study
JW1077	*E. coli*	BW25113 Δ*fabH*::*kan*	(Baba et al., 2006)
JW2442	*E. coli*	BW25113 Δ*eutD*::kan	
JW2440	*E. coli*	BW25113 Δ*eutN*::*kan*	

JW2294	*E. coli*	BW25113 Δ*pta*::*kan*	
JW3595	*E. coli*	BW25113 Δ*rfaF*::*kan*	
JW3818	*E. coli*	BW25113 Δ*rfaH*::*kan*	
JW4334	*E. coli*	BW25113 Δ*holD*::*kan*	
JW1859	*E. coli*	BW25113 Δ*cmoA*::*kan*	
JW3397	*E. coli*	BW25113 Δ*yhgN*::*kan*	
JW3929	*E. coli*	BW25113 Δ*argE*::*kan*	
JW3932	*E. coli*	BW25113 Δ*argH*::*kan*	

Table S5 Primers used in this study.

Primer	Details
GCo2700 – 5'…. CCGCGCTGGTTCGTTTCtagTGCAGCAAT GGCAACAACGTTG	Forward primer- vector fragment to amplify pBR322 for Gibson assembly with PCR product bearing wild-type *fabH* sequence
GCo2701 – 5'…. ACGTTGTTGCCATTGCTGCActaGAAACGA ACCAGCGCG	Reverse primer- vector fragment to amplify pBR322 for Gibson assembly with PCR product bearing wild-type *fabH* sequence
GCo2702 – 5'…. AGCTGTCAAACATGAGAATTATAACCGAA AAGTGACTGAGCGTACa	Forward primer- *fabH* coding sequence for Gibson assembly with pBR322
GCo2703 – 5'…. CTCAGTCACTTTTCGGTTATAATTCTCAT GTTTGACAGCTTATCATCGATAAG	Reverse primer- *fabH* coding sequence for Gibson assembly with pBR322
GCo2708 – 5'…. agctgtcaaacatgagaattAATACGCCACGGAG GCG	Forward primer- *eutN* coding sequence for Gibson assembly with pBR322
GCo2709 – 5'…. acgttgttgccattgctgcattaTTTGTGGAAAATTAC CTGACCGCC	Reverse primer- *eutN* coding sequence for Gibson assembly with pBR322
GCo2710 – 5'…. ACCCGCCTCCGTGGCGTATTaattctcatgtttg acagcttatcatcgataag	Forward primer- vector fragment to amplify pBR322 for Gibson assembly with PCR product bearing wild-type *eutN* sequence
GCo2711 – 5'…. AGGTAATTTTCCACAAAtaatgcagcaatggcaa caacgt	Reverse primer- vector fragment to amplify pBR322 for Gibson assembly

	with PCR product bearing wild-type *eutN* sequence
GCo2744 – 5'…. GCCGGGAAGCTAGAGTAAGTAGT	pBR322 forward (used to confirm gene sequence)
GCo2745 – 5'…. AAAAACATATGATTATTGAACGTTGTCGTG	Forward primer- *eutD* coding sequence with an NdeI site for cloning into pPlacRppH6
GCo2746 – 5'…. TTTTTTTCTAGAGTTCATTCAACCAGTGTTTGTAAAC	Reverse primer- *eutD* coding sequence with an XbaI site for cloning into pPlacRppH6
GCo2747 – 5'…. CGCGTTGGCCGATTCATTAATGCAGC	pPlacRppH6 forward (used to confirm gene sequence)
GCo2748 – 5'…. GCTTCGCAACGTTCAAATCCGCTCC	pPlacRppH6 reverse (used to confirm gene sequence)
ACTCCTACGGGAGGCAGCAGT	Forward primer- 16S coding sequence for qPCR
ATTACCGCGGCTGCTGGC	Reverse primer- 16S coding sequence for qPCR
GGGAGGCACCGAGTTTGAA	Forward primer- *eutR* coding sequence for qPCR
GGGAGGCACCGA GTTTGAA	Reverse primer- *eutR* coding sequence for qPCR

Table S6 Plasmid names and descriptions.

Plasmid name	Description
pGC735	*eutN* cloned into pBR322 by Gibson assembly
pGC737	pET28
pGC738	*fabH* cloned into pBR322 by Gibson assembly

| pPlac-eutD | *eutD* under the control of a *lac*UV5 promoter |
| pBR322 | pBR322 |

Table S7 Primer sequences to verify kan cassette insertion within target genes.

GCo296 AGCTTTGAAGTATTACGCCCGGAC	**Forward PCR primer *eutD***
GCo2797 CCTTTCGAGCACGGTCTACG	**Forward sequencing primer *eutD***
GCo2798 AACACCAACCAGCTTGACGC	**Reverse PCR primer *eutD***
GCo2799 AGGCCCCGGGTTTCGATC	**Reverse sequencing primer *eutD***
GCo2800 CTTCCCGATCGGCCTGAAAGG	**Forward PCR primer *eutN***
GCo2801 CTGTAATCACCCTGTGTGACGTCG	**Forward sequencing primer *eutN***
GCo2802 ACTTTGGCTGCCGCAACGGC	**Reverse PCR primer *eutN***
GCo2803 CGCGAAAACGCCCATCTCATG	**Reverse sequencing primer *eutN***
GCo2804 AATTGCCGCTCGCCTGGA	**Forward PCR primer *fabH***

GCo2805 GGTTTTGAGCTGCTGGACGG	**Forward sequencing primer *fabH***
GCo2806 GCCGCAGAAGCTTCAGCAAAC	**Reverse PCR primer *fabH***
GCo2807 GGATAGCTCGCCGCCATATCA	**Reverse sequencing primer *fabH***
GCo2808 GTCCTGCGGTGGTTATCCCAA	**Forward PCR primer *pta***
GCo2809 AAGAACTGGTTATCGCGCAAG	**Forward sequencing primer *pta***
GCo2810 AAGTGGGATGGCGCAATTCAT	**Reverse PCR primer *pta***
GCo2811 GCAGCGCAGTTAAGCAAGATAA	**Reverse sequencing primer *pta***
GCo2812 AATCACCACCGTTCAGGCTGC	**Forward PCR primer *fabF***
GCo2813 GCCACCAGGCGTAAGTGAAC	**Forward sequencing primer *fabF***
GCo2814 CCGTTAATTAAGAACATACCGGCTCC	**Reverse PCR primer *fabF***

GCo2815 AGTGTGGCAGCATGTTCACTACG	**Reverse sequencing primer *fabF***
GCo2832 AACTGAAAGGCCGCTATCAGG	**Forward PCR primer** ***rfaF***
GCo2833 GGTTACGACAAACCGTTCAAAACC	**Forward sequencing primer *rfaF***
GCo2834 GCCAGGAAGGAATCTGTGCGAA	**Reverse PCR primer** ***rfaF***
GCo2835 CCACCCAGTCAAACTTAATCCCTG	**Reverse sequencing primer** ***rfaF***
GCo2836 GGCGTTCATCTTTGCGATGCTG	**Forward PCR primer** ***rfaH***
GCo2837 CTGACGGTATAACGCAAACCGG	**Forward sequencing primer** ***rfaH***
GCo2838 TCTCACGCCAAAGCCATCATCC	**Reverse PCR primer** ***rfaH***
GCo2839 CCATATTTTGCAACGTATTGCGCAC	**Reverse sequencing primer** ***rfaH***
GCo2840 CTCTTGGCACATCTTTCACCATACA	**Forward PCR primer** ***yhgN***
GCo2841	**Forward sequencing primer *yhgN***

CGAAGCCATCAGTAATGCGACTT	
GCo2842 CCGAAACGCTGAAAACCTGGAC	**Reverse PCR primer** *yhgN*
GCo2843 GGATGATTGCGTTCCAGCTGTT	**Reverse sequencing primer** *yhgN*
GCo2844 TGTGCGCTGTTGCTGATGGT	**Forward PCR primer** *cmoA*
GCo2845 TTTGGTATATGCCCTGGGAGTTGG	**Forward sequencing primer** *cmoA*
GCo2846 GGTAGAAACTCCACCGCATTTGA	**Reverse PCR primer** *cmoA*
GCo2847 TTAAACAACCCGTGCTGCTGCT	**Reverse sequencing primer** *cmoA*
GCo2848 TGTGACGCAGCAAGACTTCACT	**Forward PCR primer** *holD*
GCo2849 ATGCAGACATGGAATGCTCCTCAA	**Forward sequencing primer** *holD*
GCo2850 ACCACTTGCGTAATCGCAAACG	**Reverse PCR primer** *holD*
GCo2851 TTGCCGTTTTGCGTTAACTGAAAGT	**Reverse sequencing primer** *holD*

References

Agarwal, I., Farnow, C., Jiang, J., Kim, K.S., Leet, D.E., Solomon, R.Z., Hale, V.A., and Goutte, C. (2018). HOP-1 Presenilin Deficiency Causes a Late-Onset Notch Signaling Phenotype That Affects Adult Germline Function in Caenorhabditis elegans. Genetics *208*, 745-762.

Antignano, F., Mullaly, S.C., Burrows, K., and Zaph, C. (2011). Trichuris muris infection: a model of type 2 immunity and inflammation in the gut. J Vis Exp.

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K.A., Tomita, M., Wanner, B.L., and Mori, H. (2006). Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst Biol *2*, 2006 0008.

Bancroft, A.J., Else, K.J., and Grencis, R.K. (1994). Low-level infection with Trichuris muris significantly affects the polarization of the CD4 response. European journal of immunology *24*, 3113-3118.

Boeck, M.E., Huynh, C., Gevirtzman, L., Thompson, O.A., Wang, G., Kasper, D.M., Reinke, V., Hillier, L.W., and Waterston, R.H. (2016). The time-resolved transcriptome of C. elegans. Genome research *26*, 1441-1450.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics *30*, 2114-2120.

Chen, W.W., Freinkman, E., Wang, T., Birsoy, K., and Sabatini, D.M. (2016). Absolute Quantification of Matrix Metabolites Reveals the Dynamics of Mitochondrial Metabolism. Cell *166*, 1324-1337 e1311.

Cheng, K.C.-C., Klancer, R., Singson, A., and Seydoux, G. (2009). Regulation of MBK-2/DYRK by CDK-1 and the pseudophosphatases EGG-4 and EGG-5 during the oocyte-to-embryo transition. Cell *139*, 560-572.

Cherepanov, P.P., and Wackernagel, W. (1995). Gene disruption in Escherichia coli: TcR and KmR cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance determinant. Gene *158*, 9-14.

Conine, C.C., Batista, P.J., Gu, W., Claycomb, J.M., Chaves, D.A., Shirayama, M., and Mello, C.C. (2010). Argonautes ALG-3 and ALG-4 are required for spermatogenesis-specific 26G-RNAs and thermotolerant sperm in Caenorhabditis elegans. Proceedings of the National Academy of Sciences *107*, 3588-3593.

D'Elia, R., Behnke, J.M., Bradley, J.E., and Else, K.J. (2009). Regulatory T cells: a role in the control of helminth-driven intestinal pathology and worm survival. The journal of immunology *182*, 2340-2348.

Dalfo, D., Michaelson, D., and Hubbard, E.J. (2012). Sensory regulation of the C. elegans germline through TGF-beta-dependent signaling in the niche. Curr Biol *22*, 712-719.

Deana, A., Celesnik, H., and Belasco, J.G. (2008). The bacterial enzyme RppH triggers messenger RNA degradation by 5′ pyrophosphate removal. Nature *451*, 355-358.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Donohue, T.M., Jr., Tuma, D.J., and Sorrell, M.F. (1983). Acetaldehyde adducts with proteins: binding of [14C]acetaldehyde to serum albumin. Arch Biochem Biophys *220*, 239-246.

Fahmy, M.A. (1954). An investigation on the life cycle of Trichuris muris. Parasitology *44*, 50-57.

Foth, B.J., Tsai, I.J., Reid, A.J., Bancroft, A.J., Nichol, S., Tracey, A., Holroyd, N., Cotton, J.A., Stanley, E.J., and Zarowiecki, M. (2014). Whipworm genome and dual-species transcriptome analyses provide molecular insights into an intimate host-parasite interaction. Nature genetics *46*, 693-700.

Gracida, X., and Eckmann, C.R. (2013). Fertility and germline stem cell maintenance under different diets requires nhr-114/HNF4 in C. elegans. Curr Biol *23*, 607-613.

Hayes, K.S., Bancroft, A.J., Goldrick, M., Portsmouth, C., Roberts, I.S., and Grencis, R.K. (2010). Exploitation of the intestinal microflora by the parasitic nematode Trichuris muris. Science *328*, 1391-1394.

Holm, J.B., Sorobetea, D., Kiilerich, P., Ramayo-Caldas, Y., Estelle, J., Ma, T., Madsen, L., Kristiansen, K., and Svensson-Frej, M. (2015). Chronic Trichuris muris Infection Decreases Diversity of the Intestinal Microbiota and Concomitantly Increases the Abundance of Lactobacilli. PLoS One *10*, e0125495.

Houlden, A., Hayes, K.S., Bancroft, A.J., Worthington, J.J., Wang, P., Grencis, R.K., and Roberts, I.S. (2015). Chronic Trichuris muris Infection in C57BL/6 Mice Causes Significant Changes in Host Microbiota and Metabolome: Effects Reversed by Pathogen Clearance. PLoS One *10*, e0125945.

Hubbard, E.J.A., and Schedl, T. (2019). Biology of the Caenorhabditis elegans Germline Stem Cell System. Genetics *213*, 1145-1188.

Irnov, I., Wang, Z., Jannetty, N.D., Bustamante, J.A., Rhee, K.Y., and Jacobs-Wagner, C. (2017). Crosstalk between the tricarboxylic acid cycle and peptidoglycan synthesis in Caulobacter crescentus through the homeostatic control of alpha-ketoglutarate. PLoS Genet *13*, e1006978.

Jenkins, T.P., Rathnayaka, Y., Perera, P.K., Peachey, L.E., Nolan, M.J., Krause, L., Rajakaruna, R.S., and Cantacessi, C. (2017). Infections by human gastrointestinal helminths are associated with changes in faecal microbiota diversity and composition. PLoS One *12*, e0184719.

Kaval, K.G., and Garsin, D.A. (2018). Ethanolamine Utilization in Bacteria. mBio *9*.

Kocsisova, Z., Kornfeld, K., and Schedl, T. (2019). Rapid population-wide declines in stem cell number and activity during reproductive aging in C. elegans. Development *146*.

Kolde, R. (2015). pheatmap: Pretty Heatmaps. R package version 1.0. 8.

Korta, D.Z., Tuck, S., and Hubbard, E.J. (2012). S6K links cell fate, cell cycle and nutrient response in C. elegans germline stem/progenitor cells. Development *139*, 859-870.

Koyama, K. (2016). Bacteria-induced hatching of Trichuris muris eggs occurs without direct contact between eggs and bacteria. Parasitol Res *115*, 437-440.

Lee, S.C., Tang, M.S., Lim, Y.A., Choy, S.H., Kurtz, Z.D., Cox, L.M., Gundra, U.M., Cho, I., Bonneau, R., Blaser, M.J.*, et al.* (2014). Helminth colonization is associated with increased diversity of the gut microbiota. PLoS Negl Trop Dis *8*, e2880.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Martin, P.K., Marchiando, A., Xu, R., Rudensky, E., Yeung, F., Schuster, S.L., Kernbauer, E., and Cadwell, K. (2018). Autophagy proteins suppress protective type I interferon signalling in response to the murine gut microbiota. Nat Microbiol *3*, 1131-1141.

Maruyama, R., Velarde, N.V., Klancer, R., Gordon, S., Kadandale, P., Parry, J.M., Hang, J.S., Rubin, J., Stewart-Michaelis, A., and Schweinsberg, P. (2007). EGG-3 regulates cell-surface and cortex rearrangements during egg activation in Caenorhabditis elegans. Current Biology *17*, 1555-1560.

Michaelson, D., Korta, D.Z., Capua, Y., and Hubbard, E.J. (2010). Insulin signaling promotes germline proliferation in C. elegans. Development *137*, 671-680.

Mok, D.Z., Sternberg, P.W., and Inoue, T. (2015). Morphologically defined sub-stages of C. elegans vulval development in the fourth larval stage. BMC Dev Biol *15*, 26.

Pacold, M.E., Brimacombe, K.R., Chan, S.H., Rohde, J.M., Lewis, C.A., Swier, L.J., Possemato, R., Chen, W.W., Sullivan, L.B., Fiske, B.P.*, et al.* (2016). A PHGDH inhibitor reveals coordination of serine synthesis and one-carbon unit fate. Nat Chem Biol *12*, 452-458.

Panesar, T. (1989). The moulting pattern in Trichuris muris (Nematoda: Trichuroidea). Canadian journal of zoology *67*, 2340-2343.

Parry, J.M., Velarde, N.V., Lefkovith, A.J., Zegarek, M.H., Hang, J.S., Ohm, J., Klancer, R., Maruyama, R., Druzhinina, M.K., and Grant, B.D. (2009). EGG-4 and EGG-5 link events of the oocyte-to-embryo transition with meiotic progression in C. elegans. Current Biology *19*, 1752-1757.

Pekar, O., Ow, M.C., Hui, K.Y., Noyes, M.B., Hall, S.E., and Hubbard, E.J.A. (2017). Linking the environment, DAF-7/TGFbeta signaling and LAG-2/DSL ligand expression in the germline stem cell niche. Development *144*, 2896-2906.

Penrod, J.T., and Roth, J.R. (2006). Conserving a volatile metabolite: a role for carboxysome-like organelles in Salmonella enterica. J Bacteriol *188*, 2865-2874.

Pepper, A.S., Lo, T.W., Killian, D.J., Hall, D.H., and Hubbard, E.J. (2003). The establishment of Caenorhabditis elegans germline pattern is controlled by overlapping proximal and distal somatic gonad signals. Dev Biol *259*, 336-350.

Ramanan, D., Bowcutt, R., Lee, S.C., Tang, M.S., Kurtz, Z.D., Ding, Y., Honda, K., Gause, W.C., Blaser, M.J., Bonneau, R.A.*, et al.* (2016). Helminth infection promotes colonization resistance via type 2 immunity. Science *352*, 608-612.

Rausch, S., Midha, A., Kuhring, M., Affinass, N., Radonic, A., Kuhl, A.A., Bleich, A., Renard, B.Y., and Hartmann, S. (2018). Parasitic Nematodes Exert Antimicrobial Activity and Benefit From Microbiota-Driven Support for Host Immune Regulation. Front Immunol *9*, 2282.

Roof, D.M., and Roth, J.R. (1989). Functions required for vitamin B12-dependent ethanolamine utilization in Salmonella typhimurium. J Bacteriol *171*, 3316-3323.

Rosa, B.A., Snowden, C., Martin, J., Fischer, K., Kupritz, J., Beshah, E., Supali, T., Gankpala, L., Fischer, P.U., Urban, J.F., Jr.*, et al.* (2021). Whipworm-Associated Intestinal Microbiome Members Consistent Across Both Human and Mouse Hosts. Front Cell Infect Microbiol *11*, 637570.

Rowley, C.A., Sauder, A.B., and Kendall, M.M. (2020). The ethanolamine-sensing transcription factor EutR promotes virulence and transmission during Citrobacter rodentium intestinal infection. Infection and immunity *88*, e00137-00120.

Roy, D., Kahler, D.J., Yun, C., and Hubbard, E.J.A. (2018). Functional Interactions Between rsks-1/S6K, glp-1/Notch, and Regulators of Caenorhabditis elegans Fertility and Germline Stem Cell Maintenance. G3 (Bethesda) *8*, 3293-3309.

Salinas, G., and Risi, G. (2018). Caenorhabditis elegans: nature and nurture gift to nematode parasitologists. Parasitology *145*, 979-987.

Schachter, J., Alvarinho de Oliveira, D., da Silva, C.M., de Barros Alencar, A.C.M., Duarte, M., da Silva, M.M.P., Ignacio, A., and Lopes-Torres, E.J. (2020). Whipworm Infection Promotes Bacterial Invasion, Intestinal Microbiota Imbalance, and Cellular Immunomodulation. Infect Immun *88*.

Sepulveda-Crespo, D., Reguera, R.M., Rojo-Vazquez, F., Balana-Fouce, R., and Martinez-Valladares, M. (2020). Drug discovery technologies: Caenorhabditis elegans as a model for anthelmintic therapeutics. Med Res Rev *40*, 1715-1753.

Shaham, S. (2006). WormBook. Methods in cell biology.

Simon-Manso, Y., Lowenthal, M.S., Kilpatrick, L.E., Sampson, M.L., Telu, K.H., Rudnick, P.A., Mallard, W.G., Bearden, D.W., Schock, T.B., Tchekhovskoi, D.V.*, et al.* (2013). Metabolite profiling of a NIST Standard Reference Material for human plasma (SRM 1950): GC-MS, LC-MS, NMR, and clinical laboratory analyses, libraries, and web-based resources. Anal Chem *85*, 11725-11731.

Smith, C.A., O'Maille, G., Want, E.J., Qin, C., Trauger, S.A., Brandon, T.R., Custodio, D.E., Abagyan, R., and Siuzdak, G. (2005). METLIN: a metabolite mass spectral database. Ther Drug Monit *27*, 747-751.

Sorrell, M.F., and Tuma, D.J. (1987). The functional implications of acetaldehyde binding to cell constituents. Ann N Y Acad Sci *492*, 50-62.

Spanier, B., Wallwitz, J., Zapoglou, D., Idrissou, B.M.G., Fischer, C., Troll, M., Petzold, K., and Daniel, H. (2018). The Reproduction Rate of Peptide Transporter PEPT-1 Deficient C. elegans Is Dependent on Dietary Glutamate Supply. Front Mol Biosci *5*, 109.

Starich, T.A., Bai, X., and Greenstein, D. (2020). Gap junctions deliver malonyl-CoA from soma to germline to support embryogenesis in Caenorhabditis elegans. Elife *9*.

Thomason, L.C., Costantino, N., and Court, D.L. (2007). E. coli Genome Manipulation by P1 Transduction. Current Protocols in Molecular Biology *79*, 1.17.11-11.17.18.

Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J.*, et al.* (2020). SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods *17*, 261-272.

Vos, T., Allen, C., Arora, M., Barber, R.M., Bhutta, Z.A., Brown, A., Carter, A., Casey, D.C., Charlson, F.J., and Chen, A.Z. (2016). Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. The lancet *388*, 1545-1602.

White, E.C., Houlden, A., Bancroft, A.J., Hayes, K.S., Goldrick, M., Grencis, R.K., and Roberts, I.S. (2018). Manipulation of host and parasite microbiotas: Survival strategies during chronic nematode infection. Sci Adv *4*, eaap7399.

Yeates, T.O., Crowley, C.S., and Tanaka, S. (2010). Bacterial microcompartment organelles: protein shell structure and evolution. Annu Rev Biophys *39*, 185-205.

Yuen, L.H., Saxena, N.S., Park, H.S., Weinberg, K., and Kool, E.T. (2016). Dark Hydrazone Fluorescence Labeling Agents Enable Imaging of Cellular Aldehydic Load. ACS Chem Biol *11*, 2312-2319.