Identification of new *Dickeya dadantii* virulence factors secreted

by the type 2 secretion system.

Guy Condemine* and Bastien Le Derout

Univ Lyon, Université Lyon 1, INSA de Lyon, CNRS UMR 5240 Microbiologie Adaptation

et Pathogénie, F-69622 Villeurbanne, France.

* Corresponding author: guy.condemine@insa-lyon.fr

Short title: *Dickeya dadantii* virulence factors

ABSTRACT

Dickeya are plant pathogenic bacteria able to provoke disease on a wide range of plants. A type

2 secretion system named Out is necessary for bacterial virulence. Its study in *D. dadantii*

showed that it secretes a wide range of pectinolytic enzymes. However, the full repertoire of

exoproteins it can secrete has not been identified. No secretion signal present on the protein

allows the identification of substrates of a type 2 secretion system. To identify new Out

substrates, we analyzed *D. dadantii* transcriptome data obtained in plant infection condition and

searched for genes strongly induced encoding a protein with a signal sequence. We identified

four new Out-secreted proteins: the expansin YoaJ, VirK and two proteins of the DUF 4879

family, SvfA and SvfB. We showed that SvfA and SvfB are required for full virulence of *D.*

dadantii and showed that Svf proteins are present with a variable number of copies, up to three

in *D. fanghzongdai*, in other *Pectobacteriaceae*. This work opens the way to the study of the

role in virulence of non-pectinolytic proteins secreted by the Out pathway in *Pectobacteriaceae*.

IMPORTANCE

The plant pathogen *Dickeya* rely on a type 2 secretion system named Out for their pathogenicity.

Importance of plant cell wall degrading enzymes secreted by this system has been well studied.

However, existence and role of other Out-secreted proteins has barely been investigated. By

mining *D. dadantii* transcriptome data, we identified four new Out-secreted proteins. We

showed that two of them, SvfA and SvfB, are necessary for the full virulence of the bacteria.

These findings show that identification of all the proteins secreted by the *Dickeya* Out system

is necessary for a better knowledge of the virulence of these bacteria.

INTRODUCTION

Soft rot *Pectobacteriaceae*, *Dickeya* and *Pectobacterium*, are plant pathogenic bacteria that can provoke disease on a more than 35% of angiosperm plant orders, including both monocot and dicot plants (1). Among those, there is a wide range of plants of agronomic interest such as potato, rice, chicory, cabbage or ornementals on which they can cause severe losses. Symptoms are usually soft rot but these bacteria can provoke blackleg or wilting on aerial parts of potato. Recently, disease on woody plants caused by *Dickeya* have been reported (2). There is no efficient way to fight the disease. There are actually twelve species of *Dickeya* described, isolated either from infected plants (*D. chrysanthemi* from *Chrysanthemum morifolium*, *D. dadantii* subsp. *dadantii* from *Pelargonium capitum*, *D. dadantii* subsp. *diffenbachiae* from *Dieffenbachia* sp., *D. dianthicola* from *Dianthus caryophillus*, *D. zeae* from *Zea mays*, *D. oryzae* from *Oryza sativa*, *D. paradisiaca* from *Musa paradisiaca*, *D. solani* from *Solanum tuberosum*, *D. fangzhongdai* from *Pyrus pyrifolia*, *D. poaceiphila* from *Saccharum officinarum*) (3)(4)(5)(6)(7) or from river of lake water (*D. aquatica*, *D. lacustris* and *D. undicola*) (8)(9)(10). The role of protein secretion systems on the onset of the disease provoked by these bacteria has been recognized long ago (11). In contrast to many plant pathogenic bacteria, the type three Hrp secretion system is not the main determinant for *Pectobacteriaceae* virulence (12). The main virulence factor for these bacteria is a type 2 secretion system (T2SS) named Out. It allows the secretion of enzymes that degrade the components of the plant cell wall, leading to the soft rot symptom distinctive of the disease. The first secreted proteins to be identified were pectinases and a cellulase which are easily detectable by simple enzymatic tests (13) (11). The pectinolytic secretome of the model strain *D. dadantii* 3937 has been studied in detail by cloning the genes of these easily detectable enzymes. *D. dadantii* secretes nine pectate lyases, one pectin methylesterase, one pectin acetylesterase and one rhamnogalacturonate lyase (14). A proteomic analysis of the secreted proteins by 2D gel electrophoresis allowed the

identification of two other secreted proteins, the feruloyl esterase FaeD and a protein with

homology with a *Xanthomonas campestris* avirulence protein AvrL (15). A search in *D.*

dadantii of homologues of proteins secreted by *Pectobacterium atrosepticum* (16) led to the

characterization of the metal binding protein IbpS (17). Since all the pectinolytic enzymes

studied in *D. dadantii* are present in most of the *Dickeya* species they are probably not

responsible for the host preference observed for these bacteria (18). We hypothesized that

additional T2SS-secreted proteins specific for some species might exist and play a role in the

host preference. To identify such proteins, we analyzed *D. dadantii* transcriptome data, looking

for genes induced in plant infection conditions coding proteins with a signal sequence. We

identified several proteins secreted by the Out machinery and showed that two proteins of the

DUF4879 family, SvfA and SvfB are *D. dadantii* virulence factors.

RESULTS
81

Identification of new Out-secreted proteins

To have a more complete knowledge of the proteins secreted by the *D. dadantii* Out T2SS that

could be involved in the pathogenicity process, we searched for candidate genes in recently

published transcriptome data (19)(20). We selected genes strongly induced during plant

infection and coding for a protein possessing a signal sequence. We retained the genes

Dda3937_01687, *Dda3937_00585* (thereafter named SvfA and SvfB, respectively) and

Dda3937_00081 (also named *yoaJ*). We also retained VirK, a protein with a signal sequence

identified among the genes controlled by PecS, a regulator of many virulence factors (21). The

proteins were tagged with a His-tag and their secretion analyzed in the *D. dadantii* wild type

strain and an OutD mutant. Proteins SvfA, SvfB, YoaJ and VirK were found in the supernatant

of the wild type strain but not of the mutant, proving their secretion by the Out machinery (Fig.

1). Production of these proteins from a plasmid may explain why secretion was not total in the

wild type strain.

YoaJ is a PecS-regulated gene (21) and it was found among the most induced genes during

Arabidopsis infection or culture in the presence of plant extracts (19). It encodes a protein with

homology to expansins. These proteins are able to non-enzymatically loosen cell wall cellulose.

They are found in all plants where they have a role in cell wall extension and also in many plant

pathogenic microorganisms (22). Their role in virulence has been shown in *Ralstonia*

solanacearum, in *P*. *atrosepticum*, *P*. *Brasiliense* and in the plant pathogen *Erwinia*

tracheiphila (23) (24). The role of *D*. *dadantii* expansin is probably the same. VirK is a protein

of unknown function that has homologues in several plant pathogenic bacteria such as *R*.

solanacearum, *Agrobacterium tumefaciens*, *Lonsdalea* and *Xanthomonas*. VirK is controlled

by PecS and induced during Arabidopsis infection or culture in the presence of plant extracts

(21) (20). No symptom for the *D*. *dadantii virK* mutant was observed whatever the plant tested

(21). The two proteins SvfA and SvfB are studied in the next paragraphs.

SvfA and SvfB are virulence factors

svfA and *svfB* are among the most induced genes during *D*. *dadantii* Arabidopsis infection or

culture of the bacteria in the presence of plant extracts (19). They are also strongly expressed

during maceration of potato by *D*. *dianthicola* and *D*. *solani* (25). The two *D*. *dadantii* proteins

share 43% identity and 58% similarity in amino acid composition (Fig. 2). SvfA is 187 amino

acid long (165 for the mature form, 17.5 kDa) and SvfB is 198 amino acid long (177 for the

mature form, 18.9 kDa). These proteins belong to the DUF 4879 family of proteins. These

proteins have no known function. A protein of this family with low homology with SvfB, YolA,

is among the most highly secreted protein of *Bacillus subtilis* (26). YolA is also present in *B*.

cereus and in the insect pathogen *B. thuringiensis*. *Bacillus* YolA is shorter than SvfA and SvfB,

missing the N-terminal more variable part (Fig. 2).

SvfA and SvfB mutants have been constructed and their pathogenicity has been tested on

potato. The *svfA* mutant was significantly less aggressive than the wild type while the *svfB*

mutant was not affected (Fig. 3A). Virulence of the *svfA* mutant could be restored by

introduction of a plasmid bearing the wild type *svfA* (Fig. 3B). Virulence of the double *svfA*

svfB mutant was further reduced showing that the role of SvfB is additive to that of SvfA (Fig.

3A). Thus, genes *Dda3937_01687*and *Dda3937_00585* were named *svfA* and *svfB* for secreted

virulence factor A and B.

All attempts to overproduce produce SvfA and SvfB to purify them and study more precisely

their function were unsuccessful because their production was toxic to the bacteria that

overproduce them.

Expression of SvfA and SvfB

To try to identify the function of SvfA and SvfB we analyzed the conditions in which their

genes are expressed. We tested the effect of galacturonate and polygalacturonate, two

compounds that are inducers of the expression of the main virulence factors, the pectate lyases,

and of glucose, which represses it. We also analyzed the effect of mutation in genes controlling

several aspects of *D. dadantii* virulence. KdgR represses the pectinase, pectin degradation and

the *out* genes (27). Its inducer is 2-keto-3-deoxygluconate, a polygalacturonate and

galacturonate catabolic derivative. PecS controls the pectinase, other secreted protein, the *out*

genes and those involved in resistance to oxidative stress (28). PecT is a regulator of the pectate

lyase, motility and exopolysaccharide synthesis genes (29). Pir regulates hyperinduction of

pectate lyses in response to plant signal (30). Finally, GacA, the regulator of the two-component

regulation system GacA-GacS, is a global regulator required for disease expression in response

to the metabolic status of the bacteria (31). Expression of *svfA* was slightly induced by

polygalacturonate but not by galacturonate (Fig. 4A). However, expression of the gene was not

modified in a *kdgR* background indicating that induction by polygalacturonate is not mediated

by *kdgR*. Growth in the presence of chicory chunks strongly induced s*vfA* expression as

expected from transcriptomic data showing induction in the presence of plant extract. A high

concentration of glucose led to a strong induction of the gene expression (Fig. 4A). This

regulation is mediated by the catabolite repressor protein CRP since a mutation in the *crp* gene

derepressed *svfA* expression. Thus, Crp is a repressor of *svfA*. Although it had not been detected

as a PecS-regulated gene (21), its expression is elevated in a *pecS* background. A *pir* mutation

derepressed also a little *svfA* expression. Neither PecT nor GacA significantly regulate *svfA*

expression (Fig. 4A).

Regulation of *svfB* is quite similar to that of *svfA*: it was not induced by galacturonate,

polygalacturonate or regulated by KdgR, it was induced by glucose and repressed by Crp, and

it was repressed by PecS (Fig. 4B). However, few differences can be noted: in contrast to what

is observed with *svfA*, no induction by plant pieces was observed and PecT was a repressor of

svfB expression while pir did not seem to control it (Fig. 4B).

Occurrence of new secreted proteins in other Dickeya species

Presence of *svfA*, *svfB*, *virK* and *yoaJ* was searched in all the *Dickeya* type strains, and in a few

Pectobacterium strains (Table 1). Presence and number of proteins of the DUF 4879 family is

variable among *Dickeya* species. *svfA* is present in all strains except *D. zeae*, *D. chrysanthemi*,

D. poaceiphila and *D. paradisiaca*. *svfB* is absent in *D. chrysanthemi*, *D. poaceiphila* and *D.*

paradisiaca, *D. undicola* and *D. aquatica*. A third gene located next to *svfA* and resulting

probably from a duplication is found in *D. fangzhongdai* and *D. undicola*. It has 53% homology

with *D. dadantii* SvfA and 34% with *D. dadantii* SvfB. It was named *svfC*. SvfC is shorter that

SvfA and SvfB (126 amino acid for the mature protein, 13.2 kDa) and has the same size as *B.*

cereus YolA (Fig. 2). It possesses a signal sequence, indicating that it could also be secreted by

the Out system. Thus, the number of genes of this family in *Dickeya* strains varies from 0 to 3.

Homologues of the *svf* genes can be found in some *Pectobacterium* strains (Table 1). For

example, two copies are present in *P. carotovorum* subsp *carotovorum*. However, even in a

given species the gene may be present or not (presence of a homologue of *svfB* in 10 out of the

23   *P.*   *brasiliense*   strains   present   in   the   ASAP   data   bank

(https://asap.ahabs.wisc.edu/asap/home.php). Outside *Pectobacteriaceae*, homologues of *svfB*

can be found in a few gammaproteobacteriaceae, i.d. in some *Photorhabdus*, *Luteibacter* and

Pseudoalteromonas strains. *yoaJ* is present in all *Dickeya* and *Pectobacterium* strains except

D. poaceiphila and *D. paradisiaca*. *virK* is absent in *D. aquatica* and in all *Pectobacterium*

strains.

We examined also the presence or absence of genes of other non-pectinolytic proteins known

to be secreted by a T2SS in *Dickeya* or *Pectobacterium*: *ibpS*, *nipE*, *xynA*, *avrL/avrM* (Table

1). IbpS is a metal binding protein that prevents ROS-induced killing of bacteria (17). NipE is

a toxin that provoke plant cell death (32). XynA is a xylanase that was identified in a corn strain

of *Erwinia chrysanthemi* (33). AvrL is a protein with homology with a *Xanthomonas campestris*

avirulence protein AvrL (15). Two very similar proteins, AvrL and AvrM, are encoded by *D.*

dadantii 3937. AvrL was named Svx in *P. atrosepticum* where a role in virulence has been

proven (34). However, its function in *Dickeya* has not been studied. While IbpS and NipE are

present in almost all the species, except *D. paradisiaca* and *D. poaceiphila* for NipE, a variation

in the presence and number of XynA and AvrL was observed (Table 1). Thus, the repertoire of

T2SS-secreted protein known to be important for virulence is very variable from species to

species.

199
DISCUSSION

The T2SS of *Dickeya* and *Pectobacterium* is the main virulence factor of these bacteria. The

knowledge of the repertoire of secreted proteins is necessary to better understand the

mechanisms of virulence of these bacteria. This analysis has been undertaken with the model

strain *D. dadantii* 3937 and partially with *Pectobacterium atrosepticum* (35)(16). *Dickeya* and

Pectobacterium are characterized by their ability to degrade pectin and new species are

identified by this characteristic on the semi selective Crystal Violet Pectate medium. They all

secrete enzymes capable of degrading this polysaccharide (pectate lyases, polygalacturonases,

pectin methylesterases) which are produced by all these bacteria. However, recent work shows

that other proteins are secreted by the Out T2SS (15)(16). In this work we used published

transcriptome data to identify new substrates of the *Dickeya dadantii* T2SS. The most induced

genes in a transcriptome experiment of *D. dadantii* infecting *A. thaliana* are known virulence

genes (*pelI, prtA, rhiE, paeY, rhaD, ibpS*, etc...) (19). However, in this top list some genes have

no known function. The presence of a signal sequence in their product suggested that these

proteins could be substrates of the T2SS necessary for the infection process. We showed here

that the proteins SvfA, SvfB and YoaJ present in the top list are substrates of the Out T2SS.

YoaJ belongs to the family of expansins, proteins that loosen cellulose fibers. Their role as a

virulence factor has been shown in *P. brasiliense* and *P. atrosepticum* (23). It probably has the

same function in *D. dadantii*. No function could be predicted for SvfA and SvfB which belong

to the DUF 4879 family of proteins. However, a reduction of virulence of a *svfA* mutant and a

svfA svfB double mutant on potato could be observed proving a role of the proteins in the

bacterial pathogenicity. Although an additive effect of the mutations was observed, they could

not have exactly the same function. The mutants should be tested on various hosts to detect

possible differences. It can be supposed that each protein would be more active on one type or

one family of plant. Presence of three proteins of this family in *D. fanghzongdai* could explain

its wide host range. Presence of homologues of SvfA and SvfB in *Photorhabdus* and in *B.*

thuringiensis strains, two insect pathogens, indicates that the role of these proteins is not

restricted to plant virulence but may participate to a common process of bacterial pathogeny

such as cell lysis. We showed also here that the PecS-regulated protein VirK is also secreted by

Out. No role on virulence had been found for this protein with the chicory leaf model of

infection (21). Other models should be tested to find the role of this protein.

Regulation of expression of the *svfA* and *svfB* genes is atypical for a *D. dadantii* gene involved

in pathogeny. While expression of most of the virulence factors is induced in the presence of

pectin or its derivatives through the repressor KdgR and repressed by glucose, that of *svfA* and

svfB is opposite: it is activated by glucose and not controlled by KdgR. Expression of *svfA* is

induced in the presence of plant tissue. This pattern of regulation has been described for *ibpS*,

which is also strongly induced in *A. thaliana* (17). This could correspond to conditions

encountered during the early phases of infection: pectin has not yet been degraded and glucose

and saccharose are plentiful in plant tissues. *svfA* and *ibpS* could be among the earliest gene to

be induced at the onset of infection, before the genes involved in pectin degradation. However,

regulation of these genes by PecS and PecT shows that *svfA* and *svfB* are fully integrated in the

network of regulators that controls *D. dadantii* virulence.

This work has extended our knowledge of the Out-dependent secretome of *D. dadantii*, showing

that besides pectinases several other proteins are secreted. If the number of pectinolytic

enzymes secreted is almost identical in the various *Dickeya* species, the number of additional

secreted proteins varies markedly. Among the proteins we analyzed (Table 1), *D. paradisiaca*

has only one (VirK) while *D. fanghzongdai* has ten. All the intermediate combinations can be

found in the various species. There seems to be less variations in the *Pectobacterium* strains

that we surveyed. It is tempting to speculate that the presence/absence of these proteins could

be one of the origin of the host preference of the *Dickeya* strains, providing additional virulence

factors necessary to infect certain hosts. Articles that compare *Dickeya* strains to understand

what makes their difference in their host range or aggressivity often focus only on the presence

of the six known types of secretion systems without analyzing what proteins could be secreted

(18)(36)(37). An exhaustive analysis of the secreted proteins would be more informative.

Is there other T2SS-secreted proteins to be identified in *Dickeya* strains? No specific signal is

present on T2SS-secreted proteins that would allow their identification. 2D gels which were

used in previous studies performed on *D. dadantii* and *P. atrosepticum* to identify these proteins

have a limited sensitivity (15)(34). More sensitive methods such as liquid chromatography-

tandem mass spectrometry (LC-MS/MS) can now be used (38). However, they give many false

positive results since periplasmic and cytoplasmic proteins are often found in the culture

supernatant. The approach we used here allowed the identification of four new secreted

proteins. However, all these methods have a drawback. They can only detect proteins that are

produced. The rhamnogalacturonate lyase RhiE could only be detected when the bacteria were

cultivated in the presence of rhamnose (39). YoaJ and VirK were not induced in *D. dianthicola*

grown on potato (25). Another problem is that a protein may not exist in the strain tested. An

analysis of the secretome of several *Dickeya* strains grown in several conditions will be

necessary to have a global view of all the virulence factors that can be secreted by *Dickeya*

species and evaluate their potential role in pathogenicity.

MATERIAL AND METHODS
273

Bacterial strains and growth conditions.

Bacterial strains, phages, plasmids and oligonucleotides used in this study are described in Table 2. *D.*

dadantii and *E. coli* cells were grown at 30 and 37°C respectively in LB medium or M63 minimal medium

supplemented with a carbon source (0.2%, w/v unless otherwise indicated). When required antibiotics

were added at the following concentrations: ampicillin, 100 mg l^{-1}, kanamycin and chloramphenicol, 25

mg l^{-1}. Media were solidified with 1.5% (w/v) agar. Transduction with phage ΦEC2 was performed

according to Résibois *et al*. (40)

Mutant construction.

To construct strain A6418 that contains a *svfA-uidA* fusion a 1.3 kb DNA fragment containing

svfA was amplified with primers 17176H+ and 17176A. The resulting fragment was inserted into

the pGEM-T plasmid (Promega). A XbaI site was created by site directed mutagenesis with the

primers 17176XbaF and 17176XbaR into the *svfA* coding sequence and a *uidA*-kanR cassette was

inserted into this XbaI site. To construct strain A6467 that contains a *svfB-uidA*-kanR fusion and

a CmR cassette a 2000bp DNA fragment containing *svfB* was amplified with the primers

15544L2+ and 15544L2-. The resulting fragment was inserted into the pGEM-T plasmid. A XmaI

site was created by site directed mutagenesis into *svfB* coding sequence with the primers

15544XmaF and 15544XmaR and a *uidA*-kanR cassette was inserted into this created XmaI site

unique. All the constructs were recombined into the *D*. *dadantii* chromosome according to Roeder

and Collmer (41). Recombinations were checked by PCR. His-tagged versions of the proteins

SvfA, SvfB, YolA, VirK were constructed by amplifying the corresponding genes with the primers

17176H+ and 17176H-, 15544H+ and 15544H-, 14642H+ and 14642H-, VirKH+ and VirKH-,

respectively. The resulting DNA fragments were cloned into plasmid pGEMT.

Secretion assays and Western blots.

D. dadantii strains containing the plasmid to test were grown overnight in LB medium in the

presence of the appropriate antibiotic. Culture supernatant containing secreted proteins was

separated from cells by centrifugation at 10,000g for 3 min, and both fractions were loaded onto

12% polyacrylamide gel electrophoresis (SDS-PAGE). The proteins were next transferred onto

Immobilon P membrane (Merck) and probed with Ni-NTA-HRP.

Pathogenicity tests.

Bacteria were grown overnight in LB medium, centrifuged and resuspended at OD_{600} 1 in M63

medium. Potatoes were surface sterilized with 70% ethanol and dried. A hole was made with a

pipette tip and 10μl of bacteria were deposited in the hole which was covered with mineral oil.

Potatoes were placed over a wet paper in a tray contained in a plastic bag to maintain moisture.

After 48h at 30°C, the weight of rotten tissue was measured.

Enzymatic assays.

β-glucuronidase assays were performed on toluenized extracts of cells grown to exponential

phase using the method of Bardonnet *et al* (42). with *p*-nitrophenyl-β-D-glucuronate as the

substrate.

ACKNOWLEDGMENTS

We thank Lison Massardier and Florence Ruaudel for technical work and Nicole Cotte-Pattat

for reading the manuscript. This work was supported by funding from CNRS, University Lyon

1 and Agence Nationale de la Recherche (ANR-19-CE35-0016).

REFERENCES

1. Ma B, Hibbing ME, Kim H-S, Reedy RM, Yedidia I, Breuer J, Breuer J, Glasner JD, Perna

NT, Kelman A, Charkowski AO. 2007. Host range and molecular phylogenies of the soft

rot enterobacterial genera *Pectobacterium* and *Dickeya*. *Phytopathology* 97:1150–1163.

2. Fujikawa T, Hatomi H, Ota N. 2020. Draft genome sequences of seven strains of *Dickeya*

dadantii, a quick decline-causing pathogen in fruit trees, isolated in Japan. *Microbiol*

Resour Announc 9:e00609-20.

3. Samson R, Legendre JB, Christen R, Saux MF-L, Achouak W, Gardan L. 2005. Transfer

of *Pectobacterium chrysanthemi* (Burkholder et al. 1953) Brenner *et al*. 1973 and

Brenneria paradisiaca to the genus *Dickeya* gen. nov. as *Dickeya chrysanthemi* comb.

nov. and *Dickeya paradisiaca* comb. nov. and delineation of four novel species, *Dickeya*

dadantii sp. nov., *Dickeya dianthicola* sp. nov., *Dickeya dieffenbachiae* sp. nov. and

Dickeya zeae sp. nov. *Int J Syst Evol Microbiol* 55:1415–1427.

4. Brady CL, Cleenwerck I, Denman S, Venter SN, Rodríguez-Palenzuela P, Coutinho TA,

De Vos P. 2012. Proposal to reclassify *Brenneria quercina* (Hildebrand and Schroth 1967)

Hauben *et al*. 1999 into a new genus, *Lonsdalea* gen. nov., as *Lonsdalea quercina* comb.

nov., descriptions of *Lonsdalea quercina* subsp. *quercina* comb. nov., *Lonsdalea quercina*

subsp. *iberica* subsp. nov. and *Lonsdalea quercina* subsp. *britannica* subsp. nov.,

emendation of the description of the genus *Brenneria*, reclassification of *Dickeya*

dieffenbachiae as *Dickeya dadantii* subsp. *dieffenbachiae* comb. nov., and emendation of

the description of *Dickeya dadantii*. *Int J Syst Evol Microbiol* 62:1592–1602.

5.   van der Wolf JM, Nijhuis EH, Kowalewska MJ, Saddler GS, Parkinson N, Elphinstone

JG, Pritchard L, Toth IK, Lojkowska E, Potrykus M, Waleron M, de Vos P, Cleenwerck

I, Pirhonen M, Garlant L, Hélias V, Pothier JF, Pflüger V, Duffy B, Tsror L, Manulis S.

2014. *Dickeya solani* sp. nov., a pectinolytic plant-pathogenic bacterium isolated from

potato *(Solanum tuberosum)*. *Int J Syst Evol Microbiol* 64:768–774.

6.   Tian Y, Zhao Y, Yuan X, Yi J, Fan J, Xu Z, Hu B, De Boer SH, Li X. 2016. *Dickeya*

fangzhongdai sp. nov., a plant-pathogenic bacterium isolated from pear trees (*Pyrus*

pyrifolia). *Int J Syst Evol Microbiol* 66:2831–2835.

7.   Wang X, He S-W, Guo H-B, Han J-G, Thin KK, Gao J, Wang Y, Zhang X-X. 2020.

Dickeya oryzae sp. nov., isolated from the roots of rice. *Int J Syst Evol Microbiol* 70:4171–

4178.

8.   Parkinson N, DeVos P, Pirhonen M, Elphinstone J. 2014. *Dickeya aquatica* sp. nov.,

isolated from waterways. *Int J Syst Evol Microbiol* 64:2264–2266.

9.   Hugouvieux-Cotte-Pattat N, Jacot-des-Combes C, Briolay J. 2019. *Dickeya lacustris* sp.

nov., a water-living pectinolytic bacterium isolated from lakes in France. *Int J Syst Evol*

Microbiol 69:721–726.

10.   Oulghazi S, Pédron J, Cigna J, Lau YY, Moumni M, Van Gijsegem F, Chan K-G, Faure

D. 2019. *Dickeya undicola* sp. nov., a novel species for pectinolytic isolates from surface

waters in Europe and Asia. *Int J Syst Evol Microbiol* 69:2440–2444.

11.   Ji J, Hugouvieux-Cotte-Pattat N, Robert-Baudouy J. 1989. Molecular cloning of the *outJ*

gene involved in pectate lyase secretion by *Erwinia chrysanthemi*. *Mol Microbiol* 3:285–

293.

12.   Yang C-H, Gavilanes-Ruiz M, Okinaka Y, Vedel R, Berthuy I, Boccara M, Chen JW-T,

Perna NT, Keen NT. 2002. *hrp* genes of *Erwinia chrysanthemi* 3937 are important

virulence factors. *Mol Plant-Microbe Interact* 15:472–480.

13.   Andro T, Chambost JP, Kotoujansky A, Cattaneo J, Bertheau Y, Barras F, Van Gijsegem

F, Coleno A. 1984. Mutants of *Erwinia chrysanthemi* defective in secretion of pectinase

and cellulase. *J Bacteriol* 160:1199–1203.

14.   Hugouvieux-Cotte-Pattat N, Condemine G, Shevchik VE. 2014. Bacterial pectate lyases,

structural and functional diversity. *Environ Microbiol Rep* 6:427–440.

15.   Kazemi-Pour N, Condemine G, Hugouvieux-Cotte-Pattat N. 2004. The secretome of the

plant pathogenic bacterium *Erwinia chrysanthemi*. *Proteomics* 4:3177–3186.

16.   Coulthurst SJ, Lilley KS, Hedley PE, Liu H, Toth IK, Salmond GPC. 2008. DsbA plays a

critical and multifaceted role in the production of secreted virulence factors by the

phytopathogen *Erwinia carotovora* subsp. *atroseptica*. *J Biol Chem* 283:23739–23753.

17.   Liu L, Gueguen-Chaignon V, Gonçalves IR, Rascle C, Rigault M, Dellagi A, Loisel E,

Poussereau N, Rodrigue A, Terradot L, Condemine G. 2019. A secreted metal-binding

protein protects necrotrophic phytopathogens from reactive oxygen species. *Nat Commun*

10:4853.

18.   Potrykus M, Golanowska M, Hugouvieux-Cotte-Pattat N, Lojkowska E. 2014. Regulators

involved in *Dickeya solani* virulence, genetic conservation, and functional variability. *Mol*

Plant-Microbe Interact 27:700–711.

19.   Pédron J, Chapelle E, Alunni B, Van Gijsegem F. 2018. Transcriptome analysis of the

Dickeya dadantii PecS regulon during the early stages of interaction with *Arabidopsis*

thaliana. *Mol Plant Pathol* 19:647–663.

20.   Jiang X, Zghidi-Abouzid O, Oger-Desfeux C, Hommais F, Greliche N, Muskhelishvili G,

Nasser W, Reverchon S. 2016. Global transcriptional response of *Dickeya dadantii* to

environmental stimuli relevant to the plant infection. *Environ Microbiol* 18:3651–3672.

21.   Hommais F, Oger-Desfeux C, Van Gijsegem F, Castang S, Ligori S, Expert D, Nasser W,

Reverchon S. 2008. PecS Is a global regulator of the symptomatic phase in the

phytopathogenic bacterium *Erwinia chrysanthemi* 3937. *J Bacteriol* 190:7508–7522.

22.   Tancos MA, Lowe-Power TM, Peritore-Galve FC, Tran TM, Allen C, Smart CD. 2018.

Plant-like bacterial expansins play contrasting roles in two tomato vascular pathogens.

Mol Plant Pathol 19:1210–1221.

23.   Narváez-Barragán DA, Tovar-Herrera OE, Torres M, Rodríguez M, Humphris S, Toth IK,

Segovia L, Serrano M, Martínez-Anaya C. 2020. Expansin-like Exl1 from *Pectobacterium*

is a virulence factor required for host infection, and induces a defence plant response

involving ROS, and jasmonate, ethylene and salicylic acid signalling pathways in

Arabidopsis thaliana. *Sci Rep* 10:7747.

24.   Rocha J, Shapiro LR, Kolter R. 2020. A horizontally acquired expansin gene increases

virulence of the emerging plant pathogen *Erwinia tracheiphila*. *Sci Rep* 10:21743.

25. Raoul des Essarts Y, Pédron J, Blin P, Van Dijk E, Faure D, Van Gijsegem F. 2019.

Common and distinctive adaptive traits expressed in *Dickeya dianthicola* and *Dickeya*

solani pathogens when exploiting potato plant host. *Environ Microbiol* 21:1004–1018.

26. Tjalsma H, Antelmann H, Jongbloed JDH, Braun PG, Darmon E, Dorenbos R, Dubois J-

YF, Westers H, Zanen G, Quax WJ, Kuipers OP, Bron S, Hecker M, van Dijl JM. 2004.

Proteomics of protein secretion by *Bacillus subtilis* : Separating the "secrets" of the

secretome. *Microbiol Mol Biol Rev* 68:207–233.

27. Condemine G, Robert-Baudouy J. 1987. Tn *5* insertion in *kdgR*, a regulatory gene of the

polygalacturonate pathway in *Erwinia chrysanthemi*. *FEMS Microbiol L*ett 42:39–46.

28. Reverchon S, Nasser W, Robert-Baudouy J. 1994. *pecS*: a locus controlling pectinase,

cellulase and blue pigment production in *Erwinia chrysanthemi*. *Mol Microbiol* 11:1127–

1139.

29. Surgey N, Robert-Baudouy J, Condemine G. 1996. The *Erwinia chrysanthemi pecT* gene

regulates pectinase gene expression. *J Bacteriol* 178:1593–1599.

30. Nomura K, Nasser W, Kawagishi H, Tsuyumu S. 1998. The *pir* gene of *Erwinia*

chrysanthemi EC16 regulates hyperinduction of pectate lyase virulence genes in response

to plant signals. *Proc Natl Acad Sci* 95:14034–14039.

31. Lebeau A, Reverchon S, Gaubert S, Kraepiel Y, Simond-Côte E, Nasser W, Van Gijsegem

F. 2008. The GacA global regulator is required for the appropriate expression of *Erwinia*

chrysanthemi 3937 pathogenicity genes during plant infection. *Environ Microbiol*

10:545–559.

32.   Laasik E, Põllumaa L, Pasanen M, Mattinen L, Pirhonen M, Mäe A. 2014. Expression of

nipP.w of *Pectobacterium wasabiae* is dependent on functional *flgKL* flagellar genes.

Microbiology 160:179–186.

33.   Keen NT, Boyd C, Henrissat B. 1996. Cloning and characterization of a xylanase gene

from corn strains of *Erwinia chrysanthemi*. *Mol Plant Microbe Interact* 9:651-657.

34.   Corbett M, Virtue S, Bell K, Birch P, Burr T, Hyman L, Lilley K, Poock S, Toth I,

Salmond G. 2005. Identification of a new quorum-sensing-controlled virulence factor in

Erwinia carotovora subsp. *atroseptica* secreted via the type II targeting pathway. *Mol*

Plant-Microbe Interact 18:334–342.

35.   Mattinen L, Nissinen R, Riipi T, Kalkkinen N, Pirhonen M. 2007. Host-extract induced

changes in the secretome of the plant pathogenic bacterium *Pectobacterium atrosepticum*.

Proteomics 7:3527–3537.

36.   Alič Š, Pédron J, Dreo T, Van Gijsegem F. 2019. Genomic characterisation of the new

Dickeya fangzhongdai species regrouping plant pathogens and environmental isolates.

BMC Genomics 20:34.

37.   Xu P, Wang H, Qin C, Li Z, Lin C, Liu W, Miao W. 2021. Analysis of the taxonomy and

pathogenic factors of *Pectobacterium aroidearum* L6 using whole-genome sequencing

and comparative genomics. *a*12:679102.

38.   Burtnick MN, Brett PJ, DeShazer D. 2014. Proteomic Analysis of the *Burkholderia*

pseudomallei type II secretome reveals hydrolytic enzymes, novel proteins, and the

deubiquitinase TssM. *Infect Immun* 82:3214–3226.

39. Laatu M, Condemine G. 2003. Rhamnogalacturonate lyase RhiE is secreted by the Out

system in *Erwinia chrysanthemi*. *J Bacteriol* 185:1642–1649.

40. Resibois A, Colet M, Faelen M, Schoonejans E, Toussaint A. 1984. φEC2, a new

generalized transducing phage of *Erwinia chrysanthemi*. Virology 137:102–112.

41. Roeder DL, Collmer A. 1985. Marker-exchange mutagenesis of a pectate lyase isozyme

gene in *Erwinia chrysanthemi*. *J Bacteriol* 164:51–56.

42. Bardonnet N. 1992. ′uidA-antibiotic-resistance cassettes for insertion mutagenesis, gene

fusions and genetic constructions. *FEMS Microbiol Lett* 93:243–247.

Fig 1: Identification of new secreted proteins. Wild-type and *outD* mutant strains containing

plasmid bearing the gene of the protein to test were grown overnight in LB medium. The

supernatant (S) and cellular (C) fractions were separated by SDS-PAGE. After blotting, the

proteins were detected with Ni-NTA-HRP.

Fig. 2: Alignment of Svf proteins. The sequences of *D*. *dadantii* SvfA (Dda3937_01687), SvfB

(Dda3937_00585) and *D*. *fanghzongdai* SvfC (CVE23_15565), *B*. *cereus* WP-193674364.1

and *Photorhabdus asymbiotica* CAQ86327.1 without their signal sequence were aligned with

Clustal omega. Identical residues are indicated by a star and chemically equivalent residues by

a double dot

Fig. 3. Virulence of *svfA* and *svfB* mutants. A. Potatoes were infected with the wild type, and

the *svfA*, the *svfB* and the *svfA svfB* mutants. Rotten tissue was weighed after 48 h. B.

Complementation of the *svfA* mutation. Potatoes were infected with the wild type strain and the

svfA mutant containing the empty plasmid pBBRGm and the *svfA* mutant containing the pBBR

plasmid bearing *svfA*. Rotten tissue was weighed after 48h. Statistical tests were performed

using the Wilcoxon- Mann-Whitney test. The *p*-value were compared with an alpha risk of 4%.

$p < 0.001 = ***, p < 0.005 = **, p < 0.01 = *$.

Fig 4 : Expression of *svfA and svfB* in various growth conditions. A. The *D*. *dadantii* strain

A6418 containing the *svfA-uidA* fusion and derivative strains containing an additional

regulatory mutation were grown in M63 medium in the presence of the indicated compounds

(Y = glycerol, G = glucose, A = galacturonate, PGA = polygalacturonate, E = chicory chunks).

Strains with additional mutations were grown with glycerol as a carbon source except the *crp*

mutant that was grown with 0.2% glucose. ß-glucuronidase activity was measured with *p*-

nitrophenyl-ß-D-glucuronate. B. The *D. dadantii* strain A6467 containing the *svfB-uidA* fusion

and derivative strains containing an additional regulatory mutation were grown in M63 medium

in the presence of the indicated compounds (Y = glycerol, G = glucose, A = galacturonate, PGA

= polygalacturonate, E = chicory chunks). Strains with additional mutations were grown with

glycerol as a carbon source except the *crp* mutant that was grown with 0.2% glucose ß-

glucuronidase activity was measured with *p*-nitrophenyl-ß-D-glucuronate. Activities are

expressed in µmoles of *p*-nitrophenol produced per minute and per milligram of bacterial dry

weight $\pm$ standard deviation. Data are expressed as the mean (n = 6) from six independent

experiments. Statistical tests were performed using the Wilcoxon- Mann-Whitney test. The *p*-

value were compared with an alpha risk of 4%. $p < 0.001$=***,

$p < 0.005$=**, $p < 0.01$=*.

Table 1: Presence of Out-secreted proteins in various *Dickeya* and *Pectobacterium* strains.

Table 2: Oligonucleotides and strains used in this study.

.

Table 1 : Presence of Out-secreted proteins in various *Dickeya* and *Pectobacterium* strains.

Strain	SvfA	SvfB	SvfC	YoaJ	VirK	IbpS	NipE	XynA	AvrL
D. dadantii	1	1	0	1	1	1	1	1	2
D. diffenbachiae	1	1	0	1	1	1	1	0	2
D. fangzhongdai	1	1	1	1	1	1	1	1	2
D. solani	1	1	0	1	1	1	1	1	2
D. zeae	0	1	0	1	1	1	1	1	1
D. oryzae	1	1	0	1	1	1	1	1	0
D. dianthicola	0	1	0	1	1	1	1	0	1
D. undicola	1	0	1	1	1	1	1	0	1
D. lacustris	1	0	0	0	0	1	1	0	2
D. aquatica	1	0	0	1	0	1	1	0	2
D. chrysanthemi	0	0	0	1	1	1	1	0	1
D. paradisiaca	0	0	0	0	1	0	0	0	0
D. poaceiphila	0	0	0	0	1	1	0	1	0
P. atrosepticum	0	0	0	1	0	1	1	0	1
P. carotovorum	1	1	0	1	0	1	1	0	1
P. parmentieri	0	0	0	1	0	1	1	0	1
P. polaris	0	0	0	1	0	1	1	0	1

The strains used in this study are *D. dadantii* 3937, *D. aquatica* 174/2, *D. chrysanthemi*

ATCC 11663, *D. dadantii* subsp *dieffenbachiae* NCPPB 2976, *D. dianthicola* NCPPB 453, *D.*

fanghzongdai DSM 101947, *D. lacustris* S29, *D. oryzae* ZYY5, *D. paradisiaca* ATCC 33242,

D. poaceiphila NCPPB 569, *D. solani* IPO 2222, *D. undicola* 2B12, *D. zeae* NCPPB 2538, *P.*

atrosepticum ATCC 33260, *P. carotovorum* subsp. *carotovorum* ATCC 15713, *P.*

parmentieri RNS08.42.1A and *P. polaris* NIBIO 1006. The presence and number of proteins

detected in each strain is indicated.

517   Table 2: Oligonucleotides and strains used in this study

519   Oligonucleotides

Line	Name	Sequence
520	17176H+	CCTCCTGAGATTAGAGAGAG
521	17176A	CGCGCCGGTGTTTTTCTTGCG
522	17176H-	GGTTAGTGATGGTGATGGTGATGCTGAATATTGAGCGACGTGC
523	17176XbaF	GAACAGGATGTGTCCTCTTCTAGACACAAAGCGCTGCGTGCG
524	17176XbaR	CGCACGCAGCGCTTTGTGTCTAGAAGAGGACACATCCTGTTC
525	15544L2+	CTAAGAATCAGTCAGTTTGCG
526	15544L2-	AGGCAGATAACGCTACTCGCC
527	15544H+	ATTTATACTCGCCACCGATGC
528	15544XmaF	GCCTTGCGAGCCCCGCTCCCGGGCTGTCTCGGGTTACCGTG
529	15544XmaR	CACGGTAACCCGAGACAGCCCGGGAGCGGGGGCTCGCAAGGC
530	15544H-	GGTTAGTGATGGTGATGGTGATGTTTAATATAGGTCTGTGTGAAC
531	14642H+	GGTGAGGAATAATTCTGGCC
532	14642H-	GGTTAGTGATGGTGATGGTGATGAGGCAGTTGTACTTTTCCAG
533	VirKH+	TGCCGTATGTGATAGTCACG
534	VirKH-	CCTTAGTGATGGTGATGGTGATGTTGCTTGAAGAAGCGGATATC

535
536
537   *Dickeya dadantii* strains
538

Line	Strain	Genotype	Source
539	3937	Wild type	Laboratory collection
540	A5653	*outD* ::CmR	Laboratory collection
541	A6417	*svfB* ::CmR	This work
542	A6418	*svfA* ::*uidA*-kanR	This work
543	A3838	*kdgR* ::Mu-CmR	(19)
544	A3845	*pecS*::Mu-CmR	(28)
545	A3846	*pecT*::CmR	(29)
546	A3849	*pir*::CmR	Laboratory collection
547	A4237	*gacA*::CmR	(31)
548	A6434	*kdgR*::Mu-CmR *svfA*::*uidA*-kanR	This work
549	A6435	*pecS*::Mu-CmR *svfA* :*uidA*-kanR	This work
550	A6436	*pecT*::CmR *svfA*::*uidA*-kanR	This work
551	A6436	*pir*::CmR *svfA*::*uidA*-kanR	This work

552	A6437	*gac*A::CmR *svfA*::*uidA*-kanR	This work
553	A6467	*svfB*::*uidA*-kanR	This work
554	A6469	*kdgR*::Mu-CmR *svfB*::*uidA*-kanR	This work
555	A6470	*pecS*::Mu-CmR *svfB*::*uidA*-kanR	This work
556	A6471	*pecT*::CmR *svfB*::*uidA*-kanR	This work
557	A6472	*pir*::CmR *svfB*::*uidA*-kanR	This work
558	A6473	*gacA*::CmR *svfB*::*uidA*-kanR	This work
559			
560			
561			
562			
563			

571   Fig. 1

```
SvfA          -------AQTS----DEPVQTVITALD--SPFVDYPLSAGSEQDVSSSEHKALRAPAPAL 47
SvfC          ----------------------------------------------------APAPAL 6
Bacillus      ----------------------------------------------------APAPPL 6
SvfB          GYSQTINNTDEAGPPIELG---LTAFD-SL-ESNSPLSAARDTS-RASTSGALRAPAPAL 54
Photorhabdus  --------QTEQLPKVPAAQKVLIEQSQLLPNVKSPIR-AEERD-LKIEDGSVHAPAPGL 50
                                                                  ****  *

SvfA          SSVQVYAVYSSLKGGWQAVPT-NTLSL-SGYAGGTLRIAVLEVGYGGNRIGWLNGGQTS- 104
SvfC          SGLRIKVLSGVYGGTWQYAPV-NAVSIGPGYAGGTLQIAVVETGYGGNRIGWINGEQKK- 64
Bacillus      TSLNVVKVESQL-GGVEFIGA-NNLSTVKDHGGSYLYIYTNEMGYGRNPIAQMSGQKLKK 64
SvfB          SRVTVYAVGSSN-CGWEYMTSIGQLSTTCDHGGAQLRVAVQEIGYGNNPVAWMNGGVLPR 113
Photorhabdus  TNMWVYAVGSTN-CGWEYTSNL--FATTCDHGGQQLRAAVLEIGYGYSSFAWMNGGLLPN 107
              : : :  : .        :          .:   .:.* *    . * *** . .. :.*

SvfA          --PYQVNPVCVVSGRYTESCPAGSIVSGWMAYFNADNMSSVTFRYQSTSTNFPNRTLSTS 162
SvfC          --PDSVKLACLVKGEMTDNCPRGATGAGWIAYFSANYQTSVTFRYQSTSANFPYKTLSTS 122
Bacillus      VDS---KMI---------DINGDRTVDGWEYKWDASGQQNGQFKYQNTSTNAPWNTLFTS 112
SvfB          SANYQTDGICIVGNQYTFPCPAGYTVVGWMYYYNLDGTDNGQFKFQDTSTNAPRNTLFTQ 173
Photorhabdus  SAMYSSKTVCITNGYYTWPCTAGQTVVGYLHEYNLDGNQNGTFRYQNTSTNSPWNTMSVQ 167
                            .         *:  :. .  . *::*.**:* * .*: ..

SvfA          LNIQ 166
SvfC          LTIK 126
Bacillus      LNIK 116
SvfB          TYIK 177
Photorhabdus  INIL 171
                *
```

Fig . 2

581   Fig. 3

A

B

583   Fig. 4