Identification of new *Dickeya dadantii* virulence factors secreted

by the type 2 secretion system.

Guy Condemine* and Bastien Le Derout

Univ Lyon, Université Lyon 1, INSA de Lyon, CNRS UMR 5240 Microbiologie Adaptation

et Pathogénie, F-69622 Villeurbanne, France.

* Corresponding author: guy.condemine@insa-lyon.fr

Short title: *Dickeya dadantii* virulence factors

ABSTRACT

Dickeya are plant pathogenic bacteria able to provoke disease on a wide range of plants. A type

2 secretion system (T2SS) named Out is necessary for bacterial virulence. Its study in *D.*

dadantii showed that it secretes a wide range of plant cell wall degrading enzymes, including

pectinases and a cellulase. However, the full repertoire of exoproteins it can secrete has

probably not yet been identified. Secreted proteins are first addressed to the periplasm before

their secretion by Out. No secretion signal present on the protein allows the identification of

substrates of a T2SS. To identify new Out substrates, we analyzed *D. dadantii* transcriptome

data obtained in plant infection condition and searched for genes strongly induced encoding a

protein with a signal sequence. We identified four new Out-secreted proteins: the expansin

YoaJ, the putative virulence factor VirK and two proteins of the DUF 4879 family, SvfA and

SvfB. We showed that SvfA and SvfB are required for full virulence of *D. dadantii* and showed

that *svf* genes are present in a variable number of copies in other *Pectobacteriaceae*, up to three

in *D. fanghzongdai*. This work opens the way to the study of the role of non-pectinolytic

proteins secreted by the Out pathway in *Pectobacteriaceae*.

IMPORTANCE

The plant pathogen *Dickeya* rely on a type 2 secretion system named Out for their pathogenicity.

Importance of plant cell wall degrading enzymes secreted by this system has been well studied.

However, existence and role of other Out-secreted proteins has barely been investigated. By

mining *D. dadantii* transcriptome data, we identified four new Out-secreted proteins. We

showed that two of them, SvfA and SvfB, are necessary for the full virulence of the bacteria.

These findings show that identification of all the proteins secreted by the *Dickeya* Out system

is necessary for a better knowledge of the virulence of these bacteria.

INTRODUCTION

Soft rot *Pectobacteriaceae* (SRP), *Dickeya* and *Pectobacterium*, are plant pathogenic bacteria that can provoke disease on more than 35% of angiosperm plant orders, including both monocot and dicot plants (1). Among those, there is a wide range of plants of agronomic interest such as potato, rice, chicory, cabbage or ornementals on which they can cause severe losses. Symptoms are usually soft rot but these bacteria can provoke blackleg or wilting on aerial parts of potato. Recently, diseases on woody plants caused by *Dickeya* have been reported (2). There is no efficient way to fight these bacterial diseases. There are actually twelve species of *Dickeya* described, isolated either from infected plants (type strain of *D. chrysanthemi* isolated from *Chrysanthemum morifolium*, *D. dadantii* subsp. *dadantii* from *Pelargonium capitum*, *D. dadantii* subsp. *diffenbachiae* from *Dieffenbachia* sp., *D. dianthicola* from *Dianthus caryophillus*, *D. zeae* from *Zea mays*, *D. oryzae* from *Oryza sativa*, *D. paradisiaca* from *Musa paradisiaca*, *D. solani* from *Solanum tuberosum*, *D. fangzhongdai* from *Pyrus pyrifolia*, *D. poaceiphila* from *Saccharum officinarum*) (3)(4)(5)(6)(7) or from river of lake waters (*D. aquatica*, *D. lacustris* and *D. undicola*) (8)(9)(10). The role of protein secretion systems on the onset of the disease provoked by these bacteria has been recognized long ago (11). In contrast to many plant pathogenic bacteria, the type three Hrp secretion system is not the main determinant for SRP virulence (12). The main virulence factor for these bacteria is a type 2 secretion system (T2SS) named Out. It allows the secretion of enzymes that degrade the components of the plant cell wall, leading to the soft rot symptom distinctive of the disease. The first Out-secreted proteins to be identified were a set of pectinases and a cellulase which are easily detectable by simple enzymatic tests (13) (11). The pectinolytic secretome of the model strain *D. dadantii* 3937 has been studied in detail by cloning the genes of these easily

detectable enzymes. *D. dadantii* secretes by the Out machinery nine pectate lyases, one pectin

methylesterase, one pectin acetylesterase and one rhamnogalacturonate lyase (14). A proteomic

analysis of the secreted proteins by 2D gel electrophoresis allowed the identification of two

other secreted proteins, the feruloyl esterase FaeD and a protein with homology with a

Xanthomonas campestris avirulence protein AvrL (15). A search in *D. dadantii* of homologues

of proteins secreted by the Out T2SS of *Pectobacterium atrosepticum* (16) recently led to the

characterization of the metal binding protein IbpS (17). There is no strict host specificity for

Dickeya species, however some of them show a preference for some plant species. Since all the

pectinolytic enzymes studied in *D. dadantii* are present in most of other *Dickeya* species these

enzymes are probably not responsible for the host preference observed for these bacteria (18).

We hypothesized that additional T2SS-secreted proteins specific for some species might exist

and play a role in the host preference. To identify such proteins, we analyzed previously

published *D. dadantii* transcriptome data, looking for genes induced in plant infection

conditions and encoding proteins with a signal sequence. We identified several proteins

secreted by the Out machinery and showed that two proteins of the DUF4879 family, SvfA and

SvfB are *D. dadantii* virulence factors.

MATERIAL AND METHODS
85

Bacterial strains and growth conditions.

Bacterial strains, phages, plasmids and oligonucleotides used in this study are described in Table 2. *D.*

dadantii and *E. coli* cells were grown at 30 and 37°C respectively in LB medium or M63 minimal medium

supplemented with a carbon source (0.2%, w/v unless otherwise indicated). When required antibiotics

were added at the following concentrations: ampicillin, 100 mg l^{-1}, kanamycin and chloramphenicol, 25

mg l^{-1}. Media were solidified with 1.5% (w/v) agar. Transduction with phage ΦEC2 was performed

according to Résibois *et al.* (40)

Mutant construction.

To construct strain A6418 that contains a *svfA-uidA* fusion a 1.3 kb DNA fragment containing

svfA was amplified with primers 17176H+ and 17176A. The resulting fragment was inserted into

the pGEM-T plasmid (Promega). A XbaI site was created by site directed mutagenesis with the

primers 17176XbaF and 17176XbaR into the *svfA* coding sequence and a *uidA*-kanR cassette was

inserted into this XbaI site. To construct strain A6467 that contains a *svfB-uidA*-kanR fusion and

a CmR cassette a 2000bp DNA fragment containing *svfB* was amplified with the primers

15544L2+ and 15544L2-. The resulting fragment was inserted into the pGEM-T plasmid. A XmaI

site was created by site directed mutagenesis into *svfB* coding sequence with the primers

15544XmaF and 15544XmaR and a *uidA*-kanR cassette was inserted into this created unique XmaI

site. All the constructs were recombined into the *D. dadantii* chromosome according to Roeder

and Collmer (41). Recombinations were checked by PCR. His-tagged versions of the proteins

SvfA, SvfB, YoaJ and VirK were constructed by amplifying the corresponding genes with the

primers 17176H+ and 17176H-, 15544H+ and 15544H-, 14642H+ and 14642H-, VirKH+ and

VirKH-, respectively. The resulting DNA fragments were cloned into plasmid pGEMT.

Secretion assays and Western blots.

D. dadantii strains containing the plasmid to test were grown overnight in LB medium in the

presence of the appropriate antibiotic. Culture supernatant containing secreted proteins was

separated from cells by centrifugation at 10,000 g for 3 min, and both fractions were loaded onto

12% polyacrylamide gel electrophoresis (SDS-PAGE). The proteins were next transferred onto

Immobilon P membrane (Merck) and probed with Ni-NTA-HRP.

Pathogenicity tests.

Bacteria were grown overnight in LB medium, centrifuged and resuspended at OD_{600} 1 in M63

medium. Potatoes were surface sterilized with 70% ethanol and dried. A hole was made with a

pipette tip and 10 μl of bacteria were deposited in the hole which was covered with mineral oil.

Potatoes were placed over a wet paper in a tray contained in a plastic bag to maintain moisture.

After 48 h at 30°C, the weight of rotten tissue was measured.

Enzymatic assays.

β-glucuronidase assays were performed on toluenized extracts of cells grown to exponential

phase using the method of Bardonnet *et al* (42) with *p*-nitrophenyl-β-D-glucuronate as the

substrate.

RESULTS

Identification of new Out-secreted proteins

To have a more complete knowledge of the proteins secreted by the *D*. *dadantii* Out T2SS that

could be involved in the pathogenicity process, we searched for candidate genes in recently

published transcriptome data (19)(20). We selected genes strongly induced during plant

infection and coding for a protein possessing a signal sequence. We retained the genes

Dda3937_01687, *Dda3937_00585* (thereafter named SvfA and SvfB, respectively) and

Dda3937_00081 (also named *yoaJ*). We also retained VirK, a protein of unknown function

with a signal sequence identified among the genes controlled by the transcriptional regulator

PecS of many virulence factors (21). Each protein was tagged with a His-tag and its secretion

was analyzed in the *D*. *dadantii* wild type strain and an *outD* mutant in which the Out machinery

is not functional. The proteins SvfA, SvfB, YoaJ and VirK were detected in the supernatant of

the wild type strain but not of the mutant, demonstrating their secretion by the Out machinery

(Fig. 1). Production of these proteins from a gene cloned on a multicopy plasmid may explain

why secretion was not total in the wild type strain.

YoaJ is a PecS-regulated gene (21) and it was found among the most induced genes during

Arabidopsis infection or culture in the presence of plant extracts (19). It encodes a protein with

homology to expansins. These proteins are able to non-enzymatically loosen cell wall cellulose.

They are found in all plants where they have a role in cell wall extension and also in many plant

pathogenic microorganisms (22). Their role in virulence has been shown in *Ralstonia*

solanacearum, in *P. atrosepticum*, *P. Brasiliense* and in the plant pathogen *Erwinia*

tracheiphila (23) (24). The role of *D. dadantii* expansin is probably the same. VirK is a protein

of unknown function that has homologues in several plant pathogenic bacteria such as *R.*

solanacearum, *Agrobacterium tumefaciens, Lonsdalea* and *Xanthomonas*. VirK is controlled

by PecS and induced during Arabidopsis infection or culture in the presence of plant extracts

(21) (20). No symptom for the *D. dadantii virK* mutant was observed whatever the plant tested

(21). The two proteins SvfA and SvfB are studied in the next paragraphs.

SvfA and SvfB are virulence factors

svfA and *svfB* are among the most induced *D. dadantii* genes during Arabidopsis infection or

during culture of the bacteria in the presence of plant extracts (19). They are also strongly

expressed during maceration of potato tubers by *D. dianthicola* and *D. solani* (25). The two *D.*

dadantii proteins share 43% identity and 58% similarity in amino acid composition (Fig. 2).

SvfA is 187 amino acid long (165 for the mature form, 17.5 kDa) and SvfB is 198 amino acid

long (177 for the mature form, 18.9 kDa). These proteins belong to the DUF 4879 family of

proteins. Proteins of this family have no known function. YolA,a protein of the DUF 4879

family showing low homology with SvfB, is among the most highly secreted protein of *Bacillus*

subtilis (26). YolA is also present in *B. cereus* and in the insect pathogen *B. thuringiensis*. *B.*

subtilis YolA is shorter than SvfA and SvfB, missing the N-terminal more variable part (Fig.

2).

svfA and *svfB* mutants have been constructed and their pathogenicity has been tested on potato.

The *svfA* mutant was significantly less aggressive than the wild type strain while the *svfB* mutant

was not significantly affected (Fig. 3A). Virulence of the *svfA* mutant could be restored by

introduction of a plasmid bearing the wild type *svfA* gene (Fig. 3B). Virulence of the double

svfA svfB mutant was further reduced showing that the role of SvfB is additive to that of SvfA

(Fig. 3A). Thus, genes *Dda3937_01687*and *Dda3937_00585* were named *svfA* and *svfB* for

s̲ecreted v̲irulence f̲actor A and B.

All our attempts to overproduce the proteins SvfA and SvfB in order to purify them and to study

more precisely their function were unsuccessful because their production was toxic to the

bacterial cells engineered to overproduce them.

Expression of *svfA* and *svfB*

To try to identify the function of SvfA and SvfB, we analyzed the conditions in which their

genes are expressed. We tested the effect of galacturonate and polygalacturonate, two

compounds that are inducers of the expression of the main virulence factors, the pectate lyases,

and of glucose, which represses it. We also analyzed the effect of mutations in genes controlling

several aspects of *D. dadantii* virulence. KdgR represses the pectinase, pectin degradation and

out genes (27). Its inducer is 2-keto-3-deoxygluconate, a polygalacturonate and galacturonate

catabolic derivative. PecS controls genes encoding the pectinases, diverse secreted protein, the

Out machinery and proteins involved in resistance to oxidative stress (28). PecT is a regulator

of the pectate lyase, motility and exopolysaccharide synthesis genes (29). Pir regulates

hyperinduction of pectate lyses in response to plant extracts (30). GacA, the regulator of the

two-component regulatory system GacA-GacS, is a global regulator required for disease

expression in response to the metabolic status of the bacteria (31). Expression of *svfA* was

slightly induced by polygalacturonate but not by galacturonate (Fig. 4A). However, expression

of this gene was not modified in a *kdgR* background indicating that induction by

polygalacturonate is not mediated by KdgR. Growth in the presence of chicory chunks strongly

induced s*vfA* expression as expected from transcriptomic data showing induction in the

presence of plant extract. A high concentration of glucose led to a strong induction of *svfA*

expression (Fig. 4A). This regulation is mediated by the catabolite repressor protein CRP since

a mutation in the *crp* gene derepressed *svfA* expression. Thus, Crp is a repressor of *svfA*.

Although it had not been previously identified as a PecS-regulated gene (21), *svf*A expression

is increased in a *pecS* background. A *pir* mutation provoked a weak derepression of *svfA*

expression. Neither PecT nor GacA significantly regulate *svfA* expression (Fig. 4A).

Regulation of *svfB* shows some similarity to that of *svfA*: it was not induced by galacturonate,

polygalacturonate or regulated by KdgR, it was induced by glucose and repressed by Crp, and

it was repressed by PecS (Fig. 4B). However, a few differences can be noted: in contrast to

what is observed with *svfA*, no induction by plant pieces was observed for *svfB* and PecT was

a repressor of *svfB* expression while Pir did not seem to control it (Fig. 4B).

Occurrence of the new secreted proteins in other Dickeya species

Presence of *svfA*, *svfB*, *virK* and *yoaJ* was searched in the genome of all the *Dickeya* type strains,

and in a few *Pectobacterium* strains (Table 1). Presence and number of proteins of the DUF

4879 family is variable among *Dickeya* species. The gene *svfA* is present in all strains except

D. zeae, *D. chrysanthemi*, *D. poaceiphila* and *D. paradisiaca*. The gene *svfB* is present in most

species but is absent in *D. chrysanthemi*, *D. poaceiphila* and *D. paradisiaca*, *D. undicola* and

D. aquatica. A third gene located next to *svfA* and probably resulting from a duplication is

found in *D. fangzhongdai* and *D. undicola*. It was named *svfC*. It has 53% homology with *D.*

dadantii SvfA and 34% with *D. dadantii* SvfB. SvfC is shorter that SvfA and SvfB (126 amino

acid for the mature protein, 13.2 kDa) and has the same size as *B. cereus* YolA (Fig. 2). It

possesses a signal sequence, indicating that it could also be secreted by the Out system. Thus,

the number of genes of the DUF 4879 family in *Dickeya* strains varies from 0 to 3. Homologues

of the *svf* genes can also be found in some *Pectobacterium* strains (Table 1). For example, two

copies are present in *P. carotovorum* subsp *carotovorum*. However, even in a given species, the

gene may be present or not (presence of a homologue of *svfB* in 10 out of the 23 *P. brasiliense*

strains present in the ASAP data bank (https://asap.ahabs.wisc.edu/asap/home.php). Outside

Pectobacteriaceae, homologues of *svfB* can be found in a few Gammaproteobacteriaceae, i.d.

in some *Photorhabdus*, *Luteibacter* and *Pseudoalteromonas* strains. *yoaJ* is present in all

Dickeya and *Pectobacterium* strains except *D. poaceiphila* and *D. paradisiaca*. *virK* is present

in all Dickeya strains except in *D. aquatica* and absent in all *Pectobacterium* strains tested.

We also examined the presence or absence of genes of other non-pectinolytic proteins known

to be secreted by a T2SS in *Dickeya* or *Pectobacterium*: *ibpS*, *nipE*, *xynA*, *avrL/avrM* (Table

1). IbpS is a metal binding protein that prevents ROS-induced killing of bacteria (17). NipE is

a toxin that provoke plant cell death (32). XynA is a xylanase that was identified in a corn strain

of *Dickeya zeae* previously named *Erwinia chrysanthemi* (33). AvrL is homologous to the

Xanthomonas campestris avirulence protein AvrL (15). Two very similar proteins, AvrL and

AvrM, are encoded by *D. dadantii* 3937. AvrL was named Svx in *P. atrosepticum* where a role

in virulence has been proven (34). However, its function in *Dickeya* has not been studied. IbpS

is present in almost all the species, except *D. paradisiaca*. NipE is absent in *D. paradisiaca* and

D. poaceiphila.   Presence of XynA is variable in *Dickeya* strains and it is absent in

Pectobacterium. A variation in the presence and number of AvrL can be observed in Dickeya

strains (Table 1). Thus, the repertoire of T2SS-secreted protein known to be important for

virulence is very variable from species to species.

251
DISCUSSION

The T2SS of *Dickeya* and *Pectobacterium* is a major virulence factor of these bacteria. The

knowledge of the repertoire of secreted proteins is necessary to better understand the precise

mechanisms of virulence of these bacteria. These analyses have been undertaken with the model

strain *D. dadantii* 3937 and partially with *Pectobacterium atrosepticum* (35)(16). *Dickeya* and

Pectobacterium are characterized by their ability to degrade pectin and they are identified by

this characteristic on the semi selective Crystal Violet Pectate medium. They all secrete

enzymes   capable   of   degrading   pectin   (pectate   lyases,   polygalacturonases,   pectin

methylesterases). However, recent works show that other proteins are secreted by the Out T2SS

(15)(16). In the present work we used published transcriptome data to identify new potential

substrates of the *D. dadantii* T2SS. The most highly induced genes in a transcriptome

experiment of *D. dadantii* infecting *A. thaliana* are known virulence genes (*pelI, prtA, rhiE,*

paeY, rhaD, ibpS, etc...) (19). However, in this top list some genes have no known function.

The presence of a signal sequence in their product suggested that these proteins could be

substrates of the T2SS necessary for the infection process. We showed here that the proteins

SvfA, SvfB and YoaJ produced by genes present in the top list of those induced in Arabidopsis

are substrates of the Out T2SS. YoaJ belongs to the family of expansins, proteins that loosen

cellulose fibers. Their role as a virulence factor has been shown in *P. brasiliense* and *P.*

atrosepticum (23) and it probably has the same function in *D. dadantii* and other *Dickeya*

species. No function could be predicted for SvfA and SvfB which belong to the DUF 4879

family of proteins. However, a reduction of virulence of a *svfA* mutant and a *svfA svfB* double

mutant on potato could be observed, proving a role of the proteins in the bacterial pathogenicity.

Although an additive effect of the mutations was observed, they could not have exactly the

same function. The mutants should be tested on various hosts to detect potential differences. It

can be supposed that each protein would be more active on one type or one family of plant.

Presence of three DUF 4879 proteins in *D. fanghzongdai* could explain its wide host range,

from orchid to pear trees. Presence of homologues of SvfA and SvfB in *Photorhabdus* and in

B. thuringiensis strains, two insect pathogens, indicates that the role of these proteins is not

restricted to plant virulence but may participate to a common process of bacterial pathogeny.

We also showed here that the PecS-regulated protein VirK is secreted by Out. No role on

virulence had been observed for this protein with the chicory leaf model of infection (21). Other

models should be tested to find the role of this protein.

Regulation of expression of the *svfA* and *svfB* genes is atypical for a *D. dadantii* gene involved

in pathogeny. While expression of most of the virulence factors is induced in the presence of

pectin or its derivatives through the repressor KdgR and repressed by glucose, that of *svfA* and

svfB is opposite: it is activated by glucose and not controlled by KdgR. Expression of *svfA* is

induced in the presence of plant tissue. This pattern of regulation has been described for *ibpS*,

which is also strongly induced in *A. thaliana* (17). This could correspond to conditions

encountered during the early phases of infection: pectin has not yet been degraded and glucose

and saccharose are plentiful in plant tissues. *svfA* and *ibpS* could be among the earliest gene to

be induced at the onset of infection, before the genes involved in pectin degradation. However,

regulation of these genes by PecS and PecT shows that *svfA* and *svfB* are fully integrated in the

network of regulators that controls *D. dadantii* virulence.

This work has extended our knowledge of the Out-dependent secretome of *D. dadantii*, showing

that besides pectinases several other proteins are secreted. If the number of pectinolytic

enzymes secreted is almost identical in the various *Dickeya* species, the number of additional

secreted proteins varies markedly. Among the proteins analyzed (Table 1), *D. paradisiaca* has

only one (VirK) while *D. fanghzongdai* has ten. All the intermediate combinations can be found

in the various species. There seems to be less variations in the *Pectobacterium* strains surveyed.

It is tempting to speculate that the presence/absence of these proteins could influence the host

preference of some *Dickeya* species, providing additional virulence factors favorable to infect

certain hosts. Works that compare *Dickeya* strains to understand what makes difference in their

host range or aggressivity often focus only on the presence of the six known types of secretion

systems without analyzing what proteins could be secreted (18)(36)(37). An exhaustive analysis

of the secreted proteins would be more informative.

Is there other T2SS-secreted proteins to be identified in *Dickeya* strains? No specific signal is

present on T2SS-secreted proteins that would allow their identification. 2D gels which were

used in previous studies performed on *D. dadantii* and *P. atrosepticum* to identify their

secretome have a limited sensitivity (15)(34). More sensitive methods such as liquid

chromatography-tandem mass spectrometry (LC-MS/MS) can now be used (38). However,

they give many false positive results since periplasmic and cytoplasmic proteins are often found

in the culture supernatant. The approach we used here allowed the identification of four new

secreted proteins. However, all these methods have a drawback. They can only detect proteins

in conditions where they are produced. For instance, the rhamnogalacturonate lyase RhiE could

only be detected when the bacteria were cultivated in the presence of rhamnose (39). The genes

encoding YoaJ and VirK were not induced in *D. dianthicola* grown on potato (25). Another

problem is that a protein may not exist in the strain tested. An analysis of the secretome of

several *Dickeya* strains grown in several conditions will be necessary to have a global view of

all the additional virulence factors that can be secreted by *Dickeya* species and evaluate their

potential role in pathogenicity.

ACKNOWLEDGMENTS

We thank Lison Massardier and Florence Ruaudel for technical work and Nicole Cotte-Pattat

for reading the manuscript. This work was supported by funding from CNRS, University Lyon

1 and Agence Nationale de la Recherche (ANR-19-CE35-0016).

REFERENCES

1.   Ma B, Hibbing ME, Kim H-S, Reedy RM, Yedidia I, Breuer J, Breuer J, Glasner JD, Perna

NT, Kelman A, Charkowski AO. Host range and molecular phylogenies of the soft rot

enterobacterial genera *Pectobacterium* and *Dickeya*. Phytopathology. 2007; 97:1150–63.

2.   Fujikawa T, Hatomi H, Ota N. Draft genome sequences of seven strains of *Dickeya*

dadantii, a quick decline-causing pathogen in fruit trees, isolated in Japan. Microbiology

Resource Announcements. 2020; 9:e00609-20.

3.   Samson R, Legendre JB, Christen R, Saux MF-L, Achouak W, Gardan L. Transfer of

Pectobacterium chrysanthemi (Burkholder et al. 1953) Brenner *et al*. 1973 and *Brenneria*

paradisiaca to the genus *Dickeya* gen. nov. as *Dickeya chrysanthemi* comb. nov. and

Dickeya paradisiaca comb. nov. and delineation of four novel species, *Dickeya dadantii*

sp. nov., *Dickeya dianthicola* sp. nov., *Dickeya dieffenbachiae* sp. nov. and *Dickeya zeae*

sp. nov. International Journal of Systematic and Evolutionary Microbiology. 2005;

55:1415–27.

4.   Brady CL, Cleenwerck I, Denman S, Venter SN, Rodríguez-Palenzuela P, Coutinho TA,

De Vos P. Proposal to reclassify *Brenneria quercina* (Hildebrand and Schroth 1967)

Hauben *et al*. 1999 into a new genus, *Lonsdalea* gen. nov., as *Lonsdalea quercina* comb.

nov., descriptions of *Lonsdalea quercina* subsp. *quercina* comb. nov., *Lonsdalea quercina*

subsp. *iberica* subsp. nov. and *Lonsdalea quercina* subsp. *britannica* subsp. nov.,

emendation of the description of the genus *Brenneria*, reclassification of *Dickeya*

dieffenbachiae as *Dickeya dadantii* subsp. *dieffenbachiae* comb. nov., and emendation of

the description of *Dickeya dadantii*. International Journal of Systematic and Evolutionary

Microbiology. 2012; 62:1592–1602.

5.   van der Wolf JM, Nijhuis EH, Kowalewska MJ, Saddler GS, Parkinson N, Elphinstone

JG, Pritchard L, Toth IK, Lojkowska E, Potrykus M, Waleron M, de Vos P, Cleenwerck

I, Pirhonen M, Garlant L, Hélias V, Pothier JF, Pflüger V, Duffy B, Tsror L, Manulis S.

Dickeya solani sp. nov., a pectinolytic plant-pathogenic bacterium isolated from potato

(*Solanum tuberosum*). International Journal of Systematic and Evolutionary

Microbiology. 2014; 64:768–74.

6.   Tian Y, Zhao Y, Yuan X, Yi J, Fan J, Xu Z, Hu B, De Boer SH, Li X. *Dickeya*

fangzhongdai sp. nov., a plant-pathogenic bacterium isolated from pear trees (*Pyrus*

pyrifolia). International Journal of Systematic and Evolutionary Microbiology. 2016;

66:2831–35.

7. Wang X, He S-W, Guo H-B, Han J-G, Thin KK, Gao J, Wang Y, Zhang X-X. *Dickeya*

oryzae sp. nov., isolated from the roots of rice. International Journal of Systematic and

Evolutionary Microbiology. 2020; 70:4171–78.

8. Parkinson N, DeVos P, Pirhonen M, Elphinstone J. *Dickeya aquatica* sp. nov., isolated

from waterways. International Journal of Systematic and Evolutionary Microbiology.

2014; 64:2264–6.

9. Hugouvieux-Cotte-Pattat N, Jacot-des-Combes C, Briolay J. *Dickeya lacustris* sp. nov., a

water-living pectinolytic bacterium isolated from lakes in France. International Journal of

Systematic and Evolutionary Microbiology. 2019; 69:721–26.

10. Oulghazi S, Pédron J, Cigna J, Lau YY, Moumni M, Van Gijsegem F, Chan K-G, Faure

D. *Dickeya undicola* sp. nov., a novel species for pectinolytic isolates from surface waters

in Europe and Asia. International Journal of Systematic and Evolutionary Microbiology.

2019; 69:2440–4.

11. Ji J, Hugouvieux-Cotte-Pattat N, Robert-Baudouy J. Molecular cloning of the *outJ* gene

involved in pectate lyase secretion by *Erwinia chrysanthemi*. Molecular Microbiology.

1989; 3:285–93.

12. Yang C-H, Gavilanes-Ruiz M, Okinaka Y, Vedel R, Berthuy I, Boccara M, Chen JW-T,

Perna NT, Keen NT. *hrp* genes of *Erwinia chrysanthemi* 3937 are important virulence

factors. Molecular Plant-Microbe Interactions. 2002; 15:472–480.

13.   Andro T, Chambost JP, Kotoujansky A, Cattaneo J, Bertheau Y, Barras F, Van Gijsegem

F, Coleno A. Mutants of *Erwinia chrysanthemi* defective in secretion of pectinase and

cellulase. Journal of Bacteriology. 1984; 160:1199–203.

14.   Hugouvieux-Cotte-Pattat N, Condemine G, Shevchik VE. Bacterial pectate lyases,

structural and functional diversity. Environmental Microbiology Reports. 2014; 6:427–

40.

15.   Kazemi-Pour N, Condemine G, Hugouvieux-Cotte-Pattat N. The secretome of the plant

pathogenic bacterium *Erwinia chrysanthemi*. Proteomics. 2004; 4:3177–3186.

16.   Coulthurst SJ, Lilley KS, Hedley PE, Liu H, Toth IK, Salmond GPC. DsbA plays a critical

and multifaceted role in the production of secreted virulence factors by the phytopathogen

Erwinia carotovora subsp. *atroseptica*. Journal of Biological Chemistry. 2008;

283:23739–53.

17.   Liu L, Gueguen-Chaignon V, Gonçalves IR, Rascle C, Rigault M, Dellagi A, Loisel E,

Poussereau N, Rodrigue A, Terradot L, Condemine G. A secreted metal-binding protein

protects necrotrophic phytopathogens from reactive oxygen species. Nature

Communications. 2019; 10:4853.

18.   Potrykus M, Golanowska M, Hugouvieux-Cotte-Pattat N, Lojkowska E. Regulators

involved in *Dickeya solani* virulence, genetic conservation, and functional variability.

Molecular Plant-Microbe Interactions 2014; 27:700–11.

19.   Pédron J, Chapelle E, Alunni B, Van Gijsegem F. Transcriptome analysis of the *Dickeya*

dadantii PecS regulon during the early stages of interaction with *Arabidopsis thaliana*.

Molecular Plant Pathology. 2018; 19:647–63.

20.   Jiang X, Zghidi-Abouzid O, Oger-Desfeux C, Hommais F, Greliche N, Muskhelishvili G,

Nasser W, Reverchon S. Global transcriptional response of *Dickeya dadantii* to

environmental stimuli relevant to the plant infection. Environmental Microbiology. 2016;

18:3651–72.

21.   Hommais F, Oger-Desfeux C, Van Gijsegem F, Castang S, Ligori S, Expert D, Nasser W,

Reverchon S. PecS Is a global regulator of the symptomatic phase in the phytopathogenic

bacterium *Erwinia chrysanthemi* 3937. *Journal of Bacteriology*. 2008; 190:7508–22.

22.   Tancos MA, Lowe-Power TM, Peritore-Galve FC, Tran TM, Allen C, Smart CD. Plant-

like bacterial expansins play contrasting roles in two tomato vascular pathogens.

Molecular Plant Pathology. 2018; 19:1210–21.

23.   Narváez-Barragán DA, Tovar-Herrera OE, Torres M, Rodríguez M, Humphris S, Toth IK,

Segovia L, Serrano M, Martínez-Anaya C. Expansin-like Exl1 from *Pectobacterium* is a

virulence factor required for host infection, and induces a defence plant response involving

ROS, and jasmonate, ethylene and salicylic acid signalling pathways in *Arabidopsis*

thaliana. Scientific Reports. 2020; 10:7747.

24.   Rocha J, Shapiro LR, Kolter R. A horizontally acquired expansin gene increases virulence

of the emerging plant pathogen *Erwinia tracheiphila*. Scientific Reports. 2020; 10:21743.

25.   Raoul des Essarts Y, Pédron J, Blin P, Van Dijk E, Faure D, Van Gijsegem F. Common

and distinctive adaptive traits expressed in *Dickeya dianthicola* and *Dickeya solani*

pathogens when exploiting potato plant host. Environmental Microbiology. 2019.

21:1004–18.

26.   Tjalsma H, Antelmann H, Jongbloed JDH, Braun PG, Darmon E, Dorenbos R, Dubois J-

YF, Westers H, Zanen G, Quax WJ, Kuipers OP, Bron S, Hecker M, van Dijl JM.

Proteomics of protein secretion by *Bacillus subtilis*: Separating the "secrets" of the

secretome. Microbiology and Molecular Biology Reviews. 2004; 68:207–33.

27. Condemine G, Robert-Baudouy J. Tn*5* insertion in *kdgR*, a regulatory gene of the

polygalacturonate pathway in *Erwinia chrysanthemi*. FEMS Microbiology Letters. 1987;

42:39–46.

28. Reverchon S, Nasser W, Robert-Baudouy J. *pecS*: a locus controlling pectinase, cellulase

and blue pigment production in *Erwinia chrysanthemi*. Molecular Microbiology. 1994;

11:1127–39.

29. Surgey N, Robert-Baudouy J, Condemine G. The *Erwinia chrysanthemi pecT* gene

regulates pectinase gene expression. Journal of Bacteriology. 1996; 178:1593–99.

30. Nomura K, Nasser W, Kawagishi H, Tsuyumu S. The *pir* gene of *Erwinia chrysanthemi*

EC16 regulates hyperinduction of pectate lyase virulence genes in response to plant

signals. Proceedings of the National Academy of Sciences. 1998; 95:14034–39.

31. Lebeau A, Reverchon S, Gaubert S, Kraepiel Y, Simond-Côte E, Nasser W, Van Gijsegem

F. The GacA global regulator is required for the appropriate expression of *Erwinia*

chrysanthemi 3937 pathogenicity genes during plant infection. Environmental

Microbiology. 2008; 10:545–59.

32. Laasik E, Põllumaa L, Pasanen M, Mattinen L, Pirhonen M, Mäe A. Expression of *nipP.w*

of *Pectobacterium wasabiae* is dependent on functional *flgKL* flagellar genes.

Microbiology. 2014; 160:179–86.

33. Keen NT, Boyd C, Henrissat B. Cloning and characterization of a xylanase gene from corn

strains of *Erwinia chrysanthemi*. Molecular Plant-Microbe Interactions. 1996; 9:651-7.

34. Corbett M, Virtue S, Bell K, Birch P, Burr T, Hyman L, Lilley K, Poock S, Toth I,

Salmond G. Identification of a new quorum-sensing-controlled virulence factor in *Erwinia*

carotovora subsp. *atroseptica* secreted via the type II targeting pathway. Molecular Plant-

Microbe Interactions. 2005; 18:334–42.

35. Mattinen L, Nissinen R, Riipi T, Kalkkinen N, Pirhonen M. Host-extract induced changes

in the secretome of the plant pathogenic bacterium *Pectobacterium atrosepticum*.

Proteomics. 2007; 7:3527–37.

36. Alič Š, Pédron J, Dreo T, Van Gijsegem F. Genomic characterisation of the new *Dickeya*

fangzhongdai species regrouping plant pathogens and environmental isolates. BMC

Genomics. 2019; 20:34.

37. Xu P, Wang H, Qin C, Li Z, Lin C, Liu W, Miao W. Analysis of the taxonomy and

pathogenic factors of *Pectobacterium aroidearum* L6 using whole-genome sequencing

and comparative genomics. Frontiers in Microbiology. 2021; *a*12:679102.

38. Burtnick MN, Brett PJ, DeShazer D. Proteomic Analysis of the *Burkholderia*

pseudomallei type II secretome reveals hydrolytic enzymes, novel proteins, and the

deubiquitinase TssM. Infection and Immunity. 2014; 82:3214–26.

39. Laatu M, Condemine G. Rhamnogalacturonate lyase RhiE is secreted by the Out system

in *Erwinia chrysanthemi*. Journal of Bacteriology. 2003; 185:1642–9.

40. Resibois A, Colet M, Faelen M, Schoonejans E, Toussaint A. ɸEC2, a new generalized

transducing phage of *Erwinia chrysanthemi*. Virology. 1984; 137:102–12.

41. Roeder DL, Collmer A. Marker-exchange mutagenesis of a pectate lyase isozyme gene in

Erwinia chrysanthemi. Journal of Bacteriology. 1985; 164:51–6.

42. Bardonnet N, Blanco C. ′uidA-antibiotic-resistance cassettes for insertion mutagenesis,

gene fusions and genetic constructions. FEMS Microbiology Letters. 1992; 93:243–7.

Fig 1: Identification of new secreted proteins by *D. dadantii*. Wild-type and *outD* mutant strains

containing plasmid bearing the gene of the selected protein were grown overnight in LB

medium. The supernatant (S) and cellular (C) fractions were separated by SDS-PAGE. After

blotting, the proteins were detected with Ni-NTA-HRP.

Fig. 2: Alignment of Svf proteins. The sequences of *D. dadantii* SvfA (Dda3937_01687) and

SvfB (Dda3937_00585), *D. fanghzongdai* SvfC (CVE23_15565), *B. cereus* WP-193674364.1

and *Photorhabdus asymbiotica* CAQ86327.1, without their signal sequence, were aligned with

Clustal omega. Identical residues are indicated by a star and chemically equivalent residues by

a double dot.

Fig. 3. Virulence of *svfA* and *svfB* mutants. A. Potatoes were infected with the wild type strain,

and the *svfA*, the *svfB* and the *svfA svfB* mutants. Rotten tissue was weighed after 48 h. B.

Complementation of the *svfA* mutation. Potatoes were infected with the wild type strain, the

svfA mutant containing the empty plasmid pBBRGm and the *svfA* mutant containing the

pBBRGm plasmid bearing *svfA*. Rotten tissue was weighed after 48h. Statistical tests were

performed using the Wilcoxon- Mann-Whitney test. The *p*-value were compared with an alpha

risk of 4%. $p < 0.001 = ***, p < 0.005 = **, p < 0.01 = *$.

Fig 4 : Expression of *svfA* and *svfB* in various growth conditions. A. The *D. dadantii* strain

A6418 containing the *svfA-uidA* fusion and its derivative strains containing an additional

regulatory mutation were grown in M63 medium in the presence of the indicated compounds

(Y = glycerol, G = glucose, A = galacturonate, PGA = polygalacturonate, E = chicory chunks).

Strains with additional mutations were grown with glycerol as a carbon source except the *crp*

mutant that was grown with 0.2% glucose. ß-glucuronidase activity was measured with *p*-

nitrophenyl-ß-D-glucuronate. B. Similar experiment for the *D. dadantii* strain A6467

containing the *svfB-uidA* fusion and its derivative strains containing an additional regulatory

mutation. Activities are expressed in µmoles of *p*-nitrophenol produced per minute and per

milligram of bacterial dry weight $\pm$ standard deviation. Data are expressed as the mean (n = 6)

from six independent experiments. Statistical tests were performed using the Wilcoxon- Mann-

Whitney test. The *p*-value were compared with an alpha risk of 4%. $p < 0.001$=***, $p <$

0.005=**, $p < 0.01$=*.

Table 1: Presence of Out-secreted proteins in various *Dickeya* and *Pectobacterium* strains.

Table 2: Oligonucleotides and strains used in this study.

.

Table 1 : Presence of Out-secreted proteins in various *Dickeya* and *Pectobacterium* strains.
523

Strain	*SvfA*	*SvfB*	*SvfC*	*YoaJ*	*VirK*	*IbpS*	*NipE*	*XynA*	*AvrL*
D. dadantii	1	1	0	1	1	1	1	1	2
D. diffenbachiae	1	1	0	1	1	1	1	0	2
D. fangzhongdai	1	1	1	1	1	1	1	1	2
D. solani	1	1	0	1	1	1	1	1	2
D. zeae	0	1	0	1	1	1	1	1	1
D. oryzae	1	1	0	1	1	1	1	1	0
D. dianthicola	0	1	0	1	1	1	1	0	1
D. undicola	1	0	1	1	1	1	1	0	1
D. lacustris	1	0	0	0	0	1	1	0	2
D. aquatica	1	0	0	1	0	1	1	0	2
D. chrysanthemi	0	0	0	1	1	1	1	0	1
D. paradisiaca	0	0	0	0	1	0	0	0	0
D. poaceiphila	0	0	0	0	1	1	0	1	0
P. atrosepticum	0	0	0	1	0	1	1	0	1
P. carotovorum	1	1	0	1	0	1	1	0	1
P. parmentieri	0	0	0	1	0	1	1	0	1
P. polaris	0	0	0	1	0	1	1	0	1

524
The strains used in this study are *D. dadantii* 3937, *D. aquatica* 174/2, *D. chrysanthemi*

ATCC 11663, *D. dadantii* subsp *dieffenbachiae* NCPPB 2976, *D. dianthicola* NCPPB 453, *D.*

fanghzongdai DSM 101947, *D. lacustris* S29, *D. oryzae* ZYY5, *D. paradisiaca* ATCC 33242,

D. poaceiphila NCPPB 569, *D. solani* IPO 2222, *D. undicola* 2B12, *D. zeae* NCPPB 2538, *P.*

atrosepticum ATCC 33260, *P. carotovorum* subsp. *carotovorum* ATCC 15713, *P.*

parmentieri RNS08.42.1A and *P. polaris* NIBIO 1006. The presence and number of proteins

detected by search of the corresponding gene in the genome in each strain is indicated.

Table 2: Oligonucleotides and strains used in this study

Oligonucleotides

Line	Name	Sequence
536	17176H+	CCTCCTGAGATTAGAGAGAG
537	17176A	CGCGCCGGTGTTTTTCTTGCG
538	17176H-	GGTTAGTGATGGTGATGGTGATGCTGAATATTGAGCGACGTGC
539	17176XbaF	GAACAGGATGTGTCCTCTTCTAGACACAAAGCGCTGCGTGCG
540	17176XbaR	CGCACGCAGCGCTTTGTGTCTAGAAGAGGACACATCCTGTTC
541	15544L2+	CTAAGAATCAGTCAGTTTGCG
542	15544L2-	AGGCAGATAACGCTACTCGCC
543	15544H+	ATTTATACTCGCCACCGATGC
544	15544XmaF	GCCTTGCGAGCCCCCGCTCCCGGGCTGTCTCGGGTTACCGTG
545	15544XmaR	CACGGTAACCCGAGACAGCCCGGGAGCGGGGGCTCGCAAGGC
546	15544H-	GGTTAGTGATGGTGATGGTGATGTTTAATATAGGTCTGTGTGAAC
547	14642H+	GGTGAGGAATAATTCTGGCC
548	14642H-	GGTTAGTGATGGTGATGGTGATGAGGCAGTTGTACTTTTCCAG
549	VirKH+	TGCCGTATGTGATAGTCACG
550	VirKH-	CCTTAGTGATGGTGATGGTGATGTTGCTTGAAGAAGCGGATATC

551
552
553   *Dickeya dadantii* strains
554

Line	Strain	Genotype	Source
555	3937	Wild type	Laboratory collection
556	A5653	*outD* ::CmR	Laboratory collection
557	A6417	*svfB* ::CmR	This work
558	A6418	*svfA* ::*uidA*-kanR	This work
559	A3838	*kdgR* ::Mu-CmR	(19)
560	A3845	*pecS*::Mu-CmR	(28)
561	A3846	*pecT*::CmR	(29)
562	A3849	*pir*::CmR	Laboratory collection
563	A4237	*gacA*::CmR	(31)
564	A6434	*kdgR*::Mu-CmR *svfA*::*uidA*-kanR	This work
565	A6435	*pecS*::Mu-CmR *svfA* :*uidA*-kanR	This work
566	A6436	*pecT*::CmR *svfA*::*uidA*-kanR	This work
567	A6436	*pir*::CmR *svfA*::*uidA*-kanR	This work

568	A6437	*gac*A::CmR *svfA*::*uidA*-kanR	This work
569	A6467	*svfB*::*uidA*-kanR	This work
570	A6469	*kdgR*::Mu-CmR *svfB*::*uidA*-kanR	This work
571	A6470	*pecS*::Mu-CmR *svfB*::*uidA*-kanR	This work
572	A6471	*pecT*::CmR *svfB*::*uidA*-kanR	This work
573	A6472	*pir*::CmR *svfB*::*uidA*-kanR	This work
574	A6473	*gacA*::CmR *svfB*::*uidA*-kanR	This work
575			
576			
577			
578			
579			

587   Fig. 1

```
SvfA          -------AQTS----DEPVQTVITALD--SPFVDYPLSAGSEQDVSSSEHKALRAPAPAL 47
SvfC          ----------------------------------------------------APAPAL 6
Bacillus      ----------------------------------------------------APAPPL 6
SvfB          GYSQTINNTDEAGPPIELG---LTAFD-SL-ESNSPLSAARDTS-RASTSGALRAPAPAL 54
Photorhabdus  --------QTEQLPKVPAAQKVLIEQSQLLPNVKSPIR-AEERD-LKIEDGSVHAPAPGL 50
                                                                  ****  *

SvfA          SSVQVYAVYSSLKGGWQAVPT-NTLSL-SGYAGGTLRIAVLEVGYGGNRIGWLNGGQTS- 104
SvfC          SGLRIKVLSGVYGGTWQYAPV-NAVSIGPGYAGGTLQIAVVETGYGGNRIGWINGEQKK- 64
Bacillus      TSLNVVKVESQL-GGVEFIGA-NNLSTVKDHGGSYLYIYTNEMGYGRNPIAQMSGQKLKK 64
SvfB          SRVTVYAVGSSN-CGWEYMTSIGQLSTTCDHGGAQLRVAVQEIGYGNNPVAWMNGGVLPR 113
Photorhabdus  TNMWVYAVGSTN-CGWEYTSNL--FATTCDHGGQQLRAAVLEIGYGYSSFAWMNGGLLPN 107
              : : :  : .      :          .:   .:.*   *   . * *** . .. :.*

SvfA          --PYQVNPVCVVSGRYTESCPAGSIVSGWMAYFNADNMSSVTFRYQSTSTNFPNRTLSTS 162
SvfC          --PDSVKLACLVKGEMTDNCPRGATGAGWIAYFSANYQTSVTFRYQSTSANFPYKTLSTS 122
Bacillus      VDS---KMI---------DINGDRTVDGWEYKWDASGQQNGQFKYQNTSTNAPWNTLFTS 112
SvfB          SANYQTDGICIVGNQYTFPCPAGYTVVGWMYYYNLDGTDNGQFKFQDTSTNAPRNTLFTQ 173
Photorhabdus  SAMYSSKTVCITNGYYTWPCTAGQTVVGYLHEYNLDGNQNGTFRYQNTSTNSPWNTMSVQ 167
                       .        .     *:  :. .   . *::*.**:* * .*: ..

SvfA          LNIQ 166
SvfC          LTIK 126
Bacillus      LNIK 116
SvfB          TYIK 177
Photorhabdus  INIL 171
                 *
```

Fig . 2

597   Fig. 3

A

B

599   Fig. 4

600