Title: Transcranial magnetic stimulation of the occipital cortex interferes with foot movements in acquired but not congenitally blind individuals

Authors: Tsuyoshi Ikegami[1,2]*, Gen Miura[1,2], Masaya Hirashima[1,2], Eiichi Naito[1,2], Satoshi Hirose[1,3]

Affiliations:

[1]Center for Information and Neural Networks (CiNet), Advanced ICT Research Institute, National Institute of Information and Communications Technology, 1-4, Yamadaoka, Suita City, Osaka, Japan

[2] Graduate School of Frontier Biosciences, Osaka University, 1-3, Yamadaoka, Suita, Osaka, Japan

[3] Otemon Gakuin University, Faculty of Psychology, 2-1-15, Nishiai, Ibaraki City, Osaka, Japan

(*corresponding author)

Corresponding author:

*Correspondence should be addressed to Tsuyoshi Ikegami (ikegami244@gmail.com).

Acknowledgments: We thank Dr. Kaoru Amano and Dr. Nobuhiro Hagura for their valuable comments on the first draft of this manuscript, Dr. Mitsuaki Takemi for his help in the TMS simulation analysis, and Dr. Keisuke Tani for his suggestion in the neck stimulation. We thank Ms. Naoko Katagiri, Dr. Kenta Yamamoto, Ms. Noriko Karasudani, Ms. Mayura Fujita, and Ms. Keiko Chiba for their help in conducting the experiments and organizing the data. We thank Mr. Shigeo Murakami and Mr. Hajime Teranishi of the Japanese Blind Football Association for his help in recruiting participants. TI was supported by JSPS KAKENHI Grant #16K12974 and JST, PRESTO #JPMJPR22S1. EN was supported by JSPS KAKENHI Grant #JP19H05723.

Competing interests:

The authors declare no competing financial interests.

Abstract

Research has shown that the occipital cortex can be reorganized and repurposed for nonvisual perception and cognitive functions following visual loss in blind individuals. However, no studies have directly examined the involvement of the visual cortex in motor function. Here, we show that a rhythmic foot movement performed by acquired blind individuals can be disrupted by transcranial magnetic stimulation (TMS) to their primary and secondary visual cortex (V1/V2). This disruptive effect was absent for congenitally blind or sighted individuals. As a control, TMS to the neck muscles of acquired blind individuals confirms that this disruptive effect was not caused solely by the sensory (tactile/auditory) sensations accompanying the stimulation. Our results provide the direct evidence that functional repurposing of the human visual cortex is not restricted to perception and cognitive functions but also extends to motor function. Moreover, our findings suggest the nessecity of the visual experience before visual loss for the functional reorganization of the visual cortex for motor function.

Significance statement

The brain can adapt remarkably after sensory loss. One striking example is that after visual loss, the visual cortex—deprived of visual input—is often reused for nonvisual perception and higher cognition. Here, we show that it can also support motor control. Using transcranial magnetic stimulation, we transiently disrupted processing in the visual cortex while participants performed rhythmic foot movements. This disruption impaired performance in people who became blind after birth, but not in sighted individuals or those blind from birth. These findings demonstrate that the visual cortex can take on a motor function after vision is lost, but only with prior visual experience.

Introduction

Sensory loss can lead to dramatic plasticity of the cerebral cortex (*1, 2*). Accumulating evidence indicates cross-modal plasticity following visual loss, where the visual cortex is reorganized to participate in the remaining sensory modalities. For example, the visual cortex of both congenitally and acquired blind individuals responds to auditory (*3-6*) and tactile stimuli (*7-11*). This cross-modal plasticity has been regarded as the neural underpinnings driving blind individuals' superior ability over sighted individuals in nonvisual perception including sound localization and tactile spatial discrimination (*12-15*). Furthermore, researchers have suggested that the visual cortex can also be reorganized to contribute to higher cognitive processes such as verbal memory, language processing, and mathematical processing (*16-20*). Thus, the visual cortex of blind

individuals takes a more prominent role in nonvisual perception and cognitive functions
compared with that of sighted individuals.

Despite this understanding, few studies have examined whether the visual cortex can be
reorganized to contribute to sensorimotor control. Previous studies have reported that the
primary visual cortex (V1) of blind individuals can be recruited for cognitive tasks
involving motor output, such as braille reading tasks (7-9). However, the recruitment of
V1 has only been attributed to their nonvisual perception or cognitive function, but not to
motor function. To our knowledge, a single study has suggested that the visual cortex may
be reorganized to participate in sensorimotor control involving the spared sensory
modalities, with blind opossums being superior to sighted ones in somatosensory-based
motor control during ladder-rung walking (21). However, direct evidence is still lacking
in humans and other animals. To directly investigate whether the visual cortex can be
reorganized for sensorimotor control, we applied transcranial magnetic stimulation
(TMS) to the occipital cortex, including the primary and secondary visual cortex (V1/V2),
of blind participants during a rhythmic foot movement.

There is another debate on differences in the cross-modal plasticity between acquired
blinds who lost their vision after birth and congenitally blind individuals who never have
experienced vision (2, 22, 23). Studies have suggested that the neuronal structure (24-26)
and activation patterns (9, 27-29) of the visual cortex as well as the behavioral
performance (30-32) differ between acquired and congenitally blinds. Particularly for
sensorimotor control, we expected that the reorganization process for recruiting the visual
cortex may differ between acquired blinds with experience in visuomotor control and
congenitally blinds without it. Therefore, we examined the above question separately for
congenital and acquired blind participants.

We observed that TMS to the occipital cortex increased the variability of the foot
movement in acquired blind participants but not in congenitally blind participants or
sighted controls. Furthermore, we demonstrated that the disruptive effect was not
observed when we applied TMS to the blinds' neck muscle of acquired blind individuals,
which induced tactile/auditory sensations similar to those induced by stimulation to the
occipital cortex without affecting neural processing. This evidence suggests that the
tactile/auditory sensations accompanying TMS alone were not sufficient to cause this
disruptive effect. Our findings suggest that a reorganization of the visual cortex in blind
individuals contributes to sensorimotor control, requiring some visual expererience

before they lose vision.

Results

Experiment-1

Twelve acquired blind individuals, 12 congenitally blind individuals, and 12 sighted individuals participated in the experiment (see Table 1 for participant characteristics). Blindfolded participants in all groups performed rhythmic movements of alternating dorsiflexion and plantar flexion of the feet (Fig. 1A) without any auditory cue. The participants were instructed to maintain their movement frequency (1 Hz) as constant as possible. We chose this task to minimize the influence of differences in visual experience on task performance. In daily life, sighted individuals often control foot movements without visual input, unlike upper limb movements. Moreover, the task requirement—rhythmic dorsiflexion–plantarflexion—can be easily understood without reference to the visual coordinate system. Therefore, task performance was expected to be minimally affected by the amount of the visual experience, allowing a fair comparison among the groups.

During the movement, 20 single TMS pulses were applied to one of 14 stimulation sites over the occipital cortex (Fig. 1B) at 2–4 s interstimulus intervals in the stimulation condition (see **Methods**), while no stimulation was applied in the no-stimulation condition. We assessed the effects of TMS on the rhythmic foot movement by quantifying the variability of the movement frequency with the SD of the cycle duration (Fig. 1A, see **Methods**). We did not observe any immediate motor effects of TMS, such as muscle twitching or movement stops. The main target area in this study was the early visual cortex. Thus, in the main text, we focused on stimulation site #11 (see Fig. 1B), which corresponds to the putative V1/V2 (pV1/V2; see Supplementary Information). Supplementary Figure S1 reports the average SD over all stimulation sites, covering broader visual areas.

Initially, we confirmed that all groups of participants successfully performed the instructed movement. The average movement frequencies over all trials in the stimulation and no-stimulation conditions were similar (0.94 ± 0.17 Hz, 1.00 ± 0.24 Hz and 0.97 ± 0.16 Hz for the acquired blind, congenitally blind and sighted groups, respectively; $F_{2,32}$ = 0.82, p = 0.45, η_p^2 = 0.05).

However, we observed distinct group differences in the effect of TMS on movement

performance. SD of the cycle duration in stimulation-to-pV1/V2 condition was higher
when compared with no stimulation condition in the acquired blind group, resulting in
the positive value of ΔSD (Increases in the SD from the no-stim condition to the
stimulation-to-pV1/V2 condition; blue box plot in Fig 2A). This increase in the SD was
not observed in the congenitally blind (green) or sighted groups (red). Wilcoxon signed-
rank test revealed a significant increase of SD in the acquired blind group (p = 0.028 with
Bonferroni correction for the three tests, $W = 64$, $r = 0.821$), but not in the congenitally
blind group (p = 0.320, $W = -28$, $r = -0.424$) or the sighted group (p = 0.622, $W = 4$, $r = 0.051$) (Fig. 2A and Table 2).
0.051) (Fig. 2A and Table 2).
150
In the acquired blind group, the increase in SD with TMS was not correlated with their
age (Speaman's correlation, r = -0.432, p = 0.160), age at the onset of visual loss (r = -0.233, p = 0.467), or duration of visual loss (r = -0.245, p = 0.442). Moreover, this increase
0.233, p = 0.467), or duration of visual loss (r = -0.245, p = 0.442). Moreover, this increase
in SD did not differ between the blind athlete (ABL[A] in Table 2) and blind nonathlete
subgroups (ABL[NA]; Mann-Whitney U test, p = 0.937 without correction, $U = 17$, $r = 0.056$). Furthermore, the increase in SD was not dependent on the presence or absence
0.056). Furthermore, the increase in SD was not dependent on the presence or absence
of light sensitivity (p = 0.202 without correction, $U = 14.5$, $r = -0.486$, Fig. S2A).
158
159 Finally, we compared the average SD over all stimulation sites with SD in the no-
160 stimulation condition for each group. Again, we found a significant increase in the
161 acquired blind group (p = 0.020 with Bonferroni correction for the three tests, $W = 62$, $r = 0.795$), but not in the congenitally blind group (p = 0.520, $W = -26$, $r = 0.333$) or the
162 = 0.795), but not in the congenitally blind group (p = 0.520, $W = -26$, $r = 0.333$) or the
163 sighted group (p = 0.677, $W = -26$, $r = -0.333$) (Fig. S1 and Table S2).
164

Experiment-2
One may argue that the increase in the SD in the acquired blind group of Experiment-1
can be caused by the sensory side effects of TMS, such as tactile sensations or a clicking
sound (Duecker and Sack, 2015). To ensure that this was not the case, we conducted
Experiment-2 by applying TMS to a neck muscle, the right spelenius capitis, which
induces tactile/auditory sensations similar to those accompanying TMS to pV1/V2,
without affecting any neural processing. We stimulated the neck muscle while 11
acquired-blind participants performed the same foot movement task as in Experiment-1
(stimulation-to-neck condition). The TMS intensity was adjusted for each participant to
feel the same level of tactile sensations as those induced by TMS to the pV1/V2 (see
Method). The variability in movement frequency under stimulation-to-neck conditions
was compared to that in the absence of stimulation (no-stimulation condition).

The average movement frequency over all trials in the stimulation-to-neck and no-

stimulation conditions was 1.012 ± 0.122 Hz, suggesting that the participants successfully

performed the instructed movement. Importantly, SD of the cycle duration in stimulation-

to-neck condition (0.087 ± 0.021) was almost identical to SD in no-stimulation condition

(0.088 ± 0.022) and there was no significant increase in the SD from the no-stim condition

to the stimulation-to-neck condition (Fig. 2B, Wilcoxon signed-rank test, p = 0.638, $W =$

-20, $r = -0.303$).

Discussion

Overview of findings

Experiment 1 demonstrated that TMS to pV1/V2 disrupts a foot movement task in

acquired blind individuals but not in congenitally blind or sighted individuals. Experiment

2 demonstrated that TMS to the neck of acquired blind individuals did not affect the

movement task, suggesting that the sensory (tactile/auditory) side effects of TMS alone

could not explain the disruptive effect observed in Experiment 1. Thus, our results provide

the direct neurobehavioral evidence that the visual cortex of blind individuals contributes

to sensorimotor control. However, this reorganization may require some visual experience

before visual loss. Numerous studies have reported that reorganization in the visual cortex

of blind individuals participates in nonvisual perception and cognitive functions (*2, 22,*

34). Our finding extends the knowledge on the reorganization of the visual cortex in blind

individuals. Specifically, the visual cortex can be reorganized not only for perceptual and

cognitive functions but also for motor function.

Potential confounders

First, we shall discuss possible confounders of our results. One may argue that the

disruptive effect of TMS (i.e., the increase in SD in Experiment 1) was attributable to

advanced foot motor control rather than to acquired blindness, because six out of 11

participants were experts in the sensorimotor control of foot movements (i.e., current or

former members of the Japanese national blind football team, ABL[A]). In fact, the foot

movement in the ABL[A] was more stable than the other groups, including the ABL[NA],

congenitally blind, or sighted groups, both under no-stimulation and stimulation-to-

pV1/V2 conditions (Table 2). This was also observed when TMS was applied to the other

stimulation sites (Table S2), indicating the strong effect of the sports experience on

movement variability; this could probably be attributed to their intensive (blind football)

exercises, which required controlling their foot much more accurately than in daily life.

Importantly, despite the different variability in each condition, a disruptive TMS effect
(ΔSD in Table 2) was similarly observed both in the ABL^A and in the ABL^{NA} subgroups.
Therefore, the disruptive effect of TMS observed in the acquired blind participants can
not be explained by the inclusion of athletes, whose SD was lower in general. Furthermore,
the movement variability in the no-stimulation condition did not account for individual
differences in the disruptive effect in the congenitally blind and sighted groups (see
Supplementary Information).
220
Another concern is that the disruptive TMS effect observed in the acquired blind group
of Experiment 1 may be attributed to sensory side effects of TMS, such as tactile (pain)
sensations or a clicking sound (35). However, Experiment 2 demonstrated it unlikely. In
Experiment 2, neck stimulation did not induce disruptive effects in acquired blind
participants, although the intensity of the tactile sensations and the volume of the clicking
sound were similar to those induced by TMS to pV1/V2. (see Methods). Thus, our results
suggest that the disruptive effect observed in Experiment 1 is primarily attributed to the
neural stimulation on the brain function rather than the sensory side effects from TMS.
Therefore, we reject the possible confounding factors and argue that the pV1/V2 of the
acquired blind participants engaged in the foot movement.
231
This neck stimulation may serve as a novel and useful control stimulus during motor tasks,
particularly for continuous movements. Typically, in motor TMS studies, the visual cortex
is often chosen as control stimulation sites because it has been considered minimally
involved in motor execution. However, recent findings suggest that the visual cortex in
sighted individuals may contribute to motor control (33), and that plasticity mechanisms
in motor and visual areas may interact (36), indicating that visual cortex stimulation may
not always be appropriate as a control condition. Furthermore, although sham stimulation
is effective for discrete movements, rapidly completed movements, it poses challenges
for continuous motor tasks such as those used in the present study. Participants may
experience prediction errors in the absence of expected tactile sensations, which can lead
them to perceive issues with the experimental procedure and, in some cases, discontinue
the task. Therefore, neck stimulation with carefully controlled levels of pain and tactile
intensity could provide a novel and effective approach for assessing sensory side effects
induced by TMS.
246

Effects of visual experience
The disruptive effect was observed in all acquired blind participants (Fig. S2B), including

five early-blind (age < 6 years), three intermediate-blind ($6 \leq$ age ≤ 16 years), and four
late-blind participants (age > 16 years), with the exception of a late blind (#8). We did not
find a significant correlation between the SD increase and age at onset of blindness.
Therefore, the amount of visual experience did not appear to affect the degree of visual
cortex engagement. This is consistent with the previous studies reporting the lack of
relationship between amount of the visual experience and perception/cognitive functions
(*20, 27, 28, 37, 38*).
256
In contrast to the acquired blind participants, the congenital blind participants did not
show any clear disruptive effect of TMS to pV1/V2 (Fig. 2A, Table 2, and S1), suggesting
that the visual experience is necessary for the engagement of the visual cortex in
sensorimotor control. Visually guided movements appear early in human development, at
around 6–8 months of age (*39-41*). This suggests that the neural connections between the
visual cortex and sensorimotor regions, crucial for visually-guided motor control (*33*),
are formed very early during development. Recent work in mice (*42*), as well as humans
(*43, 44*), suggests that normal visuomotor experiences are necessary for the typical
development of this connection, even although a part of the connection can be
inherent (*45*). In our acquired blind participants, such neural connection for visuomotor
control may be (at least somewhat) normally established at least at a very early stage of
development. Then, following visual loss, these existing neural connections may be
enhanced and reorganized (*1, 2, 46-48*) for sensorimotor control of nonvisually-guided
movements in acquired blinds' brain. On the other hand, such neural connections can not
be developed in the congenitally blind individuals without any visuomotor experience.
These might be the reason why we observed the disruptive effect of TMS to pV1/V2 on
the rhythmic feet movement in the acquired blind participants, irrespective of their age at
onset of blindness, but not congenitally blind participants.
275

Functional roles of the visual cortex
Although the current study could not identify the exact functional role of the acquired
blind participants' visual cortex in our motor task, we can hypothesize possible roles
based on the computational understanding of sensorimotor control (*49*). First, the visual
cortex almost certainly does not contribute to the generation of motor commands, since
TMS did not induce immediate motor effects such as muscle twitching or movement stop.
Instead, the increased movement variability observed may indicate that TMS affects
higher-order aspects of sensorimotor control such as temporal control processing or
feedback control processing. Specifically, we speculate that TMS affected the online

estimation of the body state (i.e., position, velocity, or movement phase of the foot), which

is transformed into motor commands through motor regions (*49-52*). This state estimation

is thought to be achieved by combining bottom-up sensory feedback and top-down

sensory prediction–predicting the sensory consequences of motor commands (*53-55*). In

the brain of sighted individuals, the visual cortex may contribute to the online estimation

in the visual domain. Recent studies in both sighted mice and humans support the role of

the visual cortex in predicting the visual consequences of movements (e.g., optical flow)

through top-down connections from motor regions to the visual cortex (*42, 56-58*). This

prediction is combined with visual feedback to achieve an online estimation of the body

state. Similarly, in acquired blind individuals, the visual cortex may contribute to the state

estimation by combining a nonvisual feedback signal and sensory prediction. This is

supported by two lines of evidence: 1) acquired blind individuals maintain or even

enhance connectivity between the visual cortex and sensorimotor regions following visual

loss (*46-48, 59*) and 2) the visual cortex in blind individuals processes nonvisual sensory

information (*1, 2*). Therefore, TMS to pV1/V2 may disrupt the online estimation of the

body state, increasing movement variability in acquired blind individuals.

Methodological limitation and future directions

There is a methodological limitation in our study. We determined the stimulation sites

(see Methods) without using a magnetic resonance imaging (MRI)-guided navigation

system (*17*). Thus, between-participant differences may exist in the anatomical location

of stimulation sites. However, we focused on V1/V2; its stimulation site (#11) was

determined based on a well-established external anatomical landmark (2.5 cm above inion

(*60-63*)). Further, we performed a post-hoc simulation analysis based on MRI anatomical

data. This simulation supported that our TMS procedure could prominently stimulate the

V1/V2 (see Supplementary Information). Therefore, we are confident that we stimulated

V1/V2 when applying TMS to stimulation site #11. On the other hand, the disruptive

TMS effect may not be restricted to early visual cortices. The analysis of the mean TMS

effect across all sites (Fig. S1, Table S1) showed a significant effect in which TMS

increased the movement variability only in the acquired blind participants. Further

research is needed to examine the possibility that a broader neuronal network, including

higher-order visual cortices, reorganizes to contribute to sensorimotor control in blind

individuals.

Conclusion

Our study provide the direct neurobehavioral evidence that the visual cortex of blind

individuals contributes to sensorimotor control. Our findings indicate that the human
brain's plasticity following sensory loss is more flexible than previously thought, i.e.,
functional repurposing of the lower sensory cortices is not restricted to perception and
cognitive functions but also extends to motor function. This plasticity may increase the
neural resources available for sensorimotor control in blind individuals, helping them
interact with the everchanging and diverse environment.
327
Methods
Participants

Eighteen acquired blind (i.e. lost vision after birth, #1–#12, #37–#42 in Table 1 and 3),
12 congenitally blind (i.e. no visual experience, #13–#24) and 12 sighted participants
(#25–#36) participated in either or both of the two experiments: Experiment-1 and
Experiment-2. All participants were healthy volunteers without a history of cognitive
impairment or psychiatric disorders. Congenital blindness was defined as total blindness
(vision limited to light perception or worse) diagnosed before one year of age, whereas
acquired blindness was defined as the same condition diagnosed at or after one year of
age. All congenitally blind participants reported that they had never experienced color or
form vision since birth. They presented no contraindications to TMS, which was assessed
using a screening questionnaire in compliance with the guidelines for noninvasive
magnetic brain stimulation in research applications (*64*). The causes of blindness for all
blind participants were related to lesions in the prechiasmatic visual system. These
experiments were approved by the ethics committee of the National Institute of
Information and Communications Technology and conducted according to the
Declaration of Helsinki. All participants provided written informed consent before
participating in the experiment and were naïve to the purpose of the study.
346
Experiment 1: Twelve acquired blind (acqureid blind group: ABL in Table 1), 12
congenitally blind (congenitally blind group: CBL) and 12 sighted (sighed group: SI)
individuals with normal or corrected-to-normal vision participated in Experiment-1. Age
was similar between the acquired blind participants (age 27.50 ± 8.37, mean $\pm$ standard
deviation (SD) across participants) and sighed participants (27.75 ± 7.24), while both
were younger significantly than the congenitally blind participants (41.00 ± 7.01; $F_{2,32} =$
11.74, $p = 1.50 \times 10^{-4}$, $\eta_p^2 = 0.42$). The acquired blind group comprised two subgroups:
athlete (ABL^A) and nonathlete (ABL^{NA}). The six blind participants in the ABL^A subgroup
were current or former members of the Japanese national blind football team. The
remaining six in the ABL^{NA} subgroup were blind individuals from the general population.

One congenitally blind participant (#12) was former members of the Japanese national blind football team, while the other participants in the CBL were from the general population. All sighted participants belonged to the general population. We determined the sample size (n = 12 per group) based on the effect of TMS on motor tasks assessed by previous studies (*65-68*).

Experiment 2: Eleven acquired blind individuals (see Table 3), including two athletes (#2, #6) and three nonatheletes (#7, #8, #12), who participated in Experiment 1, and six nonathletes who did not participate in Experiment-1 (#37–#42), took part in Experiment 2. We determined a sample size (n = 11) similar to Experiment-1.

Apparatus

Experiment 1: All participants wore eye patches and an eye mask to eliminate any possible visual stimuli ~30 min before the experiment and continued to wear them throughout the experiment. Furthermore, they were instructed to close their eyes during the experiment. This was to ensure that the sighted participants performed the task without any visual information as did blind participants. They sat in a comfortable reclining chair with their heads on a chinrest and legs extended on a leg rest. The height and location of the chin and leg rests were adjusted for each participant to comfortably make the rhythmic movements of alternating dorsiflexion and plantar flexion of their feet. An electromagnetic position sensor (Micro Sensor 1.8 Extra Flex, Polhemus Liberty, Burlington, VT) recorded the six dimensional kinematic data (position: x, y, and z; orientation: azimuth, elevation, and roll) of the first metatarsals of both feet. The obtained data were digitized with a temporal sampling ratio of 240 Hz and then low-pass filtered at a cutoff frequency of 5 Hz.

Electromyographic (EMG) activity was recorded at 2,000 Hz utilizing surface electrodes from the medial and lateral heads of the gastrocnemius and tibialis anterior of both legs using wireless EMG sensors (Wave Plus Wireless EMG, Cometa, Bareggio, Italy). This EMG recording was prepared to capture possible muscle twitches with TMS. However, since we did not observe any twitches, EMG data were not reported in this study.

TMS stimulation was applied using a 70-mm figure-eight coil and a Magstim Rapid Transcranial Magnetic Stimulator (Magstim Company, Spring Gardens, UK). The maximum stimulator output was 2.1 T. TMS pulse delivery was controlled by an in-house program running on MATLAB, version R2013b (The MathWorks, Natick, MA).

Experiment 2: The experimental apparatus was the same as Experiment1 except for the following two points. First, as all participants were blind, they neither wore eye patches nor eye masks. Second, EMG activity was not measured.

TMS protocol

Experiment 1: Fourteen stimulation sites were arranged in a 2 × 7 grid and numbered from left to right and top to bottom (Fig. 1B); these sites were identified for each participant using an elastic swimming cap with 15 markers indicating the inion position and the 14 stimulation sites. The marker positions were determined using a cap worn by a male individual with an average head size for Japanese men (head circumference: ~57 cm (*69*)) who did not participate in the experiment. The markers were placed at inion, 2.5 cm (stimulation site #11), and 5 cm (stimulation site #4) dorsal from the inion on the median line (nasion-to-inion line through the vertex), and 2, 4, and 6 cm right and left away from the median line at the same level as stimulation sites #4 and #11. Each participant wore the swimming cap on the head with 1) the inion marker placed at the participant's inion, 2) stimulation sites #4 and #11 aligned on the participant's median line, and 3) stimulation site #11 placed at 2.5 cm dorsal from the inion. Thus, stimulation site #11, the main target in this study, was well controlled among participants (on the median line and 2.5 cm dorsal from the inion). On the other hand, the other stimulation sites could not be strictly controlled.

Subsequently, the stimulation intensity for each participant was determined using the highest value that did not induce scalp pain (Table 1). Single TMS pulses were applied to each stimulation site (sites #1–#14 in order). For stimulation site #1, stimulations were applied by gradually increasing the intensity by 5% starting from 60% intensity of the maximum stimulator output until the participant reported any pain. Then, the intensity was decreased by 5% until the pain disappeared. The last intensity was defined as tentative intensity. When the participant did not report pain with a stimulation at 100% intensity, the tentative intensity was set to 100%. For other stimulation sites, we began stimulation with the tentative intensity. When the participant reported pain, we decreased it by 5% until pain disappeared and updated the tentative intensity. This procedure was repeated for all sites up to site #14.

The TMS intensity used in the experiment was comparable between the acquired blind (77.08% ± 11.97% of the maximum stimulator output), congenital (72.73% ± 16.18%),

and sighted (84.58% ± 9.64%) groups ($F_{2,32} = 2.56$, $p = 0.09$, $\eta_p^2 = 0.14$).

Experiment 2: TMS was applied to a neck muscle. The stimulation site was on the right

spelenius capitis, ~2 cm superior from the horizontal line passing through the 7th spinal

process. The TMS intensity for the neck (I_{Neck}) was adjusted for each participant to feel

the same level of tactile sensations as induced by TMS to pV1/V2 (stimulation site #11).

I_{Neck} determination consisted of four steps. In STEP 1, we determined the intensity of the

stimulation to pV1/V2 as in Experiment-1 (I_{Brain}). Then, we provided three single-pulse

stimulations to pV1/V2 with an inter-pulse interval of ~1s and asked the participant to

remember the tactile sensation induced by TMS to pV1/V2 with I_{Brain}. In STEP 2, we

applied TMS to the neck muscle to determine I_{Neck}. We gradually increased the intensity

by 5% starting from 40% of the maximum stimulator output until the participant felt a

tactile sensation comparable to that induced by TMS to pV1/V2. The last intensity was

defined as I_{Neck}. In STEP 3, to confirm that the tactile sensations induced by these two

stimulations were comparable, we again provided one single-pulse stimulation to pV1/V2

with I_{Brain} followed by one single-pulse stimulation to the neck with I_{Neck}. When the

participants confirmed that the sensations were comparable, the determination procedure

was completed. Otherwise, we moved to STEP4; when reporting that the tactile sensation

of the neck stimulation was stronger or weaker, we decreased or increased I_{Neck} by 5%

until reaching a comparable level of tactile sensation to that induced by TMS of the

pV1/V2 with I_{Brain}. Then, we went back to STEP 3 (confirmation process). STEP 4 and

STEP 3 were performed until the participants confirmed that the sensations were

comparable.

The TMS intensity of the pV1/V2 was similar between participants in Experiment 2

(I_{Brain}: 73.18% ± 20.28%) and the acquired blind group in Experiment 1 (77.083% ±

11.968%). Slightly weaker stimulations induced the same level of tactile sensations when

applying TMS to the neck muscle (I_{Neck}: 66.82% ± 19.27%). The loudness (sound

pressure level, SPL) of the clicking sound induced by TMS to the neck (69.79 ± 0.44 dB;

mean ± SD of ten measurements) was comparable to that induced by TMS to pV1/V2

(72.67 ± 0.84 dB). SPL was measured by a sensor placed around a participant's right ear

while receiving TMS to the neck with average intensity in Experiment 2 (67%) or to

pV1/V2 with average intensity in Experiment-1 (77% for the blind group).

Task procedure

Experiment 1: The participants were blindfolded and while sitting on the chair, made

rhythmic alternative movements of dorsiflexion and plantar flexion of the feet without any sound cue. To maintain the movement frequency as constant as possible, the participants practiced the movements with a sound frequency of 1 Hz for ~2 min twice: once before the determination process of the stimulation intensity and once before starting the experiment. They were instructed to maintain the consistent movement frequency throughout the experiment. No instructions were provided regarding the motion range of the foot movement.

In each experimental trial, the rhythmic movement was performed under each of the following three conditions. During the stimulation condition, 20 single TMS pulses were applied to one of the 14 stimulation sites at interstimulus intervals of 2–4 s (drawn from a uniform random distribution). The TMS coil was held tangentially to the scalp. The trial was started 3 s before the first stimulation and ended 3 s after the last. Each trial lasted 56.34–70.26 s. During the sham condition, a fake coil was placed over stimulation site #11, while another coil delivered 20 single TMS pulses in the air near the fake coil. Thus, the participant could feel the clicking sound coming from the fake coil on the scalp. The duration and timing of TMS pulses were determined according to the protocol used in the stimulation condition. During the no-stimulation condition, no coil was placed on the scalp, and TMS pulses were not generated. The trial duration was set at 50 s.

The participants completed two blocks of 15 trials, including 14 stimulation trials (one for each of the 14 stimulation sites) and one sham trial. The trial order was pseudorandomized for each block. One no-stimulation trial was performed before the first block and after the second block. Thus, the participants completed a total of 32 experimental trials.

Experiment 2: The procedure of Experiment-2 was similar to Experiment 1. The participants underwent the same ~2 min practice with a 1 Hz sound, before and after the determination process of the stimulation intensity. The participants were instructed to maintain the foot movement frequency at 1 Hz throughout the experiment without specific instructions on the motion range. Then, the participants completed two blocks of 10 trials each. For each trial, they performed the same task as Experiment-1. In each block, five odd-numbered trials were in the stimulation-to-neck condition, and five even-numbered trials were in the no-stimulation condition. In total, the participants performed 10 stimulation trials and 10 no-stimulation trials in total.

In the stimulation condition, the TMS pulse was applied to the neck muscle (the right spelenius capitis). Interstimulus interval, number of single TMS pulse, and trial's start and end points were identical to the stimulation condition in Experiment 1. As the neck is concave, we could not adhere the coil to the skin surface of the stimulation site. Thus, we held the coil so that the center position was 2–3 cm away from the skin surface. During the no-stimulation condition, no coil was placed on the neck, and the TMS pulses were not generated. The trial duration was set at 50 s, as for the no-stimulation condition in Experiment 1.

Data analysis

Experiment 1: The effects of TMS on movement were examined by evaluating movement variability. For each foot in each trial, we initially identified the duration of each movement cycle, defined as the time between the adjacent dorsiflexion peaks (Fig. 1A and see details in the next paragraph), and calculated the SD of the durations from all movement cycles. Then, we obtained the SD for each stimulation site by averaging the four SD values for left and right foot in the two stimulation trials (first and second blocks). Similarly, the SD in the no-stimulation trials was calculated as control. Our main analysis focused on stimulation site #11, pV1/V2 (see TMS protocol). The other stimulation sites might have covered broad visual areas, including early (V1/V2) and higher-order (V3, V4, and V5) visual areas, according to the literature (*61, 63, 70, 71*). To observe the overall trend of the TMS effect on early and higher-order visual cortices, we analyzed the average SDs of all stimulation sites as shown in Fig. S1.

The dorsiflexion peaks (Fig. 1A) for each foot in each trial were identified as follows. First, the time average of the elevation angle from the horizontal plane was calculated and subtracted from the entire time series to obtain the time series of a relative angle (θ in Fig. 1A). Then, the crossing times where θ changed from negative to positive were identified. Finally, the point of maximum value between each of two consecutive crossing times was identified as a peak. The identification code is available online (see Data availability).

Experiment 2: First, the SD was calculated in each trial as in Experiment 1. Then, for each stimulation-to-neck and no-stimulation condition, all SD values (10 trials for each condition) were averaged to obtain the SD.

Data exclusion

Data from the sham trials were not analyzed since almost all participants noticed the lack

537 of stimulation and/or felt something going wrong during the trials. Hence, the data were
538 probably biased by this surprising effect.
539
Experiment 1: Participant #17 was excluded from the analysis. In the middle of the 1st
block (after 9^{th} trial), the participant reoprted a discomfort with the position of the feet
and head and changed the position as well as removed the chin rest. Thus, we analyzed
data from 11 participants for the congenitally blind group. For stimulation site #14 of
participant #35 (Table 1) and sites #2, #4, #8, #13 of participants #19, data from only the
second block were utilized to calculate the SD owing to the lack of the first block data
due to a recording error. For stimulation site #5 of participant #20, the second block data
of the left foot were excluded as the foot posture changed significantly during the trial.
For stimulation sites #4 and #7 of participant #21, the second block data of the left foot
were excluded as he fell asleep during the trial. In total, 0.82% of the stimulation trials
were excluded from analysis, while all the no-stimulation trials were analyzed.
551
Experiment 2: For the left foot (participant #2, 8, 12, 37, 40, and 42) or both feet (#7, 38,
and 39) of some participants, the azimuth angle was used, which is the angular orientation
of the foot in the horizontal projection plane. This was because we had observed, upon
visual inspection, that their azimuth angle reflected the dorsiflexion peaks more clearly
than the elevation angle, given their internally-rotated leg posture. For participant #2
(Table 3), data from only the second block (5 trials for each condition) were utilized to
calculate the SD because of the lack of the first block data due to a recording error.
559

Statistical analysis
Because of the violation of normality, the SD was examined by the two-tailed Wilcoxon
signed rank test or the two-tailed Mann-Whitney U test for within- and between-
participant comparisons, respectively. Nonparametric Spearman's rank correlations were
used to assess the relationship of the SD with the following variables: age, age at the onset
of visual loss, or duration of visual loss. For the other data analysis, we analyzed
parametric one-way ANOVA for the three-group comparisons or two-tailed unpaired t-
tests for two-group comparisons. The Bonferroni correction was used for multiple
comparisons. All analyses were conducted using MATLAB R2024a (The MathWorks,
Natick, MA). The significance level was set at 0.05 in all the analyses. For the Wilcoxon
signed rank test, the *p* value, the test statistic *W*, and the effect size r were reported (*72*).
For the Mann-Whitney U test, the *p* value, the test statistic *U*, and the effect size r were
reported (*72*). We reported effect size with partial eta squared (η_p^2) for the ANOVAs,

Cohen's d_z for the paired *t*-test, and Cohen's d for the unpaired *t*-test (*73*). All data are
presented as mean ± SD across participants.

Experiment 1: To examine the effects of TMS on stimulation site # 11 (pV1/V2), the
Wilcoxon signed-rank test was used to test diffences in the SD between the no-stimulation
and stimulation conditions for each group. The same Wilcoxon signed-rank test was
conducted for the average SD of all stimulation sites (Fig. S1). Because the acquired blind
group showed a significant increase in SD on pV1/V2, the Spearman's rank correlation
analyses were performed to examine the relationship of the SD increase (Δ SD in Table
2; SD in stimulation-to-pV1/V2 condition minus SD in no-stimulation condition) with
age, age at onset of visual loss, and duration of visual loss. Furthermore, to evaluate the
relationship between SD increase and the participants' sports experience, the ABL[A] and
ABL[NA] subgroups were compared using the Mann-Whitney U test. To examine whether
the presence of light sensitivity affects SD increase, two subgroups of acquired blind
participants with and without light sensitivity were compared using the Spearman's
rankSpearman's rank (Fig. S2A). To examine differences in age, TMS intensity, and
movement frequency between groups, a one-way ANOVA was performed. When a
significant main effect was observed, the effect of group was examined using a unpaired
t-test.

Experiment 2: The effects of TMS to the neck muscle were compared by the SD in the
no-stimulation condition and the stimulation-to-neck condition using the Wilcoxon
signed rank.

Data availability
Data (time of the dorsiflexion peak for all participants and a sample foot movement
trajectory) and codes to reproduce Fig. 2 and S1 and the related analyses are available
from https://github.com/ikegami244/Blind-TMS. The code to identify the dorsiflexion
peaks is also available from the same repository.

Code to replicate the simulation reported in the Supplementary Information is available
at   https://github.com/satoshi-hirose/Blind-TMS.TMS-simulation-with-simNIBS.   The
anatomical MRI data are not publicly available due to ethical restrictions but are available
from the corresponding author on reasonable request.

References

References

1.   L. B. Merabet, J. F. Rizzo, A. Amedi, D. C. Somers, A. Pascual-Leone, What blindness can
tell us about seeing again: merging neuroplasticity and neuroprostheses. *Nat Rev Neurosci*
6, 71-77 (2005).

2.   L. B. Merabet, A. Pascual-Leone, Neural reorganization following sensory loss: the
opportunity of change. *Nat Rev Neurosci* **11**, 44-52 (2010).

3.   T. Kujala *et al.*, Auditory and somatosensory event-related brain potentials in early blind
humans. *Exp Brain Res* **104**, 519-526 (1995).

4.   R. Weeks *et al.*, A positron emission tomographic study of auditory localization in the
congenitally blind. *J Neurosci* **20**, 2664-2672 (2000).

5.   M. B. Amadeo, V. S. Stormer, C. Campus, M. Gori, Peripheral sounds elicit stronger
activity in contralateral occipital cortex in blind than sighted individuals. *Scientific reports*
9, 11637 (2019).

6.   P. Vetter *et al.*, Decoding Natural Sounds in Early "Visual" Cortex of Congenitally Blind
Individuals. *Curr Biol* **30**, 3039-3044 e3032 (2020).

7.   N. Sadato *et al.*, Activation of the primary visual cortex by Braille reading in blind subjects.
Nature **380**, 526-528 (1996).

8.   L. G. Cohen *et al.*, Functional relevance of cross-modal plasticity in blind humans. *Nature*
389, 180-183 (1997).

9.   H. Burton *et al.*, Adaptive changes in early and late blind: a fMRI study of Braille reading.
J Neurophysiol **87**, 589-607 (2002).

10.   P. Pietrini *et al.*, Beyond sensory images: Object-based representation in the human
ventral pathway. *Proc Natl Acad Sci U S A* **101**, 5658-5663 (2004).

11.   M. Ptito *et al.*, TMS of the occipital cortex induces tactile sensations in the fingers of blind
Braille readers. *Exp Brain Res* **184**, 193-200 (2008).

12.   N. Lessard, M. Pare, F. Lepore, M. Lassonde, Early-blind human subjects localize sound
sources better than sighted subjects. *Nature* **395**, 278-280 (1998).

13.   B. Roder *et al.*, Improved auditory spatial tuning in blind humans. *Nature* **400**, 162-166
(1999).

14.   D. Goldreich, I. M. Kanics, Tactile acuity is enhanced in blindness. *J Neurosci* **23**, 3439-
3445 (2003).

15.   P. Voss *et al.*, Early- and late-onset blind individuals show supra-normal auditory abilities
in far-space. *Curr Biol* **14**, 1734-1738 (2004).

16.   A. Amedi, N. Raz, P. Pianka, R. Malach, E. Zohary, Early 'visual' cortex activation
correlates with superior verbal memory performance in the blind. *Nat Neurosci* **6**, 758-
766 (2003).

17.   A. Amedi, A. Floel, S. Knecht, E. Zohary, L. G. Cohen, Transcranial magnetic stimulation
of the occipital pole interferes with verbal processing in blind subjects. *Nat Neurosci* **7**,
1266-1270 (2004).

18.   M. Bedny, A. Pascual-Leone, D. Dodell-Feder, E. Fedorenko, R. Saxe, Language
processing in the occipital cortex of congenitally blind adults. *Proc Natl Acad Sci U S A*
108, 4429-4434 (2011).

19.   S. Kanjlia, C. Lane, L. Feigenson, M. Bedny, Absence of visual experience modifies the
neural basis of numerical thinking. *Proc Natl Acad Sci U S A* **113**, 11172-11177 (2016).

20.   M. Amalric, I. Denghien, S. Dehaene, On the role of visual experience in mathematical
development: Evidence from blind mathematicians. *Dev Cogn Neurosci* **30**, 314-323
(2018).

21.   M. Englund, S. Faridjoo, C. S. Iyer, L. Krubitzer, Available Sensory Input Determines
Motor Performance and Strategy in Early Blind and Sighted Short-Tailed Opossums.
iScience **23**, 101527 (2020).

22.   I. Fine, J. M. Park, Blindness and Human Brain Plasticity. *Annu Rev Vis Sci* **4**, 337-356
(2018).

23.   S. Cornell Kärnekull, A. Arshamian, M. E. Nilsson, M. Larsson, From Perception to
Metacognition: Auditory and Olfactory Functions in Early Blind, Late Blind, and Sighted
Individuals. *Frontiers in Psychology* **7**, (2016).

24.   Q. Li *et al.*, Cortical thickness development of human primary visual cortex related to the
age of blindness onset. *Brain Imaging Behav* **11**, 1029-1036 (2017).

25.   J. Jiang *et al.*, Thick Visual Cortex in the Early Blind. *The Journal of Neuroscience* **29**,
2205-2211 (2009).

26.   W. Qin, Y. Xuan, Y. Liu, T. Jiang, C. Yu, Functional Connectivity Density in Congenitally
and Late Blind Subjects. *Cereb Cortex* **25**, 2507-2516 (2015).

27.   M. Bedny, A. Pascual-Leone, S. Dravida, R. Saxe, A sensitive period for language in the
visual cortex: distinct patterns of plasticity in congenitally versus late blind adults. *Brain*
Lang **122**, 162-170 (2012).

28.   H. Burton, Visual cortex activity in early and late blind people. *J Neurosci* **23**, 4005-4011
(2003).

29.   C. S. Carter, M. M. Botvinick, J. D. Cohen, The contribution of the anterior cingulate
cortex to executive processes in cognition. *Rev Neurosci* **10**, 49-57 (1999).

30.   S. M. Abel, J. C. Figueiredo, A. Consoli, C. M. Birt, B. C. Papsin, The effect of blindness
on horizontal plane sound source identification. *International journal of audiology* **41**,
285-292 (2002).

31.   F. Gougoux *et al.*, Neuropsychology: pitch discrimination in the early blind. *Nature* **430**,

309 (2004).

32.   C. Y. Wan, A. G. Wood, D. C. Reutens, S. J. Wilson, Early but not late-blindness leads to
enhanced auditory perception. *Neuropsychologia* **48**, 344-348 (2010).

33.   J. P. Gallivan, C. S. Chapman, D. J. Gale, J. R. Flanagan, J. C. Culham, Selective
Modulation of Early Visual Cortical Activity by Movement Intention. *Cereb Cortex* **29**,
4662-4678 (2019).

34.   E. Castaldi, C. Lunghi, M. C. Morrone, Neuroplasticity in adult human visual cortex.
Neurosci Biobehav Rev **112**, 542-552 (2020).

35.   F. Duecker, A. T. Sack, Rethinking the role of sham TMS. *Front Psychol* **6**, 210 (2015).

36.   I. D. Sari, C. Lunghi, Different Forms of Plasticity Interact in Adult Humans. *Eneuro* **10**,
(2023).

37.   P. Voss, F. Gougoux, M. Lassonde, R. J. Zatorre, F. Lepore, A positron emission
tomography study during auditory localization by late-onset blind individuals.
Neuroreport **17**, 383-388 (2006).

38.   C. Holig, J. Focker, A. Best, B. Roder, C. Buchel, Crossmodal plasticity in the fusiform
gyrus of late blind individuals during voice recognition. *Neuroimage* **103**, 374-382 (2014).

39.   A. L. Woodward, Infants selectively encode the goal object of an actor's reach. *Cognition*
69, 1-34 (1998).

40.   O. Braddick, J. Atkinson, B. Hood, Monocular vs binocular control of infants' reaching.
INVESTIGATIVE OPHTHALMOLOGY and VISUAL SCIENCE **37**, 1331-1331 (1996).

41.   Y. Kanakogi, S. Itakura, Developmental correspondence between action prediction and
motor ability in early infancy. *Nat Commun* **2**, 341 (2011).

42.   M. Leinweber, D. R. Ward, J. M. Sobczak, A. Attinger, G. B. Keller, A Sensorimotor
Circuit in Mouse Cortex for Visual Flow Predictions. *Neuron* **96**, 1204 (2017).

43.   C. Campus *et al.*, Sensitive period for the plasticity of alpha activity in humans. *Dev Cogn*
Neurosci **49**, 100965 (2021).

44.   O. Yizhar, Z. Tal, A. Amedi, Loss of action-related function and connectivity in the blind
extrastriate body area. *Front Neurosci* **17**, 973525 (2023).

45.   L. Bola, P. Vetter, M. Wenger, A. Amedi, Decoding Reach Direction in Early "Visual"
Cortex of Congenitally Blind Individuals. *J Neurosci* **43**, 7868-7878 (2023).

46.   M. Bedny, Evidence from Blindness for a Cognitively Pluripotent Cortex. *Trends Cogn*
Sci **21**, 637-648 (2017).

47.   M. J. Van Ackeren, F. M. Barbero, S. Mattioni, R. Bottini, O. Collignon, Neuronal
populations in the occipital cortex of the blind synchronize to the temporal dynamics of
speech. *Elife* **7**, (2018).

48.   S. Abboud, L. Cohen, Distinctive Interaction Between Cognitive Networks and the Visual

717   Cortex in Early Blind Individuals. *Cereb Cortex* **29**, 4725-4742 (2019).

718 49. S. H. Scott, Optimal feedback control and the neural basis of volitional motor control. *Nat*

719   *Rev Neurosci* **5**, 532-546 (2004).

720 50. E. Todorov, M. I. Jordan, Optimal feedback control as a theory of motor coordination. *Nat*

721   *Neurosci* **5**, 1226-1235 (2002).

722 51. J. Diedrichsen, R. Shadmehr, R. B. Ivry, The coordination of movement: optimal feedback

723   control and beyond. *Trends Cogn Sci* **14**, 31-39 (2010).

724 52. T. Takei, S. G. Lomber, D. J. Cook, S. H. Scott, Transient deactivation of dorsal premotor

725   cortex or parietal area 5 impairs feedback control of the limb in macaques. *Curr Biol* **31**,

726   1476-1487 e1475 (2021).

727 53. D. M. Wolpert, J. R. Flanagan, Motor prediction. *Curr Biol* **11**, R729-732 (2001).

728 54. R. Shadmehr, M. A. Smith, J. W. Krakauer, Error correction, sensory prediction, and

729   adaptation in motor control. *Annu Rev Neurosci* **33**, 89-108 (2010).

730 55. T. Ikegami, G. Ganesh, Shared Mechanisms in the Estimation of Self-Generated Actions

731   and the Prediction of Other's Actions by Humans. *eNeuro* **4**, (2017).

732 56. G. B. Keller, T. Bonhoeffer, M. Hubener, Sensorimotor mismatch signals in primary visual

733   cortex of the behaving mouse. *Neuron* **74**, 809-815 (2012).

734 57. A. B. Saleem, A. Ayaz, K. J. Jeffery, K. D. Harris, M. Carandini, Integration of visual

735   motion and locomotion in mouse visual cortex. *Nat Neurosci* **16**, 1864-1869 (2013).

736 58. B. Buaron, D. Reznik, R. Gilron, R. Mukamel, Voluntary Actions Modulate Perception

737   and Neural Representation of Action-Consequences in a Hand-Dependent Manner.

738   *Cereb Cortex* **30**, 6097-6107 (2020).

739 59. O. Collignon *et al.*, Impact of blindness onset on the functional organization and the

740   connectivity of the occipital cortex. *Brain* **136**, 2769-2783 (2013).

741 60. G. Beckers, S. Zeki, The consequences of inactivating areas V1 and V5 on visual motion

742   perception. *Brain* **118 (Pt 1)**, 49-60 (1995).

743 61. A. Cowey, V. Walsh, Magnetically induced phosphenes in sighted, blind and blindsighted

744   observers. *Neuroreport* **11**, 3269-3273 (2000).

745 62. R. Laycock, D. P. Crewther, P. B. Fitzgerald, S. G. Crewther, Evidence for fast signals and

746   later processing in human V1/V2 and V5/MT+: A TMS study of motion perception. *J*

747   *Neurophysiol* **98**, 1253-1262 (2007).

748 63. N. Salminen-Vaparanta, V. Noreika, A. Revonsuo, M. Koivisto, S. Vanni, Is selective

749   primary visual cortex stimulation achievable with TMS? *Hum Brain Mapp* **33**, 652-665

750   (2012).

751 64. S. Rossi, M. Hallett, P. M. Rossini, A. Pascual-Leone, T. M. S. C. G. Safety of, Safety,

752   ethical considerations, and application guidelines for the use of transcranial magnetic

stimulation in clinical practice and research. *Clin Neurophysiol* **120**, 2008-2039 (2009).

65. H. Foltys *et al.*, Motor control in simple bimanual movements: a transcranial magnetic
stimulation and reaction time study. *Clin Neurophysiol* **112**, 265-274 (2001).

66. J. J. Orban de Xivry, S. E. Criscimagna-Hemminger, R. Shadmehr, Contributions of the
motor cortex to adaptive control of reaching depend on the perturbation schedule. *Cereb*
Cortex **21**, 1475-1484 (2011).

67. J. J. Orban de Xivry *et al.*, Stimulation of the human motor cortex alters generalization
patterns of motor learning. *J Neurosci* **31**, 7102-7110 (2011).

68. F. Mawase, S. Uehara, A. J. Bastian, P. Celnik, Motor Learning Enhances Use-Dependent
Plasticity. *J Neurosci* **37**, 2673-2685 (2017).

69. M. Kouchi, M. Mochimaru. (National Institute of Advanced Industrial Science and
Technology., 2008).

70. A. Pascual-Leone, V. Walsh, Fast backprojections from the motion to the primary visual
area necessary for visual awareness. *Science* **292**, 510-512 (2001).

71. T. Amemiya, B. Beck, V. Walsh, H. Gomi, P. Haggard, Visual area V5/hMT+ contributes
to perception of tactile motion direction: a TMS study. *Scientific reports* **7**, 40937 (2017).

72. D. S. Kerby, The Simple Difference Formula: An Approach to Teaching Nonparametric
Correlation. *Comprehensive Psychology* **3**, 11.IT.13.11 (2014).

73. D. Lakens, Calculating and reporting effect sizes to facilitate cumulative science: a
practical primer for t-tests and ANOVAs. *Front Psychol* **4**, 863 (2013).

773
774

Tables

Table 1. Characteristics of the participants in Experiment 1

Participant #	Group	Age (y)	Sex	Onset of blindness (y)	Cause of Blindness	Light sensitivity at present	TMS intensity (%)
1	ABL[A]	26	m	2	Retinoblastoma	none	80
2	ABL[A]	27	m	14	retinal choroidal degeneration	yes	75
3	ABL[A]	39	m	20	morning glory anomaly	none	85
4	ABL[A]	28	m	23	Uveitis	yes	90
5	ABL[A]	20	m	2	Retinoblastoma	none	75
6	ABL[A]	40	m	25	retinitis pigmentosa	yes	85
7	ABL[NA]	25	m	14	chorioretinal atrophy	yes	70
8	ABL[NA]	25	m	5	Glaucoma	none	65
9	ABL[NA]	42	m	17	uveitis + glaucoma	occasional	70
10	ABL[NA]	19	m	2	behcet's disease	none	100
11	ABL[NA]	20	m	3	retinal cell death	none	75
12	ABL[NA]	19	m	15	Glaucoma	none	55
13	CBL	29	m	0	unknown	none	80
14	CBL	40	m	0	cataract	none	95
15	CBL	36	m	0	anophthalmia + persistent fetal vasculature	none	60
16	CBL	32	m	0	amaurosis	yes	75
17	CBL	36	f	0	crystalline lens tissue hyperplasia + retinal detachment	yes	80
18	CBL	35	m	0	Leber congenital amaurosis	none	65
19	CBL	48	m	0	retinoblastoma	none	90
20	CBL	43	m	0	unknown	yes	80
21	CBL	42	m	0	retinoblastoma	none	55
22	CBL	48	f	0	Retinopathy of prematurity	yes	80
23	CBL	48	m	0	Corneal staphyloma	none	80
24	CBL	49	f	0	Retinopathy of prematurity	none	40
25	SI	22	m	-	-	-	85
26	SI	22	m	-	-	-	85
27	SI	23	m	-	-	-	95
28	SI	22	m	-	-	-	90
29	SI	34	m	-	-	-	70
30	SI	35	m	-	-	-	85
31	SI	40	m	-	-	-	80
32	SI	26	m	-	-	-	75

33	SI	23	m	-	-	-	70
34	SI	27	m	-	-	-	100
35	SI	20	m	-	-	-	95
36	SI	39	m	-	-	-	85

778 *ABL, acquired blind; CBL, congenitally blind; SI, sighted; y, year; m, male. Superscripts*

779 *A and NA indicate athlete and nonathlete blind participants, respectively.*

781 **Table 2. SD values in no-stimulation condition and stimulation-to-pV1/V2 condition**

782 **and the increase in SD for each group in Experiment 1**

	$SD_{no\text{-}stim}$	$SD_{pV1/V2}$	ΔSD
ABL (n = 12)	**0.072 ± 0.045**	**0.085 ± 0.043**	**0.013 ± 0.017**
ABL^A (n = 6)	0.049 ± 0.013	0.063 ± 0.026	0.013 ± 0.019
ABL^{NA} (n = 6)	0.095 ± 0.055	0.108 ± 0.045	0.013 ± 0.018
CBL (n = 11)	**0.118 ± 0.077**	**0.104± 0.049**	**-0.013 ± 0.042**
SI (n = 12)	**0.106 ± 0.048**	**0.104 ± 0.043**	**-0.002 ± 0.019**

783 *ABL, acquired blind; CBL, congenitally blind; SI, sighted. Superscripts A and NA indicate*

784 *athlete and nonathlete blind participants, respectively. $\Delta SD = SD_{pV1/V2} - SD_{no\text{-}stim}$. All*

785 *data are reported as mean ± standard deviation across participants. The units are seconds.*

787 **Table 3. Characteristics of the participants in Experiment 2**

Partici pant #	Group	Age (y)	Sex	Onset of blindn ess (y)	Cause of blindness	Light sensiti vity at presen t	TMS intensity for brain (%)	TMS intensity for neck (%)
2	ABL^A	33	m	14	retinal choroidal degeneration	yes	90	70
6	ABL^A	46	m	25	retinitis pigmentosa	yes	100	100
7	ABL^{NA}	30	m	14	chorioretinal atrophy	yes	80	90
8	ABL^{NA}	30	m	5	glaucoma	none	75	60
12	ABL^{NA}	23	m	15	glaucoma	none	55	55
37	ABL^{NA}	32	m	2	retinoblastom a		100	65
38	ABL^{NA}	27	f	10	detachment of retina + cataract	yes	50	45
39	ABL^{NA}	43	m	34	medical error in cataract surgery	none	75	75

40	ABL[NA]	23	m	2	persistent fetal vasculature	none	50	50
41	ABL[NA]	36	m	11	detachment of retina	none	45	40
42	ABL[NA]	44	f	30	retinoblastoma	none	85	85

788 *ABL, acquired blind; y, year; m, male; f, femal. Superscripts A and NA indicate athlete*
789 *and nonathlete blind participants, respectively.*
790
791

792 **Table 4. SD values in no-stimulation condition (no-stim) and stimulation-to-neck**
793 **condition and the increase in SD in Experiment 2**

	$SD_{no-stim}$	SD_{neck}	ΔSD
ABL (n = 11)	0.088 ± 0.022	0.087 ± 0.021	-0.002 ± 0.009

794 *ABL, acquired blind. $\Delta SD = SD_{neck} - SD_{no-stim}$. All data are reported as mean ± standard*

795 *deviation across participants. The units are seconds.*

796
797

Fig. 1. The movement task and stimulation sites: A) The rhythmic foot movement of a
representative participant for a 10-s period is shown by the elevation angle, θ of right foot
(dark gray line) and left foot (light gray line). For each foot in each trial, θ is defined as
the angle relative to the average angle of the trial (θ = 0, dashed line). Each light-green
vertical line indicates the timing of a single transcranial magnetic stimulation (TMS)
pulse delivered at random. The cycle duration was defined as the time between the
adjacent dorsiflexion peaks. The SD of the cycle durations was calculated for each foot
in each trial (see **Methods**). B) Fourteen stimulation sites arranged in a 2 × 7 grid were
identified for each participant using the inion and the median line as reference points (see
Methods). Stimulation site #11 corresponds to putative primary and secondary visual
cortex (pV1/V2).

Fig. 2. A) Experiment 1: Increases in the SD from the no-stim condition to the stimulation-to-pV1/V2 condition in which TMS was applied to stimulation site #11 for acquired blind (blue), congenitally blind (green), and sighted (red) participants. Box plots show the median and interquartile ranges of the increase in the SD for each group. Each dot represents data from an individual participant. For the blind group, open dots and closed dots indicate the subgroups of athletes (ABL^A) and nonathletes (ABL^{NA}), respectively. The asterisk indicates $p < 0.05$ with Bonfferoni correction. B) Experiment 2: Increases in the SD from the no-stim condition to the stimulation-to-neck condition in which TMS was applied over the right spelenius capitis (see **Methods**) for acquired blind participants. Box plots show the median and interquartile ranges of the increase in the SD for each group. Each dot represents data from an individual participant. For the blind group, open dots and closed dots indicate the subgroups of athletes (ABL^A) and nonathletes (ABL^{NA}), respectively.

 SUPPLEMENTARY INFORMATION

TMS simulation to identify the stimulation site

We conducted a simulation analysis to investigate whether the TMS used in Experiment

1 could stimulate the early visual cortices (V1 and/or V2). Specifically, we used

anatomical MRI data from each participant to simulate the electrical field induced in the

visual cortex by TMS at stimulation site #11.

Participants

Fifteen (eight acquired blind and seven sighted) out of the 24 individuals from the

acquired and sighted groups who participated in Experiment 1 participated in this

simulation study. The rest of the participants were excluded because of contraindications

to MRI (one sighted and four blind) or video recording problems in the TMS experiment

(four sighted).

Procedures

We acquired T1-weighted anatomical brain images of the participants (MP-RAGE;

repetition time [TR]: 1900 ms; echo time [TE]: 2.48 ms; inversion time [TI]: 900 ms; flip

angle: 9°, field-of-view [FOV]: 256 mm × 256 mm; voxel size: 1.0 mm × 1.0 mm × 1.0

mm) with a 3.0-T SIEMENS scanner (Trio Tim, with a 32-channel head coil). Then, we

performed the following simulation analysis for each participant to identify the electrical

field induced by TMS to stimulation site #11.

Simulation

We conducted the simulation for two trials on stimulation site #11 for each participant,

following the procedure introduced by Saturnino et al. (2019). We first segmented the

MRI volume into the different head tissues, and the tissue surfaces were identified. Then,

we simulated the electrical field induced by stimulation by the TMS coil placed at 2.5 cm

dorsal from the inion (stimulation site #11). From the simulated electrical field, the

intensity of the electrical field (norm of the electrical field vector; V/m) in the cortical

gray matter was mapped on the cortical surface (Fig. S3).

The simulation for each participant requires three parameters: the location and orientation

of the TMS coil and the stimulus intensity. For the location, we first identified the inion

(Iz), the nasion (Nz), and the vertex (Cz) on the scalp surface based on the morphometrical

structure of the head surface. Then, we identified the point 2.5 cm above Iz on the circle

passing through Iz, Nz, and Cz. For the orientation, we identified the TMS coil orientation
in the actual TMS experiment by visual inspection of the video recording of the
experiment by one of the authors (TI). The coil orientation in each of the two trials for
stimulation site #11 was identified in 5º increments. The stimulus intensity used in the
simulation was identical to that used in the TMS experiment. We used the default values
for the other simulation parameters, including the tissues' electrical conductance.

We used the headreco pipeline (Nielsen et al., 2018) with Statistical Parametric Mapping
(SPM; https://www.fil.ion.ucl.ac.uk/spm/software/spm12) and Computational Anatomy
Toolbox for SPM (https://neuro-jena.github.io/cat) with default parameters for the
segmentation. We performed the simulation with simNIBS software (simNIBS 3.2;
https://simnibs.github.io/simnibs). The anatomical identification of the cortical area was
based on the human cortical parcellation provided by the Human Connectome Project
(HCP-MMP1.0; Glasser et al., 2016).

Results
We performed two analyses with the simulation data. First, we examined whether the
intensity peak of the simulated electrical field was observed in V1/V2 (Table S3). The
peak of the electrical field was observed within V1 or V2 in 14 out of 15 participants
(Table S3) in both trials. An exception was observed in Participant #2, whose peak was
in V3A (7.4 mm apart from the edge of V1).

Next, we checked whether the intensity of the electrical field induced in V1/V2 was
sufficient to affect neural processing in these regions. We regarded an electrical field >
10 V/m as sufficient to affect neural processing. The threshold was set according to
previous studies showing that neurons fire in response to an artificially induced electrical
field with ~ 10 V/m intensity (Park et al., 2013, Bonmassar et al., 2012, Chan and
Nicholson 1986). The proportion of the V1/V2 surface area where the induced electrical
field was above the threshold (stimulated region) is 81% ± 10% (mean ± standard
deviation; see Table S3).

These two results suggest that the TMS procedure used in Experiment-1 substantially
stimulated V1/V2, with sufficient stimulus intensity.

The movement variability in the no-stimulation condition did not
account for individual differences in the disruptive effect in the

congenitally blind and sighted groups

To deny the possibility that the disruptive effect of TMS (i.e., the increase in SD in Experiment 1) was attributable to advanced foot motor control rather than to acquired blindness, we have shown that both athletes (ABL^A) and non-athletes (ABL^{NA}) in the acquired blind group exhibited disruptive TMS effects (i.e., increases in SD). To further strengthen this argument, we evaluated the relationship between the original motor performance, assessed using SD in the non-stimulation condition, and the TMS effect.

Data from 23 participants in the congenitally blind and sighted groups were included in this analysis. Participants were divided into two subgroups—high SD (N = 12) and low SD (N = 11)—based on their SD values in the first no-stimulation condition conducted at the beginning of the experiment. The TMS effect was assessed by subtracting the SD of the second non-stimulation condition conducted at the end of the experiment, from the average SD of the two TMS stimulation-to-V1/V2 conditions. By using statistically independent values for subgroup assignment (SD of the first non-stimulation condition) and for the calculation of the dependent variable (SD of the second non-stimulation condition), we avoided the statistical bias known as regression to the mean.

As a result, no significant TMS effect was observed in either the high-SD ($t(11) = -0.80$, $p = 0.44$) or low-SD ($t(10) = 0.06$, $p = 0.96$) subgroups. Furthermore, no evident relationship was observed between SD in the initial non-stimulation condition (x-axis in Figure S4) and the TMS effect (y-axis).

References

Saturnino, G. B., Puonti, O., Nielsen, J. D., Antonenko, D., Madsen, K. H., & Thielscher,
A. (2019). SimNIBS 2.1: a comprehensive pipeline for individualized electric field
modelling for transcranial brain stimulation. Brain and human body modeling, Springer,
3-25.

Nielsen, J. D., Madsen, K. H., Puonti, O., Siebner, H. R., Bauer, C., Madsen, C. G.,
Saturnino, G. B., & Thielscher, A. (2018). Automatic skull segmentation from MR
images for realistic volume conductor models of the head: Assessment of the state-of-the-
art. NeuroImage, 174, 587–598.

Park, H. J., Bonmassar, G., Kaltenbach, J. A., Machado, A. G., Manzoor, N. F., & Gale,
J. T. (2013). Activation of the central nervous system induced by micro-magnetic
stimulation. Nature communications, 4(1), 1-9.

Bonmassar, G., Lee, S. W., Freeman, D. K., Polasek, M., Fried, S. I., & Gale, J. T. (2012).
Microscopic magnetic stimulation of neural tissue. Nature communications, 3(1), 1-10.

Chan, C. Y., & Nicholson, C. (1986). Modulation by applied electric fields of Purkinje
and stellate cell activity in the isolated turtle cerebellum. The Journal of physiology,
371(1), 89-114.

Fig. S1. The average SD over all stimulation sites (all sites) was compared with the SD
in the no-stimulation condition. The increases from the SD of the no-stim condition to the
average SD over all stimulation sites for acquired blind (blue), congenitally blind (green),
and sighted (red) groups in Experiment-1. Box plots show the median and interquartile
ranges of the increase in the SD for each group. Each dot represents data from an
individual participant. For the blind group, open dots and closed dots indicate the
subgroups of athletes (ABL[A]) and nonathletes (ABL[NA]), respectively. The asterisk
indicates $p < 0.05$ with Bonfferoni correction.

Fig. S2. SD of the cycle duration for the blind participants in the no-stimulation condition

(x-axis) and the stimulation-to-pV1/V2 condition (y-axis) of Experiment 1. A) For each

group of acquired (blue) and congenitally (green) blind participants, two subgroups with

and without light sensitivity (filled and open circles, respectivly) were compared. The

dashed line indicates y = x. In the acquired blind participants, TMS to pV1/V2 increased

the SD in the cycle duration relative to the no-stimulation condition in all participants

exept for one (#8, see Table 1). The increase in SD did not differ between the two

subgroups. In contrast, no clear trend was not observed among the congenitally blind

participants nor between the twosubgroups. B) Early-, intermediate-, late-acquired blind

participants (light, intermediate, and dark blue circles, respectively), and congenitally

blind participants (green circle) are compared. The dashed line indicates y = x. No clear

difference was observed among the three subgroups of the acquired blind participants.

Norm of the electrical field vector(V/m)

0   53

978
979   Fig. S3. Distribution of the electrical field intensity (norm of the electrical field vector; V/m) in
980   the cortical gray matter in a typical participant (participant #6). The intensity in the cortical gray
981   matter was mapped onto the cortical surface. The figure shows a posterior view of the cortex.
982

Fig. S4.

Relationship between performance in the non-stimulation condition and TMS effects: The performance in the non-stimulation condition was assessed using the SD from the first non-stimulation condition. TMS effects were evaluated by subtracting the SD of the second non-stimulation condition from the average of the two sets of TMS stimulation-to-V1/V2 conditions.

Table S1. SD of the cycle duration in all stimulation sites (#1-14) and the no-stimulation condition (no-stim) for the participants of Experiment-1 and the congenitally blind participants.

Gr SD #	SI (n = 12)	ABL (n = 12)	ABL^A (n = 6)	ABL^{NA} (n = 6)	CBL (n = 11)
no-stim	0.106±0.048	0.072±0.045	0.049±0.013	0.095±0.055	0.118±0.077
1	0.104±0.033	0.087±0.038	0.077±0.032	0.098±0.043	0.091±0.038
2	0.101±0.042	0.096±0.054	0.086±0.039	0.105±0.069	0.101±0.045
3	0.108±0.046	0.078±0.039	0.061±0.019	0.095±0.048	0.110±0.065
4	0.108±0.040	0.087±0.042	0.065±0.019	0.108±0.049	0.100±0.042
5	0.104±0.033	0.081±0.035	0.066±0.029	0.097±0.036	0.101±0.039
6	0.110±0.050	0.083±0.036	0.066±0.025	0.099±0.040	0.110±0.044
7	0.111±0.045	0.085±0.039	0.065±0.026	0.105±0.042	0.110±0.048
8	0.105±0.052	0.079±0.038	0.060±0.024	0.097±0.041	0.102±0.066
9	0.107±0.047	0.084±0.048	0.061±0.023	0.107±0.057	0.103±0.054
10	0.117±0.053	0.082±0.051	0.062±0.028	0.102±0.063	0.107±0.047
11 (pV1/V2)	0.104±0.043	0.085±0.043	0.063±0.026	0.108±0.045	0.104±0.049
12	0.107±0.043	0.085±0.042	0.065±0.019	0.104±0.052	0.100±0.041
13	0.105±0.042	0.080±0.032	0.061±0.016	0.099±0.034	0.108±0.055
14	0.098±0.037	0.083±0.028	0.074±0.025	0.093±0.030	0.119±0.065

Gr, group; SI, sighted; ABL, acquired blind; CBL, congenitally blind. Superscripts A and NA indicate athlete and nonathlete blind participants, respectively. All data are reported as mean ± standard deviation across participants. The units are seconds.

1000 **Table S2. Simulation results**

Participant #	Group	Peak location	proportion of stimulated region in V1/V2 (%)
1	ABL[A]	Left V1/Right V1*	88.5
2	ABL[A]	Left V3A	54.9
4	ABL[A]	Left V1	81.3
6	ABL[A]	Left V1	80.7
7	ABL[NA]	Left V1	76.5
8	ABL[NA]	Left V1	83.5
10	ABL[NA]	Right V1	91.4
11	ABL[NA]	Left V2	88.7
28	SI	Left V2/Right V1*	86.6
30	SI	Right V1	86.8
31	SI	Left V2	77.7
33	SI	Left V1	81.0
34	SI	Left V1	87.8
35	SI	Left V1/Left V2*	90.3
36	SI	Left V1	80.8

1001 * Asterisks indicate that the peaks for the first and second trials were located in different

1002 anatomical areas, where results were indicated as follows: first trial peak location / second

1003 trial peak location.

1004