1 **Title:** Transcranial magnetic stimulation of the occipital cortex interferes with foot
2 movements in acquired but not congenitally blind individuals
3

4 **Authors:** Tsuyoshi Ikegami[1,2]*, Gen Miura[1,2], Masaya Hirashima[1,2], Eiichi Naito[1,2],
5 Satoshi Hirose[1,3]
6
Affiliations:
[1]Center for Information and Neural Networks (CiNet), Advanced ICT Research Institute,
National Institute of Information and Communications Technology, 1-4, Yamadaoka,
Suita City, Osaka, Japan
[2] Graduate School of Frontier Biosciences, Osaka University, 1-3, Yamadaoka, Suita,
Osaka, Japan
[3] Otemon Gakuin University, Faculty of Psychology, 2-1-15, Nishiai, Ibaraki City, Osaka,
Japan
(*corresponding author)
16
17 **Corresponding author:**
18 *Correspondence should be addressed to Tsuyoshi Ikegami (ikegami244@gmail.com).
19

Acknowledgments: We thank Dr. Kaoru Amano and Dr. Nobuhiro Hagura for their
valuable comments on the first draft of this manuscript, Dr. Mitsuaki Takemi for his help
in the TMS simulation analysis, and Dr. Keisuke Tani for his suggestion in the neck
stimulation. We thank Ms. Naoko Katagiri, Dr. Kenta Yamamoto, Ms. Noriko Karasudani,
Ms. Mayura Fujita, and Ms. Keiko Chiba for their help in conducting the experiments and
organizing the data. We thank Mr. Shigeo Murakami and Mr. Hajime Teranishi of the
Japanese Blind Football Association for his help in recruiting participants. TI was
supported by JSPS KAKENHI Grant #16K12974 and JST, PRESTO #JPMJPR22S1. EN
was supported by JSPS KAKENHI Grant #JP19H05723.
29

30 **Competing interests:**
31 The authors declare no competing financial interests.
32

Abstract

Research has shown that the occipital cortex can be reorganized and repurposed for nonvisual perception and cognitive functions following visual loss in blind individuals. However, no studies have directly examined the involvement of the visual cortex in motor function. Here, we show that a rhythmic foot movement performed by acquired blind individuals can be disrupted by transcranial magnetic stimulation (TMS) to their primary and secondary visual cortex (V1/V2). This disruptive effect was absent for congenitally blind or sighted individuals. As a control, TMS to the neck muscles of acquired blind individuals confirms that this disruptive effect was not caused solely by the sensory (tactile/auditory) sensations accompanying the stimulation. Our results provide the direct evidence that functional repurposing of the human visual cortex is not restricted to perception and cognitive functions but also extends to motor function. Moreover, our findings suggest the nessecity of the visual experience before visual loss for the functional reorganization of the visual cortex for motor function.

Introduction

Sensory loss can lead to dramatic plasticity of the cerebral cortex (Merabet et al., 2005, Merabet and Pascual-Leone, 2010). Accumulating evidence indicates cross-modal plasticity following visual loss, where the visual cortex is reorganized to participate in the remaining sensory modalities. For example, the visual cortex of both congenitally and acquired blind individuals responds to auditory (Kujala et al., 1995, Weeks et al., 2000, Amadeo et al., 2019, Vetter et al., 2020) and tactile stimuli (Sadato et al., 1996, Cohen et al., 1997, Burton et al., 2002, Pietrini et al., 2004, Ptito et al., 2008). This cross-modal plasticity has been regarded as the neural underpinnings driving blind individuals' superior ability over sighted individuals in nonvisual perception including sound localization and tactile spatial discrimination (Lessard et al., 1998, Roder et al., 1999, Goldreich and Kanics, 2003, Voss et al., 2004). Furthermore, researchers have suggested that the visual cortex can also be reorganized to contribute to higher cognitive processes such as verbal memory, language processing, and mathematical processing (Amedi et al., 2003, Amedi et al., 2004, Bedny et al., 2011, Kanjlia et al., 2016, Amalric et al., 2018). Thus, the visual cortex of blind individuals takes a more prominent role in nonvisual perception and cognitive functions compared with that of sighted individuals.

Despite this understanding, few studies have examined whether the visual cortex can be reorganized to contribute to sensorimotor control. Previous studies have reported that the primary visual cortex (V1) of blind individuals can be recruited for cognitive tasks

involving motor output, such as braille reading tasks (Sadato et al., 1996, Cohen et al., 1997, Burton et al., 2002). However, the recruitment of V1 has only been attributed to their nonvisual perception or cognitive function, but not to motor function. To our knowledge, a single study has suggested that the visual cortex may be reorganized to participate in sensorimotor control involving the spared sensory modalities, with blind opossums being superior to sighted ones in somatosensory-based motor control during ladder-rung walking (Englund et al., 2020). However, direct evidence is still lacking in humans and other animals. To directly investigate whether the visual cortex can be reorganized for sensorimotor control, we applied transcranial magnetic stimulation (TMS) to the occipital cortex, including the primary and secondary visual cortex (V1/V2), of blind participants during a rhythmic foot movement.

There is another debate on differences in the cross-modal plasticity between acquired blinds who lost their vision after birth and congenitally blind individuals who never have experienced vision (Fine and Park, 2018, Cornell Kärnekull et al., 2016, Merabet and Pascual-Leone, 2010). Studies have suggested that the neuronal structure (Li et al., 2017, Jiang et al., 2009, Qin et al., 2015) and activation patterns (Bedny et al., 2012, Burton et al., 2002, Burton, 2003, Carter et al., 1999) of the visual cortex as well as the behavioral performance (Abel et al., 2002, Gougoux et al., 2004, Wan et al., 2010) differ between acquired and congenitally blinds. Particularly for sensorimotor control, we expected that the reorganization process for recruiting the visual cortex may differ between acquired blinds with experience in visuomotor control and congenitally blinds without it. Therefore, we examined the above question separately for congenital and acquired blind participants.

We observed that TMS to the occipital cortex increased the variability of the foot movement in acquired blind participants but not in congenitally blind participants or sighted controls. Furthermore, we demonstrated that the disruptive effect was not observed when we applied TMS to the blinds' neck muscle of acquired blind individuals, which induced tactile/auditory sensations similar to those induced by stimulation to the occipital cortex without affecting neural processing. This evidence suggests that the tactile/auditory sensations accompanying TMS alone were not sufficient to cause this disruptive effect. Our findings suggest that a reorganization of the visual cortex in blind individuals contributes to sensorimotor control, requiring some visual experiperience before they lose vision.

Results

Experiment-1

Eleven acquired blind individuals, 13 congenitally blind individuals, and 12 sighted individuals participated in the experiment (see Table 1 for participant characteristics). Blindfolded participants in all groups performed rhythmic movements of alternating dorsiflexion and plantar flexion of the feet (Fig. 1A) without any sound cue. The participants were instructed to maintain their movement frequency (1 Hz) as constant as possible. We selected this rhythmic and alternating lower limb movement because it can be performed easily without vision.

During the movement, 20 single TMS pulses were applied to one of 14 stimulation sites over the occipital cortex (Fig. 1B) at 2–4 s interstimulus intervals in the stimulation condition (see **Methods**), while no stimulation was applied in the no-stimulation condition. We assessed the effects of TMS on the rhythmic foot movement by quantifying the variability of the movement frequency with the SD of the cycle duration (Fig. 1A, see **Methods**). We did not observe any immediate motor effects of TMS, such as muscle twitching or movement stops. The main target area in this study was the early visual cortex. Thus, in the main text, we focused on stimulation site #11 (see Fig. 1B), which corresponds to the putative V1/V2 (pV1/V2; see Supplementary Information). Supplementary Figure S1 reports the average SD over all stimulation sites, covering broader visual areas.

Initially, we confirmed that all groups of participants successfully performed the instructed movement. The average movement frequencies over all trials in the stimulation and no-stimulation conditions were similar (0.93 ± 0.17 Hz, 1.04 ± 0.23 Hz and 0.97 ± 0.16 Hz for the acquired blind, congenitally blind and sighted groups, respectively; $F_{2,32}$ = 0.94, p = 0.40, η_p^2 = 0.06).

However, we observed distinct group differences in the effect of TMS on movement performance. SD of the cycle duration in stimulation-to-pV1/V2 condition was higher when compared with no stimulation condition in the acquired blind group (blue in Fig 2A), resulting in the positive value of Δ SD (Increases in the SD from the no-stim condition to the stimulation-to-pV1/V2 condition; blue box plot in Fig 2B). This increase in the SD was not observed in the congenitally blind (green) or sighted groups (red). Wilcoxon signed-rank test revealed a significant increase of SD in the acquired blind group (p = 0.041 with Bonferroni correction for the three tests, *W* = 54, *r* = 0.818), but

not in the congenitally blind group (p = 0.470, W = -22, r = -0.282) or the sighted group

(p = 0.622, W = 4, r = 0.051) (Fig. 2 and Table 2).

In the acquired blind group, the increase in SD with TMS was not correlated with their

age (Speaman's correlation, r = -0.461, p = 0.153), age at the onset of visual loss (r = -

0.242, p = 0.473), or duration of visual loss (r = -0.267, p = 0.424). Moreover, this increase

in SD did not differ between the blind athlete (ABL^A in Table 2, dashed blue lines in Fig.

2) and blind nonathlete subgroups (ABL^{NA}, solid blue lines; Mann-Whitney U test, p =

0.931 without correction, U = 17.5, r = 0.067). Furthermore, the increase in SD was not

dependent on the presence or absence of light sensitivity (p = 0.178 without correction,

U = 11, r = -0.533, Fig. S2A).

Finally, we compared the average SD over all stimulation sites with SD in the no-

stimulation condition for each group. Again, we found a significant increase in the

acquired blind group (p = 0.041 with Bonferroni correction for the three tests, W = 64, r

= 0.970), but not in the congenitally blind group (p = 0.677, W = 18, r = 0.231) or the

sighted group (p = 0.677, W = -26, r = -0.333) (Fig. S1 and Table S2).

Experiment-2

One may argue that the increase in the SD in the acquired blind group of Experiment-1

can be caused by the sensory side effects of TMS, such as tactile sensations or a clicking

sound (Duecker and Sack, 2015). To ensure that this was not the case, we conducted

Experiment-2 by applying TMS to a neck muscle, the right spelenius capitis, which

induces tactile/auditory sensations similar to those accompanying TMS to pV1/V2,

without affecting any neural processing. We stimulated the neck muscle while 11

acquired-blind participants performed the same foot movement task as in Experiment-1

(stimulation-to-neck condition). The TMS intensity was adjusted for each participant to

feel the same level of tactile sensations as those induced by TMS to the pV1/V2 (see

Method). The variability in movement frequency under stimulation-to-neck conditions

was compared to that in the absence of stimulation (no-stimulation condition).

The average movement frequency over all trials in the stimulation-to-neck and no-

stimulation conditions was 1.012 ± 0.122 Hz, suggesting that the participants successfully

performed the instructed movement. Importantly, SD of the cycle duration (Figure 3A) in

stimulation-to-neck condition (0.087 ± 0.021) was almost identical to SD in no-

stimulation condition (0.088 ± 0.022) and there was no significant increase in the SD

from the no-stim condition to the stimulation-to-neck condition (Fig. 3B, Wilcoxon
signed-rank test, p = 0.638, W = -20, r = -0.303).
178
Discussion
Overview of findings
With Experiment-1, we demonstrated that TMS to pV1/V2 disrupts a foot movement task
in acquired blind individuals but not in congenitally blind or sighted individuals. With
Experiment-2, we demonstrated that TMS to the neck of acquired blind individuals did
not affect the movement task, suggesting that the sensory (tactile/auditory) side effects of
TMS alone could not explain the disruptive effect observed in Experiment-1. Thus, our
results provide the direct neurobehavioral evidence that the visual cortex of blind
individuals contributes to sensorimotor control. However, this reorganization may require
some visual experience before visual loss. Numerous studies have reported that
reorganization in the visual cortex of blind individuals participates in nonvisual
perception and cognitive functions (Merabet and Pascual-Leone, 2010, Fine and Park,
2018, Castaldi et al., 2020). Our finding extends the knowledge on the reorganization of
the visual cortex in blind individuals. Specifically, the visual cortex can be reorganized
not only for perceptual and cognitive functions but also for motor function.
194
Potential confounders
First, we shall discuss possible confounders for our results. The disruptive TMS effect
(i.e., increase in the SD) in Experiment-1 could be observed only in the acquired blind
group because five out of 11 participants were experts in the sensorimotor control of foot
movements (i.e., current or former members of the Japanese national blind football team,
ABL[A]). In fact, the foot movement in the ABL[A] was more stable than the other groups,
including the ABL[NA], congenitally blind, or sighted groups, both under no-stimulation
and stimulation-to-pV1/V2 conditions (Table 2). This was also observed when TMS was
applied to any other stimulation sites (Table S2), indicating the strong effect of the sports
experience on movement variability; this could probably be attributed to their intensive
(blind football) exercises, which required controlling their foot much more accurately
than daily life. Importantly, despite the different variability in each condition, a disruptive
TMS effect (Δ SD in Table 2) was similarly observed both in the ABL[A] and in the ABL[NA]
subgroups. Therefore, the disruptive effect of TMS observed in the acqured blind
participants can not be explained by the inclusion of athletes, whose SD was lower in
general.

Another concern is that the disruptive TMS effect observed in the acquired blind group
of Experiment-1 may be attributed to sensory side effects of TMS, such as tactile (pain)
sensations or a clicking sound (Duecker and Sack, 2015). However, Experiment-2
demonstrated it unlikely. In Experiment-2, neck stimulation did not induce disruptive
effects in acquired blind participants, although the intensity of the tactile sensations and
the volume of the clicking sound were similar to those induced by TMS to pV1/V2. (see
Methods). Thus, our results suggest that the disruptive effect observed in Experiment-1
is primarily attributed to the neural stimulation on the brain function rather than the
sensory side effects from TMS. Therefore, we reject the possible confounding factors and
argue that the pV1/V2 of the acquired blind participants engaged in the foot movement.
222

Effects of visual experience
The disruptive effect was observed in all acquired blind participants (Fig. S2B), including
four early-blind (age < 6 years), three intermediate-blind (6 ≤ age ≤ 16 years), and four
late-blind participants (age > 16 years), with the exception of a late blind (#8). We did not
find a significant correlation between the SD increase and age at onset of blindness.
Therefore, the amount of visual experience did not appear to affect the degree of visual
cortex engagement. This is consistent with the previous studies reporting the lack of
relationship between amount of the visual experience and perception/cognitive functions
(Burton, 2003, Voss et al., 2006, Bedny et al., 2012, Holig et al., 2014, Amalric et al.,
2018).

In contrast to the acquired blind participants, the congenital blind participants did not
show any clear disruptive effect of TMS to pV1/V2 (Fig. 2, Table 2, and S1), suggesting
that the visual experience is necessary for the engagement of the visual cortex in
sensorimotor control. Visually guided movements appear early in human development, at
around 6–8 months of age (Woodward, 1998, Braddick, 1996, Kanakogi and Itakura,
2011). This suggests that the neural connections between the visual cortex and
sensorimotor regions, crucial for visually-guided motor control (Gallivan et al., 2019),
are formed very early during development. Recent work in mice (Leinweber et al., 2017),
as well as humans (Campus et al., 2021, Yizhar et al., 2023), suggests that normal
visuomotor experiences are necessary for the typical development of this connection,
even although a part of the connection can be inherent (Bola et al., 2023). In our acquired
blind participants, such neural connection for visuomotor control may be (at least
somewhat) normally established at least at a very early stage of development. Then,

following visual loss, these existing neural connections may be enhanced and reorganized
(Merabet et al., 2005, Merabet and Pascual-Leone, 2010, Bedny, 2017, Van Ackeren et
al., 2018, Abboud and Cohen, 2019) for sensorimotor control of nonvisually-guided
movements in acquired blinds' brain. On the other hand, such neural connections can not
be developed in the congenitally blind individuals without any visuomotor experience.
These might be the reason why we observed the disruptive effect of TMS to pV1/V2 on
the rhythmic feet movement in the acquired blind participants, irrespective of their age at
onset of blindness, but not congenitally blind participants.
255
Functional roles of the visual cortex
Although the current study could not identify the exact functional role of the acquired
blind participants' visual cortex in our motor task, we can hypothesize possible roles
based on the computational understanding of sensorimotor control (Scott, 2004). First,
the visual cortex almost certainly does not contribute to the generation of motor
commands, since TMS did not induce immediate motor effects such as muscle twitching
or movement stop. Instead, the increased movement variability observed may indicate
that TMS affects higher-order aspects of sensorimotor control such as temporal control
processing or feedback control processing. Specifically, we speculate that TMS affected
the online estimation of the body state (i.e., position, velocity, or movement phase of the
foot), which is transformed into motor commands through motor regions (Todorov and
Jordan, 2002, Scott, 2004, Diedrichsen et al., 2010, Takei et al., 2021). This state
estimation is thought to be achieved by combining bottom-up sensory feedback and top-
down sensory prediction–predicting the sensory consequences of motor commands
(Wolpert and Flanagan, 2001, Shadmehr et al., 2010, Ikegami and Ganesh, 2017). In the
brain of sighted individuals, the visual cortex may contribute to the online estimation in
the visual domain. Recent studies in both sighted mice and humans support the role of the
visual cortex in predicting the visual consequences of movements (e.g., optical flow)
through top-down connections from motor regions to the visual cortex (Keller et al., 2012,
Saleem et al., 2013, Leinweber et al., 2017, Buaron et al., 2020). This prediction is
combined with visual feedback to achieve an online estimation of the body state. Similarly,
in acquired blind individuals, the visual cortex may contribute to the state estimation by
combining a nonvisual feedback signal and sensory prediction. This is supported by two
lines of evidence: 1) acquired blind individuals maintain or even enhance connectivity
between the visual cortex and sensorimotor regions following visual loss (Collignon et
al., 2013, Bedny, 2017, Van Ackeren et al., 2018, Abboud and Cohen, 2019) and 2) the
visual cortex in blind individuals processes nonvisual sensory information (Merabet et al.,

2005, Merabet and Pascual-Leone, 2010). Therefore, TMS to pV1/V2 may disrupt the
online estimation of the body state, increasing movement variability in acquired blind
individuals.

Methodological limitation and future directions

There is a methodological limitation in our study. We determined the stimulation sites
(see Methods) without using a magnetic resonance imaging (MRI)-guided navigation
system (Amedi et al., 2004). Thus, between-participant differences may exist in the
anatomical location of stimulation sites. However, we focused on V1/V2; its stimulation
site (#11) was determined based on a well-established external anatomical landmark (2.5
cm above inion; Beckers and Zeki, 1995, Cowey and Walsh, 2000, Laycock et al., 2007,
Salminen-Vaparanta et al., 2012). Further, we performed a post-hoc simulation analysis
based on MRI anatomical data. This simulation supported that our TMS procedure could
prominently stimulate the V1/V2 (see Supplementary Information). Therefore, we are
confident that we stimulated V1/V2 when applying TMS to stimulation site #11. On the
other hand, the disruptive TMS effect may not be restricted to early visual cortices. The
analysis of the mean TMS effect across all sites (Fig. S1, Table S1) showed a significant
effect in which TMS increased the movement variability only in the acquired blind
participants. Further research is needed to examine the possibility that a broader neuronal
network, including higher-order visual cortices, reorganizes to contribute to sensorimotor
control in blind individuals.

Conclusion

Our study provide the direct neurobehavioral evidence that the visual cortex of blind
individuals contributes to sensorimotor control. Our findings indicate that the human
brain's plasticity following sensory loss is more flexible than previously thought, i.e.,
functional repurposing of the lower sensory cortices is not restricted to perception and
cognitive functions but also extends to motor function. This plasticity may increase the
neural resources available for sensorimotor control in blind individuals, helping them
interact with the everchanging and diverse environment.

Methods

Participants

Seventeen acquired blind (i.e. lost vision after birth, #1–#11, #37–#42 in Table 1 and 3),
13 congenitally blind (i.e. no visual experience, #12–#24) and 12 sighted participants
(#25–#36) participated in either or both of the two experiments: Experiment-1 and

Experiment-2; all participants were healthy volunteers without a history of cognitive impairment or psychiatric disorders. They presented no contraindications to TMS, which was assessed using a screening questionnaire in compliance with the guidelines for noninvasive magnetic brain stimulation in research applications (Rossi et al., 2009). The causes of blindness for all blind participants were related to lesions in the prechiasmatic visual system. These experiments were approved by the ethics committee of the National Institute of Information and Communications Technology and conducted according to the Declaration of Helsinki. All participants provided written informed consent before participating in the experiment and were naïve to the purpose of the study.

Experiment-1: Eleven acquired blind (acqureid blind group: ABL in Table 1), 13 congenitally blind (congenitally blind group: CBL) and 12 sighted (sighed group: SI) individuals with normal or corrected-to-normal vision participated in Experiment-1. Age was similar between the acquired blind participants (age 28.18 ± 8.42, mean $\pm$ standard deviation (SD) across participants) and sighed participants (27.75 ± 7.24), while both were younger significantly than the congenitally blind participants (39.16 ± 9.06; $F_{2,32} = 7.25$, $p = 0.003$, $\eta_p^2 = 0.312$). The acquired blind group comprised two subgroups: athlete (ABL^A) and nonathlete (ABL^{NA}). The six blind participants in the ABL^A subgroup were current or former members of the Japanese national blind football team. The remaining six in the ABL^{NA} subgroup were blind individuals from the general population. One congenitally blind participant (#12) was former members of the Japanese national blind football team, while the other participants in the CBL were from the general population. All sighted participants belonged to the general population. We determined the sample size (n = 11 or 12 per group) based on the effect of TMS on motor tasks assessed by previous studies (Foltys et al., 2001, Orban de Xivry et al., 2011a, Orban de Xivry et al., 2011b, Mawase et al., 2017).

Experiment-2: Eleven acquired blind individuals (see Table 3), including two athletes (#2, #6) and three nonatheletes (#6, #7, #11), who participated in Experiment-1, and six nonathletes who did not participate in Experiment-1 (#37–#42), took part in Experiment-2. We determined a sample size (n = 11) similar to Experiment-1.

Apparatus

Experiment-1: All participants wore eye patches and an eye mask to eliminate any possible visual stimuli ~30 min before the experiment and continued to wear them throughout the experiment. Furthermore, they were instructed to close their eyes during

the experiment. This was to ensure that the sighted participants performed the task without any visual information as did blind participants. They sat in a comfortable reclining chair with their heads on a chinrest and legs extended on a leg rest. The height and location of the chin and leg rests were adjusted for each participant to comfortably make the rhythmic movements of alternating dorsiflexion and plantar flexion of their feet. An electromagnetic position sensor (Micro Sensor 1.8 Extra Flex, Polhemus Liberty, Burlington, VT) recorded the six dimensional kinematic data (position: x, y, and z; orientation: azimuth, elevation, and roll) of the first metatarsals of both feet. The obtained data were digitized with a temporal sampling ratio of 240 Hz and then low-pass filtered at a cutoff frequency of 5 Hz.

Electromyographic (EMG) activity was recorded at 2,000 Hz utilizing surface electrodes from the medial and lateral heads of the gastrocnemius and tibialis anterior of both legs using wireless EMG sensors (Wave Plus Wireless EMG, Cometa, Bareggio, Italy). This EMG recording was prepared to capture possible muscle twitches with TMS. However, since we did not observe any twitches, EMG data were not reported in this study.

TMS stimulation was applied using a 70-mm figure-eight coil and a Magstim Rapid Transcranial Magnetic Stimulator (Magstim Company, Spring Gardens, UK). The maximum stimulator output was 2.1 T. TMS pulse delivery was controlled by an in-house program running on MATLAB, version R2013b (The MathWorks, Natick, MA).

Experiment-2: The experimental apparatus was the same as Experiment-1 except for the following two points. First, as all participants were blind, they neither wore eye patches nor eye masks. Second, EMG activity was not measured.

TMS protocol

Experiment-1: Fourteen stimulation sites were arranged in a 2 × 7 grid and numbered from left to right and top to bottom (Fig. 1B); these sites were identified for each participant using an elastic swimming cap with 15 markers indicating the inion position and the 14 stimulation sites. The marker positions were determined using a cap worn by a male individual with an average head size for Japanese men (head circumference: ~57 cm; Kouchi and Mochimaru, 2008) who did not participate in the experiment. The markers were placed at inion, 2.5 cm (stimulation site #11), and 5 cm (stimulation site #4) dorsal from the inion on the median line (nasion-to-inion line through the vertex), and 2, 4, and 6 cm right and left away from the median line at the same level as stimulation

sites #4 and #11. Each participant wore the swimming cap on the head with 1) the inion

marker placed at the participant's inion, 2) stimulation sites #4 and #11 aligned on the

participant's median line, and 3) stimulation site #11 placed at 2.5 cm dorsal from the

inion. Thus, stimulation site #11, the main target in this study, was well controlled among

participants (on the median line and 2.5 cm dorsal from the inion). On the other hand, the

other stimulation sites could not be strictly controlled.

Subsequently, the stimulation intensity for each participant was determined using the

highest value that did not induce scalp pain (Table 1). Single TMS pulses were applied to

each stimulation site (sites #1–#14 in order). For stimulation site #1, stimulations were

applied by gradually increasing the intensity by 5% starting from 60% intensity of the

maximum stimulator output until the participant reported any pain. Then, the intensity

was decreased by 5% until the pain disappeared. The last intensity was defined as

tentative intensity. When the participant did not report pain with a stimulation at 100%

intensity, the tentative intensity was set to 100%. For other stimulation sites, we began

stimulation with the tentative intensity. When the participant reported pain, we decreased

it by 5% until pain disappeared and updated the tentative intensity. This procedure was

repeated for all sites up to site #14.

The TMS intensity used in the experiment was comparable between the acquired blind

(77.27% ± 12.52% of the maximum stimulator output), congenital (72.92% ± 15.44%),

and sighted (84.58% ± 9.64%) groups ($F_{2,32}$ = 2.56, p = 0.09, η_p^2 = 0.14).

Experiment-2: TMS was applied to a neck muscle. The stimulation site was on the right

spelenius capitis, ~2 cm superior from the horizontal line passing through the 7th spinal

process. The TMS intensity for the neck (I_{Neck}) was adjusted for each participant to feel

the same level of tactile sensations as induced by TMS to pV1/V2 (stimulation site #11).

I_{Neck} determination consisted of four steps. In STEP 1, we determined the intensity of the

stimulation to pV1/V2 as in Experiment-1 (I_{Brain}). Then, we provided three single-pulse

stimulations to pV1/V2 with an inter-pulse interval of ~1s and asked the participant to

remember the tactile sensation induced by TMS to pV1/V2 with I_{Brain}. In STEP 2, we

applied TMS to the neck muscle to determine I_{Neck}. We gradually increased the intensity

by 5% starting from 40% of the maximum stimulator output until the participant felt a

tactile sensation comparable to that induced by TMS to pV1/V2. The last intensity was

defined as I_{Neck}. In STEP 3, to confirm that the tactile sensations induced by these two

stimulations were comparable, we again provided one single-pulse stimulation to pV1/V2

with I_{Brain} followed by one single-pulse stimulation to the neck with I_{Neck}. When the
participants confirmed that the sensations were comparable, the determination procedure
was completed. Otherwise, we moved to STEP4; when reporting that the tactile sensation
of the neck stimulation was stronger or weaker, we decreased or increased I_{Neck} by 5%
until reaching a comparable level of tactile sensation to that induced by TMS of the
pV1/V2 with I_{Brain}. Then, we went back to STEP 3 (confirmation process). STEP 4 and
STEP 3 were performed until the participants confirmed that the sensations were
comparable.
435
The TMS intensity of the pV1/V2 was similar between participants in Experiment-2
(I_{Brain}: 73.18% ± 20.28%) and the acquired blind group in Experiment-1 (77.27% ±
12.52%). Slightly weaker stimulations induced the same level of tactile sensations when
applying TMS to the neck muscle (I_{Neck}: 66.82% ± 19.27%). The loudness (sound
pressure level, SPL) of the clicking sound induced by TMS to the neck (69.79 ± 0.44 dB;
mean ± SD of ten measurements) was comparable to that induced by TMS to pV1/V2
(72.67 ± 0.84 dB). SPL was measured by a sensor placed around a participant's right ear
while receiving TMS to the neck with average intensity in Experiment-2 (67%) or to
pV1/V2 with average intensity in Experiment-1 (77% for the blind group).
445

Task procedure
Experiment-1: The participants were blindfolded and while sitting on the chair, made
rhythmic alternative movements of dorsiflexion and plantar flexion of the feet without
any sound cue. To maintain the movement frequency as constant as possible, the
participants practiced the movements with a sound frequency of 1 Hz for ~2 min twice:
once before the determination process of the stimulation intensity and once before starting
the experiment. They were instructed to maintain the consistent movement frequency
throughout the experiment. No instructions were provided regarding the motion range of
the foot movement.
455

In each experimental trial, the rhythmic movement was performed under each of the
following three conditions. During the stimulation condition, 20 single TMS pulses were
applied to one of the 14 stimulation sites at interstimulus intervals of 2–4 s (drawn from
a uniform random distribution). The TMS coil was held tangentially to the scalp. The trial
was started 3 s before the first stimulation and ended 3 s after the last. Each trial lasted
56.34–70.26 s. During the sham condition, a fake coil was placed over stimulation site
#11, while another coil delivered 20 single TMS pulses in the air near the fake coil. Thus,

the participant could feel the clicking sound coming from the fake coil on the scalp. The

duration and timing of TMS pulses were determined according to the protocol used in the

stimulation condition. During the no-stimulation condition, no coil was placed on the

scalp, and TMS pulses were not generated. The trial duration was set at 50 s.

The participants completed two blocks of 15 trials, including 14 stimulation trials (one

for each of the 14 stimulation sites) and one sham trial. The trial order was

pseudorandomized for each block. One no-stimulation trial was performed before the first

block and after the second block. Thus, the participants completed a total of 32

experimental trials.

Experiment-2: The procedure of Experiment-2 was similar to Experiment-1. The

participants underwent the same ~2 min practice with a 1 Hz sound, before and after the

determination process of the stimulation intensity. The participants were instructed to

maintain the foot movement frequency at 1 Hz throughout the experiment without

specific instructions on the motion range. Then, the participants completed two blocks of

10 trials each. For each trial, they performed the same task as Experiment-1. In each block,

five odd-numbered trials were in the stimulation-to-neck condition, and five even-

numbered trials were in the no-stimulation condition. In total, the participants performed

10 stimulation trials and 10 no-stimulation trials in total.

In the stimulation condition, the TMS pulse was applied to the neck muscle (the right

spelenius capitis). Interstimulus interval, number of single TMS pulse, and trial's start

and end points were identical to the stimulation condition in Experiment-1. As the neck

is concave, we could not adhere the coil to the skin surface of the stimulation site. Thus,

we held the coil so that the center position was 2–3 cm away from the skin surface. During

the no-stimulation condition, no coil was placed on the neck, and the TMS pulses were

not generated. The trial duration was set at 50 s, as for the no-stimulation condition in

Experiment-1.

Data analysis

Experiment-1: The effects of TMS on movement were examined by evaluating movement

variability. For each foot in each trial, we initially identified the duration of each

movement cycle, defined as the time between the adjacent dorsiflexion peaks (Fig. 1A

and see details in the next paragraph), and calculated the SD of the durations from all

movement cycles. Then, we obtained the SD for each stimulation site by averaging the

four SD values for left and right foot in the two stimulation trials (first and second blocks). Similarly, the SD in the no-stimulation trials was calculated as control. Our main analysis focused on stimulation site #11, pV1/V2 (see TMS protocol). The other stimulation sites might have covered broad visual areas, including early (V1/V2) and higher-order (V3, V4, and V5) visual areas, according to the literature (Cowey and Walsh, 2000, Pascual-Leone and Walsh, 2001, Salminen-Vaparanta et al., 2012, Amemiya et al., 2017). To observe the overall trend of the TMS effect on early and higher-order visual cortices, we analyzed the average SDs of all stimulation sites as shown in Fig. S1.

The dorsiflexion peaks (Fig. 1A) for each foot in each trial were identified as follows. First, the time average of the elevation angle from the horizontal plane was calculated and subtracted from the entire time series to obtain the time series of a relative angle (θ in Fig. 1A). Then, the crossing times where θ changed from negative to positive were identified. Finally, the point of maximum value between each of two consecutive crossing times was identified as a peak. The identification code is available online (see Data availability).

Experiment-2: First, the SD was calculated in each trial as in Experiment-1. Then, for each stimulation-to-neck and no-stimulation condition, all SD values (10 trials for each condition) were averaged to obtain the SD.

Data exclusion

Data from the sham trials were not analyzed since almost all participants noticed the lack of stimulation and/or felt something going wrong during the trials. Hence, the data were probably biased by this surprising effect.

Experiment-1: Participant #17 was excluded from the analysis. In the middle of the 1st block (after 9th trial), the participant reoprted a discomfort with the position of the feet and head and changed the position as well as removed the chin rest. Thus, we analyzed data from 11 participants for the congenitally blind group. For stimulation site #14 of participant #35 (Table 1) and sites #2, #4, #8, #13 of participants #19, data from only the second block were utilized to calculate the SD owing to the lack of the first block data due to a recording error. For stimulation site #5 of participant #20, the second block data of the left foot were excluded as the foot posture changed significantly during the trial. For stimulation sites #4 and #7 of participant #21, the second block data of the left foot were excluded as he fell asleep during the trial. In total, 0.82% of the stimulation trials were excluded from analysis, while all the no-stimulation trials were analyzed.

Experiment-2: For the left foot (participant #2, 8, 12, 37, 40, and 42) or both feet (#7, 38, and 39) of some participants, the azimuth angle was used, which is the angular orientation of the foot in the horizontal projection plane. This was because we had observed, upon visual inspection, that their azimuth angle reflected the dorsiflexion peaks more clearly than the elevation angle, given their internally-rotated leg posture. For participant #2 (Table 3), data from only the second block (5 trials for each condition) were utilized to calculate the SD because of the lack of the first block data due to a recording error.

Statistical analysis

Because of the violation of normality, the SD was examined by the two-tailed Wilcoxon signed rank test or the two-tailed Mann-Whitney U test for within- and between-participant comparisons, respectively. Nonparametric Spearman's rank correlations were used to assess the relationship of the SD with the following variables: age, age at the onset of visual loss, or duration of visual loss. For the other data analysis, we analyzed parametric one-way ANOVA for the three-group comparisons or two-tailed unpaired t-tests for two-group comparisons. The Bonferroni correction was used for multiple comparisons. All analyses were conducted using MATLAB R2024a (The MathWorks, Natick, MA). The significance level was set at 0.05 in all the analyses. For the Wilcoxon signed rank test, the p value, the test statistic W, and the effect size r were reported (Kerby, 2014). For the Mann-Whitney U test, the p value, the test statistic U, and the effect size r were reported (Kerby, 2014). We reported effect size with partial eta squared (η_p^2) for the ANOVAs, Cohen's d_z for the paired *t*-test, and Cohen's d for the unpaired *t*-test (Lakens, 2013). All data are presented as mean ± SD across participants.

Experiment-1: To examine the effects of TMS on stimulation site # 11 (pV1/V2), the Wilcoxon signed-rank test was used to test diffences in the SD between the no-stimulation and stimulation conditions for each group. The same Wilcoxon signed-rank test was conducted for the average SD of all stimulation sites (Fig. S1). Because the acquired blind group showed a significant increase in SD on pV1/V2, the Spearman's rank correlation analyses were performed to examine the relationship of the SD increase (Δ SD in Table 2; SD in stimulation-to-pV1/V2 condition minus SD in no-stimulation condition) with age, age at onset of visual loss, and duration of visual loss. Furthermore, to evaluate the relationship between SD increase and the participants' sports experience, the ABL[A] and ABL[NA] subgroups were compared using the Mann-Whitney U test. To examine whether

the presence of light sensitivity affects SD increase, two subgroups of acquired blind participants with and without light sensitivity were compared using the Spearman's rankSpearman's rank (Fig. S2A). To examine differences in age, TMS intensity, and movement frequency between groups, a one-way ANOVA was performed. When a significant main effect was observed, the effect of group was examined using a unpaired t-test.

Experiment-2: The effects of TMS to the neck muscle were compared by the SD in the no-stimulation condition and the stimulation-to-neck condition using the Wilcoxon signed rank.

Data availability

Data (time of the dorsiflexion peak for all participants and a sample foot movement trajectory) and codes to reproduce Fig. 2, 3, and S1 and the related analyses are available from https://github.com/ikegami244/Blind-TMS. The code to identify the dorsiflexion peaks is also available from the same repository.

Code to replicate the simulation reported in the Supplementary Information is available at https://github.com/satoshi-hirose/Blind-TMS.TMS-simulation-with-simNIBS. The anatomical MRI data are not publicly available due to ethical restrictions but are available from the corresponding author on reasonable request.

References

References

ABBOUD, S. & COHEN, L. 2019. Distinctive Interaction Between Cognitive Networks and the
Visual Cortex in Early Blind Individuals. *Cereb Cortex,* 29, 4725-4742.

ABEL, S. M., FIGUEIREDO, J. C., CONSOLI, A., BIRT, C. M. & PAPSIN, B. C. 2002. The effect
of blindness on horizontal plane sound source identification. *Int J Audiol,* 41, 285-92.

AMADEO, M. B., STORMER, V. S., CAMPUS, C. & GORI, M. 2019. Peripheral sounds elicit
stronger activity in contralateral occipital cortex in blind than sighted individuals. *Sci Rep,*
9, 11637.

AMALRIC, M., DENGHIEN, I. & DEHAENE, S. 2018. On the role of visual experience in
mathematical development: Evidence from blind mathematicians. *Dev Cogn Neurosci,* 30,
314-323.

AMEDI, A., FLOEL, A., KNECHT, S., ZOHARY, E. & COHEN, L. G. 2004. Transcranial
magnetic stimulation of the occipital pole interferes with verbal processing in blind
subjects. *Nat Neurosci,* 7, 1266-70.

AMEDI, A., RAZ, N., PIANKA, P., MALACH, R. & ZOHARY, E. 2003. Early 'visual' cortex
activation correlates with superior verbal memory performance in the blind. *Nat Neurosci,*
6, 758-66.

AMEMIYA, T., BECK, B., WALSH, V., GOMI, H. & HAGGARD, P. 2017. Visual area V5/hMT+
contributes to perception of tactile motion direction: a TMS study. *Sci Rep,* 7, 40937.

BEDNY, M. 2017. Evidence from Blindness for a Cognitively Pluripotent Cortex. *Trends Cogn*
Sci, 21, 637-648.

BEDNY, M., PASCUAL-LEONE, A., DODELL-FEDER, D., FEDORENKO, E. & SAXE, R. 2011.
Language processing in the occipital cortex of congenitally blind adults. *Proc Natl Acad*
Sci U S A, 108, 4429-34.

BEDNY, M., PASCUAL-LEONE, A., DRAVIDA, S. & SAXE, R. 2012. A sensitive period for
language in the visual cortex: distinct patterns of plasticity in congenitally versus late blind
adults. *Brain Lang,* 122, 162-70.

BOLA, L., VETTER, P., WENGER, M. & AMEDI, A. 2023. Decoding Reach Direction in Early
"Visual" Cortex of Congenitally Blind Individuals. *J Neurosci,* 43, 7868-7878.

BUARON, B., REZNIK, D., GILRON, R. & MUKAMEL, R. 2020. Voluntary Actions Modulate
Perception and Neural Representation of Action-Consequences in a Hand-Dependent
Manner. *Cereb Cortex,* 30, 6097-6107.

BURTON, H. 2003. Visual cortex activity in early and late blind people. *J Neurosci,* 23, 4005-11.

BURTON, H., SNYDER, A. Z., CONTURO, T. E., AKBUDAK, E., OLLINGER, J. M. &
RAICHLE, M. E. 2002. Adaptive changes in early and late blind: a fMRI study of Braille
reading. *J Neurophysiol,* 87, 589-607.

CAMPUS, C., SIGNORINI, S., VITALI, H., DE GIORGIS, V., PAPALIA, G., MORELLI, F. &
GORI, M. 2021. Sensitive period for the plasticity of alpha activity in humans. *Dev Cogn*
Neurosci, 49, 100965.

CARTER, C. S., BOTVINICK, M. M. & COHEN, J. D. 1999. The contribution of the anterior
cingulate cortex to executive processes in cognition. *Rev Neurosci,* 10, 49-57.

CASTALDI, E., LUNGHI, C. & MORRONE, M. C. 2020. Neuroplasticity in adult human visual
cortex. *Neurosci Biobehav Rev,* 112, 542-552.

COHEN, L. G., CELNIK, P., PASCUAL-LEONE, A., CORWELL, B., FALZ, L., DAMBROSIA,
J., HONDA, M., SADATO, N., GERLOFF, C., CATALA, M. D. & HALLETT, M. 1997.
Functional relevance of cross-modal plasticity in blind humans. *Nature,* 389, 180-3.

COLLIGNON, O., DORMAL, G., ALBOUY, G., VANDEWALLE, G., VOSS, P., PHILLIPS, C.
& LEPORE, F. 2013. Impact of blindness onset on the functional organization and the
connectivity of the occipital cortex. *Brain,* 136, 2769-83.

CORNELL KäRNEKULL, S., ARSHAMIAN, A., NILSSON, M. E. & LARSSON, M. 2016. From
Perception to Metacognition: Auditory and Olfactory Functions in Early Blind, Late Blind,
and Sighted Individuals. *Frontiers in Psychology,* 7.

COWEY, A. & WALSH, V. 2000. Magnetically induced phosphenes in sighted, blind and
blindsighted observers. *Neuroreport,* 11, 3269-73.

DIEDRICHSEN, J., SHADMEHR, R. & IVRY, R. B. 2010. The coordination of movement:
optimal feedback control and beyond. *Trends Cogn Sci,* 14, 31-9.

ENGLUND, M., FARIDJOO, S., IYER, C. S. & KRUBITZER, L. 2020. Available Sensory Input
Determines Motor Performance and Strategy in Early Blind and Sighted Short-Tailed
Opossums. *iScience,* 23, 101527.

FINE, I. & PARK, J. M. 2018. Blindness and Human Brain Plasticity. *Annu Rev Vis Sci,* 4, 337-
356.

FOLTYS, H., SPARING, R., BOROOJERDI, B., KRINGS, T., MEISTER, I. G., MOTTAGHY, F.
M. & TOPPER, R. 2001. Motor control in simple bimanual movements: a transcranial
magnetic stimulation and reaction time study. *Clin Neurophysiol,* 112, 265-74.

GOLDREICH, D. & KANICS, I. M. 2003. Tactile acuity is enhanced in blindness. *J Neurosci,* 23,
3439-45.

GOUGOUX, F., LEPORE, F., LASSONDE, M., VOSS, P., ZATORRE, R. J. & BELIN, P. 2004.
Neuropsychology: pitch discrimination in the early blind. *Nature,* 430, 309.

IKEGAMI, T. & GANESH, G. 2017. Shared Mechanisms in the Estimation of Self-Generated
Actions and the Prediction of Other's Actions by Humans. *eNeuro,* 4.

JIANG, J., ZHU, W., SHI, F., LIU, Y., LI, J., QIN, W., LI, K., YU, C. & JIANG, T. 2009. Thick
Visual Cortex in the Early Blind. *The Journal of Neuroscience,* 29, 2205-2211.

KANJLIA, S., LANE, C., FEIGENSON, L. & BEDNY, M. 2016. Absence of visual experience
modifies the neural basis of numerical thinking. *Proc Natl Acad Sci U S A,* 113, 11172-
11177.
KELLER, G. B., BONHOEFFER, T. & HUBENER, M. 2012. Sensorimotor mismatch signals in
primary visual cortex of the behaving mouse. *Neuron,* 74, 809-15.
KERBY, D. S. 2014. The Simple Difference Formula: An Approach to Teaching Nonparametric
Correlation. *Comprehensive Psychology,* 3, 11.IT.3.1.
KOUCHI, M. & MOCHIMARU, M. 2008. Anthropometric database of Japanese head 2001.
National Institute of Advanced Industrial Science and Technology.
KUJALA, T., ALHO, K., KEKONI, J., HAMALAINEN, H., REINIKAINEN, K., SALONEN, O.,
STANDERTSKJOLD-NORDENSTAM, C. G. & NAATANEN, R. 1995. Auditory and
somatosensory event-related brain potentials in early blind humans. *Exp Brain Res,* 104,
519-26.
LAKENS, D. 2013. Calculating and reporting effect sizes to facilitate cumulative science: a
practical primer for t-tests and ANOVAs. *Front Psychol,* 4, 863.
LAYCOCK, R., CREWTHER, D. P., FITZGERALD, P. B. & CREWTHER, S. G. 2007. Evidence
for fast signals and later processing in human V1/V2 and V5/MT+: A TMS study of
motion perception. *J Neurophysiol,* 98, 1253-62.
LEINWEBER, M., WARD, D. R., SOBCZAK, J. M., ATTINGER, A. & KELLER, G. B. 2017. A
Sensorimotor Circuit in Mouse Cortex for Visual Flow Predictions. *Neuron,* 96, 1204.
LESSARD, N., PARE, M., LEPORE, F. & LASSONDE, M. 1998. Early-blind human subjects
localize sound sources better than sighted subjects. *Nature,* 395, 278-80.
LI, Q., SONG, M., XU, J., QIN, W., YU, C. & JIANG, T. 2017. Cortical thickness development of
human primary visual cortex related to the age of blindness onset. *Brain Imaging Behav,*
11, 1029-1036.
MAWASE, F., UEHARA, S., BASTIAN, A. J. & CELNIK, P. 2017. Motor Learning Enhances Use-
Dependent Plasticity. *J Neurosci,* 37, 2673-2685.
MERABET, L. B. & PASCUAL-LEONE, A. 2010. Neural reorganization following sensory loss:
the opportunity of change. *Nat Rev Neurosci,* 11, 44-52.
MERABET, L. B., RIZZO, J. F., AMEDI, A., SOMERS, D. C. & PASCUAL-LEONE, A. 2005.
What blindness can tell us about seeing again: merging neuroplasticity and
neuroprostheses. *Nat Rev Neurosci,* 6, 71-7.
ORBAN DE XIVRY, J. J., CRISCIMAGNA-HEMMINGER, S. E. & SHADMEHR, R. 2011a.
Contributions of the motor cortex to adaptive control of reaching depend on the
perturbation schedule. *Cereb Cortex,* 21, 1475-84.
ORBAN DE XIVRY, J. J., MARKO, M. K., PEKNY, S. E., PASTOR, D., IZAWA, J., CELNIK, P.

& SHADMEHR, R. 2011b. Stimulation of the human motor cortex alters generalization
patterns of motor learning. *J Neurosci,* 31, 7102-10.

PASCUAL-LEONE, A. & WALSH, V. 2001. Fast backprojections from the motion to the primary
visual area necessary for visual awareness. *Science,* 292, 510-2.

PIETRINI, P., FUREY, M. L., RICCIARDI, E., GOBBINI, M. I., WU, W. H., COHEN, L.,
GUAZZELLI, M. & HAXBY, J. V. 2004. Beyond sensory images: Object-based
representation in the human ventral pathway. *Proc Natl Acad Sci U S A,* 101, 5658-63.

PTITO, M., FUMAL, A., DE NOORDHOUT, A. M., SCHOENEN, J., GJEDDE, A. & KUPERS,
R. 2008. TMS of the occipital cortex induces tactile sensations in the fingers of blind
Braille readers. *Exp Brain Res,* 184, 193-200.

QIN, W., XUAN, Y., LIU, Y., JIANG, T. & YU, C. 2015. Functional Connectivity Density in
Congenitally and Late Blind Subjects. *Cereb Cortex,* 25, 2507-16.

RODER, B., TEDER-SALEJARVI, W., STERR, A., ROSLER, F., HILLYARD, S. A. & NEVILLE,
H. J. 1999. Improved auditory spatial tuning in blind humans. *Nature,* 400, 162-6.

ROSSI, S., HALLETT, M., ROSSINI, P. M., PASCUAL-LEONE, A. & SAFETY OF, T. M. S. C.
G. 2009. Safety, ethical considerations, and application guidelines for the use of
transcranial magnetic stimulation in clinical practice and research. *Clin Neurophysiol,* 120,
2008-2039.

SADATO, N., PASCUAL-LEONE, A., GRAFMAN, J., IBANEZ, V., DEIBER, M. P., DOLD, G.
& HALLETT, M. 1996. Activation of the primary visual cortex by Braille reading in blind
subjects. *Nature,* 380, 526-8.

SALEEM, A. B., AYAZ, A., JEFFERY, K. J., HARRIS, K. D. & CARANDINI, M. 2013. Integration
of visual motion and locomotion in mouse visual cortex. *Nat Neurosci,* 16, 1864-9.

SALMINEN-VAPARANTA, N., NOREIKA, V., REVONSUO, A., KOIVISTO, M. & VANNI, S.
2012. Is selective primary visual cortex stimulation achievable with TMS? *Hum Brain*
Mapp, 33, 652-65.

SCOTT, S. H. 2004. Optimal feedback control and the neural basis of volitional motor control.
Nat Rev Neurosci, 5, 532-46.

SHADMEHR, R., SMITH, M. A. & KRAKAUER, J. W. 2010. Error correction, sensory prediction,
and adaptation in motor control. *Annu Rev Neurosci,* 33, 89-108.

TAKEI, T., LOMBER, S. G., COOK, D. J. & SCOTT, S. H. 2021. Transient deactivation of dorsal
premotor cortex or parietal area 5 impairs feedback control of the limb in macaques. *Curr*
Biol, 31, 1476-1487 e5.

TODOROV, E. & JORDAN, M. I. 2002. Optimal feedback control as a theory of motor
coordination. *Nat Neurosci,* 5, 1226-35.

VAN ACKEREN, M. J., BARBERO, F. M., MATTIONI, S., BOTTINI, R. & COLLIGNON, O.

2018. Neuronal populations in the occipital cortex of the blind synchronize to the temporal
dynamics of speech. *Elife,* 7.
VETTER, P., BOLA, L., REICH, L., BENNETT, M., MUCKLI, L. & AMEDI, A. 2020. Decoding
Natural Sounds in Early "Visual" Cortex of Congenitally Blind Individuals. *Curr Biol,* 30,
3039-3044 e2.
VOSS, P., LASSONDE, M., GOUGOUX, F., FORTIN, M., GUILLEMOT, J. P. & LEPORE, F.
2004. Early- and late-onset blind individuals show supra-normal auditory abilities in far-
space. *Curr Biol,* 14, 1734-8.
WAN, C. Y., WOOD, A. G., REUTENS, D. C. & WILSON, S. J. 2010. Early but not late-blindness
leads to enhanced auditory perception. *Neuropsychologia,* 48, 344-8.
WEEKS, R., HORWITZ, B., AZIZ-SULTAN, A., TIAN, B., WESSINGER, C. M., COHEN, L. G.,
HALLETT, M. & RAUSCHECKER, J. P. 2000. A positron emission tomographic study of
auditory localization in the congenitally blind. *J Neurosci,* 20, 2664-72.
WOLPERT, D. M. & FLANAGAN, J. R. 2001. Motor prediction. *Curr Biol,* 11, R729-32.
YIZHAR, O., TAL, Z. & AMEDI, A. 2023. Loss of action-related function and connectivity in the
blind extrastriate body area. *Front Neurosci,* 17, 973525.
756
757

Tables

Table 1. Characteristics of the participants in Experiment-1

Participant #	Group	Age (y)	Sex	Onset of blindness (y)	Cause of Blindness	Light sensitivity at present	TMS intensity (%)
1	ABL[A]	26	m	2	retinoblastoma	none	80
2	ABL[A]	27	m	14	retinal choroidal degeneration	yes	75
3	ABL[A]	39	m	20	morning glory anomaly	none	85
4	ABL[A]	28	m	23	uveitis	yes	90
5	ABL[A]	40	m	25	retinitis pigmentosa	yes	85
6	ABL[NA]	25	m	14	chorioretinal atrophy	yes	70
7	ABL[NA]	25	m	5	glaucoma	none	65
8	ABL[NA]	42	m	17	uveitis + glaucoma	occasional	70
9	ABL[NA]	19	m	2	behcet's disease	none	100
10	ABL[NA]	20	m	3	retinal cell death	none	75
11	ABL[NA]	19	m	15	glaucoma	none	55
12	CBL	20	m	0	retinoblastoma	none	75
13	CBL	29	m	0	unknown	none	80
14	CBL	40	m	0	cataract	none	95
15	CBL	36	m	0	anophthalmia + persistent fetal vasculature	none	60
16	CBL	32	m	0	amaurosis	yes	75
17	CBL	36	f	0	crystalline lens tissue hyperplasia + retinal detachment	yes	80
18	CBL	35	m	0	leber congenital amaurosis	none	65
19	CBL	48	m	0	retinoblastoma	none	90
20	CBL	43	m	0	unknown	yes	80
21	CBL	42	m	0	retinoblastoma	none	55
22	CBL	48	f	0	retinopathy of prematurity	yes	80
23	CBL	48	m	0	corneal staphyloma	none	80
24	CBL	49	f	0	retinopathy of prematurity	none	40
25	SI	22	m	-	-	-	85
26	SI	22	m	-	-	-	85
27	SI	23	m	-	-	-	95
28	SI	22	m	-	-	-	90
29	SI	34	m	-	-	-	70
30	SI	35	m	-	-	-	85
31	SI	40	m	-	-	-	80
32	SI	26	m	-	-	-	75

33	SI	23	m	-	-	-		70
34	SI	27	m	-	-	-		100
35	SI	20	m	-	-	-		95
36	SI	39	m	-	-	-		85

ABL, acquired blind; CBL, congenitally blind; SI, sighted; y, year; m, male. Superscripts A and NA indicate athlete and nonathlete blind participants, respectively.

Table 2. SD values in no-stimulation condition and stimulation-to-pV1/V2 condition and the increase in SD for each group in Experiment-1

	$SD_{no\text{-}stim}$	$SD_{pV1/V2}$	ΔSD
ABL (n = 11)	**0.076 ± 0.045**	**0.089 ± 0.042**	**0.014 ± 0.018**
ABL^A (n = 5)	0.053 ± 0.010	0.067 ± 0.027	0.015 ± 0.021
ABL^{NA} (n = 6)	0.095 ± 0.055	0.108 ± 0.045	0.013 ± 0.018
CBL (n = 12)	**0.111 ± 0.078**	**0.099 ± 0.050**	**-0.012 ± 0.041**
SI (n = 12)	**0.106 ± 0.048**	**0.104 ± 0.043**	**-0.002 ± 0.019**

ABL, acquired blind; CBL, congenitally blind; SI, sighted. Superscripts A and NA indicate athlete and nonathlete blind participants, respectively. $\Delta SD = SD_{pV1/V2} - SD_{no\text{-}stim}$. All data are reported as mean ± standard deviation across participants. The units are seconds.

Table 3. Characteristics of the participants in Experiment-2

Participant #	Group	Age (y)	Sex	Onset of blindness (y)	Cause of blindness	Light sensitivity at present	TMS intensity for brain (%)	TMS intensity for neck (%)
2	ABL^A	33	m	14	retinal choroidal degeneration	yes	90	70
5	ABL^A	46	m	25	retinitis pigmentosa	yes	100	100
6	ABL^{NA}	30	m	14	chorioretinal atrophy	yes	80	90
7	ABL^{NA}	30	m	5	glaucoma	none	75	60
11	ABL^{NA}	23	m	15	glaucoma	none	55	55
37	ABL^{NA}	32	m	2	retinoblastoma		100	65
38	ABL^{NA}	27	f	10	detachment of retina + cataract	yes	50	45
39	ABL^{NA}	43	m	34	medical error in cataract surgery	none	75	75

40	ABL[NA]	23	m	2	persistent fetal vasculature	none	50	50
41	ABL[NA]	36	m	11	detachment of retina	none	45	40
42	ABL[NA]	44	f	30	retinoblastoma	none	85	85

771 *ABL, acquired blind; y, year; m, male; f, femal. Superscripts A and NA indicate athlete*

772 *and nonathlete blind participants, respectively.*

775 **Table 4. SD values in no-stimulation condition (no-stim) and stimulation-to-neck**

776 **condition and the increase in SD in Experiment-2**

	$SD_{no\text{-}stim}$	SD_{neck}	ΔSD
ABL (n = 11)	0.088 ± 0.022	0.087 ± 0.021	-0.002 ± 0.009

ABL, acquired blind. $\Delta SD = SD_{neck} - SD_{no\text{-}stim}$. All data are reported as mean ± standard

deviation across participants. The units are seconds.

Figures

Fig. 1. The movement task and stimulation sites: A) The rhythmic foot movement of a
representative participant for a 10-s period is shown by the elevation angle, θ of right foot
(dark gray line) and left foot (light gray line). For each foot in each trial, θ is defined as
the angle relative to the average angle of the trial (θ = 0, dashed line). Each light-green
vertical line indicates the timing of a single transcranial magnetic stimulation (TMS)
pulse delivered at random. The cycle duration was defined as the time between the
adjacent dorsiflexion peaks. The SD of the cycle durations was calculated for each foot
in each trial (see **Methods**). B) Fourteen stimulation sites arranged in a 2 × 7 grid were
identified for each participant using the inion and the median line as reference points (see
Methods). Stimulation site #11 corresponds to putative primary and secondary visual
cortex (pV1/V2).

Fig. 2. A) SD of the cycle duration for acquired blind (blue), congenitally blind (green), and sighted (red) participants in Experiment 1. The SD was compared between the no-stimulation condition (no-stim) and the stimulation-to-pV1/V2 condition (pV1/V2) in which TMS was applied to stimulation site #11. Each line represents data from an individual participant. For the acquired blind participants, dashed blue lines and solid blue lines indicate subgroups of athletes (ABL^A) and nonathletes (ABL^{NA}), respectively. B) Increases in the SD from the no-stim condition to the stimulation-to-pV1/V2 condition. Box plots show the median and interquartile ranges of the increase in the SD for each group. Each dot represents data from an individual participant. For the blind group, open dots and closed dots indicate the subgroups of athletes (ABL^A) and nonathletes (ABL^{NA}), respectively. The asterisk indicates $p < 0.05$ with Bonfferoni correction.

Fig. 3. SD of the cycle duration for acquired blind participants in Experiment 2. The SD
was compared between the no-stimulation condition (no-stim) and the stimulation-to-
neck condition in which TMS was applied over the right spelenius capitis (see **Methods**).
Each line represents data from an individual participant. Dashed lines and solid lines
indicate subgroups of athletes (ABL[A]) and nonathletes (ABL[NA]), respectively. B)
Increases in the SD from the no-stim condition to the stimulation-to- neck condition. Box
plots show the median and interquartile ranges of the increase in the SD for each group.
Each dot represents data from an individual participant. For the blind group, open dots
and closed dots indicate the subgroups of athletes (ABL[A]) and nonathletes (ABL[NA]),
respectively.

 SUPPLEMENTARY INFORMATION

TMS simulation to identify the stimulation site

We conducted a simulation analysis to investigate whether the TMS used in Experiment-

1 could stimulate the early visual cortices (V1 and/or V2). Specifically, we used

anatomical MRI data from each participant to simulate the electrical field induced in the

visual cortex by TMS at stimulation site #11.

Participants

Fifteen (eight acquired blind and seven sighted) out of the 23 individuals from the

acquired and sighted groups who participated in Experiment-1 participated in this

simulation study. The rest of the participants were excluded because of contraindications

to MRI (one sighted and four blind) or video recording problems in the TMS experiment

(four sighted).

Procedures

We acquired T1-weighted anatomical brain images of the participants (MP-RAGE;

repetition time [TR]: 1900 ms; echo time [TE]: 2.48 ms; inversion time [TI]: 900 ms; flip

angle: 9°, field-of-view [FOV]: 256 mm × 256 mm; voxel size: 1.0 mm × 1.0 mm × 1.0

mm) with a 3.0-T SIEMENS scanner (Trio Tim, with a 32-channel head coil). Then, we

performed the following simulation analysis for each participant to identify the electrical

field induced by TMS to stimulation site #11.

Simulation

We conducted the simulation for two trials on stimulation site #11 for each participant,

following the procedure introduced by Saturnino et al. (2019). We first segmented the

MRI volume into the different head tissues, and the tissue surfaces were identified. Then,

we simulated the electrical field induced by stimulation by the TMS coil placed at 2.5 cm

dorsal from the inion (stimulation site #11). From the simulated electrical field, the

intensity of the electrical field (norm of the electrical field vector; V/m) in the cortical

gray matter was mapped on the cortical surface (Fig. S3).

The simulation for each participant requires three parameters: the location and orientation

of the TMS coil and the stimulus intensity. For the location, we first identified the inion

(Iz), the nasion (Nz), and the vertex (Cz) on the scalp surface based on the morphometrical

structure of the head surface. Then, we identified the point 2.5 cm above Iz on the circle

passing through Iz, Nz, and Cz. For the orientation, we identified the TMS coil orientation
in the actual TMS experiment by visual inspection of the video recording of the
experiment by one of the authors (TI). The coil orientation in each of the two trials for
stimulation site #11 was identified in 5º increments. The stimulus intensity used in the
simulation was identical to that used in the TMS experiment. We used the default values
for the other simulation parameters, including the tissues' electrical conductance.
866
We used the headreco pipeline (Nielsen et al., 2018) with Statistical Parametric Mapping
(SPM; https://www.fil.ion.ucl.ac.uk/spm/software/spm12) and Computational Anatomy
Toolbox for SPM (https://neuro-jena.github.io/cat) with default parameters for the
segmentation. We performed the simulation with simNIBS software (simNIBS 3.2;
https://simnibs.github.io/simnibs). The anatomical identification of the cortical area was
based on the human cortical parcellation provided by the Human Connectome Project
(HCP-MMP1.0; Glasser et al., 2016).
874
Results
We performed two analyses with the simulation data. First, we examined whether the
intensity peak of the simulated electrical field was observed in V1/V2 (Table S3). The
peak of the electrical field was observed within V1 or V2 in 14 out of 15 participants
(Table S3) in both trials. An exception was observed in Participant #2, whose peak was
in V3A (7.4 mm apart from the edge of V1).
881
Next, we checked whether the intensity of the electrical field induced in V1/V2 was
sufficient to affect neural processing in these regions. We regarded an electrical field >
10 V/m as sufficient to affect neural processing. The threshold was set according to
previous studies showing that neurons fire in response to an artificially induced electrical
field with ~ 10 V/m intensity (Park et al., 2013, Bonmassar et al., 2012, Chan and
Nicholson 1986). The proportion of the V1/V2 surface area where the induced electrical
field was above the threshold (stimulated region) is 81% ± 10% (mean ± standard
deviation; see Table S3).
890
891 These two results suggest that the TMS procedure used in Experiment-1 substantially
892 stimulated V1/V2, with sufficient stimulus intensity.
893

References

Saturnino, G. B., Puonti, O., Nielsen, J. D., Antonenko, D., Madsen, K. H., & Thielscher,
A. (2019). SimNIBS 2.1: a comprehensive pipeline for individualized electric field
modelling for transcranial brain stimulation. Brain and human body modeling, Springer,
3-25.

Nielsen, J. D., Madsen, K. H., Puonti, O., Siebner, H. R., Bauer, C., Madsen, C. G.,
Saturnino, G. B., & Thielscher, A. (2018). Automatic skull segmentation from MR
images for realistic volume conductor models of the head: Assessment of the state-of-the-
art. NeuroImage, 174, 587–598.

Park, H. J., Bonmassar, G., Kaltenbach, J. A., Machado, A. G., Manzoor, N. F., & Gale,
J. T. (2013). Activation of the central nervous system induced by micro-magnetic
stimulation. Nature communications, 4(1), 1-9.

Bonmassar, G., Lee, S. W., Freeman, D. K., Polasek, M., Fried, S. I., & Gale, J. T. (2012).
Microscopic magnetic stimulation of neural tissue. Nature communications, 3(1), 1-10.

Chan, C. Y., & Nicholson, C. (1986). Modulation by applied electric fields of Purkinje
and stellate cell activity in the isolated turtle cerebellum. The Journal of physiology,
371(1), 89-114.

Supplementary figures

Fig. S1. SD of the cycle duration for acquired blind (blue), congenitally blind (green),
and sighted (red) groups in Experiment-1. The average SD over all stimulation sites (all
sites) was compared with the SD in the no-stimulation condition (no-stim). Each line
represents data from an individual participant. For the blind group, dashed blue lines and
solid blue lines indicate the subgroups of athletes (ABL[A]) and nonathletes (ABL[NA]),
respectively. B) Increases in the SD from the no-stim condition to the stimulation-to-
pV1/V2 condition. Box plots show the median and interquartile ranges of the increase in
the SD for each group. Each dot represents data from an individual participant. For the
blind group, open dots and closed dots indicate the subgroups of athletes (ABL[A]) and

nonathletes (ABL^{NA}), respectively. The asterisk indicates $p < 0.05$ with Bonfferoni

correction.

935
936
Fig. S2. SD of the cycle duration for the blind participants in the no-stimulation condition
(x-axis) and the stimulation-to-pV1/V2 condition (y-axis) of Experiment 1. A) For each
group of acquired (blue) and congenitally (green) blind participants, two subgroups with
and without light sensitivity (filled and open circles, respectivly) were compared. The
dashed line indicates y = x. In the acquired blind participants, TMS to pV1/V2 increased
the SD in the cycle duration relative to the no-stimulation condition in all participants
exept for one (#8, see Table 1). The increase in SD did not differ between the two
subgroups. In contrast, no clear trend was not observed among the congenitally blind
participants nor between the twosubgroups. B) Early-, intermediate-, late-acquired blind
participants (light, intermediate, and dark blue circles, respectively), and congenitally
blind participants (green circle) are compared. The dashed line indicates y = x. No clear
difference was observed among the three subgroups of the acquired blind participants.
949

Norm of the electrical field vector(V/m)

0   53

950
951 Fig. S3. Distribution of the electrical field intensity (norm of the electrical field vector; V/m) in
952 the cortical gray matter in a typical participant (participant #6). The intensity in the cortical gray
953 matter was mapped onto the cortical surface. The figure shows a posterior view of the cortex.
954
955

Table S1. SD of the cycle duration in all stimulation sites (#1-14) and the no-stimulation condition (no-stim) for the participants of Experiment-1 and the congenitally blind participants.

Gr SD #	SI (n = 12)	ABL (n = 11)	ABL^A (n = 5)	ABL^{NA} (n = 6)	CBL (n = 12)
no-stim	0.106±0.048	0.076±0.045	0.053±0.010	0.095±0.055	0.111±0.078
1	0.104±0.033	0.086±0.039	0.072±0.031	0.098±0.043	0.092±0.036
2	0.101±0.042	0.097±0.057	0.087±0.043	0.105±0.069	0.099±0.043
3	0.108±0.046	0.081±0.039	0.064±0.019	0.095±0.048	0.104±0.065
4	0.108±0.040	0.090±0.042	0.069±0.018	0.108±0.049	0.095±0.043
5	0.104±0.033	0.085±0.034	0.071±0.028	0.097±0.036	0.095±0.041
6	0.110±0.050	0.085±0.036	0.069±0.027	0.099±0.040	0.106±0.045
7	0.111±0.045	0.088±0.039	0.069±0.027	0.105±0.042	0.105±0.049
8	0.105±0.052	0.082±0.037	0.064±0.024	0.097±0.041	0.097±0.065
9	0.107±0.047	0.088±0.048	0.065±0.023	0.107±0.057	0.097±0.054
10	0.117±0.053	0.086±0.052	0.067±0.028	0.102±0.063	0.101±0.049
11 (pV1/V2)	0.104±0.043	0.089±0.042	0.067±0.026	0.108±0.045	0.099±0.050
12	0.107±0.043	0.088±0.043	0.069±0.019	0.104±0.052	0.095±0.042
13	0.105±0.042	0.083±0.032	0.064±0.016	0.099±0.034	0.103±0.055
14	0.098±0.037	0.086±0.028	0.077±0.026	0.093±0.030	0.113±0.064

Gr, group; SI, sighted; ABL, acquired blind; CBL, congenitally blind. Superscripts A and NA indicate athlete and nonathlete blind participants, respectively. All data are reported as mean ± standard deviation across participants. The units are seconds.

Table S2. Simulation results

Participant #	Group	Peak location	proportion of stimulated region in V1/V2 (%)
1	ABL^A	Left V1/Right V1*	88.5
2	ABL^A	Left V3A	54.9
4	ABL^A	Left V1	81.3
5	ABL^A	Left V1	80.7
6	ABL^{NA}	Left V1	76.5
7	ABL^{NA}	Left V1	83.5
9	ABL^{NA}	Right V1	91.4

10	ABL[NA]	Left V2	88.7
28	SI	Left V2/Right V1*	86.6
30	SI	Right V1	86.8
31	SI	Left V2	77.7
33	SI	Left V1	81.0
34	SI	Left V1	87.8
35	SI	Left V1/Left V2*	90.3
36	SI	Left V1	80.8

967   * Asterisks indicate that the peaks for the first and second trials were located in different
968   anatomical areas, where results were indicated as follows: first trial peak location / second
969   trial peak location.
970