Short title: Rudimentary basal spikelets in wheat

High expression of *VRT2* increases the number of rudimentary basal

spikelets in wheat

Anna E. Backhaus[1,2], Ashleigh Lister[2], Melissa Tomkins[1], Nikolai M. Adamski[1], James

Simmonds[1], Iain Macaulay[2], Richard J. Morris[1], Wilfried Haerty[2], Cristobal Uauy[1]

[1] John Innes Centre, Norwich Research Park, Norwich NR4 7UH, United Kingdom

[2] Earlham Institute, Norwich Research Park, Norwich NR4 7UZ, United Kingdom

One sentence summary: Large transcriptional gradients exist *within* a wheat spike and are

associated with rudimentary basal spikelet development, resulting in the characteristic

lanceolate shape of wheat spikes.

Author Contribution: (Contributions are based on the CRediT Contributor Roles Taxonomy
(https://casrai.org/credit/)
AEB: Conceptualization, Methodology, Data curation, Formal analysis, Investigation,
Visualization, Writing – original draft
AL: Investigation, Methodology
MT: Software, Formal analysis, Writing – review & editing
NMA: Resources, Writing – review & editing
JS: Data curation, Formal analysis, Resources, Writing – review & editing
IM: Methodology, Supervision
RJM: Conceptualization, Methodology, Supervision, Writing – review & editing
WH: Conceptualization, Funding acquisition, Supervision, Writing – review & editing
CU: Conceptualization, Funding acquisition, Methodology, Project administration,
Supervision, Writing – review & editing
26

Funding: This work was supported by the UK Biotechnology and Biological Sciences

Research Council (BBSRC) through the grant BB/S016945/1, the Designing Future Wheat

(BB/P016855/1) and Genes in the Environment (BB/P013511/1) Institute Strategic

Programmes and by the John Innes Foundation. Additional funding was provided by the

European Research Council (866328).

Author for contact: Cristobal Uauy (cristobal.uauy@jic.ac.uk)

Abstract

Spikelets are the fundamental building blocks of *Poaceae* inflorescences and their development and branching patterns determine the various inflorescence architectures and grain yield of grasses. In wheat, the central spikelets produce the most and largest grains, while spikelet size gradually decreases acro- and basipetally, giving rise to the characteristic lanceolate shape of wheat spikes. The acropetal gradient correlates with the developmental age of spikelets, however the basal spikelets are developed first and the cause of their small size and rudimentary development is unclear. Here, we adapted G&T-seq, a low-input transcriptomics approach, to characterise gene expression profiles within spatial sections of individual spikes before and after the establishment of the lanceolate shape. We observed larger differences in gene expression profiles between the apical, central and basal sections of a single spike than between any section belonging to consecutive developmental timepoints. We found that *SVP* MADS-box transcription factors, including *VRT-A2*, are expressed highest in the basal section of the wheat spike and display the opposite expression gradient to flowering E-class *SEP1* genes. Based on multi-year field trials and transgenic lines, we show that higher expression of *VRT-A2* in the basal sections of the spike is associated with increased numbers of rudimentary basal spikelets. Our results, supported by computational modelling, suggest that the delayed transition of basal spikelets from vegetative to floral developmental programmes results in the lanceolate shape of wheat spikes. This study highlights the value of spatially resolved transcriptomics to gain new insights into developmental genetics pathways of grass inflorescences.

Introduction

The arrangement of flowers in individual plants of the same species is highly conserved and follows a systematic and rhythmic pattern. This systematic appearance of flowers is not surprising, as floral architectures are determined by the regular initiation of flower primordia on the flanks of the apical meristem and their rate of initiation and developmental fate are under strong genetic control (Prusinkiewicz et al., 2007). The unifying feature of floral architecture in grasses (*Poaceae*) is the formation of all flowers (termed florets) within spikelets (Kellogg et al., 2013). Spikelets are the fundamental building blocks of grass inflorescences and their development and branching patterns determine the various inflorescence architectures of grasses (e.g., spikes, panicles). Wheat (*Triticum aestivum*) forms a spike shaped inflorescence, in which sessile spikelets are directly attached to the inflorescence axis (or rachis) in a distichous phyllotaxis (Koppolu and Schnurbusch, 2019). Upon floral transition, the vegetative meristem ceases to initiate leaf primordia and transitions into the inflorescence meristem (IM). During the Double Ridge stage (DR) of wheat spike development, the IM initiates a lower leaf ridge and an upper spikelet ridge (or primordia) during each iteration. Within the inflorescence the upper ridges differentiate into spikelet meristems, while the lower ridges are suppressed upon flowering (Bommert and Whipple, 2018). DR initiation will continue at the IM until the terminal spikelet stage, when IM forms a final spikelet (Koppolu and Schnurbusch, 2019). Spikelet initiation and development has been extensively studied in wheat and other monocot crops, such as rice (*Oryza sativa*), maize (*Zea mays*), and barley (*Hordeum vulgare*), as the number of spikelets per spike is a major determining factor for grain number and thus yield per spike.

Not all spikelets across the wheat spike, however, produce the same amount of grain. The central spikelets produce the most and largest grains, while spikelet size gradually decreases acro- and basipetally. Within a single spike, the most apical and basal spikelets might produce no or only one grain while the central spikelets of the same spike set 3-5 grains. Bonnett (1966) documented that this distinct lanceolate shape of the wheat spike is first established during the Glume Primordia (GP) stage (just after the DR stage). This asynchronous development among the spikelets is maintained throughout the development of the spike. The gradual decrease in spikelet size from the central to apical section of the spike can be explained by the continuous development of new spikelet ridges from the apical inflorescence meristem: the most apical spikelets are the youngest and had the least time to develop. However, basal spikelets are initiated first and it is unclear why they remain smaller than their central counterparts. In the

mature spike the most basal one or two spikelets are often only formed in a rudimentary
manner, with small glumes present but all floral structures remaining immature.

Efforts to understand the genetics of wheat spikelet initiation and development have focused
on members of the MADS-box transcription factor (TF) family, which play central roles in the
flowering gene models (Zhao et al., 2006). Li et al. (2019) showed that MADS-box genes of
the   *SQUAMOSA*-clade,   *VERNALISATION1*   *(VRN1)*, *FRUITFULL2* (*FUL2*)
and *FRUITFULL3* (*FUL3*), have overlapping functions in controlling the timing of the
transitions from the vegetative to IM as well as the formation of the terminal spikelet.
In *vrn1ful2*-null mutants, the IM remained indeterminate causing the mutants to form more
spikelets per spike. However, all lateral spikelets were replaced by leafy shoots in the *vrn1ful2*
double and *vrn1ful2ful3* triple mutants (Li et al., 2019). These mutants had increased
expression of genes belonging to the *SHORT VEGETATIVE PHASE* (*SVP*) family of MADS-
box genes, including *VEGETATIVE TO REPRODUCTIVE TRANSITION 2* (*VRT2*).
Subsequent studies determined that overexpression of *VRT2* led to reversion of basal spikelets
to spikes and the downregulation of other MADS-box genes required for floral development,
including members of the *SEPALLATA1* (*SEP1*) clade (Li et al., 2021). Together, these studies
exemplify the importance of the temporal sequence of flowering gene expression for the correct
development of the wheat spike.

Attempts to unravel the genetic network controlling wheat spike development have focused on
these temporal changes in expression patterns across consecutive developmental stages. For
example, Li et al. (2018) and Feng et al. (2017) performed transcriptome profiling using pooled
samples of multiple complete spikes from six (vegetative to floret differentiation) and four
(double ridge to young floret) developmental stages, respectively. In a few cases, studies have
examined the expression patterns of individual genes (via quantitative reverse transcription
(qRT)-PCR) and found gene expression gradients along the spike. For example, Debernardi et
al. (2017) demonstrated that *APETALA2* (*AP2*) is expressed higher in the apical section of
wheat spikes than in central or basal sections. This *AP2* expression gradient was associated
with morphological changes along the same spike. This study alongside work in barley
(Youssef et al., 2017), suggests that gene expression gradients *within* individual developmental
stages could be important to further unravel the genetic control of spike development.
However, despite its potential biological significance, spatial transcriptome profiles along the
spike have yet to be investigated in wheat.

In this study, we aimed to characterise gene expression profiles along the spike during the establishment of the lanceolate shape of the wheat spike from DR to GP. We adapted G&T-seq (Genome and Transcriptome sequencing), a low-input sequencing approach to sequence the transcriptome of the sections. Recently, Giolai et al. (2019) adapted the protocol to identify expression differences across single leaves of *Arabidopsis* (GaST -seq), demonstrating that the G&T-seq method can be readily used for sequencing of hand harvested, small input plant material without the need of previous tissue dissociation or treatment. G&T-seq is thus comparable to methods using laser-micro dissection followed by sequencing to achieve spatially resolved transcriptome wide sequencing data. In comparison, the available transcriptome sequencing methods at higher resolutions (such as single cell RNA-seq or fluorescence-activated cell sorting (FACS)) are not spatially resolved as the complete tissue is dissolved into single cells for barcoding or selection prior to sequencing (Rich-Griffin et al., 2020).

We sequenced the apical, central, and basal sections of individual spikes before (DR) and after (GP) the establishment of the lanceolate shape. Gene expression profiles differed most strongly between spatial sections of the same spike, as opposed to temporal sections (any two sections from different timepoints). Members of the *SVP* gene family were expressed most highly in the basal sections with expression decreasing upwards from the base (acropetally), while members of the *SEP1* gene family showed the opposite expression pattern, i.e. most highly expressed in apical sections, with expression decreasing towards the base (basipetally). The increased number of rudimentary basal spikelets due to *VRT-A2* misexpression supports the hypothesis that high expression levels of *SVPs* in the basal section delays spikelet establishment, leading to their rudimentary shape in the mature spike. This study highlights that spikelets within the same spike experience significantly different flowering signals due to their consecutive development and spatial position within the spike. Acknowledging these differences can help us gain a better understanding of the genetic flowering pathway of grass inflorescences.

Results

Low-input sequencing enables spatial analysis of the wheat spike transcriptome

To investigate transcriptional differences between the apical, central, and basal section of developing wheat spikes, we adapted the low-input G&T sequencing (G&T-seq) method for RNA-seq of small plant tissue sections. G&T has been developed for single-cell RNA and DNA sequencing of mammalian systems (Macaulay et al., 2015) and was previously adapted for *Arabidopsis thaliana* (GaST-seq; Giolai et al., 2019). We collected four individual developing wheat (cv Paragon) spikes at both the double ridge (DR) and glume primordia (GP) stage and hand-dissected them into apical, central, and basal sections (Figure 1A).

Figure 1: (A) Summary diagram of low-input G&T tissue collection and sequencing approach. Grey circles indicate the 24 individual libraries prepared for sequencing from each individual tissue section dissected from individual spikes. (B) Reads per library after trimming and quality controls (see Methods). Stacked bars indicate the number of reads aligned (blue) and not aligned (red) by HISAT to the RefSeqv1.0 genome. (C) Number of expressed genes (>10 read counts) per library based on tissue section and Waddington developmental stage (DR: Double Ridge; GP: Glume Primordia). In (B and C), the X-axis indicates the ID of each sample which is composed of the tissue section (A: apical, C: central, B: basal) and plant number (1-8) as indicated in (A). Detailed quality control data for each library is provided in Supplemental Table S1.

On average, samples had 28,799,626 reads (coefficient of variation (CV) 43%), of which 90%

(CV 8.5%) aligned to the genome post adaptor trimming (Figure 1B, Table 1). Furthermore,

the number of aligned reads and the number of expressed genes per library was largely

homogenous among the spatial sections and Waddington stages (Table 1). On average, 47,313

genes per library were expressed (>10 read counts) and we found no difference ($P > 0.56$) in

the number of expressed genes across spatial (apical, central, basal) or between temporal (DR,

GP) conditions (Figure 1C). We excluded three libraries with low average number of expressed

genes (difference greater than five times the standard deviation; Figure 1C, Supplemental Table

S1) and two libraries because they were strong outliers in the principal component analysis

(PCA; Supplemental Figure S1A). In total, 19 RNA-seq libraries (DR: 3 apical, 4 central, 3

basal; GP: 2 apical, 3 central, 4 basal) passed our selection criteria and were used in the

subsequent analyses. We identified 91,646 genes being expressed across these 19 libraries.

Table 1: Average number of reads aligned to the RefSeqv1.0 genome and expressed genes
(>10 read counts) in the three tissue sections and two Waddington developmental stages (DR:
Double Ridge; GP: Glume Primordia) (n = 4 biological replicates per tissue section *
developmental stage).

	Reads aligned		Genes expressed	
	DR	**GP**	**DR**	**GP**
Apical	26,342,934	23,913,657	44,488	41,152
Central	29,823,023	25,854,540	45,913	50,074
Basal	25,174,482	23,627,494	49,513	52,740

Transcriptome-wide differences are largest between the apical and basal sections of the spike

To investigate global differences among the 19 RNA-seq libraries, we performed a principal

component analysis (PCA; Figure 2A). The first two PCs explained 19% and 16% of the overall

variance present in the libraries. We observed that the two PCs separated libraries by the spatial

position (apical, central, basal) rather than developmental stage (DR, GP). There was a clear

separation between libraries originating from apical and basal spike sections, while libraries

from central sections were dispersed between these two clusters (Figure 2A). We investigated

PC1 to PC6 and found that none of these combinations clustered libraries by developmental
stage (Supplemental Figure S1B). Given that we sequenced developing spike sections of single
plants, as opposed to the more commonly employed pooling of multiple biological samples,
we found as expected some degree of heterogeneity between samples from the same location
and stage (Figure 2A).

Figure 2: (A) Principal component analysis (PCA) on the 19 transcriptome libraries from
apical (blue), central (yellow) and basal (green) sections of Double Ridge (circles) and Glume
Primordia (triangles) spikes. Black bordered triangle is plant 8 (GP) in which the basal section
clustered closer with central-GP sections than the other basal-GP sections. (B) UpSet plot
showing the number of differentially expressed genes (DEGs) between spatial sections and
Waddington stages. Black border indicates Plant 8, which is an outlier of basal GP replicates.

To investigate this variation further, we quantified changes in gene expression across biological
replicates by calculating CVs for each gene (see Methods). The median CV for a gene across

the biological replicates was 39% (Supplemental Figure S2) with a Q1-Q3 interquartile range

between 24% and 62%. We also calculated the CV per gene for published datasets from Li et

al (2018) and Feng et al (2017). Both studies sequenced developing wheat spikes at similar

developmental stages, pooling many spikes per sample. Li et al. (2018) pooled between 100 to

200 spikes of winter wheat (KN9204) per sample, while Feng et al. (2017) reported pooling of

10 to 50 spikes (cv. Chinese Spring) per sample. In both studies the median CV of a gene was

lower (14% and 21%, respectively) than in our study. The larger CVs in our data could be

explained by the biological variation that exists between individual plants, which may have

been reduced by the pooling of many spikes in both Li et al. (2018) and Feng et al. (2017).

We first analysed differentially expressed genes (DEGs) between the DR and GP stage and

between apical, central and basal sections across the two Waddington developmental stages.

The number of DEGs between DR and GP (215 genes) was smaller than the number of DEGs

identified between the spatial positions, which ranged from 280 DEGs between central and

apical sections to 914 DEGs between the apical and basal sections (Figure 2B). Next, we

compared the apical, basal and central sections within each Waddington developmental stage.

We identified more DEGs by comparing the spatial sections within either Waddington stage

individually than in the combined analysis. The number of DEGs between apical and basal

sections at each stage (DR: 2,438; GP: 1,315) were similar to the number of DEGs between

central and basal sections (DR: 2,022; GP: 1,186). The number of DEGs between these sections

at DR, however, was nearly double the number of DEGs at GP. In contrast, the number of

DEGs between apical and central sections was similar at both stages (DR: 1,296; GP: 1,118),

suggesting that the basal section of the spike is most different in the earlier developmental

stage. Only 11% of the DEGs were shared between DR and GP in the apical to basal

comparison, 7% between the central to basal DEGs, and 5% between apical to central DEGs.

In total, we identified 5,353 unique genes as differentially expressed between any of the three

sections at either Waddington stage (Supplemental Table S2). Overall, the number of DEGs

was largest between the apical and basal sections, reflecting the strong spatial clustering

observed in the PCA graph, but most genes that were differentially expressed across the spike

did not maintain this gradient over the two developmental stages. In summary, despite the high

biological variation in gene expression in our data compared to previous pooled whole-spike

studies, we could detect transcriptome wide differences between the spatial sections of

developing wheat spikes.

The *SVP* MADS-box transcription factors have opposing expression profiles to flowering E-class genes

To further investigate the differences in expression across the spike and to identify genes with similar expression patterns, we performed hierarchical and k-means clustering. We restricted the clustering to the 5,353 genes identified as differentially expressed across the spike at either one or both Waddington stages (Figure 3A). We identified seven non-redundant clusters, each containing between 8% to 21% of the 5,353 DEGs (Figure 3A, Supplemental Figure 3A). Both hierarchical and k-means clustering produced highly similar results (Supplemental Figure S3B). We identified 1,894 genes (35% of DEGs) to be more highly expressed in the apical section, either across both timepoints (503 genes, cluster 1), or only at DR (751 genes, cluster 7) or GP (640 genes, cluster 3). In the central section, 1,362 genes (25%) had higher relative expression at either DR (917 DEGs, Cluster 5) or GP (445 DEGs, Cluster 6). In the basal section, we observed 2,097 genes (39%) being more highly expressed. Cluster 4 contained the most DEGs (1,170) and was characterized by an upregulation of expression in the basal section at both Waddington stages, although this upregulation was higher at the DR stage. Another 927 genes were upregulated in the basal section, but only at the GP stage (cluster 2).

Figure 3: (A) Normalized expression matrix and K-means clustering of the 5,353 genes differentially expressed across the spike at either one or both Waddington stages. Colours (blue to red) show relative log2 expression of genes after normalisation. (B) Expression pattern of the 1,170 genes allocated to Cluster 4 (left), and of MADS-box transcription factors in the same cluster (right). Colours indicate how well the gene expression pattern fits the average expression pattern (black line). Red = best fit, Blue = least good fit. (C) Expression pattern of the 640 genes and MADS-box transcription factors of Cluster 3 as arranged in B. Norm = normalized and scaled gene expression. RefSeq1.1 gene IDs and raw expression values of genes shown in the right-hand panels are presented in Supplemental Table S2.

To further characterize the clusters, we independently tested for enrichment of TF families and
gene ontology (GO)-terms (all GO-terms and TF families in Supplemental Table S3 and S4,
respectively). Genes that were more highly expressed in the apical section were enriched for
the GO terms "reproductive structure development" (GO:0048608) and "floral organ
development" (GO:0048437; cluster 1) as well as for HD-Zip_IV and SRS TF families ($P <$
0.0001; cluster 1). In cluster 3 (highly expressed in the apical section at GP; Figure 3B) we
found no significant enrichment of GO-terms ($P < 0.03$), but a significant enrichment of
MADS_II TFs ($P = 0.013$). For clusters defined by an increased expression in the central
sections (clusters 5 and 6) we detected an enrichment for GO-terms related to polyphosphate
processes (GO:0006797/0006779), but a significant enrichment of the C2C2_CO-like TFs at
DR ($P = 0.036$; cluster 5). Genes with higher expression in the basal section of the spike (cluster
4; Figure 3C) were enriched for a number of GO terms related to photosynthesis, (e.g.
GO:0015979) and "negative regulation of flower development" (GO:0009910), as well as for
MADS_II TFs ($P = 0.084$). Cluster 2, which was characterized by higher expression in the
basal section only at GP, was enriched for "Jasmonic acid response" (GO:0009753) and Tify
and C2C2_CO-like TFs ($P < 0.04$).

We were interested in further characterizing the expression patterns of the MADS-box TFs as
they were significantly enriched in two of the seven clusters and are important in floral
transition and development (Becker and Theißen, 2003; Feng et al., 2017). We detected 14
differentially expressed MADS-box TFs in our study, five of which were more highly
expressed in the apical section, four of these only at GP stage (cluster 3) and one being
consistently expressed across both Waddington stages (cluster 1). In contrast, five MADS-box
TFs were more highly expressed in the basal section at both Waddington stages (cluster 4) and
another two were more highly expressed in the basal section only at GP (cluster 2). An
additional two MADS-box genes were part of the remaining clusters (Supplemental Table S4).

In the apical/GP cluster 3 we noticed that all MADS-box genes belonged to the *Triticum*
aestivum SEPALLATA1 (*TaSEP1*) group (Figure 3C). All three homoeologs of *TaSEP1-4*
(*TraesCS7A02G122000, TraesCS7B02G020800, TraesCS7D02G120500)* and the D-genome
copy of *TaSEP1-5* (*TraesCS7D02G120600*) were part of this cluster. The *SEP* genes were
expressed at relatively low levels at DR (Supplemental Table S2), but were significantly
upregulated at GP, with their transcript levels being highest in the apical section. The increased
expression of *TaSEP1-4* at GP was in agreement with their previously reported expression
patterns in tetraploid wheat by Li et al. (2021).

In the contrasting cluster 4 (upregulation in basal sections), we noticed the presence of multiple

MADS-box genes belonging to the *SVP* family (Figure 3B, right-hand panel), which consists

of three genes in wheat (*SVP1*, *VRT2* and *SVP3*). Members of this family are important for the

transition from vegetative to floral meristem identity in cereals (Trevaskis et al., 2007). All

three homoeologs of *VRT2* (*TraesCS7A02G175200, TraesCS7B02G080300,*

TraesCS7D02G176700) and the B-genome copy of *SVP1* (*TraesCS6B02G343900*) were

present in cluster 4. The cluster also contained *TaFLC-A1 (TraesCS7A02G260900),* although

it was expressed higher in DR-central sections compared to the *SVPs* and had a linear

expression gradient at GP. All *SVPs* had very similar expression patterns, being strongly

expressed in basal sections only. Expression of *SVPs* was higher in all DR sections compared

to the equivalent section in GP. Constitutive over-expression of *SVP*-family members in wheat

and barley has been shown to delay or even reverse floral development (Trevaskis et al., 2007;

Li et al., 2021). This led to hypothesis that the rudimentary development of basal spikelets was

associated with an increase in *VRT2* expression levels.

SVP expression is higher in basal and peduncle sections and increased across all sections in *T. polonicum VRT-A2b isogenic lines*

316
317

To validate the expression pattern of *VRT2* in the individual spike analysis, we performed qRT-

PCR on independently collected, pooled spike sections from cv Paragon, carrying the wildtype

VRT-A2a allele (Figure 4A, blue curves). We included a later timepoint, Terminal Spikelet

(TS), which is about 10 days after GP to study how *VRT2* expression changes in later stages.

At TS stage, the central spikelets have developed multiple florets primordia. We also included

a small part of the peduncle (stem) section just below the spike as an additional spatial section.

We focused the expression analysis on the A-genome homoeolog, *VRT-A2*, as its role in spike,

glume and grain development of wheat was recently characterised (Adamski et al., 2021; Liu

et al., 2021).

Figure 4: Relative expression (2^{ddCT}) of *VRT-A2* (A) and *SEP1-4* (B) in the different sections of the spike across three timepoints in near isogenic lines (NILs) carrying either the wildtype *VRT-A2a* (blue) or the *VRT-A2b* allele from *Triticum turgidum* ssp. *polonicum* (orange). The data are shown as mean ± SE of gene expression compared with control gene Actin. N = 3 biological replicates (See Supplemental Table S5 for expression data and Supplemental Table S6 for statistical analysis of gene expression differences).

We identified a significant interaction effect between Waddington stage and spatial section; we thus analysed the three Waddington stages separately (Figure 4). At DR, we were limited to dissecting the spike into apical, basal and peduncle sections, as the small size of the spike meristem did not allow precise dissection of the central section when using multiple (pooled) spikes. At DR, we found *VRT-A2* marginally expressed, with significantly lower expression levels in the apical section compared to the basal ($P = 0.003$) and peduncle sections ($P = 0.001$). Although expression in the peduncle was higher than in the basal section at DR, this was not significant ($P = 0.116$). At GP, *VRT-A2* expression was borderline detectable in the apical, central and basal sections, but expression was significantly higher in the peduncle with respect

to the three spike tissues ($P < 0.001$ for all three comparisons). Lastly, *VRT-A2* expression at

TS stage was just detectable and significantly different between all sections ($P = 6.6E\text{-}06$).

Overall, expression decreased significantly from DR to GP/TS Waddington stages in the apical

($P = 0.00015$), basal ($P = 0.0074$), and peduncle ($P = 0.012$) sections consistent with the

previously reported strong downregulation of *VRT-A2* in the early wheat spike development

(Li et al., 2021; Adamski et al., 2021; Liu et al., 2021). This is also consistent with the observed

downregulation of *VRT-A2* orthologs upon floral transition in barley (Trevaskis et al., 2007)

and rice (Harrop et al., 2018). As observed in the low-input RNA-seq data, the qRT-PCR data

confirmed the strong basipetal gradient in *VRT-A2* expression across the spike at DR and

revealed that its expression was even higher within the peduncle.

We hypothesised that the higher expression in the basal section of the wheat spike compared

to the central and apical sections is associated with the rudimentary development of the basal

spikelets. To test the effect of higher *VRT-A2* expression on basal spikelet development, we

analysed the effect of the *Triticum turgidum* ssp. *polonicum VRT-A2b* allele on the expression

gradient of *VRT-A2* and spike morphology. Adamski et al. (2021) showed that *VRT-A2* in *T.*

polonicum, a tetraploid subspecies of wheat, carries a sequence re-arrangement in its first

intron. This results in the higher expression of the *T. polonicum VRT-A2b* allele, with respect

to the wildtype *VRT-A2a* allele, during early spike development. We performed qRT-PCR on

a cv Paragon NIL carrying the *VRT-A2b* allele and compared *VRT-A2* expression against the

Paragon wildtype NIL described above (Figure 4). Consistent with the results of Adamski et

al. (2021), we detected significantly higher expression of *VRT-A2b* compared to the wildtype

allele across most of the tissue sections (see Supplemental Table S6 for individual

comparisons), and a progressive decrease in *VRT-A2b* expression over time ($P = 0.031$). In

contrast to the wildtype NILs, ANOVA did not identify a significant interaction effect between

spatial section and Waddington stage in *VRT-A2b* NILs ($P = 0.18$). We thus examined the

overall expression patterns and found that across all three developmental stages *VRT-A2b*

expression differences were significant ($P < 0.0001$). These results suggest that the basipetal

expression gradient in the spike is maintained in the NILs with the *T. polonicum VRT-A2b*

allele.

We also tested the effect of *VRT-A2* expression levels on *SEP1* expression in the tissue sections

of *VRT-A2* NILs. We confirmed that *SEP1-4* expression is only marginally detectable at DR

and differences in expression between the apical, basal, and peduncle section are hardly

detectable with only marginally higher expression in the apical sections ($P = 0.015$) at this

stage. At GP, *SEP1-4* expression is significantly higher towards the tip of the spike ($P < 0.0001$)
consistent with the low-input RNA-seq data. Furthermore, *SEP1-4* expression was significantly
lower in *VRT-A2b* NILs compared to the wildtype allele across all spike sections ($P = 0.008$),
confirming that higher *VRT-A2* expression can negatively affect *SEP1-4* expression. Similar
trends were observed at TS, where expression was significantly lower in basal sections ($P < 0.0001$) and the gradient across the spike was maintained in *VRT-A2b* NILs, but expression
was overall lower ($P < 0.003$). *SEP1-4* is not expected to be expressed in vegetative tissue such
as the peduncle, therefore the lack of expression in this tissue across the three stages indicates
that no floral tissue was accidently sampled as peduncle (Fig. 4B).

Misexpression of *VRT-A2b* in *T. polonicum* increases rudimentary basal spikelet numbers
To evaluate if the higher expression of *VRT-A2* in basal spikelets affects their development, we
examined the *VRT-A2* NILs (BC_4 and BC_6) sown as winter crops in four environments. In each
field trial, we evaluated the number of rudimentary basal spikelets (RBS), that is spikelets
which are reduced in size and do not contain mature grains (Figure 5A). The number of RBS
was significantly increased in NILs carrying the *VRT-A2b* allele in all four environments ($P < 0.0001$, except Morley 2017 $P < 0.01$; Figure 5B; Supplemental Table S7). The *VRT-A2a* NILs
and the recurrent parent Paragon had on average 1.85 RBS, whereas *VRT-A2b* NILs produced
on average 2.91 RBS. A similar difference in RBS between the NILs was observed in
glasshouse conditions (*VRT-A2b* effect of +1.6 RBS; Supplemental Table S8; Figure 5C).

Figure 5: Phenotypic difference between *VRT-A2a* (blue) and *VRT-A2b* (orange) on
rudimentary basal spikelet numbers (RBS). (A) Mature spikes from the field (left) and
dissected basal spikelets at anthesis (right) from the glasshouse. Numbers indicate position
along the spike starting at the base. Scale bar = 1 cm. (B) Number of RBS per spike from 10-
ear samples collected in the field at maturity at Morley (2017, 2019 and 2020) and Church
Farm (CF, 2020). (C) Number of RBS recorded in the glasshouse for the NILs (left panel) and
for seven critical recombinant lines (R1-R4, R6-R8; see Supplemental Table S8 for graphical
genotype of these lines from Adamski et al. (2021). (D) RBS per spike recorded for the
transgenic lines carrying zero, low (1-5) or high (9-35) copy-number insertions of *VRT-A2b* in
cv. Fielder. In B-D, the box represents the middle 50% of data with the borders of the box
representing the 25th and 75th percentile. The horizontal line in the middle of the box
represents the median. Whiskers represent the minimum and maximum values, unless a point
exceeds 1.5 times the interquartile range in which case the whisker represents this value and
values beyond this are plotted as single points (outliers). *P* values: o ≤0.1; ** ≤ 0.01; *** ≤
0.001.

Furthermore, we also recorded the number of RBS in seven homozygous BC_6 recombinant
lines used to fine-map *VRT-A2b* by Adamski et al. (2021). The RBS phenotype was mapped in
complete linkage with the 50.3 kbp interval containing *VRT-A2* (Figure 5C; Supplemental
Table S8). This genetic and phenotypic data suggests that the increase in RBS is a pleiotropic
effect of the *T. polonicum VRT-A2b* allele and supports the hypothesis that misexpression of
VRT-A2 negatively affects spikelet development in the base of the spike. In Paragon, the first
(sometimes second) rudimentary basal spikelet fully develops the floral organs of florets one
and two (e.g. lemma, palea, stamen, and ovary), however these are severely reduced in size and
delayed in development compared to the florets of central spikelets just before flowering
(~Waddington stage 8-10; Supplemental Figure S4). At this stage, the further growth and
development of these basal florets is stopped and in the mature spike only the glumes of RBS

are visible. In NILs carrying the *VRT-A2b* allele the development of the most basal spikelet is
very similar to the wildtype. However, the second, third and sometimes fourth spikelet also
display similar signs of reduced development, leading to the larger number of rudimentary
basal spikelets (Figure 5B, C; Supplemental Figure S4).

To validate the phenotypic effect of *VRT-A2*, we analysed transgenic wheat lines transformed
with the complete genomic *T. polonicum VRT-A2b* sequence (including the native promoter
and the intron 1 re-arrangement). Transgenic T_1 lines were classified based on the transgene
copy number which was previously shown by Adamski et al. (2021) to be highly correlated
with *VRT-A2* expression levels in multiple tissues. We phenotyped lines with zero (n = 2
independent events; 5 plants each), low (1-5 transgene copies; n = 4 independent events; 5
plants each) and high (9-35 transgene copies, n = 2 independent events; 5 plants each) transgene
copy number. We identified a significant and stepwise increase in the number of RBS with
transgenic copy number, from 0.8 RBS (zero copy) to 1.6 RBS (low copy; $P = 0.078$ vs zero
copy) and to 4.3 RBS (high copy; $P < 0.0001$ vs zero copy) (Figure 5D; Supplemental Table
S9). The low copy number lines had an average increase of 0.8 RBS with respect to the zero
copy number lines, equivalent to the average difference between the *VRT-A2a* and *VRT-A2b*
NILs in the field (*VRT2-A2b* effect of +1.1 RBS). The high copy number lines produced on
average 4.3 RBS, which is higher than the *VRT-A2b* NILs and similar to the number of RBS
observed in *T. polonicum* (3.75 ± 0.62 RBS; n = 16 spikes). The dosage-dependent effects
observed in the transgenic lines provide further evidence that elevated expression of *VRT-A2*
leads to increased number of rudimentary basal spikelets in polyploid wheat.

DISCUSSION

High-resolution spatial transcriptomics in crops

We hypothesised that the establishment of the lanceolate shape in wheat spikes could be manifested in gene expression differences between the apical, central and basal sections of a developing spike, as has been shown using qRT-PCR for individual genes in wheat (*AP2*; Debernardi (2017)) and barley (*VRS2*, Youssef et al. (2017)). However, currently available transcriptome data (e.g., Li et al. (2018) and Feng et al. (2017)) lack the spatial resolution *within* each individual developmental stage to answer this question. This focus on 'between stage' comparisons (as opposed to within a single stage) is perhaps related to the technical challenges of dissecting and sectioning young meristems. Given the relatively small size of these spike meristems (0.2 mm length at Transition Stage; 3 mm length at Terminal Spikelet stage), RNA-seq methods require bulking of multiple individuals (usually between 30 and 50 different plants) to accumulate enough tissue for a single RNA-seq sample. If one sought to further section each meristem, this would require even further bulking. While laborious, this is achievable; however, under this scenario, the challenge is to properly stage ~100 plants to an equivalent developmental stage. Furthermore, it can be technically challenging to section these young spikes each time into the exact same apical, central and basal sections. Consequently, the spatial resolution in gene expression within a wheat spike at individual developmental stages has remained largely uncharacterised to date.

To address this challenge, we adapted the G&T method for micro-scale spatial-transcriptomics workflow (Macaulay et al., 2015; Giolai et al., 2019), to conduct RNA-seq of the apical, central and basal sections of individual, hand-dissected wheat spikes. This highly-automated workflow requires low tissue input and allowed us to combine 24 Nextera libraries into a single Illumina NovaSeq lane. For 19 out of the 24 samples the method worked successfully, determined by >20,000 expressed genes per library and the clustering among biological replicates. We found that the number of expressed genes per library was on average similar to the number of genes reported for bulked whole spike RNA-seq samples (Feng et al., 2017; Li et al., 2018). This is consistent with the fact that the hand-dissected sections are composed of a large mixture of different tissues (e.g., rachis, spikelet, and floret primordia) and cell types, which in the equivalent maize ears have distinct expression profiles (Xu et al., 2021). Compared to previous bulk RNA-seq studies in developing wheat spikes, the variation observed here (measured as CV) was high among biological replicates (Supplemental Figure S2). This variation is likely

caused by both biological variation (e.g., inherent variation of individual plants) and technical variation (e.g. inaccuracies in sectioning and in the developmental staging of the plant/spike) as well as the number of replicates in our analysis. A minimum of six replicates has been proposed for bulked RNA-seq (Schurch et al., 2016). Our results suggest that the RNA-seq from these small sections would benefit from a higher number of biological replicates, which should be feasible considering the high-throughput method employed for RNA extraction and library preparation, the low tissue input requirement, and the possibility to pool multiple biological replicates per sequencing lane. Despite some limitations, we could identify over 5,000 DEGs between the spatial sections for subsequent functional analysis.

In addition to G&T-Seq, several other technologies have been proposed for obtaining high resolution transcriptional profiles of plant tissues, for example, single cell RNA-seq (McFaline-Figueroa et al., 2020; Rich-Griffin et al., 2020), FACS, and the isolation of nuclei tagged in specific cell types (INTACT). These methods, however, are not spatially resolved as the complete tissue is dissolved into single cells for barcoding or selection (Rich-Griffin et al., 2020). Thus, these current methodologies do not allow, for example, to investigate whether the cell type composition of spikelets differs across the inflorescence. This would only be possible if spikelets were 'harvested' individually, for example through laser capture microdissection (LCM) before dissolving the tissue further into individual cells. Thiel et al. (2021) recently combined LCM followed by RNA-seq of the distinct lower/leaf ridge and upper/spikelet ridge of barley spikes. This allowed them to identify precise spatio-temporal expression patterns of many genes related to architecture and yield in barley spikes with unprecedented resolution. Looking ahead, increased resolution of Spatial Transcriptomics (currently 100 μm; (Giacomello et al., 2017)), which quantifies full transcriptomes while maintaining tissue integrity, offers the true prospect of direct localisation and quantification of gene expression. Our results argue strongly for the need of these transcriptome-wide and spatially resolved approaches to advance our biological understanding of fundamental developmental processes in plants.

The composite nature of spikes

Early morphological studies of wheat spike development described that the stronger elongation of central spikelets during their initial establishment (glume primordia stage) first causes the lanceolate shape of the wheat spike (Bonnett, 1966). The continuous formation of primordia at

the tip of the spike means that at any given growth stage, spikelets in different developmental

stages will be present across the spike (Bonnett, 1966). In this study, we detected more

differentially expressed genes between the three spatial sections of the spike (apical, central

and basal) than between the two investigated developmental stages (Double Ridge and Glume

Primordia). We identified 215 DEGs between the two developmental stages, consistent with

Li et al. (2018) who identified 206 DEGs between consecutive stages across a time course of

six inflorescence development stages. Feng et al. (2017) identified 753 DEGs between the

Double Ridge and Floret Primordia stage, which are further apart in development than the

stages used in this study. They also detected fewer DEGs when comparing early stages than

between more developed spikes. By contrast, we identified 1,315 and 2,438 unique genes to be

differentially expressed between the apical and basal section at DR and GP, respectively. The

higher number of DEGs between spatial sections could be due to the developmental gradients

occurring in the three spatial sections, which are revealed by the spatial sampling. These

differences would be blurred when comparing whole inflorescences between stages due to the

mixture of tissue types and spikelets at different developmental stages. A possible improvement

for future transcriptome studies could be the collection of only central sections of the

developing spikes or complete spatial sampling as conducted here.

The composite nature of the inflorescence tissues has been acknowledged by studies in maize

(ears and tassels), where new meristems are initiated in a stepwise manner. Leiboff and Hake

(2019) quantified the meristematic tissue composition of maize and sorghum tassels. For

example, maize tassels in the second stage are mainly composed of spikelet pair meristems,

but also contain some meristems in spikelet and inflorescence state. They concluded that the

changes in these tissue compositions over time correlated well with the independently staged

transcriptional changes of the tassels. Eveland et al. (2014) showed that the range of

developmental ages across the maize ear, if acknowledged, can be used as an advantage in

RNA-seq studies. They sequenced the tip, middle, and basal sections of 10-mm long ears

independently, aiming to analyse the expression patterns in specific developmental meristem

types enriched in these sections (inflorescence, spikelet, and floral meristems, respectively).

The dissection of the ear therefore allowed them to study gene expression specifically for each

meristematic tissue type rather than for all meristem types in intact ears. In this study, we

observed that apically expressed genes are enriched for GO-terms related to "shoot system

development" and "maintenance of floral organ identity". This is consistent with the hypothesis

that the apical part of the inflorescence is younger and undergoing early phases of spikelet
development initiation compared to the central inflorescence section.

Delayed transition of basal spikelets from vegetative to floral developmental programmes
We detected transcriptional gradients across the spike, with the basal section deviating most
strongly from the rest of the spike. We noticed that both *SVP* and *CENTRORADIALIS* (*CEN*)
genes remained highly expressed in the basal section of the spike, whereas their expression was
lower in the central and apical sections. *In-situ* hybridisation of these genes also showed that
their expression is strongest in vegetative tissue and basal spikelets in early spike development
(Li et al., 2021). In contrast, *SEP1-4* and *SEP1-5* genes were expressed in the opposite gradient
and showed the strongest expression in apical and central sections of the spike at Glume
Primordia stage. Recent studies allow us to interpret these gradients in the context of the early
steps of vegetative to floral growth transition. In wheat (Li et al., 2021; Adamski et al., 2021;
Liu et al., 2021), rice (Sentoku et al., 2005; Lee et al., 2008), and barley (Trevaskis et al., 2007),
SVPs have been characterised to be associated with vegetative growth and are downregulated
upon floral transition.

In wheat, the double *SVP* mutant *vrt2svp1* leads to the formation of axillary inflorescences (Li
et al., 2021). Similarly, overexpression of *CEN-D2* (*TaTFL1-2D*) in wheat extends the duration
of the Double Ridge stage (Wang et al., 2017), whereas loss-of-function mutations in barley
CEN suggest they repress floral development under short-day conditions (Bi et al., 2019).
Double knockout mutants of the MADS-box *SQUAMOSA* genes *vrn1ful2* highlighted that
these two genes act as transcriptional repressors of *SVP* and *CEN* genes in early wheat spike
development (Li et al., 2019). Furthermore, through a series of genetic and biochemical studies,
Li et al. (2021) showed that the downregulation of *SVP* genes is necessary for the formation of
flowering promoting MADS-box protein complexes including VRN1, FUL2 and SEP proteins.
Hence the coordinated downregulation of *SVPs*, and possibly *CEN* genes, along with the
upregulation of *SEP* genes is required for normal floral transition and spikelet development in
wheat. Previous studies in rice have found similar expression patterns, as well as mutant effects,
of *SVPs* and *SEPs* suggesting a conserved function in flowering transition across the two
species (Ren et al., 2016; Wu et al., 2018).

Based on our results, the floral developmental programme across the wheat spike appears to be
most advanced in its apical and central sections, while being delayed in the basal sections. We

hypothesise that this is due to elevated *VRT2* expression at the base of the spike, which hinders
the progression of the flowering programme via *SEP* class flowering genes. Likewise, the
higher expression levels of the wheat *CEN2* and *CEN5* homologs at the base are consistent
with a delay in floral transition that could interfere with the development of the spikelet
primordia. Therefore, although the basal spikelet primordia are initiated first chronologically,
their developmental age in terms of the floral programme is delayed with respect to the more
recently formed central and apical spikelet primordia. This could explain in part why the
spikelet primordia in the basal region of the spike elongate less and develop slower than central
spikelets despite being initiated first (Bonnett, 1966). Likewise, the less advanced floral
developmental programme could also explain why the overexpression of *SVPs* in barley
(*HvBM10*) leads to complete floral reversion in basal but not apical spikelets (Trevaskis et al.,
2007).

We hypothesise that *SVPs* need to be downregulated upon floral transition to allow timely
establishment and progression of the early spikelet primordia. Failure to do so would delay
their development and result in their final rudimentary shape in the mature spike. In line with
this hypothesis, we observed increased RBS in genotypes with prolonged and increased *VRT2*
expression in a dosage-dependent manner. In our qRT-PCR data, we also observe increased
expression of *SVPs* alongside reduced expression of *SEPs* in *VRT-A2b* lines at Double Ridge
and Glume Primordia stage. However, we cannot exclude the possibility that the increase in
VRT2 expression in the *VRT-A2b* lines could also affect basal spikelet development at a later
stage of spike developmental (e.g., from Terminal Spikelet stage to anthesis). We are currently
using quantitative live imaging to compare cellular growth dynamics of spikelets at different
stages of spike development between the *VRT2* NILs.

The finding that the expected downregulation of *SVP*s and *CEN*s does not follow the
chronological age of the tissues suggests that other gradients across the spike might influence
spikelet development. Debernardi et al. (2017) showed that in tetraploid wheat *AP2-5* and
miR172 have consistent and opposing expression gradients across the spike at three
consecutive developmental stages. The persistent expression gradient of *AP2-5* supports the
idea that expression patterns across the spike, beyond the ones caused by age differences of
spikelets, exist. Furthermore, they proposed a model illustrating that the phenotypic effect of
mutants across the spike differs due to the existing gradient of expression of this gene
(Debernardi et al., 2017). Other examples of mutants with different phenotypic effects across
an inflorescence include *vrn1ful2* (Li et al., 2019) in wheat, *tassel sheath1* (*tsh1*, (Whipple et

al., 2010)) and *ramosa2* (*ra2,* (Bortiri et al., 2006)) in maize, *SEPALLATA* double mutant
Osmads5Osmads34 in rice (Zhu et al. 2021), as well as *many noded dwarf1* (*mnd1,* (Walla et
al., 2020)), *frizzy panicle* (Poursarebani et al., 2015), and *vrs2* in barley, which was also found
to be consistently differentially expressed across the spike (Youssef et al., 2016). *VRS2* has
been shown to maintain a basal to apical expression pattern across three, post awn initiation
developmental timepoints in barley (Youssef et al., 2016). The study of *vrs2* mutants revealed
that *VRS2* is furthermore engaged with the basal-apical patterns of auxin, cytokinin, and
gibberellin across the spike. While hormonal gradients across the spike in early development
have not been studied in great detail in wheat, they have been shown to play crucial roles in
floral induction and development in *Arabidopsis* (Reinhardt et al., 2000). Their patterns across
the spike should be investigated in future studies addressing developmental differences across
the spike.

A model for the regulation of leaf and spikelet ridge outgrowth in the base of the spike
Recently, Meir et al. (2021) proposed that in shoot apical meristems of tomato, similar to
processes during embryonic development, transient programmes are required to inhibit a
preceding setup (i.e. vegetative growth), before a new developmental program (flowering) can
be initiated. We propose that the altered gene expression and development of the basal spikelets
could be a consequence of their initiation during the transient phase between vegetative and
floral network shifts and thus being exposed to mixed signals of development. Upon floral
transition, the lower (leaf) ridge is supressed, while the growth of spikelet ridges from the
previously supressed axillary meristems is activated. Development of lower ridges subtending
all branching events is suppressed in grass inflorescences upon flowering transition (Whipple
et al., 2010). Li et al. (2019) noticed that this suppression was disrupted in the double *vrn1ful2*
and triple *vrn1ful2ful3*-null mutants, which fail to down-regulate *SVP* genes. In these mutants,
the upper spikelet meristems generate vegetative structures resembling tillers that are
subtended by bracts or leaves originating from the lower leaf ridge.

We observed that genes that were highly expressed in the basal section of the inflorescence
(cluster 4) have previously been shown to be expressed specifically in the lower/bract ridge
and before or at vegetative to floral transition. This is also supported by the GO-term
enrichment of photosynthesis related terms in cluster 4. Our tissue sections do not allow us to
distinguish lower and upper ridge tissues, however, the two ridges have been separately

collected and sequenced via LCM in barley (Thiel et al., 2021). In this barley dataset, we found
a higher expression of *HvVRT2 (HORVU7Hr1G036130)* and *FLOWERING LOCUS C*
(HvFLC; HORVU7Hr1G054320) in the lower ridge compared to the upper ridge, whereas
HvSVP1 (*HORVU6Hr1G077300*) was also marginally more highly expressed (Supplemental
Figure S5). Furthermore, the barley *MND1* gene (*HORVU7Hr1G113480*) has recently been
shown to be expressed in leaf primordia and during the Double Ridge stage in the basal region
of the spike in barley (Walla et al., 2021), while it is most highly expressed in the vegetative
meristem and lower/leaf ridge in the LCM data (Thiel et al 2021). We observed that in our data,
the wheat *MND1* orthologs (*TraesCS7A02G506400*, *TraesCS7B02G413900*,
TraesCS7D02G494500) were significantly more highly expressed in the basal section than the
apical section at both DR and GP stage (Supplemental Table S2). The suppressed leaf ridge (or
bract) has been proposed to act as a signalling centre, regulating the fate of the upper spikelet
meristem ridge (Whipple, 2017). Insufficient bract suppression during the formation of the
basal spikelets might therefore negatively affect initiation and development of spikelets.

At DR, the widest point of the spike is indeed as expected the base and not the central section
(Figure 1A). The lower ridge is however much less developed in the central section and can be
hardly seen in the apical ridges. Interestingly, mutants failing to repress the lower ridge growth,
such as *third outer glume1* (*trd1*), the barley ortholog of maize *tsh1*, develop large bracts from
the lower ridge in basal spikelets, unlike apical spikelets, which do not develop bracts from
their lower ridges regardless of the absence of *TRD*. This is reminiscent of the gradient in the
strength of the phenotypic effects observed from the top to the base of the inflorescence in
multiple *Poaceae* mutants (discussed above). We therefore hypothesise that the basal
meristems develop into smaller spikelets and larger bract primordia due to a slow suppression
of "vegetative growth signals" (e.g., *SVPs*) and a concomitant slow upregulation of "floral
growth signals" (e.g., *SEPs*) upon floral transition. To investigate how a change from
vegetative to floral signalling might affect the development of individual meristems, we
modelled the genetic interaction of *SVPs* and *SEPs,* as proposed by Li et al. (2021), in the
spatial context of a growing spike (Supplemental File S1). Under the assumptions that SVP
supresses *SEP* expression, *SVP* expression is downregulated upon flowering, and that SEP
promotes spikelet outgrowth, the model could recapitulate (a) the observed opposing gradients
in expression of *SVPs* and *SEPs* along the spike, and (b) the formation of a lanceolate shaped
wheat spike with reduced spikelet elongation and stronger bract growth in the most basal

spikelets. Thus, whilst this hypothesis will require further investigation and testing, modelling
supports its plausibility.

Materials and Methods

Plant materials

Hexaploid wheat (*Triticum aestivum*) germplasm used in this study includes wildtype
hexaploid wheat cultivar Paragon and *P1/VRT2 germplasm* described in Adamski et al. (2021)
including *P1* NILs, recombinants, and T_1 transgenic lines carrying the *T. polonicum VRT-A2b*
copy under the native promoter. *T. polonicum* accession T1100002 was obtained from the John
Innes   Centre   Germplasm   Resources   Unit   (https://www.seedstor.ac.uk/search-
infoaccession.php?idPlant=27422). For field experiments, we used between two to four sibling
BC_4/ BC_6 NILs differing for the *VRT-A2b* allele.

Low input RNA sequencing

Paragon seedlings were grown in a single batch in a controlled environment growth chambers
in 24-cell seed trays under long-day (16 h light/8 h dark) photoperiods at 300 µmol m^{-2} s^{-1},
with a day temperature of 20 °C and a night temperature of 15 °C. Inflorescences for Double
Ridge (DR) stage were collected 18 days after sowing, while inflorescences for Glume
Primordia (GP) stage were collected 22 days after sowing. All plants were grown in "John
Innes Cereal Mix" (40% Medium Grade Peat, 40% Sterilized Soil, 20% Horticultural Grit, 1.3
kg·m^{-3} PG Mix 14-16-18 + Te Base Fertiliser, 1 kg·m^{-3} Osmocote Mini 16-8-11 2 mg + Te
0.02% B, Wetting Agent, 3 kg·m^{-3} Maglime, 300 g·m^{-3} Exemptor).

Four individual spikes per developmental stage (DR and GP) were dissected into apical,
central, and basal sections (1:1:1 ratio) using a stereo microscope (Leica MZ16). Sections were
immediately placed into 96-well plates (on ice) containing 10 µL of RLT plus (Qiagen, Hilden,
Germany). All instruments and surfaces were cleaned with 80% ethanol, RNAse-free water
and lastly RNAse-out solution after each sample to reduce cross-contamination and RNA
degradation. Samples were stored at -80 °C until cDNA preparation, using the G&T-seq
method as previously described (Macaulay et al., 2015). cDNA was normalised to 0.2 ng/ µL
before Nextera (Illumina, San Diego, CA, USA) library preparation using a Mosquito HV

liquid handler (STP, Royston, UK) in a total reaction volume of 4 μL as described in Mora-

Castilla et al. (2016). Libraries were pooled by volume and sequenced on a single lane of a

NovaSeq 6000 (NVS200S2 flow cell, 100 bp paired-end reads).

Bioinformatic analysis

For the RNA-seq analysis, we used the RefSeqv1.0 genome assembly and the RefSeqv1.1 gene

annotation (https://urgi.versailles.inra.fr/download/iwgsc/IWGSC_RefSeq_Assemblies/v1.0/;

IWGSC et al. (2018)). Reads were trimmed and adapters were removed using trim-galore

v.0.4.2 (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) with settings: "--

paired --fastqc --a GGTATCAACGCAGAGT --clip_R1 20 --clip_R2 20 --trim-n". Minimum

length of reads retained was set to 50 bp. Reads were aligned to the RefSeqv1.0 genome

assembly using HISAT2 v. 2.1.0 (https://daehwankimlab.github.io/hisat2/; Kim et al. (2019))

with the following parameters: "--pen-noncansplice 20 --mp 1,0 --rna-strandness RF".

Alignment files were converted to BAM format, sorted, indexed, filtered, and purged of all

none-primary alignments (0x100 flag) using samtools (v. 1.9; Li et al. (2009)). HTSeq v.0.6.1

(https://htseq.readthedocs.io/en/master/; Anders et al. (2015)) was used to count the read

numbers mapped to the RefSeqv1.1 gene models.

HT-read count normalization and differential expression analyses were performed using the

DESeq2   v.1.28.1   R   packages

(https://bioconductor.org/packages/release/bioc/html/DESeq2.html; Love et al., 2014; RStudio

1.2.5001). Genes with an average expression below 10 HT-count, and which were not

expressed (i.e. ≤ 10 HT counts) in at least three libraries, were removed from the analysis.

Correlation between expressed genes and Waddington stage and/or section was tested by

ANOVA. Raw read data from Li et al. (2018) and Feng et al. (2017) were pseudo-aligned

using Kallisto Sleuth pipeline (https://scilifelab.github.io/courses/rnaseq/labs/kallisto) and the

coefficient of variation was calculated for each gene (by condition) using R (RStudio 1.2.5001)

ddply (plyr 1.8.6). Differentially expressed genes (DEGs) between the two Waddington stages

(DR and GP) were calculated with the design "~plant + section", while DEGs between the

three sections (apical, central, basal) were determined using the design "~section +

waddington:plant + waddington". DEGs among the sections within each Waddington stage

were determined with the design "~plant + section". For each gene, an adjusted *P*-value was

computed by DESeq2 (using the using the Benjamini and Hochberg method (Benjamini and

Hochberg, 1995)), and those with an adjusted *P*-value of ≤ 0.05 were considered differentially

expressed. DeSeq2 also computed Log2FoldChanges as well as the associated uncertainty

(lfcSE, see Love et al. (2014) for further detail). The "contrast" function was used to determine

pairwise comparison P-values. The full set of expression data and comparisons is presented in

Supplemental Data Set S1. Enrichment of GO-terms was performed using the online tool

"PLAZA" (https://bioinformatics.psb.ugent.be/plaza; Van Bel et al. (2017)) using the

recommended settings, and all enriched GO-terms of Biological function (BF) and Cellular

Compartment (CC) were retained. In brief, PLAZA determines the overrepresentation of a

certain GO-term in a gene set compared to the genome-wide background frequency (= all

expressed genes in this experiment; submitted manually). The significance of over- or

underrepresentation is determined using the hypergeometric distribution and the Bonferroni

method is applied to correct for multiple testing. Note that enrichment folds are reported in

log2 fold scale. Enrichment of TF families (Genes that were annotated as TFs were obtained

from https://opendata.earlham.ac.uk/wheat/under_license/toronto/Ramirez-

Gonzalez_etal_2018-06025-Transcriptome-Landscape/data/data_tables/ (Ramirez-Gonzalez

et al., 2018)) and MADS-box TFs (based on Schilling et al. (2020)) was performed in R using

the phyper() function from stats package v.4.0.1 to test for Hypergeometric Distribution. All

DEGs were scaled and centred using R-base function "scale". All cluster analysis was

performed on scaled data using R (stats) functions kmeans and hclust, followed by visualisation

through pheatmap v.1.0.12 (https://cran.rstudio.com/web/packages/pheatmap/index.html).

Correlation to centroid cluster shape of each gene expression pattern was calculated using the

"cor" function from R stats.

Quantitative real-time PCR analysis

P1 NILs were grown in controlled growth chambers in 24-cell seed trays under the same

conditions as used in the low input RNA-seq experiment (see above). For each biological

replicate, we pooled 30 inflorescences for DR stage, 15 for GP stage, and nine for Terminal

Spikelet stage (n = 4 biological replicates per stage). Inflorescences from NILs were dissected

using a stereo microscope (Leica MZ16). Inflorescences were dissected into apical, central,

basal and peduncle sections (1:1:1:1 ratio). At Double Ridge stage, inflorescences were only

dissected into apical, basal and peduncle section as the inflorescences were too small to be

accurately dissected into four sections for all 30 plants per biological replicate. Each section

was immediately placed into 1.5-mL tubes on dry ice and tubes were snap frozen in liquid

nitrogen as soon as all plants for the sample were collected. Samples were stored at −80°C until

needed. Inflorescences were collected within 2-3 hours, 9 hours after the lights came on in the

growth chamber. Tissue was homogenized in a TissueLyser II (Cat No.: 85300, QIAGEN)

using 3-mm steel beads (Cat No.: 69997, Qiagen); tubes were shaken for 20-s at 28 Hz with

dry ice.

All RNA extractions were performed using the RNeasy Plant Mini Kit (Cat No.: 74904,

Qiagen) with RLT buffer according to the manufacturer's protocol followed by RNA ethanol

precipitation   (https://projects.iq.harvard.edu/files/hlalab/files/ethanol-precipitation-of-

rna_hla.pdf). DNA digestion was performed using the RQ1 RNase-free DNase set (Cat No.:

M6101, Promega) according to the manufacturer's protocol. RNA was reverse transcribed

using M-MLV reverse transcriptase (Cat No.: 28025013, Thermofisher) according to the

manufacturer's protocol. For the qRT-PCR reactions, LightCycler 480 SYBR Green I Master

Mix (Roche Applied Science, UK) was used according to the manufacturer's protocol. The

reactions were run in a LightCycler 480 instrument (Roche Applied Science, UK) under the

following conditions: 10 min at 95 °C; 40 cycles of 10 sec at 95 °C, 15 sec at 62 °C, 30 sec at

72 °C; dissociation curve from 60 °C to 95 °C to confirm primer specificity. All reactions were

performed with three technical replicates per sample and using *TaActin* as the reference gene

(Uauy et al., 2006). Relative gene expression was calculated using the $2^{-\Delta\Delta Ct}$ method (Livak

and Schmittgen, 2001) with a common calibrator so that values are comparable across genes,

tissues, and developmental stages. All primers used in qRT-PCR came from Adamski et al.,

2021 can be found in Supplemental Table S10.

All qRT-PCR data was normalised using a log2 transformation. A three-way ANOVA

including Waddington stage, section, and genotype yielded significant two-way interactions.

The differences between sections of the genotypes were therefore further analysed individually

for each Waddington stage and genotype. For each of the two genotypes we individually

performed Tukey multiple comparison tests to determine differences between the sections

within each developmental stage by Tukey multiple comparison test. Differences between the

two genotypes were also analysed individually for each Waddington stage. Furthermore, the

differences between the genotypes were investigated individually for each section within the

Waddington stage if the interaction term was significant (in GP and TS). For all analysis see

Supplemental Table S6.

Field experiments and phenotyping

VRT-A2 NILs were evaluated in four field experiments. Three trials were located at The Morley Agricultural Foundation trials site, Morley St Botolph, UK (52°33'15.1"N 1°01'59.2"E) in 2017, 2018 and 2020 and one trial was sown in 2020 at the John Innes Experimental trials site in Norwich, UK (52°37'50.7"N 1°10'39.7"E). In Morley (2017) we analysed two BC_4 lines of *VRT-A2a* and three BC_4 lines of *VRT-A2b*. In Morley (2019) we analysed two BC_6 and one BC_4 line per *VRT-A2* allele and in Morley and Church Farm 2020 we analysed two BC_6 and two BC_4 lines for each *VRT-A2* allele. All experiments were drilled as yield-scale plots (6 m x 1.2 m) and sown by grain number for comparable plant densities aiming for 275 seeds m^{-2}. The trials were arranged in a randomised complete block design (RCBD) with five replicates per sibling line per location. Developmental and plant architecture traits were evaluated throughout the growing period. A 10-ear grab sample was collected from each plot pre-harvest for the assessment of rudimentary basal spikelet (RBS) numbers and other phenotypes (recorded in Adamski et al. 2021). RBS were defined as spikelets carrying no grain at maturity and counted for each spike individually. To determine the differences between the $P1^{POL}$ and $P1^{WT}$ NILs, we performed analysis of variance (ANOVA) on the multiple field trials phenotypic data. For the analysis of individual trials, we used a two-way ANOVA including Genotype + Block performed in R ('car' package version 3.0-10; RStudio 1.2.5001).

Glasshouse phenotyping

We evaluated the BC_4 NILs and BC_6F_3 recombinant lines, as well as *T. polonicum* accession T1100002, under standard glasshouse conditions. 18-20 plants per genotype were grown in 1 L pots containing John Innes Cereal Mix under long day conditions (16 h light, 8 h dark). The genotypes of all plants were confirmed using KASP marker *SP1Pol* (Adamski et al., 2021).We counted the number of rudimentary basal spikelets (RBS) for all tillers of all biological replicates at maturity. To evaluate the differences in RBS between genotypes, we performed a two-way ANOVA analysis and post-hoc multi-pairwise comparisons Sidak test ('car' package version 3.0-10; RStudio 1.2.5001).

Phenotyping of transgenic lines

T_1 lines from Adamski et al. (2021) differing for the copy number of the *VRT-A2b* transgenic construct (zero = 0 copies; low = 1-5 copies; high = 9-35 copies) were grown in 1 L pots with

John Innes Cereal Mix under 16 h light at 20°C and 8 h dark at 15°C in controlled environment

growth chambers. We measured RBS number for the main tiller of all plants at maturity. To

determine differences in RBS between the three transgenic classes, we performed analysis of

variance (two-way ANOVA; 'car' package version 3.0-10). We performed Dunnett tests to

compare the low and high copy lines against the zero copy number controls (RStudio 1.2.5001).

Modelling

The computational model of wheat spike shape formation was developed using the multi-agent

programming language and modelling environment, Netlogo (Wilensky, 1999). Gene

interactions were modelled as previously described (Li et al., 2021). The model can be accessed

via the interactive web-version of the model (Supplemental File S1).

In brief, both spikelets and leaves are initiated with rates that depend on the levels of SEP. Leaf

initiation rates are suppressed by SEP, whereas spikelet initiation requires SEP. The maximum

initiation rates are the same for both spikelets and leaves but different before ($r_{vegetative}$) and

after ($r_{flowering}$) flowering. Once initiated, the leaves and spikelets grow at a rate defined by the

parameters r_{leaf} and $r_{spikelet}$, respectively. Leaf growth does not depend on SVP or SEP levels,

whereas spikelets only increase in size every iteration if their SEP level is above a given

threshold ($SEP_{growth\_threshold}$). Expression of both *SVP* and *SEP* only occurs at meristem

initiation. After this, the levels of SVP and SEP cannot increase, although SEP is degraded.

SVP is not degraded, solely because at this point, nothing is dependent on SVP levels, whilst

spikelet growth depends upon SEP levels.

SVP expression rates start to decrease, once flowering is triggered, according to:

854
$$r_{SVP}(t+1) = r_{SVP}(t) f_{red}$$

where $r_{SVP}(t)$ is the rate of *SVP* expression at that time step, and f_{red} is a rate reduction factor.

SEP expression depends upon the levels of SVP in the meristem in which the initiation points

are located, depending on a Hill function (Alon, 2007),

$r_{SEP}(SVP) = r_{SEP,max}\left(\frac{K_D{}^n}{K_D{}^n + SVP^n}\right),$

where $r_{SEP,max}$ is the maximum rate of *SEP* expression, K_D is the binding constant, and *n* is the

Hill coefficient. The resulting curves for *SVP* and *SEP* expression are shown in Supplemental

Figure S6.

SVP levels are initiated with the current value of r_{SVP}. SEP levels are initiated using r_{SEP}, and
reduce by degradation rate, δ_{SEP}, following

$$\frac{dSEP}{dt} = SEP \cdot \delta_{SEP}.$$

Data Availability
The raw RNA-seq read libraries used in this study are available from NCBI BioProject
PRJNA749586.

Acknowledgments
We thank Phil Robinson from the Scientific Photography at JIC for taking high-quality images
of spikes and spikelets and Tobin Florio (http://flozbox-science.com/) for scientific
illustrations for Figure 1. We thank the JIC Field Experimentation and Horticultural Services
teams for technical support in field and glasshouse experiments and the JIC Bioimaging facility
and staff for their services and technical support.

Competing Interest
The authors declare no competing interest.

Additional Files
Supplemental Figure S1: Principal Component analysis (PCA) of RNA-seq libraries.

Supplemental Figure S2: Comparison of Coefficient of variation (CV) of gene expression in
wheat RNA-seq data sets.

Supplemental Figure S3: Expression patterns of al DEGs clusters (n=7) as identified by k-
means clustering.

Supplemental Figure S4: Dissected Floret 1 and 2 of basal and central spikelets of BC_6 NILs
before anthesis.

Supplemental Figure S5: Expression of barley genes in Thiel et al. (2021), which are orthologous to wheat genes highly expressed in basal spike sections.

Supplemental Figure S6: Example of Netlogo simulation outcome with default parameters.

Supplemental Table S1: Quality control measurements of all 24 RNA-seq libraries.

Supplemental Table S2: Summary of normalised gene expression, statistical analyses and k-means clustering for all differentially expressed genes ($Padj \leq 0.05$).

Supplemental Table S3: Enrichment of Gene Ontology (GO)-terms in the seven identified clusters of DEGs.

Supplemental Table S4: Enrichment of Transcription factor families and MADS-box transcription factor genes in the seven identified clusters of DEGs.

Supplemental Table S5: Relative expression of *VRT-A2* and *SEP1-4* measured in Paragon NILs with either the wildtype (*VRT-A2a*) or *T. polonicum* allele (*VRT-A2b*).

Supplemental Table S6: Statistical analysis of qRT-PCR data from *VRT-A2* NIL spike sections (Supplemental Table S5).

Supplemental Table S7: Field evaluations for rudimentary basal spikelets (RBS) in *VRT-A2* NILs.

Supplemental Table S8: Graphical genotype from Adamski et al (2021) and RBS phenotype of BC$_6$ recombinant inbred lines (RILs) for *VRT-A2*.

Supplemental Table S9: Rudimentary basal spikelet phenotypic data from *VRT-A2* transgenic lines.

Supplemental Table S10: List of primers used in qRT-PCR in this study.

Supplemental Dataset S1: Summary of normalised gene expression, statistical analyses and k-means clustering for all expressed genes.

Supplemental File S1: Interactive web-version of the wheat spike model.

References

Adamski NM, Simmonds J, Brinton JF, Backhaus AE, Chen Y, Smedley M, Hayta S, Florio T, Crane P, Scott P, Pieri A, Hall O, Barclay JE, Clayton M, Doonan JH, Nibau C, Uauy C (2021) Ectopic expression of *Triticum polonicum VRT-A2* underlies elongated glumes and grains in hexaploid wheat in a dosage-dependent manner. The Plant Cell **33**:2296-2319

Alon U (2007) An introduction to systems biology: design principles of biological circuits. Chapman & Hall/CRC, London

Anders S, Pyl PT, Huber W (2015) HTSeq--a python framework to work with high-throughput sequencing data. Bioinformatics **31**:166-169

Becker A, Theißen G (2003) The major clades of MADS-box genes and their role in the development and evolution of flowering plants. Molecular Phylogenetics and Evolution **29**: 464-489

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. J. R. Stat. Soc. Ser. B Stat. Method. **57**: 289–300

Bi X, van Esse W, Mulki MA, Kirschner G, Zhong J, Simon R, von Korff M (2019) *CENTRORADIALIS* interacts with *FLOWERING LOCUS T-like* genes to control floret development and grain number. Plant Physiology **180:** 1013-1030

Bommert P, Whipple C (2018) Grass inflorescence architecture and meristem determinacy. Seminars in Cell & Developmental Biology **79:** 37-47

Bonnett OT (1966) Inflorescences of maize, wheat, rye, barley, and oats: their initiation and development. Bulletin (University of Illinois (Urbana-Champaign campus) Agricultural Experiment Station) **721**

Bortiri E, Chuck G, Vollbrecht E, Rocheford T, Martienssen R, Hake S (2006) *ramosa2* Encodes a LATERAL ORGAN BOUNDARY Domain Protein That Determines the Fate of Stem Cells in Branch Meristems of Maize. The Plant Cell **18:** 574-585

Debernardi JM, Lin H, Chuck G, Faris JD, Dubcovsky J (2017) microRNA172 plays a crucial role in wheat spike morphogenesis and grain threshability. Development **144:** 1966-1975

Eveland AL, Goldshmidt A, Pautler M, Morohashi K, Liseron-Monfils C, Lewis MW, Kumari S, Hiraga S, Yang F, Unger-Wallace E (2014) Regulatory modules controlling maize inflorescence architecture. Genome Research **24**: 431-443

Feng N, Song G, Guan J, Chen K, Jia M, Huang D, Wu J, Zhang L, Kong X, Geng S, Liu J, Li A, Mao L (2017) Transcriptome Profiling of Wheat Inflorescence Development from Spikelet Initiation to Floral Patterning Identified Stage-Specific Regulatory Genes. Plant Physiology **174:** 1779-1794

Giacomello S, Salmén F, Terebieniec BK, Vickovic S, Navarro JF, Alexeyenko A, Reimegård J, McKee LS, Mannapperuma C, Bulone V (2017) Spatially resolved transcriptome profiling in model plant species. Nature Plants **3**: 17061

Giolai M, Verweij W, Lister A, Heavens D, Macaulay I, Clark MD (2019) Spatially resolved transcriptomics reveals plant host responses to pathogens. Plant Methods **15**: 114

Harrop TWR., Ud Din I, Gregis V, Osnato M, Jouannic S, Adam H, Kater MM (2016) Gene expression profiling of reproductive meristem types in early rice inflorescences by laser microdissection. Plant Journal **86**: 75-88

IWGSC, Appels R, Eversole K, Stein N, Feuillet C, Keller B, Rogers J, Pozniak CJ, Choulet F, et_al. (2018) Shifting the limits in wheat research and breeding using a fully annotated reference genome. Science **361**: eear7191

Kellogg E, Camara P, Rudall P, Ladd P, Malcomber S, Whipple C, Doust A (2013) Early
inflorescence development in the grasses (Poaceae). Frontiers in Plant Science **4:** 250

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. Nature Biotechnology **37:** 907-915

Koppolu R, Schnurbusch T (2019) Developmental pathways for shaping spike inflorescence
architecture in barley and wheat. Journal of Integrative Plant Biology **61:** 278-295

Lee S, Choi SC, An G (2008) Rice SVP-group MADS-box proteins, OsMADS22 and OsMADS55,
are negative regulators of brassinosteroid responses. Plant Journal **54:** 93-105

Leiboff S, Hake S (2019) Reconstructing the transcriptional ontogeny of maize and sorghum supports
an inverse hourglass model of inflorescence development. Current Biology **29:** 3410-3419. e3413

Li C, Lin H, Chen A, Lau M, Jernstedt J, Dubcovsky J (2019) Wheat *VRN1*, *FUL2* and *FUL3* play
critical and redundant roles in spikelet development and spike determinacy. Development **146:**
dev175398

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
Subgroup GPDP (2009) The Sequence Alignment/Map format and SAMtools. Bioinformatics **25:**
2078-2079

Li K, Debernardi JM, Li C, Lin H, Zhang C, Jernstedt J, von Korff M, Zhong J, Dubcovsky J
(2021) Interactions between SQUAMOSA and SHORT VEGETATIVE PHASE MADS-box proteins
regulate meristem transitions during wheat spike development. The Plant Cell **33:** 3621–3644

Li Y, Fu X, Zhao M, Zhang W, Li B, An D, Li J, Zhang A, Liu R, Liu X (2018) A Genome-wide
View of Transcriptome Dynamics During Early Spike Development in Bread Wheat. Scientific Reports
8: 15338

Liu J, Chen Z, Wang Z, Zhang Z, Xie X, Wang Z, Chai L, Song L, Cheng X, Feng M, Wang X,
Liu Y, Hu Z, Xing J, Su Z, Peng H, Xin M, Yao Y, Guo W, Sun Q, Liu J, Ni Z (2021) Ectopic
expression of *VRT-A2* underlies the origin of *Triticum polonicum* and *Triticum petropavlovskyi* with
long outer glumes and grains. Molecular Plant **14:** 1472-1488

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biology **15:** 550

Macaulay IC, Haerty W, Kumar P, Li YI, Hu TX, Teng MJ, Goolam M, Saurat N, Coupland P,
Shirley LM, Smith M, Van der Aa N, Banerjee R, Ellis PD, Quail MA, Swerdlow HP, Zernicka-
Goetz M, Livesey FJ, Ponting CP, Voet T (2015) G&T-seq: parallel sequencing of single-cell
genomes and transcriptomes. Nature Methods **12:** 519-522

McFaline-Figueroa JL, Trapnell C, Cuperus JT (2020) The promise of single-cell genomics in
plants. Current Opinion in Plant Biology **54:** 114-121

Meir Z, Aviezer I, Chongloi GL, Ben-Kiki O, Bronstein R, Mukamel Z, Keren-Shaul H, Jaitin D,
Tal L, Shalev-Schlosser G, Harel TH, Tanay A, Eshed Y (2021) Dissection of floral transition by
single-meristem transcriptomes at high temporal resolution. Nature Plants **7:** 800-813

Mora-Castilla S, To C, Vaezeslami S, Morey R, Srinivasan S, Dumdie JN, Cook-Andersen H,
Jenkins J, Laurent LC (2016) Miniaturization technologies for efficient single-cell library preparation
for next-generation sequencing. Journal of laboratory automation **21:** 557-567

Poursarebani N, Seidensticker T, Koppolu R, Trautewig C, Gawroński P, Bini F, Govind G,
Rutten T, Sakuma S, Tagiri A, Wolde GM, Youssef HM, Battal A, Ciannamea S, Fusca T,
Nussbaumer T, Pozzi C, Börner A, Lundqvist U, Komatsuda T, Salvi S, Tuberosa R, Uauy C,
Sreenivasulu N, Rossini L, Schnurbusch T (2015) The Genetic Basis of Composite Spike Form in
Barley and 'Miracle-Wheat'. Genetics **201:** 155-165

Prusinkiewicz P, Erasmus Y, Lane B, Harder LD, Coen E (2007) Evolution and Development of Inflorescence Architectures. Science **316**: 1452-1456

Ramirez-Gonzalez RH, Borrill P, Lang D, Harrington SA, Brinton J, Venturini L, Davey M, Jacobs J, van Ex F, Pasha A, Khedikar Y, Robinson SJ, Cory AT, Florio T, Concia L, Juery C, Schoonbeek H, Steuernagel B, Xiang D, Ridout CJ, Chalhoub B, Mayer KFX, Benhamed M, Latrasse D, Bendahmane A, Wulff BBH, Appels R, Tiwari V, Datla R, Choulet F, Pozniak CJ, Provart NJ, Sharpe AG, Paux E, Spannagl M, Brautigam A, Uauy C (2018) The transcriptional landscape of polyploid wheat. Science **361**: eaar6089

Reinhardt D, Mandel T, Kuhlemeier C (2000) Auxin Regulates the Initiation and Radial Position of Plant Lateral Organs. The Plant Cell **12:** 507-518

Ren D, Rao Y, Leng Y, Li Z, Xu Q, Wu L, Qiu Z, Xue D, Zeng D, Hu J (2016) Regulatory role of OsMADS34 in the determination of glumes fate, grain yield, and quality in rice. Frontiers in plant science **7**: 1853

Rich-Griffin C, Stechemesser A, Finch J, Lucas E, Ott S, Schäfer P (2020) Single-cell transcriptomics: a high-resolution avenue for plant functional genomics. Trends in Plant Science **25**: 186-197

Schilling S, Kennedy A, Pan S, Jermiin LS, Melzer R (2020) Genome-wide analysis of MIKC-type MADS-box genes in wheat: pervasive duplications, functional conservation and putative neofunctionalization. New Phytologist **225**: 511-529

Schurch NJ, Schofield P, Gierliński M, Cole C, Sherstnev A, Singh V, Wrobel N, Gharbi K, Simpson GG, Owen-Hughes T (2016) How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use? RNA **22:** 839-851

Sentoku N, Kato H, Kitano H, Imai R (2005) *OsMADS22*, an STMADS11-like MADS-box gene of rice, is expressed in non-vegetative tissues and its ectopic expression induces spikelet meristem indeterminacy. Molecular Genetics and Genomics **273:** 1-9

Thiel J, Koppolu R, Trautewig C, Hertig C, Kale S, Erbe S, Mascher M, Himmelbach A, Rutten T, Esteban E (2021) Transcriptional landscapes of floral meristems in barley. Science Advances **7**: eabf0832

Trevaskis B, Tadege M, Hemming MN, Peacock WJ, Dennis ES, Sheldon C (2007) *Short Vegetative Phase-Like* MADS-Box Genes Inhibit Floral Meristem Identity in Barley. Plant Physiology **143:** 225-235

Uauy C, Distelfeld A, Fahima T, Blechl A, Dubcovsky J (2006) A NAC gene regulating senescence improves grain protein, zinc, and iron content in wheat. Science **314**: 1298-1301

Van Bel M, Diels T, Vancaester E, Kreft L, Botzki A, Van de Peer Y, Coppens F, Vandepoele K (2017) PLAZA 4.0: an integrative resource for functional, evolutionary and comparative plant genomics. Nucleic Acids Research **46**: D1190-D1196

Waddington S, Cartwright P, Wall P (1983) A quantitative scale of spike initial and pistil development in barley and wheat. Annals of Botany **51:** 119-130

Walla A, Wilma van Esse G, Kirschner GK, Guo G, Brünje A, Finkemeier I, Simon R, von Korff M (2020) An Acyl-CoA *N*-Acyltransferase Regulates Meristem Phase Change and Plant Architecture in Barley1. Plant Physiology **183**: 1088-1109

Wang Y, Yu H, Tian C, Sajjad M, Gao C, Tong Y, Wang X, Jiao Y (2017) Transcriptome Association Identifies Regulators of Wheat Spike Architecture. Plant Physiology **175**: 746-757

Whipple CJ (2017) Grass inflorescence architecture and evolution: the origin of novel signaling centers. New Phytologist **216**: 367-372

Whipple CJ, Hall DH, DeBlasio S, Taguchi-Shiobara F, Schmidt RJ, Jackson DP (2010) A
Conserved Mechanism of Bract Suppression in the Grass Family The Plant Cell **22:** 565-578

Wu D, Liang W, Zhu W, Chen M, Ferrándiz C, Burton RA, Dreni L, Zhang D (2018) Loss of
LOFSEP transcription factor function converts spikelet to leaf-like structures in rice. Plant Physiology
176: 1646-1664

**Xu X, Crow M, Rice BR, Li F, Harris B, Liu L, Demesa-Arevalo E, Lu Z, Wang L, Fox N, Wang
X, Drenkow J, Luo A, Char SN, Yang B, Sylvester AW, Gingeras TR, Schmitz RJ, Ware D, Lipka
AE, Gillis J, Jackson D** (2021) Single-cell RNA sequencing of developing maize ears facilitates
functional analysis and trait candidate gene discovery. Developmental Cell **56**: 557-568.e556

**Youssef HM, Eggert K, Koppolu R, Alqudah AM, Poursarebani N, Fazeli A, Sakuma S, Tagiri
A, Rutten T, Govind G, Lundqvist U, Graner A, Komatsuda T, Sreenivasulu N, Schnurbusch T**
(2017) VRS2 regulates hormone-mediated inflorescence patterning in barley. Nature Genetics **49**: 157-
161

Zhao T, Ni Z, Dai Y, Yao Y, Nie X, Sun Q (2006) Characterization and expression of 42 MADS-box
genes in wheat (*Triticum aestivum* L.). Molecular Genetics and Genomics **276**: 334