The complex genetics of symbiotic extended phenotypes across environments in a model mutualism

Rebecca T. Batstone,[a,*] **Hanna Lindgren,**[b] **Cassandra M. Allsup,**[b] **Laura A. Goralka,**[b] **Alex B. Riley,**[b] **Michael A. Grillo,**[c] **Amy Marshall-Colon,**[a,b] **Katy D. Heath,**[a,b,*]

Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, 1206 West Gregory Drive, Urbana, IL 61801 USA[a]; Department of Plant Biology, University of Illinois at Urbana-Champaign, 286 Morrill Hall, 505 South Goodwin Avenue, Urbana, IL 61801 USA[b]; Department of Biology, Loyola University Chicago, 1032 W. Sheridan Road, Chicago, IL 60618 USA[c]

R. Batstone and H. Lindgren contributed equally. Author order was determined based on overall contribution, particularly time spent analyzing the data, as well as writing and compiling feedback.

Compiled October 27, 2021
This is a draft manuscript, pre-submission

* Co-corresponding authors:
R. Batstone, rtbatstone@gmail.com
K. Heath, kheath@iliinois.edu.

Authorship statement: KDH and AMC conceived of the work, acquired funding, and designed the experiments. MAG collected soil, performed trapping experiments, and isolated strains. ABR performed preliminary experiments and extracted DNA. HL and CMA led experimental work and data collection with assistance from LAG, ABR, KDH, RTB, and AMC. RTB, HL, and KDH performed data analysis and interpretation. KDH, RTB, and HL drafted the article. All authors participated in critical revisions and approved the final version for submission.

ABSTRACT   A goal of modern biology is to develop the genotype-phenotype (G→P) map, a predictive understanding of how genomic information generates trait variation that forms the basis of both natural and managed communities. As microbiome research advances, however, it has become clear that many of these traits are symbiotic extended phenotypes, being governed by genetic variation encoded not only by the host's own genome, but also by the genomes of myriad cryptic symbionts. Building a reliable G→P map therefore requires accounting for the multitude of interacting genes and even genomes involved in symbiosis. Here we use naturally-occurring genetic variation in 191 strains of the model microbial symbiont *Ensifer meliloti* in four mapping experiments to study the genomic architecture of a key symbiotic extended phenotype – partner quality, or the fitness benefit conferred to a host by a particular symbiont genotype, within and across environmental contexts and host genotypes. We define three novel categories of loci in rhizobium genomes that must be accounted for if we want to build a reliable G→P map of partner quality; namely, 1) loci whose identities depend on the environment, 2) those that depend on the host genotype with which rhizobia interact, and 3) universal loci that are likely important in all or most environments.

IMPORTANCE   Given the rapid rise of research on how microbiomes can be harnessed to improve host health, understanding the contribution of microbial genetic variation to host phenotypic variation is pressing, and will better enable us to predict the evolution of (and select more precisely for) symbiotic extended phenotypes that impact host health. We uncover extensive context-dependency in both the identity and functions of symbiont loci that control host growth, which makes predicting the genes and pathways important for determining symbiotic outcomes under different conditions more challenging. Despite this context-dependency, we also resolve a core set of universal loci that are likely important in all or most environments, and thus, serve as excellent targets both for genetic engineering and future coevolutionary studies of symbiosis.

KEYWORDS: GWAS, mapping, *Medicago truncatula*, *Ensifer meliloti*, symbiosis, partner quality, rhizobium, GxE, GxG, genotype-phenotype map, G→P map, root nodule, nodulation, symbiotic nitrogen fixation

INTRODUCTION

We live in a symbiotic world. It is increasingly recognized that many important traits, including human metabolism, insect diet and defense, and plant nutrient foraging, are actually *symbiotic extended phenotypes* governed (at least in part) by cryptic variation in their microbial symbionts (e.g., 1, 2, 3, 4, 5, 6, 7; reviewed by 8, 9, 10, 11, 12). Incorporating the complex genetics arising from these interactions, however, has lagged behind understanding their ecological outcomes. Characterizing the genetic basis of these cross-domain relationships represents a symbiotic extension of the genotype-phenotype (G→P) map (13, 14, 15), wherein genetic variation present in symbionts (G) presents as phenotypic variation (P) expressed by the host. In symbiosis, as in all organisms, the G→P map is a crucial step towards a predictive understanding of evolution (e.g., a trait's genomic architecture influences the rates and trajectories of its response to selection), and for bioengineering, where we must know not only which genes to target and exactly how to edit them, but also anticipate how consistent the knock-on effects will be across diverse environments and/or genetic backgrounds (16, 17).

A key lesson from studies of the G→P map within single organisms is that phenotypic variation fundamentally arises in non-additive ways as genomic variation is filtered through "layer(s) of context-dependence" (14) including genotype-by-environment interactions, epistasis, and pleiotropy (15, 18, 19, 20, 21, 22, 23). The role of context-dependency in the G→P is particularly relevant for symbiotic extended phenotypes, given the potential for higher-order interactive genetic effects. For example, interacting genomes can generate genotype-by-genotype interactions between host and symbiont (intergenomic epistasis, or G × G: 24, 25, 26, 27, 28, 29) and even G × G × E whereby trait values depend not only on the interaction of alleles in both partners, but on the environmental context (30, 31, 32). Such complex context-dependent effects are critical for evolution; for example, G × G × E for fitness is the statistical expression of the coevolutionary selection mosaic (27, 33, 34). Additionally, genes that consistently affect symbiotic extended phenotypes arising in plant-microbe interactions, independent of these layers of context dependency, are important targets for breeding and bioengineering (9, 16, 35, 36, 37). Fine-scale mapping of the loci contributing to symbiotic extended phenotypes across partner genotype and environmental contexts can therefore generate a more holistic picture of both types of genetic effects (i.e., the genes that universally contribute and those that are context-dependent), and thus, provides insights both for natural (co)evolutionary processes and sustainable agriculture.

The symbiosis between nitrogen-fixing rhizobia and leguminous plants has been a key model for addressing both additive and interactive genetic effects in symbiosis (i.e., G × G, G × E: 38, 39, 40, 41, 42), with resequencing projects, germplasm collections, and many genetic and genomic tools, particularly in *Medicago-Ensifer* interactions (43, 44, 45, 46, 47). These symbioses are also major drivers of the terrestrial nitrogen (N) cycle, contributing up to 21 million tonnes of fixed N each year (48, 49). In their specialized root nodules, legumes trade photosynthetically-derived fixed carbon in exchange for N fixed by the rhizobia, making legumes keystone members of natural plant communities and sustainable agriculture (reviewed by 50, 51). Rhizobial genomes themselves have been studied as models of bacterial genome evolution, reflecting the dynamic tension between the core set of genes, often on the main chromosome, and the more flexible genes, often found on mobile plasmids, islands, and other elements (52, 53, 54, 55, 56). Symbiosis plasmids in particular often show abundant recombination (45, 57, 58), allowing genome-wide association studies (GWAS) to detect asso-

ciations between individual genomic variants and traits of interest (45, 56, 59, 60, 61).
This symbiosis is thus well poised for understanding the genetic basis of important
symbiotic extended phenotypes.

Decades of functional genetic studies have resolved a number of symbiosis genes,
mostly those that disrupt symbiosis when knocked out (reviewed by 62, 63), enabling
increasingly well-resolved models for how interactions establish and how trade is co-
ordinated (64). Nevertheless, mapping efforts in legume-rhizobium symbiosis, which
leverage standing genetic variation, indicate that a much broader set of genes and ge-
netic pathways contribute to the quantitative variation in partner quality that we know
to be important in natural and managed systems (45, 46, 59, 65, 66). Thus quantita-
tive genetic mapping efforts can generate novel candidate genes for symbiosis while
shedding light on the G→P map of symbiotic extended phenotypes in nature.

Here we study the complex genetic basis and layers of context-dependency in
the G→P map of an ecologically-important symbiotic extended phenotype: partner
quality, or the fitness benefit a legume host receives when in association with a par-
ticular rhizobium strain, which not only impacts plant fitness, but also the nutritional
composition of leaves, as high-quality rhizobia fix more atmospheric N that plants can
incorporate into their tissues. Using a panel of 191 strains of the model rhizobium *En-*
sifer meliloti isolated from natural populations in the native range, we 1) ask whether
the genetic architecture of rhizobium partner quality differs across environments and
across host genotypes, or alternatively if variation at a core set of rhizobium genes
control partner quality in all contexts; 2) study how the three genomic elements of *E.*
meliloti contribute to partner quality variation across two host genotypes and in mul-
tiple environments, validating the functional division of labour of these elements in
nature; and finally, 3) suggest candidate metabolic functions and genetic loci that are
responsible for both context-dependency as well as universally-beneficial effects that
are consistent across environments.

RESULTS

Our phenotypic and genetic analyses suggest important roles for context-dependency
in the genetic control of symbiotic extended phenotypes. Overall, we found 1,453
variants (**Supp. Dataset S1**, 67) in 746 (**Supp. Dataset S2**, 67) unique coding genes
associated with shoot biomass, our focal rhizobium partner quality phenotype, while
5,402 variants (**Supp. Dataset S3**, 67) and 1,770 coding genes (**Supp. Dataset S4**, 67)
were associated with at least one partner quality phenotype analyzed in this study (i.e.,
shoot biomass, chlorophyll A, plant height, leaf number); an additional 572 variants fell
into 369 distinct non-coding regions (**Supp. Dataset S3**, 67).

Three categories of G→P associations emerged from our analyses (**Fig. 1**; see
Supp. Fig. S1 in 68 for other traits). First were conditional associations that depended
on the environment (G × E: comparing experiments within host lines, **Fig. 1A**); 510
(68%) were found in only one of the four experiments. Second were "G × G genes", the
51 (7%) genes that were associated with partner quality in both experiments with one
host genotype but were found in neither experiment with the other host genotype (**Fig.**
1A). Specifically, 18 A17-specific genes were found in both experiments with this host
(but never with DZA), while 33 DZA-specific genes were found in both experiments with
this host (but never with A17). Last, but not least, were "universal" genes, the 60 (8%)
genes that were found to be associated with rhizobium partner quality independent
of host genotype or environment (union in the center of **Fig. 1A**). Specifically, while
only five of these genes were associated with partner quality in all four experiments,
55 additional genes were found in three of the four experiments and thus are strong

Batstone et al.

FIG 1 **Association mapping of partner quality reveals strong signatures of environmental-dependence. Venn diagram showing number of rhizobium (*E. meliloti*) genes significantly associated with plant (*M. truncatula*) aboveground biomass (one metric of symbiotic partner quality) in A) each of four separate mapping experiments with either host line DZA (I and III in green) or host line A17 (II and IV in pink). Green and pink unions represent genes contributing to plant biomass in both experiments with either DZA or A17, respectively, while the mauve oval in the center represents universal genes found to contribute to plant aboveground biomass in at least three of the four experiments. B) Genes significantly associated with cross-experiment plasticity in plant aboveground biomass for host genotype DZA in green (experiments I vs. III), A17 in pink (experiments II vs. IV), or independently associated with plasticity in both hosts (central mauve triangle).**

candidates for contributing to partner quality in many conditions. We discuss these three types of genetic effects in turn, at the phenotypic, genomic (three elements in the tripartite genome), gene, and variant level, then conclude with the general implications of our findings for plant-microbe interactions and symbiosis evolution more broadly.

Genotype-by-environment interactions in the G→P map of symbiotic partner quality. Environmental dependence was a pervasive pattern at both the phenotypic, genomic, and individual gene levels, suggesting that different sets of rhizobium genes contribute to symbiotic partner quality under different environmental conditions. At the phenotypic level, ANOVA indicated abundant genetic variance in partner quality traits, including both significant main effects of strain as well as strong strain-by-experiment (G × E) interactions (see **Table 1** and within-experiment heritabilities in **Table 2**). Cross-environment genetic correlations (*r*) of strain means between experiments within a host line were generally significant (**Table 2**), consistent with strain main effects (some strains had consistently higher partner quality than others; see **Supp. Figs. S2 & S3** in 68 for all within and between experiment trait correlations).

TABLE 1   Linear mixed model genotype by environment (G x E) ANOVAs for square-root transformed traits measured on plant lines A17 (left) and DZA (right).
Numbers outside of and within parantheses in columns "A17" and "DZA" represent χ^2 values and degrees of freedom. Rack included as a random effect, while all other terms are fixed. Significance of rack determined by calculating the log likelihood ratio between models with and without the random effect of rack.

Trait	Model term	A17	DZA
Chlorophyll	Intercept	619.4(1)***	1447.65(1)***
	Strain	252.6(190)**	361.67(190)***
	Experiment	0.15(1)	0(1)
	Strain X Experiment	191.38(182)	205.16(183)
	Rack	30.86(1)***	23.91(1)***
Plant height	Intercept	373.58(1)***	355.55(1)***
	Strain	340.68(190)***	268.05(190)***
	Experiment	20.24(1)***	2.5(1)
	Strain X Experiment	308.63(185)***	177.71(183)
	Rack	38.35(1)***	54.12(1)***
Leaves	Intercept	207.69(1)***	164.09(1)***
	Strain	355.37(190)***	207.58(190)
	Experiment	7.85(1)**	0.45(1)
	Strain X Experiment	243.88(185)**	218.54(183)*
	Rack	27.51(1)***	41.67(1)***
Shoot biomass	Intercept	97.65(1)***	179.32(1)***
	Strain	544.81(190)***	499.67(190)***
	Experiment	10.08(1)**	6.93(1)**
	Strain X Experiment	330.06(185)***	262.2(183)***
	Rack	11.32(1)***	155.67(1)***

Significance: $p < 0.001$ = '***'; $p < 0.01$ = '**'; $p < 0.05$ = '*'

TABLE 2 Within-experiment broad-sense heritabilities (H^2) and cross-environment genetic correlations (r). G × E interactions were partitioned into changes in variance versus rank order (i.e., crossing), and the percent due to crossing is presented. Experiments (exps.) 1 and 3 for DZA, 2 and 4 for A17.

Trait	Plant line	H^2 (exps. 1/2)	H^2 (exps. 3/4)	r	Crossing (%)
Shoot biomass	A17	0.302***	0.262***	0.221**	99.83
Chlorophyll		0	0.093*	0.201**	0
Plant height		0.181***	0.173***	0.1	73.37
Leaves		0.173***	0.127***	0.173*	94.13
Shoot biomass	DZA	0.248***	0.253***	0.265***	87.19
Chlorophyll		0.118*	0.141***	0.297***	99.06
Plant height		0.131**	0.065*	0.175*	99.97
Leaves		0.05	0.084*	-0.071	63.97

Significance: $p < 0.001$ = '***'; $p < 0.01$ = '**'; $p < 0.05$ = '*'

Nevertheless, for most traits in both hosts, G × E was driven by changes in the rank order of strain means across environments (**Table 2** crossing (%); see **Supp. Fig. S4** in 68 for reaction norms). Rhizobium strains varied considerably in cross-environment plasticity for shoot biomass, with some strains having consistently low/high partner quality (i.e., resulting in small/large plant biomass, respectively) independent of the environment, while others responded strongly to the environmental differences between experiments (**Supp. Fig. S5**, 68).

At the genomic-level, our evidence indicates that G × E was driven by small-scale shifts in the identity, and estimated allelic effects, of individual loci across environments rather than large shifts in which genomic elements have contributed to partner quality variation (**Supp. Fig. S6**, 68). The vast majority of genes contributing to partner quality variation were located on the symbiotic elements (megaplasmid pSymA or chromid pSymB; **Fig. 2**; **Supp. Dataset S2**, 67; see **Supp. Fig. S7** in 68 for other traits), although some environmentally-dependent loci were found on the chromosome (i.e., G × E and plasticity). Within host lines, 330 total genes were mapped with host line DZA (186 and 144 in experiments 1 and 3, respectively), while only 180 total genes mapped to host line A17 (104 and 76 in experiments 2 and 4, respectively; **Fig. 1A**).

At the gene-level, for experiment 1 in host line DZA, the INTERPRO terms "Transcription regulator *hth*, *lac*I" and "Lambda repressor-like, DNA-binding domain" were marginally significantly enriched (p = 0.00101 and 0.00482, respectively), as was the "Oxidative phosphorylation" KEGG pathway (p = 0.00959; **Supp. Dataset S5**, 67), whereas for experiment 3 in host line DZA, the "Benzoate degradation" KEGG pathway was significantly enriched (FDR-corrected p = 0.00880), while the GO terms "3,4-dihydroxybenzoate catabolic process" and "transmembrane transport" were marginally significantly enriched (p = 0.00204 and 0.00376, respectively; **Supp. Dataset S5**, 67). For experiment 2 in host line A17, the GO terms "transcription, DNA-templated" were significantly enriched (FDR corrected p = 0.00143), as were the related UNIPROT keywords for "transcription regulation" (FDR corrected p = 0.00310) and "DNA-binding" (p = 0.00310; **Supp. Dataset S5**, 67), whereas for experiment 4 in host line A17, the INTERPRO term "Amidohydrolase 1" was marginally significantly enriched (p = 0.0307; **Supp. Dataset S5**, 67). Overall, many of the underlying molecular processes important for driving variation in partner quality appear to be environmentally-dependent, making it harder to predict which genes and pathways will be important for determining symbiotic outcomes under different conditions.

At the variant-level, for each host (DZA or A17), we used correlations of the es-

FIG 2 Loci associated with partner quality are mostly limited to the symbiosis plasmids. Circos plot showing positions of genes (dots) significantly associated with shoot biomass. Each ring represents a different gene category, outermost to innermost: 1) G x E, 2) G x G, 3) partially universal and universal, 4) plasticity, while 5) depicts a histogram based on the total number of significant genes across 100 kbp-sized windows. The x- and y-axes for rings 1-4 represent genomic position (Mbp) and average absolute effect sizes of variants within each gene, respectively. The colours reflect categories in the Venn Diagrams: for rings 1, 2, and 4, genes associated with DZA-only traits are represented by shades of green, on A17-only with shades of purple, and both hosts in mauve (ring 4). For ring 3, genes associated with both hosts in more than three environments are represented in mauve (i.e., "partially universal"), and universal genes in black. Relevant loci are highlighted in blue, with abbreviations for clusters on the outer circle as specified in Supp. Fig. S7 (67).

timated allelic effects to assess the degree to which the individual effects of rhizo-
bium alleles on partner quality were consistent in direction and/or magnitude, or
whether they depended on the environment. When variants were significantly asso-
ciated with partner quality in both experiments, they tended to have inconsistent ef-
fects on host line DZA, while being more consistent on A17 (dotted coloured lines in
Fig. 3A,B, respectively). In fact, on A17, all of the nearly-universal variants (**Fig. 3B**
black dots) had concordant (i.e., same-sign) effects on shoot biomass between experi-
ments, whereas the opposite was true for DZA (**Fig. 3A** black dots) - all nearly-universal
variants had discordant (i.e., opposite-sign) effects between experiments. For host
line DZA, 12 particularly interesting variants had significant but opposing effects on
plant biomass across the two experiments (**Fig. 3A**). Such associations might point to
interesting environmentally-dependent genes, including the glycan metabolism gene
*kdul*l1 (SMa0214/NP_435359.2), and the ornithine decarboxylase gene *spe*C (Smed_5195/
YP_001313905.1; **Supp. Dataset S1**, 67). Regardless of host line, however, the vast ma-
jority of variants in our studies had conditionally-neutral effects, even those with large
magnitude (**Fig. 3**; see **Supp. Fig. S8** in 68 for other traits), and for both hosts, we re-
jected the global null hypothesis that allelic effects were the same across experiments
(both $p < 0.0001$; **Supp. Fig. S9**, 68).

Next we mapped the among-strain differences in plasticity (**Supp. Fig. S5**, 68)
to identify specific loci contributing to G × E. At the genomic-level, most (63%) plas-
ticity loci were found on pSymB (**Fig. 2**; **Supp. Fig. S6B**, 68). Of 576 genes, 400
(69%) were also associated with shoot biomass variation *within* at least one of the
four experiments when each was mapped independently, indicating abundant over-
lap in the genetic architecture of both within- and among-environment partner qual-
ity variation (**Supp. Dataset S6**, 67). Plasticity loci were marginally significantly en-
riched for the INTERPRO terms "Ti-type conjugative transfer relaxase *tra*A" ($p = 0.0292$)
and "*mob*A/*mob*L protein" ($p = 0.0292$), as well as several INTROPRO terms containing
"tetratricopeptide-repeat" (p range: 0.006 to 0.0310), among others (**Supp. Dataset
S5**, 67). Of 576 total plasticity genes, only 93 (16%) were mapped in both host geno-
types (**Fig. 1B**, **Fig. 2**; **Supp. Dataset S7**, 67). Perhaps most interesting are the 14 of
these plasticity genes that were *not* associated with variation within any of the four
experiments when mapped separately (**Supp. Dataset S6**, 67); these loci are particu-
larly strong candidates for understanding how G x E in rhizobia scales up to alter host
growth, and none to our knowledge have known functional roles in symbiosis.

**The genetic architecture of rhizobium partner quality depends on host geno-
type.** Given our four-experiment design, we used the conservative approach of only
considering G × G genes to be those that were associated with partner quality in *both*
experiments with one host genotype and *neither* of the other. We found 33 genes that
contributed to partner quality but only in DZA, and 18 that contributed but only in A17
(**Fig. 1**; **Supp. Dataset S2**, 67). These genes split evenly across the two symbiosis plas-
mids of the genome (pSymA: N = 24; pSymB: N = 27). Below, we discuss our results at
the gene-level for each host line separately.

In the list of A17-only G x G genes, analyses of UNIPROT keywords indicated that
"Selenium" was marginally significantly overrepresented ($p = 0.00771$), while GO terms
for "transferase activity, transferring glycosyl groups" were marginally overrepresented
($p = 0.0417$; **Supp. Dataset S5**, 67). Notably, these genes include both *exo*U (SMb20948/
NP_437611.1) and *exo*W (SMb21690/NP_437613.1; **Supp. Dataset S2**, 67), two glu-
cosyltransferases critical for adding the 6^{th} and 7^{th} sugars (respectively) on the suc-
cinoglycan molecule, an exopolysaccharide required for infection thread formation
and thus host plant invasion for effective symbiosis (69, 70, 71). Non-functional *exo*

FIG 3 Allelic effects are more consistent between experiments for A17, but not for DZA. Correlations between the estimated allelic effects of individual *E. meliloti* variants on plant shoot biomass (from GWAS) in each of two experiments for either A) host DZA (green) or B) A17 (pink). Only allelic effects that were significant in one (lighter colours) or both (dark points) environments are shown, while black dots represent nearly universal variants, i.e., associated with the same trait in three experiments. Linear relationships and R^2 values are depicted for all variants (solid coloured line) or variants significant in both experiments (dotted coloured line). Counts of significant variants for one or both environments appear in the corners of each quadrant within or outside the parentheses, respectively.

genes are known to result in less-efficient, though sometimes not entirely deficient,
symbiosis ([72]). Together with this prior knowledge, our finding that variants in two
succinoglycan biosynthesis enzymes affected host biomass production in host A17
suggests a strong role for succinoglycan-mediated host invasion in mediating symbi-
otic benefits in natural populations. Future studies of coevolution between succino-
glycan structure and host plant detection would likely be fruitful. Also notable is *sel*A
(SMa0011/NP_435251.1; **Supp. Dataset S2**, [67]), a transferase required for the biosyn-
thesis of selenocysteine, a less-used amino acid incorporated into select proteins in
only about 20% of bacterial genomes ([73], [74], [75]). To our knowledge, nothing is known
about the role of selenocysteine in rhizobia or in legume-rhizobium symbiosis.

By contrast, the list of DZA-only G × G genes was dominated by GO terms "tran-
scription factor activity, sequence-specific DNA binding" (p = 0.0247) and "cellular amino
acid metabolic process" (p = 0.0425; **Supp. Dataset S5**, [67]). Notably this list con-
tains three *lys*R transcriptional regulators (**Supp. Dataset S2**, [67]) of at least 90 in
the genome: SMa2287 (NP_436477.2) on pSymA plus SM_b20494 (NP_437016.1) and
SM_b21434 (NP_437803.1) on pSymB are not well-studied, and had weak (SM_b20494)
or undetectable effects in a plasmid insertion mutagenesis screen compared to the
named *lsr*A and *lsr*B ([76]). This class of loci is well-known to control expression of
genes for symbiosis, both mutualistic and pathogenic ([77]). Beyond those discovered
in knockdown studies, our results suggest additional symbiotic roles for natural vari-
ation at these *lys*R regulators in *E. meliloti*. Interestingly, one metabolic process gene
(*gdh*A: SMa0228/NP_435368.1 on pSymA; **Supp. Dataset S2**, [67]), part of one bacterial
pathway for assimilation of ammonium into glutamate ([78]), was identified in a com-
parative genomics study of five *Ensifer* (formerly *Sinorhizobium*) species as specific to
E. meliloti ([79]). The role of *gdh*A in rhizobial symbiosis is not well-known, though in-
terestingly it was found to increase ammonium assimilation when the *E. coli* copy was
expressed transgenically in tobacco ([80]).

Universal associations highlight transport functions and secretion systems.
Despite these layers of context-dependency, we did find significant main effects of
strain for all phenotypes (**Table 1**). Concomitantly we resolved a number of loci that
were consistently associated with shoot biomass, being mapped in either three ex-
periments (55 nearly-universal genes) or even all four experiments (5 universal genes)
(**Fig. 1**). At the genomic and gene-levels, the set of 60 universal/nearly-universal genes
were split between pSymA and pSymB (38 and 22 loci, respectively; **Fig. 2** mauve
dots; **Supp. Dataset S2**, [67]) and featured marginally-enriched INTERPRO terms for
"Tetratricopeptide-like helical" (p < 0.001) and "Tetratricopeptide repeat-containing do-
main" (p = 0.00160) as well as UNIPROT keywords "Transmembrane" (p = 0.0190) and
"Transmembrane helix" (p = 0.0407), though none of these terms were significant after
FDR-correction (**Supp. Dataset S5**, [67]).

Of the five truly universal genes (mapped in all 4 experiments; **Supp. Dataset
S2**, [67]), most do not have known functions in symbiosis. This includes a *cax* gene
(SMa0675/ NP_435603.2) putatively involved in calcium/proton exchange. Though *cax*
genes are widespread in bacteria and eukaryotes ([81]), and the importance of calcium
both in nodule establishment and trade of benefits is known ([82], [83]), it is difficult to
hypothesize on the function of this particular gene or its genetic variants in symbio-
sis currently. Given their as-yet unknown functions and lack of context-dependency in
our studies, the five universal candidates might hold the most potential for novel func-
tional information and consistent phenotypic effects, which might make them ideal
candidates both for validation and for symbiosis improvement.

Like the five universal genes, the "nearly-universal" set of genes associated with

shoot biomass in three (of four) experiments highlights the existence of segregating
natural variation in a number of interesting metabolic pathways, only some of which
have established roles in symbiosis. We found five loci annotated as involved in trans-
membrane transport (**Supp. Dataset S2**, 67), including *kdp*A (SMa2333/NP_436501.1),
*pot*E (SMa0678/NP_435605.1), and *msb*A1 (SM_b20813/NP_437093.2) – in addition to
the nodulation protein *glm*S/*nod*M (SMa0878/NP_435728.1) and the universally-associated
cax transporter (discussed above). The role of potassium transporters such as *kdp*A
in osmoregulation during symbiosis is not well understood (84). The *pot*E locus codes
for a putrescine/ornithine antiporter, while another nearby nearly-universal locus
(SMa0682/NP_435607.4) is a predicted amino acid decarboxylase in the putrescine
biosynthesis pathway, potentially suggesting a role for variation in putrescine metabolism,
which is known to vary in *E. meliloti* (45), in symbiotic partner quality (45, 85).

We additionally compared our consistent (universal and nearly-universal) candi-
date gene lists with GWAS associations from other research groups in different ge-
nomic backgrounds (e.g., 45, 66), and discovered important roles for natural variation
in rhizobial glucosamine processing, T4SS transferase loci (*tra*A and *tra*G), and a hand-
ful of others in determining symbiotic partner quality in nature. See **Supp. Results**
(68) & **Supp. Dataset S2** (67) for more details.

DISCUSSION

Understanding our symbiotic world requires a genetically-accurate appreciation of
the symbiotic extended phenotypes upon which selection acts. While evolutionary
ecology has long recognized the importance of genetic variation in symbiosis while ig-
noring the underlying mechanisms, functional geneticists have traditionally resolved
mechanisms without taking into account natural variation. Synthesis of these two per-
spectives has started to resolve ecologically-relevant quantitative variation at the nu-
cleotide level (45, 46, 47, 60, 86, 87). Here we quantify multiple symbiotic extended
phenotypes in four mapping experiments using a model plant-microbe symbiosis and
find that the genetic architecture of symbiotic partner quality is complex, underlain
by networks of numerous interacting loci and environmental-dependence. We find
that some loci in the microbial symbiont have strong and consistent effects on host
growth across environments, contributing to the overall differences in mutualistic part-
ner quality that have been the focus of many empirical and theoretical studies of mu-
tualism to date (reviewed by 40, 88, 89, 90, 91). Nonetheless, the vast majority of loci
identified in our study were significantly associated with variation in partner quality in
specific environments (G × E effects), or with specific host genotypes (G × G effects).
We first discuss the roles of environmentally-dependent loci, versus universal loci that
are found consistently. Next we discuss the coevolutionary implications of genotype-
dependence (G x G interactions), then wrap up with a call for further synthesis with
metabolic network models towards systems genetics of symbiosis.

Environmental context-dependency in the G→P map of symbiosis. Ecological
effects of context-dependency in mutualism have been recognized for a long time, i.e.,
traits and mutualism benefits often shift across environmental conditions such as nu-
trient availability or light environments (92, 93, 94), although not ubiquitously for all
traits (e.g., 95, 96). More recent studies have begun to document the evolutionary
changes that can result from these ecological effects, e.g., divergence of host and/or
symbiont symbiosis traits across strong ecological gradients (59, 97, 98, 99, 100). Evolu-
tionary change in response to environments implies that the loci underlying selected
traits have differential effects on fitness across environments. Here we identify the
loci that generate important trait variation both within and among environments, the

trait variation upon which selection acts in nature.

The majority of context-dependent partner quality genes we identified might be
viewed much like the conditionally-neutral variation so often found in studies of lo-
cal adaptation ([101], [102], [103], [104], [105]), contributing strongly to variation in partner
quality in some contexts but having a range of weaker effects (both in the same or op-
posing direction) in another context. For example, by experimentally-evolving strains
of *E. meliloti* on five *M. truncatula* lines, Batstone et al. ([66]) found that local adaptation
was largely governed by conditional neutrality (beneficial on local host, neutral on non-
local hosts) or mildly deleterious effects on non-local hosts, likely due to drift in the
local context. Yet, in nature where rhizobium population sizes are much larger and
more diverse, and gene flow is present, the extent to which local adaptation occurs
has rarely been tested but might be unlikely ([28], [106], [107]), with the exception of pop-
ulations differentiated by strong ecological gradients (e.g., [97]). Moreover, the host
genotypes used in our study, DZA and A17, are unlikely to share an evolutionary his-
tory with our strains, and so it remains unclear whether stronger trade-offs (and less
conditional neutrality) would be present if our strains shared an evolutionary history
with the host lines being tested.

Despite widespread conditional neutrality, a handful of interesting variants had
strong effects on host growth, but in *opposing directions* across experiments within a
single host genetic background. These sorts of antagonistic effects can favour differ-
ent variants in different environments, and thus, potentially help explain the mainte-
nance of mutualism variation in nature ([90], [108], [109], [110]). Nevertheless we note that
these sorts of G × E variants were rare in our study, despite strong rank-order effects
among strain means at the organismal level (**Table 2**); moreover, because our "envi-
ronments" were simply different greenhouse experiments, relating such antagonistic
effects to adaptation in the wild will require the type of *in situ* studies that are common
in plants (e.g., [99], [102], [111]) but more difficult in soil microbes. We nevertheless found
several genes and pathways that are ideal for functional follow-up studies aimed at
identifying consistent associations, or loci which might benefit from "fine-tuning" sym-
biotic benefits towards improving plant health. The strengths of GWAS (capturing ge-
netic variation, ecological relevance) also lead to weaknesses (imprecision due to con-
founding population structure, false negatives and false positives). Thus our strongest
recommendations for functional hypotheses for validation in follow up studies are the
universal genes identified here that overlap with candidate genes identified in sepa-
rate GWAS using different genomic backgrounds and environments (e.g., [45], [66]).

Overall we find that the loci underlying quantitative variation in symbiotic extended
phenotypes often (but not always) depend on the environment, and therefore that a
nuanced understanding of how complex traits interact with environmental variables
will be necessary for many of the lofty goals in plant microbiome and symbiosis re-
search ([9], [35]). This point has been made before ([112]), as the presence of G × E as
studied at the phenotypic level has been recognized for decades; what is novel here is
our ability to interrogate this variation at the genomic level and for multiple host lines
(see below).

Coevolutionary implications of genotype-dependence. Uncovering the com-
plex genetics of how two (or more) genomes interact with each other to generate
trait variation is an important step to better understanding how these traits (co)evolve
([6]) and how to better manipulate traits in the future to address societal challenges
([37], [113]). For example, identifying the loci underlying mutualistic traits allows us to
address longstanding debates within mutualism theory ([114]), including how readily
conflict evolves ([115]), and how genetic variation is maintained despite host selection

for the 'best' symbiont (90).

Previous mapping efforts in the legume-rhizobium system have focused on the HapMap collections, which maximize diversity using a range-wide sample. Our study focuses on the segregating natural variation at a smaller geographic scale, a scale at which pSymA and pSymB segregate (107); thus the G $\times$ G-driven variation we find here would be available to local evolutionary and/or coevolutionary processes. G $\times$ G interactions for fitness outcomes have long been of interest in the legume-rhizobium symbiosis (25, 116, 28, 97, 117) and other interactions (118, 119, 120) because such statistical interactions generate the fitness variation that drives coevolution, but also have implications for breeding and agricultural production whereby the functional effects of symbiont variation depend on the crop genotype (121, 122). Recent molecular genetic and transcriptomic approaches on a handful of genotypes have begun to resolve the mechanistic underpinnings of G $\times$ G (123, 124, 125, 126, 127, 128), while biparental or GWAS mapping approaches (45, 46, 129) provide broader insight into the genetic architecture underlying G $\times$ G (i.e., the number, average effect size, and consistency of loci). The picture emerging from ours and others' studies is that G $\times$ G, like symbiotic partner quality itself, has a complex, polygenic basis and will require both statistically sophisticated and metabolically-informed models to unravel.

Symbiotic extended phenotypes as quantitative traits. In the age of rapid microbiome sequencing and expanding efforts to characterize the loci in plant genomes that contribute to microbiome variation among cultivars or genotypes (e.g., rice, *Lotus*, *Medicago*), our results present an important juxtaposition, as the abundant and context-dependent genetic variation characterized in detail here occurs within a set of 191 strains with > 98% average nucleotide identity (well above the typical threshold for delineating and enumerating operational taxonomic units, or OTUs, in metagenomic studies). At the same time, functional genetic studies have made much progress identifying loci critical to symbiosis establishment and downstream processes by creating knock-out or knock-down mutants and comparing their associated symbiotic phenotypes to a wildtype strain (e.g., *dnf* mutants in legumes, Ané et al. 130; fix+/fix- mutants of rhizobia, Wheatley et al. 131), but these loci are often viewed as on-or-off switches for symbiosis more generally, or cooperation more specifically in models of mutualism theory (132, 133, 134, 135). Our study demonstrates that the vast majority of loci within the symbiont genome act more like dials than on-and-off switches, generating the quantitative variation in symbiotic extended phenotypes observed in nature.

While our approach allows us to zero-in on particular loci underlying G $\times$ E and G $\times$ G interactions, *in silico* modelling of both plant (136) and rhizobium (137) metabolism that links together genetic information with metabolic pathways could then be used to simulate key symbiotic processes under a wide range of conditions (138). Ultimately combining transcriptomic, genomic, and metabolomic datasets will be required for a synthetic understanding of how nucleotide variation percolates up through shared symbiotic metabolic networks (17, 139, 140). Efforts to reintegrate research on symbiosis genetics and (co)evolution with plant-microbiome work (reviewed in 37, 113) will be fruitful in revealing additional intraspecific variation driving patterns of genotype-dependence and coevolution and resolving mechanisms of host control of the microbiome.

MATERIALS AND METHODS

Full details are available in the Supplemental Materials (**Supp. Methods**, 68). We performed four greenhouse experiments to estimate partner quality phenotypes in *E. meliloti*. In each experiment, plants from one of two host lines (either A17 or DZA)

were grown in single inoculation with each of 191 *E. meliloti* strains, with three to four replicates per strain per experiment (six to eight total replicates for each plant line x strain combination, N = 2,825 plants total). Experiments were planted in 2018, in March (I: DZA), May (II: A17), Sept (III: DZA), and Nov (IV: A17). Uninoculated controls (40 per experiment) were included to gauge contamination, which was minimal overall, and limited to the first two experiments (**Supp. Fig. S10**, 68). We measured multiple proxies of partner quality, namely leaf chlorophyll A content, plant height, number of leaves, and above-ground dried shoot biomass, although we focus on the latter in the main text (see Supplemental Materials for all others). We conducted multiple pheno-typic analyses to determine how much variation in partner quality was due to strain, experiment, and the interaction of both, among other questions.

We sequenced the entire genomes of all 191 *E. meliloti* strains, called single nu-cleotide polymorphisms (SNPs, henceforth referred to as variants), and performed genome-wide association tests that accounted for rhizobium population structure and included only unlinked variants. We determined which loci were significantly associ-ated with partner quality using a permutation method, and binned these loci into three categories based on the context-dependency of their phenotypic effects, and thus, their contribution to the layers of the G→P map for each of our symbiotic extended phenotypes. First, "nearly-universal genes" were those found to have significant ef-fects in at least three of the four experiments for a particular trait ("universal genes" were mapped in all four experiments). Second, we used a conservative approach to call "G × G genes" as those mapped in both experiments for one host genotype but neither of the experiments for the other host genotype (i.e., "DZA G × G" genes were significant in both experiments I and III with DZA but neither II nor IV with host A17). Third were genes significantly associated with partner quality in a single experiment, and never in another (i.e., "G × E" genes). Finally, we conducted candidate gene func-tional analyses to understand how loci within different categories differed from one another functionally, or whether they were part of the same networks/metabolic path-ways.

Data Availability. Strains and plant lines are available upon request. All raw data and analysis code are available on GitHub (see "Complex_genetics" folder). Once raw sequence reads and assemblies are archived and made available on NCBI, accession numbers will be added to this manuscript.

SUPPLEMENTAL MATERIAL

Detailed methods, additional results, and twelve (12) supplementary figures are avail-able on Zenodo (68) (doi: https://doi.org/10.5281/zenodo.5550958). Seven (7) supple-mentary datasets are available on Dryad (68) (doi: https://doi.org/10.5061/dryad.5dv41ns6r). All supplemental material, in addition to the main text, is also available at bioRχiv (doi: https://doi.org/10.1101/2021.08.03.454976).

ACKNOWLEDGMENTS

We thank greenhouse staff (esp. Debbie Black). We acknowledge funding from the National Science Foundation (IOS-1645875 and NPGI-1401864), a Carl R. Woese Insti-tute for Genomic Biology postdoctoral fellowship to R.T. Batstone, strain sequencing by Joint Genome Institute (CSP-1223795), as well as analytical advice from B. Epstein.

REFERENCES

1. **Klironomos JN**. 2003. Variation in plant response to native and exotic arbuscular mycorrhizal fungi. Ecology 84 (9):2292–2301.

2. **Oliver KM, Moran NA, Hunter MS**. 2005. Variation in resistance to parasitism in aphids is due to symbionts not host genotype. Proc Natl Acad Sci 102 (36):12795–12800.

3. **Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, Griffin NW, Lombard V, Henrissat B, Bain JR, et al.**. 2013. Gut microbiota from twins discordant for obesity modulate metabolism in mice. Science 341 (6150).

4. **Brock DA, Jones K, Queller DC, Strassmann JE**. 2016. Which phenotypic traits of Dictyostelium discoideum farmers are conferred by their bacterial symbionts? Symbiosis 68 (1-3):39–48.

5. **Gilbert JA, Lynch SV**. 2019. Community ecology as a framework for human microbiome research. Nat Med 25 (6):884–889.

6. **O'Brien AM, Jack CN, Friesen ML, Frederickson ME**. 2021. Whose trait is it anyways? Coevolution of joint phenotypes and genetic architecture in mutualisms. Proc Royal Soc B 288 (1942):20202483.

7. **Afkhami ME, Friesen ML, Stinchcombe JR**. 2021. Multiple Mutualism Effects generate synergistic selection and strengthen fitness alignment in the interaction between legumes, rhizobia and mycorrhizal fungi. Ecol Lett .

8. **Brundrett MC**. 2002. Coevolution of roots and mycorrhizas of land plants. New Phytol 154 (2):275–304.

9. **Friesen ML, Porter SS, Stark SC, von Wettberg EJ, Sachs JL, Martinez-Romero E**. 2011. Microbially mediated plant functional traits. Annu Rev Ecol Evol Syst 42:23–46.

10. **Hurst GD**. 2017. Extended genomes: symbiosis and evolution. Interface Focus 7 (5):20170001.

11. **Hawkes CV, Bull JJ, Lau JA**. 2020. Symbiosis and stress: how plant microbiomes affect host evolution. Philos Trans Royal Soc B 375 (1808):20190590.

12. **Moeller AH, Sanders JG**. 2020. Roles of the gut microbiota in the adaptive evolution of mammalian species. Philos Trans Royal Soc B 375 (1808):20190597.

13. **Hansen TF**. 2006. The evolution of genetic architecture. Annu Rev. Ecol. Evol. Syst. 37:123–157.

14. **Orgogozo V, Morizot B, Martin A**. 2015. The differential view of genotype–phenotype relationships. Front Genet 6:179.

15. **Mackay TF, Huang W**. 2018. Charting the genotype–phenotype map: lessons from the Drosophila melanogaster Genetic Reference Panel. Wiley Interdiscip Rev Dev Biol 7 (1):e289.

16. **Checcucci A, diCenzo GC, Ghini V, Bazzicalupo M, Becker A, Decorosi F, Döhlemann J, Fagorzi C, Finan TM, Fondi M, et al.**. 2018. Creation and characterization of a genomically hybrid strain in the nitrogen-fixing symbiotic bacterium *Sinorhizobium meliloti*. ACS Synth Biol 7 (10):2365–2378.

17. **Marshall-Colón A, Kliebenstein DJ**. 2019. Plant networks as traits and hypotheses: moving beyond description. Trends Plant Sci 24 (9):840–852.

18. **Cheverud JM**. 1982. Phenotypic, genetic, and environmental morphological integration in the cranium. Evolution p 499–516.

19. **Cheverud JM**. 1996. Developmental integration and the evolution of pleiotropy. Am Zoöl 36 (1):44–50.

20. **Hansen TF**. 2013. Why epistasis is important for selection and adaptation. Evolution 67 (12):3501–3511.

21. **Lee SH, Ripke S, Neale BM, Faraone SV, Purcell SM, Perlis RH, Mowry BJ, Thapar A, Goddard ME, Witte JS, et al.**. 2013. Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. Nat Genet 45 (9):984–995.

22. **Campbell-Staton SC, Cheviron ZA, Rochette N, Catchen J, Losos JB, Edwards SV**. 2017. Winter storms drive rapid phenotypic, regulatory, and genomic shifts in the green anole lizard. Science 357 (6350):495–498.

23. **Saltz JB, Hessel FC, Kelly MW**. 2017. Trait correlations in the genomics era. Trends Ecol & Evol 32 (4):279–290.

24. **Frank S**. 1992. Models of plant-pathogen coevolution. Trends Genet 8 (6):213–219.

25. **Parker MP**. 1995. Plant fitness variation caused by different mutualist genotypes. Ecology 76 (5):1525–1535.

26. **Lambrechts L, Fellous S, Koella JC**. 2006. Coevolutionary interactions between host and parasite genotypes. Trends Parasitol 22 (1):12–16.

27. **Wade MJ**. 2007. The co-evolutionary genetics of ecological communities. Nat Rev Genet 8 (3):185–195.

28. **Heath KD**. 2010. Intergenomic epistasis and coevolutionary constraint in plants and rhizobia. Evolution 64 (5):1446–1458.

29. **Parker BJ, Hrček J, McLean AH, Godfray HCJ**. 2017. Genotype specificity among hosts, pathogens, and beneficial microbes influences the strength of symbiont-mediated protection. Evolution 71 (5):1222–1231.

30. **Piculell BJ, Hoeksema JD, Thompson JN**. 2008. Interactions of biotic and abiotic environmental factors in an ectomycorrhizal symbiosis, and the potential for selection mosaics. BMC Biol 6 (1):1–11.

31. **Heath KD, Stock A, Stinchcombe J**. 2010. Mutualism variation in the nodulation response to nitrate. J Evol Biol 23 (11):2494–2500.

32. **Wendling CC, Fabritzek AG, Wegner KM**. 2017. Population-specific genotype x genotype x environment interactions in bacterial disease of early life stages of pacific oyster larvae. Evol Appl 10 (4):338–347.

33. **Thompson JN**. 2005. The geographic mosaic of coevolution. University of Chicago Press.

34. **Thompson JN**. 2009. The coevolutionary process. University of Chicago press.

35. **Busby PE, Soman C, Wagner MR, Friesen ML, Kremer J, Bennett A, Morsy M, Eisen JA, Leach JE, Dangl JL**. 2017. Research priorities for harnessing plant microbiomes in sustainable agriculture. PLoS Biol 15 (3):e2001793.

36. **Porter SS, Sachs JL**. 2020. Agriculture and the disruption of plant–microbial symbiosis. Trends Ecol & Evol 35 (5):426–439.

37. **Batstone RT**. 2021. Genomes within Genomes: nested symbiosis and its implications for plant evolution. New Phytol (in review).

38. **Cook DR**. 1999. *Medicago truncatula*–a model in the making! Curr Opin Plant Biol 2 (4):301–304.

39. **Heath KD, Grillo MA**. 2016. Rhizobia: tractable models for bacterial evolutionary ecology: Models for bacterial evolutionary ecology. Environ Microbiol 18 (12).

40. **Sachs JL, Quides KW, Wendlandt CE**. 2018. Legumes versus rhizobia: a model for ongoing conflict in symbiosis. New Phytol 219 (4):1199–1206.

41. **diCenzo GC, Zamani M, Checcucci A, Fondi M, Griffitts JS, Finan TM, Mengoni A**. 2019. Multidisciplinary approaches for studying rhizobium–legume symbioses. Can J Microbiol 65 (1):1–33.

42. **Burghardt LT**. 2020. Evolving together, evolving apart: measuring the fitness of rhizobial bacteria in and out of symbiosis with leguminous plants. New Phytol 228 (1):28–34.

43. **Young ND, Cannon SB, Sato S, Kim D, Cook DR, Town CD, Roe BA, Tabata S**. 2005. Sequencing the genespaces of *Medicago truncatula* and *Lotus japonicus*. Plant Physiol 137 (4):1174–1181.

44. **Epstein B, Branca A, Mudge J, Bharti AK, Briskine R, Farmer AD, Sugawara M, Young ND, Sadowsky MJ, Tiffin P**. 08 2012. Population Genomics of the Facultatively Mutualistic Bacteria *Sinorhizobium meliloti* and *S. medicae*. PLoS Genet 8:1–10. doi: 10.1371/journal.pgen.1002868.

Batstone et al.

45. **Epstein B, Abou-Shanab RA, Shamseldin A, Taylor MR, Guhlin J, Burghardt LT, Nelson M, Sadowsky MJ, Tiffin P**. 2018. Genome-wide association analyses in the model Rhizobium *Ensifer meliloti*. MSphere 3 (5).

46. **Stanton-Geddes J, Paape T, Epstein B, Briskine R, Yoder J, Mudge J, Bharti AK, Farmer AD, Zhou P, Denny R, et al.**. 2013. Candidate genes and genetic architecture of symbiotic and agronomic traits revealed by whole-genome, sequence-based association genetics in *Medicago truncatula*. PloS One 8 (5):e65688.

47. **Burghardt LT, Epstein B, Guhlin J, Nelson MS, Taylor MR, Young ND, Sadowsky MJ, Tiffin P**. 2018. Select and resequence reveals relative fitness of bacteria in symbiotic and free-living environments. Proc Natl Acad Sci 115 (10):2425–2430.

48. **Graham PH, Vance CP**. 2003. Legumes: importance and constraints to greater use. Plant Physiol 131 (3):872–877.

49. **Foyer CH, Lam HM, Nguyen HT, Siddique KH, Varshney RK, Colmer TD, Cowling W, Bramley H, Mori TA, Hodgson JM, et al.**. 2016. Neglecting legumes has compromised human health and sustainable food production. Nat Plants 2 (8):1–10.

50. **Fustec J, Lesuffleur F, Mahieu S, Cliquet JB**. 2010. Nitrogen rhizodeposition of legumes. A review. Agron for Sustain Dev 30 (1):57–66.

51. **Duchene O, Vian JF, Celette F**. 2017. Intercropping with legume for agroecological cropping systems: Complementarity and facilitation processes and the importance of soil microorganisms. A review. Agric Ecosyst & Environ 240:148–161.

52. **Harrison PW, Lower RP, Kim NK, Young JPW**. 2010. Introducing the bacterial 'chromid': not a chromosome, not a plasmid. Trends Microbiol 18 (4):141–148.

53. **Young ND, Zhou P, Silverstein KA**. 2016. Exploring structural variants in environmentally sensitive gene families. Curr Opin Plant Biol 30:19–24.

54. **Dicenzo GC, Finan TM**. 2017. The divided bacterial genome: structure, function, and evolution. Microbiol Mol Biol Rev 81 (3):e00019–17.

55. **Misra HS, Maurya GK, Kota S, Charaka VK**. 2018. Maintenance of multipartite genome system and its functional significance in bacteria. J Genet 97 (4):1013–1038.

56. **Cavassim MIA, Moeskjær S, Moslemi C, Fields B, Bachmann A, Vilhjálmsson BJ, Schierup MH, Young JPW, Andersen SU**. 2020. Symbiosis genes show a unique pattern of introgression and selection within a Rhizobium leguminosarum species complex. Microb Genom 6 (4).

57. **Nelson M, Guhlin J, Epstein B, Tiffin P, Sadowsky MJ**. 2018. The complete replicons of 16 Ensifer meliloti strains offer insights into intra-and inter-replicon gene transfer, transposon-associated loci, and repeat elements. Microb Genom 4 (5).

58. **Epstein B, Tiffin P**. 2021. Comparative genomics reveals high rates of horizontal transfer and strong purifying selection on rhizobial symbiosis genes. Proc Royal Soc B 288 (1942):20201804.

59. **Klinger CR, Lau JA, Heath KD**. 2016. Ecological genomics of mutualism decline in nitrogen-fixing bacteria. Proc Royal Soc B 283 (1826):20152563.

60. **Porter SS, Chang PL, Conow CA, Dunham JP, Friesen ML**. 2017. Association mapping reveals novel serpentine adaptation gene clusters in a population of symbiotic Mesorhizobium. The ISME J 11 (1):248–262.

61. **Power RA, Parkhill J, de Oliveira T**. 2017. Microbial genome-wide association studies: lessons from human GWAS. Nat Rev Genet 18 (1):41–50.

62. **Wang D, Yang S, Tang F, Zhu H**. 2012. Symbiosis specificity in the legume–rhizobial mutualism. Cell Microbiol 14 (3):334–342.

63. **Roy S, Liu W, Nandety RS, Crook A, Mysore KS, Pislariu CI, Frugoli J, Dickstein R, Udvardi MK**. 2020. Celebrating 20 years of genetic discoveries in legume nodulation and symbiotic nitrogen fixation. The Plant Cell 32 (1):15–41.

64. **diCenzo GC, Tesi M, Pfau T, Mengoni A, Fondi M**. 2020. Genome-scale metabolic reconstruction of the symbiosis between a leguminous plant and a nitrogen-fixing bacterium. Nat Commun 11 (1):1–11.

65. **Porter SS, Faber-Hammond J, Montoya AP, Friesen ML, Sackos C**. 2019. Dynamic genomic architecture of mutualistic cooperation in a wild population of Mesorhizobium. The ISME J 13 (2):301–315.

66. **Batstone RT, O'Brien AM, Harrison TL, Frederickson ME**. 2020. Experimental evolution makes microbes more cooperative with their local host genotype. Science 370 (6515):476–478.

67. **Batstone R, Lindgren H, Allsup CM, Goralka LA, Riley AB, Grillo MA, Marshall-Colon A, Heath KD**. 2021. Supplemental datasets for "The complex genetics of symbiotic extended phenotypes across multiple environments in a model mutualism". https://doi.org/10.5061/dryad.5dv41ns6r.

68. **Batstone R, Lindgren H, Allsup CM, Goralka LA, Riley AB, Grillo MA, Marshall-Colon A, Heath KD**. 2021. Supplemental material for "The complex genetics of symbiotic extended phenotypes across multiple environments in a model mutualism". https://doi.org/10.5281/zenodo.5550958.

69. **Becker A, Kleickmann A, Küster H, Keller M, Arnold W, Pühler A**. 1993. Analysis of the *Rhizobium meliloti* genes exoU, exoV, exoW, exoT, and exoI involved in exopolysaccharide biosynthesis and nodule invasion: exoU and exoW probably encode glucosyltransferases. Mol Plant-Microbe Interactions: MPMI 6 (6):735–744.

70. **Reuber TL, Walker GC**. 1993. Biosynthesis of succinoglycan, a symbiotically important exopolysaccharide of *Rhizobium meliloti*. Cell 74 (2):269–280.

71. **Nogales J, Muñoz S, Olivares J, Sanjuán J**. 2006. *Sinorhizobium meliloti* genes involved in tolerance to the antimicrobial peptide protamine. FEMS Microbiol Lett 264 (2):160–167.

72. **Mendis HC, Madzima TF, Queiroux C, Jones KM**. 2016. Function of succinoglycan polysaccharide in *Sinorhizobium meliloti* host plant invasion depends on succinylation, not molecular weight. MBio 7 (3):e00606–16.

73. **Copeland PR**. 2005. Making sense of nonsense: the evolution of selenocysteine usage in proteins. Genome Biol 6 (6):1–5.

74. **Peng T, Lin J, Xu YZ, Zhang Y**. 2016. Comparative genomics reveals new evolutionary and ecological patterns of selenium utilization in bacteria. The ISME J 10 (8):2048–2059.

75. **Wells M, Stolz JF**. 2020. Microbial selenium metabolism: a brief history, biogeochemistry and ecophysiology. FEMS Microbiol Ecol 96 (12):fiaa209.

76. **Luo L, Yao SY, Becker A, Rüberg S, Yu GQ, Zhu JB, Cheng HP**. 2005. Two new *Sinorhizobium meliloti* LysR-type transcriptional regulators required for nodulation. J Bacteriol 187 (13):4562–4572.

77. **Schell MA**. 1993. Molecular biology of the *LysR* family of transcriptional regulators. Annu Rev Microbiol 47 (1):597–626.

78. **Merrick M, Edwards R**. 1995. Nitrogen control in bacteria. Microbiol Rev 59 (4):604–622.

79. **Sugawara M, Epstein B, Badgley BD, Unno T, Xu L, Reese J, Gyaneshwar P, Denny R, Mudge J, Bharti AK, et al.**. 2013. Comparative genomics of the core and accessory genomes of 48 *Sinorhizobium* strains comprising five genospecies. Genome Biol 14 (2):1–20.

80. **Ameziane R, Bernhard K, Lightfoot D**. 2000. Expression of the bacterial *gdh*A gene encoding a NADPH glutamate dehydrogenase in tobacco affects plant growth and development. Plant Soil 221 (1):47–

57.

81. **Shigaki T, Rees I, Nakhleh L, Hirschi K**. 2006. Identification of three distinct phylogenetic groups of CAX cation/proton antiporters. J Mol Evol 63 (6):815–825.

82. **Oldroyd GE, Downie JA**. 2004. Calcium, kinases and nodulation signalling in legumes. Nat Rev Mol Cell Biol 5 (7):566–576.

83. **Clarke VC, Loughlin PC, Day DA, Smith P**. 2014. Transport processes of the legume symbiosome membrane. Front Plant Sci 5:699.

84. **Domínguez-Ferreras A, Muñoz S, Olivares J, Soto MJ, Sanjuán J**. 2009. Role of potassium uptake systems in *Sinorhizobium meliloti* osmoadaptation and symbiotic performance. J Bacteriol 191 (7):2133–2143.

85. **Becerra-Rivera VA, Dunn MF**. 2019. Polyamine biosynthesis and biological roles in rhizobia. FEMS Microbiol Lett 366 (7):fnz084.

86. **Branca A, Paape TD, Zhou P, Briskine R, Farmer AD, Mudge J, Bharti AK, Woodward JE, May GD, Gentzbittel L, et al.**. 2011. Whole-genome nucleotide diversity, recombination, and linkage disequilibrium in the model legume *Medicago truncatula*. Proc Natl Acad Sci 108 (42):E864–E870.

87. **Friesen ML, von Wettberg EJ, Badri M, Moriuchi KS, Barhoumi F, Chang PL, Cuellar-Ortiz S, Cordeiro MA, Vu WT, Arraouadi S, et al.**. 2014. The ecological genomic basis of salinity adaptation in Tunisian *Medicago truncatula*. BMC Genom 15 (1):1160.

88. **Simms EL, Taylor DL**. 2002. Partner choice in nitrogen-fixation mutualisms of legumes and rhizobia. Integr Comp Biol 42 (2):369–380.

89. **Denison RF**. 2000. Legume sanctions and the evolution of symbiotic cooperation by rhizobia. The Am Nat 156 (6):567–576.

90. **Heath KD, Stinchcombe JR**. 2014. Explaining mutualism variation: a new evolutionary paradox? Evolution 68 (2):309–317.

91. **Stoy KS, Gibson AK, Gerardo NM, Morran LT**. 2020. A need to consider the evolutionary genetics of host–symbiont mutualisms. J Evol Biol 33 (12):1656–1668.

92. **Chamberlain SA, Bronstein JL, Rudgers JA**. 2014. How context dependent are species interactions? Ecol Lett 17 (7):881–890.

93. **Bronstein JL**. 2015. Mutualism. Oxford University Press, USA.

94. **Hoeksema JD, Bruna EM**. 2015. Context-Dependent Outcomes of Mutualistic Interactions, p 181–202. *In* Bronstein JL (ed), Mutualism. Oxford University Press, Oxford, UK.

95. **Regus JU, Gano KA, Hollowell AC, Sachs JL**. 2014. Efficiency of partner choice and sanctions in Lotus is not altered by nitrogen fertilization. Proc Royal Soc B: Biol Sci 281 (1781):20132587.

96. **Grillo MA, Stinchcombe JR, Heath KD**. 2016. Nitrogen addition does not influence pre-infection partner choice in the legume–rhizobium symbiosis. Am J Bot 103 (10):1763–1770.

97. **Porter SS, Stanton ML, Rice KJ**. 2011. Mutualism and adaptive divergence: co-invasion of a heterogeneous grassland by an exotic legume-rhizobium symbiosis. PloS One 6 (12):e27935.

98. **Weese DJ, Heath KD, Dentinger BT, Lau JA**. 2015. Long-term nitrogen addition causes the evolution of less-cooperative mutualists. Evolution 69 (3):631–642.

99. **Rúa MA, Antoninka A, Antunes PM, Chaudhary VB, Gehring C, Lamit LJ, Piculell BJ, Bever JD, Zabinski C, Meadow JF, et al.**. 2016. Home-field advantage? Evidence of local adaptation among plants, soil, and arbuscular mycorrhizal fungi through meta-analysis. BMC Evol Biol 16 (1):1–15.

100. **Piculell BJ, José Martínez-García P, Nelson CD, Hoeksema JD**. 2019. Association mapping of ectomycorrhizal traits in loblolly pine (Pinus taeda L.). Mol Ecol 28 (8):2088–2099.

101. **Anderson JT, Willis JH, Mitchell-Olds T**. 2011. Evolutionary genetics of plant adaptation. Trends Genet 27 (7):258–266.

102. **Anderson JT, Lee CR, Rushworth CA, Colautti RI, Mitchell-Olds T**. 2013. Genetic trade-offs and conditional neutrality contribute to local adaptation. Mol Ecol 22 (3):699–708.

103. **Des Marais DL, Hernandez KM, Juenger TE**. 2013. Genotype-by-environment interaction and plasticity: exploring genomic responses of plants to the abiotic environment. Annu Rev Ecol Evol Syst 44:5–29.

104. **Wadgymar SM, Lowry DB, Gould BA, Byron CN, Mactavish RM, Anderson JT**. 2017. Identifying targets and agents of selection: innovative methods to evaluate the processes that contribute to local adaptation. Methods Ecol Evol 8 (6):738–749.

105. **Mee JA, Yeaman S**. 2019. Unpacking conditional neutrality: genomic signatures of selection on conditionally beneficial and conditionally deleterious mutations. The Am Nat 194 (4):529–540.

106. **Harrison TL, Wood CW, Borges IL, Stinchcombe JR**. 2017. No evidence for adaptation to local rhizobial mutualists in the legume *Medicago lupulina*. Ecol Evol 7 (12):4367–4376.

107. **Riley A, Grillo M, Epstein B, Tiffin P, Heath K**. 2021. Partners in space: Discordant population structure between legume hosts and rhizobium symbionts in their native range. https://www.biorxiv.org/content/10.1101/2021.06.30.449460.

108. **Frederickson ME**. 2013. Rethinking mutualism stability: cheaters and the evolution of sanctions. The Q Rev Biol 88 (4):269–295.

109. **Pahua V, Stokes P, Hollowell A, Regus J, Gano-Cohen K, Wendlandt C, Quides K, Lyu J, Sachs J**. 2018. Fitness variation among host species and the paradox of ineffective rhizobia. J Evol Biol 31 (4):599–610.

110. **Vaidya P, Stinchcombe JR**. 2020. The Potential for Genotype-by-Environment Interactions to Maintain Genetic Variation in a Model Legume–Rhizobia Mutualism. Plant Commun 1 (6):100114.

111. **Popovic D, Lowry DB**. 2020. Contrasting environmental factors drive local adaptation at opposite ends of an environmental gradient in the yellow monkeyflower (Mimulus guttatus). Am J Bot 107 (2):298–307.

112. **Mackay TF, Stone EA, Ayroles JF**. 2009. The genetics of quantitative traits: challenges and prospects. Nat Rev Genet 10 (8):565–577.

113. **Wagner MR**. 2021. Prioritizing host phenotype to understand microbiome heritability in plants. New Phytol (Accepted article).

114. **Batstone RT, Burghardt LT, Heath KD**. 2021. The genetic basis of cooperation and conflict in natural populations of a model symbiont. https://www.biorxiv.org/content/10.1101/2021.07.19.452989.

115. **Jones EI, Afkhami ME, Akçay E, Bronstein JL, Bshary R, Frederickson ME, Heath KD, Hoeksema JD, Ness JH, Pankey MS, et al.**. 2015. Cheaters must prosper: reconciling theoretical and empirical perspectives on cheating in mutualism. Ecol Lett 18 (11):1270–1284.

116. **Burdon J, Thrall P**. 1999. Spatial and temporal patterns in coevolving plant and pathogen associations. The Am Nat 153 (S5):S15–S33.

117. **Barrett LG, Broadhurst LM, Thrall PH**. 2012. Geographic adaptation in plant–soil mutualisms: tests using *Acacia* spp. and rhizobial bacteria. Funct Ecol 26 (2):457–468.

118. **Vale PF, Little TJ**. 2010. CRISPR-mediated phage resistance and the ghost of coevolution past. Proc Royal Soc B 277 (1691):2097–2103.

119. **Hoeksema JD**. 2010. Ongoing coevolution in mycorrhizal interactions. New Phytol 187 (2):286–300.

120. **Chong RA, Moran NA**. 2016. Intraspecific genetic variation in hosts affects regulation of obligate heritable symbionts. Proc Natl Acad Sci 113 (46):13114–13119.

121. **Wallace JG, Rodgers-Melnick E, Buckler ES**. 2018. On the road to breeding 4.0: unraveling the good, the bad, and the boring of crop quantitative genomics. Annu Rev Genet 52:421–444.

122. **Clouse KM, Wagner MR**. 2021. Plant genetics as a tool for manipulating crop microbiomes: Opportunities and challenges. Front Bioeng Biotechnol 9.

123. **Heath KD, Burke PV, Stinchcombe JR**. 2012. Coevolutionary ge-netic variation in the legume-rhizobium transcriptome. Mol Ecol 21 (19):4735–4747.

124. **Guefrachi I, Nagymihaly M, Pislariu CI, Van de Velde W, Ratet P, Mars M, Udvardi MK, Kondorosi E, Mergaert P, Alunni B**. 2014. Ex-treme specificity of NCR gene expression in *Medicago truncatula*. BMC Genom 15 (1):1–16.

125. **Burghardt LT, Guhlin J, Chun CL, Liu J, Sadowsky MJ, Stupar RM, Young ND, Tiffin P**. 2017. Transcriptomic basis of genome by genome variation in a legume-rhizobia mutualism. Mol Ecol 26 (21):6122–6135.

126. **Wang Q, Liu J, Zhu H**. 2018. Genetic and molecular mechanisms un-derlying symbiotic specificity in legume-rhizobium interactions. Front Plant Sci 9:313.

127. **Walker L, Lagunas B, Gifford ML**. 2020. Determinants of host range specificity in legume-rhizobia Symbiosis. Front Microbiol 11:3028.

128. **Fagorzi C, Bacci G, Huang R, Cangioli L, Checcucci A, Fini M, Perrin E, Natali C, diCenzo GC, Mengoni A**. 2021. Nonadditive Transcrip-tomic Signatures of Genotype-by-Genotype Interactions during the Initiation of Plant-Rhizobium Symbiosis. MSystems 6 (1):e00974–20.

129. **Gorton AJ, Heath KD, Pilet-Nayel ML, Baranger A, Stinchcombe JR**. 2012. Mapping the genetic basis of symbiotic variation in legume-rhizobium interactions in *Medicago truncatula*. G3: Genes| Genomes| Genet 2 (11):1291–1303.

130. **Ané JM, Kiss GB, Riely BK, Penmetsa RV, Oldroyd GE, Ayax C, Lévy J, Debellé F, Baek JM, Kalo P, et al.**. 2004. *Medicago truncatula dmi*1 required for bacterial and fungal symbioses in legumes. Science 303 (5662):1364–1367.

131. **Wheatley RM, Ford BL, Li L, Aroney ST, Knights HE, Ledermann R, East AK, Ramachandran VK, Poole PS**. 2020. Lifestyle adaptations of Rhizobium from rhizosphere to symbiosis. Proc Natl Acad Sci 117 (38):23823–23834.

132. **Bull JJ, Rice W**. 1991. Distinguishing mechanisms for the evolution of co-operation. J Theor Biol 149 (1):63–74.

133. **West S, Kiers ET, Pen I, Denison R**. 2002. Sanctions and mutualism stability: when should less beneficial mutualists be tolerated? J Evol Biol 15 (5):830–837.

134. **West SA, Kiers ET, Simms EL, Denison RF**. 2002. Sanctions and mu-tualism stability: why do rhizobia fix nitrogen? Proc Royal Soc B 269 (1492):685–694.

135. **Foster KR, Kokko H**. 2006. Cheating can stabilize cooperation in mu-tualisms. Proc Royal Soc B 273 (1598):2233–2239.

136. **Pfau T, Christian N, Masakapalli SK, Sweetlove LJ, Poolman MG, Ebenhöh O**. 2018. The intertwined metabolism during symbiotic ni-trogen fixation elucidated by metabolic modelling. Sci Reports 8 (1):1–11.

137. **Checcucci A, Bazzicalupo M, Mengoni A, Viti C, Dziewit L, Finan TM, Galardini M, Fondi M, et al.**. 2016. Metabolic modelling re-veals the specialization of secondary replicons for niche adaptation in *Sinorhizobium meliloti*. Nat Commun 7 (1):1–10.

138. **diCenzo GC, Finan TM**. 2018. Techniques for large-scale bacterial genome manipulation and characterization of the mutants with re-spect to in silico metabolic reconstructions, p 291–314. *In* Metabolic Network Reconstruction and Modeling. Springer.

139. **Civelek M, Lusis AJ**. 2014. Systems genetics approaches to under-stand complex traits. Nat Rev Genet 15 (1):34–48.

140. **Kliebenstein D**. 2004. Secondary metabolites and plant/environment interactions: a view through *Arabidopsis thaliana* tinged glasses. Plant, Cell & Environ 27 (6):675–684.