Effect of EMS induced mutation in rice cultivar Nagina 22 on salinity tolerance

Arun Shankar[A], OP Choudhary[AB] Dharminder Bhatia [AB] and Kuldeep-Singh[ABC]

A: Punjab Agricultural Univerty, Ludhiana (Punjab), India; Present address: Rama University, Kanpur (Uttar Pradesh), India

AB: Punjab Agricultural Univerty, Ludhiana (Punjab), India Email: opchoudhary@pau.edu, d.bhatia@pau.edu

ABC: Punjab Agricultural Univerty, Ludhiana (Punjab), India; Present address: Director NBPGR, New Delhi, India Email: kuldeep.singh4@icar.gov.in

Abstract

Salinized hydroponic culture experiment with three salinity levels (EC control, 6 and 12 dS/m) was performed to screen salt tolerant mutants of aerobic rice cultivar Nagina 22 and to study the nature of salt tolerance from a total of 432 EMS induced M4 mutants. Plants were harvested 30 days after sowing. Growth parameters viz. root weight, shoot weight, root length, shoot length, Na and K concentrations in shoot and roots were measured. Combined Factor scores of growth parameters was computed by Principle Component Analysis using Minitab software.

At EC 12 dS/m 10 mutants out of 432 were able to survive. At moderate salinity, some mutant lines produced higher shoot weight compared to their respective control showing inverse trend and the effectiveness of EMS induced mutation in inducing salinity tolerance to these mutants. At high salinity only10 mutants survived (remained green) up to the time of 30 days harvest. These mutants performed well in terms of overall growth recording 2.1-2.5 times higher factor score and 8-14 times higher shoot weight compared to the N 22 check. One mutant N22-L-1010 almost completely excluded Na at xylem parenchyma level. Two other mutants N22-L-1013 and N22-L-806 maintained Na exclusion compared to the N22 check. N22 check and mutant N22-L-1009 maintained similar degree of Na exclusion though the N22 check died because it cannot maintain adequate K in shoot.

We conclude that EMS has induced salinity tolerance in some mutants. The study can be advanced further to characterize the putative mutants through molecular genetics approaches.

Introduction

Soil salinity is a major abiotic threat limiting plant growth and development causing yield loss in many crops (Hasegawa et al., 2000; Qadir et al., 2007). Salt-affected soils contain excessive levels of water-soluble salts, especially sodium chloride (NaCl) (Tanji, 2002). Sodium chloride upon ionization produces sodium (Na^+) and chloride (Cl^-) ions causing ionic and osmotic stress in higher plants.(Mansour and Salama, 2000; Chinnusamy et al., 2005). Soil salinity is a very severe problem particularly in semiarid, and coastal rice-producing areas of the tropics (Lee et al., 2003). The rice production in the South Western districts of the state of Punjab, India which lies in semiarid zone is severely affected by soil salinity. The prime cause of salinity in this region is the brackish nature of underground water. Moreover, salinity is usually coupled with water logging due to shallow depth of water table. Hence the farmers prefer to grow rice as compared to other crops it can withstand standing water.

Considered moderately sensitive to salinity (Akbar et al., 1972; Mass and Hoffman, 1977; Mori et al., 1987) rice (*Oryza sativa* L. spp. *indica*) is one of the five main carbohydrate crops responsible for feeding the world's population, and is especially important in Asian countries. Rice has previously been reported as being salt susceptible in both the seedling and reproductive stages (Zeng et al., 2001; Moradi and Ismail, 2007), leading to a reduction in yield of more than 50% when exposed to 6.65 dS m^{-1} ECe (Zeng and Shannon, 2000). Change in farming systems to alleviate salinity may be one crucial approach, however, it is likely to be a long and difficult process because it will require the use of new land and will not address the problem of growing crops in land that is already compromised (Yamaguchi and Blumwald 2005). Therefore, it is crucial to improve tolerance to salinity in rice by breeding high yielding salinity tolerant varieties.

Differential salt tolerance exist among different genera and species, and also within the different parts of the same species (Flowers and Hajibagheri 2001; Ismail, 2003). Studying the response of cultivars of any species to salinity provides a convenient and useful tool for understanding the basic mechanisms involved in salt tolerance (Tammam et al., 2008). Breeding rice varieties having in-built salt tolerance is realized as valuable, resource conserving, more economic and socially acceptable approach. Salt tolerance is a multigenic trait enabling plants to grow and sustain economic yield even when the salt concentration is high particularly NaCl (Hurkman, 1992; Deepa Sankar et al., 2011). Flowers and Yeo (1995) reported five ways to develop salt-tolerant crops: (1) developing halophytes as alternative crops; (2) use of interspecific hybridization to enhance the tolerance of current crops; (3) utilize the already present variation with respect to salt tolerance in existing crops; (4) develop variation within crops by using recurrent selection, mutagenesis or tissue culture, and (5) breeding for economic yield rather than tolerance. Conventional breeding of cultivars for salt tolerance is time consuming requiring multiple crosses for many generations and labor intensive (Peleman and Voort, 2003; Hwa and Yang, 2008). Moreover, traditional breeding methods are not suitable for screening a large sample size for exploiting variations in tolerance.

Chemical mutagens viz. EMS, DEB and sodium azide; irradiation viz. Gamma rays, X rays and fast neutrons have been used to generate large number of functional variations in rice. (Talebi et al, 2012). Chemicals generate mostly point mutations so they are ideal for producing missense and nonsense mutations providing a sequence of functional mutations. Hence, induced mutation technique to create variability among rice germplasm (Mba et al, 2007) and screening of salt resistant cultivars under solution culture (Gregorio et al., 1997)

could become a viable alternative towards tolerance to soil salinity as screening of large number of germplasm under field conditions is difficult due to stress heterogeneity, presence of other soil-related stresses, and the significant influence of environmental factors (Gregorio et al., 1997). Most of the studies focus on the effects of sodium on the plant growth and the importance of control mechanisms for sodium uptake in salt-resistant plants, but little emphasis has been put on root selectivity for potassium viz a viz sodium (Asch et al 2000) which is discussed in this paper. Moreover, large numbers of papers has been published regarding salinity tolerance mechanisms yet very few salt tolerant cultivars has been released offering only slight improvement over parent lines (Flowers and Yeo, 1995; Glenn et al., 1999).

Keeping the above factors in mind, mutation was induced by EMS (Ethyl methane sulfonate) in aerobic rice cultivar Nagina 22 (which is also drought tolerant cultivar). A solution culture experiment was performed to screen salt tolerant mutants by studying their morphological parameters such as root, shoot length and dry weight and to study the underlying mechanism of salt tolerance by studying Na and K concentrations in root and shoot of the promising and worse mutant. Knowledge obtained from such studies will be helpful to characterize the putative mutants through molecular genetics approaches.

Material and Methods

Salinized hydroponics culture experiment with three salinity levels (EC control, 6 and 12 dS/m) was conducted to screen salt tolerant mutant from a total of 432 EMS induced M_4 mutants of drought tolerant rice cultivar Nagina 22. Plants were harvested 30 days after sowing. Root weight, shoot weight, root length, shoot length, Na and K concentrations in shoot and roots were measured. The experiment was conducted at the Greenhouse of School of Agricultural Biotechnology, Punjab Agricultural University Ludhiana situated at 300 56/ N latitude and 750 52/ E longitude and at an altitude of 247 m above mean sea level. Mutations in 'Nagina 22' were induced using 0.5% Ethyl methanesulfonate (EMS). These mutants were produced at IARI, New Delhi and distributed to all the collaborating centres in the country including Punjab Agricultural University. Nagina 22 (N22) was used as check.

Creation of salinity and experimental set up

In control treatment no salts were added. Salts of NaCl and $CaCl_2.2H_2O$ were added in 2:1 on equivalent basis to get the desired EC level. The electrical conductivity of the solution was monitored by the use of conductivity meter. Plastic trays (holding 8 litres of half Hoagland's solution up to the brim) of the size 45 cm x 32 cm x 8 cm were used to hold nutrient solution. A thermocol sheet of the size 45 cm x 32 cm was cut and small holes of 2 cm diameter were made at distance of 5.5 cm (fig.1). A plastic net was fixed on the lower side of the thermocol sheet to hold the seed in the respective holes which remained in touch with the Hoagland's solution in the trays. Five seeds were placed in each hole. The Hoagland's solution was replaced after 3 days. Electrical conductivity was checked periodically during the growth phase and maintained at targeted level. The composition of Hoagland solution is mentioned in Table 5.

Harvesting, collection and processing of shoot samples

Shoot and root length was recorded after 30 days of sowing and plants were harvested. The harvested shoot samples were washed with acidified deionized water, and then with distilled water and finally with double distilled water. The washed shoot samples were air dried and then dried in oven at 65 ± 2^0C. Thereafter, the shoots were weighed for dry matter yield and stored in paper bags. They were digested in diacid mixture of HNO_3 and $HClO_4$ in the ratio 3:1. The digested samples were analyzed for Na and K by flame photometer.

Partition quotient:

The dry weight of root and shoot was calculated as a percentage of total plant weight and mineral uptake of each organ was calculated as a percentage of total plant mineral uptake. Using these values, the normalized partitioning of that mineral within the plant was calculated by dividing each organ's percentage mineral uptake by its percentage DW which will be denoted as partition quotient (PQ) (Waters and Grusak, 2008).

Statistical analysis:

Combined Factor scores of growth parameters was computed by Principle Component Analysis using Minitab software. The correlation analysis was done using data analysis tool in MS excel.

Results and Discussion

The overall growth of screened mutants was quite good at control and EC 6 dS/m but the survival of most of the mutants was poor at EC 12 dS/m where only 10 mutants out of 432 were able to survive. According to Table 1, mutants were categorized based on shoot weight and factor score as good at moderate salinity (Sl. No. 1-10 and 17-24), good at high salinity (Sl. No. 1-10), worse at moderate salinity (Sl. No. 25-28) and worse at high salinity (Sl. No. 12-16). The N22 check was susceptible at moderate and high salinity. The combined factor score based on root length, root weight, shoot length and shoot weight (Table 1) was a good indicator of overall growth of mutants at different salinity levels. The combined factor score declined with salinity in all the mutants including N 22 check. This indicates that salinity generally imposes reduction to overall growth of plants. At moderate salinity, mutant lines (Sl. 17 to 24) produced higher shoot weight compared to their respective controls thereby showing inverse trend. These mutants performed well in terms of shoot length, root length and root weight (data not shown) as these mutants recorded 91-96 % relative factor score and more than twice shoot weight compared to their respective controls. Mutant lines N22-L-1240, N22-L-1195, N22-L-1247 and N22-L-1151 were the worse mutants at moderate salinity in terms of absolute shoot weight and factor score. N22 check was also susceptible compared to the tolerant mutants at moderate salinity performing better only compared to some worst mutants. These results are in

accordance with Danaguiraman et al. (2003) who reported that, tolerant genotypes of rice
recorded a higher germination percentage, root length, shoot length and vigour index.

According to Table 2, K uptake gradually decreased with increase in salinity level
except for the mutants showing inverse trend at moderate salinity (Sr. No. 17-24) while shoot
Na uptake generally increased with salinity in all the mutants. Shoot Na:K ratio increased in all
the mutants compared to their respective controls. However, increase was the maximum in the
succeptible mutants (Sr. No.25-28) showing 1.2-2 times increase compared to the N22 check.
Conversely, the good mutants (Sr. No. 1-10 and 17-24) recorded 30-76% lower Na:K ratio
compared to N22 check. Shoot weight was highly negatively correlated with shoot Na:K ratio
at moderate and high salinity (r^2= -0.82 and r^2= -0.74, respectively). The relationship between
potassium and sodium uptake by the rice plant and tolerance to salinity stress has been reported
by Qadar, (1988); Pandey & Srivastava, (1991); Bohra & Dörffling, (1993).

Plants use three main mechanisms to survive salinity stress: osmotic tolerance, Na^+
exclusion and Na^+ tissue tolerance (Munns and Tester 2008). Earlier Cl^- was thought to be
main cause of salt injuries but later it was confirmed that excessive uptake of Na^+ is the chief
cause of salt injuries to plants (Clarkson and Hanson 1980). Salt injuries include injuries
related to osmotic effects (especially in the initial growth stages (Lauchli and Grattan, 2007).
Hence, exclusion of the sodium ion (Na^+) from shoot and increased absorption of K^+ to
maintain a good Na^+: K^+ balance in the shoot are very important traits of tolerant genotypes
(Roshandel and Flowers 2009).

Blumwald (2000) reported that ion ratios in plants are altered by the influx of Na^+
through K^+ pathways. The similarity of the hydrated ionic radii of Na^+ and K^+ makes it difficult
to discriminate between them and this is the basis of Na^+ toxicity. However, at moderate
salinity, most of the mutants marinated fairly lower Na:K ratio (Table 2) except for the
sensitive mutants (Sr. 25-28) leading to their better performance compared to the sensitive
mutants. Kader and Lindberg (2008) reported that the transient uptake of Na^+, which occurs
only in the tolerant cultivar (Na exclusion), and the fast compartmentalization of Na^+ into the
vacuole (tissue tolerance), probably are the most important cellular trait for Na^+ tolerance in
rice. Later, Munns and Tester (2008) in their review article suggested that Na exclusion in
shoots is associated with partitioning of Na to roots while tissue tolerance is associated with
compartmentalization of Na^+ into the vacuole and Na transport can be controlled by tolerant
plants at moderate salinity. Hence, the susceptibility of worse mutants to moderate salinity
could be explained largely by their higher Na:K ratio compared to the N22 check. Roshandel
and Flowers (2009) suggested that salt-tolerant cultivars of rice accumulate less Na^+ and more
K^+ in comparison to susceptible ones. Results of this study reveal that at moderate salinity,
plants adopt sodium avoidance mechanism by excluding sodium to maintain a low Na:K ratio.
Also Munns (2002) has suggested that Na to K selectivity of plants roots minimizing the entry
of Na into plants and maintaining effective uptake of K together to the shoots plays crucial role
in salinity tolerance to plants.

At high salinity, only 10 mutants survived (remained green) up to the time of 30 days
harvest (Sr. No. 1-10). The N 22 check did not survive. These mutants performed well in terms
of overall growth recording 2.1-2.5 times higher factor score and 8-14 times higher shoot
weight compared to the N 22 check. Five of the worse mutants (Sr. No. 12-16, Table 1) at high
salinity recorded 61-89 % lower shoot weight compared to N22 check. At high salinity, the
shoot Na:K ratio ranged between 0.36 to 10.92 for the good mutants (Sr. No. 1-10, Table 2)
and 10.98-37.69 for the worst mutants (Sr. No. 12-16, Table 2). The N22 check recorded a

value of 5.72. The other worse mutants died because they cannot maintain Na exclusion at the xylem parenchyma level. One mutant N22-L-1010 almost completely excluded Na at xylem parenchyma level. Two other mutants N22-L-1013 and N22-L-806 maintained Na exclusion compared to the N22 check. N22 check and mutant N22-L-1009 maintained similar degree of Na exclusion though the N22 check died because it cannot maintain adequate K in shoot from roots at the xylem parenchyma level. When salinity results from an excess of NaCl, the most common type of salt stress, the concentrations of sodium (Na^+) and chloride (Cl^-) in the plant increase, and concentrations of potassium (K^+) and calcium (Ca^+) are reduced. This is known as salt-specific or ion-excess effect of salinity. Increases in salt concentrations more than optimum value can lead to toxic accumulation of ions such as Cl^- and, in particular, Na^+ exerting ion toxicity at cellular level with varying concentrations according to the capacity of salt tolerance of plants (Diedhiou, 2006).

Exclusion of Na from leaves is facilitated by two mechanisms: first, an outward active transport of Na at the plasma membrane of root cortical cells resulting into extrusion of Na to the outward medium (Jeschke, 1984, Pitman and Saddler, 1967). Second, Resorption of Na ions from the xylem sap and its accumulation by xylem parenchyma cells in the roots and stem base reducing amount of Na reaching the leaves (Lauchli, 1984). In this study, three mutants N22-L-1010, N22-L-1009 and N22-L-998 maintained lesser relative Na: K Partition Quotient (PQ) ratio compared to N22 check indicating Na to K discrimination at the xylem parenchyma level. Notably, most of the other good mutants survived in spite of having higher relative Na: K PQ ratio than N22 check. This is in accordance to the Na tissue tolerance mechanism. All the worse mutants maintained higher relative Na: K PQ ratio compared to N22 check showing poor Na: K discrimination. Sodium accumulation at toxic levels results into necrosis of leaves, shortening of lifespan of leaves, growth and yield reductions associated to the capacity of Na to compete with K for binding sites essential for cellular function (Tester and Davenport, 2003).

According to Table 3, PQ of K in roots at high salinity decreased for six mutants and N22 check compared to their respective controls while four mutants N22-L-806, N22-L-1010, N22-L-1013 and N22-L-834 recorded an increase. The N22 check recorded lowest PQ of K in shoots while all the good mutants recorded 1.5-2.5 times higher PQ of K in shoots compared to the N 22 check. Notably, mutants characterized as worse recorded higher (4.3-8.5 times) PQ of K in shoots compared to N22 check. Under control EC, the N22 check recorded higher PQ of Na in shoots compared to all other mutants (good or worse). Regarding PQ of Na in shoots, all the mutants accumulated either higher or similar Na compared to N22 check except one mutant line, N22-L-1010 which recorded 90% lower shoot Na accumulation compared to N22 check. However, the worse mutants on the other hand recorded 2.3 to 13.9 times higher shoot Na accumulation except one mutant N22-L-931(1.7 times) while the good mutants recorded (0.1-1.8 times). Among the good mutants, N22-L-1009, N22-L-1010, N22-L-1013 and N22-L-998 recorded 22, 89, 22 and 11 % lesser Na in shoots compared to N22 check. N22 check died at shoot Na PQ value of 0.9. Considering this value as the threshold value all the worse mutants accumulated higher shoot Na PQ leading to their poor performance at high salinity.

The PQ of Na in shoots was lower in good mutants compared to worse mutants. Intracellular K^+/Na^+ balance is an important factor in salt tolerance, because selectively taking up and maintaining a high level of K^+ restricts the entry of Na^+ (Blumwald 2000). According to Table 4, one mutant N22-L-1010 almost completely excluded Na at xylem parenchyma level through excluding Na from shoots. Two other mutants N22-L-1013 and N22-L-806 also maintained shoot Na exclusion compared to the N22 check. This mode of Na exclusion has

been studied in *Triticeae* (Greenway and Munns, 1980; Gorham et al., 1985) and barley (Forster et al., 1994).

The characteristic of Na exclusion mechanism for salinity tolerance is lower Na accumulation in shoots by partioning excess Na to roots and this does not involve higher K uptake (Munns and Testor, 2008). As evident from Table 4, the mutants N22-L-1013, N22-L-806 and N22-L-1010 excluded Na in the shoots to survive. At the same time it was not accompanied by K accumulation in shoots indicating these mutants are true Na excluders. The worst mutants were not able to exclude Na from shoots hence they could not survive. The N22 check was not able to maintain adequate Na:K ratio and performed poorly.

Conclusions

Based on our studies we conclude that EMS has induced variation in mutants with respect to their response to different levels of salinity and that can be exploited to identify the tolerant as well as susceptible mutants to salinity. N22-L-1013, N22-L-806 and N22-L-1010 were identified as typical Na excluders while rest of the tolerant mutants showed tissue tolerance mechanism.

Acknowledgment
Authors are highly acknowledging the financial support provided by the Indian Council of Agricultural research (ICAR) in conducting this research.

References

Akbar M, Yabuno Y, Nakao S (1972) Breeding for saline resistant varieties of rice. I. Variability for salt-tolerance among some rice varieties. Jpn. J. Breed 22: 277-284.

Asch F, Dingkuhn M, Dorffing K and Meizan K (2000) Leaf K/Na ratio predicts salinity induced yield loss in irrigated rice. Euphytica 113: 109-118.

Blumwald E (2000) Sodium transport and salt tolerance in plants. Curr Opin In Cell Biol 12: 431–434.

Bohra JS & Dörffling K (1993) Potassium nutrition of rice (Oryza sativa L.) varieties under NaCl salinity. Plant and Soil 152: 299–303.

Chi-Min Hwa & Xian-Ci Yang (2008) The AtMKK3 pathway mediates ABA and salt signaling in Arabidopsis Acta Physiologiae Plantarum volume 30, pages277–286

Chinnusamy, V.; Jagendorf, A.; Zhu, J.K. 2005. Understanding and improving salt tolerance in plants. Crop Science 45: 437-448.

Clarkson DT and Hanson JB (1980) The mineral nutrition of higher plants Ann. Rev. Plant Physiol 31:239-298,

Deepa Sankar, P., N. Subbaraman and S. Lakshmi Narayanan. 2009. Assessment of embryogenic calli derived from temperature sensitive genic male sterility based rice hybrids for salt tolerance. J Ecotoxicol Environ Monit 19(3): 219-224.

Djanaguiraman M, Ramadass R, Durga-Devi D (2003) Effect of salt stress on germination and seedling growth in rice genotypes. Madras Agric J 90: 50-53.

Flowers TJ , Yeo AR. (1995) Breeding for salinity resistance in crop plants—where next? Austr J Plant Physiol 22:875–884.

Flowers TJ and Hajiabagheri MA (2001) Salinity tolerance in Hordeum vulgare : Ion concentrations in root cells of cultivars differing in salt tolerance. Plant Soil 231: 1-9.

Forster BP (1994) Cytogenetic manipulations in the Triticeae. In: Yeo AR, Flowers TJ, eds, Soil mineral stresses: approaches to crop improvement, Vol. 21. Berlin: Springer-Verlag, 61–82.

Glenn EP, Brown JJ, Blumwald E (1999) Salt tolerance and crop potential of halophytes. Critical Reviews Plant Sci 8:227–255.

Gorham J, McDonnell E, Budrewicz E, Wyn Jones RG (1985) Salt tolerance in the Triticeae: growth and solute accumulation in leaves of Thinopyrum bessarabicum. J Exp Botany 36:1021–1031.

Greenway H & Munns R (1980) Mechanisms of salt tolerance in non-halophytes. Annual Review of Plant Physiology31:149–190.

Gregorio G.B, Dharmawansa S and Mendoza RD (1997) Screening rice for salinity tolerance. IRRI Discussion Paper Series NO.22 Int Rice Res Inst   Manila, Philippines. pp.2-23.

Hasegawa PM, Bressan RA, Zhu JK, Bohnert HJ (2000) Plant cellular and molecular responses to high salinity. Annual Review Plant Physiol and Mol Biol 51: 463-499.

Hurkman WJ (1992) Effect of salt stress on plant gene expression : A review. Plant and Soil, 146: 145-151.

Ismail AM (2003) Effect of salinity on physiological responses of selected lines/variety of wheat. Acta Agron 51: 1-9.

Jeschke W D (1984) K+–Na+ exchange at cellular membranes, intracellular compartmentation
of cations, and salt tolerance. In Salinity tolerance in plants: Strategies for crop
improvement. Eds. J D Staples and G H Toenniessen. pp. 37–66. Wiley, New York.

Kader MA & Lindberg S (2008) Cellular traits for sodium tolerance in rice (Oryza sativa L).
Plant Biotechnol. 25: 247-255.

Läuchli A. (1984) Salt exclusion: an adaptation of legumes for crops and pastures under saline
conditions. In: Staples RC, ed. Salinity tolerance in plants: strategies for crop
improvement. New York, NY: Wiley, 171–187.

Läuchli A & Grattan SR (2007) Plant growth and development under salinity stress in
Advances in molecular-breeding towards salinity and drought tolerance, M. A. Jenks, P.
A. Hasegawa, and S. M. Jain, eds., Springer-Verlag, New York, 1–31

Lee KS, Choi WY, Ko JC, Kim TS and Gregoria GB (2003) Salinity tolerance of japonica and
indica rice (Oryza sativa L.) at the seedling stage. Planta 216:1043–1046.

Maas EV, Hoffman GJ (1977) Crop salt tolerance–current assessment. J Irrig drain Division
of the American Society of Civil Engineering 103:115-–134.

Mansour MMF and Salama KHA (2000) Cellular basis of salinity tolerance in plants.
Environ Exp Botany 52: 113-122.

Mba C, Afza R, Jain SM et al (2007) Induced mutations for enhancing salinity tolerance in
rice. In: Jenks MA, Hasegawa PM, Jain SM (eds) Advances in molecular breeding
towards drought and salt tolerant crops. Springer, Berlin, pp 413–454

Moradi F, Ismail AM (2007) Responses of photosynthesis, chlorophyll fluorescence and ROS-
scavenging systems to salt stress during seedling and reproductive stages in rice. Ann
Bot 99:1161-1173.

Mori I, Kinoshita T (1987). Salt tolerance in rice callus clones. Rice Genet. Newsl. 4: 112-113

Munns, R., and Tester, M. (2008) Mechanisms of salinity tolerance. Ann. Rev. Plant Biol. 59:
651–681.

Pandey UK & Srivastava RDL (1991) Leaf potassium as an index of salt tolerance in paddy.
Nat Acad Sci Let 14: 161–164.

Peleman JD, van der Voort J R (2003) Breeding by design. Trend Plant Sci 8:330-334

Pitman MG, Saddler HDW (1967) Active sodium and potassiumtransport in cells of barley
roots. Proc Natl Acad Sci USA 57:44-49

Qadar A, (1988) Potassium status of the rice shoot as Index for salt tolerance. Ind J Plant
Physiol 31: 388–393.

Qadir M, Oster JD, Schubert S, Noble AD, Sahrawat KL (2007) Phytoremediation of sodic and
saline-sodic soils. Adv Agron 96: 197-247.

Munns R (2002) Comparative physiology of salt and water stress Plant Cell Environ 25:239-
250

Roshandel P & Flowers T (2009) The ionic effects of NaCl on physiology and gene
expression in rice genotypes differing in salt tolerance. Plant Soil 315: 135–147.

Yamaguchi T & Blumwald E (2005) Developing salt-tolerant crop plants: challenges and
opportunities. Trends Plant Sci 10: 615-620

Talebi AB & Shahrokhifar B (2012) Ethyl methane sulphonate (EMS) induced mutagenesis in
malaysian rice (cv. MR219) for lethal dose determination. Am J Plant Sci 3:1661–1665.

Tammam AA, Abou Alhamd MF and Hemeda M (2008) Study of salt tolerance in wheat
(Triticum aestium L.) cultivar Banysoif 1. Austr J Crop Sci 1:115-125.

Tanji KK (2002) Salinity in the soil environment. p. 21-51. In: Lauchli A.; Luttge U., eds.
Salinity environment-plant - molecules. Kluwer Academic, Dordrecht, The
Netherlands.

Tester M & Davenport R (2003) Na+ tolerance and Na+ transport in higher plants. Ann Bot
91:503–527.

Waters B M and Grusak M A (2008) Whole-plant mineral partitioning throughout the life cycle
in Arabidopsis thaliana ecotypes Columbia, Landsberg erecta, Cape Verde Islands, and
the mutant line ysl1ysl3. New Phytol 177: 389–405.

Zeng L & Shannon MC (2000) Salinity effects on seedling growth and yield components of
rice. Crop Sci 40:996-1003

Table 1: Growth parameters and factor score of N22 mutants at three salinity levels

Sl. No.	Mutant ID	Low Salinity (EC control)			Moderate salinity (EC 6 dS m^{-1})			High salinity (EC 12 dS m^{-1})		
		Shoot weight (mg)	Root weight (mg)	Factor Score	Shoot weight (mg)	Root weight (mg)	Factor Score	Shoot weight (mg)	Root weight (mg)	Factor Score
1	N22-L-1009	1022	269	97.1	702	209	79.3	212	37	40.5
2	N22-L-991	1641	506	94.2	1013	236	76.4	220	75	41.1
3	N22-L-806	913	340	95.8	818	135	76.2	208	5	36.0
4	N22-L-1010	1230	298	94.8	846	139	81.1	318	15	40.5
5	N22-L-1013	1195	298	94.0	861	294	81.1	185	4	36.9
6	N22-L-1007	1200	262	91.8	863	232	74.2	185	75	41.1
7	N22-L-1012	1181	235	91.3	968	220	78.8	211	75	34.2
8	N22-L-927	1454	510	83.4	970	127	66.1	145	51	36.5
9	N22-L-1011	827	262	91.8	1079	230	77.5	190	150	35.8
10	N22-L-998	907	296	95.2	1297	283	80.2	248	53	37.5
11	**N22 check**	512	266	53.9	135	61	39.7	18	27	16.3
12	N22-L-834	1022	133	73.2	273	79	57.4	4	26	19.1
13	N22-L-954	906	297	78.1	360	274	63.1	2	27	19.7
14	N22-L-931	1429	388	83.7	563	211	66.9	6	14	21.8
15	N22-L-893	752	231	80.6	345	206	63.9	7	21	23.7
16	N22-L-939	1123	343	86.2	385	186	71.7	3	46	21.9
17	N22-L-1109	356		64.9	930		62.0			
18	N22-L-1111	293		61.0	918		57.0			
19	N22-L-1094	306		62.5	886		59.5			
20	N22-L-1098	394		68.5	825		64.2			
21	N22-L-1092	391		67.0	809		61.9			
23	N22-L-1108	295		69.1	751		63.0			
24	N22-L-1090	316		63.6	750		58.9			
25	N22-L-1240	313		52.2	59		35.7			
26	N22-L-1195	125		46.0	52		36.0			
27	N22-L-1247	396		50.7	49		36.6			
28	N22-L-1151	438		52.0	47		36.8			

388   Table 2: Shoot Na and K uptake of some mutants at low, moderate and high salinity levels
389

Sl. No.	Mutant ID	Low Salinity (EC control)				Moderate salinity (EC 6 dS m⁻¹)		High salinity (EC 12 dS m⁻¹)			
		Shoot Na uptake	Shoot K uptake	Root Na uptake	Root K uptake	Shoot Na uptake	Shoot K uptake	Shoot Na uptake	Shoot K uptake	Root Na uptake	Root K uptake
1	N22-L-1009	11.2	275	1.6	182	73.7	214	35.4	86	28.3	11.9
2	N22-L-991	23.3	315	3.8	268	187	330	530	62	63.6	12.8
3	N22-L-806	14.7	379	2.2	154	112	249	337	163	52.7	12.8
4	N22-L-1010	18.5	451	1.6	143	87.1	308	59.8	167	361	46.8
5	N22-L-1013	15.3	424	2.7	176	68.0	316	353	104	150	8.3
6	N22-L-1007	23.4	308	1.1	128	138	283	596	54.6	84.8	11.6
7	N22-L-1012	18.0	417	1.6	145	147	315	333	86.5	55.4	15.1
8	N22-L-927	20.1	429	5.4	228	93.1	342	529	94.8	22.8	11.5
9	N22-L-1011	17.9	455	2.2	150	69.1	323	304	82.7	36.9	18.3
10	N22-L-998	13.6	331	2.2	149	162	409	87	126	53.3	10.6
11	**N22 check**	13.3	428	1.1	181	81	99.3	22.3	3.9	42.9	3.8
12	N22-L-834	36.8	413	4.8	54	41.2	48.0	60.3	1.6	40.8	17.0
13	N22-L-954	15.2	418	5.0	137	69.1	90.0	69.0	3.3	35.3	6.1
14	N22-L-931	54.3	1292	14.7	351	79.9	124	52.7	4.8	65.2	1.3
15	N22-L-893	20.1	382	6.2	117	42.8	70.0	55.4	4.2	48.9	1.0
16	N22-L-939	10.9	356	3.3	109	45.8	90.1	66.3	2.9	20.1	9.0
17	N22-L-1109	20.6	390	-	-	259	789	-	-	-	-
18	N22-L-1111	10.3	200	-	-	360	666	-	-	-	-
19	N22-L-1094	14.1	370	-	-	282	764	-	-	-	-
20	N22-L-1098	15.2	308	-	-	281	870	-	-	-	-
21	N22-L-1092	15.2	417	-	-	73.6	377	-	-	-	-
23	N22-L-1108	17.4	253	-	-	238	701	-	-	-	-
24	N22-L-1090	4.4	62.3	-	-	237	633	-	-	-	-
25	N22-L-1240	7.5	280	-	-	140	87	-	-	-	-
26	N22-L-1195	7.1	140	-	-	87	84	-	-	-	-
27	N22-L-1247	13.5	428	-	-	103	102	-	-	-	-
28	N22-L-1151	7.0	99	-	-	101	96	-	-	-	-

Table 3: Partition quotient of Na in shoot and root of some mutants at high salinity and their respective controls

Sl. No.	Mutants	EC control				EC 12 (dS/m)			
		PQ Na roots	PQ K roots	PQ Na shoot	PQ K shoot	PQ Na roots	PQ K roots	PQ Na shoot	PQ K shoot
1	N22-L-1009	0.6	191	1.1	0.8	3.0	81.7	0.7	1.0
2	N22-L-991	0.6	195	1.1	0.7	0.4	67.4	1.2	1.1
3	N22-L-806	0.5	106	1.2	1.0	5.8	310	0.9	0.9
4	N22-L-1010	0.4	124	1.1	0.9	19.0	486	0.1	0.8
5	N22-L-1013	0.8	147	1.1	0.9	14.1	350	0.7	0.9
6	N22-L-1007	0.2	164	1.2	0.9	0.4	60.5	1.2	1.2
7	N22-L-1012	0.5	155	1.1	0.9	0.5	56.6	1.2	1.2
8	N22-L-927	0.8	134	1.1	0.9	0.2	41.7	1.3	1.2
9	N22-L-1011	0.5	103	1.2	1.0	0.2	41.1	1.6	1.5
10	N22-L-998	0.6	126	1.1	0.9	2.2	44.1	0.8	1.1
11	**N22 check**	0.2	86.8	1.4	1.1	1.1	82.6	0.9	1.3
12	N22-L-834	1.0	100.0	1.0	1.0	0.5	105	4.5	0.6
13	N22-L-954	1.0	99.9	1.0	1.0	0.4	69.7	9.6	5.1
14	N22-L-931	1.0	100.0	1.0	1.0	0.8	30.2	1.5	2.6
15	N22-L-893	1.0	100.0	1.0	1.0	0.6	24.9	2.1	3.3
16	N22-L-939	1.0	100.0	1.0	1.0	0.2	80.5	12.5	4.0

422 Table 4: Partitioning of Na and K in shoots compared to their respective controls
423

Mutant lines	Relative Na PQ shoots: roots	Relative K PQ shoots: roots
N22-L-1009	0.127	2.921
N22-L-991	1.636	4.544
N22-L-806	0.065	0.308
N22-L-1010	0.002	0.226
N22-L-1013	0.036	0.421
N22-L-1007	0.500	3.619
N22-L-1012	1.091	3.656
N22-L-927	4.727	4.269
N22-L-1011	3.333	3.755
N22-L-998	0.198	3.498
N22 check	0.117	1.242
N22-L-834	9.000	0.569
N22-L-954	24.000	7.310
N22-L-931	1.875	8.609
N22-L-893	3.500	13.253
N22-L-939	62.500	4.969

Table 5: Composition of Hoagland's solution used:

The following Stock solutions were made:
1. Ca $(NO_3)_2.4H_2O$ - 236.1 g/l
2. KNO_3 - 101.1 g/l
3. KH_2PO_4 - 136.1 g/l
4. $MgSO_4.7H_2O$ - 246.5 g/l
5. Trace elements (made up to 1 L)
H_3BO_3 - 2.8 g
$MnCl_2.4H_2O$ - 1.8 g
$ZnSO_4.7H_2O$ - 0.2 g
$CuSO_4.5H_2O$ - 0.1 g
$NaMoO4$ 0.025 g

6. FeEDTA
EDTA.2Na - 10.4 g
$FeSO_4.7H_2O$ - 7.8 g
KOH - 56.1 g
To make up 1 L of KOH, pH was adjusted to ~5.5 using H_2SO_4. Then EDTA.2Na and $FeSO_4.7H_2O$ were added.

To make 1 L half Hoagland's solution from these stocks, following quantities were added
7 ml Ca $(NO_3)_2$ stock
5 ml KNO_3
2ml KH_2PO_4
2 ml $MgSO_4$
1 ml Trace elements
1 ml FeEDTA
to 1 L water

Fig.1. A Set of Sheets ready for Hydroponics trial

Fig.2. Tolerant mutants compared with N22 check at high salinity.

Fig. 3 N22 mutants at three salinity levels

Fig. 4 Partition quotient (PQ) of Na and K of some mutants in roots shoot at control EC

Partition quotient (PQ) of Na in some mutants in roots and shoot at high salinity

Partition quotient (PQ) of K in some mutants in roots at high salinity

Fig. 5 Partition quotient (PQ) of K in some mutants in shoot at high salinity