1 ***But Mouse, you are not alone*: On some severe acute respiratory syndrome**
2 **coronavirus 2 variants infecting mice**
3

Michael J. Kuiper, PhD, is a Biomolecular Modeller who leads the Modelling and Simulations Team in at
the CSIRO Data61, Dockands, Victoria, Australia

Laurence OW Wilson, PhD, is a Research Scientist who leads the Digital Genome Engineering Team at
the CSIRO Transformational Bioinformatics Group, North Ryde, New South Wales, Australia

Shruthi Mangalaganesh, is studying Bachelor of Biomedical Science and Bachelor of Commerce at
Monash University, Clayton, Victoria, Australia, and pursued an internship at the CSIRO Australian
Centre for Disease Preparedness, Geelong, Victoria, Australia

Carol Lee, BAnVetBioSci(Hons), PhD, is CERC Postdoctoral Research Fellow in the CSIRO
Transformational Bioinformatics Group, North Ryde, New South Wales, Australia

Daniel Reti, BE(Hons), BS, worked as a Research Technician at the CSIRO Transformational
Bioinformatics Group, North Ryde, New South Wales, Australia, and is currently Front-End Software
Engineer at the Centre for Population Genomics, joint initiative of the Garvan Institute of Medical
Research, Sydney, New South Wales, Australia and the Murdoch Children's Research Institute,
Melbourne, Victoria, Australia

Seshadri S Vasan, DPhil, is a Senior Principal Research Consultant and COVID-19 Project Leader at the
CSIRO Australian Centre for Disease Preparedness, Geelong, Victoria, Australia, and Honorary Professor
in the Department of Health Sciences at the University of York, York, UK

Address correspondence to Professor Seshadri S Vasan, CSIRO Australian Centre for Disease
Preparedness, 5 Portarlington Road, Geelong, 3220, Victoria, Australia or email vasan.vasan@csiro.au
23
24 NB: MJK and LOWW contributed equally to this paper.
25
26 **Running Title:** On some SARS-CoV-2 variants infecting mice

27
Abstract

In silico predictions combined with *in vitro*, *in vivo* and *in situ* observations collectively suggest that

mouse adaptation of the SARS-CoV-2 virus requires an aromatic substitution in position 501 or position

498 (but not both) of the spike protein's receptor binding domain. This effect could be enhanced by

mutations in positions 417, 484, and 493 (especially K417N, E484K, Q493K and Q493R), and to a lesser

extent by mutations in positions 486 and 499 (such as F486L and P499T). Such enhancements due to

more favourable binding interactions with residues on the complementary angiotensin-converting

enzyme 2 (ACE2) interface, are however, unlikely to sustain mouse infectivity on their own based on

theoretical and experimental evidence to date. Our current understanding thus points to the Alpha, Beta

and Gamma variants of concern infecting mice, while Delta and 'Delta Plus' lack a similar biomolecular

basis to do so. This paper identifies eleven countries (Brazil, Chile, Djibouti, Haiti, Malawi, Mozambique,

Reunion, Suriname, Trinidad and Tobago, Uruguay and Venezuela) where targeted local field

surveillance of mice is encouraged because they may have come in contact with humans who had the

virus with adaptive mutation(s). It also provides a systematic methodology to analyze the potential for

other animal reservoirs and their likely locations.

Keywords

AlphaFold; Animal reservoir; COVID-19; *in silico*; *in vitro*; *in vivo*; mouse adaptation; SARS-CoV-2; variants

Introduction

The 'novel coronavirus disease' (COVID-19) has resulted in significant global morbidity and mortality on

a scale similar to the influenza pandemic of 1918.[1] The ongoing pandemic has been sustained through

the activities of human beings, who are the largest reservoir of the causative 'severe acute respiratory

syndrome virus 2' (SARS-CoV-2). This RNA virus in a new host (human beings) is evolving rapidly,

accumulating mutations, and existing as a cloud of variants with quasispecies diversity.[2] Last year, the

world witnessed the risk of this virus acquiring additional reservoirs (such as minks) and new mutations

of consequence (such as 'Cluster 5'), which could increase transmissibility and lead to a potentially

weaker antibody response.[3]

Both the original form of the virus (known as 'D614') and the subsequent more transmissible 'G614'

variant which has replaced it almost entirely in circulation,[4,5] did not infect mice because their

'angiotensin-converting enzyme 2' (ACE2) receptor did not bind the viral spike protein effectively to

allow entry into cells. Since mouse (*Mus musculus*) is a popular animal model of infection, the virus had

to be adapted through techniques such as sequential passaging in mouse lung tissues and modifying the

receptor binding domain (RBD).[6-8] Other strategies to infecting mice with the original form of the virus or

the G614 included transgenic mice expressing human ACE2 (hACE2), and sensitizing the mouse

respiratory tract through transduction with adenovirus or adeno-associated virus expressing hACE2.[9-12]

Recently, virus 'variants of concern' (VOC) originating from Brazil, South Africa and the UK (which

contain the common mutation N501Y in the RBD) were shown to infect mice.[13,14] This ability of SARS-

CoV-2 variants of concern to infect mice is unsettling because of the potential to establish additional

reservoirs in a species that is in close contact with people and companion animals, especially as its

population is hard to vaccinate or control.

Methodology

In this paper, we have combined our structural predictions from biomolecular modelling with available

experimental evidence to understand how specific mutations in VOC and mouse adapted strains,

especially in the RBD, have enabled this virus to infect mice. To do this, we compared the spatial

interactions of these key mutations with the corresponding regions in both mouse ACE2 (mACE2) and

hACE2 using homology molecular models based on cryogenic electron microscopy (cryo-EM) data of the

ACE2/RBD interface (protein database 'pdb' entry number 6M17).[15] We have assumed the coordinates

of ACE2 and RBD in the 6M17 structure to be reasonable starting positions for our respective models,

with root-mean-square deviation (RMSD) scripts alignment based on C-Alpha carbon protein backbones.

As the protein prediction software *AlphaFold* has been recently been released,[16] we also recalculated

RBD structures, finding *AlphaFold* predictions to be in excellent agreement with our initial homology

models, with less than 0.9 Å RMSD average alignment. Amino acid side chain conformations of the RBD

models were adapted from *AlphaFold* predictions as they were deemed more reliable.

By introducing the ACE2 models and variants into molecular dynamics simulations to optimize the

interactions, similar to our recent work,[17,18] we qualitatively identified the key interactions of side chain

residues at the ACE2/RBD interface. Further details are provided under 'Supplementary Methods for

Molecular Modelling'. We then compared our *in-silico* findings with experimental results reported by

different research groups with mouse adapted strains/isolates[6-8,13,21-29] and VOC[14,30-34] (**Tables 1 and 2**).

Taken together, we were able to gain valuable insights into the likely effects of different mutations of

consequence.

Subsequently, we queried the world's largest public database called 'GISAID' (Global Initiative on

Sharing All Influenza Data) and looked for these mutations, and equivalent mutations of consequence.

This 'Big Data', comprising of circa 2.4 million and 4 million SARS-CoV-2 genome sequences as of 21 July

and 4 October 2021 respectively, formed the basis of our *in situ* analysis.[29] Over 99.9% of these

sequences are from viruses sequenced from human hosts. In brief, sequences were aligned to the back

to the SARS-CoV-2 reference (EPI_ISL_402124, denoted as the WT allele) in order to generate a file

containing all mutations in the variant call format (VCF), which is a concise way for storing gene

sequence variations. We then calculated the ratio of the frequency at which the WT allele versus the

mutant allele were observed across a rolling 14-day window. Given the data is discrete, highly variable in

size across countries, and contains background noise, such a window of time is essential from our

experience to reduce distortions and glean meaningful insights. For example, if 14 WT and 28 mutant

sequences were observed within a specified 14-day window, then the frequencies were 1/day and 2/day

respectively. The ratio of mutant:WT frequency is therefore 2, indicating the mutant allele is appearing

twice as frequently as the WT allele within that period. This was used to create a heatmap for the key

mutations, both individually and in combination (as discussed below). For clarity, we only considered

countries where the mutant:WT ratio exceeded 1.2 across the entire time the mutant was sampled or

within any given 14-day window. To reduce the noise further from low-sampled countries, we also

instituted a minimum threshold of 10 WT and 10 mutant samples over at least 14 days to suggest

possible spread locally. For this reason, a country where a mutant had only been recorded on a single

day, or where only 9 mutant samples were recorded overall, was not included in our analysis. Our

heatmap scale has also been truncated at 2.00 for ease of visual comparison.

Results and Discussion

***In silico* results**

In silico comparison of the interface residues of the RBD/ACE2 complex in human and mouse models

reveals 30 ACE2 residues in close contact, of which 19 are conserved between mACE2 and hACE2

(approximately 63% identity for the contacting ACE2 residues). The RBD mutations associated with the

mouse adaptations listed in **Table 1** can be grouped loosely into 3 regions by their positions on the

ACE2/RBD interface as follows: Region 1 (RBD positions 498, 499 and 501) are centered around the

highly conserved ACE2 residue tyrosine 41 (Y41); Region 2 (RBD positions 417, 493) centered around

ACE2 residue 34; and Region 3 (RBD positions 484 and 486) close to a cluster of ACE2 residues 78 to 82.

ACE2 residues within 5 Å (chosen to account for molecular fluctuations) of each RBD adaptation are

listed in **Table 3**, with dissimilar human/mouse ACE2 residues highlighted in yellow. **Figure 1** further

illustrates these residues visually, and how they are relatively positioned in three regions. **Figure 2** aligns

the human and mouse ACE2 to highlight the key differences at the contact points with the RDB (shown

in yellow).

Modelling the N501Y mutation at the ACE2/RBD interface reveals a close interaction with the highly

conserved Y41 residue in ACE2, through attractive, non-covalent bonding between aromatic amino acids

known as π-stacking interactions. In mACE2, π-stacking can be enhanced by the proximal substitution of

histidine (H) in place of lysine (K) which is present at position 353 of hACE2. Our modelling also shows

similar π-stacking enhancement with the conserved Y41, through the substitution of RBD glutamine at

position 498 (Q498) to either histidine (H) or tyrosine (Y). These aromatic π-stacking interactions appear

to be reasonably strong as N501Y and Q498H can each sustain mouse adaptation on its own in

experiments.[6,13,23,28] Counterintuitively, our modelling predicts that simultaneous aromatic mutations at

RBD positions 498 and 501 is detrimental to mouse adaptation due to local π-stacking distortion to the

binding interface. This could explain why very few simultaneous aromatic mutations at these positions

have been observed. In over 2.4 million entries on GISAID as of 21 July 2021, we detected only a single

co-occurrence, that of an adapted isolate from mouse lung homogenate where N501Y occurred with

glutamate 498 to arginine (Q498R).[29] However, when we recently re-run our *in situ* model with over 4

million human-origin GISAID entries, we detected 0, 2 and 14 instances of Y501 co-existing with Y498,

H498 and R498 respectively. The two instances of Y501-H498 are low quality and/or low coverage

sequences; the 14 instances of Y501-R498 have been reported from France, Netherlands, South Africa,

Spain, UK and USA between March and September 2021. Although arginine is not aromatic, it is

frequently associated with π-stacking interactions with inherent conformational flexibility compared to

tyrosine or histidine, therefore it is likely that the Y501-R498 combination is more tolerated.[37]

In Region 1 (**Table 3**) we also note the proline 499 to threonine (P499T) substitution in two infectious

clones, presumably engineered to enhance Y498. Modelling provides the following insight on this: the

change from P to T will relax the backbone constraints of proline and allow conformational

rearrangement of threonine to contact the conserved ACE2 residues Y41 and L45. However, we don't

find any strong molecular modelling basis for this mutation to sustain mouse adaptation on its own or

evolve naturally alongside adaptive mutations at positions 498 or 501; this is borne out by experimental

evidence to date. In other words, *in silico* predictions combined with *in vitro* and *in vivo* evidence

collectively suggest that mouse adaptation requires an aromatic substitution in either position 501 or

position 498 (but not both); while additional mutations, especially in Region 2 and Region 3 of the RBD

as summarized below, enhancing ACE2 binding interactions and specificity in mice. These predictions

are further supported by *AlphaFold* that assigns very high confidence scores (>93 in a scale of 0-100) for

the structural predictions involving these key mutations, demonstrating the applications of this

improved protein prediction software especially during pandemics such as COVID-19 and future

'Disease-X'.

From **Table 1**, we see that mouse adapted strains sometimes carry the Q493K/R mutation (polar

glutamine to basic lysine or arginine). Modelling predicts this as being enabled through favourable salt-

bridge interactions with both glutamic acid 35 (E35) or aspartic acid 38 (D38) both of which are

conserved in hACE2 as well as mACE2 (c.f. **Table 3**, Region 2). The K417N substitution (lysine to

asparagine), which is another experimental observation from **Table 1**, is also predicted by modelling to

be advantageous for mouse adaptation due to favourable amide hydrogen bond interactions with

interfacial mACE2 residues asparagine 31 (N31) and glutamine 34 (Q34); such amide hydrogen bonding

is not possible in hACE2 as it has non-amide lysine (K) and histidine (H) residues. With the Gamma

variant of concern, it is unclear whether the K417T enhances the role of N501Y in mouse adaptation in a

similar manner. The 'Delta Plus' variant of concern has the K417N mutation, but there is no molecular

modelling basis to believe that it can infect mice without an aromatic change in position 498 or 501 as

described above. It would be worthwhile to further investigate any interfering role of glycosylation at

this interface region, because hACE2 contains N-linked glycosylation at asparagine 90 (N90) whereas

mACE2 does not (its analogous residue is threonine T90 according to Uniprot references Q8R0I0 and

Q9BYF1).

In Region 3, the K484 residue is not positioned directly at the interface and not observed to interact

strongly with any ACE2 residues; however, our model shows occasional salt bridges can be formed with

relatively close glutamic acid residues in positions 35 and 75 that are conserved in both hACE2 and

mACE2. Our simulations show that the distance between K484 and these glutamic acid residues

fluctuate dynamically from 3 to 20 Å, with salt bridges more likely when distances are around 3 Å. Thus,

the E484K, which is present in the Beta and Gamma variants of concern, and more recently in some

Alpha isolates as well, is likely to have an enhancing role through transient salt bridges. The same cannot

be said about E484Q seen in the Delta variant of concern because salt bridge formation is unlikely with

the polar glutamine (Q) residue. With no accompanying aromatic change in positions 498 or 501, we

believe that the E484Q in Delta, and additionally the K417N in 'Delta Plus', cannot sustain mouse

infectivity on their own based on current biomolecular understanding. The residues 75 to 82 in mACE2

are significantly different from hACE2 (**Table 3**), therefore any mutation in the corresponding RBD

interface is worth investigation. We could find one from experimental observations, the engineered

substitution F486L,[27] and consider it to have at best an enhancing role. As it was observed

simultaneously with Q498Y (which is likely to sustain mouse infectivity on its own), the contribution of

F486L to the overall mouse adaptation remains to be ascertained.

Comparison with *in vitro*, *in vivo* and *in situ* observations

Early *in silico* predictions based on comparative structural analysis of ACE2 suggested that mouse has a

very low probability of being infected.[35,36] Although correct about mouse, those analyses also made

inconsistent and erroneous predictions that ferrets wouldn't be susceptible, pigs would be susceptible,

etc., thus exposing the need for experimental inputs into the model. Therefore, this paper takes into

account a range of experimental observations to cross-check our *in silico* predictions through

biomolecular modelling. Wan *et al.* reasoned that "*mouse or rat ACE2 contains a histidine at the 353*

position which does not fit into the virus-receptor interaction as well as a lysine does". [35] While this is

true of the original Wuhan strain containing asparagine 501 (N501) in the RBD, our modelling indicates

why the tyrosine 501 mutation enables mouse infectivity, even on its own. In hACE2, lysine 353 (K353)

creates a salt bridge with conserved aspartic acid 38 (D38). In mACE2, lysine 353 is replaced by the

aromatic histidine (H353) to complete the salt bridge, as well as contribute to π-stacking with Y501

variant. The N501Y mutation will also lead to favorable π-stacking with the highly conserved tyrosine 41

(Y41) residue in mammalian ACE2, as suggested by Starr *et al.*[38] with deep scanning of RBD mutations

and hACE2 affinity assays. These authors highlighted enhanced affinity of F501 (as it had the highest

score), followed by Y501, V501, W501 and T501, in that order. But Y501 and T501 require only a single

nucleotide change and have been observed more frequently *in situ* (**Figure S1a**) – compared to F501,

V501 and W501 which require 2, 2 and 3 nucleotide changes respectively. It is unsurprising that the

latter variants requiring two or more changes were rarely observed *in situ* regardless of their high *in*

vitro affinity scores from Starr *et al.*[38]

From the above and **Table 4**, we see that *in silico* analysis can provide valuable insights to interpret and

bridge *in vitro*, *in vivo* and *in situ* observations on the RBD position 501. A similar analysis is possible with

the alternative essential mutation for mouse adaptation at RBD position 498, where the *in vitro* affinity

enhancement order is H498, Y498, F498 and W498 according to Starr *et al.*[38] Of these, H498 (on its own)

and Y498 (with enhancing RBD mutations) have been shown to result in mouse adaptation *in vivo* (**Table**

1);[7,8,23-28] Q498R was also reported once, unusually in combination with N501Y, isolated from mouse

lung after 30 passages. In humans, *in situ* observations of these variants have been limited to R498 (57

occurrences) and H498 (8 occurrences) so far. Thus, it is clear from *in vitro*, *in vivo* and *in situ* analyses

(**Table 4**) that H498, R498 and Y498 are possible but not yet common. This is consistent with our *in silico*

predictions because H498 and Y498 are aromatic (enabling π-stacking with ACE2 Y41; similar to Y501),

while R498 has conformational flexibility and can still be associated with π-stacking interactions.[37] H498

and R498 observed *in situ* require a single nucleotide change from Q498, while Y498 requires two

nucleotide changes (or one change from H498).

Similar insights are also possible for the enhancing RBD mutations (c.f. **Table 1** and **Table 4**). From *in vivo*

and *in situ* observations, we see that K417M is less common than K417N or K417T (**Figures S2a** and **S2b**),

although *in vitro* studies did not predict any enhancement.[38] *In silico* predictions show that all these

require a single nucleotide change, but that N417 (and Q417) would benefit from amide hydrogen

bonding. E484K is present in Beta and Gamma (**Figure S2c**), and increasingly in Alpha VOC, while E484Q

is present in the Kappa variant of interest that is related to the Delta VOC. In comparison to these two

substitutions and notwithstanding higher *in vitro* affinity scores, R484 and T484 are infrequently

observed *in situ*, which is consistent with our *in silico* predictions because they each require 2 nucleotide

changes from E484, or one change from K484. The F486L and P499T were originally engineered *in vivo*,

but have had sporadic *in situ* presence in human populations. *In silico* predictions suggest that the F486L

mutation (accessible by three possible ways of a single nucleotide change) can aid mACE2 adaptation,

due to the human-mouse differences in ACE2 at the 78-82 region; the P499T is also a single nucleotide

change (but only one way from P to T) and predicted to be rare in comparison. Finally, *in silico*

predictions for Q493 substituted by K, L or R (each a single nucleotide change) are borne out *in vivo* and

in situ, although their affinity scores from *in vitro* experiments are low. The affinity scores from Starr *et*

al.[38] were developed for hACE2 (not mACE2), so we expected a greater correlation than what has been

observed *in situ* in human populations, but perhaps it is still early in the pandemic to assess this

definitively. It also looks likely that there are factors other than enhancement of ACE2 binding that

determine susceptibility – as seen from Starr *et al.*[38] and **Table 4**; from our own inconclusive attempts at

correlating free energy binding affinities using *in silico* methods (not shown); and from a more detailed

in silico model by Piplani *et al.*[39] which counter-intuitively predicts lower binding free energy for dogs

compared to more susceptible animals such as monkeys, hamsters, ferrets, cats and tigers.[40]

Some mutations in the essential as well as enhancing positions can lead to other mutations. For

example, N501Y, the key mutation common to the Alpha, Beta, and Gamma variants of concern, can

lead to F501 with a further nucleotide change. Similarly, the enhancing E484K mutation can also lead to

R484 or T484 with a further nucleotide change. We examined whether *in situ* observations are

consistent or contrary to our *in silico* predictions. Indeed, F501 was observed in Sweden (28 April 2021),

Germany (7 May 2021), Mexico (22 June 2021) and Colombia (30 June 2021), once in each of these four

countries, while the Y501 has been observed in these countries since 12 March 2020, 21 October 2020,

31 January 2021 and 19 September 2020 respectively. UK reported E484R in August 2020, followed by

Angola in April 2021; Brazil and USA in May 2021; South Korea in June 2021; and South Africa in July

2021. In each case, the E484K was detected prior to E484R – the former mutation circulating in UK,

Angola, Brazil, USA, South Korea and South Africa from April, August, April, March, December and

August 2020 respectively. Similarly, E484T was only recently detected in the USA in June 2021, 15

months after the first report of E484K in that country. All eleven instances are thus consistent with our

prediction – whether this link is causal or a coincidence is worthy of investigation with local

epidemiological data. While bioinformatics tools can provide useful insights, out of 53 COVID-positive

cases only one sample is on average sent for virus genome sequencing (as of 4 October 2021), with huge

variations across time and locations, and lots of missing meta-data.[41,42] This means we are more

confident about ruling in (e.g. when a variant has been detected in a location) than ruling out the

possibility of a mutation circulating purely based on *in silico* data, even if the latter is statistically large (4

million as of 4 October 2021).

Our analysis is not just of theoretical interest; it has huge practical applications because mice can be

kept as pets, or come into contact with other pets like cats which are known to be susceptible. Also,

mouse plague can occur in area of COVID-19 outbreaks or endemicity, as is currently the case in New

South Wales and adjacent states of Australia.[43] In order to help public health and animal health

professionals, Figures 3a and 3b show the eleven countries where key essential and enhancing

mutations listed in Tables 1 and 4 (viz. N501Y, E484K and K417N/T) have co-occurred. Supplementary

Table S1 gives the raw data, down to regional counts for these combinations. We believe that this

information will help locate areas at risk (especially in Brazil, Chile, Djibouti, Haiti, Malawi, Mozambique,

Reunion, Suriname, Trinidad and Tobago, Uruguay and Venezuela) for appropriate mitigation measures.

Conclusion and further analyses

Assessing the risk of viruses adapting to new hosts requires careful interpretation of all available data

from *in silico*, *in vivo*, *in vitro* and *in situ* sources. Understanding host adaptation at a molecular level, via

modelling helps reconcile seemingly conflicting, experimental, and clinical observations while a

pandemic is still in progress. We have demonstrated this with the SARS-CoV-2 virus adapting to mice.

Our conclusions come with humility, as they are based on best available evidence up to this point, but

allowing us and others to refine when more evidence becomes available. Armed with our collective

understanding from different approaches, and bolstered by bioinformatics and emerging artificial

intelligence technologies such as *AlphaFold*, we have shown how to position ourselves better to predict

and mitigate virus host adaptations, not just for this pandemic but also for future Disease-X. Further

analyses pertaining to COVID-19 should focus on the role of mutations beyond the spike and RBD;

improved computational modelling of binding affinities to explore correlations with susceptibility, if any;

and study other hosts like rats and other potential reservoir species (even those that previously

exhibited low receptor activities) which will be hard to vaccinate or control.[35,36,44]

Supplementary Methods for Molecular Modelling

Molecular simulations were performed using NAMD2.14[45] with CHARM36m[46] forcefield employing a

'TIP3' water model. The SARS-CoV-2 spike/ACE2 model was a homology model based on

the pdb structure 6M17.[15] Variant models of the SARS-CoV-2 spike RBD domain (residues 330 to 530)

containing the mouse adapted mutations were constructed by mutating residues in the NAMD build

scripts, but later found to be in very close agreement with the same model constructed using

AlphaFold[16] (less than 0.9 Å RMSD difference to C-Alpha backbone atoms). A truncated mACE2

consisting of residues 19 to 600 was built using Swiss modeller,[47] and similarly found to be in close

agreement with the equivalent *AlphaFold* model. Amino acid side chain conformations predicted by

AlphaFold were used in all initial RBD model starting conformations. Glycosylation of the spike and

mACE2 protein was manually constructed using Visual Molecular Dynamics (VMD). Simulations were

run with Periodic Boundary Conditions 'PBCs' using the 'NPT' isothermal-isobaric ensemble at 310K and

1 bar pressure employing Langevin dynamics. The PBCs were constant in the XY dimensions. Long-range

Coulomb forces were computed with the Particle Mesh Ewald method with a grid spacing of 1 Å. 2 fs

timesteps were used with non-bonded interactions calculated every 2 fs and full electrostatics every 4 fs

while hydrogens were constrained with the 'SHAKE' algorithm. The cut-off distance was 12 Å with a

switching distance of 10 Å and a pair-list distance of 14 Å. Pressure was controlled to 1 atmosphere

using the Nosé-Hoover Langevin piston method employing a piston period of 100 fs and a piston decay

of 50 fs. Trajectory frames were captured every 100 ps. Eleven variant models were constructed

representing the mouse-adapted variants observed in **Table 1** as well as the original Wuhan RBD model

with both mouse and human ACE2. Models were simulated for 300 nanoseconds. Trajectories were

visualized with VMD and Nanome. Modelling data shall be made available on the CSIRO Data access

portal (https://data.csiro.au/).

Acknowledgements

This work was supported by funding (Principal Investigator: S.S.V.) from the Australia's Department of

Finance, CSIRO Future Science Platforms, National Health and Medical Research Council (MRF2009092),

and United States Food and Drug Administration (FDA) Medical Countermeasures Initiative contract

(75F40121C00144). The article reflects the views of the authors and does not represent the views or

policies of the funding agencies including the FDA. We are grateful for support from our colleagues at

the Australian Centre for Disease Preparedness (https://www.grid.ac/institutes/grid.413322.5)

(especially Simran Chahal, Trevor Drew, Alexander McAuley and Nagendrakumar Singanallur) and the

Transformational Bioinformatics Group (especially Denis Bauer, Yatish Jain, Brendan Hosking and Aidan

Tay). The title is from the poem 'To a Mouse: On Turning her up in her Nest, with the Plough, November

1785' by Scotland's national poet Robert Burns, in which he says that the mouse is not alone in proving

foresight may be vain as the best-laid schemes of mice and men go oft awry (But, Mousie, thou art no

thy-lane, In proving foresight may be vain: The best-laid schemes o' Mice an' Men Gang aft agley).

Author contributions

Conceptualization, methodology, and funding acquisition, S.S.V.; in silico analysis, M.J.K.; in vitro

analysis, M.J.K. and S.S.V.; in vivo analysis, S.M. and S.S.V.; in situ analysis, L.O.W.W., D.R. and C.L.;

writing – original draft preparation, S.S.V, M.J.K. and S.M.; writing – review and editing, all authors.

References

1.   Dong E, Du H, Gardern L. An interactive web-based dashboard to track COVID-19 in real time.

Lancet Infectious Diseases. 2020;20(5):533-534. Available at https://doi.org/10.1016/S1473-

3099(20)30120-1 (accessed 1 December 2020).

2.   Bauer DC, Tay AP, Wilson, LOW, et al. Supporting pandemic response using genomics and

bioinformatics: A case study on the emergent SARS-CoV-2 outbreak. *Transboundary & Emerging*

Diseases. 2020; 67(4):1453-1462. Available at https://doi.org/10.1111/tbed.13588 (accessed 1

December 2020).

3.   Lassaunière R, Fonager J, Rasmussen, M, et al. Working paper on SARS-CoV-2 spike mutations

arising in Danish mink, their spread to humans and neutralization data. 2020. Available at

https://files.ssi.dk/Mink-cluster-5-short-report_AFO2 (accessed 1 December 2020).

4.   Korber B, Fischer WM, Gnanakaran S, et al. Tracking changes in SARS-CoV-2 spike: Evidence that

D614G increases infectivity of the COVID-19 virus. *Cell.* 2020; 182(4):812-827. Available at

https://doi.org/10.1016/j.cell.2020.06.043 (accessed 1 December 2020).

5.   Zhang J, Cai Y, Xiao T, et al. Structural impact on SARS-CoV-2 spike protein by D614G

substitution. *Science.* 2021; 372(6541):525-530. Available at

https://doi.org/10.1126/science.abf2303 (accessed 1 May 2021).

6.   Gu H, Chen Q, Yang G, et al. Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine

efficacy. *Science.* 2020; 369(6511):1603-1607. Available at

https://doi.org/10.1126/science.abc4730 and

https://www.biorxiv.org/content/10.1101/2020.05.02.073411v1 (accessed 1 December 2020).

7.   Dinnon KH, Leist SR, Schäfer A, et al. A mouse-adapted model of SARS-CoV-2 to test COVID-19

countermeasures. *Nature.* 2020; 586:560-566. Available at https://doi.org/10.1038/s41586-020-

2708-8 (accessed 1 December 2020).

8.   Leist SR, Dinnon KH, Schäfer A, et al. A mouse-adapted SARS-CoV-2 induces acute lung injury

370   and mortality in standard laboratory mice. *Cell*. 2020; 183:1070-1085. Available at

371   https://doi.org/10.1016/j.cell.2020.09.050 (accessed 1 December 2020).

372   9.   Yuan L, Tang Q, Cheng T, Xia N. Animal models for emerging coronavirus: progress and new

373   insights. *Emerging Microbes & Infections*. 2020; 9(1):949-961. Available at

374   https://doi.org/10.1080/22221751.2020.1764871 (accessed 15 May 2021).

375   10.   Muñoz-Fontela C, Dowling WE, Funnell SGP, et al. Animal models for COVID-19. *Nature*. 2020;

376   586: 509-515. Available at https://doi.org/10.1038/s41586-020-2787-6 (accessed 1 December

377   2020).

378   11.   Zeiss CJ, Compton S, Veenhuis, RT. Animal models of COVID-19. I. Comparative virology and

379   disease pathogenesis. *ILAR Journal*. 2021; ilab007. Available at

380   https://doi.org/10.1093/ilar/ilab007 (accessed 1 May 2021).

381   12.   Veenhuis RT, Zeiss CJ. Animal models of COVID-19 II. Comparative immunology. *ILAR Journal*.

382   2021; ilab010. Available at https://doi.org/10.1093/ilar/ilab010 (accessed 1 May 2021).

383   13.   Rathnasinghe R, Jangra S, Cupic A, et al. The N501Y mutation in SARS-CoV-2 spike leads to

384   morbidity in obese and aged mice and is neutralized by convalescent and post-vaccination

385   human sera. *medRxiv*. 2021; 2021.01.19.21249592. Available at

386   https://doi.org/10.1101/2021.01.19.21249592 (accessed 1 February 2021).

387   14.   Yao W, Wang Y, Ma D, et al. Circulating SARS-CoV-2 variants B.1.1.7, 501Y.V2, and P.1 have

388   gained ability to utilize rat and mouse Ace2 and altered in vitro sensitivity to neutralizing

389   antibodies and ACE2-Ig. *bioRxiv*. 2021; 2021.01.27.428353. Available at

390   https://doi.org/10.1101/2021.01.27.428353 (accessed 1 May 2021).

15.   Yan R, Zhang Y, Li Y, et al. Structural basis for the recognition of SARS-CoV-2 by full-length

human ACE2. *Science*. 2020; 367:1444-1448. Available at

https://doi.org/10.1126/science.abb2762 (accessed 1 December 2020).

16.   Jumper, J., Evans, R., Pritzel, A. *et al.* Highly accurate protein structure prediction with

AlphaFold. Nature (2021). Available at https://doi.org/10.1038/s41586-021-03819-2 (accessed

20 July 2021).

17.   McAuley AJ, Kuiper MJ, Durr PA, et al. Experimental and *in silico* evidence suggests vaccines are

unlikely to be affected by D614G mutation in SARS-CoV-2 spike protein. *Npj Vaccines*. 2020;

5:96. Available at https://doi.org/10.1038/s41541-020-00246-8 (accessed 1 December 2020).

18.   Riddell S, Goldie S, McAuley A, et al. Live virus neutralisation of the 501Y.V1 and 501Y.V2 SARS-

CoV-2 variants following INO-4800 vaccination of ferrets. *bioRxiv*. 2021; 2021.04.17.440246.

Available at https://doi.org/10.1101/2021.04.17.440246 (accessed 1 May 2021).

19.   Tai W, He L, Zhang X. Characterization of the receptor-binding domain (RBD) of 2019 novel

coronavirus: implication for development of RBD protein as a viral attachment inhibitor and

vaccine. Cellular & Molecular Immunology. 2020; 17:613-620. Available at

https://doi.org/10.1038/s41423-020-0400-4 (accessed 1 December 2020).

20.   Grantham, R. Amino acid difference formula to help explain protein evolution. *Science*. 1974;

185(4154): 862-864. Available at https://doi.org/10.1126/science.185.4154.862 (accessed 1

December 2020).

21.   Sun S, Gu H, Cao L, et al. Characterization and structural basis of a lethal mouse-adapted SARS-

CoV-2. bioRxiv. 2020; 2020.11.10.377333. Available at

https://doi.org/10.1101/2020.11.10.377333 (accessed 1 December 2020).

22.   Fagre A, Lewis J, Eckley M, et al. SARS-CoV-2 infection, neuropathogenesis and transmission

among deer mice: Implications for spillback to New World rodents. *PloS Pathogens*. 2021; 17(5):

e1009585. Available at https://doi.org/10.1371/journal.ppat.1009585 (accessed 24 May 2021).

23.   Wang J, Shuai L, Wang C, et al. Mouse-adapted SARS-CoV-2 replicates efficiently in the upper

and lower respiratory tract of BALB/c and C57BL/6J mice. *Protein & Cell*. 2020; 11:776-782.

Available at https://doi.org/10.1007/s13238-020-00767-x (accessed 1 December 2020).

24.   Liu Z, Zheng H, Lin H, et al. Identification of common deletions in the spike protein of Severe

Acute Respiratory Syndrome Coronavirus 2. *Journal of Virology*. 2020; 94(17):e00790-20.

Available at https://doi.org/10.1128/JVI.00790-20 (accessed 1 December 2020).

25.   Zhou H, Chen X, Hu T, et al. A novel bat coronavirus closely related to SARS-CoV-2 contains

natural insertions at the S1/S2 cleavage site of the spike protein. *Current Biology*. 2020;

30:2196-2203. Available at https://doi.org/10.1016/j.cub.2020.05.023 (accessed 1 December

2020).

26.   Huang K, Zhang Y, Hui X, et al. Q493K and Q498H substitutions in spike promote adaptation of

SARS-CoV-2 in mice. *EbioMedicine*. 2021; 67:103381. Available at

https://doi.org/10.1016/j.ebiom.2021.103381 (accessed 24 May 2021).

27.   Muruato A, Vu MN, Johnson BA, et al. Mouse adapted SARS-CoV-2 protects animals from lethal

SARS-CoV challenge. *bioRxiv*. 2021; 2021.05.03.442357. Available at

https://doi.org/10.1101/2021.05.03.442357 (accessed 15 May 2021).

28.   Zhang Y, Huang K, Wang T, et al. SARS-CoV-2 rapidly adapts in aged BALB/c mice and induces

typical pneumonia. *Journal of Virology*. 2021; 95(11):e02477-20. Available at

https://doi.org/10.1128/JVI.02477-20 (accessed 24 May 2021).

29.   Van Noorden, R. Scientists call for fully open sharing of coronavirus genome data. *Nature*. 2021;

590:195-196. Available at https://doi.org/10.1038/d41586-021-00305-7 (accessed 1 March

2021).

30.   Rambaut A, Loman N, Pybus O, et al. Preliminary genomic characterisation of an emergent

SARS-CoV-2 lineage in the UK defined by a novel set of spike mutations. *Virological*. 2020.

Available at https://virological.org/t/preliminary-genomic-characterisation-of-an-emergent-sars-

cov-2-lineage-in-the-uk-defined-by-a-novel-set-of-spike-mutations/563 (accessed 1 March

2021).

31.   Li Q, Nie J, Wu J, et al. SARS-CoV-2 501Y.V2 variants lack higher infectivity but do have immune

escape. *Cell*. 2021; 184(9):2362-2371. Available at https://doi.org/10.1016/j.cell.2021.02.042

(accessed 1 May 2021).

32.   Montagutelli X, Prot M, Levillayer L, et al. The B1.351 and P.1 variants extend SARS-CoV-2 host

range to mice. *bioRxiv*. 2021; 2021.03.18.436013. Available at

https://doi.org/10.1101/2021.03.18.436013 (accessed 1 May 2021).

33.   Faria NR, Claro IM, Candido D, et al. Genomic characterisation of an emergent SARS-CoV-2

lineage in Manaus: preliminary findings. *Virological*. 2021. Available at

https://virological.org/t/genomic-characterisation-of-an-emergent-sars-cov-2-lineage-in-

manaus-preliminary-findings/586 (accessed 1 March 2021).

34.   Lucaci AG, Zehr JD, Shank SD, et al. RASCL: Rapid assessment of SARS-COV-2 clades enabled

through molecular sequence analysis and its application to B.1.617.1 and B.1.617.2. *Virological*.

2021. Available at https://virological.org/t/rascl-rapid-assessment-of-sars-cov-2-clades-enabled-

through-molecular-sequence-analysis-and-its-application-to-b-1-617-1-and-b-1-617-2/709

(accessed 29 May 2021).

35.   Wan Y, Shang J, Graham R, et al. Receptor recognition by novel coronavirus from Wuhan: an

analysis based on decade-long structural studies of SARS coronavirus. *Journal of Virology.* 2020;

94(7):e00127-20. Available at https://doi.org/10.1128/JVI.00127-20 (accessed on 1 December

2020).

36.   Damas J, Hughes GM, Keough C, et al. Broad host range of SARS-CoV-2 predicted by comparative

and structural analysis of ACE2 in vertebrates. *PNAS.* 2020; 117(36):22311-22322. Available at

https://doi.org/10.1073/pnas.2010146117 (accessed 1 December 2020).

37.   Flocco MM, Mowbray SL. Planar stacking interactions of arginine and aromatic side-chains in

proteins. *Journal of Molecular Biology*. 1994; 235(2):709-717. Available at

https://doi.org/10.1006/jmbi.1994.1022 (accessed 1 December 2020).

38.   Starr TN, Greaney AJ, Hilton SK, et al. Deep mutational scanning of SARS-CoV-2 receptor binding

domain reveals constraints on folding and ACE2 binding. *Cell.* 2020; 182(5):1295-1310.e20.

Available at https://doi.org/10.1016/j.cell.2020.08.012 (accessed 1 December 2020).

39.   Piplani S, Singh PK, Winkler DA, Petrovsky, N. *In silico* comparison of SARS-CoV-2 spike protein-

ACE2 binding affinities across species and implications for virus origin. *Scientific Reports.* 2021;

11:13063. Available at https://doi.org/10.1038/s41598-021-92388-5 (accessed 1 October 2021).

40.   Muñoz-Fontela C, Dowling, WE, Funnell, SGP, et al. Animal models for COVID-19. *Nature.* 2020;

586:509-515. Available at https://doi.org/10.1038/s41586-020-2787-6 (accessed 1 December

2020).

41.   Bauer DC, Metke-Jimenez A, Maurer-Stroh S, et al. Interoperable medical data: the missing link

for understanding COVID-19. *Transboundary & Emerging Diseases*. 2021; 68(4):1753-1760.

Available at https://onlinelibrary.wiley.com/doi/10.1111/tbed.13892 (accessed 1 July 2021).

42.   Priyadarshini, S. Massive coronavirus sequencing efforts urgently need patient data. *Nature*

India (special issue #13 on COVID-19 crisis). 2020; 10.1038/nindia.2020.75:11-13. Available at

https://www.natureasia.com/en/nindia/pdf/special-issues/13/Nature-India-COVID-19-Crisis.pdf

and https://go.nature.com/2y7kUIw (accessed 1 December 2020).

43.   NSW Government. Help for regional communities impacted by the mouse plague. 2021.

Available at https://www.nsw.gov.au/initiative/mouse-control-support-program (accessed 24

May 2021).

44.   Zhao X, Chen D, Szabla R, et al. Broad and differential animal angiotensin-converting enzyme 2

receptor usage by SARS-CoV-2. *Journal of Virology*. 2020; 94(18):e00940-20. Available at

https://doi.org/10.1128/JVI.00940-20 (accessed 1 December 2020).

45.   Phillips JC, Braun R, Wang W, et al. Scalable molecular dynamics with NAMD. *Journal of*

Computational Chemistry. 2005; 26:1781-1802. Available at https://doi.org/10.1002/jcc.20289

(accessed 1 December 2020).

46.   Huang J, Rauscher S, Nawrocki G, et al. CHARMM36m: an improved force field for folded and

intrinsically disordered proteins. *Nature Methods.* 2017; 14:71-73. Available at:

https://doi:10.1038/nmeth.4067 (accessed 1 December 2020).

47.   Waterhouse A, Bertoni M, Bienert S, et al. SWISS-MODEL: homology modelling of protein

structures and complexes. *Nucleic Acids Research*. 2018; 46:W296-W303. Available at

https://doi.org/10.1093/nar/gky427 (accessed 1 December 2020).

Mice-infecting SARS-CoV-2 'strains'/isolates	Key mutations in the spike RBD (residues 331-524)[19]							Color legend – mutations scored on Grantham Distance,[20] in the scale of 5 (conservative) to 215 (radical), similar to Riddell et al.[18]											
	K417	E484	F486	Q493	Q498	P499	N501	22	24	38	43	53	56	73	94	95	99	143		
Mouse Adapted (MA) strains								**Authors' synthesis/comments**												
MASCp6[6]							N→Y	L37F (nsp6), P84S (nsp10), and D128Y (N) were the other observed nonsynonymous mutations.												
MASCp25[21]				Q→K			N→Y	MASCp6 passaged serially to develop MASCp25 and MASCp36, which accumulated Q493K and subsequently K417N in the RBD.												
MASCp36[21]	K→N			Q→K			N→Y	Highly virulent in aged mice and reported stable/mature RBD binding. It also had I1258V (nsp3), S301L (nsp5), and R32C (N).												
IC-MA1[7]					Q→Y	P→T		Spike Q498Y and P499T were engineered for compatibility with Q42 of mouse ACE2.												
IC-MA10[8]				Q→K	Q→Y	P→T		Q493K evolved in MA10 and interacts with mACE2's N31; others were T285I (nsp4), K2R (nsp7), E23G (nsp8), and F7S (ORF6).												
WA1-MA-P11[13]							N→Y	In addition to N501Y, there were H655Y, T7I (M), L84S (ORF8), S194T (N) and the +KLRS (216-219) insertion also observed by others and predicted to form a solvent-accessible loop.[22]												
HRB26M[23] (HRB26-P14)					Q→H			HRB26-P5 had Q498H, N969S, and the 675QTQTN679 deletion observed by others.[24,25] P14 also had A81T (nsp8).												
Hu-1-WBP-1[26]				Q→K	Q→H			Q493K and Q498H by P5, interacting respectively with mACE2's N31 via H-bond and Y41 via π-stacking. Nonsynonymous Q493K and T77I (nsp9) mutations increased frequency from P8 to P11.												
IC-MA4[27]			F→L		Q→Y			F486L and Q498Y were engineered for infecting mice.												
IC-CMA1-3[27]	K→N						N→Y	CMA1-3 all had N501Y. CMA2 also had D128Y; CMA3 also had D128Y, L37F, and P87S. Passaging of CMA3 resulted in K417N, H655Y, and E8V (E).												
LG[28]					Q→H			Other mutations present were N74K, H655Y, I6V (ORF10).												
IA-N501Y-MA30	K→M	E→K		Q→R	Q→R		N→Y	Homogenate from BALB/c mouse lung after 30 passages. GISAID entry EPI_ISL_1666328.[29]												
Variants of Concern (VOC) infecting mice																				
Alpha (501Y.V1 or B.1.1.7[14,30])		E→K					N→Y	First reported in the UK, September 2020. Spike also has A570D, P681H, T716I, S982A, and D1118H, in addition to HV69-70 and Y144 deletions. **E484K has been detected in this VOC in 15 countries as of 29 April 2021 (436 entries on GISAID).**												
Beta (501Y.V2 or B.1.351[14,31,32])	K→N	E→K					N→Y	First reported in South Africa, May 2020. V2-1 had D80A, D215G, E484K, N501Y, D614G and A701V in the Spike. L18F and K417N arose in V2-2. V2-3 had 242-244 deletion.												
Gamma (501Y.V3 or P.1[14,32,33])	K→T	E→K					N→Y	First reported in Brazil, November 2020. L18F, T20N, P26S, D138Y, R190S, K417T, E484K, N501Y, H655Y and T1027I in Spike.												
VOC not known to infect mice																				
Delta (452R.V3 or B.1.617.2[34])	K→N							First reported in India, October 2020. Also has L452R in the RBD; G142D, R158G, T478K, P681R and D950N in Spike, along with deletions 156-157. **K417N has been detected in this VOC (called 'Delta Plus') in 12 countries as of 25 June 2021.** The related Kappa variant of interest (B.1.617.1) has L452R and E484Q in RBD (Grantham Distance scores of 102 and 29 respectively).												

Table 1 Key mutations in 14 'mouse adapted strains' and 3 'variants of concern' of SARS-CoV-2 known to infect mice, with further characterization as outlined in **Table 2**.

Species and age of mice (Study reference)	Strain, dose, shedding and transmission*	Evidence of productive SARS-CoV-2 replication in the respiratory tract	Clinical outcomes, histopathological evidence of disease
Aged (9 months old) and young (6 weeks old) BALB/c mice studied. (Gu et al, 2020)[6]	**Strain name:** MASCp6 **Dose:** Intranasal inoculation with 1.6×10^4 PFU **Shedding:** Viral shedding observed in the feces at day 3,5 and 7 after inoculation. **Transmission:** No information	High amounts of viral RNAs in the lungs and tracheas were detected at 3, 5 and 7 days after inoculation in all aged mice, with peak viral RNA loads of ~10^{10} copies/g at 3 days after inoculation. Viral RNAs were also detected in heart, liver, spleen, and brain. Similar tissue distribution of SARS-CoV-2 RNA was also seen in the MASCp6 infected young mice.	**Clinical:** No visible clinical symptoms or body weight loss were observed in both the young and aged mice. No fatalities recorded. **Histopathical:** Led to interstitial pneumonia and inflammatory responses in both young and aged mice, however the lung damage in the aged mice was more severe.
Young (8-week-old) and aged (9 months old) male and female BALB/c mice and C57BL/6 mice studied. (Sun et al, 2021)[21]	**Strain name:** MASCp36 - developed from MASCp6 (Gu et al, 2020)[6] **Dose:** Intranasal inoculation with varying doses including: 1.2 PFU, 12 PFU, 120 PFU, 1200 PFU, 12000 PFU **Shedding:** No information **Transmission:** No information	qRT-PCR results of aged BALB/c and C57BL/6 mice challenged with 12,000 PFU of MASCp36 showed that high levels of SARS-CoV-2 subgenomic RNAs were persistent in the lung and tracheas till 4 dpi. A similar tissue distribution of SARS-CoV-2 was observed in the young BALB/c and C57BL/6 mice.	**Clinical:** All the aged BALB/c and C57BL/6 mice challenged with high doses (1200 or 12,000 PFU) of MASCp36 developed typical respiratory symptoms and exhibited features like ruffled fur, hunched back, and reduced activity. Young BALB/c and C57BL/6 mice were resistant to MASCp36 challenge, and only one animal that received 12,000 PFU challenge died during the observation period. **Histopathological:** Necrotizing pneumonia and extensive diffuse alveolar damages observed on 4 dpi in aged mice. Milder pathology observed in the young mice.
Young adult (12 weeks old) and aged (One year old) BALB/c mice studied. (Dinnon et al, 2020)[7]	**Strain name:** IC-MA1 **Dose:** Intranasal inoculation with 10^5 PFU **Shedding:** No information **Transmission:** No information	High-titre virus replication was also observed in lung tissue at 2 dpi. This cleared by 4 dpi in the young adult mice but continued to persist at 4 dpi in the aged mice, suggesting that there was increased viral replication in the aged mice. MA1 was also observed in the upper airway and viral antigen was present in nasal turbinate epithelium at 2 dpi in young adult mice; viral antigen was found in conducting airway epithelium, interstitium and nasal epithelium in aged mice at 4 dpi.	**Clinical:** No overt clinical signs such as weight loss observed in young adult mice but a significant decrease in body weight was observed in aged mice. Whole-body plethysmography on mice indicated impaired lung function at 2 dpi, with the extent of impairment significantly worse in the aged mice. **Histopathological:** Interstitial congestion, epithelial damage, inflammatory infiltrate and peribronchial lymphocytic inflammation surrounding airways at 2 dpi. Histopathological effects seen in the aged mice are more severe than those observed in young adult mice.
Young (10-week-old) and aged (1 year old) BALB/c mice studied. (Leist et al, 2020)[8]	**Strain name:** IC-MA10 **Dose:** Doses ranging from 10^2, 10^3, 10^4 or 10^5 PFU were tested. However, 10^4 PFU and 10^3 PFU were determined as the optimal doses for analysis in young and aged mice respectively. **Shedding:** Both old and young mice exhibited viral titers in the nasal cavity over the first 3 days of infection. **Transmission:** No information	Virus replication in the lungs of young mice peaked 1–2 dpi and was absent in most surviving mice by 7 dpi. Viral replication in the upper respiratory tract of young mice remained high on 1–3 dpi but was undetectable in most mice by 5 dpi. A similar trend was observed in the aged mice, but the viral replication was detectable for a much longer period.	**Clinical:** Young mice rapidly lost weight and reached maximum weight loss at day 4 post inoculation. On day 5, the collective mortality rate was approximately 15%. Significant weight loss was also observed in aged mice. Mortality rates of 20% and 60% were recorded for infection with 10^4 and 10^5 PFU, respectively in the young mice. **Histopathological:** At the time of necropsy, acute stage lung damage was noted in the young mice. Analysis at 2, 4, and 7 dpi revealed early multifocal damage to conducting airway epithelia and hallmarks of diffuse alveolar damage (DAD) was observed. The aged mice had severe DAD and higher acute lung injury scores. Overall, a dose- and age-related increase in pathogenesis was observed.
C57Bl6, BALB/c and 129S1/SVMJ mice were studied Young (6-8-weeks) or aged (52 weeks old) C57Bl6 mice were used to study the effect of obesity, obesity-associated diabetes, and advanced age on clinical outcomes. (Rathnasinghe et al, 2021)[13]	**Strain name:** WA1-MA-P11 **Dose:** Intranasal inoculation with 2.5×10^4 PFU **Shedding:** No information **Transmission:** No information	Detectable virus titres in lungs and nasal turbinates of 129S1/SVMJ mice and lungs of male and female C57Bl6 and BALB/c mice, obtained at different days post infection	**Clinical:** Weight loss detected in the 129S1/SVMJ mice but not in the C57Bl6 and BALB/c mice. **Effect of increased age, comorbidities:** Both obesity/diabetes and advanced age in mice result in higher morbidity during SARS-CoV-2 infection.
Groups of young (4–6-week-old) female BALB/c mice, young (4–6-week-old female) C57BL/6J mice and aging (8–9-month-old) male BALB/c mice studied. (Wang et al, 2020)[23]	**Strain name:** HRB26M **Dose:** Intranasal inoculation with $10^{4.4}$ PFU in a volume of 50 µL. **Shedding:** No information **Transmission:** No direct evidence of transmission provided. However, the presence of efficient viral replication in the upper and lower respiratory tracts indicates that transmission could be possible.	In young BALB/c mice, viral RNA was detected in the nasal turbinates on day 3, 5, and 7 p.i. (post infection) and the infectious virus was detected on day 3 and 5 p.i. The viral RNA was also detected in the heart, liver, kidney, and spleen on day 3 p.i. In C57BL/6J mice, high viral loads in the nasal turbinates and lungs on days 3, 5 and 7 p.i. Similar observations made in the aging mice. However, when compared to the young mice, the average PFU titres in the aging mice were 3 times higher and 10 times higher in the nasal turbinates and lungs respectively.	**Clinical:** Similar to the young mice, the aging adult mice showed transient weight loss on days 2 and 3 p.i. and recovered thereafter. **Histopathical:** Mild pathological changes were observed in the respiratory tract of young BALB/c mice infected. Moderate-to-severe pathological changes in the lungs after infection in the aging mice.
Young (4–6-week-old) BALB/c mice studied. [Female 12-month-old BALB/c mice only used for the initial generation of the mouse adapted strain, not for any analysis of the mouse adapted strain]. (Huang et al, 2021)[26]	**Strain name:** Hu-1-WBP-1 **Dose:** Intranasally infected with 10-fold serial dilutions of the WBP-1 virus (i.e., 10^2 to 10^5 PFU). Subsequently, detailed analysis on mice was done by infecting mice with 50 µL of 2 LD_{50} [The LD_{50} of WBP-1 is $10^{3.8}$ PFU]. **Shedding:** No information **Transmission:** No information	In mice given the 50 µL of 2 LD_{50}: High viral loads were observed in the trachea and lung of infected mice, with lung tissues having the highest number of viral RNA copies. Trace amounts of viral nucleic acid were detectable in turbinate, heart, and spleen.	**Clinical:** In the groups of mice infected with the 10-fold serial dilutions: All groups of infected mice began to lose weight at 2 dpi and the weight loss was dose dependent. Bristled fur and depressed spirits in mice was also observed. The group of mice receiving a dose of 10^5 PFU WBP-1 virus all died (5/5) by 5 dpi. Two out of five mice survived in the group administered 10^4 PFU virus. In the mice given the 50 µL of 2 LD_{50}: Clinical signs of infection were observed in infected mice at 2 dpi. **Histopathological:** In mice given the 50 µL of 2 LD_{50}, Acute stage lung damage was noted. Moderate interstitial pneumonia with thickened alveolar septa occurred at 3 dpi, which progressed to severe interstitial pneumonia on 5 dpi. Inflammatory cell infiltration was observed in the lung tissue.
10-week-old female BALB/c mice. (Muruato et al, 2021)[27]	**Strain name:** IC-MA4 **Dose:** Mice infected with 10^5 PFU **Shedding:** No information **Transmission:** No information	Evidence of robust viral replication in mouse lungs.	**Clinical:** No information **Histopathological:** No information
10-week-old female BALB/c mice. (Muruato et al, 2021)[27]	**Strain name:** IC-CMA1, IC-CMA2, IC-CMA3 **Dose:** Mice intranasally inoculated with 10^5 PFU **Strain name:** IC-CMA3p20 [IC-CMA3 after passage 20] **Strain name:** IC-CMA3p20 [IC-CMA3 after passage 20] **Dose:** 10^5 PFU was the main dose administered, but doses 10^4 and 10^6 PFU also administered to some mice.	All three mutants (CMA1, CMA2, CMA3) produced approximately, 10^5 PFU per lobe at day 2 post infection. However, no virus was detected at day 4, suggesting rapid clearance by the host. In mice given IC-CMA3p20, robust viral replication was observed. There was a significantly greater viral load in the lung compared to IC-CMA3p20 at 2 dpi.	**Clinical:** None of the three mutants (CMA1, CMA2, CMA3) induced major disease, although both CMA2 and CMA3 caused more weight loss than CMA1. Mice infected with IC-CMA3p20 had a 10% weight loss by day 3 and showed signs of disease including ruffled fur and hunched posture. **Histopathological:** Mice infected with CMA3p20 had significant lung infiltration and inflammation characterized by peribronchiolar, perivascular cuffing, and perivasculitis by day 2 post infection. At 4 dpi, collapsed airways and interstitial pneumonia was observed in the mice.
Young (2 months old) and aged (12-month-old) BALB/c mice (Zhang et al, 2021)[28]	**Strain name:** LG **Dose:** 1×10^5 $TCID_{50}$ in a volume of 50 µl. **Shedding:** No transmission **Transmission:** No information	Effective viral replication in the lungs and trachea of the young and aged mice, although there was more viral replication in the aged mice.	**Clinical:** Weight of the aged mice was slightly decreased, but there were no evident changes in the weight of young mice. **Histopathological:** Lung lesions of the aged BALB/c mice were obvious, but those of young mice were not, with only slight lymphocyte exudation.

508 **Table 2.** Description of the characteristics of the mouse adapted strains generated from the *in vivo*
509 studies discussed in **Table 1**. *Strain = Mouse adapted strain name; Dose = Dose of SARS-CoV-2 mouse
510 adapted strain provided to the mice stated in plaque forming units (PFU) or the tissue culture infective
511 Dose (TCID50); Shedding & Transmission = Any evidence of viral shedding and viral transmission
512 observed in the mice.

	SARS-CoV-2 RBD mouse adaptations	human/mouse ACE2 interface residues within 5 Å of mutation
Region 1	**N501 Y**	38D/D, 41Y/Y, **353K/H**, 355D/D, 357R/R
	P499 T	41Y/Y, 45L/L, **329E/A**, 330N/N
	Q498 H/R/Y	41Y/Y, 42Q/Q, 45L/L, 49N/N, **61N/K**, 357R/R
Region 2	**Q493 K/R**	**34H/Q**, 35E/E, 38D/D, 42Q/Q
	K417 M/N/T	**31K/N**, **34H/Q**
Region 3	**E484 K/Q**	35E/E, 75E/E
	F486 L	75E/E, 76Q/Q, **78 T/K**, **79 L/T**, **82 M/S**

518 **Table 3.** SARS-CoV-2 spike RBD mutations associated with mouse adaptation and their close contact
519 residues in respective ACE2 proteins. Positions with dissimilar human/mouse residues are highlighted in
520 yellow.

	Essential position in RBD		Enhancing position(s) in RBD for mouse adaptation				
Analysis	Q498	N501	K417	E484	F486	Q493	P499
In silico [1]	H Y F W [5]	Y H F W [5]	N Q [8]	K Q	L	K L R	T
In vitro [2]	H Y F W	F Y V W T	-	R K T Q	-	M A Y F K L V	-
In vivo [3]	H R Y	Y	M N T	K Q	L	K R	T
In situ [4]	R H [6]	Y T H F [7]	N T M [9]	K Q R T [10]	L [11]	K L R [12]	T [13]

1. Predicted by our modelling. For E484, F486, Q493, and P499 our work only reconfirmed *in vivo* observations.

2. Arranged in descending order of affinity binding from *in vitro* experiments reported by Starr *et al.*[38]

3. Observed *in vivo* with studies summarized in Table 1 (alphabetical order). Note L486 and P499 were engineered.

4. Observed on GISAID as of 21 July 2021 after removing noise from low sample countries. Only key trends are presented.

5. These aromatic residues are predicted to have similar interactions with Y41 of ACE2.

6. F W Y not observed yet. Since early 2021, R observed in several countries, and H in Slovenia and USA (albeit the latter are of low quality and/or low coverage).

7. V W not observed yet. H observed in several countries (mid 2020), T in Spain's Canary Islands (early 2021), and F in Colombia, Germany, Mexico and Sweden (mid 2021). Y observed in several countries (**Figure S1**).

8. Modelled mutations predict favorable amide hydrogen bonding. N and Q mutations each require a single nucleotide change but there are 2 ways to get from N to K.

9. Q observed in the USA (August 2021). N (mutation G22813T) and T observed several countries (**Figures S2a** and **S2b**). Interestingly, N (alternative mutation G22813C) has a sporadic presence in UK and USA (since late 2020).

10. K observed in several countries (**Figure S2c**). Sporadic presence of Q (several countries from March 2020 to August 2021), R (Angola, Brazil, South Africa, South Korea, UK, USA, since late 2020), and T (USA in mid 2021) observed.

11. Sporadic presence of different synonymous mutations (T23018C in several countries; T23020G in

USA and one instance in Turkey) observed since late 2020.
12. A, F, M, V, Y not observed yet. Sporadic presence observed for K (Italy since early 2020), L (Trinidad and Tobago in mid 2021), and R (several countries mid 2020).
13. Sporadic presence since late 2020 in Bulgaria, Canada, Netherlands and USA.

525 **Table 4.** Comparison of *in silico*, *in vitro*, *in vivo* and *in situ* observations of key mutations in **Table 1**.

Figure 1. The RBD/ACE2 interface. The receptor binding domain (RBD) is shown in cyan, while the ACE2

is shown in yellow. Pink spheres indicate relative positions of mouse adapting mutations on the RBD,

while the blue spheres represent interface residues that differ between human and mouse ACE2

sequences, shown in green and orange respectively.

```
SP|Q9BYF1|ACE2_HUMAN  MSSSSWLLLS  LVAVTAAQST  IEEQAKTFLD  KFNHEAEDLF  YQSSLASWNY  NTNITEENVQ  60
SP|Q8R0I0|ACE2_MOUSE  MSSSSWLLLS  LVAVTTAQSL  TEENAKTFLN  NFNQEAEDLS  YQSSLASWNY  NTNITEENAQ  60
                      **********  *****:****    **:*****:  :**:*****   *********  ********.*

SP|Q9BYF1|ACE2_HUMAN  NMNNAGDKWS  AFLKEQSTLA  QMYPLQEIQN  LTVKLQLQAL  QQNGSSVLSE  DKSKRLNTIL  120
SP|Q8R0I0|ACE2_MOUSE  KMSEAAAKWS  AFYEEQSKTA  QSFSLQEIQT  PIIKRQLQAL  QQSGSSALSA  DKNKQLNTIL  120
                      :*.:*. ***  ** :***. *  * : *****.    :* ***** **.***.**  **.*:*****

SP|Q9BYF1|ACE2_HUMAN  NTMSTIYSTG  KVCNPDNPQE  CLLLEPGLNE  IMANSLDYNE  RLWAWESWRS  EVGKQLRPLY  180
SP|Q8R0I0|ACE2_MOUSE  NTMSTIYSTG  KVCNPKNPQE  CLLLEPGLDE  IMATSTDYNS  RLWAWEGWRA  EVGKQLRPLY  180
                      **********  *****.****  ********:*  ***.* ***.  ******.**:  **********

SP|Q9BYF1|ACE2_HUMAN  EEYVVLKNEM  ARANHYEDYG  DYWRGDYEVN  GVDGYDYSRG  QLIEDVEHTF  EEIKPLYEHL  240
SP|Q8R0I0|ACE2_MOUSE  EEYVVLKNEM  ARANNYNDYG  DYWRGDYEAE  GADGYNYNRN  QLIEDVERTF  AEIKPLYEHL  240
                      **********  ****:*:***  ********.:  *.***:*.*.  *******:**   *********

SP|Q9BYF1|ACE2_HUMAN  HAYVRAKLMN  AYPSYISPIG  CLPAHLLGDM  WGRFWTNLYS  LTVPFGQKPN  IDVTDAMVDQ  300
SP|Q8R0I0|ACE2_MOUSE  HAYVRRKLMD  TYPSYISPTG  CLPAHLLGDM  WGRFWTNLYP  LTVPFAQKPN  IDVTDAMMNQ  300
                      ***** ***:  :******* *  **********  *********  *****.****  ********:.*
                                                                   |329                      |353
SP|Q9BYF1|ACE2_HUMAN  AWDAQRIFKE  AEKFFVSVGL  PNMTQGFWEN  SMLTDPGNVQ  KAVCHPTAWD  LGKGDFRILM  360
SP|Q8R0I0|ACE2_MOUSE  GWDAERIFQE  AEKFFVSVGL  PHMTQGFWAN  SMLTEPADGR  KVVCHPTAWD  LGHGDFRIKM  360
                      .***:***:*  **********  *:******. *  ****:*.: :  *.********  **:***** *
                                                       |387
SP|Q9BYF1|ACE2_HUMAN  CTKVTMDDFL  TAHHEMGHIQ  YDMAYAAQPF  LLRNGANEGF  HEAVGEIMSL  SAATPKHLKS  420
SP|Q8R0I0|ACE2_MOUSE  CTKVTMDNFL  TAHHEMGHIQ  YDMAYARQPF  LLRNGANEGF  HEAVGEIMSL  SAATPKHLKS  420
                      *******:**  **********  ******.***  **********  **********  **********
```

Figure 2   Sequence alignment of human and mouse ACE2 highlighting contact points with the SARS-CoV-2 spike receptor binding domain in yellow. Differences between contact points are highlighted in green (human) and orange (mouse), while common contact residues are highlighted in cyan.

-

a)

b)

Figure 3. Significant occurrences on GISAID of **a)** N501Y, E484K and K417N (nucleotide change G22813T)

triple mutations and **b)** N501Y, E484K and K417T triple mutations, at the virus Receptor Binding Domain

(RBD) since the start of the COVID-19 pandemic (31 December 2019). These countries are encouraged to

perform targeted field surveillance.

Figure S1. Significant occurrences on GISAID of the essential N501Y mutation (nucleotide change

A23063T), at the virus Receptor Binding Domain (RBD) since the start of the COVID-19 pandemic (31

December 2019).

a)

b)

c)

Figure S2. Significant occurrences on GISAID of the enhancing mutations **a)** K417N (nucleotide change

G22813T), **b)** K417T (nucleotide change A22812C), **c)** E484K (nucleotide change G23012A), at the virus

Receptor Binding Domain (RBD), since the start of the COVID-19 pandemic (31 December 2019).

```
SP|Q9BYF1|ACE2_HUMAN   MSSSSWLLLS LVAVTAAQST IEEQAKTFLD KFNHEAEDLF YQSSLASWNY NTNITEENVQ   60
SP|Q8R0I0|ACE2_MOUSE   MSSSSWLLLS LVAVTTAQSL TEENAKTFLN NFNQEAEDLS YQSSLASWNY NTNITEENAQ   60
                       ********** ****:***  :**  ****:  :**:******* ********** *******  *

SP|Q9BYF1|ACE2_HUMAN   NMNNAGDKWS AFLKEQSTLA QMYPLQEIQN LTVKLQLQAL QQNGSSVLSE DKSKRLNTIL  120
SP|Q8R0I0|ACE2_MOUSE   KMSEAAAKWS AFYEEQSKTA QSFSLQEIQT PIIKRQLQAL QQSGSSALSA DKNKQLNTIL  120
                       :*.:*.  ** ** :***::* * :.*****: *  *:***** **.***.** **.*:****

SP|Q9BYF1|ACE2_HUMAN   NTMSTIYSTG KVCNPDNPQE CLLLEPGLNE IMANSLDYNE RLWAWESWRS EVGKQLRPLY  180
SP|Q8R0I0|ACE2_MOUSE   NTMSTIYSTG KVCNPKNPQE CLLLEPGLDE IMATSLDYNS RLWAWEGWRA EVGKQLRPLY  180
                       ********** *****.**** *******.* ***.****:* ******.**: **********

SP|Q9BYF1|ACE2_HUMAN   EEYVVLKNEM ARANHYEDYG DYWRGDYEVN GVDGYDYSRG QLIEDVEHTF EEIKPLYEHL  240
SP|Q8R0I0|ACE2_MOUSE   EEYVVLKNEM ARANNYYDYG DYWRGDYEAE GADGYNYNRN QLIEDVERTF AEIKPLYEHL  240
                       ********** ****:* *** ********.  *.***:*.*. *******:** .*********

SP|Q9BYF1|ACE2_HUMAN   HAYVRAKLMN AYPSYISPIG CLPAHLLGDM WGRFWTNLYS LTVPFGQKPN IDVTDAMVDQ  300
SP|Q8R0I0|ACE2_MOUSE   HAYVRRKLMD TYPSYISPTG CLPAHLLGDM WGRFWTNLYP LTVPFAQKPN IDVTDAMMNQ  300
                       *****.***: :****** * ********** *********  *****.*** *******::*

                                                     |329                      |353
SP|Q9BYF1|ACE2_HUMAN   AWDAQRIFKE AEKFFVSVGL PNMTQGFWEN SMLTDPGNVQ KAVCHPTAWD LGKGDFRILM  360
SP|Q8R0I0|ACE2_MOUSE   GWDAERIFQE AEKFFVSVGL PHMTQGFWAN SMLTEPADGR KVVCHPTAWD LGHGDFRIKM  360
                       .***:***:* ********** *:******.* ****:*.    *.******** **:****:*

                                             |387
SP|Q9BYF1|ACE2_HUMAN   CTKVTMDDFL TAHHEMGHIQ YDMAYAAQPF LLRNGANEGF HEAVGEIMSL SAATPKHLKS  420
SP|Q8R0I0|ACE2_MOUSE   CTKVTMDDFL TAHHEMGHIQ YDMAYARQPF LLRNGANEGF HEAVGEIMSL SAATPKHLKS  420
                       ********** ********** ******.*** ********** ********** **********
```

Spike N501Y, E484K, K417N(T)

Spike N501Y, E484K, and K417T(C)

N501Y

E484K

Species and age of mice (Study reference)	Strain, dose, shedding and transmission*	Evidence of productive SARS-CoV-2 replication in the respiratory tract	Clinical outcomes, histopathological evidence of disease
Aged (9 months old) and young (6 weeks old) BALB/c mice studied. (Gu et al, 2020)[6]	**Strain name:** MASCp6 **Dose:** Intranasal inoculation with 1.6×10^4 PFU **Shedding:** Viral shedding observed in the feces at day 3,5 and 7 after inoculation. **Transmission:** No information	High amounts of viral RNAs in the lungs and tracheas were detected at 3, 5 and 7 days after inoculation in all aged mice, with peak viral RNA loads of ~10^{10} copies/g at 3 days after inoculation. Viral RNAs were also detected in heart, liver, spleen, and brain. Similar tissue distribution of SARS-CoV-2 RNA was also seen in the MASCp6 infected young mice.	**Clinical:** No visible clinical symptoms or body weight loss were observed in both the young and aged mice. No fatalities recorded. **Histopathological:** Led to interstitial pneumonia and inflammatory responses in both young and aged mice, however the lung damage in the aged mice was more severe.
Young (8-week-old) and aged (9 months old) male and female BALB/c mice and C57BL/6 mice studied. (Sun et al, 2021)[21]	**Strain name:** MASCp36 - developed from MASCp6 (Gu et al, 2020)[6] **Dose:** Intranasal inoculation with varying doses including: 1.2 PFU, 12 PFU, 120 PFU, 1200 PFU, 12000 PFU **Shedding:** No information **Transmission:** No information	qRT-PCR results of aged BALB/c and C57BL/6 mice challenged with 12,000 PFU of MASCp36 showed that high levels of SARS-CoV-2 subgenomic RNAs were persistent in the lung and tracheas till 4 dpi. A similar tissue distribution of SARS-CoV-2 was observed in the young BALB/c and C57BL/6 mice.	**Clinical:** All the aged BALB/c and C57BL/6 mice challenged with high doses (1200 or 12,000 PFU) of MASCp36 developed typical respiratory symptoms and exhibited features like ruffled fur, hunched back, and reduced activity. Young BALB/c and C57BL/6 mice were resistant to MASCp36 challenge, and only one animal that received 12,000 PFU challenge died during the observation period. **Histopathological:** Necrotizing pneumonia and extensive diffuse alveolar damages observed on 4 dpi in aged mice. Milder pathology observed in the young mice.
Young adult (12 weeks old) and aged (One year old) BALB/c mice studied. (Dinnon et al, 2020)[7]	**Strain name:** IC-MA1 **Dose:** Intranasal inoculation with 10^5 PFU **Shedding:** No information **Transmission:** No information	High-titre virus replication was also observed in lung tissue at 2 dpi. This cleared by 4 dpi in the young adult mice but continued to persist at 4 dpi in the aged mice, suggesting that there was increased viral replication in the aged mice. MA1 was also observed in the upper airway and viral antigen was present in nasal turbinate epithelium at 2 dpi in young adult mice; viral antigen was found in conducting airway epithelium, interstitium and nasal epithelium in aged mice at 4 dpi.	**Clinical:** No overt clinical signs such as weight loss observed in young adult mice but a significant decrease in body weight was observed in aged mice. Whole-body plethysmography on mice indicated impaired lung function at 2 dpi, with the extent of impairment significantly worse in the aged mice. **Histopathological:** Interstitial congestion, epithelial damage, inflammatory infiltrate and peribronchial lymphocytic inflammation surrounding airways at 2 dpi. Histopathological effects seen in the aged mice are more severe than those observed in young adult mice.
Young (10-week-old) and aged (1 year old) BALB/c mice studied. (Leist et al, 2020)[8]	**Strain name:** IC-MA10 **Dose:** Doses ranging from 10^2, 10^3, 10^4 or 10^5 PFU were tested. However, 10^4 PFU and 10^3 PFU were determined as the optimal doses for analysis in young and aged mice respectively. **Shedding:** Both old and young mice exhibited viral titers in the nasal cavity over the first 3 days of infection. **Transmission:** No information	Virus replication in the lungs of young mice peaked 1–2 dpi and was absent in most surviving mice by 7 dpi. Viral replication in the upper respiratory tract of young mice remained high on 1–3 dpi but was undetectable in most mice by 5 dpi. A similar trend was observed in the aged mice, but the viral replication was detectable for a much longer period.	**Clinical:** Young mice rapidly lost weight and reached maximum weight loss at day 4 post inoculation. On day 5, the collective mortality rate was approximately 15%. Significant weight loss was also observed in aged mice. Mortality rates of 20% and 60% were recorded for infection with 10^4 and 10^5 PFU, respectively in the young mice. **Histopathological:** At the time of necropsy, acute stage lung damage was noted in the young mice. Analysis at 2, 4, and 7 dpi revealed early multifocal damage to conducting airway epithelia and hallmarks of diffuse alveolar damage (DAD) was observed. The aged mice had severe DAD and higher acute lung injury scores. Overall, a dose- and age-related increase in pathogenesis was observed.
C57Bl6, BALB/c and 129S1/SVMJ mice were studied Young (6-8-weeks) or aged (52 weeks old) C57Bl6 mice were used to study the effect of obesity, obesity-associated diabetes, and advanced age on clinical outcomes. (Rathnasinghe et al, 2021)[13]	**Strain name:** WA1-MA-P11 **Dose:** Intranasal inoculation with 2.5×10^4 PFU **Shedding:** No information **Transmission:** No information	Detectable virus titres in lungs and nasal turbinates of 129S1/SVMJ mice and lungs of male and female C57Bl6 and BALB/c mice, obtained at different days post infection	**Clinical:** Weight loss detected in the 129S1/SVMJ mice but not in the C57Bl6 and BALB/c mice. **Effect of increased age, comorbidities:** Both obesity/diabetes and advanced age in mice result in higher morbidity during SARS-CoV-2 infection.
Groups of young (4–6-week-old) female BALB/c mice, young (4–6-week-old female) C57BL/6J mice and aging (8–9-month-old) male BALB/c mice studied. (Wang et al, 2020)[23]	**Strain name:** HRB26M **Dose:** Intranasal inoculation with $10^{4.4}$ PFU in a volume of 50 µL. **Shedding:** No information **Transmission:** No direct evidence of transmission provided. However, the presence of efficient viral replication in the upper and lower respiratory tracts indicates that transmission could be possible.	In young BALB/c mice, viral RNA was detected in the nasal turbinates on day 3, 5, and 7 p.i. (post infection) and the infectious virus was detected on day 3 and 5 p.i. The viral RNA was also detected in the heart, liver, kidney, and spleen on day 3 p.i. In C57BL/6J mice, high viral loads in the nasal turbinates and lungs on days 3, 5 and 7 p.i. Similar observations made in the aging mice. However, when compared to the young mice, the average PFU titres in the aging mice were 3 times higher and 10 times higher in the nasal turbinates and lungs respectively.	**Clinical:** Similar to the young mice, the aging adult mice showed transient weight loss on days 2 and 3 p.i. and recovered thereafter. **Histopathological:** Mild pathological changes were observed in the respiratory tract of young BALB/c mice infected. Moderate-to-severe pathological changes in the lungs after infection in the aging mice.
Young (4–6-week-old) BALB/c mice studied. [Female 12-month-old BALB/c mice only used for the initial generation of the mouse adapted strain, not for any analysis of the mouse adapted strain]. (Huang et al, 2021)[26]	**Strain name:** Hu-1-WBP-1 **Dose:** Intranasally infected with 10-fold serial dilutions of the WBP-1 virus (i.e., 10^2 to 10^5 PFU). Subsequently, detailed analysis on mice was done by infecting mice with 50 µL of 2 LD$_{50}$ [The LD$_{50}$ of WBP-1 is $10^{3.8}$ PFU]. **Shedding:** No information **Transmission:** No information	In mice given the 50 µL of 2 LD$_{50}$: High viral loads were observed in the trachea and lung of infected mice, with lung tissues having the highest number of viral RNA copies. Trace amounts of viral nucleic acid were detectable in turbinate, heart, and spleen.	**Clinical:** In the groups of mice infected with the 10-fold serial dilutions: All groups of infected mice began to lose weight at 2 dpi and the weight loss was dose dependent. Bristled fur and depressed spirits in mice was also observed. The group of mice receiving a dose of 10^5 PFU WBP-1 virus all died (5/5) by 5 dpi. Two out of five mice survived in the group administered 10^4 PFU virus. In the mice given the 50 µL of 2 LD$_{50}$: Clinical signs of infection were observed in infected mice at 2 dpi. **Histopathological:** In mice given the 50 µL of 2 LD$_{50}$, Acute stage lung damage was noted. Moderate interstitial pneumonia with thickened alveolar septa occurred at 3 dpi, which progressed to severe interstitial pneumonia on 5 dpi. Inflammatory cell infiltration was observed in the lung tissue.
10-week-old female BALB/c mice). (Muruato et al, 2021)[27]	**Strain name:** IC-MA4 **Dose:** Mice infected with 10^5 PFU **Shedding:** No information **Transmission:** No information	Evidence of robust viral replication in mouse lungs.	**Clinical:** No information **Histopathological:** No information
10-week-old female BALB/c mice. (Muruato et al, 2021)[27]	**Strain name:** IC-CMA1, IC-CMA2, IC-CMA3 **Dose:** Mice intranasally inoculated with 10^5 PFU **Strain name:** IC-CMA3p20 [IC-CMA3 after passage 20] **Strain name:** IC-CMA3p20 [IC-CMA3 after passage 20] **Dose:** 10^5 PFU was the main dose administered, but doses 10^4 and 10^6 PFU also administered to some mice.	All three mutants (CMA1, CMA2, CMA3) produced approximately, 10^5 PFU per lobe at day 2 post infection. However, no virus was detected at day 4, suggesting rapid clearance by the host. In mice given IC-CMA3p20, robust viral replication was observed. There was a significantly greater viral load in the lung compared to IC-CMA3p20 at 2 dpi.	**Clinical:** None of the three mutants (CMA1, CMA2, CMA3) induced major disease, although both CMA2 and CMA3 caused more weight loss than CMA1. Mice infected with IC-CMA3p20 had a 10% weight loss by day 3 and showed signs of disease including ruffled fur and hunched posture. **Histopathological:** Mice infected with CMA3p20 had significant lung infiltration and inflammation characterized by peribronchiolar, perivascular cuffing, and perivasculitis by day 2 post infection. At 4 dpi, collapsed airways and interstitial pneumonia was observed in the mice.
Young (2 months old) and aged (12-month-old) BALB/c mice (Zhang et al, 2021)[28]	**Strain name:** LG **Dose:** 1×10^5 TCID$_{50}$ in a volume of 50 µl. **Shedding:** No transmission **Transmission:** No information	Effective viral replication in the lungs and trachea of the young and aged mice, although there was more viral replication in the aged mice.	**Clinical:** Weight of the aged mice was slightly decreased, but there were no evident changes in the weight of young mice. **Histopathological:** Lung lesions of the aged BALB/c mice were obvious, but those of young mice were not, with only slight lymphocyte exudation.