But Mouse, you are not alone: On some severe acute respiratory syndrome coronavirus 2 variants infecting mice

Michael J. Kuiper, PhD, is a Biomolecular Modeller who leads the Modelling and Simulations Team in at
the CSIRO Data61, Dockands, Victoria, Australia

Laurence OW Wilson, PhD, is a Research Scientist who leads the Digital Genome Engineering Team at
the CSIRO Transformational Bioinformatics Group, North Ryde, New South Wales, Australia

Shruthi Mangalaganesh, is studying Bachelor of Biomedical Science and Bachelor of Commerce at
Monash University, Clayton, Victoria, Australia, and pursued an internship at the CSIRO Australian
Centre for Disease Preparedness, Geelong, Victoria, Australia

Daniel Reti, BE(Hons), BS, worked as a Research Technician at the CSIRO Transformational
Bioinformatics Group, North Ryde, New South Wales, Australia, and is currently Front-End Software
Engineer at the Centre for Population Genomics, joint initiative of the Garvan Institute of Medical
Research, Sydney, New South Wales, Australia and the Murdoch Children's Research Institute,
Melbourne, Victoria, Australia

Seshadri S Vasan, DPhil, is a Senior Principal Research Consultant and COVID-19 Project Leader at the
CSIRO Australian Centre for Disease Preparedness, Geelong, Victoria, Australia, and Honorary Professor
in the Department of Health Sciences at the University of York, York, UK

Address correspondence to Professor Seshadri S Vasan, CSIRO Australian Centre for Disease
Preparedness, 5 Portarlington Road, Geelong, 3220, Victoria, Australia or email vasan.vasan@csiro.au

NB: MJK and LOWW contributed equally to this paper.

Running Title: On some SARS-CoV-2 variants infecting mice

Abstract

In silico predictions combined with *in vitro*, *in vivo* and *in situ* observations collectively suggest that mouse adaptation of the SARS-CoV-2 virus requires an aromatic substitution in position 501 or position 498 (but not both) of the spike protein's receptor binding domain. This effect could be enhanced by mutations in positions 417, 484 and 493 (especially K417N, E484K, Q493K and Q493R), and to a lesser extent by mutations in positions 486 and 499 (such as F486L and P499T). Such enhancements due to more favourable binding interactions with residues on the complementary angiotensin-converting enzyme 2 (ACE2) interface, are however, unlikely to sustain mouse infectivity on their own based on theoretical and experimental evidence to date. Our current understanding thus points to the Alpha, Beta and Gamma

variants of concern infecting mice, while Delta and 'Delta Plus' lack a similar biomolecular basis to do so.

This paper identifies a list of countries where local field surveillance of mice is encouraged because they

may have come in contact with humans who had the virus with adaptive mutation(s). It also provides a

systematic methodology to analyze the potential for other animal reservoirs and their likely locations.

Keywords

AlphaFold; animal reservoir; COVID-19; *in silico*; *in vitro*; *in vivo*; mouse adaptation; SARS-CoV-2; variants

Introduction

The 'novel coronavirus disease' (COVID-19) has resulted in significant global morbidity and mortality on a

scale similar to the influenza pandemic of 1918.[1] The ongoing pandemic has been sustained through the

activities of human beings, who are the largest reservoir of the causative 'severe acute respiratory

syndrome virus 2' (SARS-CoV-2). This RNA virus in a new host (human beings) is evolving rapidly,

accumulating mutations, and existing as a cloud of variants with quasispecies diversity.[2] Last year, the

world witnessed the risk of this virus acquiring additional reservoirs (such as minks) and new mutations

of consequence (such as 'Cluster 5'), which could increase transmissibility and lead to a potentially weaker

antibody response.[3]

Both the original form of the virus (known as 'D614') and the subsequent more transmissible 'G614'

variant which has replaced it almost entirely in circulation,[4,5] did not infect mice because their

'angiotensin-converting enzyme 2' (ACE2) receptor did not bind the viral spike protein effectively to allow

entry into cells. Since mouse (*Mus musculus*) is a popular animal model of infection, the virus had to be

adapted through techniques such as sequential passaging in mouse lung tissues and modifying the

receptor binding domain (RBD).[6-8] Other strategies to infecting mice with the original form of the virus or

the G614 included transgenic mice expressing human ACE2 (hACE2), and sensitizing the mouse respiratory

tract through transduction with adenovirus or adeno-associated virus expressing hACE2.[9-12] Recently, virus

'variants of concern' (VOC) originating from Brazil, South Africa and the UK (which contain the common

mutation N501Y in the RBD) were shown to infect mice.[13,14] This ability of SARS-CoV-2 variants of concern

to infect mice is unsettling because of the potential to establish additional reservoirs in a species that is

in close contact with people and companion animals, especially as its population is hard to vaccinate or

control.

Methodology

In this paper, we have combined our structural predictions from biomolecular modelling with available

experimental evidence to understand how specific mutations in VOC and mouse adapted strains,

especially in the RBD, have enabled this virus to infect mice. To do this, we compared interactions of these

key mutations with the corresponding regions in both mouse ACE2 (mACE2) and hACE2 using molecular

models based on crystallographic data of the ACE2/RBD interface (protein database 'pdb' entry number

6M17)[15]. As the improved protein prediction software *Alphafold* has just been made available, we used

these new artificial intelligence methods to further validate our model structures.[16] By introducing the

ACE2 models and variants into molecular dynamics simulations, similar to our recent work,[17,18] we

qualitatively identified the key interactions of side chain residues at the ACE2/RBD interface. Further

details are provided under Supplementary Methods for Molecular Modelling. We then compared our *in-*

silico findings with experimental results reported by different research groups with mouse adapted

strains/isolates[6-8,13,21-29] and VOC[14,30-34] (**Table 1**). Taken together, we were able to gain valuable insights

into the likely effects of different mutations of consequence.

We then queried the world's largest public database called 'GISAID' (Global Initiative on Sharing All

Influenza Data) and looked for these mutations, and equivalent mutations of consequence, in the circa

2.4 million SARS-CoV-2 genome sequences available as of 21 June 2021.[29] In brief, sequences were aligned

to the back to the SARS-CoV-2 reference (EPI_ISL_402124, denoted as the WT allele) in order to generate

a file containing all mutations in the variant call format (VCF), which is a concise way for storing gene

sequence variations. We then calculated the ratio of the frequency at which the WT allele versus the

mutant allele were observed across a rolling 14-day window. Given the data is discrete, highly variable in

size across countries, and contains background noise, such a window of time is essential from our

experience to reduce distortions and glean meaningful insights. For example, if 14 WT and 28 mutant

sequences were observed within a specified 14-day window, then the frequencies were 1/day and 2/day

respectively. The ratio of mutant:WT frequency is therefore 2, indicating the mutant allele is appearing

twice as frequently as the WT allele within that period. This was used to create a heatmap for the key

mutations, both individually and in combination (as discussed below). For clarity, we only considered

countries where the mutant:WT ratio exceeded 1.2 across the entire time the mutant was sampled or

within any given 14-day window. To reduce the noise further from low-sampled countries, we also

instituted a minimum threshold of 10 WT and 10 mutant samples over at least 14 days to suggest possible

spread locally. For this reason, a country where a mutant had only been recorded on a single day, or where

only 9 mutant samples were recorded overall, was not included in our analysis. Our heatmap scale has

also been truncated at 2.00 for ease of visual comparison.

Results and Discussion

***In silico* results**

In silico comparison of the interface residues of the RBD/ACE2 complex in human and mouse models

reveals 30 ACE2 residues in close contact, of which 19 are conserved between mACE2 and hACE2

(approximately 63% identity for the contacting ACE2 residues). The RBD mutations associated with the

mouse adaptations listed in **Table 1** can be grouped loosely into 3 regions by their positions on the

ACE2/RBD interface as follows: Region 1 (RBD positions 498, 499 and 501) are centered around the highly

conserved ACE2 residue tyrosine 41 (Y41); Region 2 (RBD positions 417, 493) centered around ACE2

residue 34; and Region 3 (RBD positions 484 and 486) close to a cluster of ACE2 residues 78 to 82. ACE2

residues within 5 Å (chosen to account for molecular fluctuations) of each RBD adaptation are listed in

Table 2, with dissimilar human/mouse ACE2 residues highlighted in yellow. **Figure 1** further illustrates

these residues visually, and how they are relatively positioned in three regions. **Figure 2** aligns the human

and mouse ACE2 to highlight the key differences at the contact points with the RDB (shown in yellow).

Modelling the N501Y mutation at the ACE2/RBD interface reveals a close interaction with the highly

conserved Y41 residue in ACE2, through attractive, non-covalent bonding between aromatic amino acids

known as π-stacking interactions. In mACE2, π-stacking can be enhanced by the proximal substitution of

histidine (H) in place of lysine (K) which is present at position 353 of hACE2. Our modelling also shows

similar π-stacking enhancement with the conserved Y41, through the substitution of RBD glutamine at

position 498 (Q498) to either histidine (H) or tyrosine (Y). These aromatic π-stacking interactions appear

to be reasonably strong as N501Y and Q498H can each sustain mouse adaptation on its own in

experiments.[6,13,23,28] Counterintuitively, our modelling predicts that simultaneous aromatic mutations at

RBD positions 498 and 501 is detrimental to mouse adaptation due to local π-stacking distortion to the

binding interface. This could explain why we have not observed simultaneous aromatic mutations at these

positions in over 2.4 million entries on GISAID as of 21 July 2021. We did detect one adapted isolate on

GISAID from mouse lung homogenate where N501Y occurs with glutamate 498 to arginine (Q498R),[29]

which strictly speaking is not aromatic, however arginine is frequently associated with π-stacking

interactions.[37] Thus, the Q498R mutation is more tolerated with N501Y due to arginine's inherent

conformational flexibility compared to tyrosine or histidine.

In Region 1 (**Table 2**) we also note the proline 499 to threonine (P499T) substitution in two infectious

clones, presumably engineered to enhance Y498. Modelling provides the following insight on this: the

change from P to T will relax the backbone constraints of proline and allow conformational rearrangement

of threonine to contact the conserved ACE2 residues Y41 and L45. However, we don't find any strong

molecular modelling basis for this mutation to sustain mouse adaptation on its own or evolve naturally

alongside adaptive mutations at positions 498 or 501; this is borne out by experimental evidence to date.

In other words, *in silico* predictions combined with *in vitro* and *in vivo* evidence collectively suggest that

mouse adaptation requires an aromatic substitution in either position 501 or position 498 (but not both);

while additional mutations, especially in Region 2 and Region 3 of the RBD as summarized below,

enhancing ACE2 binding interactions and specificity in mice. These predictions are further supported by

AlphaFold that assigns very high confidence scores (>93 in a scale of 0-100) for the structural predictions

involving these key mutations, demonstrating the applications of this improved protein prediction

software especially during pandemics such as COVID-19 and future 'Disease-X'.

From **Table 1**, we see that mouse adapted strains sometimes carry the Q493K/R mutation (polar

glutamine to basic lysine or arginine). Modelling predicts this as being enabled through favourable salt-

bridge interactions with both glutamic acid 35 (E35) or aspartic acid 38 (D38) both of which are conserved

in hACE2 as well as mACE2 (c.f. **Table 2**, Region 2). The K417N substitution (lysine to asparagine), which is

another experimental observation from **Table 1**, is also predicted by modelling to be advantageous for

mouse adaptation due to favourable amide hydrogen bond interactions with interfacial mACE2 residues

asparagine 31 (N31) and glutamine 34 (Q34); such amide hydrogen bonding is not possible in hACE2 as it

has non-amide lysine (K) and histidine (H) residues. With the Gamma variant of concern, it is unclear

whether the K417T enhances the role of N501Y in mouse adaptation in a similar manner. The 'Delta Plus'

variant of concern has the K417N mutation, but there is no molecular modelling basis to believe that it

can infect mice without an aromatic change in position 498 or 501 as described above. It would be

worthwhile to further investigate any interfering role of glycosylation at this interface region, because

hACE2 contains N-linked glycosylation at asparagine 90 (N90) whereas mACE2 does not (its analogous

residue is threonine T90 according to Uniprot references Q8R0I0 and Q9BYF1).

In Region 3, the K484 residue is not positioned directly at the interface and not observed to interact

strongly with any ACE2 residues; however, our model shows occasional salt bridges can be formed with

relatively close glutamic acid residues in positions 35 and 75 that are conserved in both hACE2 and mACE2.

Our simulations show that the distance between K484 and these glutamic acid residues fluctuate

dynamically from 3 to 20 Å, with salt bridges more likely when distances are around 3 Å. Thus, the E484K,

which is present in the Beta and Gamma variants of concern, and more recently in some Alpha isolates as

well, is likely to have an enhancing role through transient salt bridges. The same cannot be said about

E484Q seen in the Delta variant of concern because salt bridge formation is unlikely with the polar

glutamine (Q) residue. With no accompanying aromatic change in positions 498 or 501, we believe that

the E484Q in Delta, and additionally the K417N in 'Delta Plus', cannot sustain mouse infectivity on their

own based on current biomolecular understanding. The residues 75 to 82 in mACE2 are significantly

different from hACE2 (**Table 2**), therefore any mutation in the corresponding RBD interface is worth

investigation. We could find one from experimental observations, the engineered substitution F486L,[27]

and consider it to have at best an enhancing role. As it was observed simultaneously with Q498Y (which

is likely to sustain mouse infectivity on its own), the contribution of F486L to the overall mouse adaptation

remains to be ascertained.

Comparison with *in vitro*, *in vivo* and *in situ* observations

Early *in silico* predictions based on comparative structural analysis of ACE2 suggested that mouse has a very low probability of being infected.[35,36] Although correct about mouse, those analyses also made inconsistent and erroneous predictions that ferrets wouldn't be susceptible, pigs would be susceptible, etc., thus exposing the need for experimental inputs into the model. Therefore, this paper takes into account a range of experimental observations to cross-check our *in silico* predictions through biomolecular modelling. Wan *et al.* reasoned that "*mouse or rat ACE2 contains a histidine at the 353 position which does not fit into the virus-receptor interaction as well as a lysine does*".[35] While this is true of the original Wuhan strain containing asparagine 501 (N501) in the RBD, our modelling indicates why the tyrosine 501 mutation enables mouse infectivity, even on its own. In hACE2, lysine 353 (K353) creates a salt bridge with conserved aspartic acid 38 (D38). In mACE2, lysine 353 is replaced by the aromatic histidine (H353) to complete the salt bridge, as well as contribute to π-stacking with Y501 variant. The N501Y mutation will also lead to favorable π-stacking with the highly conserved tyrosine 41 (Y41) residue in mammalian ACE2, as suggested by Starr *et al.*[38] with deep scanning of RBD mutations and hACE2 affinity assays. These authors highlighted enhanced affinity of F501 (as it had the highest score), followed by Y501, V501, W501 and T501, in that order. But Y501 and T501 require only a single nucleotide change and have been observed more frequently *in situ* (**Figures 3a,3b**) – compared to F501, V501 and W501 which require 2, 2 and 3 nucleotide changes respectively. It is unsurprising that the latter variants requiring two or more changes were rarely observed *in situ* regardless of their high *in vitro* affinity scores from Starr *et al.*[38]

From the above and **Table 3**, we see that *in silico* analysis can provide valuable insights to interpret and bridge *in vitro*, *in vivo* and *in situ* observations on the RBD position 501. A similar analysis is possible with the alternative essential mutation for mouse adaptation at RBD position 498, where the *in vitro* affinity

enhancement order is H498, Y498, F498 and W498 according to Starr *et al.*[38] Of these, H498 (on its own)

and Y498 (with enhancing RBD mutations) have been shown to result in mouse adaptation *in vivo* (**Table**

1);[7,8,23-28] Q498R was also reported once, unusually in combination with N501Y, isolated from mouse lung

after 30 passages. However, *in situ* observations of these variants have been limited to R498 (15

occurrences) and H498 (3 occurrences) so far. It is clear from *in vitro*, *in vivo* and *in situ* analyses (**Table 3**)

that H498, R498 and Y498 are possible but not yet common. This is consistent with our *in silico* predictions

because H498 and Y498 are aromatic (enabling π-stacking with ACE2 Y41; similar to Y501), while R498 has

conformational flexibility and can still be associated with π-stacking interactions.[37] H498 and R498

observed *in situ* require a single nucleotide change from Q498, while Y498 requires two nucleotide

changes (or one change from H498).

Similar insights are also possible for the enhancing RBD mutations (c.f. **Table 1** and **Table 3**). From *in vivo*

and *in situ* observations, we see that K417M is less common than K417N or K417T (**Figure 4a,4b**), although

in vitro studies did not predict any enhancement.[38] *In silico* predictions show that all these require a single

nucleotide change, but that N417 (and Q417) would benefit from amide hydrogen bonding. E484K is

present in Beta and Gamma, and increasingly in Alpha VOC, while E484Q is present in the Kappa variant

of interest that is related to the Delta VOC. In comparison to these two substitutions and notwithstanding

higher *in vitro* affinity scores, R484 and T484 are infrequently observed *in situ*, which is consistent with

our *in silico* predictions because they each require 2 nucleotide changes from E484, or one change from

K484. The F486L and P499T were originally engineered *in vivo*, but have had sporadic *in situ* presence in

human populations. *In silico* predictions suggest that the F486L mutation (accessible by three possible

ways of a single nucleotide change) can aid mACE2 adaptation, due to the human-mouse differences in

ACE2 at the 78-82 region; the P499T is also a single nucleotide change (but only one way from P to T) and

predicted to be rare in comparison. Finally, *in silico* predictions for Q493 substituted by K, L or R (each a

single nucleotide change) are borne out *in vivo* and *in situ*, although their affinity scores from *in vitro*

experiments are low. The affinity scores from Starr *et al.*[38] were developed for hACE2 (not mACE2), so we

expected a greater correlation than what has been observed *in situ* in human populations, but perhaps it

is still early in the pandemic to assess this definitively.

Some mutations in the essential as well as enhancing positions can lead to other mutations. For example,

N501Y, the key mutation common to the Alpha, Beta, and Gamma variants of concern, can lead to F501

with a further nucleotide change. Similarly, the enhancing E484K mutation can also lead to R484 or T484

with a further nucleotide change. We examined whether *in situ* observations are consistent or contrary

to our *in silico* predictions. Indeed, F501 was observed once in Germany (25 May 2021) and Mexico (22

June 2021), while the Y501 has been observed in these countries since 22 October 2020 and 31 January

2021 respectively. UK reported E484R in August 2020, followed by USA, Angola, Brazil and South Korea in

February, April, May and June 2021 respectively. In each case, the E484K was detected prior to E484R –

the former mutation circulating in UK, USA, Angola, Brazil and South Korea from April, March, August,

April and December 2020 respectively. Similarly, E484T was only recently detected in the USA in June

2021, 15 months after the first report of E484K in that country. All eight instances are thus consistent with

our prediction – whether this link is causal or a coincidence is worthy of investigation with local

epidemiological data. While bioinformatics tools can provide useful insights, out of 78 COVID-positive

cases only one sample is on average sent for virus genome sequencing, with huge variations across time

and locations, and lots of missing meta-data.[39,40] This means we are more confident about ruling in (e.g.

when a variant has been detected in a location) than ruling out the possibility of a mutation circulating

purely based on *in silico* data, even if the latter is statistically large (2.4 million as of 21 June 2021).

Our analysis is not just of theoretical interest; it has huge practical applications because mice can be kept

as pets, or come into contact with other pets like cats which are known to be susceptible. Also, mouse

plague can occur in area of COVID-19 outbreaks or endemicity, as is currently the case in New South Wales

and adjacent states of Australia.[41] In order to help public health and animal health professionals, **Figure**

5a,5b shows the countries where key essential and enhancing mutations listed in **Tables 1 and 3** (viz.

N501Y, E484K and K417N/T) have co-occurred. **Supplementary Table S1** gives the raw data, down to

regional counts for these combinations. We believe that this information will help locate areas at risk for

appropriate mitigation measures.

Conclusion and further analyses

Assessing the risk of viruses adapting to new hosts requires careful interpretation of all available data

from *in silico*, *in vivo*, *in vitro* and *in situ* sources. Understanding host adaptation at a molecular level, via

modelling helps reconcile seemingly conflicting, experimental, and clinical observations while a pandemic

is still in progress. We have demonstrated this with the SARS-CoV-2 virus adapting to mice. Our

conclusions come with humility, as they are based on best available evidence up to this point, but allowing

us and others to refine when more evidence becomes available. Armed with our collective understanding

from different approaches, and bolstered by bioinformatics and emerging artificial intelligence

technologies such as *AlphaFold*, we have shown how to position ourselves better to predict and mitigate

virus host adaptations, not just for this pandemic but also for future Disease-X. Further analyses pertaining

to COVID-19 should focus on the role of mutations beyond the spike and RBD; and explore other hosts

like rats and other potential reservoir species (even those that previously exhibited low receptor

activities[35,36,42]) which will be hard to vaccinate or control.

Supplementary Methods for Molecular Modelling

Molecular simulations were performed using NAMD2.14[43] with CHARM36m[44] forcefield employing a

'TIP3' water model. The SARS-CoV-2 spike/ACE2 model was a homology model based on the pdb structure

6M17.[15] Models consisted of the SARS-CoV-2 spike RBD domain (residues 330 to 530), while the truncated

mACE2 protein consisted of residues 19 to 600 built using Swiss Modeller.[45] Glycosylation of the spike and

mACE2 protein was manually constructed using Visual Molecular Dynamics (VMD). Simulations were

run with Periodic Boundary Conditions 'PBCs' using the 'NPT' isothermal-isobaric ensemble at

310K and 1 bar pressure employing Langevin dynamics. The PBCs were constant in the XY

dimensions. Long-range Coulomb forces were computed with the Particle Mesh Ewald method

with a grid spacing of 1 Å. 2 fs timesteps were used with non-bonded interactions calculated

every 2 fs and full electrostatics every 4 fs while hydrogens were constrained with the 'SHAKE'

algorithm. The cut-off distance was 12 Å with a switching distance of 10 Å and a pair-list distance

of 14 Å. Pressure was controlled to 1 atmosphere using the Nosé-Hoover Langevin piston method

employing a piston period of 100 fs and a piston decay of 50 fs. Trajectory frames were captured

every 100 ps. Variants were built from the main RBD/mouse ACE2 model by mutating residues to cover

the main mouse adaptation mutations listed in **Tables 1 and 3**. Models were simulated for 300

nanoseconds. Trajectories were visualized with VMD and Nanome.

Acknowledgements

This work was supported by funding from the Australian government's Department of Finance, National

Health and Medical Research Council, and the CSIRO Future Science Platforms (Principal Investigator:

S.S.V.). We are grateful for support from our colleagues at the Australian Centre for Disease Preparedness

(https://www.grid.ac/institutes/grid.413322.5) (especially Simran Chahal, Trevor Drew and Alexander

McAuley) and the Transformational Bioinformatics Group (especially Denis Bauer, Yatish Jain, Brendan

Hosking and Aidan Tay). The title is from the poem '*To a Mouse: On Turning her up in her Nest, with the*

Plough, November 1785' by Scotland's national poet Robert Burns, in which he says that the mouse is not

alone in proving foresight may be vain as the best-laid schemes of mice and men go oft awry (*But, Mousie,*

thou art no thy-lane, In proving foresight may be vain: The best-laid schemes o' Mice an' Men Gang aft

agley).

Author contributions

Conceptualization, methodology, and funding acquisition, S.S.V.; *in silico* analysis, M.J.K.; *in vitro* analysis,

M.J.K. and S.S.V.; *in vivo* analysis, S.M. and S.S.V.; *in situ* analysis, L.O.W.W. and D.R.; writing – original

draft preparation, S.S.V, M.J.K. and S.M.; writing – review and editing, all authors.

References

1.   Dong E, Du H, Gardern L. An interactive web-based dashboard to track COVID-19 in real time.

Lancet Infectious Diseases. 2020;20(5):533-534. Available at https://doi.org/10.1016/S1473-

3099(20)30120-1 (accessed 1 December 2020).

2.   Bauer DC, Tay AP, Wilson, LOW, et al. Supporting pandemic response using genomics and

bioinformatics: A case study on the emergent SARS-CoV-2 outbreak. *Transboundary & Emerging*

Diseases. 2020; 67(4):1453-1462. Available at https://doi.org/10.1111/tbed.13588 (accessed 1

December 2020).

3.   Lassaunière R, Fonager J, Rasmussen, M, et al. Working paper on SARS-CoV-2 spike mutations

arising in Danish mink, their spread to humans and neutralization data. 2020. Available at

https://files.ssi.dk/Mink-cluster-5-short-report_AFO2 (accessed 1 December 2020).

4.   Korber B, Fischer WM, Gnanakaran S, et al. Tracking changes in SARS-CoV-2 spike: Evidence that

D614G increases infectivity of the COVID-19 virus. *Cell.* 2020; 182(4):812-827. Available at

https://doi.org/10.1016/j.cell.2020.06.043 (accessed 1 December 2020).

5.   Zhang J, Cai Y, Xiao T, et al. Structural impact on SARS-CoV-2 spike protein by D614G

substitution. *Science.* 2021; 372(6541):525-530. Available at

https://doi.org/10.1126/science.abf2303 (accessed 1 May 2021).

6.   Gu H, Chen Q, Yang G, et al. Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine

efficacy. *Science.* 2020; 369(6511):1603-1607. Available at

https://doi.org/10.1126/science.abc4730 and

https://www.biorxiv.org/content/10.1101/2020.05.02.073411v1 (accessed 1 December 2020).

7.   Dinnon KH, Leist SR, Schäfer A, et al. A mouse-adapted model of SARS-CoV-2 to test COVID-19

countermeasures. *Nature.* 2020; 586:560-566. Available at https://doi.org/10.1038/s41586-020-

2708-8 (accessed 1 December 2020).

8.   Leist SR, Dinnon KH, Schäfer A, et al. A mouse-adapted SARS-CoV-2 induces acute lung injury

and mortality in standard laboratory mice. *Cell.* 2020; 183:1070-1085. Available at

https://doi.org/10.1016/j.cell.2020.09.050 (accessed 1 December 2020).

9.   Yuan L, Tang Q, Cheng T, Xia N. Animal models for emerging coronavirus: progress and new

insights. *Emerging Microbes & Infections.* 2020; 9(1):949-961. Available at

https://doi.org/10.1080/22221751.2020.1764871 (accessed 15 May 2021).

10.   Muñoz-Fontela C, Dowling WE, Funnell SGP, et al. Animal models for COVID-19. *Nature.* 2020;

586: 509-515. Available at https://doi.org/10.1038/s41586-020-2787-6 (accessed 1 December

2020).

11.   Zeiss CJ, Compton S, Veenhuis, RT. Animal models of COVID-19. I. Comparative virology and

disease pathogenesis. *ILAR Journal.* 2021; ilab007. Available at

https://doi.org/10.1093/ilar/ilab007 (accessed 1 May 2021).

12.   Veenhuis RT, Zeiss CJ. Animal models of COVID-19 II. Comparative immunology. *ILAR Journal.*

2021; ilab010. Available at https://doi.org/10.1093/ilar/ilab010 (accessed 1 May 2021).

13.   Rathnasinghe R, Jangra S, Cupic A, et al. The N501Y mutation in SARS-CoV-2 spike leads to

morbidity in obese and aged mice and is neutralized by convalescent and post-vaccination

human sera. *medRxiv.* 2021; 2021.01.19.21249592. Available at

https://doi.org/10.1101/2021.01.19.21249592 (accessed 1 February 2021).

14.   Yao W, Wang Y, Ma D, et al. Circulating SARS-CoV-2 variants B.1.1.7, 501Y.V2, and P.1 have

gained ability to utilize rat and mouse Ace2 and altered in vitro sensitivity to neutralizing

antibodies and ACE2-Ig. *bioRxiv.* 2021; 2021.01.27.428353. Available at

https://doi.org/10.1101/2021.01.27.428353 (accessed 1 May 2021).

15.   Yan R, Zhang Y, Li Y, et al. Structural basis for the recognition of SARS-CoV-2 by full-length

human ACE2. *Science.* 2020; 367:1444-1448. Available at

https://doi.org/10.1126/science.abb2762 (accessed 1 December 2020).

16.   Jumper, J., Evans, R., Pritzel, A. *et al.* Highly accurate protein structure prediction with

AlphaFold. *Nature* (2021). Available at https://doi.org/10.1038/s41586-021-03819-2 (accessed

20 July 2021).

17.   McAuley AJ, Kuiper MJ, Durr PA, et al. Experimental and *in silico* evidence suggests vaccines are

unlikely to be affected by D614G mutation in SARS-CoV-2 spike protein. *npj Vaccines.* 2020;

5:96. Available at https://doi.org/10.1038/s41541-020-00246-8 (accessed 1 December 2020).

18.   Riddell S, Goldie S, McAuley A, et al. Live virus neutralisation of the 501Y.V1 and 501Y.V2 SARS-

362   CoV-2 variants following INO-4800 vaccination of ferrets. *bioRxiv*. 2021; 2021.04.17.440246.

363   Available at https://doi.org/10.1101/2021.04.17.440246 (accessed 1 May 2021).

364   19.   Tai W, He L, Zhang X. Characterization of the receptor-binding domain (RBD) of 2019 novel

365   coronavirus: implication for development of RBD protein as a viral attachment inhibitor and

366   vaccine. Cellular & Molecular Immunology. 2020; 17:613-620. Available at

367   https://doi.org/10.1038/s41423-020-0400-4 (accessed 1 December 2020).

368   20.   Grantham, R. Amino acid difference formula to help explain protein evolution. *Science*. 1974;

369   185(4154): 862-864. Available at https://doi.org/10.1126/science.185.4154.862 (accessed 1

370   December 2020).

371   21.   Sun S, Gu H, Cao L, et al. Characterization and structural basis of a lethal mouse-adapted SARS-

372   CoV-2. bioRxiv. 2020; 2020.11.10.377333. Available at

373   https://doi.org/10.1101/2020.11.10.377333 (accessed 1 December 2020).

374   22.   Fagre A, Lewis J, Eckley M, et al. SARS-CoV-2 infection, neuropathogenesis and transmission

375   among deer mice: Implications for spillback to New World rodents. *PLoS Pathogens*. 2021; 17(5):

376   e1009585. Available at https://doi.org/10.1371/journal.ppat.1009585 (accessed 24 May 2021).

377   23.   Wang J, Shuai L, Wang C, et al. Mouse-adapted SARS-CoV-2 replicates efficiently in the upper

378   and lower respiratory tract of BALB/c and C57BL/6J mice. *Protein & Cell*. 2020; 11:776-782.

379   Available at https://doi.org/10.1007/s13238-020-00767-x (accessed 1 December 2020).

380   24.   Liu Z, Zheng H, Lin H, et al. Identification of common deletions in the spike protein of Severe

381   Acute Respiratory Syndrome Coronavirus 2. *Journal of Virology*. 2020; 94(17):e00790-20.

382   Available at https://doi.org/10.1128/JVI.00790-20 (accessed 1 December 2020).

383   25.   Zhou H, Chen X, Hu T, et al. A novel bat coronavirus closely related to SARS-CoV-2 contains

384   natural insertions at the S1/S2 cleavage site of the spike protein. *Current Biology*. 2020;

30:2196-2203. Available at https://doi.org/10.1016/j.cub.2020.05.023 (accessed 1 December

2020).

26.   Huang K, Zhang Y, Hui X, et al. Q493K and Q498H substitutions in spike promote adaptation of

SARS-CoV-2 in mice. *EBioMedicine.* 2021; 67:103381. Available at

https://doi.org/10.1016/j.ebiom.2021.103381 (accessed 24 May 2021).

27.   Muruato A, Vu MN, Johnson BA, et al. Mouse adapted SARS-CoV-2 protects animals from lethal

SARS-CoV challenge. *bioRxiv.* 2021; 2021.05.03.442357. Available at

https://doi.org/10.1101/2021.05.03.442357 (accessed 15 May 2021).

28.   Zhang Y, Huang K, Wang T, et al. SARS-CoV-2 rapidly adapts in aged BALB/c mice and induces

typical pneumonia. *Journal of Virology.* 2021; 95(11):e02477-20. Available at

https://doi.org/10.1128/JVI.02477-20 (accessed 24 May 2021).

29.   Van Noorden, R. Scientists call for fully open sharing of coronavirus genome data. *Nature.* 2021;

590:195-196. Available at https://doi.org/10.1038/d41586-021-00305-7 (accessed 1 March

2021).

30.   Rambaut A, Loman N, Pybus O, et al. Preliminary genomic characterisation of an emergent

SARS-CoV-2 lineage in the UK defined by a novel set of spike mutations. *Virological.* 2020.

Available at https://virological.org/t/preliminary-genomic-characterisation-of-an-emergent-sars-

cov-2-lineage-in-the-uk-defined-by-a-novel-set-of-spike-mutations/563 (accessed 1 March

2021).

31.   Li Q, Nie J, Wu J, et al. SARS-CoV-2 501Y.V2 variants lack higher infectivity but do have immune

escape. *Cell.* 2021; 184(9):2362-2371. Available at https://doi.org/10.1016/j.cell.2021.02.042

(accessed 1 May 2021).

32.   Montagutelli X, Prot M, Levillayer L, et al. The B1.351 and P.1 variants extend SARS-CoV-2 host

range to mice. *bioRxiv.* 2021; 2021.03.18.436013. Available at

https://doi.org/10.1101/2021.03.18.436013 (accessed 1 May 2021).

33.   Faria NR, Claro IM, Candido D, et al. Genomic characterisation of an emergent SARS-CoV-2

lineage in Manaus: preliminary findings. *Virological.* 2021. Available at

https://virological.org/t/genomic-characterisation-of-an-emergent-sars-cov-2-lineage-in-

manaus-preliminary-findings/586 (accessed 1 March 2021).

34.   Lucaci AG, Zehr JD, Shank SD, et al. RASCL: Rapid assessment of SARS-COV-2 clades enabled

through molecular sequence analysis and its application to B.1.617.1 and B.1.617.2. *Virological.*

2021. Available at https://virological.org/t/rascl-rapid-assessment-of-sars-cov-2-clades-enabled-

through-molecular-sequence-analysis-and-its-application-to-b-1-617-1-and-b-1-617-2/709

(accessed 29 May 2021).

35.   Wan Y, Shang J, Graham R, et al. Receptor recognition by novel coronavirus from Wuhan: an

analysis based on decade-long structural studies of SARS coronavirus. *Journal of Virology.* 2020;

94(7):e00127-20. Available at https://doi.org/10.1128/JVI.00127-20 (accessed on 1 December

2020).

36.   Damas J, Hughes GM, Keough C, et al. Broad host range of SARS-CoV-2 predicted by comparative

and structural analysis of ACE2 in vertebrates. *PNAS.* 2020; 117(36):22311-22322. Available at

https://doi.org/10.1073/pnas.2010146117 (accessed 1 December 2020).

37.   Flocco MM, Mowbray SL. Planar stacking interactions of arginine and aromatic side-chains in

proteins. *Journal of Molecular Biology.* 1994; 235(2):709-717. Available at

https://doi.org/10.1006/jmbi.1994.1022 (accessed 1 December 2020).

38.   Starr TN, Greaney AJ, Hilton SK, et al. Deep mutational scanning of SARS-CoV-2 receptor binding

430   domain reveals constraints on folding and ACE2 binding. *Cell.* 2020; 182(5):1295-1310.e20.

431   Available at https://doi.org/10.1016/j.cell.2020.08.012 (accessed 1 December 2020).

432   39.   Bauer DC, Metke-Jimenez A, Maurer-Stroh S, et al. Interoperable medical data: the missing link

433   for understanding COVID-19. *Transboundary & Emerging Diseases.* 2021; 68(4):1753-1760.

434   Available at https://onlinelibrary.wiley.com/doi/10.1111/tbed.13892 (accessed 1 July 2021).

435   40.   Priyadarshini, S. Massive coronavirus sequencing efforts urgently need patient data. *Nature*

436   *India (special issue #13 on COVID-19 crisis).* 2020; 10.1038/nindia.2020.75:11-13. Available at

437   https://www.natureasia.com/en/nindia/pdf/special-issues/13/Nature-India-COVID-19-Crisis.pdf

438   and https://go.nature.com/2y7kUlw (accessed 1 December 2020).

439   41.   NSW Government. Help for regional communities impacted by the mouse plague. 2021.

440   Available at https://www.nsw.gov.au/initiative/mouse-control-support-program (accessed 24

441   May 2021).

442   42.   Zhao X, Chen D, Szabla R, et al. Broad and differential animal angiotensin-converting enzyme 2

443   receptor usage by SARS-CoV-2. *Journal of Virology.* 2020; 94(18):e00940-20. Available at

444   https://doi.org/10.1128/JVI.00940-20 (accessed 1 December 2020).

445   43.   Phillips JC, Braun R, Wang W, et al. Scalable molecular dynamics with NAMD. *Journal of*

446   *Computational Chemistry.* 2005; 26:1781-1802. Available at https://doi.org/10.1002/jcc.20289

447   (accessed 1 December 2020).

448   44.   Huang J, Rauscher S, Nawrocki G, et al. CHARMM36m: an improved force field for folded and

449   intrinsically disordered proteins. *Nature Methods.* 2017; 14:71-73. Available at:

450   https://doi:10.1038/nmeth.4067 (accessed 1 December 2020).

451   45.   Waterhouse A, Bertoni M, Bienert S, et al. SWISS-MODEL: homology modelling of protein

452   structures and complexes. *Nucleic Acids Research.* 2018; 46:W296-W303. Available at

453   https://doi.org/10.1093/nar/gky427 (accessed 1 December 2020).

Mice-infecting SARS-CoV-2 'strains'/isolates	Key mutations in the spike RBD (residues 331-524)[19]							Color legend – mutations scored on Grantham Distance,[20] in the scale of 5 (conservative) to 215 (radical), similar to Riddell et al.[18]
	K417	E484	F486	Q493	Q498	P499	N501	22 24 38 43 53 56 73 94 95 99 143
Mouse Adapted (MA) strains								Authors' synthesis/comments
MASCp6[6]							N→Y	L37F (nsp6), P84S (nsp10), and D128Y (N) were the other observed nonsynonymous mutations.
MASCp25[21]				Q→K			N→Y	MASCp6 passaged serially to develop MASCp25 and MASCp36, which accumulated Q493K and subsequently K417N in the RBD.
MASCp36[21]	K→N			Q→K			N→Y	Highly virulent in aged mice and reported stable/mature RBD binding. It also had I1258V (nsp3), S301L (nsp5), and R32C (N).
IC-MA1[7]					Q→Y	P→T		Spike Q498Y and P499T were engineered for compatibility with Q42 of mouse ACE2.
IC-MA10[8]				Q→K	Q→Y	P→T		Q493K evolved in MA10 and interacts with mACE2's N31; others were T285I (nsp4), K2R (nsp7), E23G (nsp8), and F7S (ORF6).
WA1-MA-P11[13]							N→Y	In addition to N501Y, there were H655Y, T7I (M), L84S (ORF8), S194T (N) and the +KLRS (216-219) insertion also observed by others and predicted to form a solvent-accessible loop.[22]
HRB26M[23] (HRB26-P14)					Q→H			HRB26-P5 had Q498H, N969S, and the 675QTQTN679 deletion observed by others.[24,25] P14 also had A81T (nsp8).
Hu-1-WBP-1[26]				Q→K	Q→H			Q493K and Q498H by P5, interacting respectively with mACE2's N31 via H-bond and Y41 via π-stacking. Nonsynonymous Q493K and T77I (nsp9) mutations increased frequency from P8 to P11.
IC-MA4[27]			F→L		Q→Y			F486L and Q498Y were engineered for infecting mice.
IC-CMA1-3[27]	K→N						N→Y	CMA1-3 all had N501Y. CMA2 also had D128Y; CMA3 also had D128Y, L37F, and P87S. Passaging of CMA3 resulted in K417N, H655Y, and E8V (E).
LG[28]					Q→H			Other mutations present were N74K, H655Y, I6V (ORF10).
IA-N501Y-MA30	K→M	E→K		Q→R	Q→R		N→Y	Homogenate from BALB/c mouse lung after 30 passages. GISAID entry EPI_ISL_1666328.[29]
Variants of Concern (VOC) infecting mice								
Alpha (501Y.V1 or B.1.1.7[14,30])		E→K					N→Y	First reported in the UK, September 2020. Spike also has A570D, P681H, T716I, S982A, and D1118H, in addition to HV69-70 and Y144 deletions. **E484K has been detected in this VOC in 15 countries as of 29 April 2021 (436 entries on GISAID).**
Beta (501Y.V2 or B.1.351[14,31,32])	K→N	E→K					N→Y	First reported in South Africa, May 2020. V2-1 had D80A, D215G, E484K, N501Y, D614G and A701V in the Spike. L18F and K417N arose in V2-2. V2-3 had 242-244 deletion.
Gamma (501Y.V3 or P.1[14,32,33])	K→T	E→K					N→Y	First reported in Brazil, November 2020. L18F, T20N, P26S, D138Y, R190S, K417T, E484K, N501Y, H655Y and T1027I in Spike.
VOC not known to infect mice								
Delta (452R.V3 or B.1.617.2)[34]	K→N							First reported in India, October 2020. Also has L452R in the RBD; G142D, R158G, T478K, P681R and D950N in Spike, along with deletions 156-157. **K417N has been detected in this VOC (called 'Delta Plus') in 12 countries as of 25 June 2021.** The related Kappa variant of interest (B.1.617.1) has L452R and E484Q in RBD (Grantham Distance scores of 102 and 29 respectively).

457 **Table 1** Key mutations in 14 'mouse adapted strains' and 3 'variants of concern' of SARS-CoV-2 known to
458 infect mice
459

	SARS-CoV-2 RBD mouse adaptations	human/mouse ACE2 interface residues within 5 Å of mutation
Region 1	N501 Y	38D/D, 41Y/Y, **353K/H**, 355D/D, 357R/R
	P499 T	41Y/Y, 45L/L, **329E/A**, 330N/N
	Q498 H/R/Y	41Y/Y, 42Q/Q, 45L/L, 49N/N, **61N/K**, 357R/R
Region 2	Q493 K/R	**34H/Q**, 35E/E, 38D/D, 42Q/Q
	K417 M/N/T	**31K/N**, **34H/Q**
Region 3	E484 K/Q	35E/E, 75E/E
	F486 L	75E/E, 76Q/Q, **78 T/K**, **79 L/T**, **82 M/S**

463 **Table 2.** SARS-CoV-2 spike RBD mutations associated with mouse adaptation and their close contact

464 residues in respective ACE2 proteins. Positions with dissimilar human/mouse residues are highlighted in

465 yellow.

	Essential position in RBD		Enhancing position(s) in RBD for mouse adaptation				
Analysis	Q498	N501	K417	E484	F486	Q493	P499
In silico [1]	H Y F W [5]	Y H F W [5]	N Q [8]	K Q	L	K L R	T
In vitro [2]	H Y F W	F Y V W T	-	R K T Q	-	M A Y F K L V	-
In vivo [3]	H R Y	Y	M N T	K Q	L	K R	T
In situ [4]	R H [6]	Y T H F [7]	N T M [9]	K Q R T [10]	L [11]	K L R [12]	T [13]

1. Predicted by our modelling. For E484, F486, Q493, and P499 our work only reconfirmed *in vivo* observations.
2. Arranged in descending order of affinity binding from *in vitro* experiments reported by Starr *et al.*[38]
3. Observed *in vivo* with studies summarized in Table 1 (alphabetical order). Note L486 and P499 were engineered.
4. Observed on GISAID as of 21 July 2021 after removing noise from low sample countries. Only key trends are presented.
5. These aromatic residues are predicted to have similar interactions with Y41 of ACE2.
6. F W Y not observed yet. R (France, Germany, Netherlands) observed in mid 2021. H (Slovenia, USA) observed since early 2021.
7. V W not observed yet. H observed in Norway (early 2021) and F in Mexico (mid 2021). Y and T observed in several countries (**Figure 3a,b**).
8. Modelled mutations predict favorable amide hydrogen bonding. N and Q mutations each require a single nucleotide changes but there are 2 ways to get from N to K.
9. Q not observed yet. N (mutation G22813T) and T observed several countries (**Figure 4a,b**). Interestingly, N (alternative mutation G22813C) has sporadic presence in UK and USA (since late 2020).
10. K and Q observed in several countries (Figures 7,8). Sporadic presence of T (USA in mid 2021), and R (Angola and South Korea since mid 2021) observed.
11. Sporadic presence of different synonymous mutations (T23018C in Netherlands and Columbia; T23020G in USA) observed since late 2020.
12. A, F, M, V, Y not observed yet. Sporadic presence observed for K (Italy since early 2020), L (Trinidad and Tobago in mid 2021) and R (Czech Republic and Romania mid 2021).
13. Sporadic presence since late 2020 in Netherlands, Bulgaria and Canada.

470   **Table 3** Comparison of *in silico*, *in vitro*, *in vivo* and *in situ* observations of key mutations in **Table 1**.

Figure 1. The RBD/ACE2 interface. The receptor binding domain (RBD) is shown in cyan, while the ACE2

is shown in yellow. Pink spheres indicate relative positions of mouse adapting mutations on the RBD, while

the blue spheres represent interface residues that differ between human and mouse ACE2 sequences,

shown in green and orange respectively.

```
SP|Q9BYF1|ACE2_HUMAN MSSSSWLLLS LVAVTAAQST IEEQAKTFLD KFNHEAEDLF YQSSLASWNY NTNITEENVQ 60
SP|Q8R0I0|ACE2_MOUSE MSSSSWLLLS LVAVTTAQSL TEENAKTFLN NFNQEAEDLS YQSSLASWNY NTNITEENAQ 60
                     ********** *****:***   **:*****: :**:***** ********** ********.*

SP|Q9BYF1|ACE2_HUMAN NMNNAGDKWS AFLKEQSTLA QMYPLQEIQN LTVKLQLQAL QQNGSSVLSE DKSKRLNTIL 120
SP|Q8R0I0|ACE2_MOUSE KMSEAAAKWS AFYEEQSKTA QSFSLQEIQT PIIKRQLQAL QQSGSSALSA DKNKQLNTIL 120
                     :*.:*. *** ** :***. * *: *****.   :* ***** **.***.** **.*:*****

SP|Q9BYF1|ACE2_HUMAN NTMSTIYSTG KVCNPDNPQE CLLLEPGLNE IMANSLDYNE RLWAWESWRS EVGKQLRPLY 180
SP|Q8R0I0|ACE2_MOUSE NTMSTIYSTG KVCNPKNPQE CLLLEPGLDE IMATSTDYNS RLWAWEGWRA EVGKQLRPLY 180
                     ********** *****.**** ********:* ***.* ***. ******.**: **********

SP|Q9BYF1|ACE2_HUMAN EEYVVLKNEM ARANHYEDYG DYWRGDYEVN GVDGYDYSRG QLIEDVEHTF EEIKPLYEHL 240
SP|Q8R0I0|ACE2_MOUSE EEYVVLKNEM ARANNYNDYG DYWRGDYEAE GADGYNYNRN QLIEDVERTF AEIKPLYEHL 240
                     ********** ****:*:*** ********.: *.***:*.*. *******:** *********

SP|Q9BYF1|ACE2_HUMAN HAYVRAKLMN AYPSYISPIG CLPAHLLGDM WGRFWTNLYS LTVPFGQKPN IDVTDAMVDQ 300
SP|Q8R0I0|ACE2_MOUSE HAYVRRKLMD TYPSYISPTG CLPAHLLGDM WGRFWTNLYP LTVPFAQKPN IDVTDAMMNQ 300
                     ***** ***: :******* * ********** ********* *****.**** ********::*
                                                     |329                          |353
SP|Q9BYF1|ACE2_HUMAN AWDAQRIFKE AEKFFVSVGL PNMTQGFWEN SMLTDPGNVQ KAVCHPTAWD LGKGDFRILM 360
SP|Q8R0I0|ACE2_MOUSE GWDAERIFQE AEKFFVSVGL PHMTQGFWAN SMLTEPADGR KVVCHPTAWD LGHGDFRIKM 360
                     .***:***:* ********** *:******* * ****:*.: : *.******** **:***** *
                                                     |387
SP|Q9BYF1|ACE2_HUMAN CTKVTMDDFL TAHHEMGHIQ YDMAYAAQPF LLRNGANEGF HEAVGEIMSL SAATPKHLKS 420
SP|Q8R0I0|ACE2_MOUSE CTKVTMDNFL TAHHEMGHIQ YDMAYARQPF LLRNGANEGF HEAVGEIMSL SAATPKHLKS 420
                     *******:** ********** ******.*** ********** ********** **********
```

Figure 2 Sequence alignment of human and mouse ACE2 highlighting contact points with the SARS-CoV-2 spike receptor binding domain in yellow. Differences between contact points are highlighted in green (human) and orange (mouse), while common contact residues are highlighted in cyan.

495 -

497 a)

499   b)

Figure 3. Significant occurrences on GISAID of **a)** N501Y mutation (nucleotide change A23063T), and **b)**

N501T mutation (nucleotide change A23064C), at the virus Receptor Binding Domain (RBD) since the start

of the COVID-19 pandemic (31 December 2019).

504   a)

b)

Figure 4. Significant occurrences on GISAID of **a)** K417N mutation (nucleotide change G22813T), and **b)**

K417T mutation (nucleotide change A22812C), at the virus Receptor Binding Domain (RBD), since the start

of the COVID-19 pandemic (31 December 2019).

514   a)

b)

Figure 5. Significant co-occurrences on GISAID of **a)** N501Y, E484K and K417N triple mutations, and **b)**

N501Y, E484K and K417T triple mutations, at the virus Receptor Binding Domain (RBD), since the start of

the COVID-19 pandemic (31 December 2019). See **Supplementary Table S1** for raw data.

520