Role of polycomb repressive complex 2 in regulation of human transcription factor gene expression

Jay C. Brown
Department of Microbiology, Immunology and Cancer Biology
University of Virginia School of Medicine
Charlottesville, Virginia, 22908

Corresponding author:
Dr. Jay C. Brown
Department of Microbiology, Immunology and Cancer Biology
University of Virginia School of Medicine Box 800734
Charlottesville, Virginia 22908
USA

Email: jcb2g@virginia.edu
Phone: 01-434-924-1814

ABSTRACT

The study described here was pursued with the idea that information about control of

human gene expression might result from comparing the promoter regions of a highly

expressed gene population with that of a related population that is weakly expressed.

Systematic differences in the two promoter populations would be candidates for a role in

causing the distinct expression levels observed. The results reported here were obtained

with human genes encoding transcription factors (TF). Among TF genes, those with wide

tissue transcription are strongly expressed while those expressed in one or only a few

tissues are weakly transcribed in most cases. The promoter regions of the two populations

were compared using the results of ChIP-seq experiments. Results demonstrated that the

promoter region of broadly expressed TF genes is enriched in binding sites for POLR2A, a

component of RNA polymerase II while promoters of tissue targeted genes are enriched in

EZH2, a subunit of polycomb repressive complex 2 (PRC2). It was rare to observe

promoters with binding sites for both POLR2A and EZH2 or for neither one. The findings

are interpreted to indicate that strong expression of broadly expressed TF genes is due to

the presence of RNA polymerase II at the promoter while weak expression of tissue

targeted promoters results from the presence of PRC2. Finally, transcription factor families

were compared in the proportion of broadly expressed and tissue targeted genes. The

results demonstrated that most families possess both broadly expressed and tissue

targeted members. For instance, this was the case with 16 of 20 TF families. The results

are interpreted to indicate that while individual TFs such as EZH2 may be specific for

broadly expressed or tissue targeted genes, this is not a property of most TF families. Most

have both broadly expressed and tissue targeted members.

1. Introduction

Studies on the control of gene expression have benefitted from recent advances in DNA sequencing and from identification of biochemical systems that result in up- or down-regulation of transcription. Whole genome sequences are now available for a wide variety of organisms including humans, and well-characterized elements of regulatory control include promoters, epigenetic signaling, CpG islands, structured chromosome domains, mRNA splicing and many others [1-4]. In view of the variety of regulatory pathways found in higher vertebrates, it is unreasonable to assume that there is an overall regulatory system that applies to all genes. More likely is the idea that distinct regulatory regimes apply to distinct gene populations with the effects of individual regulatory systems integrated to create the overall developmental program observed.

In view of the environment of regulatory studies described above, it is reasonable to suggest that progress in understanding regulatory control might be made by comparing the control elements of two populations of related genes that differ in their level of expression. The idea is that such a comparison might lead to information about the identity of the regulatory features involved, at least in a restricted group of genes.

Here I describe the results of a study employing the strategy described above and focused on the genes encoding human transcription factors (TF). Transcription factors expressed in a wide range of tissues (broadly expressed genes) were found to be highly expressed compared to most tissue targeted TF genes where transcription is weaker. TF expression levels were assessed by the results of RNA-seq experiments while ChIP-seq studies were examined to identify protein binding sites in the gene promoter.

The results are interpreted to support the view that polycomb repressive complex 2

(PRC2) plays a central role in regulating the expression of human TF genes, particularly

those of weakly expressed tissue targeted genes. PRC are multi-subunit molecular

complexes able to bind the promoter region of a target gene and suppress the gene's

expression [5-6]. First discovered in Drosophila in 1947 [7], PRC functions have been

actively studied ever since. PRC or components of PRC are found in all metazoans and

in some more primitive species including fungi. Two forms of PRC complex are known,

PRC1 and PRC2, and both can introduce covalent modifications into the histone

components of chromatin. PRC1 ubiquitinates H2 at lysine119 [8] while PRC2

methylates H3 at lysine27 [9-12]. PRC1 and PRC2 are thought to function together to

repress gene expression and/or to report on the expression due to other control factors.

2. Materials and methods

2.1 Gene databases

2.1.1 All human transcription factor genes (Supplementary Table 1; 1018 genes)

This database was derived from the list of Vaquerizas et al. [13] with a modification

involving zinc finger transcription factor genes. ZNF genes were examined individually

using GeneCards (https://www.genecards.org) and NCBI Gene

(https://www.ncbi.nlm.nih.gov). A ZNF gene was retained in the database only if its role

as a transcription factor was supported by an experimental study.

2.1.2 Sample of all human genes (SupplementaryTable 2, 183 genes)

A sample of all human protein coding genes was accumulated to serve as a control for

analysis of transcription factor genes. Included genes begin with ACO1 and proceed

rightward on chromosome 9 for a total of 183 protein coding genes.

2.1.3 Sample of testis-specific genes (Supplementary Table 3, 204 genes)

This database was curated by Brown [14].

2.1.4 Database of all human transcription factor genes showing predominant structural

features present in the TF protein (Supplementary Table 4, 1018 genes)

Structural element annotations were obtained from GeneCards and NCBI Gene.

2.2 RNA-seq and ChIP-seq results for transcription factor and control genes

Transcription levels for all genes reported here were obtained from the GTEx Portal of

RNA-seq results as reported in the UCSC Genome Browser (version hg38

(https://genome.ucsc.edu/)). A gene was annotated as "broadly expressed" (B in

Supplementary Table 1) if it was found to be expressed in 90% or more of the 52

tissues reported by GTEx. Otherwise, the gene was annotated as "tissue targeted" (T).

Borderline genes were rare (10 of 1018 genes). They are indicated by B/T or T/B in

Supplementary Table 1 depending on the category they most resemble. Transcription

levels for broadly expressed genes were the average of the top 5 tissues reported by

GTEx. For tissue targeted genes, reported values were those of the tissue specific gene

or the average of the top 5 tissues in cases where more than one tissue was in the

group.

Transcription factor binding sites in the promoter region of transcription factor genes

were derived from ChIP-seq studies reported in the ENCODE project database of cis-

Regulatory Elements (3 November 2018 version) by way of the UCSC Genome

Browser. Visual scanning of promoter binding sites was done with UCSC Genome

Browser. An EZH2 entry was included in Supplementary Table 1 if it was present in

50% or more of the studies reported by ENCODE. Values for the level of EZH2

promoter binding were those reported by the Integrated Genomics Viewer

(https://igv.org/app/); the cells and study were GM12878 and ENCFF273NDY,

respectively. Only binding sites in the promoter region are reported here; sites in non-

promoter regions were ignored.

2.3 Transcription factor families and protein structural features

These were derived from GeneCards and from NCBI Gene.

2.4 Data handling

Data were manipulated with RStudio and Excel. Graphic rendering was done with

SigmaPlot v14.5.

3. Results

3.1 Broadly expressed and tissue targeted human TF genes

The project was carried out beginning with a database of 1018 human transcription

factor genes (Supplementary Table 1). The database was that of Vaquerizas et al. [13]

with the modifications described in Materials and Methods. Each gene was classified as

either broadly expressed or tissue targeted using information about its tissue expression

in the RNA-seq results reported by GTEx Portal via the UCSC Genome Browser

(https://genome.ucsc.edu/). Of the 1018 database genes, 589 (58%) were classified as

broadly expressed and 419 (41%) as tissue targeted (Supplementary Table 1, B/T

column).

3.2 Gene expression in broadly expressed and tissue targeted TF databases

It was expected that broadly expressed TF genes would be expressed at a higher level

than that of tissue targeted TF genes because of the need for expression of tissue

targeted TF's to be suppressed in some tissues. This expectation was tested by rank

ordering each population according to the expression level of each gene. The two

populations were then plotted together so that the overall expression level could be

compared. The results (Fig. 1) showed that tissue targeted TF genes are expressed at a

lower level than broadly expressed ones, particularly among the genes with the lowest

transcription levels. Above a rank of ~180, expression of the two populations was more

equal with the tissue targeted population exceeding the broadly expressed genes at the

highest expression levels (Fig. 1). It was concluded that comparison of the two

populations was appropriate for the studies contemplated here.

Broadly Expressed and Tissue Targeted Transcription Factor Gene Populations Used for Analysis

Figure 1. Expression level of broadly expressed compared to tissue targeted human

transcription factors. The expression levels of genes in the two populations were

ordered by level and compared by plotting on the same graph. Note that the level of

tissue targeted TF expression is less compared to broadly expressed genes, particularly

in low expression range.

3.3 TF binding sites in the promoter region of TF genes

TF binding sites were identified using ChIP-seq results available from ENCODE. At first,

identified sites in graphical form were examined visually in the UCSC Genome Browser.

Images revealed that binding sites for POLR2A and EZH2 were asymmetrically

distributed in broadly expressed compared to tissue targeted TF gene promoters. For

instance, among 589 broadly expressed TF genes, 499 (85%) had one or more

POLR2A binding sites in the promoter while among 419 tissue targeted TF genes, 246

(59%) had binding sites for EZH2 (Fig. 2a). It was rare to find EZH2 binding sites in

broadly expressed gene promoters (7%) or POLR2A in tissue targeted genes (12%; see

Fig. 2a). It was also rare to find TF gene promoters that lacked both POLR2A and

EZH2. Of the 1018 genes examined, only 54 (5%) lacked binding sites for both (Fig.

2a).

High Proportion of EZH2 Binding Sites in Promoters
of Tissue Targeted Human Transcription Factor Genes

(a) All database transcription factor genes

(b) Sample of all human genes

(c) Sample of all testis-specific genes

Figure 2. POLR2A and EZH2 binding sites in the promoter regions of broadly

expressed and tissue targeted genes. Results are shown for (a) all database human

transcription factor genes; (b) an unselected population of human protein coding genes;

and (c) a sample of human testis-specific genes. Note that EZH2 binding sites

(indicative of PRC2 binding) are prominent in tissue targeted TF genes, but not in

broadly expressed TF genes (a), a mixed population of human protein coding genes (b)

or in testis-specific human genes (c).

3.4 Control: POLR2A and EZH2 binding sites in the promoters of unselected human

protein coding genes

As a control, an experiment was performed to test whether the asymmetry observed

between POLR2A and EZH2 in the promoter region of TF genes would also be

observed in an unselected population of human protein coding genes. Is asymmetry

observed only in TF genes, or does it exist more widely? To test the idea, the

experiment described above was repeated beginning with an unselected population of

all human protein coding genes rather than with TF genes only (Supplementary Table

2). The results showed that among 141 unselected, broadly expressed genes, 127

(91%) had POLR2A in the promoter while 8 (6%) had EZH2, a result comparable to the

TF gene population. In contrast, in the tissue targeted population the results were 40%

and 29% for POLR2A and EZH2, respectively, a result quite different from the TF genes

(Fig. 2b). The results are interpreted to support the view that the POLR2A/EZH2

asymmetry is found in TF genes only and is not found in an unselected gene population.

3.5 Control: POLR2A and EZH2 binding sites in testis-specific genes

A further control study was carried out to examine POLR2A and EZH2 binding in the

promoters of a population of tissue targeted genes, a population of 204 testis-specific

genes (Supplementary Table 3). ChIP-seq results, as described above, were examined

in the promoter regions of the testis-specific genes. It was expected that EZH2 binding

sites would be observed in a high proportion of genes if testis-specific genes were

regulated in the same way found for tissue targeted TF genes. Quite a different

outcome was detected (Fig. 2c).

Promoters of testis-specific genes were not found to be enriched in either POLR2A or

EZH2. Instead, the promoters had a mixture of transcription factor binding sites in which

no predominant species could be identified (see Supplementary Table 3). For instance,

of 195 testis-specific genes that yielded a result, 135 (69%) had neither a POLR2A nor

an EZH2 site. POLR2A and EZH2 were found in 22% and 9%, respectively (Fig. 2c).

The result is interpreted to indicate that while EZH2 binding sites are prevalent in the

promoters of tissue targeted TF genes, the same is not found in other populations of

tissue targeted genes.

3.6 EZH2 represses TF gene expression as a component of the polycomb repressive

complex 2

Binding of EZH2 to the promoter regions of tissue targeted TF genes as described

above was interpreted to indicate that EZH2 acts to repress TF gene expression as a

component of PRC2, a complex known to be involved in gene repression [5-6]. EZH2 is

a prominent component of PRC2, and PCR2 introduces repressive modifications into

chromatin. To test the idea that EZH2/PCR2 acts to repress TF gene expression, the

expression level of tissue targeted TF genes was plotted against the amount of EZH2

reported in the TF gene promoter (Fig. 3). Creation of the plot was complicated by the

fact that as EZH2 is repressive, it was difficult to find an appropriate population of tissue

targeted TF genes. Most were expressed to only a background level. Fig. 3 shows the

results of all 24 of the genes with transcription levels above background. The regression

line in the plot indicates that the amount of promoter bound EZH2 is corelated with

lower rather than higher gene expression. The results support the conclusion that EZH2

is repressive for its target TF gene population. The results are also compatible with the

idea that EZH2 acts as a part of the PRC2 complex.

Figure 3. Plot showing transcription factor expression level as a function of EZH2

protein bound to the promoter. Note that increased EZH2 binding correlates with

decreased TF gene expression. Data points corresponding to a few individual TFs are

noted on the plot.

3.7 Preferred levels of EZH2 bound at target promoters

The availability of quantitative values for the level of EZH2/PRC2 in the promoters of

tissue targeted TF genes enabled a determination of whether the collection of values is

a continuous function or whether there exist preferred levels. The issue was examined

beginning with all 246 database TF genes that were judged to be tissue targeted and

also to be enriched in EZH2 binding in the promoter (see Supplementary Table 1, EZH2

column). A histogram was plotted relating the amount of EZH2 bound at the promoter

and the number of times a similar value was found among the 246 genes (Fig. 4). The

plot shows evidence of preferred binding levels in promoter associated EZH2. At least

four preferred complexes were identified (at 4,8,12 and 16 EZH2 units), and spacing

was found to be ~4 in the units provided by the ChIP-seq analysis (signal p-value).

Figure 4. Histogram showing TF gene count as a function of the measured level of

EZH2 protein bound at the promoter. Analysis was carried out with all 246 tissue

targeted database TFs with EZH2 bound at the promoter. Note that there are preferred

levels of EZH2 protein bound.

The above result was not expected. Preferred spacing was not observed when similar

histogram analysis was performed with gene expression values from the same dataset.

The observation (Fig. 4) is interpreted to indicate that EZH2/PRC2 complexes are

added to promoter regions in units containing two or more EZH2 molecules. Preferred

spacing arises when more than one multi-EZH2/PRC2 complex attaches to a promoter.

3.8 Transcription factor families

The information above indicating that promoter binding sites for EZH2 are located

preferentially in tissue targeted TF genes raises the question of whether there may be

families of TFs whose members have a selectivity for controlling tissue targeted TF

genes only, or broadly expressed TF genes only. Could binding to tissue targeted

promoters be a property of a TF family and not only of an individual TF? To explore the

above question, the TF family of each TF employed here was recorded and the families

were characterized according to the number of broadly expressed and tissue targeted

genes they contain. All results are shown in Supplementary Table 4 with a sample of 20

families shown graphically in Fig. 5.

Figure 5. Number of broadly expressed and tissue targeted TFs in selected TF families. See Supplementary Table 4 for all results. Note that most families contain both broadly expressed and tissue targeted TF genes. Families shown: ATF, activating TF; CREB, cAMP response element binding TF; E2F, E2 factors; ETV, FGF-regulated TFs; FOX, forkhead box; HOX, homeodomain; IRF, interferon-regulatory factor; KLF, Krüppel-like TF's; LHX, LIM homeobox; NKX, NKX homeodomain box; NR, nuclear receptor TF's; PAX, paired homeobox; POU, Pit-Oct-Unc TF's; SP, SRY-related HMG box; SP, Sp; TBX, T-box; TCF, TCF/LEF; ZBTB, ZBTB family; ZNF, zinc finger TF's; ZSCAN, zinc finger and scan domain TF's.

Of the 20 families, the greatest number (16) contain both broadly expressed and tissue

targeted members. The remaining four families have members with only broadly

expressed (ATF and CREB) and only tissue targeted (NKX and PAX) genes. Members

of the largest family (ZNF; 109 genes) are predominantly broadly expressed genes (101

broadly expressed vs. 8 tissue targeted genes). Forkhead (FOX) and homeobox (HOX)

families have the highest number of tissue targeted genes (31 and 37, respectively; see

Fig. 5). The results support the conclusion that most human TF families contain both

broadly expressed and tissue targeted member genes.

A similar analysis was performed with TFs grouped according to the structural motif

found in the TF protein. Eight prominent structural motifs found among TF proteins were

identified and the number of broadly expressed and tissue targeted TFs was noted in

each one (Fig. 6). The results resembled those observed in family groups. Of the eight

structural groups, both broadly expressed, and tissue targeted TF were found to be

present in all but one (ZNF-BTB; see Fig. 6). Homeobox were found to have the highest

number of tissue targeted TFs and ZNF the largest number of broadly expressed.

Number of Broadly Expressed and Tissue Targeted Genes
in Selected Transcription Factor Structure Groups

Figure 6. Number of broadly expressed and tissue targeted TFs in selected groups

sharing the indicated protein structural feature(s). See Supplementary Table 4 for all

results. Note that all structure groups contain both broadly expressed and tissue

targeted TF genes. Structural groups shown: bHLH, basic helix-loop-helix; bZIP, basic

leucine zipper domain; ETS, erythroblast transformation specific domain; Forkhead,

forkhead box; Homeobox, homeobox sequence; NR, multiple structural domains; ZNF,

zinc finger domain; ZNF-BTB, zinc finger plus bric-à-brac motifs.

4. Discussion

4.1 Experimental strategy; comparing weakly and strongly expressing gene populations

The experimental strategy adopted here suggests itself as a pathway to make further

progress in understanding the control of gene expression. The idea is to start with two

populations of related genes that differ in their level of expression. The results of ChIP-

seq experiments are then used to compare TF binding sites in the promoter regions with

the goal of identifying sites that differ in the two gene populations. In principle, such a

study could begin with any two populations of genes that differ in their level of

expression. In practice, however, the best outcomes might be expected with genes

resembling each other in features such as function, biochemical pathway, or

evolutionary time the genes entered the human genome. In view of the wide range of

regulatory systems that could impact a gene, the chance of identifying genes using the

same control system might be best in genes that resemble each other in other respects.

4.2 Broadly expressed compared to tissue targeted genes

The distinction between broadly expressed and tissue targeted genes is a fundamental

one in developmental and evolutionary biology [15-18]. For a gene that might benefit the

fitness of the human genome, for instance, if the gene can be broadly expressed then

its pathway to incorporation into the genome is simpler than if tissue specific expression

is required. In the case of tissue specific expression, the gene needs to enter the

genome and be expressed, but further evolutionary steps are required to ensure that

the gene is not expressed in off-target tissues. In view of the quite different nature of

broadly expressed and tissue targeted genes, it is not surprising that the two gene types

have different mechanisms to control their expression as described here. Tissue

targeted TF genes were found to have a mechanism, involving PRC2, that is less

abundant in broadly expressed TF genes. In fact, it is reasonable to expect that there

are other yet to be discovered mechanisms that distinguish gene expression in broadly

expressed compared to tissue targeted genes.

4.3 High abundance of EZH2/PRC2 in the promoters of tissue targeted TF

An unexpected outcome of the study described here was the high proportion of

EZH2/PRC2 found in the promoters of tissue targeted TF. The presence of repressive

signals in the promoters was expected [19], but in view of the many types of repressive

signals available in the human genome, it was expected that a greater variety would be

observed. Instead, of 419 tissue targeted TF examined here, 246 (59%) were found to

have EZH2/PRC2 binding sites in the promoter. The high proportion suggests

repression by PCR2 is particularly well suited for repression of tissue targeted TF.

Clues about the close link between PCR2 and tissue targeted TF might arise from

considering X-chromosome inactivation, a prominent focus of PRC function. PRC2 is

well known for its role in suppressing all gene expression from one X chromosome in

human females. Targeted to one of the two X chromosomes by an untranslated RNA

(Xist), PRC complexes bind along the entire chromosome to silence the genes

permanently, creating the Barr body [20-22]. Permanent repression is not observed,

however, in all X inactivation situations. Temporary, reversible inactivation has been

observed in mouse and in mouse ES cells [23-24]. The case for reversible inactivation

therefore needs to be considered open in the case of human TF cells.

PRC2 deposition to create the Barr body differs in another significant way from

deposition at promoter regions. Whereas PRC complexes coat an entire X chromosome

to form a Barr body, deposition sites are widely distributed in the human genome to

suppress tissue targeted TF gene expression [19]. Most tissue targeted TF genes are

expected to be separate sites of PRC2 deposition. This situation creates a more

formidable problem for targeting complexes at TF genes than it does with an entire X

chromosome. Understanding of PCR2 targeting to non-X chromosome sites is further

complicated by the ability of polycomb complexes to bind not only to chromatin, but also

to RNA molecules as they are being synthesizing [25].

4.4 Families of human TF

Evidence that a TF can act selectively on a tissue targeted gene (e.g. EZH2) or a

broadly expressed one (e.g. POLR2A) as described here raises the possibility that such

selectivity might apply to TF groups as well as to individual TFs. Could all members of a

TF family, for instance, be specialized to target only broadly expressed genes? Studies

to address this issue generally supported the view that most TF groups contain

members able to recognize broadly expressed genes and others that recognize tissue

targeted ones. Little evidence was observed for broadly expressed/tissue targeted

specificity among either TF families or among TFs with the same protein structural

motifs (Figs 5 and 6; Supplementary Table 4). Despite the lack of overall specificity

observed, some examples of selectivity were noted. For instance, zinc finger TFs

exhibited a strong preference for broadly expressed TF genes while homeodomain TFs

showed a preference for tissue targeted genes (Fig. 6).

I suggest that future studies of gene regulatory control might benefit from a focus on TF

groups such as NKX and PAX in which all members have tissue targeted expression. It

is expected that the promoter regions of these genes will have binding sites for both

activating elements able to activate genes in the targeted tissue and repressive

elements for non-targeted tissues. Examination of the promoters in such genes might

facilitate the identification of the features that influence gene expression.

Data Availability

All data employed in this study are contained in Supplementary Tables 1-4 from which it

can be freely downloaded.

Acknowledgements

I gratefully acknowledge Karsten Siller for advice on computational aspects of this

project. This manuscript was submitted as a pre-print in the link

https://biorxiv.org/content/10.1101/2021.08.04.455052v1 [26].

References

1.   Lambert SA, Jolma A, Campitelli LF, Das PK, Yin Y, Albu M, et al. The Human

Transcription Factors. Cell. 2018;172(4):650-65.

https://doi.org/10.1016/j.cell.2018.01.029

2.   Schubeler D. Function and information content of DNA methylation. Nature.

2015;517(7534):321-6.

https://doi.org/10.1038/nature14192

3.   Szalaj P, Plewczynski D. Three-dimensional organization and dynamics of the

genome. Cell Biol Toxicol. 2018;34(5):381-404.

https://doi.org/10.1007/s10565-018-9428-y

4.   Cavalli G, Heard E. Advances in epigenetics link genetics to the environment and

disease. Nature. 2019;571(7766):489-99.

https://doi.org/10.1038/s41586-019-1411-0

5.   Blackledge NP, Rose NR, Klose RJ. Targeting Polycomb systems to regulate

gene expression: modifications to a complex story. Nat Rev Mol Cell Biol.

2015;16(11):643-9.

https://doi.org/10.1038/nrm4067

6.   Schuettengruber B, Bourbon HM, Di Croce L, Cavalli G. Genome Regulation by

Polycomb and Trithorax: 70 Years and Counting. Cell. 2017;171(1):34-57.

https://doi.org/10.1016/j.cell.2017.08.002

7.   Lewis EB. A gene complex controlling segmentation in Drosophila. Nature.

1978;276(5688):565-70.

https://doi.org/10.1038/276565a0

8.   Cao R, Tsukada Y, Zhang Y. Role of Bmi-1 and Ring1A in H2A ubiquitylation and

Hox gene silencing. Mol Cell. 2005;20(6):845-54.

https://doi.org/10.1016/j.molcel.2005.12.002

9.   Cao R, Wang L, Wang H, Xia L, Erdjument-Bromage H, Tempst P, et al. Role of

histone H3 lysine 27 methylation in Polycomb-group silencing. Science.

2002;298(5595):1039-43.

https://doi.org/10.1126/science.1076997

10.   Czermin B, Melfi R, McCabe D, Seitz V, Imhof A, Pirrotta V. Drosophila enhancer

of Zeste/ESC complexes have a histone H3 methyltransferase activity that marks

chromosomal Polycomb sites. Cell. 2002;111(2):185-96.

https://doi.org/10.1016/S0092-8674(02)00975-3

11.   Kuzmichev A, Nishioka K, Erdjument-Bromage H, Tempst P, Reinberg D.

Histone methyltransferase activity associated with a human multiprotein complex

containing the Enhancer of Zeste protein. Genes Dev. 2002;16(22):2893-905.

https://doi.org/10.1101/gad.1035902

12.   Muller J, Hart CM, Francis NJ, Vargas ML, Sengupta A, Wild B, et al. Histone

methyltransferase activity of a Drosophila Polycomb group repressor complex. Cell.

2002;111(2):197-208.

https://doi.org/10.1016/S0092-8674(02)00976-5

13.   Vaquerizas JM, Kummerfeld SK, Teichmann SA, Luscombe NM. A census of

human transcription factors: function, expression and evolution. Nat Rev Genet.

2009;10(4):252-63.

https://doi.org/10.1038/nrg2538

14.   Brown JC. Control of human testis-specific gene expression. PLoS One.

2019;14(9):e0215184.

https://doi.org/10.1371/journal.pone.0215184

15.   Hounkpe BW, Chenou F, de Lima F, De Paula EV. HRT Atlas v1.0 database:

redefining human and mouse housekeeping genes and candidate reference transcripts

by mining massive RNA-seq datasets. Nucleic Acids Res. 2021;49(D1):D947-D55.

https://doi.org/10.1093/nar/gkaa609

16.   Eisenberg E, Levanon EY. Human housekeeping genes, revisited. Trends in

Genetics. 2013;29(10):569-74.

https://doi.org/10.1016/j.tig.2013.05.010

17.   Sonawane AR, Platig J, Fagny M, …Kuijjer ML. Understanding Tissue-Specific

Gene Regulation. Cell Rep. 2017;21(4):1077.

https://doi.org/10.1016/j.celrep.2017.10.001

18.   Liu X, Yu X, Zack DJ, Zhu H, Qian J. TiGER: a databbase for tissue-specific

gene expression and regulation. BMC Bioinformatics. 2008;9:271.

https://doi.org/10.1186/1471-2105-9-271

19.   Lee TI, Jenner RG, Boyer LA, Guenther MG, et al.Control of development

regulators by Polycomb in human embryonic stem cells. Cell. 2006;125(2):301-313.

https://doi.org/10.1016/j.cell.2006.02.043

20.   Kohlmaier A, Savarese F, Lachner M, Martens J, Jenuwein T, Wutz A. A

chromosomal memory triggered by Xist regulates histone methylation in X inactivation.

PLoS Biol. 2004;2(7):E171.

https://doi.org/10.1371/journal.pbio.0020171

21.   Silva J, Mak W, Zvetkova I, Appanah R, Nesterova TB, Webster Z, et al.

Establishment of histone h3 methylation on the inactive X chromosome requires

transient recruitment of Eed-Enx1 polycomb group complexes. Dev Cell. 2003;4(4):481-

95.

https://doi.org/10.1016/S1534-5807(03)00068-6

22.   Plath K, Fang J, Mlynarczyk-Evans SK, Cao R, Worringer KA, Wang H, et al.

Role of histone H3 lysine 27 methylation in X inactivation. Science.

2003;300(5616):131-5.

https://doi.org/10.1126/science.1084274

23.   Okamoto I, Otte AP, Allis CD, Reinberg D, Heard E. Epigenetic dynamics of

imprinted X inactivation during early mouse development. Science.

2004;303(5658):644-9.

https://doi.org/10.1126/science/1092727

24.   Brookes E, Santiago I, Hebenstreit D, Morris KJ, Carroll, T. et al. Polycomb

associates genome-wide with a specific RNA polymerase II variant, and regulates

metabolic genes in ESCs. Cell Stem Cell. 2012;10(2):157-70.

https://doi.org/10.1016/j.stem.2011.12.017

25.   Beltran M, Yates CM, Skalska L, Dawson M, Reis FP, et al. The interaction of

PRC2 with RNA or chromatin is mutually antagonistic. Genome Res. 2016;26(7):896-

907.

https://doi.org/10.1101/gr.197632.115

26.   Brown JC. Role of polycomb repressive complex 2 in regulation of human

transcription factor gene expression.

https://biorxiv.org/content/10.1101/2021.08.04.455052v1