Default Mode Network Connectivity Predicts Individual Differences in Long-Term

Forgetting: Evidence for Storage Decay, not Retrieval Failure

Yinan Xu[1,*], Chantel Prat[1,2], Florian Sense[3,4], Hedderik van Rijn[3,4], and Andrea Stocco[1,2]

[1] Department of Psychology, University of Washington, Seattle, WA USA

[2] Neuroscience Program, University of Washington, Seattle, WA USA

[3] Department of Experimental Psychology, University of Groningen, Groningen, NL

[4] Behavioral and Cognitive Neuroscience, University of Groningen, Groningen, NL

[*] Now at the Department of Psychology, University of Houston

Author Note

Correspondence should be sent to Andrea Stocco,

Campus Box 351525, University of Washington,

Seattle, WA 98195

Email: stocco@uw.edu

Abstract

Despite the importance of memories in everyday life and the progress made in understanding how they are encoded and retrieved, the neural processes by which declarative memories are maintained or forgotten remain elusive. Part of the problem is that it is empirically difficult to measure the rate at which memories fade, even between repeated presentations of the source of the memory. Without such a ground-truth measure, it is hard to identify the corresponding neural correlates. This study addresses this problem by comparing individual patterns of functional connectivity against behavioral differences in forgetting speed derived from computational phenotyping. Specifically, the individual-specific values of the speed of forgetting in long-term memory (LTM) were estimated for 33 participants using a formal model fit to accuracy and response time data from an adaptive fact learning task. Individual speeds of forgetting were then used to examine participant-specific patterns of resting-state fMRI connectivity, using machine learning techniques to identify the most predictive and generalizable features. Our results show that individual speeds of forgetting are associated with resting-state connectivity within the default mode network (DMN) as well as between the DMN and cortical sensory areas. Cross-validation showed that individual speeds of forgetting were predicted with high accuracy ($r = .78$) from these connectivity patterns alone. These results support the view that DMN activity and the associated sensory regions are actively involved in maintaining memories and preventing their decline, suggesting that forgetting is better understood as a result of storage decay, rather than retrieval failure.

Keywords: cognitive modeling, long-term memory, resting-state fMRI, functional connectivity

Introduction

Despite considerable progress in memory research, we still struggle to understand how and why memories are lost and *forgotten*. In fact, researchers disagree even about when and how forgetting takes place. Consider, for example, the three-stage model of a memory's lifecycle, which is a staple of cognitive research [1]. In this idealized depiction, memories are first encoded as engrams in the hippocampus (synaptic consolidation); then undergo a period of consolidation (system consolidation); and they are finally retrieved at some later time—which can vary from minutes to years. Because forgetting is only inferred after a failed retrieval attempt, it is difficult to pinpoint at which of the earlier stages it really occurred. Specifically, researchers disagree on whether forgetting is a process that is best thought of as occurring during the consolidation stages [2,3] or during the retrieval stage [4–6] .

According to the *storage decay* class of theories, forgetting is a process that mostly affects maintenance and consolidation of memories. In this view, engrams are intrinsically labile and accumulate progressive damage over time. Such damage could be due, for example, to biological decay processes, such as the spontaneous loss of synapses [3,7,8]. It could also be due to the engrams' synapses being recruited to encode new traces (a form of catastrophic interference: [9]). Finally, it has been suggested that spontaneous brain activity might play a pivotal role in consolidating memories [10], and that its disruption might underpin abnormal forgetting in amnestic neurodegeneration [11,12].

According to the *retrieval failure* class of theories, engrams are relatively stable over long intervals, and forgetting is instead driven by an inability to access and retrieve them. As in the case of storage decay, multiple processes have been hypothesized to explain retrieval failure. For example, memories might become less accessible because of the accumulation of more

memories that compete for retrieval (retrieval interference). Most theories assume that initial encoding of memories takes place in the hippocampus, and that controlled memory retrieval is underpinned by the prefrontal cortex; thus, it is possible that loss of synaptic connections between prefrontal cortices and the hippocampus render some memories inaccessible. Evidence for these theories comes from the fact that direct neurostimulation of the engram can re-awaken long-forgotten memories, a fact that has been reported in humans [13] and, more recently, in animal models of Alzheimer's Disease [14].

Implications and Predictions of Storage Decay vs. Retrieval Failure

The distinction between these two families of theories is not merely academic; it has important implications for both memory research and clinical practice. Consider, for example, Alzheimer's Disease (AD), a dramatic form of abnormal forgetting. If the storage decay theory is correct, the promising new clinical treatments that remove amyloid beta deposits (e.g., [15,16]) would stop the progression of the disease but would not restore memory function and would not bring back lost memories, as the underlying neural substrate has already been compromised. However, according to the retrieval failure theory, at least some lost memories might be retrievable, and function could be partially restored. Knowing which theory is correct, therefore, is important to guide treatment and future clinical research.

The current understanding of the mechanisms by which memories are encoded, maintained, and retrieved makes it possible to distinguish between these two theories. Most theories assume that the initial engrams in the hippocampus work as a hub connecting to the distributed sensory representations that form a memory. Activation of this engram results in the re-experience of the associated memory, which is associated with a re-enactment of the original neural activation in sensory regions [17,18]. Replays of this activity, either through active

retrieval or spontaneous brain activity, result in consolidation. This internal re-activation of the memories is supported by the coordinated activity of the so-called Default Mode Network (DMN), a large-scale brain network that includes the hippocampus, the medial frontal cortex, and the precuneus [19]. The DMN is believed to play a key role in propagating the re-activation of the original engrams, mediating the interactions between the hippocampus and the neocortical regions during consolidation [20]. In fact, the DMN's activity at rest is explained as a way of rehearsing and maintaining memory as a form of "priors" for later processing [10,21,22].

The controlled retrieval of memories, on the other hand, is supported by the activity of multiple dorsal and frontal regions. Dorsal regions, in particular, are involved in voluntary memory retrieval as well as working memory, and provide a natural connection between the two forms of memory [23]. Ventral lateral prefrontal regions are specifically involved in different facets of retrieval [24], including the selection of competing memories [25] or the maintenance of retrieval cues [26–28]. Ventral and medial prefrontal regions, finally, seem to play a critical role in monitoring the retrieval process, as evidenced by their involvement in meta-memory judgments and feeling-of-knowing [29–32] and in confabulation in frontal patients [33,34].

Since consolidation and retrieval processes have different neural substrates, the storage decay and retrieval failure theories also make different predictions. According to the storage decay framework, individual differences in forgetting should be accompanied by differences in the functional connectivity between the DMN [20,35] and the sensory association areas, where long-term memories might be stored as the end-product of consolidation [36–38].

According to the retrieval failure framework, however, forgetting occurs because of an inability to properly access previously stored memories. Because the retrieval process is controlled by prefrontal cortices, the theory predicts that forgetting would be associated with the

functional connectivity between the prefrontal and parietal regions that oversee controlled

retrieval and the encoding and consolidation regions of the DMN [39].

Obstacles to the Testing the Theories

Since the two theories make different predictions, it should be possible to empirically test

them by examining which features of individual anatomy and functional connectivity co-vary

with individual differences in forgetting. One important obstacle, however, remains: it is

extremely difficult to *quantify* forgetting in the first place. As noted above, forgetting is typically

operationalized as failing to correctly recall or recognize an item in a memory test: performance

failure is the only ground truth available. Unfortunately, this carries significant limitations. First,

common behavioral measures such as the proportion of correctly remembered items are prone to

statistical misinterpretations [40,41]. Second, performance in memory tests is also affected by

the specifics of the paradigms used. For example, memory performance differs greatly when the

same materials are tested using recognition or recall; recognition is generally easier, but recall,

being more effortful, leads to better consolidation and retention.

This retrieval accuracy-based operationalization does not easily account for possible

temporal dynamics of forgetting. If forgetting is conceptualized as a process, a more precise

assessment of an engram's accessibility is needed even if a memory test results in a successful

retrieval.

The goal of this study is to overcome these limitations through *computational*

phenotyping [42,43], that is, by fitting a formal model of cognitive processes to individual task

data, and using the model's parameters to quantify individual differences. Model parameters

obtained through computational phenotyping are easier to interpret and more reliable than raw

behavioral measures [43–45], such as accuracies or response times. In this study, individual

differences in forgetting were quantified by fitting a computational model of memory consolidation to individual learning data in a sample of healthy individuals for whom functional connectivity data were available. By analyzing which connectivity features best predict individual differences in forgetting, we can empirically compare the storage decay and retrieval failure theories.

A Computational Theory of Forgetting

As referred to above in the context of proportion of correct responses, purely behavioral measures of psychological constructs often have poor reliability [41], which leads, in turn, to poor brain-behavioral associations [46]. One way to obviate this limitation is to fit realistic computational models to individual behaviors, and use the fitted model parameters as dependent variables [47,48]. When the model captures the dynamics of the underlying neural process, its parameters can be more reliable and interpretable than pure behavioral measures. This has been shown, for example, with the threshold and drift parameters of accumulator models in perceptual decision-making (White et al., 2014, 2016) and in learning rate parameters in reinforcement learning models of learning [45]. Finally, using realistic models of long-term memory avoids the pitfalls that were originally pointed out by Loftus [40,49] in applying statistical linear models to non-linear effects in memory performance.

In this study, the estimation of the speed of forgetting was based on a well-established model of episodic memory [44,50–53]. This model, coupled with an adaptive fact-learning task that dynamically adjusts the presentation schedules of the stimuli, yields individual speeds of forgetting for individuals that can be measured within a short time (< 15 mins) but can be reliably extrapolated over very long intervals [44,53,54].

In the model, the probability of retrieving a memory m at time t is proportional to its activation $A(m, t)$, a scalar quantity that reflects a memory's availability. As in the multiple trace theory [55,56], each memory m is associated with multiple episodic *traces*, each of which corresponds to a specific engram that was created when m was retrieved or (re)encoded, and each of which decays over time.

Figure 1: *Activation over time of a memory that has been encoded in three different episodes at times t = 1, 20, and 75 seconds. Each episode leaves an independent memory trace (colored lines) with a characteristic decay rate. The total activation of the memory is the sum of the decaying traces (gray line).*

The activation of each trace decays as a power function of time, and the activation of m is the log sum of the decaying activations of all its traces:

$$A(m,\ t) = \log \textstyle\sum_i [t - t(i)]^{-d(i)} \tag{1}$$

where $t(i)$ and $d(i)$ are, respectively, the time of the creation of the i-the trace and its characteristic decay rate. A trace's decay rate, in turn, is related to the m's activation at the moment each trace is created, thus capturing the fact that traces with higher initial activation decay faster than traces with lower activation [51]. Specifically:

$$d(i) = e^{A(m,\ t(i))} + \varphi$$

with $A(m,\ t(i))$ being the activation of m at the moment the i-th trace was encoded. Thus, if the times at which a memory m has been encoded, re-encoded, and retrieved are known and controlled for, its activation depends only on φ, which is the only free parameter in the above equations and, by determining the decay of each memory, was taken to represent an individual-specific *speed of forgetting* (SoF).

Previous studies have shown that individual values of φ are stable across time and materials and predictive of real-world outcomes such as learning and test scores [44,53,57–59], thus meeting the psychometric criteria of reliability and validity [60]. Finally, at least one previous study has established that speed of forgetting, thus measured, reliably correlates with individual differences in neural activity [61].

Experimental Predictions

To distinguish between the retrieval failure and storage decay hypotheses, this study applied a statistical learning approach [62] to resting-state fMRI data from healthy participants. Specifically, a regularized linear model's hyperparameter was tuned to minimize the cross-validated prediction error of an individual's speed of forgetting from their resting-state connectivity. The specific patterns of connectivity that were identified to successfully predict individual differences in forgetting would shed light onto the biological nature of the phenomenon by indicating whether the involved networks align best with the retrieval or the storage decay hypothesis.

Resting-state activity fMRI was chosen because of its spatial precision and its reliability within participants [63]. It has been speculated that brain activity at rest reflects the invariant functional architecture of the brain [22,64]; small variations within this architecture are characteristic to every individual, to the point that it can be used to "fingerprint" individual participants [65,66]. Consistent with this finding, individual differences in the resting-state functional connectivity between different regions have been shown to be predictive of performance differences in a variety of cognitive abilities, including working memory [67], executive functions [68,69], motor learning [70], perceptual discrimination [71], and most importantly, recognition memory [72]. Thus, resting-state functional connectivity provides an ideal target to identify the biological underpinnings of individual differences in long-term memory retention and forgetting.

Specifically, we hypothesized that the speed of forgetting would reliably co-vary with the functional connectivity between different sub-networks of the DMN. This prediction is common to both hypotheses and follows from the central role played by the DMN in consolidation.

In addition, we also hypothesized that individual differences in the speed of forgetting will be reflected in the connectivity between the DMN and other networks. These patterns of connectivity would allow us to distinguish between the storage decay and the retrieval failure hypotheses. If the storage decay framework is correct, then we expect to see that individual differences in forgetting are best predicted by patterns of connectivity between the DMN, sensory, and associative cortical areas [38,56,73], including higher-level visual areas such as those associated with the ventral and dorsal visual attention networks.

If, on the other hand, the retrieval failure framework is correct, then we would expect to find that individual differences in forgetting are best predicted by the connectivity between the DMN and brain networks associated with frontal and parietal control regions.

Materials and Methods

Participants

A total of 33 monolingual English-speaking participants (19 females) aged between 18 to 33 years old were recruited from a pool of University of Washington undergraduates for whom resting-state fMRI data had already been acquired in a previous, unrelated experiment. All participants provided informed consent and were compensated with a $25 gift card for their participation in the study. All of the recruitment and testing procedures were approved by the University of Washington's Institutional Review Board.

Adaptive Fact-Learning Task

Individual speeds of forgetting were estimated using the adaptive fact-learning task described in [44]. The stimuli consisted of 25 Swahili-English word pairs selected from a previous study [74,75]. The task dynamically interleaved study trials and test trials. On study trials, a Swahili word (e.g., "samaki") and its corresponding English equivalent ("fish") were presented simultaneously on the screen, and participants typed in the English word to proceed.

On test trials, only the Swahili word was presented on the screen as a cue (e.g., "samaki"), and

participants were again asked to respond by typing the corresponding English word ("fish") in an

empty textbox. Trials were self-paced, with no cap on the time allowed for a response.

Consecutive trials are separated by a 600 ms ISI after a correct response and a participant paced

ISI following an incorrect response during which feedback was shown on screen. The order of

repetitions and moment of introduction for each item were determined by the adaptive

scheduling algorithm outlined below and described in more detail elsewhere [54,57]. The task

lasted approximately 12 minutes.

Figure 2. *Visual illustration of one complete study trial and the onset of a test trial from the*

adaptive vocabulary learning task. Note that study items are repeated based on an estimate of

when their memory activation would drop below a predefined threshold.

The scheduling algorithm was designed to optimally interleave the stimuli so that items

would be repeated before the internal cognitive model predicted that they would be forgotten as

determined by the continuously updated speed of forgetting (φ) for each item. In the model, each

memory *m* is a pair associate (see [76]) that links a new Swahili word ("samaki") to a known

English word ("fish"). In response to a study probe, participants are given the Swahili word

("samaki") as a cue to retrieve *m* ("samaki" - "fish"). Only when *m* is retrieved can participants

type in the correct answer ("fish"). In turn, φ is estimated from the time *T* it takes participants to

respond (first key press), which is related to each pair's activation by the equation:

$$T = t_0 + e^{-A(m,\ t)}$$

where $A(m,\ t)$ is the current memory activation (Eq. 1) and t_0 represents a fixed offset

(300 ms) for non-memory related processes, such as the time spent for encoding the visual

stimulus and making a motor response [77,78]. Thus, responses that are either incorrect or

slower than expected result in upwards adjustments of φ, while responses that are faster than

expected result in downwards adjustments of the parameters.

rs-fMRI Data Collection and Preprocessing

Each resting-state run lasted 7 minutes and consisted of 210 echo-planar images (EPI)

collected with TR = 2,000 ms, TE = 25 ms, FA = 72 degrees. Each image in the series consisted

of 36 oblique 3-mm bi-dimensional slices with no gap in-between, with an in-plane resolution of

80x80 3x3 mm voxels. Each functional series was preprocessed using the AFNI [79] software

package. Specifically, each resting-state time series was de-spiked, corrected for differences in

slice acquisition time, realigned to the mean image in the series, normalized to the MNI

ICBM152 template, and spatially smoothed with a 3D 8-mm FWHM Gaussian filter. Artifacts

induced by subject movement were removed by regressing out the six motion parameters and

their temporal derivatives from each voxel's time series. In addition to the functional data, a

high-resolution anatomical MP-RAGE was also collected to aid in the normalization process.

Brain Parcellation

To calculate functional connectivity, each participant's brain was divided into discrete

regions using the network parcellation originally proposed by Yeo et al [80]. Figure 3 depicts the

17 brain networks identified by Yeo, using the color scheme proposed in the original paper.

Figure 3. *The 17 networks in the Yeo et al (2011) parcellation. Subcortical regions, not included*

in the parcellation, are marked in black.

To test the storage decay and retrieval failure hypothesis, each network was assigned to

one of three possible groups: the DMN (consisting of Default A, B, and C), the Retrieval group

(consisting of networks including key prefrontal and parietal regions, such as Control A and B,

Dorsal Attention A, Salience B, Limbic B, and Temporal Parietal) and Storage group (all

networks encompassing the sensory areas: Visual A and B, Somatomotor A and B, Dorsal

Attention B, Salience A, Limbic A, and Control C).

Figure 4: *The grouping of the networks from the Yeo et al (2011) parcellation according to their involvement in consolidation (DMN, black) storage (blue) or retrieval (brown) processes.*

Connectome Extraction

Each participant's global pattern of brain connectivity, known as the *connectome* [81], was extracted using a standard procedure [82,83]. For each of the 17x17 pairs of regions in the parcellation, the Pearson correlation coefficient between the respective time series was computed.

One problem with this canonical connectome is that the raw correlation coefficients are almost all positive and partially driven by common, correlated factors. To remove these confounds, we followed the procedure of Cole and collagues [82], and recalculated the correlation matrices using *partial* correlation coefficients, so that, when computing the correlation between two networks, the correlations between each network in the pair and the remaining 15 networks were partialled out. The coefficients data from the individual partial-correlation connectomes were the functional connectivity data that was fed into the statistical analysis (see below).

Statistical Analysis

Functional connectivity data were analyzed with the Least Absolute Shrinkage and Selection Operator [LASSO: 84], a statistical learning method [85,86] that combines linear regression and feature selection in a single framework. Like linear regression, LASSO estimates a vector of regression weights $\boldsymbol{\beta^*}$ that minimizes the squared sum of the differences between the observations $\mathbf{y}$ and their predicted values, that is $\text{argmin}_{\boldsymbol{\beta}} \|\mathbf{y} - \boldsymbol{\beta}\mathbf{X}\|_2$ (where the notation $\|v\|_n$ represents the Ln norm of the vector $\mathbf{v}$). Unlike traditional regression, the difference is further penalized in proportion to the sum of the absolute values of the regressor weights (their L1 norm), so that:

$$\boldsymbol{\beta}^* = \text{argmin}_{\boldsymbol{\beta}}(\|\mathbf{y} - \boldsymbol{\beta}X\|_2 + \lambda\|\boldsymbol{\beta}\|_1)$$

The use of the L1 norm allows the least informative regressors to shrink to zero and, effectively, be dropped from the model—thus combining shrinkage and feature selection [84]. In turn, this reduces the complexity of the model, favoring generalizability and reducing overfitting. The hyperparameter λ represents the tradeoff between model simplicity (encouraged by the $\|\boldsymbol{\beta}\|_1$ penalty) and accuracy (encouraged by the least-squares minimization term $\|\mathbf{y} - \boldsymbol{\beta}\mathbf{X}\|_2$). As with all machine learning approaches, careful validation is needed to estimate the values of λ and to identify the proper predictors; in this project, Leave-One-Out validation was used to select the proper value of λ by minimizing the prediction error over the out-of-sample observation; the result of this process, therefore, is a set of sparse regressors that have the greatest predictive weight. Once the best value of λ was identified, a final model was then constructed by refitting the entire dataset.

Results

Speed of Forgetting

One SoF parameter was calculated for each individual by averaging across the SoF values of each word pair they studied and which was presented at least three times. The resulting values ranged from 0.253 to 0.385, with a mean of 0.305 and a standard deviation of 0.029. Both the mean and the range of values are in line with previously published studies [44,61].

Figure 5: *Distribution of the observed speeds of forgetting (N=33). Rug plot lines on the x-axis indicate values from each individual participant.*

Resulting Connectome and LASSO Model

Figure 6 shows the group-averages of the resulting individual connectome matrices. The left side shows shows a canonical distribution of *r* values, including the zones of maximum *r*

values along the diagonal (corresponding to connectivity within the same networks). This pattern

is entirely consistent with previous findings using the same parcellation scheme.

The right side of Figure 6 shows the average of the partial correlation connectomes As

expected, this second matrix is sparser and includes both negative and positive correlations

[82,87], giving a clearer picture of the underlying functional organization. The partial

connectomes were the features entered into the LASSO model.

Although LASSO reduces the number of predictors, it has been shown that its

performance is hindered when the number of regressors is much larger than the number of

participants [88], which would be the case if we considered the full $17 \times 16 / 2 = 136$ possible

connections between networks. Thus, to reduce the number of regressors, we limited the analysis

to the connections between the three default mode networks (Default A, B, and C in Figure 1),

including their connectivity with each other, and all of the other 14 networks. Thus, the total

number of regressors included in this analysis was $3 \times 14 + 3 = 45$. Notice that this does not

reduce the generality of our findings, since our theoretical hypotheses concern precisely the

relationship between the default mode networks and other networks (associative vs. control

networks).

Figure 6. *Mean raw Pearson correlations (left) and partial correlations (right) between each of the 17 networks across all participants. The mean correlation coefficient across participants was calculated by first turning each individual coefficient into a z-value using Fisher's r-to-z transformation, then averaging the individual z-values, and finally transforming the average z value back into an equivalent r coefficient. Network names are color-coded based on whether the network belongs to the Storage, Retrieval, or DMN group (Figure 4).*

Network Connectivity Predictive of Individual Speed of Forgetting

The cross-validation procedure yielded a hyperparameter value of $\lambda = 0.0044$, which resulted in only 8 regressors representing network connections being selected: it was the individual variations in these regressors that were found to be most predictive of individual speeds of forgetting. Table 1 lists these eight network connections, while Figure 7 provides a

visual breakdown of their connectivity, organized according to the different DMN subnetworks involved. All of the six networks whose connectivity to the DMN was identified as a reliable predictor of forgetting (Somatomotor A, Visual A, Salience A, Control C, Limbic A, and Dorsal Attention B: Table 1 and Figure 7) belong to the storage-related group (see Figure 4), suggesting that forgetting can be better understood within the storage decay framework. Further analysis of these connectivity patterns, described below, confirmed this hypothesis.

Figure 7: *Functional network connectivity predictive of individual speeds of forgetting. The connectivity values are organized according to the DMN subnetwork they refer to (in dark gray); The color of each network reflects the value of the corresponding β weight of its functional connectivity.*

Table 1: *List of network connections predictive of individual SOF values, together with their β weights.*

Network Connectivity	Regressor Weight (β)
Default A to Default C	0.055810
Default A to Limbic A	–0.014422
Default B to Control C	–0.022168
Default B to Salience / Ventral Attention A	0.025213
Default B to Visual Central (Visual A)	0.058361
Default C to Dorsal Attention B	–0.027954
Default C to Limbic A	0.017076
Default C to Somatomotor A	0.008430

The predictions from this model were then compared to the observed data. Figure 8A plots the predicted speeds of forgetting against the observed values estimated from the adaptive fact-learning task. The final model's predictions have a mean correlation of $r(33) = 0.77$ ($p < 0.001$) with the observed data.

Figure 8: *Observed vs. predicted speeds of forgetting across all 33 participants, using the eight regressors identified by the LASSO Model. Each dot represents a single participant; the black dashed line is the identity line, while the blue line and ribbon represent the linear regression estimate and its standard error.*

Validation of the LASSO Results

Given equally valid predictors, the penalty term might sometimes force the LASSO algorithm to select between predictors at random, providing an inadequate and partial picture of which regressors are indeed essential. If this were the case, other equally valid predictors could have been overlooked. A simple test to exclude this possibility is to remove the 8 connections from the original matrix of 45 regressors **X**, and re-run the cross-validation procedure. If the procedure converges on a different set of features, then other important regressors were

accidentally excluded. Alternatively, if no other regressors are left that can account for variability in the speed of forgetting, then it would be impossible for the cross-validation procedure to find a satisfying value of λ that allows for generalizable results.

When applied to our data, this test shows that the nested cross-validation procedure does *not* converge anymore. Specifically, it selects a large hyperparameter value ($\lambda = 0.0103$) under whose constraints all regressors are set to zero; under these conditions, the resulting linear model reduces to the intercept and fails to capture any individual differences in forgetting.

A second test consists of measuring the collinearity of the resulting model. Since the number of predictors in the model ($p = 8$) remains proportionally large when compared to the number of observations ($n = 33$), it is possible that the surviving regressors are highly collinear, which would suggest that all connectivity values are similarly related to the speed of forgetting. To exclude this hypothesis, the eight regressors in the model were examined using the Variance Inflation Factor (VIF) metric as implemented in R's *car* package [89]. The VIF of a regressor is the ratio of the variance of its estimate (that is, its β weight) in the full model over its variance in a reduced model that includes only that regressor. As a rule of thumb, VIF factors > 10 indicate collinearity [90]. This analysis revealed that all of the predictors had a VIF value < 2.5, which suggests that they contributed independently to an individual speed of forgetting.

Relative Importance of DMN Networks

Having confirmed the internal validity of the model, we can now further investigate the nature of its predictors. The connections in Table 1 can be thought of as weighted edges connecting nodes in a network, with the weights corresponding to their β values; therefore, they can be analyzed using network-level measures. In particular, it is possible to compute the relative contribution of each of the Yeo networks as the *importance* of the corresponding node, which is

the L1 norm of all of its connections to other nodes. In this case, each node is one of the Yeo

networks, and its importance is the sum of the absolute value of the regressors that connect it to

other Yeo networks in Table 1. As a first step, we computed the importance of each of the three

DMN subnetworks (Default A, B, and C in Figure 3). The results are shown in Figure 9.

Figure 9: *Relative importance of each DMN subnetwork, calculated as the L1 norm of its*

connections.

Relative Importance of Storage vs. Retrieval Networks

As noted above, all of the six non-default networks identified by the LASSO model

(Somatomotor A, Visual A, Salience A, Control C, Limbic A, and Dorsal Attention B) belong to

the storage-related group. Even accounting for the fact that our classification contains more

storage networks (eight, accounting for 24 connections to the DMN subnetworks) than retrieval

networks (six, for a total of 18 connections), the probability that *all* eight regressors would

involve only storage networks is $p < 0.003$.

A fuller picture, however, can be obtained by applying the importance measure to the

non-DMN networks in Table 1; in this case, the importance of each network is computed as the

L1 norm of its connectivity with any of the DMN subnetworks (Figure 10).

Figure 10: *Relative importance of each non-DMN subnetworks, calculated as the L1 norm of its connections.*

If the storage decay hypothesis is correct, we would expect that the storage-related networks of Figure 4 would not only be selected more frequently, but would also have significantly greater importance than the retrieval-related networks. This is indeed confirmed by Figure 11, which illustrates the distribution of the importance scores of *all* of the non-DMN networks. A Wilcoxon test further confirmed that, even when all of the non-DMN networks were included (using a regressor value of zero for those eliminated by LASSO), the mean importance of the storage network ($M = 0.021$, $SD = 0.019$) was significantly greater than zero ($t(7) = 3.179$, $p = 0.016$) and, as a consequence, greater than the mean across the retrieval networks (which is zero).

Figure 11: *Connectivity importance of storage and retrieval networks to predict memory performance.*

Discussion

In this paper, we provided evidence that an individual's speed of forgetting, operationalized as the decay intercept parameter [51] averaged over items, can be predicted using a distributed pattern of functional connectivity obtained from resting-state (or task-free) brain connectivity measures. This set of connections was identified through LASSO regression, demonstrating remarkable predictive accuracy while also reducing the full connectome to a manageable set of predictors. Functionally, these findings imply that forgetting primarily involves connectivity within the DMN and between the DMN and the storage-related networks, providing evidence in favor of the storage decay hypothesis over the retrieval failure account.

Limitations

While these findings are encouraging, it is important to recognize some of their inherent limitations—most importantly, that the data comes from a small number of fairly homogeneous participants who studied a relatively small set of foreign vocabulary items, which potentially limits external validity. Thus, future studies should expand the sample size and explore other sets of study materials.

The limited sample size also prevents us from entirely ruling out a role for retrieval failure. It is possible, for example, that the contribution of retrieval networks is significant and consistent but not visible in our limited sample. It is also possible that the relative contributions of storage decay and retrieval failure are mediated by external factors, such as age and clinical conditions. Thus, storage decay may be the main contributor to forgetting in healthy young adults (our sample's population), but retrieval failure might play a greater role in elderly adults or in patients affected by dementia. Once more, future studies will be needed to further examine these alternatives.

A related limitation concerns the interpretability of the specific connectivity results in Figure 7. It is impossible to judge whether a higher or lower beta value for a function connection translates to higher or lower forgetting, because the effect depends on the direction of the original functional connectivity. For example, if two regions were *anti*-correlated but their functional connectivity had a positive weight, an increase in functional connectivity would decrease, instead of increase, the predicted speed of forgetting. In a previous study by Yang et al. [91], the authors worked around this problem by multiplying the beta weight of a connection by the sign of its mean, group-level connectivity value, so that the sign of the product would

correctly reflect the direction of the effect. If applied to our data, this approach yields the results illustrated in Figure 12.

Figure 12: *Interpretable functional network connectivity predictive of individual speeds of forgetting, organized according to the DMN subnetwork they refer to (in dark gray). The color of each network reflects the value of the corresponding β weight multiplied by its mean functional connectivity; positive values are associated with increases (and negative values with decrease) in the Speed of Forgetting.*

The results in Figure 12 suggest that *greater* connectivity at rest between storage-related networks and the DMN are associated with increased forgetting—with the only exception of the connectivity between the salience network and the DMN. In turn, this suggests two possible interpretations. The first is that greater resting-state activity between these networks might be

associated with shallower *encoding* of the initial engram, thus setting the stage for faster forgetting. A second interpretation is that greater functional activity at rest would be associated with greater interference during the *maintenance* and consolidation stages—for example, by leading to greater interference during memory reactivation and replay. Although both hypotheses are worth pursuing in future research, we should note that this approach relies on having reliable estimates of the sign of group-level mean functional connectivity. Thus, given the limited size of our sample, the interpretation of the results in Figure 12 should be taken with caution.

Considerations

These limitations notwithstanding, we believe it is important to stress four characteristics of our findings. First, they are based on a measure that directly quantifies forgetting, a latent and unobservable process, through computational phenotyping; this approach provides an advantage over simpler behavioral tests of memory performance. As noted in the introduction, forgetting is a *process* and there is no consensus on how long after encoding a memory should be tested. Since long-term memory retention spans from seconds to decades, it is difficult to test it in a laboratory setting. Consequently, previous studies have relied on individuals exhibiting dramatic differences in memory function, such as amnesic patients (such as patients suffering from medial temporal lobe injury [92–94], while most laboratory experiments focus on relatively short retention intervals. Two participants, however, might have comparable memory performance a few minutes after studying but vastly different performances weeks later. In contrast, our findings rely on a novel metric, the speed of forgetting, that quantifies the rate at which memories are forgotten, and is better suited for making inferences as to where forgetting might happen in the brain.

Second, it is worth noting that our findings could be applied in the opposite direction: to decode an individual's speed of forgetting from distributed patterns of connectivity using a standard brain parcellation. Such an approach could be used, for example, to detect the early onset of abnormal forgetting in individuals with known memory pathologies and cognitive degenerative diseases. Most importantly, this finding suggests the possibility of decoding model parameters directly from resting state data, rather than from a combination of specific tasks. This procedure could play an important role in the development of high-fidelity idiographic models. This procedure also has the benefit of showing reflections of parameters in brain connectivity network regions, shedding light on a more holistic and comprehensive way of doing functional anatomy research on human cognition.

Third, our results are broadly consistent with the existing literature. As predicted, we did find that individual differences in forgetting covary with connectivity with and within the DMN. Not only have these sets of networks been theorized to play a role in memory replay and consolidation [10], but their connectivity with the salience network and the dorsal attention network (both of which were identified in our analysis) are altered in pathologies that negatively affect long-term memory function, such as Alzheimer's Disease [12], mild traumatic brain injury [95] and depression [96].

Finally, our results are also consistent with the findings of a previous study that previously attempted to identify the neural underpinnings of forgetting using the *same* model-based approach but with a different neuroimaging modality—EEG instead of fMRI [61]. In that study, a significant positive correlation between spectral power in the beta band (12-20 Hz) and the speed of forgetting was observed distributing at middle frontal (AF3 and AF4), bilateral

posterior (channel O1, O2), and parietal (channel P8) regions; these spatial locations are consistent with the cortical regions of the DMN subnetworks A and B (see Figures 10 and 11).

Conclusions

In summary, although further research will be needed to clarify the specific mechanisms of forgetting, our findings do provide a preliminary answer to the question of *where* memories are lost—that is, in the connections between the DMN and the sensory networks that are involved in the encoding, re-experience, and consolidation of a memory. Thus, our results imply that forgetting can be interpreted as a storage decay process, rather than a retrieval access problem, and that future clinical and neuroscientific research should focus on the storage networks as a target to better understand, improve, or support memory function.

<div align="center">**Acknowledgments**</div>

This work was supported by grant N00014-17-1-2607 from the Office of Naval Research to CP and by grant FA9550-19-1-0299 from the Air Force Office of Scientific Research to AS.

<div align="center">**References**</div>

1. Melton AW. Implications of short-term memory for a general theory of memory. J Verbal Learning Verbal Behav. 1963;2: 1–21.

2. Feld GB, Born J. Sculpting memory during sleep: concurrent consolidation and forgetting. Current Opinion in Neurobiology. 2017;44: 20–27.

3. Hardt O, Nader K, Nadel L. Decay happens: the role of active forgetting in memory. Trends Cogn Sci. 2013;17: 111–120.

4. Guskjolen A, Cembrowski MS. Engram neurons: Encoding, consolidation, retrieval, and forgetting of memory. Mol Psychiatry. 2023;28: 3207–3219.

5. Shiffrin RM. Forgetting: Trace Erosion or Retrieval Failure? Science. 1970;168: 1601–

1603.

6.   Spear NE. Forgetting as retrieval failure. Animal memory. 1971; 45–109.

7.   Davis RL, Zhong Y. The biology of forgetting—A perspective. Neuron. 2017;95: 490–503.

8.   Ryan TJ, Frankland PW. Forgetting as a form of adaptive engram cell plasticity. Nat Rev
      Neurosci. 2022;23: 173–186.

9.   Frankland PW, Köhler S, Josselyn SA. Hippocampal neurogenesis and forgetting. Trends
      Neurosci. 2013;36: 497–503.

10.   Pezzulo G, Zorzi M, Corbetta M. The secret life of predictive brains: what's spontaneous
      activity for? Trends Cogn Sci. 2021;25: 730–743.

11.   Jin M, Pelak VS, Cordes D. Aberrant default mode network in subjects with amnestic mild
      cognitive impairment using resting-state functional MRI. Magn Reson Imaging. 2012;30:
      48–61.

12.   Qi H, Liu H, Hu H, He H, Zhao X. Primary Disruption of the Memory-Related Subsystems
      of the Default Mode Network in Alzheimer's Disease: Resting-State Functional
      Connectivity MRI Study. Front Aging Neurosci. 2018;10: 344.

13.   Penfield W. Engrams in the human brain. Mechanisms of memory. Proc R Soc Med.
      1968;61: 831–840.

14.   Roy DS, Arons A, Mitchell TI, Pignatelli M, Ryan TJ, Tonegawa S. Memory retrieval by
      activating engram cells in mouse models of early Alzheimer's disease. Nature. 2016;531:
      508–512.

15.   Swanson CJ, Zhang Y, Dhadda S, Wang J, Kaplow J, Lai RYK, et al. A randomized,
      double-blind, phase 2b proof-of-concept clinical trial in early Alzheimer's disease with
      lecanemab, an anti-Aβ protofibril antibody. Alzheimers Res Ther. 2021;13: 80.

16. Cummings J, Apostolova L, Rabinovici GD, Atri A, Aisen P, Greenberg S, et al. Lecanemab: Appropriate Use Recommendations. J Prev Alzheimers Dis. 2023;10: 362–377.

17. Danker JF, Anderson JR. The ghosts of brain states past: remembering reactivates the brain regions engaged during encoding. Psychol Bull. 2010;136: 87–102.

18. Huang Q, Xiao Z, Yu Q, Luo Y, Xu J, Qu Y, et al. Replay-triggered brain-wide activation in humans. Nat Commun. 2024;15: 7185.

19. Ranganath C, Ritchey M. Two cortical systems for memory-guided behaviour. Nat Rev Neurosci. 2012;13: 713–726.

20. Kaefer K, Stella F, McNaughton BL, Battaglia FP. Replay, the default mode network and the cascaded memory systems model. Nat Rev Neurosci. 2022;23: 628–640.

21. Raichle ME, Snyder AZ. A default mode of brain function: a brief history of an evolving idea. Neuroimage. 2007;37: 1083–90; discussion 1097–9.

22. Sibert C, Hake HS, Stocco A. The Structured Mind at Rest: Low-Frequency Oscillations Reflect Interactive Dynamics Between Spontaneous Brain Activity and a Common Architecture for Task Control. Front Neurosci. 2022;16: 832503.

23. Petrides M. The mid-ventrolateral prefrontal cortex and active mnemonic retrieval. Neurobiol Learn Mem. 2002;78: 528–538.

24. Badre D, Wagner AD. Left ventrolateral prefrontal cortex and the cognitive control of memory. Neuropsychologia. 2007;45: 2883–2901.

25. Thompson-Schill SL, D'Esposito M, Aguirre GK, Farah MJ. Role of left inferior prefrontal cortex in retrieval of semantic knowledge: a reevaluation. Proc Natl Acad Sci U S A. 1997;94: 14792–14797.

26. Danker JF, Gunn P, Anderson JR. A rational account of memory predicts left prefrontal

activation during controlled retrieval. Cereb Cortex. 2008;18: 2674–2685.

27. Borst JP, Anderson JR. Using model-based functional MRI to locate working memory updates and declarative memory retrievals in the fronto-parietal network. Proc Natl Acad Sci U S A. 2013;110: 1628–1633.

28. Danker JF, Fincham JM, Anderson JR. The neural correlates of competition during memory retrieval are modulated by attention to the cues. Neuropsychologia. 2011;49: 2427–2438.

29. Lepage M, Ghaffar O, Nyberg L, Tulving E. Prefrontal cortex and episodic memory retrieval mode. Proc Natl Acad Sci U S A. 2000;97: 506–511.

30. Chua EF, Schacter DL, Sperling RA. Neural correlates of metamemory: a comparison of feeling-of-knowing and retrospective confidence judgments. J Cogn Neurosci. 2009;21: 1751–1765.

31. Modirrousta M, Fellows LK. Medial prefrontal cortex plays a critical and selective role in "feeling of knowing" meta-memory judgments. Neuropsychologia. 2008;46: 2958–2965.

32. Schnyer DM, Verfaellie M, Alexander MP, LaFleche G, Nicholls L, Kaszniak AW. A role for right medial prefrontal cortex in accurate feeling-of-knowing judgments: evidence from patients with lesions to frontal cortex. Neuropsychologia. 2004;42: 957–966.

33. Turner MS, Cipolotti L, Yousry TA, Shallice T. Confabulation: damage to a specific inferior medial prefrontal system. Cortex. 2008;44: 637–648.

34. Benson DF, Djenderedjian A, Miller BL, Pachana NA, Chang L, Itti L, et al. Neural basis of confabulation. Neurology. 1996;46: 1239–1243.

35. Raichle ME. The brain's default mode network. Annu Rev Neurosci. 2015;38: 433–447.

36. McClelland JL, McNaughton BL, O'Reilly RC. Why there are complementary learning systems in the hippocampus and neocortex: insights from the successes and failures of

connectionist models of learning and memory. Psychol Rev. 1995;102: 419.

37. O'Reilly RC, Bhattacharyya R, Howard MD, Ketz N. Complementary learning systems. Cogn Sci. 2014;38: 1229–1248.

38. Alvarez P, Squire LR. Memory consolidation and the medial temporal lobe: a simple network model. Proc Natl Acad Sci U S A. 1994;91: 7041–7045.

39. Zhang M, Nathaniel U, Savill N, Smallwood J, Jefferies E. Intrinsic connectivity of left ventrolateral prefrontal cortex predicts individual differences in controlled semantic retrieval. Neuroimage. 2022;246: 118760.

40. Loftus GR. On interpretation of interactions. Mem Cognit. 1978;6: 312–319.

41. Brady TF, Robinson MM, Williams JR, Wixted JT. Measuring memory is harder than you think: How to avoid problematic measurement practices in memory research. Psychon Bull Rev. 2023;30: 421–449.

42. Patzelt EH, Hartley CA, Gershman SJ. Computational phenotyping: Using models to understand individual differences in personality, development, and mental illness. Personal Neurosci. 2018;1: e18.

43. Schurr R, Reznik D, Hillman H, Bhui R, Gershman SJ. Dynamic computational phenotyping of human cognition. Nat Hum Behav. 2024;8: 917–931.

44. Sense F, Behrens F, Meijer RR, van Rijn H. An individual's rate of forgetting is stable over time but differs across materials. Top Cogn Sci. 2016;8: 305–321.

45. Xu Y, Stocco A. Recovering Reliable Idiographic Biological Parameters from Noisy Behavioral Data: the Case of Basal Ganglia Indices in the Probabilistic Selection Task. Comput Brain Behav. 2021; 1–17.

46. Marek S, Tervo-Clemmens B, Calabro FJ, Montez DF, Kay BP, Hatoum AS, et al.

Reproducible brain-wide association studies require thousands of individuals. Nature. 2022;603: 654–660.

47. Daw ND. Trial-by-trial data analysis using computational models. Decision making, affect, and learning: Attention and performance XXIII. 2011;23: 3–38.

48. Collins AGE. The Tortoise and the Hare: Interactions between Reinforcement Learning and Working Memory. J Cogn Neurosci. 2018;30: 1422–1432.

49. Wagenmakers E-J, Krypotos A-M, Criss AH, Iverson G. On the interpretation of removable interactions: a survey of the field 33 years after Loftus. Mem Cognit. 2012;40: 145–160.

50. Anderson JR, Schooler LJ. Reflections of the Environment in Memory. Psychol Sci. 1991;2: 396–408.

51. Pavlik PI Jr, Anderson JR. Practice and forgetting effects on vocabulary memory: An activation-based model of the spacing effect. Cogn Sci. 2005;29: 559–586.

52. Pavlik PI, Anderson JR. Using a model to compute the optimal schedule of practice. J Exp Psychol Appl. 2008;14: 101–117.

53. Sense F, Meijer RR, van Rijn H. Exploration of the Rate of Forgetting as a Domain-Specific Individual Differences Measure. Frontiers in Education. 2018;3: 112.

54. van der Velde M, Sense F, Borst JP, van Rijn H. Large-scale evaluation of cold-start mitigation in adaptive fact learning: Knowing "what" matters more than knowing "who." User Model User-adapt Interact. 2024. doi:10.1007/s11257-024-09401-5

55. Nadel L, Samsonovich A, Ryan L, Moscovitch M. Multiple trace theory of human memory: computational, neuroimaging, and neuropsychological results. Hippocampus. 2000;10: 352–368.

56. Moscovitch M, Rosenbaum RS, Gilboa A, Addis DR, Westmacott R, Grady C, et al.

Functional neuroanatomy of remote episodic, semantic and spatial memory: a unified account based on multiple trace theory. J Anat. 2005;207: 35–66.

57. van der Velde M, Sense F, Borst JP, van Rijn H. Alleviating the Cold Start Problem in Adaptive Learning using Data-Driven Difficulty Estimates. Computational Brain & Behavior. 2021;4: 231–249.

58. Sense F, Maaß S, Gluck K, van Rijn H. Within-Subject Performance on a Real-Life, Complex Task and Traditional Lab Experiments: Measures of Word Learning, Raven Matrices, Tapping, and CPR. J Cogn. 2019;2: 12.

59. Sense F, Behrens F, Meijer RR, van Rijn H. Stability of Individual Parameters in a Model of Optimal Fact Learning. Proceedings of the 13th International Conference on Cognitive Modeling. 2015. pp. 136–141.

60. Shrout PE, Lane SP. Psychometrics. In: Mehl MR, editor. Handbook of research methods for studying daily life , (pp. New York, NY, US: The Guilford Press, xxvii; 2012. pp. 302–320.

61. Zhou P, Sense F, van Rijn H, Stocco A. Reflections of idiographic long-term memory characteristics in resting-state neuroimaging data. Cognition. 2021;212: 104660.

62. James G, Witten D, Hastie T, Tibshirani R. Statistical Learning. In: James G, Witten D, Hastie T, Tibshirani R, editors. An Introduction to Statistical Learning: with Applications in R. New York, NY: Springer New York; 2013. pp. 15–57.

63. Braun U, Plichta MM, Esslinger C, Sauer C, Haddad L, Grimm O, et al. Test-retest reliability of resting-state connectivity network characteristics using fMRI and graph theoretical measures. Neuroimage. 2012;59: 1404–1412.

64. Stocco A, Sibert C, Steine-Hanson Z, Koh N, Laird JE, Lebiere CJ, et al. Analysis of the

human connectome data supports the notion of a "Common Model of Cognition" for human and human-like intelligence across domains. Neuroimage. 2021;235: 118035.

65. Finn ES, Shen X, Scheinost D, Rosenberg MD, Huang J, Chun MM, et al. Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. Nat Neurosci. 2015;18: 1664–1671.

66. Waller L, Walter H, Kruschwitz JD, Reuter L, Müller S, Erk S, et al. Evaluating the replicability, specificity, and generalizability of connectome fingerprints. Neuroimage. 2017;158: 371–377.

67. Avery EW, Yoo K, Rosenberg MD, Greene AS, Gao S, Na DL, et al. Distributed Patterns of Functional Connectivity Predict Working Memory Performance in Novel Healthy and Memory-impaired Individuals. J Cogn Neurosci. 2020;32: 241–255.

68. Reineberg AE, Andrews-Hanna JR, Depue BE, Friedman NP, Banich MT. Resting-state networks predict individual differences in common and specific aspects of executive function. Neuroimage. 2015;104: 69–78.

69. Reineberg AE, Banich MT. Functional connectivity at rest is sensitive to individual differences in executive function: A network analysis. Hum Brain Mapp. 2016;37: 2959–2975.

70. McGregor HR, Gribble PL. Functional connectivity between somatosensory and motor brain areas predicts individual differences in motor learning by observing. J Neurophysiol. 2017;118: 1235–1243.

71. Baldassarre A, Lewis CM, Committeri G, Snyder AZ, Romani GL, Corbetta M. Individual variability in functional connectivity predicts performance of a perceptual task. Proc Natl Acad Sci U S A. 2012;109: 3516–3521.

72. Persson J, Stening E, Nordin K, Söderlund H. Predicting episodic and spatial memory performance from hippocampal resting-state functional connectivity: Evidence for an anterior-posterior division of function. Hippocampus. 2018;28: 53–66.

73. Durrant S, Lewis PA. Memory consolidation: tracking transfer with functional connectivity. Curr Biol. 2009;19: R860–2.

74. Van den Broek GSE, Segers E, Van Rijn H, Takashima A, Verhoeven L. Effects of elaborate feedback during practice tests: Costs and benefits of retrieval prompts. J Exp Psychol Appl. 2019;25: 588–601.

75. Nelson TO, Dunlosky J. Norms of paired-associate recall during multitrial learning of Swahili-English translation equivalents. Memory. 1994;2: 325–335.

76. Anderson JR. Retrieval of propositional information from long-term memory. Cogn Psychol. 1974;6: 451–474.

77. Anderson JR. How Can the Mind Occur in the Physical Universe? Oxford University Press; 2007.

78. Anderson JR, Bothell D, Byrne MD, Douglass S, Lebiere C, Qin Y. An integrated theory of the mind. Psychol Rev. 2004;111: 1036–1060.

79. Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. Comput Biomed Res. 1996;29: 162–173.

80. Yeo BTT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, et al. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. J Neurophysiol. 2011;106: 1125–1165.

81. Sporns O. The human connectome: a complex network. Ann N Y Acad Sci. 2011;1224: 109–125.

82. Cole MW, Ito T, Bassett DS, Schultz DH. Activity flow over resting-state networks shapes cognitive task activations. Nat Neurosci. 2016;19: 1718–1726.

83. Shen X, Finn ES, Scheinost D, Rosenberg MD, Chun MM, Papademetris X, et al. Using connectome-based predictive modeling to predict individual behavior from brain connectivity. Nat Protoc. 2017;12: 506–518.

84. Tibshirani R. Regression shrinkage and selection via the lasso. J R Stat Soc Series B Stat Methodol. 1996. Available: https://academic.oup.com/jrsssb/article-abstract/58/1/267/7027929

85. James G, Witten D, Hastie T, Tibshirani R. An Introduction to Statistical Learning. Springer US; 2013.

86. James G, Witten D, Hastie T, Tibshirani R, Taylor J. An Introduction to Statistical Learning: with Applications in Python. Springer Nature; 2023.

87. Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME. The human brain is intrinsically organized into dynamic, anticorrelated functional networks. Proc Natl Acad Sci U S A. 2005;102: 9673–9678.

88. Jia J, Yu B. On model selection consistency of the Elastic Net when p □ n. Stat Sin. 2010;20: 595–611.

89. Fox J, Weisberg S. An R Companion to Applied Regression. SAGE Publications; 2018.

90. Fox J, Monette G. Generalized Collinearity Diagnostics. J Am Stat Assoc. 1992;87: 178–183.

91. Yang YC, Sibert C, Stocco A. Reliance on Episodic vs. Procedural Systems in Decision-Making Depends on Individual Differences in Their Relative Neural Efficiency. Computational Brain & Behavior. 2024. doi:10.1007/s42113-024-00203-x

92. Scoville WB, Milner B. Loss of recent memory after bilateral hippocampal lesions. J Neurol Neurosurg Psychiatry. 1957;20: 11–21.

93. Corkin S. What's new with the amnesic patient H.M.? Nat Rev Neurosci. 2002;3: 153–160.

94. Gabrieli JD, Cohen NJ, Corkin S. The impaired learning of semantic knowledge following bilateral medial temporal-lobe resection. Brain Cogn. 1988;7: 157–177.

95. Andrews MJ, Salat DH, Milberg WP, McGlinchey RE, Fortier CB. Poor sleep and decreased cortical thickness in veterans with mild traumatic brain injury and post-traumatic stress disorder. Mil Med Res. 2024;11: 51.

96. Mulders PC, van Eijndhoven PF, Schene AH, Beckmann CF, Tendolkar I. Resting-state functional connectivity in major depressive disorder: A review. Neurosci Biobehav Rev. 2015;56: 330–344.